p53 deletion rescues lethal microcephaly in a mouse model with neural stem cell abscission defects

Jessica Neville Little[1] and Noelle D. Dwyer[1,2]

[1] Department of Cell Biology, University of Virginia School of Medicine, Charlottesville, VA, USA

[2] Corresponding author. Email: ndwyer@virginia.edu

ABSTRACT

Building a cerebral cortex of the proper size involves balancing rates and timing of neural stem cell (NSC) proliferation, neurogenesis, and cell death. The cellular mechanisms connecting genetic mutations to brain malformation phenotypes are still poorly understood. Microcephaly may result when NSC divisions are too slow, produce neurons too early, or undergo apoptosis, but the relative contributions of these cellular mechanisms to various types of microcephaly are not understood. We previously showed that mouse mutants in *Kif20b* (formerly called *Mphosph1*, *Mpp1*, or *KRMP1*) have small cortices that show elevated apoptosis, and defects in maturation of NSC midbodies, which mediate cytokinetic abscission. Here we test the contribution of intrinsic NSC apoptosis to brain size reduction in this lethal microcephaly model. By making double mutants with the pro-apoptotic genes *Bax* and *Trp53 (p53)*, we find that apoptosis of cortical NSCs accounts for most of the microcephaly, but that there is a significant apoptosis-independent contribution as well. Remarkably, heterozygous *p53* deletion is sufficient to fully rescue survival of the *Kif20b* mutant into adulthood. In addition, the NSC midbody maturation defects are not rescued by *p53* deletion, showing that they are either upstream of p53 activation, or in a parallel pathway. Thus, this work potentially identifies a novel midbody-mediated pathway for p53 activation, and elucidates both NSC apoptosis and abscission mechanisms that could underlie human microcephaly or other brain malformations.

INTRODUCTION

Human genetics is increasingly successful at linking specific gene mutations to

congenital brain malformations and other neurodevelopmental disorders. However, the cellular

mechanisms connecting genetic mutations to brain phenotypes are still poorly understood.

There are 17 human primary microcephaly genes identified, and there are many syndromes that

feature microcephaly (Duerinckx and Abramowicz, 2018; Online Mendelian Inheritance in Man

(OMIM) database). Thus it is a heterogeneous disorder. Known microcephaly genes encode

proteins with diverse molecular functions, but many are involved in cell division.

Cell division genes may be prominent in brain malformations like microcephaly because

cortical neural stem cells (NSC) divisions have several unusual features (Dwyer et al., 2016).

NSCs are tall, thin cells that reside in the pseudostratified epithelium of the cortex, with their

apical endfeet forming the ventricular surface, and their basal processes stretching to the basal

lamina beneath the meninges. Their nuclei undergo interkinetic nuclear migration during the cell

cycle, moving basally for S-phase and to the apical membrane for M-phase and cytokinesis. In

addition, NSC divisions can produce symmetric or asymmetric daughter fates, giving rise to

more NSCs, neurons, intermediate progenitors, and glia during corticogenesis. These stem cells

must produce the right types and numbers of daughter cells within specific windows of time.

With all these complex demands, the developing cortex is particularly vulnerable to insults to

cell division.

We previously identified a mouse model of microcephaly that is recessive, perinatal

lethal, and relatively severe, with a brain about half as thick as normal during late gestation. It

carries a loss-of-function mutation in the kinesin microtubule motor gene *Kif20b*. The mutant

brains do not display NSC mitotic arrest or abnormal cleavage angles, which have been noted

in other microcephaly mutants. Instead, *Kif20b* mutant brains display defects in cytokinetic

abscission (Dwyer et al., 2011; Janisch et al., 2013). Abscission is the process of severing the

connection between mother and daughter cell, taking an hour or more after telophase (Mierzwa

and Gerlich, 2014). The cleavage furrow compacts the central spindle microtubules into the

midbody, which mediates abscission by recruiting proteins to remodel and "cut" the cytoskeleton

and membrane. We showed that Kif20b protein localizes to the central spindle and midbody in

human cell lines and mouse NSCs (Janisch, et al., 2013; 2018). Furthermore, Kif20b appears to

facilitate changes in midbody microtubule structure as the midbody "matures" during the

abscission process, and ensures timely abscission in cell lines. Suggesting that it may

accelerate cell division, *Kif20b* is elevated in some cancers ((Kanehira et al., 2007; Liu et al.,

2014; Lin et al., 2018). Interestingly, *Kif20b* evolved with the vertebrate lineage, so its subtle role

in abscission may be important for growing bigger, more complex nervous systems.

In addition to abnormal midbodies, *Kif20b* mutants (*Kif20b-/-*) also display increased

apoptosis in the embryonic cortex from E10.5 to E16.5 (Janisch et al., 2013). However, it was

unclear whether the relatively small amount of detectable apoptosis observed could account for

the severity of the microcephaly. Apoptosis appears to be relatively low in the healthy embryonic

neocortex, but was seen to be elevated in some mouse models of brain malformations

(Marthiens et al., 2013; Stottmann et al., 2013; Chen et al., 2014; Insolera et al., 2014;

Marjanovic et al., 2015; Breuss et al., 2016). The intrinsic or stress-induced apoptotic pathway

can be triggered by environmental stresses or genetic insults, such as particular mutations

(Arya and White 2015). Bax and p53 (gene *Trp53*) are expressed in embryonic brain and

appear to respond to damage. In response to apoptotic stimuli, Bax, a multi-BH domain-

containing member of the BCL2 family, can form pores across the outer mitochondrial

membrane, thereby releasing cytochrome C that can activate the caspase cascade. p53, a

tumor suppressor mutated in many human cancers, is an upstream activator of Bax

(Kastenhuber and Lowe 2017). Mouse knockouts of *Bax* and *p53* have normal brain

development with surprisingly few low penetrance defects (Knudson et al., 1995; Jacks et al.,

1994; Insolera et al., 2014).

Here, we set out to determine the relative contributions of apoptosis and abscission

dysregulation to the microcephaly of the *Kif20b* mutant, and to understand the relationship

between these phenotypes. To do this, we crossed genetic mutants of the intrinsic apoptosis

pathway to *Kif20b-/-* mice and asked whether the apoptosis, microcephaly, and abscission

defects in *Kif20b* mutants were rescued, unaffected, or potentially worsened. To our surprise,

we found that *p53* deletion prevented apoptosis and rescued brain size and structure to a

remarkable degree. A partial apoptosis inhibition by *Bax* deletion correlated with a lesser extent

of brain size rescue. Surprisingly, deletion of even one allele of *p53* is able to completely block

the apoptosis and lethality in *Kif20b* mutants. However, *Kif20b; p53* double mutant brains are

still smaller than controls at birth, suggesting Kif20b regulates cortical development through

additional mechanisms, perhaps by ensuring timely abscission. Indeed, *p53* deletion does not

rescue the defects in midbody structure seen in *Kif20b* mutant NSCs, and additional midbody

defects are revealed when *p53* is deleted. Our data provides the first evidence that abscission

defects can cause p53 activation in NSCs or any cell type, through a yet-to-be identified

pathway. Finally, our work addresses apoptosis inhibition as a potential way to ameliorate the

severity of microcephaly caused by genetic, viral, or environmental insults.

RESULTS

The intrinsic apoptotic pathway is the key driver of microcephaly in *Kif20b-/-* mice

To test whether the intrinsic apoptotic pathway mediates the elevated apoptosis and

microcephaly in the *Kif20b* mutant cortex, we utilized genetic crosses to mutants in two key

genes in this pathway, *Bax* and *p53 (Trp53)*. Importantly, homozygous deletion mutants in *Bax*

or *p53* develop properly with normal size cortices (Knudson et al., 1995; Jacks et al., 1994;

Insolera et al., 2014). First, we produced *Kif20b*; *Bax* double mutants. Bax, together with its

partner Bak, increases the permeability of the mitochondrial membrane, increasing cytochrome

c release (Westphal et al., 2011). We examined double mutant brains at age E14.5, when

apoptosis is elevated 4-fold in *Kif20b* mutants, and the cortical plate has started to form.

Interestingly, embryos carrying homozygous mutations in both *Bax* and *Kif20b* showed a partial

block of the elevated apoptosis and a partial rescue of cortical thickness (**Fig. 1A-H**).

Craniofacial defects observed in *Kif20b* mutants were not rescued by Bax mutation (data not

shown). These data suggest that apoptosis and microcephaly are correlated, but that additional

proteins are required for the full apoptotic response to *Kif20b* loss. For example, Bak is a

partner of Bax and is partially redundant (Lindsten et al., 2000).

The tumor suppressor p53 is upstream of Bax/Bak in the intrinsic apoptotic pathway.

Therefore, we tested whether *p53* deletion could fully block the apoptosis in *Kif20b* mutants by

crossing to the *Trp53-/-* mutant (Jacks et al., 1994). Strikingly, apoptosis and cortical thickness

of E14.5 *Kif20b-/-* embryos are both fully rescued by either heterozygous or homozygous

deletion of *p53* (**Fig. 2A-J**). Thus, two functional *p53* genes are required to produce the

apoptosis and microcephaly triggered by *Kif20b* loss. Furthermore, *p53* is required for the

craniofacial defects of *Kif20b-/-* embryos: either heterozygous or homozygous *p53* deletion

significantly ameliorates the small eye and snout phenotypes (**Fig. S1**). p53 protein appears to

be elevated in *Kif20b* mutant cortices, as immunoblots of E12.5 cortical lysates show p53 band

intensity increased 50% in *Kif20b-/-* samples, normalized to the NSC protein beta-catenin (**Fig.**

S2 A, B). Furthermore, immunostaining suggests this increase is specifically in the nuclei of

NSCs (**Fig S2 C, D**). Together, these data show that p53 is activated when *Kif20b* is lost, and

p53 function is required for the excess apoptosis and microcephaly in this mutant. Moreover, the

partial and full rescues of microcephaly by *Bax* and *p53* deletion, respectively, show that that

the amount of apoptosis and the severity of microcephaly are strongly correlated, suggesting

that apoptosis is the key cellular mechanism driving microcephaly in *Kif20b-/-* mice.

143 *p53* **deletion restores growth of neuronal and subventricular layers in embryonic** *Kif20b-*
144 */- cortex*

145   We previously showed that the reduced cortical thickness *of Kif20b* mutants is due to
146 thinner neuronal layers and fewer intermediate progenitors (IPs) (Janisch et al., 2013). To
147 determine if *p53* deletion could rescue cortical neurogenesis and structure as well as thickness,
148 we labeled *Kif20b-/-* single mutant and *Kif20b*; *p53* double mutant E14.5 cortical sections for
149 Pax6, Tuj1, and Tbr2, to label NSCs, neurons, and IPs, respectively. In *Kif20b-/-* single mutant
150 brains, as expected, the neuronal layer (cortical plate) and axonal layer (intermediate zone) are
151 thin (**Fig. 3A, B, G**). Remarkably, *Kif20b-/-; p53-/-* embryos have cortices that appear to have
152 normal organization, with cortical plates and intermediate zones indistinguishable from controls
153 (**Fig. 3C, G**). Additionally, *Kif20b-/-; p53-/-* embryos display normal ventricular zone thickness
154 and IP generation (**Fig. 3D-F, H, I**). Therefore, blocking apoptosis by *p53* deletion in the cortices
155 of *Kif20b-/-* brains increased cortical thickness by improving production or survival of neurons
156 and IPs. Furthermore, the rescued neurons and IPs appear to migrate normally and create a
157 normal-appearing structure.

158

159 *p53* **deletion rescues postnatal survival but not full cortical size of** *Kif20b-/-* **mice at birth**
160   In addition to reduced brain size, *Kif20b-/-* mice exhibit perinatal lethality. Remarkably,
161 while no *Kif20b* mutant mice with wild-type *p53* status survive the day of birth (postnatal day 0,
162 "P0"), *Kif20b-/-* mice with heterozygous or homozygous *p53* deletion survive postnatally at
163 expected Mendelian ratios (**Figure 4A**). Even more surprising, most *Kif20-/- ; p53+/-* mice and
164 *Kif20b-/- ; p53-/-* mice live to adulthood and are fertile. Some *Kif20b-/-; p53+/-* mice (~15%) still
165 have visible craniofacial defects including a small eye or short snout, and ~5% have
166 hydrocephalus, but the majority have normal facial structure. Double homozygotes (*Kif20b-/-;*
167 *p53-/-*) have even fewer craniofacial defects, but die prematurely at ~ 3 - 4 months of age due to
168 spontaneous tumors, as do *p53-/-* single mutants (Jacks et al., 1994).
169   The postnatal survival of *Kif20b-/-; p53-/-* mice enabled us to investigate the requirement
170 of *Kif20b* for corticogenesis without the confounding factor of excess apoptosis. First, we
171 confirmed previous reports that *p53* knockout mice have normal cortical size at P0 (**Fig. 4 B, C**)
172 (Insolera et al., 2014). Interestingly, in contrast to the full rescue of cortical thickness seen at
173 E14.5 in *Kif20b-/-; p53-/-* brains, at P0 they show a 10% decrease in cortical length and a 20%
174 reduction in cortical thickness compared to heterozygous controls (**Fig. 4 B- F**). Furthermore,
175 the reduction is in the cortical plate and intermediate zone, but not the ventricular and
176 subventricular zones (**Fig. 4G**). These data indicate that despite the dramatic improvements in

cortical growth and postnatal survival afforded by blocking p53-dependent apoptosis, a deficit in

corticogenesis remains. Thus, while elevated apoptosis largely accounts for the microcephaly of

the *Kif20b* mutant, proper cortical growth requires a *Kif20b* function that cannot be compensated

by preventing apoptosis.

***Kif20b* is required cell-autonomously for midbody maturation of cortical NSCs**

Previously we showed that *Kif20b* is expressed in germinal zones of the embryonic

brain, and that Kif20b protein localizes to midbodies of dividing embryonic cortical neural stem

cells at the ventricular surface (Janisch et al., 2013). We further showed that *Kif20b* mutants

have abnormalities in NSC midbodies: they tend to be wider and less aligned to the apical

membrane. These phenotypes could be due to a cell-autonomous requirement for Kif20b during

abscission, or due to non-cell autonomous effects through cell-cell interactions at the apical

membrane junctions, because cytokinesis in epithelia is a multicellular process (Herszterg et al.,

2014). To further probe Kif20b's function in embryonic NSC division, we used dissociated

cortical cell cultures.

Midbodies at various stages of maturation can be detected with tubulin and Aurora

Kinase B (AurKB) staining by their characteristic shapes (**Fig. 5A-C**). Early-stage midbodies are

wide (**Fig. 5A**), but become thinner as the midbody matures (Guizetti et al., 2011). At late

stages, microtubule constriction sites (abscission sites) are detectable on one or both sides of

the midbody center (**Fig. 5, B, C**, arrows) (Mierzwa and Gerlich et al., 2014). We found in

human cell lines that Kif20b is recruited to early stage midbodies, and at late stages

accumulates around the constriction sites (Janisch et al., 2018). Interestingly, *Kif20b-/-* NSC

cultures have an increased frequency of wide midbodies (**Fig. 5D**), and fewer midbodies with at

least one constriction site (**Fig. 5E**). These analyses show that *Kif20b* is required for normal

NSC midbody maturation, and that this requirement is cell-autonomous.

Some other mouse models of microcephaly show increased mitotic index in the cortex,

due to NSC mitosis delay or arrest (Marthiens et al., 2013; Chen et al., 2014; Insolera et al.,

2014; Marjanovic et al., 2015; Breuss et al., 2016). We showed previously that the *Kif20b*

mutant cortex does not have increased mitotic index (Janisch et al., 2013), but here we use the

dissociated cortical cell cultures to address this possibility with higher cellular resolution. We

assayed whether *Kif20b* mutant NSCs spent more time in mitosis or abscission than control

NSCs by determining the mitotic and midbody index in cultures. Surprisingly, among cycling

NSCs (Ki67+) in *Kif20b* mutant cultures, both the mitotic index and midbody index were not

increased but were actually slightly reduced (**Fig. S3B**). This could suggest that *Kif20b* mutant

NSCs undergo mitosis and abscission more rapidly than control cells, or that some undergo

apoptosis. To analyze relative durations of cell division phases, we categorized mitotic and

midbody stage NSCs into sub-stages by PH3 and AurKB appearance (**Fig. S3A, C**). Among

mutant NSCs at some stage of cell division, the percentages in prophase or

prometa/metaphase were not different, but the percentage in anaphase/early telophase was

slightly increased in *Kif20b* mutant cultures. These data are consistent with our previous results

that early steps of cell division are not disrupted in *Kif20b* mutant NSCs, but that cytokinesis is

affected (Janisch et al 2018).

***Kif20b* loss activates apoptosis cell-autonomously in proliferating cortical NSCs**

An attractive hypothesis is that defective midbody maturation of NSCs in *Kif20b* mutants

activates the intrinsic pathway of apoptosis in NSCs. However, our previous *in vivo* analysis

could not distinguish whether the apoptotic cells in *Kif20b* mutant brains were NSCs or neurons,

and whether apoptosis was triggered autonomously or by cell-cell interactions. Therefore, we

used the dissociated cultures to distinguish these possibilities. Indeed, in cultures of *Kif20b*

mutant cells after one day *in vitro*, apoptosis is elevated more than two-fold over controls (**Fig.**

6A, B, E; arrows, CC3+ cells), suggesting that apoptosis is triggered autonomously.

Furthermore, neurons (TuJ1+) do not display increased apoptosis, while NSCs (Nestin+) do,

and cycling NSCs (Ki67+) show a more pronounced increase (**Fig. 6 C, D, F**). This may explain

the reduced mitotic and midbody indices among mutant NSCs (**Fig. S2B**), as some arrest in the

cell cycle or die. Finally, since the elevated apoptosis in *Kif20b-/-* brains requires *p53* (**Fig. 2**),

we wondered whether dissociated *Kif20b-/-* NSC apoptosis also requires p53. Indeed, in

cultures from double mutant mice, the rate of apoptosis is similar to controls, showing that the

elevated apoptosis in isolated *Kif20b-/-* NSCs is also p53-dependent **(Figure 6G).** Together,

these data show that loss of *Kif20b* causes cell-intrinsic, p53-dependent apoptosis, specifically

in cycling NSCs but not postmitotic neurons.

***p53* deletion does not rescue impaired abscission in *Kif20b* mutant mice**

The preceding data show that in *Kif20b* mutant brains, the NSC apoptosis and

microcephaly are downstream of p53 activation. However, another important question is

whether the midbody defects seen in Kif20b mutant mice, indicating an abnormal abscission

process, are upstream or downstream of p53 activation. Defects or delays in midbody

maturation could activate p53. Alternatively, these midbody defects could be a consequence of

p53 activation, since p53 can regulate many genes and processes. To distinguish these

possibilities, we tested whether abnormal midbody phenotypes observed in *Kif20b* mutant NSCs

are rescued by *p53* deletion. First, we analyzed NSC midbody index in the dissociated cultures.

As *Kif20b-/-* NSCs *in vitro* are less frequently observed at the midbody stage **(Figure S3B),** and

we hypothesized this is due to NSC arrest or apoptosis, we predicted this phenotype would be

rescued by *p53* deletion. In fact, the midbody index of *Kif20b-/-; p53-/-* NSCs is not merely

rescued, but instead is significantly increased above controls **(Fig. 7A, B).** This is consistent

with the notion that some *Kif20b-/-* midbody-stage NSCs die, and further suggests that if these

cells are prevented from dying, they take longer to complete abscission than control cells. Next,

we analyzed NSC midbody structure in dissociated cultures. Similar to *Kif20b-/-* midbodies,

Kif20b-/-; p53-/- midbodies are significantly wider than controls (data not shown), and fewer of

them have constriction sites **(Fig. 7C).** Thus, unlike apoptosis, the midbody width and

constriction site phenotypes are not p53-dependent.

We next analyzed *Kif20b-/-; p53-/-* double mutant midbodies *in vivo* in E13.5 cortices.

Cytokinesis is more complex in the cortical neuroepithelium than *in vitro.* NSC nuclei must

migrate to the apical membrane to undergo mitosis and cytokinesis. During cytokinesis, the

cleavage furrow ingresses asymmetrically from the basal side of the cell, forming the midbody at

the apical membrane (Kosodo et al., 2008). To visualize NSC midbodies *en face*, we

immunostained cortical slab whole-mounts for AurKB and the apical junction marker zona

occludens-1 (ZO-1), and imaged the apical/ventricular surfaces **(Fig. 7D).** As *in vitro*, some

midbodies are short and wide, and others are long and thin, since midbodies narrow as they

mature (examples, **Fig. 7E**). In *Kif20b-/-* cortices*,* midbodies have a shifted width distribution,

with an increased median width compared to controls (**Fig. 7F** and Janisch and Dwyer, 2016).

Interestingly, we find that the *Kif20b-/-; p53-/-* double mutant cortices have a strikingly similar

midbody width distribution as *Kif20b* single mutants, with the same median **(Fig. 7F).** Thus, this

midbody maturation phenotype (width) i*n vivo* is not rescued by *p53* deletion. But surprisingly,

we find that midbody lengths, which are similar in controls and *Kif20b-/-* single mutants, are

significantly longer in *Kif20b-/-; p53-/-* double mutants than in either *Kif20b-/-* single mutants or

controls **(Fig. 7G)***.* Exceptionally long midbodies were observed after delayed abscission in cell

lines that are resistant to apoptosis (Gromley et al., 2003; Weiderhold et al., 2016). Thus, *Kif20b*

mutant NSCs may have delayed abscission, which can manifest as longer midbodies only if

apoptosis is prevented. An additional midbody phenotype we observe *in vivo* in *Kif20b* mutant

cortices is that about a third of NSC midbodies are not aligned parallel to the apical membrane,

compared to ~15% in control brains (Janisch et al., 2013; **Fig. 7H, I**). The cause of this

phenotype is unclear, but we used the *Kif20b; p53* double mutants to test whether it is p53-

dependent, perhaps due to the apoptotic process in dividing or neighboring cells. Interestingly,

p53 deletion did not prevent the misalignment phenotype, showing that it is not due to p53

activation or apoptosis **(Fig 7I)**. Taken together, these midbody analyses support the hypothesis

that defective midbody maturation and alignment are primary consequences of *Kif20b* loss

rather than secondary consequences of p53 activation. Further, the additional midbody

phenotypes detected in *Kif20b-/-; p53-/-* double mutant NSCs, namely increased midbody index

and midbody length, are consistent with delayed midbody maturation in *Kif20b* mutant NSCs;

however, these symptoms of delay are precluded in *Kif20b* mutants by p53 activation and

apoptosis.

DISCUSSION

Here we have tested the contribution of the intrinsic apoptosis pathway to the reduced

cortex size in a lethal microcephaly model, the *Kif20b* mouse mutant. We have shown that

apoptosis of cortical NSCs accounts for most of the microcephaly, but that there is a significant

apoptosis-independent contribution as well, likely reflecting the importance of Kif20b's role in

abscission for corticogenesis. Furthermore, we showed that the excess apoptosis is partially

dependent on Bax, and fully dependent on p53. Remarkably, heterozygous *p53* deletion is

sufficient to fully suppress the lethality of the *Kif20b* mutant, and rescues the brain size equally

as well as homozygous *p53* deletion. Lastly, we demonstrated that the NSC midbody maturation

defects are not rescued by *p53* deletion, which indicates that they are not caused by p53

activation, but may be upstream of p53. Thus, this work potentially identifies a novel midbody-

initiated pathway for p53 activation, and suggests that at least some types of microcephaly,

although severe, could be greatly ameliorated by inhibiting apoptosis.

The genetic and cellular experiments herein support the following working model for the

etiology of microcephaly in the *Kif20b* mutant **(Fig. 7J)**. Midbody maturation defects in some

Kif20b-/- NSCs cause p53 activation by an unknown molecular pathway (dashed line). p53 then

triggers the apoptotic cascade including Bax and other effectors**.** Apoptosis depletes the NSC

pool, thereby reducing the number of neuron and IP daughters produced, resulting in a small

brain. There is likely some stochasticity to this process, as not all *Kif20b-/-* NSCs have midbody

defects, and only a small percentage undergo apoptosis (Janisch et al., 2013; this work). It may

be that apoptosis is only triggered if midbody maturation (and hence abscission) is delayed

beyond a certain threshold. This would be analogous to a previous study in developing cortex

showing that apoptosis likelihood increases if prophase is delayed past a threshold (Pilaz et al.,

2016). Though both of these types of cell division delays trigger apoptosis through p53, the

molecular pathway upstream of p53 is likely distinct, since midbody maturation takes place well

after telophase, in the next G1 phase (Gershony et al., 2014). In fact, how defects in early steps

of mitosis signal to trigger p53 activation is only beginning to be elucidated, primarily in

immortalized cell lines that are apoptosis-resistant (McKinley and Cheeseman 2017; Lambrus et

al., 2016; Meitinger et al., 2016). The mechanism by which abscission delay could activate p53

or apoptosis has not been addressed in any system. Much more work is needed to determine

whether and how a midbody "error sensor" could directly or indirectly activate p53 in NSCs and

other cell types. It remains possible that *Kif20b* loss causes p53 activation and midbody defects

through two separate pathways. In either case, this mutant is a tool to elucidate a novel pathway

to p53 activation, one that appears very sensitive in cortical neural stem cells.

p53 activates apoptosis in NSCs following diverse cellular defects

Our work furthers the evidence that p53 responds to multiple intrinsic cellular defects to

acutely regulate NSC survival. Zika virus infection activates p53-dependent apoptosis in human

NSCs (Ghouzzi et al., 2016). A handful of other microcephalic mouse mutants, with impaired

DNA replication, mitosis, or cleavage furrowing, have implicated p53 in apoptosis and reduced

brain size (Bianchi et al., 2017; Houlihan et al., 2014; Insolera et al., 2014; Marjanovic et al.,

2015; Marthiens et al., 2013; Murga et al., 2009). Interestingly, inhibition of p53 in these mutants

was sometimes able to increase cortical thickness, but in other cases worsened brain

phenotypes (Marthiens et al., 2013; Murga et al., 2009). By contrast, *Kif20b-/-* embryonic brain

structure and postnatal survival were well-rescued by even heterozygous deletion of *p53*.

Heterozygous *p53* rescue has not been reported in other microcephaly mouse mutants. In the

Brca1 mutant, dwarfism but not microcephaly was rescued by heterozygous *p53* deletion (Xu et

al., 2001). Thus, the *Kif20b-/-* microcephaly model is more severe in reduction of brain size

compared to some other models, but it is also more easily rescuable by apoptosis inhibition. In

fact, *p53* heterozygous or homozygous deletion can completely rescue the perinatal lethality of

Kif20b-/- mice. This is important for eventual treatment of human microcephaly caused by

genetic mutations or viruses, to determine which subtypes and phenotypes might be treatable

by blocking apoptosis.

A p53-independent consequence of Kif20b loss accounts for the remaining cortical size

deficit at birth

The 20% deficit in cortical thickness in *Kif20b*; *p53* double mutant mice at birth indicates

that even without the excess apoptosis, *Kif20b* mutant NSCs and neurons cannot create a brain

of normal size and structure, probably due to abnormal abscission causing some other problem

in *Kif20b-/-; p53-/-* NSCs. The midbody maturation defects that are more frequent in *Kif20b*

mutant brains may cause delays in daughter cell severing from the mother cell and delamination

from the apical membrane in the case of neuronal daughters. It could also affect whether

abscission occurs on both sides of the midbody or only one. The significance of the midbody

misalignment increase is not clear, but here we showed that it is not caused by p53 activation or

apoptosis. It is possible that Kif20b may help anchor or stabilize midbodies at the apical

membrane via cell adhesion junctions until abscission is complete.

Aside from regulating abscission, Kif20b has postmitotic roles in cortical neurons that

may contribute to cortical growth. We previously showed that *Kif20b* mutant embryonic cortical

neurons in culture display a polarization defect and abnormalities in neurite growth and

branching. Thus the deficit in thickness of the cortical plate at birth may be due to reduced

neuropil. A key function of Kif20b is to enable tight microtubule packing: Kif20b has microtubule

crosslinking activity *in vitro*, and in *Kif20b* mutant neurons, wider neurites and gaps in

microtubule bundles were noted (Abaza et al., 2003; McNeely et al., 2017). Therefore, Kif20b

may help to crosslink microtubules in both NSC midbodies and neuronal axons.

Elucidating the heterogeneous mechanisms of brain malformations requires many

genetic models and culture systems

Defects in cytokinetic abscission mechanisms could underlie a range of poorly

understood microcephalies and related brain malformations. For example, human genetic

studies in two different families showed that a very severe prenatal lethal

microcephaly/anencephaly is caused by mutations in the gene encoding the key abscission

protein Cep55 (OMIM Ref #236500) (Frosk et al., 2017; Bondeson et al., 2017) Cep55

depletion in cell lines causes more severe midbody structural defects and abscission delay than

Kif20b depletion does (Zhao et al., 2006). Patients with *Kif20b* loss-of-function mutations have

not yet been identified, so it remains to be seen whether the severity of single-cell abscission

phenotypes correlates with the severity of brain malformation. Cell lines such as HeLa cells can

model some abscission defects, for example midbody maturation, but not others, such as

midbody positioning at the apical membrane, p53 activation, or apoptosis. Our studies of the

Kif20b mouse model provide human geneticists with a candidate gene and cellular markers for

syndromes involving peri- or prenatal lethality, microcephaly, craniofacial defects, or

microphthalmia. Comparing mouse models for abscission genes with other microcephalic mice

as well as human phenotypes will help us understand the heterogeneous etiologies of and

potential diagnosis and treatments for these devastating conditions.

MATERIALS AND METHODS

Mice: Mouse colonies were maintained in accordance with NIH guidelines and policies

approved by the IACUC. Embryos were harvested by caesarean section, and the morning of the

vaginal plug was considered embryonic day (E) 0.5. Littermate embryos served as controls for

all experiments. The *Kif20b^{magoo}* allele, as previously described (Janisch et al., 2013) is

maintained on both C57BL/6 and FVB/N backgrounds, and 50/50% mixed background embryos

are used for experiments. *Trp53^{tm1Tyj}* mice on C57BL/6 background were obtained from The

Jackson Laboratory (JAX stock #002101) (Jacks et al., 1994). Mixed BL6/FVB/N background

Bax^{tm1Sjk} mice were a gift from Christopher Deppmann (JAX stock #002994) (Knudson et al.,

1995). Sex of embryonic mice were not noted as sex was not a relevant biological variable for

these experiments. The specific ages of embryonic mice used is noted in figure legends for

each experiment.

Cortical cell cultures: Cells were dissociated from E12.5 cortices following a protocol adapted

from Sally Temple's lab (Qian et al., 1998). The Worthington Papain Dissociation Kit was used

to dissociate cells (Worthington Biochemical Corporation, Cat # LK003150). Cells were cultured

in DMEM with Na-Pyruvate, L-Glutamine, B-27, N2, N-acetyl-cysteine and basic Fibroblast

Growth Factor (bFGF). After 24 hours, cells were fixed by adding an equal volume of room-

temperature 8% PFA for 5 minutes to cell media, followed by removal of media and addition of -

20º cold methanol for 5 minutes.

Apical slab preparation: Apical slabs were prepared as previously described (Janisch and

Dwyer, 2016). The meninges and skull were removed to expose the brain in E13.5 embryos,

followed by fixation with 2% PFA for 20 minutes. Next, apical slabs were made by pinching off

cortices, flipping so that the apical surface was upright, and trimming to flatten the slab. Slabs

were fixed for another 2 minutes with 2% PFA followed by blocking with 5% normal goat serum

(NGS) for 1 hr. Primary antibodies were applied for 1 hr. at room temperature and then moved

to 4° overnight. The next day, after 3, 10-minute PBS washes secondary antibodies and DAPI

were applied at a concentration of 1:200 for 30 minutes. After two more 10 minute PBS washes

slabs were coverslipped with VectaShield fluorescent mounting medium (Vector Laboratories

Inc., H-1000) and imaged. *z*-stack depth was 8-20 µm and step size was 0.5 µm. Midbodies

were considered misaligned if the two halves of the midbody were not in the same *z*-plane or

within two adjacent *z*-planes.

Immunoblotting: Brain lysates were prepared with RIPA lysis buffer (150 mM NaCl, 1% NP40,

0.5% sodium deoxycholate, 0.1% SDS, 50 mM Tris-HCl pH 8) with protease and phosphatase

inhibitors. Protein concentration in lysates was determined by bicinchoninic acid (BCA) assay,

and 60 µg total protein was loaded per lane on 4-20 gradient% polyacrylamide gels. Proteins

were transferred by electroblotting onto a 0.2 µm PVDF membrane overnight at 30 mA.

Membranes were blocked in 150 mM NaCl, 100 mM Tris-HCl pH 7.5 and 0.5% Tween 20

(TBST) with 5% dried milk (blocking buffer) for 1 hour. Primary antibodies were incubated with

the membrane overnight at 4°C. After three washes, LI-COR IRDye 800 CW goat anti-rabbit IgG

and 680 RD goat anti-mouse secondary antibodies were applied (1:10000) in blocking buffer for

1 hour at room temperature. After washing with TBST for 5 minutes, 3 times, immune

complexes were visualized using a LI-COR western blot imager.

Immunostaining: To collect cryosections for IHC, age E14.5 and P0 brains were removed from

heads and fixed for 6 and 24 hours, respectively, in 4% PFA, followed by immersion in 30%

sucrose in PBS overnight. Next, whole brains were embedded in OTC (Tissue-Tek, 4583) and

cryosections were cut at 20 µm thickness and collected on Superfrost Plus slides (Fisher

Scientific, 12-550-15). Frozen sections were stored at -80 degrees. Prior to immunostaining,

cryosections were warmed to room temperature, then if antigen retrieval was needed, immersed

in 10 mM citrate buffer at 95 degrees for 20 minutes. After cooling, sections were blocked in 2%

NGS for 1 hour, followed by incubation with primary antibodies overnight at 4º C. The next day,

after PBS washes sections were incubated with AlexaFluor secondary antibodies at 1:200 and

DAPI at 1:100 for 30 minutes followed by PBS washes and coverslipping with VectaShield

fluorescent mounting medium. For IF on coverslips of dissociated cortical progenitors, a similar

protocol was used but with primary antibodies applied for 3 hours at room temperature, and

antigen retrieval was not used. Coverslips were mounted on Superfrost Plus slides with Fluoro-

Gel (Electron Microscopy Sciences, 17985-10).

Antibodies: Antibodies used in this analysis were rabbit polyclonal anti-human cleaved-
caspase 3 (Cell-Signaling 9661S, 1:250), mouse monoclonal anti-rat beta-III-tubulin (Tuj1)
(BioLegend 801201, 1:500), rat monoclonal anti-mouse Tbr2 (ebioscience 14-4875, 1:200),
rabbit polyclonal anti-mouse Pax6 (BioLegend PRB-278P, 1:200), Aurora B kinase mouse
monoclonal anti-rat (BD Biosciences 611082, 1:300), rabbit polyclonal anti-human alpha-tubulin
(Thermo Scientific RB-9281-P0, 1:300), rat monoclonal alpha-tubulin (Novus Biologicals NB600-
506, 1:300), rabbit monoclonal anti-human PH3 (Cell Signaling 3458, 1:200), chicken polyclonal
anti-mouse Nestin (Aves Labs NES, 1:600), rat monoclonal anti-human Ki67 (eBioscience 14-
5698, 1:100), mouse monoclonal anti-human p53 (Millipore 05-224, 1:250), rabbit polyclonal
anti-mouse p53 (Leica Biosystems NCL-L-p53-CM5p, 1:500), rabbit polyclonal anti-human
Beta-Catenin (Sigma-Aldrich SAB4500545 1:1000), and polyclonal rabbit anti Zo-1 (rabbit,
Invitrogen 61-7300,1:50). All antibodies were validated for the application used in multiple
previous publications.

Images were collected on either a Zeiss Axio ImagerZ1 microscope with AxioCam MRm
(Figures 1, 2, 3, 5 6 and 7A-C), a Zeiss AxioObserver fluorescent widefield inverted scope
microscope (Figures 1, 2 and 3), an inverted DeltaVision with TrueLight deconvolution
microscope and softWoRx Suite 5.5 image acquisition software (Applied Precision) (Figure 7D-
J), or a Leica MZ16F microscope with DFC300FX camera (Figure 2, H&E and Figure 4).
Control and mutant fluorescence images were collected with the same exposure times
and on the same day. All image analysis was performed in ImageJ/Fuji and any changes to
brightness and contrast were applied uniformly across images. Statistical analyses were
performed using Excel (Microsoft) or GraphPad PRISM software. The sample sizes were pre-
determined based on our labs previous experience with cortical development analyses and
others' published results. After obtaining pilot data, power analyses were performed if necessary
to determine if the number of samples obtained was high enough for the effect size seen. NSC
cultures that were unhealthy were not imaged and analyzed, but no other data was excluded
from the analysis. No randomization or blinding was used as no experimental manipulation was
applied other than genetic knockouts. Genotyping was performed after collection of embryos to
determine genetic status. Statistical tests used are specified in each figure legend. For each
sample set a statistical test of normality was performed using GraphPad PRISM software.
Parametric tests were used when sample sets had a normal distribution and non-parametric
tests were used when sample sets did not have a normal distribution. Variance was calculated
and was typically similar between groups. N's listed in figure legends indicate the number of

coverslips, brains and litters collected for each experiment. For each brain, at least three

sections were imaged, and for each coverslip, at least 5, 20X pictures or 10, 40X pictures were

analyzed.

ACKNOWLEDGEMENTS

We thank Chris Deppmann for use of the *Bax* mutant mouse line. We are grateful to Bettina

Winckler, Jing Yu, Xiaowei Lu, Ann Sutherland, Todd Stukenberg and members of their labs, as

well as Katrina McNeely for advice and discussion. We are grateful to Madison Hecht, Gabrielle

Wolfe, Mackenzie Shannon, Haley Hopkinson and Adriana Ehlers for help with cryosectioning

and data analysis. This work was supported by the National Institutes of Health (R01 NS076640

to NDD), the UVA Medical Scientist Training Program (MSTP T32 GM 7267-37) and the UVA

Cell and Molecular Biology Training Grant (2T32GM008136-31A1).

COMPETING INTERESTS: No competing interests declared.

REFERENCES

Abaza, A., Soleilhac, J.M., Westendorf, J., Piel, M., Crevel, I., Roux, A. and Pirollet, F.
(2003). M phase phosphoprotein 1 is a human plus-end-directed kinesin-related protein required
for cytokinesis. *J Biol. Chem.* **278**(30): 27844-27852.

Arya, R. and White, K. (2015). Cell death in development: Signaling pathways and core

mechanisms. *Semin Cell Dev Biol.* **39**:12-9.

503
Bianchi, F.T., Tocco, C., Pallavicini, G., Liu, Y., Verni, F., Merigliano, C., Bonaccorsi, S.,

El-Assawy, N., Priano, L., Gai, M., Berto, G.E., Chiotto, A.M., Sgro, F., Caramello, A.,

Tasca, L., Ala, U., Neri, F., Oliviero, S., Mauro, A., Geley, S., Gatti, M. and Di Cunto, F.

(2017). Citron Kinase Deficiency Leads to Chromosomal Instability and TP53-Sensitive

Microcephaly. *Cell Reports.* **18**(7): 1674-1686.

Bondeson, M.L, Ericson, K., Gudmundsson, S., Ameur, A., Ponten, F., Wesstrom, J.,

Frykholm, C. and Wilbe, M. (2017). A nonsense mutation in CEP55 defines a new locus for a

Meckel-like syndrome, an autosomal recessive lethal fetal ciliopathy. *Clin Genet.* **92**(5): 510-516.

Breuss, M. Fritz, T. Gstrein, T., Chan, K., Ushakova, L., Yu, N., Vonberg, F.W., Werner, B., Elling, U., and Keays, D.A. (2016). Mutations in the murine homologue of TUBB5 cause microcephaly by perturbing cell cycle progression and inducing p53-associated apoptosis. *Development.* **143**(7): 1126-33.

Chen, J.F., Zhang, Y., Wilde, J., Hansen, K.C., Lai, F. and Niswander, L. (2014). Microcephaly disease gene Wdr62 regulates mitotic progression of embryonic neural stem cells and brain size. *Nat Commun.* **5**:3885.

Duerinckx, S. and Abramowicz, M. (2018). The genetics of congenitally small brains. *Semin Cell Dev Biol.* **76**:76-85.

Dwyer, N.D., Manning, D.K., Moran, J.L., Mudbhary, R., Fleming, M.S., Favero, C.B., Vock, V.M., O'Leary, D.D., Walsh, C.A. and Beier, D.R. (2011). A forward genetic screen with a thalamocortical axon reporter mouse yields novel neurodevelopment mutants and a distinct Emx2 mutant phenotype. *Neural Dev.* **6**(1):3.

Dwyer, N.D., Chen, B. Chou, S.J., Hippenmeyer, S. Nguyen, L. and Ghashghaei, H.T. (2016). Neural Stem Cells to Cerebral Cortex: Emerging Mechanisms of Regulating Progenitor Behavior and Productivity. *J Neurosci.* **36**(45): 11394-11401.

Frosk, P., Arts, H.H., Philippe, J., Gunn, C.S., Brown, E.L., Chodirker, B., Simard, L., Majewski, J., Fahiminiya, S., Russell, C., Liu, Y.P., Hegele, R., Katsanis, N., Goerz, C., Del Bigio, M.R. and Davis, E.E. (2017). A truncating mutation in CEP55 is the likely cause of MARCH, a novel syndrome affecting neuronal mitosis. *J Med Genetics.* **54**(7): 490-501.

Gershony, O., Pe'er, T., Noach-Hirsh, M., Elia, N. and Tzur, A. (2014). Cytokinetic abscission is an acute G1 event. *Cell Cycle.* **13**(21): 3436-3441.

Ghouzzi, V.E., Bianchi, F.T., Molineris, I., Mounce, B.C., Berto, G.E., Rak, M., Lebon, S., Aubry, L., Tocco, C., Gai, M., Chiotto, A.M., Sgro, F., Pallavicini, G., Simon-Loriere, E.,

545 **Passemard, S., Vignuzzi, M., Gressens, P. and Di Cunto, F**. (2016). ZIKA virus elicits P53
546 activation and genotoxic stress in human neural progenitors similar to mutations involved in
547 severe forms of genetic microcephaly. *Cell Death Dis.* **7**(10):e2440.
548
549 **Gromley, A., Jurczyk, A., Sillibourne, J., Halilovic, E., Mogensen, M., Groisman, I.,**
550 **Blomberg, M. and Doxsey, S.** (2003). A novel human protein of the maternal centriole is
551 required for the final stages of cytokinesis and entry into S phase. *Journal of Cell Biology*.
552 **161**(3); 535-45.
553
554 **Guizetti, J., Schermelleh, L. Mantler, J., Maar, S., Poser, I., Leonhardt, H., Muller-Reichert,**
555 **T. and Gerlich, D.W.** (2011). Cortical constriction during abscission involves helices of ESCRT-
556 III-dependent filaments. *Science.* **331**(6024): 1616-20.
557
558 **Herszterg, S., Pinheiro, D. and Bellaiche, Y.** (2014). A multicellular view of cytokinesis in
559 epithelial tissue. *Trends Cell Biol*. **25**(5): 285-93.
560
561 **Houlihan, S. and Feng, Y.** (2014). The scaffold protein Nde1 safeguards the brain genome
562 during S phase of early neural progenitor differentiation. *Elife*. **3**:e03297.
563
564 **Insolera, R., Bazzi, H., Shao, W., Anderson, K.V., and Shi, S.H.** (2014). Cortical
565 neurogenesis in the absence of centrioles. *Nature Neuroscience*. **17**(11): 1528-35.
566
567 **Janisch, K.M., Vock, V.M., Fleming, M.S., Shrestha, A., Grimsley-Myers, C.M., Rasoul,**
568 **B.A., Neale, S.A., Cupp, T.D., Kinchen, J.M., Liem, K.F. Jr. and Dwyer, N.D.** (2013). The
569 vertebrate-specific Kinesin-6, Kif20b, is required for normal cytokinesis of polarized cortical stem
570 cells and cerebral cortex size. *Development*. **140**(23):4672- 4682.
571
572 **Jacks, T., Remington, L., Williams, B.O., Schmitt, E.M., Halachmi, S., Bronson, R.T. and**
573 **Weinberg, R.A.** (1994). Tumor spectrum analysis in p53-mutant mice. *Current Biology*. **4**(1): 1-
574 7.
575
576 **Janisch, K.M. and Dwyer, N.D.** (2016). Imaging and quantitative analysis of cytokinesis in
577 developing brains of Kinesin-6 mutant mice. *Methods in Cell Biology*. **131**:233-252.
578

Janisch, K.M., McNeely, K.C., Dardick, J.M., Lim, S.H. and Dwyer, N.D. (2018). Kinesin-6
KIF20B is required for efficient cytokinetic furrowing and timely abscission in human cells. *Mol*
Biol Cell. **29**(2): 166-179.

Kanehira, M., Katagiri, T., Shimo, A., Takata, R., Shuin, T., Miki, T., Fujioka, T. and
Nakamura, Y. (2007). Oncogenic role of MPHOSPH1, a cancer-testis antigen specific to human
bladder cancer. *Cancer Res.* **67**(7): 3276-85.

Kastenhuber, E.R. and Lowe, S.W. (2017). Putting p53 in Context. *Cell.* **170**(6): 1062-1078.

Knudson, C.M., Tung, K.S., Tourtellotte, W.G., Brown, G.A. and Korsmeyer, S.J. (1995).
Bax-deficient mice with lymphoid hyperplasia and male germ cell death. *Science* **270**(5233): 96-
9.

Kosodo, Y., Toida, K., Dubreuil, V., Alexandre, P., Schenk, J., Kiyokage, E., Attardo, A.,
Mora-Bermudez, F., Arii, T., Clarke, J.D. and Huttner, W.B. (2008). Cytokinesis of
neuroepithelial cells can divide their basal process before anaphase. *Embo J*. **27**(23):3151-
3163.

Lambrus, B.G., Daggubati, V., Uetake, Y., Scott, P.M., Clutario, K.M., Sluder, G. and
Holland, A.J. (2016). A USP28-53BP1-p53-p21 signaling axis arrests growth after centrosome
loss or prolonged mitosis. *J Cell Biol.* **214**(2): 143-53.

Lin, W.F., Lin, X.L., Fu, S.W., Yang, L., Tang, C.T., Gao, Y.J., Chen, H.Y. and Ge, Z.Z.
(2018). Pseudopod-associated protein KIF20B promotes Gli1-induced epithelial-mesenchymal
transition modulated by pseudopodial actin dynamic in human colorectal cancer. *Mol Carcinog.*
Doi: 10.1002/mc.22812

Lindsten, T., Ross, A.J., King, A., Zong, W.X., Rathmell, J.C., Shiels, H.A., Ulrich, E.,
Waymire, K.G., Mahar, P., Frauwirth, K., Chen, Y., Wei, M., Eng, V.M., Adelman, D.M.,
Simon, M.C., Ma, A., Golden, J.A., Evan, G., Korsmeyer, S.J., MacGregor, G.R. and
Thompson, C.B. (2000). The combined functions of proapoptotic Bcl-2 family members bak
and bax are essential for normal development of multiple tissues. *Mol Cell.* 6: 1389-99.

Liu, X., Zhou, Y., Liu, X., Peng, A., Gong, H., Huang, L., Ji, K., Petersen, R.B., Zheng, L.

and Huang, K. (2014). MPHOSPH1: a potential therapeutic target for hepatocellular carcinoma.

Cancer Res. **74**(22): 6623-34.

Marjanovic, M., Sanchez-Huertas, C., Terre, B., Gomez, R., Scheel, J.F., Pacheco, S.,

Knobel, P.A., Martinez-Marchal, A., Aivio, S., Palenzuela, L., Wolfrum, U., McKinnon, P.J.,

Suja, J.A., Roig, I., Costanzo, V., Luders, J. and Stracker, T.H. (2015). CEP63 deficiency

promotes p53-dependent microcephaly and reveals a role for the centrosome in meiotic

recombination. *Nature Communications*. **6**:7676.

Marthiens, V., Rujano, M.A., Pennetier, C., Tessier, S., Paul-Gilloteaux, P. and Basto, R.

(2013). Centrosome amplification causes microcephaly. *Nat Cell Biol*. **15**(7):731-740.

Meitinger, F., Anzola, J.V., Kaulich, M., Richardson, A., Stender, J.D., Benner, C., Glass,

C.K., Dowdy, S.F., Desai, A., Shiau, A.K. and Oegema, K. (2016). *J Cell Biol.* **214**(2): 155-66.

Murga, M., Bunting, S., Montana, M.F., Soria, R., Mulero, F., Canamero, M., Lee, Y.,

McKinnon, P.J., Nussenzweig, A. and Fernandez-Capetillo, O. (2009). A mouse model of

ATR-Seckel shows embryonic replicative stress and accelerated aging. *Nature Genetics*. **41**(8):

891-8.

McKinley, K.L. and Cheeseman, I.M. (2017). Large-Scale Analysis of CRISPR/Cas9 Cell-

Cycle Knockouts Reveals the Diversity of p53-Dependent Responses to Cell-Cycle Defects.

Developmental Cell. **40**(4): 405-420.

McNeely, K.C., Cupp, T.D., Little, J.N., Janisch, K.M., Shrestha, A. and Dwyer, N.D. (2017).

Mutation of Kinesin-6 Kif20b causes defects in cortical neuron polarization and morphogenesis**.**

Neural Development. **12**:5.

Mierzwa, B. and Gerlich, D.W. (2014). **Cytokinetic Abscission: Molecular Mechanisms and**

Temporal Control. *Developmental Cell*. **31**:5 (525-538).

Pilaz, L.J., McMahon, J.J., Miller, E.E., Lennox, A.L., Suzuki, A., Salmon, E. and Silver,

D.L. (2016). Prolonged Mitosis of Neural Progenitors Alters Cell Fate in the Developing Brain.

Neuron. **89**(1): 83-99.

Qian, X., Goderie, S.K., Shen, Q., Stern, J.H. and Temple, S. (1998). Intrinsic programs of

patterned cell lineages in isolated vertebrate CNS ventricular zone cells. *Development.* **125**

(3143-3152).

Stottmann, R.W., Donlin, M., Hafner, A., Bernard, A. Sinclair, D.A. and Beier, D.R. (2013). A

mutation in Tubb2b, a human polymicrogyria gene, leads to lethality and abnormal cortical

development in the mouse. *Hum Mol Genet.* **22**(20): 4053-63.

Weiderhold, K.N., Fadri-Moskwik, M., Pan, J., Nishino, M., Chuang, C., Deeraksa, A., Lin,

S.H. and Yu-Lee, L.Y. (2016). Dynamic Phosphorylation of NudC by Aurora B in Cytokinesis.

PLoS One. **11**(4):20153455.

Westphal, D., Dewson, G., Czabotar, P.E. and Kluck, R.M. (2011). Molecular biology of Bax

and Bak activation and action. *Biochemica et Biophysica Acta (BBA) – Molecular Cell Research*.

1813(4): 521-531.

Xu, X., Qiao, W., Linke, S.P., Cao, L., Li, W., Furth, P.A., Harris, C.C. and Deng, C. (2001).

Genetic interactions between tumor suppressors Brca1 and p53 in apoptosis, cell cycle and

tumorigenesis. *Nature Genetics*. **28**: 266-271.

Figure 1. *Bax* deletion partially rescues microcephaly and apoptosis in *Kif20b-/-* mice

A-D. Representative E14.5 cortical sections, and plotted mean of cortical thicknesses show the severe reduction in *Kif20b* mutants is partly restored in *Kif20b; Bax* double mutants.

E-H. Cleaved caspase-3 (CC3) staining in E14.5 cortices shows the increased apoptosis in *Kif20b* mutants is partially reduced in *Kif20b; Bax* double mutants.

For (D, H), n= 5 *Kif20b+/-; Bax+/-*, 4 *Kif20b-/-; Bax+/+*, 5 *Kif20b-/-; Bax-/-* mice, from a total of 8 litters.

* p < 0.05, ** p < 0.01, *** p < 0.001, **** p < .0001, one-way ANOVA. Error bars are +/- s.e.m. Scale bars: A: 500μm; E: 20μm.

Figure 2. *p53* deletion rescues microcephaly and apoptosis in *Kif20b-/-* mice

A-E. E14.5 cortical sections and plotted mean cortical thicknesses show the severe reduction in *Kif20b* mutants is fully rescued by heterozygous or homozygous *p53* deletion. n = 6 *Kif20b+/-; p53+/-*, 3 *Kif20b-/-; p53+/+*, 4 *Kif20b-/-; p53+/-*, 4 *Kif20b-/-; p53-/-* mice, from a total of 7 litters.

F-J. CC3 staining shows apoptosis in *Kif20b* mutants returns to control levels by heterozygous or homozygous *p53* deletion. n = 3 *Kif20b+/-; p53+/-, Kif20b-/-; p53+/+, Kif20b-/-; p53+/-, Kif20b-/-; p53-/-* mice, from a total of 6 litters.
 * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, one-way ANOVA. Error bars are +/- s.e.m. Scale bars: A: 1 mm; F: 20µm.

Figure 3. *p53* deletion restores normal production of intermediate progenitors and neurons in *Kif20b-/-* cortex

A-C. E14.5 cortical sections labeled with neuronal-specific tubulin (Tuj1, red) show the decreased neuronal layer thickness in *Kif20b* mutants is rescued in *Kif20b;p53* double mutants. cp: cortical plate; iz: intermediate zone.

D-F. E14.5 cortices labeled with Pax6 (NSCs, pink) and Tbr2 (IPs, green) show the altered layer proportions in *Kif20b* mutant are rescued by *p53* deletion. svz: subventricular zone; vz: ventricular zone.

G. The mean neuronal layer proportional thickness is halved in *Kif20b* mutants but rescued by *p53* co-deletion.

H. The proper NSC layer (Pax6+) proportionality is restored by *p53* co-deletion.

I. The density of Tbr2+ IPs is reduced in *Kif20b* mutants, but rescued in *Kif20b;p53* double mutants. n = 3 each *Kif20b+/-;p53+/-, Kif20b-/-;p53+/+, Kif20b-/-;p53-/-* mice for Tuj1 and Pax6 layer thickness. n = 6 *Kif20b+/-;p53+/-, 3 Kif20b-/-;p53+/+, 4 Kif20b-/-;p53-/-* mice for Tbr2+ cell counts, from a total of 6 litters.

* $p < 0.05$, ** $p < 0.01$, one-way ANOVA. Error bars are +/- s.e.m. Scale bars: 20 µm.

A Mice alive on P0 (101 total observed)

Genotype	Observed	Expected
Kif20b +/+ or +/-; p53 +/+ or +/-	58	~57 (9/16)
Kif20b +/+ or +/-; p53 -/-	22	~19 (3/16)
Kif20b -/-; p53 +/+	0	~6 (1/16)
Kif20b -/-; p53 +/-	15	~13 (1/8)
Kif20b -/-; p53 -/-	6	~6 (1/16)

Figure 4. *p53* deletion rescues postnatal survival but incompletely rescues cortical size at birth in *Kif20b-/-* mice

A. *Kif20b-/-; p53+/+* mice do not survive at birth at expected Mendelian ratios, but *Kif20b-/-; p53+/-* and *Kif20b-/-; p53-/-* mice survive at comparable levels to controls.

B,C. The average length of *Kif20b-/-;p53-/-* mice is slightly decreased compared to controls at P0. N = 54 *Kif20b+/+ or +/- p53+/+ or +/-* mice, 22 *Kif20b+/+ or +/- p53-/-,* and 5 *Kif20b-/-; p53-/-* mice.

D, E. Coronal sections from P0 rostral forebrains stained with hematoxylin and eosin (H&E) show reduced cortical thickness and enlarged ventricles of double mutants.

F. Cortical thickness in *Kif20b-/-;p53-/-* mice is reduced approximately 20% compared to controls in rostral, middle and caudal sections at P0.

G. Reduction is in the cortical plate (cp) and intermediate zone (iz) but not ventricular zone/subventricular zone (vz/svz) layers. For F and G N = 8 *Kif20b+/-; p53+/-* and 5 *Kif20b-/-; p53-/-* brains.

* p < 0.05, ** p < 0.01, one-way ANOVA (C) and student's t-test (F,G). Error bars are +/- s.e.m. Scale bars: B: 1 mm; D: 200 µm; D': 100 µm.

Figure 5. Kif20b loss cell-autonomously causes abscission defects in cortical NSCs.
A-C. Dissociated E12.5 NSC midbodies at 24 hrs in culture labeled with alpha-tubulin (red) and Aurora kinase B (AurkB, green). Early-stage midbodies are wider with no constriction sites (A); late-stage midbodies are thinner and have one (B) or two (C) constriction sites (arrows).
D. *Kif20b-/-* midbodies have a significantly shifted width distribution, with more wide midbodies. (Medians: Control, 0.89μm; mutant, 0.96μm).
E. The mean percentage of midbodies with at least one constriction site (detected by tubulin) is significantly reduced in *Kif20b-/-* cultures.
n (D,E): 141 control, 125 mutant midbodies; 6 coverslips and 3 embryos each, 3 litters. D, Mann-Whitney; E, Student's t-test. Error bars are +/- s.e.m. All cultures from E12.5 cortices and fixed after 24 hours. Scale bar in A: 5 μm

Figure 6. *Kif20b* **loss causes apoptosis in proliferating cortical NSCs**

A,B. Representative fields of control (left) and *Kif20b-/-* (right) dissociated cortical cultures labeled with CC3 (red, arrows) for apoptotic cells.

C,D. Representative fields labeled with Nestin (NSCs, pink) and CC3 (green). Arrows in D indicate cells co-positive for Nestin/CC3.

E. The mean percentage of apoptotic cells in *Kif20b-/-* cortical cultures is increased (>2 fold) above controls. n = 13 control, 11 mutant coverslips; 8 and 7 brains, 7 litters.

F. The percent of apoptotic neurons (CC3+/Tuj1) is not increased in *Kif20b-/-* cultures, but the percents of apoptotic NSCs and proliferating NSCs (CC3+/Nestin+; CC3+/Ki67+) are significantly increased. Tuj1/CC3 analysis: n = 7 control, 6 mutant coverslips; 4 and 3 brains, 4 litters. Nestin/CC3 analysis: n = 7 control, mutant coverslips; 4 brains each, 3 litters. Ki67/CC3 analysis: n = 6 control, 5 mutant coverslips; 3 brains each, 3 litters.

G. Apoptosis is not increased in *Kif20b;p53* double mutant cultures. n = 4 *Kif20b+/-;p53+/-*, 3 *Kif20b-/-;p53-/-* coverslips; 3 and 2 embryos, 2 litters. * p < 0.05, ** p < 0.01, *** p < 0.001, Student's t-test. Error bars are +/- s.e.m. All cultures from E12.5 cortices and fixed after 24 hours. Scale bars in A: 20 µm.

Figure 7. *p53* deletion does not rescue abscission phenotypes of *Kif20b* mutant NSCs

A. Representative image of E12.5 dissociated cortical cells, labeled with AurkB for midbodies (green, arrow), and Ki67 for proliferating NSCs (pink).

B. The mean percent midbody stage proliferating NSCs is significantly increased in *Kif20b-/-;p53-/-* cultures. n = 4 *Kif20b+/-;p53+/-*, 3 *Kif20b-/-;p53-/-* coverslips; 3 embryos each, 2 litters.

C. Of the NSCs at midbody stage, the percent with a constriction site is significantly reduced in *Kif20b;p53* cultures. n = 111 *Kif20b+/-;p53+/-*, 101 *Kif20b-/-;p53-/-* midbodies; 3 embryos each, 2 litters.

D. Representative image of E13.5 cortical slab, apical surface, labeled with ZO-1 (red, apical junctions) and AurkB (green, midbodies).

E. NSC midbodies range from short and wide (top) to long and thin (bottom).

F. The median midbody width is similarly increased in *Kif20b* single and *Kif20b;p53* double mutant cortices compared to controls. Median widths (µm) = 0.92 in *Kif20b+/-;p53+/-*, 1.0 in *Kif20b-/-;p53+/+*, 1.02 in *Kif20b-/-;p53-/-* cortices.

G. The median midbody length is similar in *Kif20b-/-* and control cortices, but is significantly increased in *Kif20b-/-;p53-/-* cortices. Median lengths (µm) = 4.82 in *Kif20b-/-;p53+/+*, 5.11 in *Kif20b+/-;p53+/-*, 5.43 in *Kif20b-/-;p53-/-* cortices. n for (F,G) = 346 *Kif20b+/-;p53+/-*, 189 *Kif20b-/-;p53+/+*, and 299 *Kif20b-/-;p53-/-* midbodies; 8, 5, and 7 cortical hemisphere slabs, 6, 4, and 5 embryos, 8 litters.

H. Schematic of aligned and misaligned midbodies relative to the apical membrane.

I. The percent misaligned midbodies is doubled in both *Kif20b-/-* and *Kif20b-/-;p53-/-* cortices compared to controls. n = 8 *Kif20b+/-;p53+/-*, 7 *Kif20b-/-;p53+/-*, 7 *Kif20b-/-;p53-/-* cortical slabs from 6, 4, and 6 embryos, respectively, 8 litters.

J. Working model. p53 is required for NSC apoptosis and microcephaly caused by *Kif20b loss* but is not required for the midbody defects. How midbody defects lead to p53 activation remains unknown (dotted arrow).

A-C are E12.5 cortical cultures fixed at 24 hrs. *p < 0.05, ** p < 0.01, Student's t-test for (B,C); K.S. test for (F,G). Error bars are +/- s.e.m. Scale bars, 10 µm.

A Kif20b+/- p53+/- **B** Kif20b-/- p53+/+ **C** Kif20b-/- p53+/- **D** Kif20b-/- p53-/-

E

Genotype	Percent with craniofacial defects
p53-/- ; Kif20b+/+ or +/-	12% (2/17)
p53+/+; Kif20b-/-	100% (3/3)
p53+/- ; Kif20b-/-	8% (1/12)
p53-/- ; Kif20b-/-	11% (1/9)

Supplemental Figure S1: *p53* deletion rescues craniofacial defects in *Kif20b* mutant mice.
A-D. Representative E14.5 embryos of the indicated genotypes.
E. E14.5 embryos observed with craniofacial defects including shortened snout,
underdeveloped eyes, and/or brain malformations of the indicated genotypes.

Supplemental Figure S2. p53 protein level is elevated in *Kif20b-/-* E12.5 brain lysates and dissociated NSCs

A. Immunoblot from E12.5 control and *Kif20b-/-* cortical lysates shows bands for p53 (53 kDa) and beta-catenin (95 kDa) as a loading control.

B. p53 band intensity, normalized to beta-catenin bands, is increased by 50% in *Kif20b-/-* samples. n = 4 blots from 4 embryos each. * p < 0.05, paired ratio t-test.

C. Representative images of E12.5 dissociated cortical cultures fixed after 24 hrs *in vitro*, and immunostained for p53 (green) and nestin (red) to mark NSCs. Scale bar: 10 μm.

D. The average percent of NSCs with detectable nudear p53 staining is increased more than two-fold in *Kif20b-/-* cultures. n= 3 cultures for each genotype, each from an independent E12.5 brain.

Supplemental Figure S3. Analyses of mitotic and midbody indices in *Kif20b* mutant NSC cultures
A. Examples of E12.5 dissociated cortical NSCs at stages of mitosis and cytokinesis, labeled with PH3 (red) and AurkB (green).
B. The mean percentages (+/- s.e.m.) of cycling NSCs (Ki67+) in mitosis or midbody stage are slightly reduced in *Kif20b-/-* cultures.
C. Categorizing the dividing NSCs from B, there is a small increase of *Kif20b-/-* NSCs in anaphase/early telophase. For B and C, n = 3 coverslips from 3 embryos each, with 1499 control and 1421 *Kif20b-/-* Ki67+ cells. * p < 0.05, Student's t-test for (B), Chi Square and Fisher's exact test for (C). All cultures dissociated from E12.5 cortices and fixed after 24 hours. Scale bar, 5 μm.