**Rarity and conservation status do not predict vertebrate population trends**

Gergana N. Daskalova[1]*, Isla H. Myers-Smith[1] and John L. Godlee[1]

**Classification:** Biological Sciences – Ecology

**Short title:** Rarity does not predict population trends

**Author affiliation:**

1. School of GeoSciences, University of Edinburgh, Edinburgh, UK

gndaskalova@gmail.com

johngodlee@gmail.com

isla.myers-smith@ed.ac.uk

* Corresponding author:

Gergana N. Daskalova

gndaskalova@gmail.com

Crew Building, King's Buildings

West Mains Road

Edinburgh EH9 1UU

Scotland

Tel: +44 7946274522

**Keywords:** population change, rarity, vertebrates, species attributes, geographic

range, habitat specificity, population size, Living Planet Database, GBIF, IUCN

**Submission type***:* Research report

**Abstract**

Species' attributes such as rarity and conservation status are often assumed to predict population change which in turn can influence conservation decision-making; however, there are very few empirical tests across taxa and biomes. We use a database of 9284 vertebrate populations from 2084 species in the UK and globally to quantify population change across rarity, Red List status, biomes and taxa. We found that rarity metrics (geographic range, population size and habitat specificity) did not predict population trends in the UK and globally, but greater fluctuations were more likely for smaller populations and species with larger ranges. There was no consistent pattern of net declines with increasing Red List status. We found higher-order taxonomic, but not geographic patterning of population change at global scales, with pronounced declines for amphibians. On a finer taxonomic scale, within amphibian, bird and reptile classes species, for which species-level phylogenies have been especially well resolved, there was no strong phylogenetic or taxanomic signal in the variation in population trends and fluctuations. Our findings suggest that rarity metrics and conservation status are poor predictors of vertebrate population trends of well monitored species, but that species with smaller populations were nevertheless more likely to fluctuate, potentially increasing their extinction risk. Thus, improved population monitoring across a diverse range of taxa is required, as species rarity and conservation status alone are insufficient to inform conservation prioritization.

**Significance statement**

Understanding which factors predispose species to population change is essential for improving conservation decision-making. The rarity and conservation status of a species are often assumed to predict population change and thus could inform conservation management. However, empirical tests of this assumption are lacking. We found that rarity and conservation status did not predict population change in the UK and globally, but species with smaller populations and larger ranges were more likely to fluctuate. We found taxonomic but not geographic patterning of population change among over 9000 populations from over 2000 vertebrate species, among which amphibian populations experienced net declines. Smaller populations were more likely to fluctuate and thus could have a higher likelihood of going extinct. Our findings suggest that the rarity and conservation status of species alone do not inform conservation prioritization, and improved monitoring of diverse taxa are required to capture global population change.

## Introduction

Species' attributes such as rarity, conservation status, phylogenetic relatedness, distribution and taxon are often assumed to predict variation in extinction risk and in population change (Julliard *et al.* 2004; Shultz *et al.* 2005; Tonini *et al.* 2016; Jetz & Pyron 2018). Population trends (Hutchings & Reynolds 2004) and population fluctuations (Melbourne & Hastings 2008) have both been frequently linked with a higher likelihood of a species' extinction. For decades, conservation efforts have focused on protecting rare species (Pigott 1977; Gunn 1980; Longton & Hedderson 2000; Povilitis 2001; Gaston & Fuller 2008), those with restricted geographic extents, small population sizes or high habitat specificity, as these species are thought to be more likely to be threatened with extinction. However, there are very few empirical tests of the influence of rarity and conservation status on population change and extinction risk (e.g., Purvis *et al.* 2000; O'Grady *et al.* 2004; Ripple *et al.* 2017), and those that exist mostly focus on either specific taxa (Jones *et al.* 2003; Cardillo *et al.* 2004; Fritz *et al.* 2009) or on population declines alone (Purvis *et al.* 2000; Fritz *et al.* 2009; Gilroy *et al.* 2016; Hefley *et al.* 2016; Ceballos *et al.* 2017), and not population change across taxa and geographic regions. Population change is the fundament process leading to community reassembly (Batt *et al.* 2017). And thus, in a period of accelerating human impacts and global change (Ehrlén & Morris 2015; Hefley *et al.* 2016), global analyses of population change are required to inform conservation policy and decision-making.

Understanding what factors best explain population change and extinction risk across species has been a key focus in population ecology and conservation biology, with possible measures including species' attributes and traits, such as rarity

(Gaston & Blackburn 1995; Owens & Bennett 2000; Purvis *et al.* 2000; Cardillo *et al.*

2004; O'Grady *et al.* 2004; Olden *et al.* 2007; Collen *et al.* 2009; Ripple *et al.* 2017).

Extinction risk, the likelihood that a species will go extinct within a given time frame

(Mace *et al.* 2008), is widely assumed to increase when species are rare (i.e., they

have small geographic ranges (Jones *et al.* 2003), small population sizes (Green

2003) and are very specialized in their habitat preferences (Gilroy *et al.* 2016). As

per population dynamics theory, species with small populations are more likely to

undergo stochastic fluctuations that could lead to extinction at any given point

(Lande 1993; Melbourne & Hastings 2008) and local population declines and

extinctions can lead to eventual global species extinction (Ceballos *et al.* 2017).

Small populations are also more vulnerable to demographic stochasticity and

inbreeding (Kareiva 1990; Hanski 1998). Allee effects, the relationship between

individual fitness and population density, further increase the likelihood of declines

due to lack of potential mates and low reproductive output once populations reach a

critically low density (Dennis *et al.* 2016; Sun 2016). Thus, small populations from

rare species are thought to be more likely to experience population declines

(McKinney 1997; Purvis *et al.* 2000; Manne & Pimm 2001; Hutchings & Reynolds

2004).

Investigating population change over time in relation to rarity can provide insights

into species' extinction risk. Rabinowitz (1981) defined 'seven forms of rarity' based

on the combinations of the extremes of three characteristics: geographic range

(small vs. large), habitat specificity (generalist vs. specialist), and local population

size (small vs. large). Rarity can be either a state in which species naturally occur, or

a state to which they are driven following external perturbations to populations

(Magurran & Henderson 2003). However, previous analyses of the influence of rarity

on population trends and extinction risk have shown mixed results. Geographic

range did not correlate with species' vulnerability to extinction in 45 bird and mammal

taxa from all continents except Antarctica (O'Grady *et al.* 2004) and the minimum

viable population size did not vary with geographic range and population size in 629

invertebrate, 529 vertebrate and 30 plant species (Brook *et al.* 2006). In contrast, in

bats, geographic range was found to be the most important predictor of extinction

risk (Jones *et al.* 2003). Thus, a generalizable relationship between rarity and

population declines and increased extinction risk across taxa and geographic regions

remains unquantified. However, simplifying the concept of rarity so that it can be

modelled as a predictor of population change remains a considerable challenge

(Magurran 2013; Batt *et al.* 2017).

Investigations of potential factors that predispose species to population change

across geographic and taxonomic scales (e.g., from country to biome level, and from

population to taxa level) are particularly valuable for guiding conservation decision-

making (O'Grady *et al.* 2004; Ehrlén & Morris 2015; Hefley *et al.* 2016; Batt *et al.*

2017). Conservation actions are predominantly implemented on smaller, country-

specific, regional or local scales (Mills *et al.* 2010; Guerrero *et al.* 2013), yet policy

may be determined for larger regions or broader taxonomic groups (Agardy 2005;

Pressey *et al.* 2007). Similarly, overall species' Red List status is determined on a

global scale based on multiple criteria (Mace *et al.* 2008), but on a local scale,

populations may vary in the direction and magnitude of experienced change (Ehrlén

& Morris 2015; Lawson *et al.* 2015; Gilroy *et al.* 2016; van Strien *et al.* 2016; Leung

*et al.* 2017). Population time-series data from the Living Planet Database (LPD),

complemented by extensive occurrence data from the Global Biodiversity

Information Facility (GBIF) and BirdLife (BirdLife International 2018), and information

on species habitat preferences and conservation status from the IUCN Red List,

provide a unique opportunity to test which species attributes explain global and

country-specific population trends across well-monitored vertebrate species.

Here, we addressed the research question: Is variation in vertebrate population

change explained by differences in species attributes, such as rarity metrics and

conservation status? We investigated two types of population change – trends and

fluctuations. We combined open source data from the LPD, GBIF and IUCN to

examine population change in the UK, a country with high-quality monitoring and

abundance data, and global population change across rarity, species' Red List

status, biomes, systems (freshwater, marine, terrestrial) and taxa. We considered

rarity to be the state in which species exist if their geographic range is small vs.

large, their local population size is low vs. high, or they are habitat generalists vs.

specialists. Thus, we quantified rarity through these three individual metrics -

geographic range extent, mean population size, habitat specificity, and additionally

species' Red List status.

We tested the following hypotheses: (1) rare species are more likely to be

experiencing negative population trends than common species on UK and global

scales; (2) declining populations are more likely for species from more threatened

Red List categories, (3) there are geographic, taxonomic and phylogenetic patterns

in vertebrate population trends and fluctuations (e.g., hotspots of declines, increases

or fluctuations in abundances in specific regions, for specific taxa and among

specific clades) and (4) rare species or those with a higher Red List status are more

likely to experience greater population fluctuations.

**Methods**

*Population data*

To quantify vertebrate population change, we extracted the abundance data for all

populations monitored at more than five different time points from the LPD (LPI

2016), resulting in 9284 populations from 2084 species for our system, biome, taxa

and conservation status analysis (global scale) and 508 populations from 237

species for our rarity analysis (UK scale, but note that habitat specificity data were

only available for 402 populations from 206 species, Table S1). Monitoring duration

differed among populations (Figures S1 and S2); however, in our analysis, we

retained all populations that had at least five time points of monitoring data. Before

analysis, we scaled the abundance of each population to a common magnitude

between zero and one, because the populations were monitored using different

metrics of abundance (e.g., population indices vs. number of individuals). This

allowed us to explore trends among populations relative to the variation experienced

across each time series.

Phylogenetic data

We obtained phylogenies for amphibian species from Jetz & Pyron (2018,

https://vertlife.org), for bird species from Jetz *et al.* (2012, https://birdtree.org), and

for reptile species from Tonini *et al.* (2016, https://vertlife.org). For each of the three

classes (*Amphibia*, *Aves* and *Reptilia*), we downloaded 100 trees and randomly

chose 10 for analysis (30 trees in total). We could not conduct phylogenetic analyses

of population change for the classes *Actinopterygii* and *Mammalia*, as species-level

phylogenies for them have not yet been resolved with high confidence (Foley *et al.*

2016; Tarver *et al.* 2016).

*Rarity metrics data*

We defined rarity following a simplified version of the 'seven forms of rarity' model

(Rabinowitz 1981), and thus consider rarity to be the state in which species exist

when they have a small geographic range, low population size, or narrow habitat

specificity. We combined publicly available data from three sources: (1) population

records for vertebrate species from the LPD (LPI 2016), (2) occurrence data from the

Global Biodiversity Information Facility (GBIF 2018) and range data from BirdLife

(BirdLife International 2018), and (3) habitat specificity and Red List classification

data for each species from the International Union for Conservation (IUCN 2017).

To estimate geographic range for species monitored in the UK, we calculated a $km^2$

occurrence area extent using a minimal convex hull approach based on occurrence

data for each species from GBIF (GBIF 2018). We filtered the GBIF data to remove

invalid records and outliers using the *CoordinateCleaner* package (Töpel *et al.*

2016). We excluded records with no decimal places in the decimal latitude or

longitude values, with equal latitude or longitude, within a one degree radius of the

GBIF headquarters in Copenhagen, within 0.0001 degrees of various biodiversity

institutions and within 0.1 degrees of capital cities. For each species, we excluded

the lower 0.02 and upper 0.98 quantile intervals of the latitude and longitude records

to account for outlier points that are records from zoos or other non-wild populations.

We drew a convex hull to most parsimoniously encompass all remaining occurrence

records using the *chull* function, and we calculated the area of the resulting polygon

using *areaPolygon* from the *geosphere* package (Hijmans 2016). We did not use

IUCN range maps, as they were not available for all of our study species, and

previous studies using GBIF occurrences to estimate range have found a positive

correlation between GBIF-derived and IUCN-derived geographic range (Phillips *et al.*

2017).  Extracting and filtering GBIF data and calculating range was computationally-

intensive and occurrence data availability was lower for certain species; thus, we did

not estimate geographic range for all species globally. Instead, we focused on

analyzing range effects for species monitored in the UK, a country with intensive and

detailed biodiversity monitoring of vertebrate species, and for birds globally, as they

are a very well-studied taxon. To estimate geographic range for bird species

monitored globally, we extracted the area of occurrence in $km^2$ for all bird species in

the LPD that had records in the BirdLife Data Zone (BirdLife International 2018).

We calculated mean population size across the monitoring duration for each

population using the raw population data. To create an index of habitat specificity,

we extracted the number of distinct habitats a species occupies based on the IUCN

habitat category for each species, accessed through the package *rredlist*

(Chamberlain 2017). We also quantified habitat specificity by surveying the number

of breeding and non-breeding habitats for each species from its online IUCN profile

(the 'habitat and ecology' section, IUCN 2017). The two approaches yielded similar

results (Figure S3, Table S2, key for the profiling method is presented in Table S4).

We obtained global conservation status data for all study species through the IUCN

Red List classification (IUCN 2017).

*Population change modelling*

As the first stage of our analysis, we modeled population change to produce a
population trend (slope and variance of population change over time). We used
state-space models that model abundance over time (freely-varying intercept) to
calculate the amount of overall population change experienced by each population
(*mu*), following Dennis *et al.* (2006) and Humbert *et al.* (2009). State-space models
account for process noise and observation error and thus deliver reliable estimates
of population change when working with large datasets where records were collected
using different approaches (Knape *et al.* 2011; Pedersen *et al.* 2011; Leung *et al.*
2017). Not accounting for process and measurement error could lead to over-
estimation of population declines (Rueda-Cediel *et al.* 2018). Positive *mu* values
indicate population increase and negative *mu* values indicate population decline. We
calculated the error around *mu* for each population (half of the difference between
the upper 95% confidence limit and the lower 95% confidence limit), with larger
errors indicating populations that have experienced more fluctuations (variance in
abundance metrics) overtime.

*Population trends across rarity metrics*

To test the influence of rarity metrics (geographic range, mean population size and
habitat specificity) on variation in population trends in the UK, we conducted a
second stage analysis using six Bayesian linear models (one per metric in the UK
and globally) based on Markov Chain Monte Carlo Bayesian inference (Hadfield
2010). The response variable was population trend (*mu* values from state-space
models), and fixed effects were geographic range (log transformed), mean
population size (log transformed) and habitat specificity (number of distinct habitats

occupied). We used weakly informative priors for all coefficients (an inverse Wishart

prior for the variances and a normal prior for the fixed effects), and we refer to

predictors as 'significant' when the 95% credible interval for the corresponding

parameter in the fitted models did not overlap zero. We did not include random

effects in the Bayesian models (e.g., a species or region random intercept), because

only 28% of the studied species had more than four monitored populations and

spatial replication was low (Table S1).

*Population trends across species' Red List status*

To investigate the relationship between population trends in the UK and globally, and

species' Red List status, we modelled population trends from the first stage of the

analysis (state-space models) as a function of Red List status (categorical variable)

using two Bayesian linear models (for the UK and global databases).

*Global population trends across latitude, biomes, systems and taxa*

To investigate the geographic and taxonomic patterns of population trends, we

estimated population trend from the first stage of the analysis (state-space models)

as a function of (1) latitude and biome (as defined by the 'biome' category in the

LPD, e.g., 'temperate broadleaf forest', Olson & Dinerstein 2002), (2) system

(freshwater, marine, terrestrial), and (3) taxon (*Actinopterygii*, bony fish; *Amphibia*,

amphibians; *Aves*, birds; *Mammalia*, mammals; *Reptilia*, reptiles). Each categorical

model from this second stage of our analyses was fitted with a zero intercept to allow

us to determine if net population trends differed from zero for each of the categories

under investigation.

*Global population trends within amphibian, bird and reptile classes*

To determine if there is a phylogenetic signal in the patterning of population change

within amphibian, bird and reptile taxa, we tested variation in population trends

among amphibian, bird and reptile species, the taxa whose phylogenies have been

resolved with high confidence (Jetz *et al.* 2012; Tonini *et al.* 2016; Jetz & Pyron

2018). We modelled population change to test the phylogenetic and species-level

taxonomic relatedness, which were both included as random effects in three (one

per class) Bayesian linear mixed effects models using the package *MCMCglmm*

(Hadfield 2010). The models did not include fixed effects. We used parameter-

expanded priors, with a variance-covariance structure that allows the slopes of

population trend (the *mu* values from the first stage analysis using state-space

models) to covary for each random effect. To account for phylogenetic uncertainty,

for each class, we ran ten models with identical structures but based on different

randomly selected phylogenetic trees. We report the mean estimates and their range

for each class. The phylogeny and taxonomy models were ran for 120 000 iterations

with a thining factor of ten and a burn-in period of 20 000 iterations. We assessed

significance of the random effects (phylogeny and species) by inspecting their

posterior distributions, with a distribution pushed up against zero indicating lack of

effect significance, since these distributions are always bounded by zero and have

only positive values.

*Population fluctuations across systems, taxa, Red List status and rarity metrics*

To test whether rare species fluctuate more than common species and whether there

is geographic and taxonomic patterning in population fluctuations, we modelled the

error around *mu* (half of the 95% confidence interval) as a response variable across

systems, taxa, Red List status and rarity metrics.

*Monitoring duration and units*

To assess the influence of monitoring duration on population trends, we used a

Bayesian linear model (Figures S1 and S2). We modelled population trend (*mu*) as a

function of monitoring duration (years) for each population, fitted with a zero

intercept, as when duration is zero, no population change has occurred. The units of

population monitoring across the LPD varied with 1368 different categorical levels

(e.g., population index, nest counts, numbers of individuals). Variation in population

trends were not explained by sampling method across the five most commonly used

units (Figure S5). We estimated population change using state-space models

specifically because of their suitability for analyses of large databases where records

were derived using different methods (Humbert *et al.* 2009; Leung *et al.* 2017).

*Comparison of modelling approaches*

We conducted the following supplementary analyses: in the second-stage Bayesian

models estimating population trends across systems, biomes, taxa and rarity

metrics, 1) we weighed *mu* values by the square of *tau*, the observation error

estimate derived from the state-space models, 2) we used slopes of linear model fits

of abundance (scaled at the population level, centered on zero and with a standard

deviation of one) instead of the *mu* estimates from state-space models, 3) we

modelled the standard error around the slope values of the linear models and the

standard deviation of the raw population data for each time-series as additional

metrics of population fluctuations. All different analytical approaches yielded very

similar results (see main text and Figures S6, S7 and S10, Table S2).

All statistical analyses were conducted using R (version 3.4.2, R Core Team 2017).

Code and data are publicly available on GitHub (link to be added at time of

publication).

**Results**

*Population trends in the UK and globally across rarity*

Across monitored vertebrate populations in the UK and globally, species with smaller

ranges, smaller population sizes, or narrower habitat specificity (i.e., rare species)

were not more prone to population declines than common species (Figure 1a-f,

Table S2).

**Figure 1. Effects of rarity metrics (geographic range, mean population size and habitat specificity) on vertebrate population trends in the UK (plots a, c, e, g) and globally (b, d, f, h).** The numbers on x axis indicate sample size for each category. Geographic range refers to the global range of UK species (a) and the global range of the bird species in the global LPD (b). The *mu* values of population change (plots a-f) are derived from state-space model fits of changes in abundance over the monitoring duration for each population. Effect sizes (plots g and h) are population trends (*mu*) calculated using linear models (light grey circles), state-space models (grey triangles), and population trends (*mu*) weighted by *tau* (black squares), the observation error estimate from the state-space models. Error bars show confidence intervals around population trend (*mu,* plots a-f) and credible intervals for *mu* across rarity metrics (plots g and h). Effect size coefficients (g and h) were standardized by dividing the effect size by the standard deviation of the corresponding input data. See Table S2 for full model outputs.

*Population trends in the UK and globally across Red List status*

Over the monitoring duration for populations in the UK, species classified as least concern and near threatened had a positive population trend (LC: slope = 0.01, CI = 0.004 to 0.02; NT: slope = 0.03, CI = 0.01 to 0.06), whereas vulnerable, endangered and critically endangered populations did not experience directional change (Figure 2b and Table S2). Globally, we found increasing, decreasing and stable populations across all Red List categories of conservation status. For example, a population of the least concern species red deer (*Cervus elaphus*) in Canada declined by 68% over 7 years going from 606 to 194 individuals and a population of the critically endangered Hawksbill sea turtle (*Eretmochelys imbricate*) from Barbados increased

by 269% over 7 years going from 89 to 328 individuals. Populations from least

concern species were more likely to increase (slope = 0.01, CI = 0.005 to 0.01),

whereas populations from vulnerable, endangered and critically endangered species

did not experience net directional change (Figure 2a). Populations from near

threatened species were more likely to decline (slope = -0.01, CI = -0.01 to -0.001).

Across all categories, there was no consistent pattern of greater population declines

in the UK and globally with increasing Red List status (Table S2). Most of the

populations in the LPD that were monitored at more than five time points were from

species categorized as least concern (Figure 2).

(a) Distribution of records

(b) Red List status effects (UK)

(c) Red List status effects (global)

**Figure 2. Distribution of population records by Red List status (plot a) and vertebrate population trends in the UK (plot b) and globally (plot c) across species' Red List status.** Colored points on the map indicate the Red List status of each population as indicated in plots b and c. Effect sizes (plots b and c) include the population trend (*mu*) calculated using linear models (circles), using state-

388 space and linear models (triangles), and population trend (*mu*) weighted by *tau*,

389 the observation error estimate from the state-space models (squares). The

390 numbers on the x axis show the sample size in each Red List category (plots b and

391 c). Error bars show 95% credible intervals (plots b and c). Effect size coefficients

392 (plots b and c) were standardized by dividing the effect size by the standard

393 deviation of the corresponding input data. See Table S2 for full model outputs.

395 *Global population trends across latitude and biomes*

396 Global vertebrate population trends did not show distinct geographic patterning, as

397 increasing, stable and declining populations were present on all continents and

398 across all latitudes (Figure 3a and b). Populations in Mediterranean forests, montane

399 grasslands, polar freshwaters, temperate wetlands, tropical forests and tropical coral

400 biomes were more likely to increase, whereas populations from the remaining

401 studied biomes were not more likely to be increasing versus decreasing (Figure 3c

402 and Table S2). The lack of biogeographic patterning in vertebrate population trends

403 was also apparent on a UK scale (Figure S6 and Table S2).

**Figure 3. Lack of biogeographic patterning in global vertebrate population trends (plot a), latitudinal distribution of population trends (plot b), and population trends across biomes (plot c).** Results include 9284 populations from 2084 species. Effect sizes (plot c) include the population trend (*mu*) calculated using linear models (light grey circles), state-space models (grey triangles), and population trend (*mu*) weighted by *tau*, the observation error estimate from the state-space models (black squares). The numbers on the x axis show the sample size for each biome category (plot c). Error bars show 95% credible intervals (plot c). Effect size coefficients (plot c) were standardized by dividing the effect size by the standard deviation of the corresponding input data. See Table S2 for full model outputs.

*Global population trends across systems and taxa*

Globally, monitored vertebrate populations experienced population increases in all

three studied systems (freshwater slope = 0.005, CI = 0.002 to 0.01; marine slope =

0.004, CI = 0.002 to 0.01; terrestrial slope = 0.003, CI = 0.001 to 0.005, Figure 2a-b,

Table S2). In freshwater and terrestrial habitats, there was a bimodal distribution of

population change (Hartigans' dip test, $D = 0.04$, $p < 0.01$, Figure S7a), likely driven

by the bimodal distribution for bird populations (Hartigans' dip test, $D = 0.04$, $p <$

$0.01$, Figure S7b). Amphibian populations had net declines (slope = -0.01, CI = -0.02

to -0.005), whereas birds, mammals and reptiles had net increases (slope = 0.004,

CI = 0.003 to 0.01; slope = 0.01, CI = 0.01 to 0.01; slope = 0.02, CI = 0.01 to 0.02).

Bony fish populations were centred on zero (slope = -0.001, CI = -0.004 to 0.002).

(a) Distribution of records

(b) Distribution of records

(c) Monitoring duration

(d) Monitoring duration

(e) Across systems

(f) Across taxa

**Figure 4. Distribution of records across (plot a) freshwater, marine and**

**terrestrial systems and (plot b) bony fish, amphibians, birds, mammals and**

**reptiles, monitoring duration (plots c and d) and their respective population**

**trend (*mu*) effect sizes (plots e and f).** The numbers on x axis indicate sample size

for each category. Effect sizes are population trends (*mu*) from linear models

(circles) calculated using state-space models (triangles), and population trend (*mu*)

weighted by *tau*, the observation error estimate from the state-space models

(squares, plots e and f). The numbers on the x axis show the sample size for each

taxa category (plots e and f). Error bars show 95% credible intervals (plots e and f).

Effect size coefficients (plots e and f) were standardized by dividing the effect size by

the standard deviation of the corresponding input data. See Table S2 for full model

outputs.

*Population fluctuations across systems, taxa, Red List status and rarity metrics*

Global population fluctuations (variation in population abundance over time) did not

show distinct geographic patterning (Figure 5a-b). Populations that experienced

more fluctuations had in general smaller mean population sizes (across all taxa,

slope = -0.004, CI = -0.01 to -0.002) and for bird species had larger geographic

ranges (slope = 0.005, CI = 0.003 to 0.01, Figure 5c). On a global scale, population

fluctuations were more common with increasing Red List threat level (LC: slope =

0.11. CI = 0.11 to 0.12; CR: slope = 0.17, CI = 0.15 to 0.19, Figure 5d, see Table S2

for all model outputs). Across the three studied systems, terrestrial populations

experienced lower population fluctuations (slope = 0.10, CI = 0.10 to 0.11, Figure

5e). Across taxa, fluctuations were most common for amphibian populations (slope =

0.22, CI = 0.20 to 0.23, Figure 5f).

**Figure 5. Global population fluctuations in vertebrate species (plot a) across latitude (plot b), rarity metrics (global scale, plot c), Red List status (global scale, plot d), systems (plot e) and taxa (plot f).** The numbers on x axis indicate sample size for each category. The response variables in the models were: 1) the standard error around the slope estimates of the linear model fits of abundance versus year (circles), 2) half of the 95% confidence interval around the *mu* value of population change (triangles), 3) half of the 95% confidence interval around *mu* weighted by *tau*, the observation error estimate from the state-space models (squares), and 4) the standard deviation of the raw data for each population time-series (plus symbols).  Larger errors indicate greater population fluctuations. Effect size coefficients (c-f) were standardized by dividing the effect size by the standard deviation of the corresponding input data. Error bars show 95% credible intervals (c-f). See Table S2 for full model outputs.

*Global population trends and fluctuations within amphibian, bird and reptile classes*

When we tested for finer scale species-level taxonomic clustering of population change (both overall population trends and population fluctuations) within amphibian, bird and reptile classes, we did not detect a phylogenetic relatedness signal in the patterning of population change for either trends or fluctuations (Figures 6 and S8, Table S2). Similarly, species identity within amphibian, bird and reptile classes also did not explain variation in population trends or fluctuations (Figures 6 and S8, Table S2). There were no distinct clusters of specific clades that were more likely undergo increases, decreases or fluctuations in populations (Figure 6).

**Figure 6. No distinct patterning in population trends (average *mu* across the population trends of each species, plots a-c) and in population fluctuations**

**(plots d-f) was found across 69 amphibian species, 806 bird species and 79**

**reptile species.** Heat maps around phylogenetic trees indicate population change

with darker blue indicating species with declining populations, green indicating

species with populations with no change and yellow indicating species with

increasing populations across monitored species in the Living Planet Database.

Density plots show mean population change across the sampled species in each

class. The phylogenies were obtained from Jetz & Pyron (2018) for ampibians

(https://vertlife.org), from Jetz *et al.* (2012) for birds (https://birdtree.org), and from

Tonini *et al.* (2016) for reptiles (https://vertlife.org). For model outputs, see Figure S8

and Table S2.

*Global population trends across monitoring duration*

Monitoring duration was weakly positively related to vertebrate population trends,

with slightly more increasing populations found for longer duration studies (Figure

S9, Table S2).

**Discussion**

Taken together, our findings do not support the application of rarity metrics as

generalizable predictors of population trends in well-monitored vertebrate

populations. Across monitored vertebrate populations in the UK and globally, rarity

metrics (geographic range, population size and habitat specificity) and species' Red

List status did not explain variation in population trends (Figures 1 and 2). There was

no consistent pattern of greater population declines with increasing Red List status in

either the UK or globally (Figures 1e, f and 2). There was no biogeographic

patterning of population trends and fluctuations, as populations increased, remained

stable and declined across different biomes on all continents (Figure 3). Among

higher-order taxonomic units (class), birds, mammals and reptiles increased, bony

fish did not experience net directional change and amphibians were the only taxon

with overall declines, as well as the taxon that experienced the most population

fluctuations (Figures 4-5). On a finer taxonomic scale, across amphibian, bird and

reptile species, there were no distinct phylogenetic clusters of closely-related species

experiencing similar population trends or fluctuations. Fluctuations were more likely

for species with smaller population sizes and broader habitat preferences (Figure 5).

Thus, although we did not find evidence that rarity metrics explain variation in

population trends, rare species were nevertheless more likely to fluctuate, potentially

increasing their extinction risk.

*Population change across rarity metrics*

The power of rarity metrics to explain population trends could be mediated by the

type of threats to which populations are exposed (Schemske *et al.* 1994), as well as

whether species are naturally rare, or have become rare due to external drivers in

recent time (de Lange & Norton 2004; Robbirt *et al.* 2006; Harrison *et al.* 2008).

Geographic range, population size and habitat specificity may be of lesser

importance in human-dominated landscapes where threats such as habitat change,

pollution and exploitation are more prevalent (Newmark 1996; Purvis *et al.* 2000;

O'Grady *et al.* 2004; Brook *et al.* 2006, Sullivan *et al.* 2016 but see Owens & Bennett

2000 and Julliard *et al.* 2004). For UK and global vertebrate populations, the lack of

rarity effects on population trends (Figure 1) could be related to urbanization,

intensive land management and/or high investment in conservation (Keith *et al.*

2015), which may be contributing to the observed increases in species classified as

near threatened by the IUCN (Figure 1f). Species with larger geographic ranges and

small populations were most likely to fluctuate (Figure 5). A higher likelihood of

fluctuations for populations with smaller mean population size (Figure 5c) is in line

with population dynamics theory, suggesting that even though rarity metrics did not

predict population trends, they might still be good indicators of extinction risk, as

greater fluctuations could in time lead to extinction (Lande 1993; Melbourne &

Hastings 2008).

*Population change across Red List status*

Across all Red List status categories, there was no consistent pattern of net declines

in the UK and globally with increasing threat level (Figure 2). Populations from

vulnerable and endangered species did not experience directional change,

suggesting that local scale conservation actions might have stabilized populations for

species which are declining on a global scale (Saterson *et al.* 2004; Vickery *et al.*

2004; Gillings *et al.* 2005; Hoffmann *et al.* 2010). The threats and mitigation actions

to which populations with more threatened Red List statuses are exposed on a local

scale are likely heterogeneous across different biomes (Pereira *et al.* 2012; Maxwell

*et al.* 2016). The finding that populations from species with higher Red List status

were more likely to fluctuate (Figure 5d) relates to one of the Red List criteria –

number of breeding individuals (Mace *et al.* 2008). Extinction risk is higher when

there are very few breeding individuals due to Allee effects (Dennis *et al.* 2016; Sun

2016). Thus, although we did not find evidence to support the hypothesis that

species with a higher Red List status are more likely to be declining, those

populations did experience higher population fluctuations, suggesting that species of

concern might still have a higher risk of extinction.

*Global population change across latitude, biomes and systems*

Global vertebrate population trends varied among taxa but did not show distinct

geographic patterning (Figures 3-4). Some biomes like Mediterranean forests were

more likely to include increasing populations, whilst populations in other biomes did

not experience directional population change (Figure 3 and Table S2). Similar

heterogeneity in the patterns of population trends across habitats has been

documented in the Netherlands (van Strien *et al.* 2016) and in temperate North

America and Europe (Leung *et al.* 2017) also using the LPD, suggesting that local

scale environmental factors or proximity to anthropogenic drivers of change may be

better predictors of population trends (Brook *et al.* 2006; Fritz *et al.* 2009, but see

Phillips *et al.* 2017). The lack of distinct geographic patterning of population trends

and fluctuations around the world (Figures 3, 4 and 6a, b and e) might be attributable

to either stochastic (random events) and deterministic (e.g., anthropogenic threats)

processes influencing inter-annual population change (Kareiva 1990; Lundberg *et al.*

2000). Thus, to attribute population trends and fluctuations to anthropogenic drivers,

we need to go beyond previous studies (Ceballos *et al.* 2017; Davidson *et al.* 2017)

and quantify the full spectrum of population change including positive, negative and

stable trends and fluctuations. Quantifying the drivers of population change is a

critical first step towards disentangling the pathways from population change to

biodiversity change in the Anthropocene (Leung *et al.* 2017; Yoccoz *et al.* 2018).

*Global population change across taxa*

Taxon-specific factors including reproductive strategy, trophic level, generation time,

life history traits (Purvis *et al.* 2000; Cardillo *et al.* 2004), or sampling effects

(Magurran *et al.* 2010) could explain why some taxa were more likely to increase

(birds, mammals and reptiles), whereas others were more likely to decline

(amphibians) or remain stable (bony fish), (Figure 4 and Table S2). However, within

the classes *Amphibia*, *Aves* and *Reptilia*, phylogenetic relatedness and species-level

taxanomic classification did not predict variation in population trends and

fluctuations. Similar lack of phylogenetic dependencies has been detected for the

population growth rates of migratory birds (Moller *et al.* 2008). Among seabirds,

however, breeding phenology varied based on phylogeny and taxonomy (Keogan *et*

*al.* 2018). Among squamates and amphibians, there is evidence that phylogenetic

relatedness predicts extinction (Tonini *et al.* 2016; Jetz & Pyron 2018), a process

that occurs over much longer time-scales than the population change we study here.

Amphibians were the only taxa to experience net population declines as well as

greater population fluctuations among monitored vertebrate species in the LPD

(Figure 4f and 5f). The global amphibian decline (Houlahan *et al.* 2000), also

demonstrated with this dataset at the temperate North American and European scale

(Leung *et al.* 2017), has been linked to the spread of a fungal disease (chytrid

fungus, *Batrachochytrium dendrobatidis*), facilitated by global warming (Pounds *et al.*

2006), as well as habitat loss and Allee effects in small populations (Green 2003).

*Study limitations*

Our study is limited by three main factors: 1) the gaps in global monitoring of

populations, 2) challenges in estimating geographic range and habitat specificity,

and 3) overrepresentation of common species. The LPD under-represents

populations outside of Europe and North America (Pereira & Cooper 2006; Proença

*et al.* 2016; McRae *et al.* 2017) and includes different species abundance measures

across populations that have been monitored for different durations (Figures S1, S2

and S6). The GBIF database includes sub-optimal representation of some species

and there might be confusion between sub-species and occurrence outliers (Yesson

*et al.* 2007; Boakes *et al.* 2010), despite being the largest international open source

database of species locality data (Yesson *et al.* 2007). In addition, we used a static

measure of geographic range, which does not account for changes in species

distributions over time. For example, species could naturally have a small range or

the small range size could be due to historic habitat loss (Magurran & Henderson

2003; Prendergast *et al.* 1993). In addition, UK populations included in the LPD are

predominantly from species with wide geographic ranges (Table S4). The

relationship between population change and geographic range could differ for highly

endemic species with restricted distribution that are not monitored or included in the

LPD dataset.

*Conclusion*

In summary, rarity metrics did not explain vertebrate population trends and

we did not find a consistent pattern of net declines with increasing Red List status in

the UK and globally. We found higher-order taxonomic but not geographic patterning

of population change around the world, yet within-classes, phylogeny and species

identity did not explain variation in population trends and fluctuations among

amphibian, bird and reptile species. Vertebrate populations experienced different

magnitudes of net change and population fluctuations, which in the future may lead

to shifts in community composition as some species become locally extinct and

others expand into new habitats. Our findings caution the use of rarity metrics for the

prediction of future global population trends, but suggest that such metrics, in

particular population size, are nevertheless indicators of population fluctuations,

which might ultimately link to increased species extinction risk. It is only with

improved monitoring of diverse taxa that we can test the drivers of global population

change during the Anthropocene to inform wildlife conservation.

**Statement of authorship**

GND and IMS conceived the idea and conducted the statistical analyses. JLG

contributed to the calculation of geographic range estimates. All authors contributed

to the integration of the LPI, GBIF and IUCN databases. GND wrote the first draft of

the manuscript and all authors contributed to revisions.

**Conflict of interest**

The authors declare no conflicts of interest.

**Data accessibility statement**

We will make all R code for the analyses and data publicly available on GitHub and

will include the relevant DOIs and links.

**Acknowledgements**

We thank the WWF and the RZSL for compiling and maintaining the Living Planet

Database. We thank Dmitry Schigel and Jan Legind at GBIF for providing access to

a direct download link for the UK species occurrence data. We are grateful to the

Critical Thinking in Ecology and Environmental Sciences 2015-16 tutorial group at

the University of Edinburgh including Hannah Louise Stevens, Ashlie Nithsdale, Lisa

Kopsieker and Claudia Ardrey who helped us conceive the idea for the analysis. We

thank Dan Greenberg for advice on state-space models and Kat Keogan and Albert

Phillimore for advice on phylogenetic models. We thank Anne Magurran, Albert

Phillimore and Mark Vellend for providing very helpful feedback on the manuscript.

Funding for this research was provided by the NERC doctoral training partnership

grant (NE/L002558/1) at the University of Edinburgh and the Carnegie Trust for the

Universities of Scotland.

**References**

Agardy, T. (2005). Global marine conservation policy versus site-level

implementation. *Mar. Ecol. Prog. Ser.*, 300, 242–248.

Batt, R.D., Morley, J.W., Selden, R.L., Tingley, M.W. & Pinsky, M.L. (2017). Gradual

changes in range size accompany long-term trends in species richness. *Ecol.*

*Lett.*, 20, 1148–1157.

BirdLife International. (2018). IUCN Red List for birds. Downloaded from

http://www.birdlife.org in March 2018.

Boakes, E.H., McGowan, P.J., Fuller, R.A., Chang-qing, D., Clark, N.E., O'Connor,

K., *et al.* (2010). Distorted views of biodiversity: spatial and temporal bias in

species occurrence data. *PLoS Biol.*, 8, e1000385.

Brook, B.W., Traill, L.W. & Bradshaw, C.J.A. (2006). Minimum viable population

sizes and global extinction risk are unrelated: MVP is unrelated to global

extinction risk. *Ecol. Lett.*, 9, 375–382.

Cardillo, M., Purvis, A., Sechrest, W., Gittleman, J.L., Bielby, J. & Mace, G.M.

(2004). Human Population Density and Extinction Risk in the World's Carnivores.

*PLoS Biol.*, 2, e197.

Ceballos, G., Ehrlich, P.R. & Dirzo, R. (2017). Biological annihilation via the ongoing

sixth mass extinction signaled by vertebrate population losses and declines.

*Proc. Natl. Acad. Sci.*, 201704949.

Chamberlain, S. (2017). rredlist: "IUCN" Red List Client. R package version 0.4.0.

https://CRAN.R-project.org/package=rredlist.

Collen, B., Loh, J., Whitmee, S., McRAE, L., Amin, R. & Baillie, J.E.M. (2009).

Monitoring Change in Vertebrate Abundance: the Living Planet Index. *Conserv.*

*Biol.*, 23, 317–327.

Davidson, A.D., Shoemaker, K.T., Weinstein, B., Costa, G.C., Brooks, T.M.,

Ceballos, G., *et al.* (2017). Geography of current and future global mammal

extinction risk. *PLOS ONE*, 12, e0186934.

Dennis, B., Assas, L., Elaydi, S., Kwessi, E. & Livadiotis, G. (2016). Allee effects and

resilience in stochastic populations. *Theor. Ecol.*, 9, 323–335.

Dennis, B., Ponciano, J.M., Lele, S.R., Taper, M.L. & Staples, D.F. (2006).

Estimating density dependence, process noise, and observation error. *Ecol.*

*Monogr.*, 76, 323–341.

Ehrlén, J. & Morris, W.F. (2015). Predicting changes in the distribution and

abundance of species under environmental change. *Ecol. Lett.*, 18, 303–314.

Foley, N.M., Springer, M.S. & Teeling, E.C. (2016). Mammal madness: is the

mammal tree of life not yet resolved? *Philos. Trans. R. Soc. B Biol. Sci.*, 371,

20150140.

Fritz, S.A., Bininda-Emonds, O.R.P. & Purvis, A. (2009). Geographical variation in

predictors of mammalian extinction risk: big is bad, but only in the tropics. *Ecol.*

*Lett.*, 12, 538–549.

Gaston, K. & Fuller, R. (2008). Commonness, population depletion and conservation

biology. *Trends Ecol. Evol.*, 23, 14–19.

Gaston, K.J. & Blackburn, T.M. (1995). Birds, body size and the threat of extinction.

*Philos. Trans. R. Soc. Lond. B Biol. Sci.*, 347, 205–212.

GBIF. (2017). Biodiversity occurrence data. http://www.gbif.org/ Downloaded Feb

2016 - Apr 2017.

GBIF. (2018). Species occurrence data.

Gillings, S., Newson, S.E., Noble, D.G. & Vickery, J.A. (2005). Winter availability of

cereal stubbles attracts declining farmland birds and positively influences

breeding population trends. *Proc. R. Soc. B Biol. Sci.*, 272, 733–739.

Gilroy, J.J., Gill, J.A., Butchart, S.H.M., Jones, V.R. & Franco, A.M.A. (2016).

Migratory diversity predicts population declines in birds. *Ecol. Lett.*, 19, 308–317.

Green, D.M. (2003). The ecology of extinction: population fluctuation and decline in

amphibians. *Biol. Conserv.*, 111, 331–343.

Guerrero, A.M., McALLISTER, R.R.J., Corcoran, J. & Wilson, K.A. (2013). Scale

Mismatches, Conservation Planning, and the Value of Social-Network Analyses:

Scale Mismatches and Conservation Planning. *Conserv. Biol.*, 27, 35–44.

Gunn, A. (1980). Why should we care about rare species? *Environ. Ethics*, 2, 17–37.

Hadfield, J.D. (2010). MCMC Methods for Multi-Response Generalized Linear Mixed

Models: The MCMCglmm R Package. *J. Stat. Softw.*, 33, 1–22.

Hanski, I. (1998). Metapopulation dynamics. *Nature*, 396, 41.

Harrison, S., Viers, J.H., Thorne, J.H. & Grace, J.B. (2008). Favorable environments

and the persistence of naturally rare species. *Conserv. Lett.*, 1, 65–74.

Hefley, T.J., Hooten, M.B., Drake, J.M., Russell, R.E. & Walsh, D.P. (2016). When

can the cause of a population decline be determined? *Ecol. Lett.*, 19, 1353–1362.

Hijmans, R.J. (2016). geosphere: Spherical Trigonometry. R package version 1.5-5.

https://CRAN.R-project.org/package=geosphere.

Hoffmann, M., Hilton-Taylor, C., Angulo, A., Bohm, M., Brooks, T.M., Butchart,

S.H.M., *et al.* (2010). The Impact of Conservation on the Status of the World's

Vertebrates. *Science*, 330, 1503–1509.

Houlahan, J.E., Findlay, C.S., Schmidt, B.R., Meyer, A.H. & Kuzmin, S.L. (2000).

Quantitative evidence for global amphibian population declines. *Nature*, 404,

752–755.

Humbert, J.-Y., Scott Mills, L., Horne, J.S. & Dennis, B. (2009). A better way to

estimate population trends. *Oikos*, 118, 1940–1946.

Hutchings, J.A. & Reynolds, J.D. (2004). Marine fish population collapses:

consequences for recovery and extinction risk. *AIBS Bull.*, 54, 297–309.

IUCN. (2017). Conservation status data. http://www.iucnredlist.org/ Downloaded Jan-

April 2017.

Jetz, W. & Pyron, R.A. (2018). The interplay of past diversification and evolutionary

isolation with present imperilment across the amphibian tree of life. *Nat. Ecol.*

*Evol.*

Jetz, W., Thomas, G.H., Joy, J.B., Hartmann, K. & Mooers, A.O. (2012). The global

diversity of birds in space and time. *Nature*, 491, 444–448.

Jones, K.E., Purvis, A. & Gittleman, J.L. (2003). Biological Correlates of Extinction

Risk in Bats. *Am. Nat.*, 161, 601–614.

Julliard, R., Jiguet, F. & Couvet, D. (2004). Common birds facing global changes:

what makes a species at risk? *Glob. Change Biol.*, 10, 148–154.

Kareiva, P. (1990). Population dynamics in spatially complex environments: theory

and data. *Phil Trans R Soc Lond B*, 330, 175–190.

Keith, D., Akçakaya, H.R., Butchart, S.H.M., Collen, B., Dulvy, N.K., Holmes, E.E., *et*

*al.* (2015). Temporal correlations in population trends: Conservation implications

from time-series analysis of diverse animal taxa. *Biol. Conserv.*, 192, 247–257.

Keogan, K., Daunt, F., Wanless, S., Phillips, R.A., Walling, C.A., Agnew, P., *et al.*

(2018). Global phenological insensitivity to shifting ocean temperatures among

seabirds. *Nat. Clim. Change*, 8, 313–318.

Knape, J., Jonzén, N. & Sköld, M. (2011). On observation distributions for state

space models of population survey data: Observation models for population data.

*J. Anim. Ecol.*, 80, 1269–1277.

Lande, R. (1993). Risks of population extinction from demographic and

environmental stochasticity and random catastrophes. *Am. Nat.*, 142, 911–927.

de Lange, P.J. & Norton, D.A. (2004). The ecology and conservation of Kunzea

sinclairii (Myrtaceae), a naturally rare plant of rhyolitic rock outcrops. *Biol.*

*Conserv.*, 117, 49–59.

Lawson, C.R., Vindenes, Y., Bailey, L. & van de Pol, M. (2015). Environmental

variation and population responses to global change. *Ecol. Lett.*, 18, 724–736.

Leung, B., Greenberg, D.A. & Green, D.M. (2017). Trends in mean growth and

stability in temperate vertebrate populations. *Divers. Distrib.*, 23, 1372–1380.

Longton, R.E. & Hedderson, T.A. (2000). What are rare species and why conserve

them? *Lindbergia*, 25, 53–61.

LPI. (2016). Living Planet Index database, 2016. www.livingplanetindex.org/

Downloaded February 2016.

Lundberg, P., Ranta, E., Ripa, J. & Kaitala, V. (2000). Population variability in space

and time. *Trends Ecol. Evol.*, 15, 460–464.

Mace, G.M., Collar, N.J., Gaston, K.J., Hilton-Taylor, C., AkçAkaya, H.R., Leader-

Williams, N., *et al.* (2008). Quantification of Extinction Risk: IUCN's System for

Classifying Threatened Species. *Conserv. Biol.*, 22, 1424–1442.

Magurran, A. (2013). *Measuring biological diversity*. John Wiley & Sons.

Magurran, A.E., Baillie, S.R., Buckland, S.T., Dick, J.M., Elston, D.A., Scott, E.M., *et*

*al.* (2010). Long-term datasets in biodiversity research and monitoring: assessing

change in ecological communities through time. *Trends Ecol. Evol.*, 25, 574–582.

Magurran, A.E. & Henderson, P.A. (2003). Explaining the excess of rare species in

natural species abundance distributions. *Nature*, 422, 714.

Manne, L.L. & Pimm, S.L. (2001). Beyond eight forms of rarity: which species are

threatened and which will be next? *Anim. Conserv.*, 4, 221–229.

Maxwell, S.L., Fuller, R.A., Brooks, T.M. & Watson, J.E. (2016). Biodiversity: The

ravages of guns, nets and bulldozers. *Nature*, 536, 143–145.

McKinney, M.L. (1997). Extinction vulnerability and selectivity: combining ecological

and paleontological views. *Annu. Rev. Ecol. Syst.*, 28, 495–516.

McRae, L., Deinet, S. & Freeman, R. (2017). The Diversity-Weighted Living Planet

Index: Controlling for Taxonomic Bias in a Global Biodiversity Indicator. *PLOS*

*ONE*, 12, e0169156.

Melbourne, B.A. & Hastings, A. (2008). Extinction risk depends strongly on factors

contributing to stochasticity. *Nature*, 454, 100–103.

Mills, M., Pressey, R.L., Weeks, R., Foale, S. & Ban, N.C. (2010). A mismatch of

scales: challenges in planning for implementation of marine protected areas in

the Coral Triangle: Mismatch of scales in conservation planning. *Conserv. Lett.*,

3, 291–303.

Moller, A.P., Rubolini, D. & Lehikoinen, E. (2008). Populations of migratory bird

species that did not show a phenological response to climate change are

declining. *Proc. Natl. Acad. Sci.*, 105, 16195–16200.

Newmark, W.D. (1996). Insularization of Tanzanian parks and the local extinction of

large mammals. *Conserv. Biol.*, 10, 1549–1556.

O'Grady, J.J., Reed, D.H., Brook, B.W. & Frankham, R. (2004). What are the best

correlates of predicted extinction risk? *Biol. Conserv.*, 118, 513–520.

Olden, J.D., Hogan, Z.S. & Zanden, M.J.V. (2007). Small fish, big fish, red fish, blue

fish: size-biased extinction risk of the world's freshwater and marine fishes. *Glob.*

*Ecol. Biogeogr.*, 16, 694–701.

Olson, D.M. & Dinerstein, E. (2002). The Global 200: Priority Ecoregions for Global

Conservation. *Ann. Mo. Bot. Gard.*, 89, 199.

Owens, I.P. & Bennett, P.M. (2000). Ecological basis of extinction risk in birds:

habitat loss versus human persecution and introduced predators. *Proc. Natl.*

*Acad. Sci.*, 97, 12144–12148.

Pedersen, M.W., Berg, C.W., Thygesen, U.H., Nielsen, A. & Madsen, H. (2011).

Estimation methods for nonlinear state-space models in ecology. *Ecol. Model.*,

222, 1394–1400.

Pereira, H.M. & Cooper, D.H. (2006). Towards the global monitoring of biodiversity

change. *Trends Ecol. Evol.*, 21, 123–129.

Pereira, H.M., Navarro, L.M. & Martins, I.S. (2012). Global Biodiversity Change: The

Bad, the Good, and the Unknown. *Annu. Rev. Environ. Resour.*, 37, 25–50.

Phillips, H.R.P., Newbold, T. & Purvis, A. (2017a). Land-use effects on local

biodiversity in tropical forests vary between continents. *Biodivers. Conserv.*, 26,

2251–2270.

Phillips, H.R.P., Newbold, T. & Purvis, A. (2017b). Land-use effects on local

biodiversity in tropical forests vary between continents. *Biodivers. Conserv.*, 26,

2251–2270.

Pigott, C.D. & Walters, S.M. (1977). The scientific basis of practical conservation:

aims and methods of conservation. In: *Proc. R. Soc. Lond. B*. The Royal Society,

pp. 59–68.

Pounds, A.J., Bustamante, M.R., Coloma, L.A., Consuegra, J.A., Fogden, M.P.L.,

Foster, P.N., *et al.* (2006). Widespread amphibian extinctions from epidemic

disease driven by global warming. *Nature*, 439, 161–167.

Povilitis, T. (2001). Toward a robust natural imperative for conservation. *Conserv.*

*Biol.*, 15, 533–535.

Prendergast, J.R., Quinn, R.M., Lawton, J.H., Eversham, B.C. & Gibbons, D.W.

(1993). Rare species, the coincidence of diversity hotspots and conservation

strategies. *Nature*, 365, 335–337.

Pressey, R.L., Cabeza, M., Watts, M.E., Cowling, R.M. & Wilson, K.A. (2007).

Conservation planning in a changing world. *Trends Ecol. Evol.*, 22, 583–592.

Proença, V., Martin, L.J., Pereira, H.M., Fernandez, M., McRae, L., Belnap, J., *et al.*

(2016). Global biodiversity monitoring: From data sources to Essential

Biodiversity Variables. *Biol. Conserv.*

Purvis, A., Gittleman, J.L., Cowlishaw, G. & Mace, G.M. (2000). Predicting extinction

risk in declining species. *Proc. R. Soc. B Biol. Sci.*, 267, 1947–1952.

R Core Team. (2017). R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria.

Rabinowitz, D. (1981). Seven forms of rarity. In: *The biological aspects of rare plants*

*conservation*. pp. 205–217.

Ripple, W.J., Wolf, C., Newsome, T. M., Hoffmann, M., Wirsing, A. J. & McCauley, D.

J. (2017). Extinction risk is most acute for the world's largest and smallest

vertebrates. *114*, 10678–10683.

Robbirt, K.M., Roberts, D.L. & Hawkins, J.A. (2006). Comparing IUCN and

Probabilistic Assessments of Threat: Do IUCN Red List Criteria Conflate Rarity

and Threat? *Biodivers. Conserv.*, 15, 1903–1912.

Rueda-Cediel, P., Anderson, K.E., Regan, T.J. & Regan, H.M. (2018). Effects of

uncertainty and variability on population declines and IUCN Red List

classifications. *Conserv. Biol.*

Saterson, K.A., Christensen, N.L., Jackson, R.B., Kramer, R.A., Pimm, S.L., Smith,

M.D., *et al.* (2004). Disconnects in Evaluating the Relative Effectiveness of

Conservation Strategies. *Conserv. Biol.*, 18, 597–599.

Schemske, D.W., Husband, B.C., Ruckelshaus, M.H., Goodwillie, C., Parker, I.M. &

Bishop, J.G. (1994). Evaluating Approaches to the Conservation of Rare and

Endangered Plants. *Ecology*, 75, 584–606.

Shultz, S., B. Bradbury, R., L. Evans, K., D. Gregory, R. & M. Blackburn, T. (2005).

Brain size and resource specialization predict long-term population trends in

British birds. *Proc. R. Soc. B Biol. Sci.*, 272, 2305–2311.

van Strien, A.J., Meyling, A.W.G., Herder, J.E., Hollander, H., Kalkman, V.J., Poot,

M.J.M., *et al.* (2016). Modest recovery of biodiversity in a western European

country: The Living Planet Index for the Netherlands. *Biol. Conserv.*, 200, 44–50.

Sun, G.-Q. (2016). Mathematical modeling of population dynamics with Allee effect.

*Nonlinear Dyn.*, 85, 1–12.

Tarver, J.E., dos Reis, M., Mirarab, S., Moran, R.J., Parker, S., O'Reilly, J.E., *et al.*

(2016). The Interrelationships of Placental Mammals and the Limits of

Phylogenetic Inference. *Genome Biol. Evol.*, 8, 330–344.

Tonini, J.F.R., Beard, K.H., Ferreira, R.B., Jetz, W. & Pyron, R.A. (2016). Fully-

sampled phylogenies of squamates reveal evolutionary patterns in threat status.

*Biol. Conserv.*, 204, 23–31.

Töpel, M., Zizka, A., Calió, M.F., Scharn, R., Silvestro, D. & Antonelli, A. (2016).

SpeciesGeoCoder: Fast Categorization of Species Occurrences for Analyses of

Biodiversity, Biogeography, Ecology, and Evolution. *Syst. Biol.*, syw064.

Vickery, J.A., Bradbury, R.B., Henderson, I.G., Eaton, M.A. & Grice, P.V. (2004). The

role of agri-environment schemes and farm management practices in reversing

the decline of farmland birds in England. *Biol. Conserv.*, 119, 19–39.

Yesson, C., Brewer, P.W., Sutton, T., Caithness, N., Pahwa, J.S., Burgess, M., *et al.*

(2007). How Global Is the Global Biodiversity Information Facility? *PLoS ONE*, 2,

e1124.

Yoccoz, N.G., Ellingsen, K.E. & Tveraa, T. (2018). Biodiversity may wax or wane

depending on metrics or taxa. *Proc. Natl. Acad. Sci.*, 115, 1681–1683.

**Supplementary Information**

 **Table S1. Number of species and populations included in analyses.**

| Scale | Analysis | Number of species | Number of populations | Number of species with >3 populations |
|---|---|---|---|---|
| **UK** | Geographic range | 167 | 381 | 32 |
| | Population size | 112 | 253 | 19 |
| | Habitat specificity | 144 | 289 | 29 |
| | Red List status | 153 | 364 | 31 |
| **Global** | System | 2074 | 9284 | 834 |
| | Biome | 2074 | 9284 | 834 |
| | Taxa | | | |
| | - *Actinopterygii* | 544 | 1626 | 151 |
| | - *Amphibia* | 78 | 193 | 21 |
| | - *Aves* | 968 | 5852 | 538 |
| | - *Mammalia* | 306 | 1158 | 94 |
| | - *Reptilia* | 110 | 322 | 16 |
| | Red List status | 1702 | 8064 | 708 |
| | Latitude | 2074 | 9284 | 834 |
| | Duration | 2074 | 9284 | 834 |

**Figure S1. Distribution of monitoring duration across (a) freshwater, (b) marine and (c) terrestrial systems. Results include 9284 populations from 2084 species.**

**Figure S2. Distribution of monitoring duration across all monitored vertebrate populations (plot a), bony fish (plot b), amphibians (plot c), birds (plot d), mammals (plot e) and reptiles (plot f).** See Table S1 for sample sizes of each group.

Habitat specificity (UK)

**Figure S3. Effect of habitat specificity (calculated by surveying the 'Habitat and Ecology' profile for each species on the IUCN Red List website) on vertebrate population trends in the UK.** The *mu* values of population change are derived from state-space model fits of changes in abundance over the monitoring duration for each population. Error bars show confidence intervals around population trend (mu). See Table S2 for full model outputs.

915 **Figure S4. Correlation matrix between rarity metrix on (a) UK and (b) global scales.** The three rarity metrics were weakly correlated on both study scales.

**Figure S5. Population change (*mu* values from state-space models) across the**

**five most commonly used units in the subset of the LPD we analyzed.** Numbers

on the x axis correspond to sample size in each category.

**Figure S6. Geographic distribution of vertebrate population change in the UK,**

**from state-space models (plot a) and linear models (plot b).** Results include 508

populations from 237 species in the UK.

**Figure. S7. Distribution of population change (*mu*) values from state-space models of vertebrate abundance over time across systems (plot a) and across taxa (plot b).** See Table S1 for sample sizes.

**Figure. S8. Distribution of posterior means (mean distributions across ten models for each taxa for each type of population change using ten random trees to account for phylogenetic uncertainty) for phylogenetic, species and residual variance across taxa.** Phylogeny effects were calculated based on a branch length covariance matrix. See Table S2 for full model outputs.

**Figure S9. Vertebrate population trends across monitoring duration.** Results

are for 9284 populations from 2084 species. Population trends (*mu*) were estimated

for all populations monitored for more than five time points using state-space models

(plot a) and linear models (plot b). Population fluctuations (plot c) were estimated

using the confidence interval (half of the 95% confidence interval) of the *mu* values

(see also Figure 5). Error bars show 95% confidence intervals.

**Figure S10. Geographic distribution of population change, calculated using**
**state-space models (plot a), linear models (plot b), and using the confidence**
**interval (half of the 95% confidence interval) of the *mu* values (plot c).** Results
include 9284 populations from 2084 species. Larger confidence intervals on (c)
correspond to populations that have experienced more fluctuations.

**Table S2. Model outputs from UK and global analyses.** Sigma is the overall

model residual variance. Net population change is estimated using *mu* values

derived from state-space models of population abundance versus time and slopes of

linear models of population abundance versus time. The weighted *mu* models

included *mu* as a response variable, weighted by *tau*, the observation error estimate

derived from the state-space models. The fluctuation models included half of the

95% confidence interval around the *mu* value of population change, the standard

error around the slopes of linear models of population abundance versus time, and

the standard deviation of the raw time-series data for each population. Larger

confidence intervals correspond to populations that experienced more fluctuations.

The weighted fluctuation models were weighted by *tau*, the observation error

estimate derived from the state-space models. To account for phylogenetic

uncertainty, we ran the phylogenetic models for amphibian, bird and reptile species

using 10 different trees for each class, and here we present the mean, min and max

values from the different model runs.

| Model name | Variable | Posterior mean | Lower 95% CI | Upper 95% CI | Effective sample size | pMCMC | Effect |
|---|---|---|---|---|---|---|---|
| Geographic range (UK) - mu | (Intercept) | -0.001 | -0.04 | 0.03 | 18,000 | 0.94 | fixed |
| | log(km2_range) | 0.001 | -0.001 | 0.004 | 18,000 | 0.41 | fixed |
| | sigma | 0.01 | 0.005 | 0.01 | 18,000 | | residual |
| Geographic range (UK) - weighted | (Intercept) | -0.01 | -0.05 | 0.03 | 17,509 | 0.58 | fixed |
| | log.range | 0.002 | -0.001 | 0.005 | 17,525 | 0.16 | fixed |
| | sigma | 0.005 | 0.004 | 0.01 | 18,435 | | residual |
| Geographic range (UK) - slope | (Intercept) | -0.01 | -0.07 | 0.06 | 18,000 | 0.85 | fixed |
| | log.range | 0.002 | -0.002 | 0.01 | 17,583 | 0.38 | fixed |
| | sigma | 0.02 | 0.02 | 0.02 | 18,000 | | residual |
| Geographic range (UK) - fluctuations CI | (Intercept) | 0.03 | -0.04 | 0.10 | 18,000 | 0.41 | fixed |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| | log.range | 0.01 | 0.003 | 0.01 | 18,000 | 0.002 | fixed |
| | sigma | 0.02 | 0.02 | 0.02 | 18,000 | | residual |
| Geographic range (UK) - fluctuations CI weighted | (Intercept) | 0.02 | -0.06 | 0.10 | 18,000 | 0.59 | fixed |
| | log.range | 0.01 | 0.003 | 0.01 | 18,000 | 0.002 | fixed |
| | sigma | 0.02 | 0.02 | 0.03 | 18,000 | | residual |
| Geographic range (UK) - fluctuations SE | (Intercept) | -0.02 | -0.09 | 0.06 | 18,000 | 0.67 | fixed |
| | log.range | 0.01 | 0.005 | 0.02 | 18,000 | 0.0002 | fixed |
| | sigma | 0.02 | 0.02 | 0.03 | 18,000 | | residual |
| Geographic range (UK) - fluctuations SD | (Intercept) | 0.15 | 0.08 | 0.23 | 17,410 | 0.0002 | fixed |
| | log.range | 0.02 | 0.01 | 0.02 | 18,056 | 0.0001 | fixed |
| | sigma | 0.03 | 0.02 | 0.03 | 18,000 | | residual |
| Mean population size (UK) - mu | (Intercept) | 0.003 | -0.02 | 0.03 | 18,710 | 0.79 | fixed |
| | log.meanpop | 0.001 | -0.003 | 0.005 | 21,192 | 0.67 | fixed |
| | sigma | 0.005 | 0.004 | 0.01 | 18,000 | | residual |
| Mean population size (UK) - weighted | (Intercept) | 0.003 | -0.02 | 0.02 | 18,000 | 0.80 | fixed |
| | log.meanpop | 0.001 | -0.003 | 0.005 | 18,000 | 0.67 | fixed |
| | sigma | 0.005 | 0.004 | 0.01 | 18,000 | | residual |
| Mean population size (UK) - slope | (Intercept) | 0.04 | -0.01 | 0.08 | 18,581 | 0.11 | fixed |
| | log(meanpop) | -0.001 | -0.01 | 0.01 | 18,717 | 0.74 | fixed |
| | sigma | 0.02 | 0.02 | 0.03 | 18,000 | | residual |
| Mean population size (UK) - fluctuations CI | (Intercept) | 0.30 | 0.24 | 0.36 | 17,427 | 0.0001 | fixed |
| | log.meanpop | -0.01 | -0.02 | 0.0004 | 18,000 | 0.05 | fixed |
| | sigma | 0.03 | 0.03 | 0.04 | 18,000 | | residual |
| Mean population size (UK) - fluctuations CI weighted | (Intercept) | 0.30 | 0.24 | 0.36 | 17,570 | 0.0001 | fixed |
| | log.meanpop | -0.01 | -0.02 | 0.0001 | 18,000 | 0.05 | fixed |
| | sigma | 0.03 | 0.03 | 0.04 | 18,000 | | residual |
| Mean population size (UK) - fluctuations SE | (Intercept) | 0.21 | 0.17 | 0.25 | 18,000 | 0.0001 | fixed |
| | log.meanpop | -0.01 | -0.02 | -0.01 | 18,000 | 0.0001 | fixed |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| | sigma | 0.02 | 0.01 | 0.02 | 18,000 | | residual |
| Mean population size (UK) - fluctuations SD | (Intercept) | 0.50 | 0.44 | 0.56 | 18,000 | 0.0001 | fixed |
| | log.meanpop | -0.01 | -0.03 | -0.005 | 18,000 | 0.01 | fixed |
| | sigma | 0.03 | 0.02 | 0.04 | 18,403 | | residual |
| Habitat specificity (UK) - mu | (Intercept) | 0.01 | -0.01 | 0.02 | 18,000 | 0.38 | fixed |
| | hab_spec | 0.001 | -0.001 | 0.002 | 18,000 | 0.28 | fixed |
| | sigma | 0.01 | 0.005 | 0.01 | 16,892 | | residual |
| Habitat specificity (UK) - weighted | (Intercept) | 0.01 | -0.01 | 0.02 | 18,000 | 0.40 | fixed |
| | hab_spec | 0.001 | -0.001 | 0.003 | 18,000 | 0.23 | fixed |
| | sigma | 0.005 | 0.004 | 0.01 | 18,727 | | residual |
| Habitat specificity (UK) - slope | (Intercept) | 0.01 | -0.01 | 0.04 | 20,246 | 0.33 | fixed |
| | hab_spec | 0.001 | -0.001 | 0.003 | 18,000 | 0.44 | fixed |
| | sigma | 0.02 | 0.02 | 0.02 | 18,436 | | residual |
| Habita specificity (UK) - profiling method | (Intercept) | 0.01 | -0.003 | 0.03 | 18,000 | 0.11 | fixed |
| | hab_spec | -0.0003 | -0.002 | 0.002 | 18,000 | 0.76 | fixed |
| | sigma | 0.01 | 0.01 | 0.01 | 18,000 | | residual |
| Habitat specificity (UK) - fluctuations CI | (Intercept) | 0.11 | 0.09 | 0.14 | 18,000 | 0.0001 | fixed |
| | hab_spec | 0.003 | 0.0003 | 0.01 | 18,000 | 0.03 | fixed |
| | sigma | 0.02 | 0.02 | 0.02 | 18,000 | | residual |
| Habitat specificity (UK) - fluctuations CI weighted | (Intercept) | 0.11 | 0.08 | 0.15 | 18,000 | 0.0001 | fixed |
| | hab_spec | 0.004 | 0.0005 | 0.01 | 18,000 | 0.02 | fixed |
| | sigma | 0.02 | 0.02 | 0.03 | 18,000 | | residual |
| Habitat specificity (UK) - fluctuations SE | (Intercept) | 0.11 | 0.08 | 0.14 | 18,000 | 0.0001 | fixed |
| | hab_spec | 0.002 | -0.001 | 0.004 | 18,000 | 0.28 | fixed |
| | sigma | 0.02 | 0.02 | 0.03 | 18,000 | | residual |
| Habitat specificity (UK) - fluctuations SD | (Intercept) | 0.37 | 0.34 | 0.40 | 18,000 | 0.0001 | fixed |
| | hab_spec | 0.001 | -0.001 | 0.004 | 18,000 | 0.32 | fixed |
| | sigma | 0.03 | 0.02 | 0.03 | 18,000 | | residual |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| Red List status (UK) - mu | Least concern | 0.01 | 0.004 | 0.02 | 18,000 | 0.001 | fixed |
| | Near threatened | 0.03 | 0.005 | 0.06 | 16,876 | 0.02 | fixed |
| | Vulnerable | 0.002 | -0.02 | 0.03 | 18,000 | 0.86 | fixed |
| | Endangered | -0.04 | -0.18 | 0.11 | 18,000 | 0.60 | fixed |
| | Critically endangered | 0.06 | -0.02 | 0.15 | 18,000 | 0.15 | fixed |
| | sigma | 0.01 | 0.005 | 0.01 | 18,000 | | residual |
| Red List status (UK) - weighted | Least concern | 0.02 | 0.01 | 0.02 | 18,000 | 0.0001 | fixed |
| | Near threatened | 0.03 | 0.002 | 0.07 | 18,416 | 0.04 | fixed |
| | Vulnerable | -0.01 | -0.04 | 0.03 | 18,000 | 0.73 | fixed |
| | Endangered | -0.04 | -0.17 | 0.10 | 18,000 | 0.59 | fixed |
| | Critically endangered | 0.04 | -0.18 | 0.28 | 18,000 | 0.72 | fixed |
| | sigma | 0.005 | 0.004 | 0.01 | 18,398 | | residual |
| Red List status (UK) - slope | Least concern | 0.02 | 0.01 | 0.03 | 18,827 | 0.002 | fixed |
| | Near threatened | 0.04 | -0.01 | 0.09 | 18,000 | 0.10 | fixed |
| | Vulnerable | -0.001 | -0.05 | 0.05 | 18,000 | 0.96 | fixed |
| | Endangered | -0.01 | -0.28 | 0.25 | 18,000 | 0.93 | fixed |
| | Critically endangered | 0.05 | -0.10 | 0.21 | 18,000 | 0.48 | fixed |
| | sigma | 0.02 | 0.02 | 0.02 | 18,000 | | residual |
| Red List status (UK) - fluctuations CI | Least concern | 0.14 | 0.13 | 0.16 | 18,000 | 0.0001 | fixed |
| | Near threatened | 0.14 | 0.09 | 0.19 | 18,000 | 0.0001 | fixed |
| | Vulnerable | 0.10 | 0.05 | 0.16 | 18,000 | 0.0002 | fixed |
| | Endangered | 0.24 | -0.05 | 0.52 | 18,000 | 0.10 | fixed |
| | Critically endangered | 0.09 | -0.07 | 0.26 | 17,435 | 0.30 | fixed |
| | sigma | 0.02 | 0.02 | 0.02 | 18,000 | | residual |
| Red List status (UK) - fluctuations CI weighted | Least concern | 0.15 | 0.13 | 0.17 | 17,566 | 0.0001 | fixed |
| | Near threatened | 0.16 | 0.09 | 0.22 | 18,000 | 0.0001 | fixed |
| | Vulnerable | 0.11 | 0.04 | 0.17 | 17,582 | 0.001 | fixed |
| | Endangered | 0.24 | -0.06 | 0.54 | 18,000 | 0.11 | fixed |
| | Critically endangered | 0.06 | -0.23 | 0.34 | 18,000 | 0.68 | fixed |
| | sigma | 0.02 | 0.02 | 0.03 | 17,700 | | residual |
| Red List status (UK) - fluctuations SE | Least concern | 0.13 | 0.11 | 0.14 | 18,000 | 0.0001 | fixed |
| | Near threatened | 0.13 | 0.08 | 0.18 | 18,600 | 0.0001 | fixed |
| | Vulnerable | 0.07 | 0.01 | 0.13 | 18,809 | 0.02 | fixed |
| | Endangered | 0.07 | -0.23 | 0.37 | 19,204 | 0.65 | fixed |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| | Critically endangered | 0.07 | -0.11 | 0.24 | 18,000 | 0.45 | fixed |
| | sigma | 0.02 | 0.02 | 0.03 | 18,000 | | residual |
| Red List status (UK) - fluctuations SD | Least concern | 0.33 | 0.33 | 0.34 | 18,000 | 0.0001 | fixed |
| | Near threatened | 0.40 | 0.37 | 0.43 | 20,507 | 0.0001 | fixed |
| | Vulnerable | 0.23 | 0.21 | 0.25 | 18,000 | 0.0001 | fixed |
| | Endangered | 0.24 | -0.02 | 0.52 | 18,000 | 0.08 | fixed |
| | Critically endangered | 0.28 | 0.16 | 0.40 | 18,000 | 0.0001 | fixed |
| | sigma | 0.02 | 0.02 | 0.02 | 18,000 | | residual |
| System (UK) - mu | Freshwater | 0.02 | 0.001 | 0.03 | 18,000 | 0.04 | fixed |
| | Marine | 0.01 | -0.001 | 0.02 | 18,905 | 0.07 | fixed |
| | Terrestrial | 0.02 | 0.004 | 0.03 | 18,135 | 0.01 | fixed |
| | sigma | 0.01 | 0.004 | 0.01 | 18,000 | | residual |
| System (UK) - weighted | Freshwater | 0.02 | -0.004 | 0.03 | 21,467 | 0.10 | fixed |
| | Marine | 0.01 | 0.0004 | 0.02 | 17,586 | 0.04 | fixed |
| | Terrestrial | 0.02 | 0.01 | 0.03 | 18,424 | 0.005 | fixed |
| | sigma | 0.005 | 0.004 | 0.01 | 18,000 | | residual |
| System (UK) - slope | Freshwater | 0.03 | 0.004 | 0.06 | 18,000 | 0.02 | fixed |
| | Marine | 0.01 | -0.005 | 0.03 | 18,064 | 0.20 | fixed |
| | Terrestrial | 0.03 | 0.01 | 0.05 | 18,705 | 0.01 | fixed |
| | sigma | 0.02 | 0.01 | 0.02 | 18,000 | | residual |
| Taxa (UK) - mu | Actinopterygii | -0.004 | -0.02 | 0.01 | 18,000 | 0.51 | fixed |
| | Amphibia | -0.002 | -0.04 | 0.03 | 18,000 | 0.90 | fixed |
| | Aves | 0.01 | 0.002 | 0.02 | 18,000 | 0.02 | fixed |
| | Mammalia | 0.03 | 0.02 | 0.05 | 18,000 | 0.0001 | fixed |
| | Reptilia | 0.04 | -0.02 | 0.10 | 18,000 | 0.16 | fixed |
| | sigma | 0.005 | 0.004 | 0.01 | 18,000 | | residual |
| Taxa (UK) - weighted | Actinopterygii | -0.01 | -0.02 | 0.01 | 18,000 | 0.29 | fixed |
| | Amphibia | -0.002 | -0.05 | 0.04 | 17,451 | 0.92 | fixed |
| | Aves | 0.02 | 0.01 | 0.03 | 18,000 | 0.003 | fixed |
| | Mammalia | 0.04 | 0.02 | 0.05 | 18,000 | 0.0001 | fixed |
| | Reptilia | 0.03 | -0.04 | 0.10 | 17,161 | 0.41 | fixed |
| | sigma | 0.004 | 0.004 | 0.01 | 18,000 | | residual |
| Taxa (UK) - slope | Actinopterygii | -0.01 | -0.04 | 0.01 | 18,000 | 0.41 | fixed |
| | Amphibia | -0.003 | -0.07 | 0.06 | 18,000 | 0.92 | fixed |
| | Aves | 0.02 | 0.0001 | 0.03 | 18,468 | 0.05 | fixed |
| | Mammalia | 0.06 | 0.04 | 0.09 | 18,000 | 0.0001 | fixed |
| | Reptilia | 0.07 | -0.05 | 0.18 | 18,000 | 0.23 | fixed |
| | sigma | 0.02 | 0.01 | 0.02 | 18,000 | | residual |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| System (global) - mu | Freshwater | 0.005 | 0.002 | 0.01 | 19,286 | 0.0001 | fixed |
| | Marine | 0.004 | 0.002 | 0.01 | 18,820 | 0.0003 | fixed |
| | Terrestrial | 0.003 | 0.001 | 0.005 | 18,000 | 0.001 | fixed |
| | sigma | 0.003 | 0.003 | 0.004 | 18,000 | | residual |
| System (global) - weighted | Freshwater | 0.01 | 0.002 | 0.01 | 18,000 | 0.001 | fixed |
| | Marine | 0.004 | 0.001 | 0.01 | 18,000 | 0.01 | fixed |
| | Terrestrial | 0.004 | 0.002 | 0.01 | 17,492 | 0.001 | fixed |
| | sigma | 0.004 | 0.004 | 0.004 | 19,112 | | residual |
| System (global) - slope | Freshwater | 0.01 | 0.01 | 0.01 | 18,000 | 0.0001 | fixed |
| | Marine | 0.004 | 0.0002 | 0.01 | 18,000 | 0.04 | fixed |
| | Terrestrial | 0.004 | 0.001 | 0.01 | 18,000 | 0.01 | fixed |
| | sigma | 0.01 | 0.01 | 0.01 | 18,000 | | residual |
| System (global) - fluctuations CI | Freshwater | 0.13 | 0.13 | 0.14 | 18,000 | 0.0001 | fixed |
| | Marine | 0.13 | 0.13 | 0.14 | 18,000 | 0.0001 | fixed |
| | Terrestrial | 0.10 | 0.10 | 0.10 | 18,000 | 0.0001 | fixed |
| | sigma | 0.02 | 0.02 | 0.02 | 18,000 | | residual |
| System (global) - fluctuations CI weighted | Freshwater | 0.15 | 0.14 | 0.16 | 18,000 | 0.0001 | fixed |
| | Marine | 0.15 | 0.14 | 0.16 | 18,000 | 0.0001 | fixed |
| | Terrestrial | 0.11 | 0.10 | 0.11 | 18,000 | 0.0001 | fixed |
| | sigma | 0.02 | 0.02 | 0.02 | 18,000 | | residual |
| System (global) - fluctuations SE | Freshwater | 0.12 | 0.12 | 0.13 | 18,000 | 0.0001 | fixed |
| | Marine | 0.12 | 0.11 | 0.12 | 18,000 | 0.0001 | fixed |
| | Terrestrial | 0.09 | 0.09 | 0.09 | 18,000 | 0.0001 | fixed |
| | sigma | 0.02 | 0.02 | 0.02 | 16,149 | | residual |
| System (global) - fluctuations SD | Freshwater | 0.55 | 0.55 | 0.56 | 18,943 | 0.0001 | fixed |
| | Marine | 0.56 | 0.56 | 0.56 | 18,000 | 0.0001 | fixed |
| | Terrestrial | 0.54 | 0.54 | 0.54 | 18,000 | 0.0001 | fixed |
| | sigma | 0.01 | 0.01 | 0.01 | 18,000 | | residual |
| Biome (global) - mu | Boreal forests/taiga | -0.001 | -0.004 | 0.002 | 17,462 | 0.70 | fixed |
| | Deserts and xeric shrublands | -0.01 | -0.03 | 0.002 | 18,000 | 0.09 | fixed |
| | Trop. and subtrop. grasslands savannas and shrublands | -0.01 | -0.02 | -0.001 | 18,000 | 0.03 | fixed |
| | Large lakes | 0.003 | -0.004 | 0.01 | 18,000 | 0.42 | fixed |
| | Mediterranean forests woodlands and scrub | 0.02 | 0.01 | 0.02 | 18,000 | 0.0002 | fixed |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| | Montane freshwaters | 0.01 | -0.01 | 0.04 | 18,000 | 0.34 | fixed |
| | Montane grasslands and shrublands | 0.02 | 0.001 | 0.04 | 18,000 | 0.03 | fixed |
| | Polar freshwaters | 0.01 | 0.001 | 0.01 | 18,000 | 0.02 | fixed |
| | Polar seas | -0.002 | -0.02 | 0.02 | 18,000 | 0.83 | fixed |
| | Temperate forests | 0.005 | 0.002 | 0.01 | 18,000 | 0.0001 | fixed |
| | Temperate wetlands and rivers | 0.005 | 0.002 | 0.01 | 18,000 | 0.001 | fixed |
| | Temperate grasslands savannas and shrublands | -0.004 | -0.01 | 0.002 | 18,000 | 0.22 | fixed |
| | Tropical wetlands and rivers | 0.01 | -0.003 | 0.01 | 18,000 | 0.21 | fixed |
| | Tropical and subtropical forests | 0.02 | 0.01 | 0.03 | 18,000 | 0.0001 | fixed |
| | Tropical coral | 0.02 | 0.01 | 0.04 | 18,000 | 0.0001 | fixed |
| | Tundra | 0.01 | -0.001 | 0.01 | 18,000 | 0.09 | fixed |
| | Xeric freshwaters and endorheic basins | -0.002 | -0.02 | 0.01 | 15,937 | 0.77 | fixed |
| | sigma | 0.003 | 0.003 | 0.003 | 17,444 | | residual |
| Biome (global) - slope | Boreal forests/taiga | -0.001 | -0.01 | 0.003 | 18,000 | 0.55 | fixed |
| | Deserts and xeric shrublands | -0.04 | -0.06 | -0.02 | 17,811 | 0.001 | fixed |
| | Trop. and subtrop. grasslands savannas and shrublands | -0.01 | -0.02 | -0.002 | 18,000 | 0.02 | fixed |
| | Large lakes | 0.004 | -0.01 | 0.02 | 18,000 | 0.50 | fixed |
| | Mediterranean forests woodlands and scrub | 0.03 | 0.01 | 0.04 | 18,000 | 0.0001 | fixed |
| | Montane freshwaters | 0.03 | -0.01 | 0.07 | 18,000 | 0.18 | fixed |
| | Montane grasslands and shrublands | 0.02 | -0.004 | 0.05 | 18,000 | 0.10 | fixed |
| | Polar freshwaters | 0.01 | -0.004 | 0.02 | 18,000 | 0.22 | fixed |
| | Polar seas | -0.01 | -0.04 | 0.02 | 18,000 | 0.50 | fixed |
| | Temperate forests | 0.01 | 0.004 | 0.01 | 18,000 | 0.0001 | fixed |
| | Temperate wetlands and rivers | 0.01 | 0.002 | 0.01 | 18,000 | 0.004 | fixed |
| | Temperate grasslands savannas and shrublands | -0.004 | -0.01 | 0.01 | 18,000 | 0.42 | fixed |
| | Tropical wetlands and rivers | 0.003 | -0.01 | 0.02 | 18,000 | 0.64 | fixed |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| | Tropical and subtropical forests | 0.02 | 0.01 | 0.04 | 17,621 | 0.0001 | fixed |
| | Tropical coral | 0.04 | 0.02 | 0.05 | 18,960 | 0.0001 | fixed |
| | Tundra | 0.01 | 0.0001 | 0.03 | 18,000 | 0.04 | fixed |
| | Xeric freshwaters and endorheic basins | 0.01 | -0.02 | 0.03 | 18,000 | 0.60 | fixed |
| | sigma | 0.004 | 0.003 | 0.004 | 18,000 | | residual |
| Biome (global) - weighted | Boreal forests/taiga | -0.002 | -0.01 | 0.003 | 17,517 | 0.42 | fixed |
| | Deserts and xeric shrublands | -0.03 | -0.05 | 0.003 | 18,000 | 0.08 | fixed |
| | Trop. and subtrop. grasslands savannas and shrublands | -0.002 | -0.02 | 0.01 | 18,000 | 0.71 | fixed |
| | Large lakes | 0.01 | -0.01 | 0.02 | 18,000 | 0.29 | fixed |
| | Mediterranean forests woodlands and scrub | 0.02 | 0.005 | 0.03 | 19,064 | 0.01 | fixed |
| | Montane freshwaters | 0.01 | -0.03 | 0.05 | 18,469 | 0.65 | fixed |
| | Montane grasslands and shrublands | 0.02 | -0.01 | 0.06 | 18,000 | 0.12 | fixed |
| | Polar freshwaters | 0.02 | 0.01 | 0.03 | 18,630 | 0.002 | fixed |
| | Polar seas | -0.01 | -0.05 | 0.02 | 17,320 | 0.44 | fixed |
| | Temperate forests | 0.01 | 0.004 | 0.01 | 17,516 | 0.0002 | fixed |
| | Temperate wetlands and rivers | 0.01 | 0.01 | 0.01 | 17,324 | 0.0001 | fixed |
| | Temperate grasslands savannas and shrublands | -0.01 | -0.02 | 0.002 | 17,573 | 0.14 | fixed |
| | Tropical wetlands and rivers | 0.01 | -0.004 | 0.02 | 18,000 | 0.13 | fixed |
| | Tropical and subtropical forests | 0.01 | -0.001 | 0.03 | 18,000 | 0.07 | fixed |
| | Tropical coral | 0.04 | 0.02 | 0.06 | 17,511.66 | 0.0001 | fixed |
| | Tundra | 0.01 | -0.004 | 0.02 | 18,000 | 0.16 | fixed |
| | Xeric freshwaters and endorheic basins | -0.03 | -0.05 | -0.0004 | 18,000 | 0.04 | fixed |
| | sigma | 0.01 | 0.01 | 0.01 | 18,000 | | residual |
| Biome (global) - fluctuations CI | Boreal forests/taiga | 0.07 | 0.07 | 0.08 | 18,000 | 0.0001 | fixed |
| | Deserts and xeric shrublands | 0.22 | 0.18 | 0.26 | 18,658 | 0.0001 | fixed |
| | Trop. and subtrop. grasslands savannas and shrublands | 0.21 | 0.19 | 0.23 | 18,000 | 0.0001 | fixed |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| | Large lakes | 0.11 | 0.09 | 0.12 | 18,000 | 0.0001 | fixed |
| | Mediterranean forests woodlands and scrub | 0.15 | 0.13 | 0.16 | 17,403 | 0.0001 | fixed |
| | Montane freshwaters | 0.14 | 0.09 | 0.19 | 18,000 | 0.0001 | fixed |
| | Montane grasslands and shrublands | 0.17 | 0.13 | 0.21 | 18,000 | 0.0001 | fixed |
| | Polar freshwaters | 0.14 | 0.13 | 0.15 | 18,000 | 0.0001 | fixed |
| | Polar seas | 0.16 | 0.11 | 0.21 | 18,148 | 0.0001 | fixed |
| | Temperate forests | 0.09 | 0.09 | 0.10 | 18,000 | 0.0001 | fixed |
| | Temperate wetlands and rivers | 0.13 | 0.12 | 0.13 | 18,000 | 0.0001 | fixed |
| | Temperate grasslands savannas and shrublands | 0.08 | 0.07 | 0.09 | 18,000 | 0.0001 | fixed |
| | Tropical wetlands and rivers | 0.20 | 0.19 | 0.22 | 18,440 | 0.0001 | fixed |
| | Tropical and subtropical forests | 0.20 | 0.18 | 0.21 | 18,000 | 0.0001 | fixed |
| | Tropical coral | 0.16 | 0.14 | 0.19 | 17,964 | 0.0001 | fixed |
| | Tundra | 0.11 | 0.09 | 0.13 | 18,000 | 0.0001 | fixed |
| | Xeric freshwaters and endorheic basins | 0.18 | 0.14 | 0.21 | 18,000 | 0.0001 | fixed |
| | sigma | 0.02 | 0.01 | 0.02 | 18,000 | | residual |
| Biome (global) - fluctuations CI weighted | Boreal forests/taiga | 0.08 | 0.07 | 0.08 | 18,000 | 0.0001 | fixed |
| | Deserts and xeric shrublands | 0.26 | 0.21 | 0.30 | 18,000 | 0.0001 | fixed |
| | Trop. and subtrop. grasslands savannas and shrublands | 0.25 | 0.23 | 0.27 | 17,477 | 0.0001 | fixed |
| | Large lakes | 0.12 | 0.09 | 0.14 | 18,000 | 0.0001 | fixed |
| | Mediterranean forests woodlands and scrub | 0.16 | 0.14 | 0.19 | 18,000 | 0.0001 | fixed |
| | Montane freshwaters | 0.23 | 0.15 | 0.31 | 17,217 | 0.0001 | fixed |
| | Montane grasslands and shrublands | 0.18 | 0.12 | 0.23 | 18,681 | 0.0001 | fixed |
| | Polar freshwaters | 0.17 | 0.15 | 0.19 | 17,751 | 0.0001 | fixed |
| | Polar seas | 0.18 | 0.12 | 0.23 | 18,000 | 0.0001 | fixed |
| | Temperate forests | 0.09 | 0.09 | 0.10 | 18,000 | 0.0001 | fixed |
| | Temperate wetlands and rivers | 0.14 | 0.13 | 0.15 | 18,000 | 0.0001 | fixed |
| | Temperate grasslands | 0.08 | 0.06 | 0.10 | 18,000 | 0.0001 | fixed |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| | savannas and shrublands | | | | | | |
| | Tropical wetlands and rivers | 0.22 | 0.20 | 0.25 | 18,000 | 0.0001 | fixed |
| | Tropical and subtropical forests | 0.21 | 0.19 | 0.24 | 18,000 | 0.0001 | fixed |
| | Tropical coral | 0.18 | 0.15 | 0.21 | 18,000 | 0.0001 | fixed |
| | Tundra | 0.11 | 0.09 | 0.14 | 18,000 | 0.0001 | fixed |
| | Xeric freshwaters and endorheic basins | 0.23 | 0.18 | 0.27 | 18,000 | 0.0001 | fixed |
| | sigma | 0.02 | 0.02 | 0.02 | 18,000 | | residual |
| Biome (global) - fluctuations SE | Boreal forests/taiga | 0.06 | 0.06 | 0.07 | 18,000 | 0.0001 | fixed |
| | Deserts and xeric shrublands | 0.21 | 0.17 | 0.25 | 18,000 | 0.0001 | fixed |
| | Trop. and subtrop. grasslands savannas and shrublands | 0.11 | 0.09 | 0.13 | 17,497 | 0.0001 | fixed |
| | Large lakes | 0.09 | 0.07 | 0.11 | 18,000 | 0.0001 | fixed |
| | Mediterranean forests woodlands and scrub | 0.16 | 0.14 | 0.18 | 18,000 | 0.0001 | fixed |
| | Montane freshwaters | 0.15 | 0.10 | 0.21 | 16,658 | 0.0001 | fixed |
| | Montane grasslands and shrublands | 0.11 | 0.07 | 0.15 | 18,000 | 0.0001 | fixed |
| | Polar freshwaters | 0.14 | 0.13 | 0.15 | 18,000 | 0.0001 | fixed |
| | Polar seas | 0.08 | 0.03 | 0.13 | 18,000 | 0.002 | fixed |
| | Temperate forests | 0.09 | 0.08 | 0.09 | 18,000 | 0.0001 | fixed |
| | Temperate wetlands and rivers | 0.11 | 0.11 | 0.12 | 18,000 | 0.0001 | fixed |
| | Temperate grasslands savannas and shrublands | 0.07 | 0.06 | 0.09 | 18,000 | 0.0001 | fixed |
| | Tropical wetlands and rivers | 0.21 | 0.19 | 0.23 | 18,000 | 0.0001 | fixed |
| | Tropical and subtropical forests | 0.15 | 0.13 | 0.17 | 18,000 | 0.0001 | fixed |
| | Tropical coral | 0.18 | 0.16 | 0.21 | 18,000 | 0.0001 | fixed |
| | Tundra | 0.13 | 0.11 | 0.14 | 17,343 | 0.0001 | fixed |
| | Xeric freshwaters and endorheic basins | 0.13 | 0.09 | 0.17 | 18,000 | 0.0001 | fixed |
| | sigma | 0.02 | 0.02 | 0.02 | 18,000 | | residual |
| Biome (global) - fluctuations SD | Boreal forests/taiga | 0.51 | 0.51 | 0.52 | 18,000 | 0.0001 | fixed |
| | Deserts and xeric shrublands | 0.60 | 0.57 | 0.62 | 17,053.04 | 0.0001 | fixed |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| | Trop. and subtrop. grasslands savannas and shrublands | 0.61 | 0.60 | 0.62 | 18,000 | 0.0001 | fixed |
| | Large lakes | 0.54 | 0.52 | 0.55 | 18,000 | 0.0001 | fixed |
| | Mediterranean forests woodlands and scrub | 0.59 | 0.58 | 0.61 | 17,567 | 0.0001 | fixed |
| | Montane freshwaters | 0.58 | 0.54 | 0.62 | 18,000 | 0.0001 | fixed |
| | Montane grasslands and shrublands | 0.59 | 0.56 | 0.62 | 18,000 | 0.0001 | fixed |
| | Polar freshwaters | 0.55 | 0.54 | 0.56 | 18,000 | 0.0001 | fixed |
| | Polar seas | 0.55 | 0.52 | 0.59 | 18,000 | 0.0001 | fixed |
| | Temperate forests | 0.53 | 0.53 | 0.54 | 18,056 | 0.0001 | fixed |
| | Temperate wetlands and rivers | 0.55 | 0.55 | 0.56 | 18,000 | 0.0001 | fixed |
| | Temperate grasslands savannas and shrublands | 0.52 | 0.51 | 0.53 | 17,393 | 0.0001 | fixed |
| | Tropical wetlands and rivers | 0.60 | 0.59 | 0.62 | 18,000 | 0.0001 | fixed |
| | Tropical and subtropical forests | 0.62 | 0.60 | 0.63 | 18,000 | 0.0001 | fixed |
| | Tropical coral | 0.61 | 0.59 | 0.63 | 18,000 | 0.0001 | fixed |
| | Tundra | 0.54 | 0.53 | 0.56 | 18,000 | 0.0001 | fixed |
| | Xeric freshwaters and endorheic basins | 0.56 | 0.53 | 0.58 | 18,000 | 0.0001 | fixed |
| | sigma | 0.01 | 0.01 | 0.01 | 18,000 | | residual |
| Taxa (global) - mu | Actinopterygii | -0.001 | -0.003 | 0.002 | 18,000 | 0.67 | fixed |
| | Amphibia | -0.01 | -0.02 | -0.005 | 18,000 | 0.003 | fixed |
| | Aves | 0.004 | 0.003 | 0.01 | 18,000 | 0.0001 | fixed |
| | Mammalia | 0.01 | 0.01 | 0.01 | 18,000 | 0.0001 | fixed |
| | Reptilia | 0.02 | 0.01 | 0.02 | 18,000 | 0.0001 | fixed |
| | sigma | 0.003 | 0.003 | 0.004 | 18,000 | | residual |
| Taxa (global) - weighted | Actinopterygii | -0.001 | -0.01 | 0.003 | 18,000 | 0.68 | fixed |
| | Amphibia | -0.01 | -0.03 | -0.002 | 18,000 | 0.03 | fixed |
| | Aves | 0.01 | 0.003 | 0.01 | 18,000 | 0.0001 | fixed |
| | Mammalia | 0.01 | 0.01 | 0.02 | 18,000 | 0.0001 | fixed |
| | Reptilia | 0.01 | -0.001 | 0.02 | 18,000 | 0.07 | fixed |
| | sigma | 0.004 | 0.004 | 0.004 | 18,000 | | residual |
| Taxa (global) - slope | Actinopterygii | -0.002 | -0.01 | 0.002 | 18,000 | 0.34 | fixed |
| | Amphibia | -0.02 | -0.04 | -0.01 | 18,000 | 0.002 | fixed |
| | Aves | 0.01 | 0.004 | 0.01 | 17,484 | 0.0001 | fixed |
| | Mammalia | 0.01 | 0.01 | 0.02 | 16,803 | 0.0001 | fixed |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| | Reptilia | 0.03 | 0.02 | 0.04 | 17,631 | 0.0001 | fixed |
| | sigma | 0.01 | 0.01 | 0.01 | 18,000 | | residual |
| Taxa (global) - fluctuations CI | Actinopterygii | 0.15 | 0.14 | 0.16 | 18,000 | 0.0001 | fixed |
| | Amphibia | 0.22 | 0.20 | 0.24 | 17,663 | 0.0001 | fixed |
| | Aves | 0.09 | 0.09 | 0.10 | 18,000 | 0.0001 | fixed |
| | Mammalia | 0.17 | 0.17 | 0.18 | 18,000 | 0.0001 | fixed |
| | Reptilia | 0.17 | 0.15 | 0.18 | 18,410 | 0.0001 | fixed |
| | sigma | 0.01 | 0.01 | 0.02 | 18,000 | | residual |
| Taxa (global) - fluctuations CI weighted | Actinopterygii | 0.17 | 0.16 | 0.18 | 18,000 | 0.0001 | fixed |
| | Amphibia | 0.28 | 0.26 | 0.31 | 18,000 | 0.0001 | fixed |
| | Aves | 0.10 | 0.09 | 0.10 | 18,000 | 0.0001 | fixed |
| | Mammalia | 0.20 | 0.19 | 0.21 | 18,000 | 0.0001 | fixed |
| | Reptilia | 0.21 | 0.19 | 0.23 | 18,000 | 0.0001 | fixed |
| | sigma | 0.02 | 0.02 | 0.02 | 18,000 | | residual |
| Taxa (global) - fluctuations SE | Actinopterygii | 0.14 | 0.13 | 0.14 | 18,000 | 0.0001 | fixed |
| | Amphibia | 0.24 | 0.23 | 0.26 | 18,539 | 0.0001 | fixed |
| | Aves | 0.08 | 0.07 | 0.08 | 18,000 | 0.0001 | fixed |
| | Mammalia | 0.15 | 0.14 | 0.16 | 16,776 | 0.0001 | fixed |
| | Reptilia | 0.19 | 0.18 | 0.20 | 18,576 | 0.0001 | fixed |
| | sigma | 0.02 | 0.02 | 0.02 | 18,836 | | residual |
| Taxa (global) - fluctuations SD | Actinopterygii | 0.56 | 0.56 | 0.57 | 18,000 | 0.0001 | fixed |
| | Amphibia | 0.62 | 0.60 | 0.63 | 17,434 | 0.0001 | fixed |
| | Aves | 0.53 | 0.53 | 0.53 | 18,000 | 0.0001 | fixed |
| | Mammalia | 0.60 | 0.59 | 0.60 | 18,000 | 0.0001 | fixed |
| | Reptilia | 0.59 | 0.58 | 0.60 | 18,000 | 0.0001 | fixed |
| | sigma | 0.01 | 0.01 | 0.01 | 18,000 | | residual |
| Amphibian population trends – mu and phylogeny | (Intercept) | -0.009 (-0.009 – -0.008) | -0.005 (-0.005 – -0.005) | 0.031 (0.027 – 0.035) | 10,047 (10,713 – 9350) | 0.59 (0.55 – 0.62) | fixed |
| | phylo | 0.001 (0.001 – 0.002) | 0.001 (0.001 – 0.001) | 0.004( 0.004 – 0.005) | 7,037 (6,171 – 7,602) | | random |
| | species | 0.001 (0.001 – 0.001) | 0.001 (0.001 – 0.001) | 0.002 (0.002 – 0.002) | 7,912 (6,541 – 8,500) | | random |
| | sigma | 0.006 (0.006 – 0.006) | 0.004 (0.004 – 0.004) | 0.007 (0.007 – 0.007) | 9,855 (9,280 – 10,041) | | residual |
| Bird population trends – mu and phylogeny | (Intercept) | 0.005 (0.004 – 0.005) | -0.009 (-0.014 – -0.005) | 0.017 (0.02 – 0.02) | 10,304 (9,515 – 12,560) | 0.45 (0.31 – 0.63) | fixed |
| | phylo | 0.001 (0.001 – 0.003) | 0.001 (0.001 – 0.001) | 0.001 (0.001 – 0.001) | 3,850 (3,305 – 4,144) | | random |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| | species | 0.001 (0.001 – 0.001) | 0.001 (0.001 – 0.001) | 0.001 (0.001 – 0.001) | 6,799 (6,358 – 7,260) | | random |
| | sigma | 0.002 (0.002 – 0.002) | 0.002 (0.002 – 0.002) | 0.003 (0.003 – 0.003) | 9,991 (9,713 – 10,345) | | residual |
| Reptile population trends – mu and phylogeny | (Intercept) | 0.004 (0.003 – 0.005) | -0.004 (-0.005 – -0.003) | 0.047 (0.043 – 0.056) | 10,030 (9,679 – 10,595) | 0.83 (0.81 – 0.85) | fixed |
| | phylo | 0.002 (0.001 – 0.003) | 0.001 (0.001 – 0.001) | 0.007 (0.005 – 0.010) | 3,971 (2,850 – 4,823) | | random |
| | species | 0.004 (0.004 – 0.004) | 0.001 (0.001 – 0.001) | 0.007 (0.007 – 0.007) | 3,418 (3,058 – 3,701) | | random |
| | sigma | 0.004 (0.004 – 0.004) | 0.001 (0.001 – 0.001) | 0.007 (0.007 – 0.007) | 3,562 (3,133 – 3,765) | | residual |
| Amphibian population fluctuations – CI and phylogeny | (Intercept) | 0.271 (0.269 – 0.271) | 0.217 (0.215 – 0.219) | 0.323 (0.321 – 0.327) | 9,905 (8,988 – 10,066) | 0.001 (0.001 – 0.001) | fixed |
| | phylo | 0.001 (0.001 - 0.002) | 0.001 (0.001 - 0.001) | 0.006 (0.005 – 0.006) | 9,754 (8,283 – 10,000) | | random |
| | species | 0.001 (0.001 – 0.001) | 0.001 (0.001 - 0.001) | 0.005 (0.005 – 0.005) | 9,822 (8,557 – 10,000) | | random |
| | sigma | 0.045 (0.0349 – 0.045) | 0.035 (0.035 – 0.035) | 0.056 (0.055 – 0.056) | 10,076 (10,000 – 10,413) | | residual |
| Bird population fluctuations – CI and phylogeny | (Intercept) | 0.114 (0.109 – 0.119) | 0.059 (0.048 – 0.070) | 0.169 (0.153 – 0.188) | 10,000 (10,000 – 10,000) | 0.001 (0.001 – 0.001) | fixed |
| | phylo | 0.005 (0.004 – 0.006) | 0.003 (0.0.002 – 0.003) | 0.007 (0.005 – 0.009) | 4,017 (3,779 – 4,336) | | random |
| | species | 0.002 (0.002 – 0.002) | 0.001 (0.001 – 0.001) | 0.003 (0.002 – 0.003) | 5,706 (4,991 – 6,092) | | random |
| | sigma | 0.010 (0.010 – 0.010) | 0.010 (0.010 – 0.010) | 0.011 (0.011 – 0.011) | 9,936 (9,223 – 11,161) | | residual |
| Reptile population fluctuations – CI and phylogeny | (Intercept) | 0.302 (0.301 – 0.306) | 0.180 (0.1625 – 0.190) | 0.424 (0.410 – 0.434) | 9,837 (9,242 – 10,009) | 0.001 (0.001 – 0.001) | fixed |
| | phylo | 0.014 (0.012 – 0.014) | 0.001 (0.001 – 0.001) | 0.056 (0.050 – 0.060) | 2,779 (2,256 – 3,213) | | random |
| | species | 0.050 (0.049 – 0.050) | 0.019 (0.016 – 0.022) | 0.082 (0.081 – 0.084) | 4,407 (3,544 – 4,730) | | random |

| | | Estimate | CI lower | CI upper | N | p | Effect |
|---|---|---|---|---|---|---|---|
| | sigma | 0.018 (0.018 – 0.018) | 0.006 (0.006 – 0.007) | 0.036 (0.035 – 0.036) | 3,985 (3,817 – 4,536) | | residual |
| Geographic range (birds) - mu | (Intercept) | -0.01 | -0.03 | 0.004 | 18,000 | 0.15 | fixed |
| | log(range) | 0.001 | -0.0000 | 0.002 | 18,000 | 0.05 | fixed |
| | sigma | 0.003 | 0.003 | 0.003 | 18,000 | | residual |
| Geographic range (birds) - weighted | (Intercept) | -0.003 | -0.03 | 0.02 | 18,000 | 0.77 | fixed |
| | log(range) | 0.001 | -0.001 | 0.002 | 17,620 | 0.41 | fixed |
| | sigma | 0.003 | 0.003 | 0.003 | 18,000 | | residual |
| Geographic range (birds) - slope | (Intercept) | -0.02 | -0.04 | 0.01 | 18,000 | 0.26 | fixed |
| | log.range | 0.001 | -0.0003 | 0.003 | 18,000 | 0.10 | fixed |
| | sigma | 0.01 | 0.01 | 0.01 | 18,881 | | residual |
| Geographic range (birds) - fluctuations CI | (Intercept) | 0.02 | -0.02 | 0.05 | 18,000 | 0.37 | fixed |
| | log(range) | 0.005 | 0.003 | 0.01 | 18,000 | 0.0001 | fixed |
| | sigma | 0.01 | 0.01 | 0.01 | 18,000 | | residual |
| Geographic range (birds) - fluctuations CI weighted | (Intercept) | 0.01 | -0.04 | 0.05 | 18,000 | 0.72 | fixed |
| | log(range) | 0.01 | 0.003 | 0.01 | 18,000 | 0.0001 | fixed |
| | sigma | 0.02 | 0.01 | 0.02 | 18,657 | | residual |
| Geographic range (birds) - fluctuations SE | (Intercept) | -0.04 | -0.08 | -0.01 | 18,000 | 0.01 | fixed |
| | log(range) | 0.01 | 0.01 | 0.01 | 18,000 | 0.0001 | fixed |
| | sigma | 0.01 | 0.01 | 0.01 | 18,000 | | residual |
| Geographic range (birds) - fluctuations SD | (Intercept) | 0.47 | 0.44 | 0.50 | 18,000 | 0.0001 | fixed |
| | log(range) | 0.004 | 0.002 | 0.01 | 18,000 | 0.0001 | fixed |
| | sigma | 0.01 | 0.01 | 0.01 | 18,000 | | residual |
| Mean population size (global) - mu | (Intercept) | 0.01 | 0.001 | 0.01 | 18,000 | 0.03 | fixed |
| | log.meanpop | 0.001 | -0.0002 | 0.001 | 18,000 | 0.14 | fixed |
| | sigma | 0.005 | 0.005 | 0.01 | 17,393 | | residual |
| Mean population size (global) - weighted | (Intercept) | 0.01 | 0.0004 | 0.01 | 17,924 | 0.04 | fixed |
| | log.meanpop | 0.001 | -0.001 | 0.002 | 18,000 | 0.37 | fixed |
| | sigma | 0.01 | 0.01 | 0.01 | 18,000 | | residual |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| Mean population size (global) - slope | (Intercept) | 0.01 | 0.0004 | 0.02 | 17,382 | 0.04 | fixed |
| | log.meanpop | 0.001 | -0.0003 | 0.002 | 17,360 | 0.14 | fixed |
| | sigma | 0.01 | 0.01 | 0.01 | 18,000 | | residual |
| Mean population size (global) - fluctuations CI | (Intercept) | 0.21 | 0.20 | 0.23 | 18,000 | 0.0001 | fixed |
| | log.meanpop | -0.004 | -0.01 | -0.002 | 18,000 | 0.0001 | fixed |
| | sigma | 0.03 | 0.03 | 0.03 | 18,000 | | residual |
| Mean population size (global) - fluctuations CI weighted | (Intercept) | 0.25 | 0.23 | 0.26 | 18,000 | 0.0001 | fixed |
| | log.meanpop | -0.01 | -0.01 | -0.003 | 18,000 | 0.0001 | fixed |
| | sigma | 0.03 | 0.03 | 0.03 | 18,000 | | residual |
| Mean population size (global) - fluctuations SE | (Intercept) | 0.22 | 0.21 | 0.23 | 18,000 | 0.0001 | fixed |
| | log.meanpop | -0.01 | -0.01 | -0.01 | 18,000 | 0.0001 | fixed |
| | sigma | 0.03 | 0.03 | 0.03 | 17,622 | | residual |
| Mean population size (global) - fluctuations SD | (Intercept) | -0.002 | -0.002 | -0.001 | 18,000 | 0.0001 | fixed |
| | log.meanpop | 0.0003 | 0.0002 | 0.0004 | 18,000 | 0.0001 | fixed |
| | sigma | 0.0002 | 0.0001 | 0.0002 | 18,000 | | residual |
| Habitat specificity (global) - mu | (Intercept) | 0.004 | 0.001 | 0.01 | 18,000 | 0.002 | fixed |
| | hab_spec | 0.0001 | -0.0001 | 0.0004 | 18,000 | 0.36 | fixed |
| | sigma | 0.004 | 0.004 | 0.004 | 18,000 | | residual |
| Habitat specificity (global) - weighted | (Intercept) | 0.003 | -0.0002 | 0.01 | 18,000 | 0.06 | fixed |
| | hab_spec | 0.0003 | 0.0000 | 0.001 | 18,000 | 0.05 | fixed |
| | sigma | 0.004 | 0.004 | 0.004 | 18,000 | | residual |
| Habitat specificity (global) - slope | (Intercept) | 0.01 | 0.002 | 0.01 | 17,426 | 0.004 | fixed |
| | hab_spec | 0.0002 | -0.0002 | 0.001 | 17,536 | 0.41 | fixed |
| | sigma | 0.01 | 0.01 | 0.01 | 18,000 | | residual |
| Habitat specificity (global) - fluctuations CI | (Intercept) | 0.11 | 0.10 | 0.11 | 17,358 | 0.0001 | fixed |
| | hab_spec | 0.001 | 0.001 | 0.002 | 18,000 | 0.0001 | fixed |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| | sigma | 0.02 | 0.02 | 0.02 | 18,000 | | residual |
| Habitat specificity (global) - fluctuations CI weighted | (Intercept) | 0.12 | 0.11 | 0.12 | 18,000 | 0.0001 | fixed |
| | hab_spec | 0.001 | 0.001 | 0.002 | 18,000 | 0.0001 | fixed |
| | sigma | 0.02 | 0.02 | 0.02 | 18,000 | | residual |
| Habitat specificity (global) - fluctuations SE | (Intercept) | 0.10 | 0.09 | 0.10 | 18,000 | 0.0001 | fixed |
| | hab_spec | 0.001 | 0.001 | 0.002 | 18,000 | 0.0001 | fixed |
| | sigma | 0.02 | 0.02 | 0.02 | 18,000 | | residual |
| Habitat specificity (global) - fluctuations SD | (Intercept) | 0.54 | 0.54 | 0.55 | 18,000 | 0.0001 | fixed |
| | hab_spec | 0.001 | 0.0002 | 0.001 | 18,563 | 0.002 | fixed |
| | sigma | 0.01 | 0.01 | 0.01 | 18,000 | | residual |
| Red List status (global) - mu | Least concern | 0.01 | 0.005 | 0.01 | 18,000 | 0.0001 | fixed |
| | Near threatened | -0.01 | -0.01 | -0.0005 | 18,000 | 0.04 | fixed |
| | Vulnerable | 0.002 | -0.003 | 0.01 | 18,294 | 0.40 | fixed |
| | Endangered | -0.002 | -0.01 | 0.01 | 18,000 | 0.60 | fixed |
| | Critically endangered | -0.01 | -0.02 | 0.0001 | 18,000 | 0.05 | fixed |
| | sigma | 0.004 | 0.003 | 0.004 | 18,000 | | residual |
| Red List status (global) - weighted | Least concern | 0.01 | 0.01 | 0.01 | 19,846 | 0.0001 | fixed |
| | Near threatened | -0.003 | -0.01 | 0.005 | 18,000 | 0.45 | fixed |
| | Vulnerable | 0.002 | -0.005 | 0.01 | 18,000 | 0.52 | fixed |
| | Endangered | -0.01 | -0.02 | 0.003 | 18,000 | 0.19 | fixed |
| | Critically endangered | -0.02 | -0.03 | -0.01 | 18,000 | 0.01 | fixed |
| | sigma | 0.004 | 0.004 | 0.004 | 18,000 | | residual |
| Red List status (global) - slope | Least concern | 0.01 | 0.01 | 0.01 | 18,000 | 0.0001 | fixed |
| | Near threatened | -0.01 | -0.02 | -0.004 | 18,000 | 0.01 | fixed |
| | Vulnerable | 0.002 | -0.01 | 0.01 | 17,507 | 0.58 | fixed |
| | Endangered | -0.001 | -0.01 | 0.01 | 18,000 | 0.91 | fixed |
| | Critically endangered | -0.01 | -0.02 | 0.01 | 17,564 | 0.44 | fixed |
| | sigma | 0.01 | 0.01 | 0.01 | 18,000 | | residual |
| Red List status (global) - fluctuations CI | Least concern | 0.11 | 0.11 | 0.12 | 18,000 | 0.0001 | fixed |
| | Near threatened | 0.14 | 0.13 | 0.15 | 18,000 | 0.0001 | fixed |
| | Vulnerable | 0.14 | 0.13 | 0.15 | 19,536 | 0.0001 | fixed |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| | Endangered | 0.16 | 0.15 | 0.17 | 18,000 | 0.0001 | fixed |
| | Critically endangered | 0.17 | 0.15 | 0.19 | 18,000 | 0.0001 | fixed |
| | sigma | 0.02 | 0.02 | 0.02 | 18,000 | | residual |
| Red List status (global) - fluctuations CI weighted | Least concern | 0.12 | 0.12 | 0.12 | 18,432 | 0.0001 | fixed |
| | Near threatened | 0.16 | 0.15 | 0.18 | 18,812 | 0.0001 | fixed |
| | Vulnerable | 0.15 | 0.14 | 0.17 | 18,832 | 0.0001 | fixed |
| | Endangered | 0.19 | 0.17 | 0.21 | 17,714 | 0.0001 | fixed |
| | Critically endangered | 0.19 | 0.16 | 0.22 | 18,469 | 0.0001 | fixed |
| | sigma | 0.02 | 0.02 | 0.02 | 18,000 | | residual |
| Red List status (global) - fluctuations SE | Least concern | 0.10 | 0.10 | 0.10 | 18,000 | 0.0001 | fixed |
| | Near threatened | 0.13 | 0.12 | 0.14 | 19,308 | 0.0001 | fixed |
| | Vulnerable | 0.11 | 0.10 | 0.12 | 18,645 | 0.0001 | fixed |
| | Endangered | 0.13 | 0.12 | 0.15 | 18,000 | 0.0001 | fixed |
| | Critically endangered | 0.15 | 0.13 | 0.17 | 18,000 | 0.0001 | fixed |
| | sigma | 0.02 | 0.02 | 0.02 | 18,555 | | residual |
| Red List status (global) - fluctuations SD | Least concern | 0.54 | 0.54 | 0.55 | 18,390 | 0.0001 | fixed |
| | Near threatened | 0.57 | 0.56 | 0.58 | 18,000 | 0.0001 | fixed |
| | Vulnerable | 0.56 | 0.55 | 0.57 | 18,214 | 0.0001 | fixed |
| | Endangered | 0.59 | 0.58 | 0.60 | 18,000 | 0.0001 | fixed |
| | Critically endangered | 0.61 | 0.59 | 0.62 | 17,295 | 0.0001 | fixed |
| | sigma | 0.01 | 0.01 | 0.01 | 18,000 | | residual |
| Latitude (global) - mu | (Intercept) | 0.004 | 0.002 | 0.01 | 18,254 | 0.0001 | fixed |
| | Decimal.Latitude | -0.0000 | -0.0000 | 0.0000 | 18,670 | 0.90 | fixed |
| | sigma | 0.003 | 0.003 | 0.004 | 18,000 | | residual |
| Duration (global) - mu | End | 0.0000 | 0.0000 | 0.0001 | 18,561 | 0.03 | fixed |
| | sigma | 0.003 | 0.003 | 0.004 | 18,000 | | residual |

**Table S3. List of species included in the UK scale analysis of population change across rarity metrics.**

| Species name | Number of populations |
|---|:---:|
| *Acrocephalus schoenobaenus* | 1 |
| *Acrocephalus scirpaceus* | 1 |
| *Agonus cataphractus* | 1 |
| *Alca torda* | 4 |
| *Anarhichas lupus* | 1 |
| *Anas acuta* | 1 |
| *Anas crecca* | 1 |
| *Anas platyrhynchos* | 2 |
| *Anser albifrons* | 4 |
| *Anser fabalis* | 1 |
| *Anthus pratensis* | 2 |
| *Ardea cinerea* | 1 |
| *Arenaria interpres* | 1 |
| *Argentina silus* | 1 |
| *Argentina sphyraena* | 1 |
| *Arnoglossus laterna* | 1 |
| *Asio flammeus* | 1 |
| *Aythya ferina* | 4 |
| *Aythya fuligula* | 2 |
| *Botaurus stellaris* | 14 |
| *Branta bernicla* | 54 |
| *Branta canadensis* | 1 |
| *Branta leucopsis* | 3 |
| *Brosme brosme* | 1 |
| *Bucephala clangula* | 1 |
| *Bufo bufo* | 1 |
| *Burhinus oedicnemus* | 1 |
| *Buteo buteo* | 1 |
| *Calidris alba* | 1 |
| *Calidris alpina* | 1 |
| *Calidris canutus* | 2 |
| *Calidris maritima* | 2 |
| *Callionymus maculatus* | 1 |

| | |
|---|---|
| *Capreolus capreolus* | 3 |
| *Carduelis cannabina* | 2 |
| *Cepphus grylle* | 2 |
| *Cervus elaphus* | 2 |
| *Cetorhinus maximus* | 1 |
| *Cettia cetti* | 1 |
| *Charadrius hiaticula* | 1 |
| *Chelidonichthys lucerna* | 2 |
| *Circus aeruginosus* | 1 |
| *Circus cyaneus* | 1 |
| *Clupea harengus* | 2 |
| *Columba oenas* | 1 |
| *Coronella austriaca* | 1 |
| *Corvus corax* | 1 |
| *Corvus corone* | 1 |
| *Corvus monedula* | 1 |
| *Crex crex* | 1 |
| *Cyclopterus lumpus* | 1 |
| *Cygnus columbianus* | 3 |
| *Cygnus cygnus* | 1 |
| *Cygnus olor* | 1 |
| *Delphinus delphis* | 2 |
| *Echiichthys vipera* | 1 |
| *Egretta garzetta* | 1 |
| *Emberiza cirlus* | 1 |
| *Emberiza citrinella* | 1 |
| *Emberiza schoeniclus* | 3 |
| *Eptesicus serotinus* | 1 |
| *Esox lucius* | 2 |
| *Falco peregrinus* | 1 |
| *Falco tinnunculus* | 1 |
| *Fulica atra* | 1 |
| *Fulmarus glacialis* | 7 |
| *Gadus morhua* | 7 |
| *Glyptocephalus cynoglossus* | 1 |
| *Haematopus ostralegus* | 2 |
| *Haliaeetus albicilla* | 1 |
| *Halichoerus grypus* | 52 |

| | |
|---|---|
| *Hippoglossus hippoglossus* | 1 |
| *Lagopus lagopus* | 3 |
| *Larus argentatus* | 1 |
| *Larus canus* | 1 |
| *Larus fuscus* | 1 |
| *Larus melanocephalus* | 1 |
| *Lepidorhombus whiffiagonis* | 2 |
| *Lepus timidus* | 1 |
| *Limosa lapponica* | 2 |
| *Limosa limosa* | 2 |
| *Lissotriton vulgaris* | 5 |
| *Lophius budegassa* | 1 |
| *Lophius piscatorius* | 2 |
| *Lullula arborea* | 1 |
| *Melanogrammus aeglefinus* | 9 |
| *Meles meles* | 1 |
| *Mergus serrator* | 1 |
| *Merlangius merlangus* | 8 |
| *Merluccius merluccius* | 2 |
| *Micromesistius poutassou* | 1 |
| *Milvus milvus* | 3 |
| *Molva molva* | 1 |
| *Morus bassanus* | 4 |
| *Muscicapa striata* | 1 |
| *Myotis nattereri* | 1 |
| *Natrix natrix* | 1 |
| *Netta rufina* | 1 |
| *Numenius arquata* | 2 |
| *Nyctalus noctula* | 1 |
| *Oenanthe oenanthe* | 1 |
| *Orcinus orca* | 1 |
| *Oriolus oriolus* | 1 |
| *Oryctolagus cuniculus* | 6 |
| *Oxyura jamaicensis* | 2 |
| *Pandion haliaetus* | 1 |
| *Parus major* | 2 |
| *Passer domesticus* | 1 |
| *Passer montanus* | 1 |

| | |
|---|---|
| *Perca fluviatilis* | 3 |
| *Perdix perdix* | 1 |
| *Phalacrocorax aristotelis* | 8 |
| *Phoca vitulina* | 28 |
| *Phrynorhombus norvegicus* | 1 |
| *Pipistrellus pipistrellus* | 1 |
| *Pipistrellus pygmaeus* | 1 |
| *Platichthys flesus* | 1 |
| *Plecotus auritus* | 1 |
| *Plectrophenax nivalis* | 2 |
| *Pleuronectes platessa* | 6 |
| *Pluvialis apricaria* | 2 |
| *Pluvialis squatarola* | 1 |
| *Podiceps cristatus* | 1 |
| *Pollachius pollachius* | 1 |
| *Pollachius virens* | 3 |
| *Prunella modularis* | 1 |
| *Puffinus mauretanicus* | 2 |
| *Pyrrhula pyrrhula* | 1 |
| *Raja brachyura* | 1 |
| *Raja clavata* | 1 |
| *Raja microocellata* | 1 |
| *Raja montagui* | 1 |
| *Rhinolophus ferrumequinum* | 2 |
| *Rhinolophus hipposideros* | 4 |
| *Rissa tridactyla* | 9 |
| *Salmo salar* | 1 |
| *Salmo trutta* | 1 |
| *Scomber scombrus* | 2 |
| *Scyliorhinus canicula* | 1 |
| *Sitta europaea* | 1 |
| *Sprattus sprattus* | 1 |
| *Stenella coeruleoalba* | 1 |
| *Stercorarius parasiticus* | 3 |
| *Sterna dougallii* | 5 |
| *Sterna hirundo* | 1 |
| *Sterna paradisaea* | 1 |
| *Sternula albifrons* | 1 |

| | |
|---|---|
| *Streptopelia turtur* | 1 |
| *Strix aluco* | 1 |
| *Sturnus vulgaris* | 1 |
| *Sylvia communis* | 1 |
| *Syngnathus rostellatus* | 1 |
| *Tachybaptus ruficollis* | 1 |
| *Tadorna tadorna* | 1 |
| *Thalasseus sandvicensis* | 2 |
| *Trachurus trachurus* | 2 |
| *Tringa nebularia* | 1 |
| *Tringa totanus* | 1 |
| *Trisopterus esmarkii* | 2 |
| *Trisopterus luscus* | 1 |
| *Trisopterus minutus* | 1 |
| *Triturus cristatus* | 5 |
| *Turdus philomelos* | 1 |
| *Tursiops truncatus* | 4 |
| *Uria aalge* | 10 |
| *Vanellus vanellus* | 3 |
| *Xiphias gladius* | 1 |

971 **Table S4. Profiling method for estimating habitat specificity for 144 species**

972 **with populations in the UK in the LPD.** We extracted the habitats in which each

973 species occurs from their IUCN profiles (http://www.iucnredlist.org/) and we followed

974 this key for consistency.

| Habitat | Considered to be the same as: | Considered to be different to: |
|---|---|---|
| **Rural park** | Suburban park, urban park, rural garden, suburban garden, urban garden | |
| **Lake** | Big lake, small lake, pond, pool, dam, oxbow lake, reservoir | |
| **Bog** | Swamp, bogland | Lagoon |
| **Coastal cliff** | Island cliff | |
| **Shingle beach** | Pebble beach, rock beach | Sandy beach |
| **Stream** | River | Weir |
| **Fruit Tree Plantation** | Fruit Garden, orchard | |
| **Thicket** | Copse, grove, small stand | |
| **Forest** | | Woodland |
| **Glade** | Forest Clearing | |
| **Broadleaf** | Deciduous | |
| **Urban** | Suburban | |
| **River margin** | Various types of river margins | |
| **Tidal Creek** | | Estuary |
| **Harbour** | Dock, jetty, pier | |
| **Bush lands** | Shrublands | |
| **Irrigation channel** | Ditch | |
| **Heath** | Moorland | |
| **Sandy beach** | Spit, dune | Shingle beach, pebble beach, rock beach |
| **Crag** | Rocky outcrop, cliff, rocky slope | |

| | |
|---|---|
| **Marsh** | Wet meadow |
| **Islet** | Island |