Rarity and conservation status do not predict vertebrate population trends

Gergana N. Daskalova[1]*, Isla H. Myers-Smith[1] and John L. Godlee[1]

1. School of GeoSciences, University of Edinburgh, Edinburgh, UK

gndaskalova@gmail.com

johngodlee@gmail.com

isla.myers-smith@ed.ac.uk

* Corresponding author:

Gergana N. Daskalova

gndaskalova@gmail.com

Crew Building, King's Buildings

West Mains Road

Edinburgh EH9 1UU

Scotland

Tel: +44 7946274522

Short running title: Rarity does not explain pop. change

Keywords: population change, rarity, vertebrates, species attributes, geographic

range, habitat specificity, population size

Submission type: Letter

Abstract: 144 words

Main text: 4589 words

References: 84

Figures and Tables: 5

Statement of authorship:

GND and IMS conceived the idea and conducted the statistical analyses. JLG

calculated geographic range estimates. All authors contributed to the integration of

the LPI, GBIF and IUCN databases. GND wrote the first draft of the manuscript and

all authors contributed to revisions.

Data accessibility statement:

We will make all R code for the analyses and data publicly available on GitHub and will

include the relevant DOIs and links.

Abstract

Species' attributes such as rarity and conservation status are often assumed to predict population change which in turn can influence conservation decision-making; however, there are very few empirical tests across taxa and biomes. We use a database of 9284 vertebrate populations from 2084 species in the UK and globally to quantify population change across rarity, Red List status, biomes and taxa. In the UK, rarity metrics (geographic range, population size and habitat specificity) did not explain trends, but greater fluctuations were more likely for smaller populations. There was no consistent pattern of net declines with increasing Red List status in the UK and globally. We found taxonomic but not geographic patterning of population change at global scales. Our findings suggest that rarity metrics and conservation status are poor predictors of vertebrate population trends, but that rare species were nevertheless more likely to fluctuate, potentially increasing their extinction risk.

Introduction

Species' attributes such as rarity, conservation status, distribution and taxon are often assumed to predict variation in extinction risk (Purvis *et al.* 2000; Olden *et al.* 2007; Fritz *et al.* 2009; Ripple *et al.* 2017) and in population change (Julliard *et al.* 2004; Shultz *et al.* 2005). Population trends (Hutchings & Reynolds 2004) and population fluctuations (Melbourne & Hastings 2008) have both been frequently linked with a higher likelihood of a species' extinction. For decades, conservation efforts have focused on protecting rare species (Pigott 1977; Gunn 1980; Longton & Hedderson 2000; Povilitis 2001; Gaston & Fuller 2008), those with restricted geographic extents, small population sizes or high habitat specificity, as these species are thought to be more likely to be threatened with extinction. However, there are very few empirical tests of the influence of rarity and conservation status on population change and extinction risk (e.g., Purvis *et al.* 2000; O'Grady *et al.* 2004; Ripple *et al.* 2017), and those that exist mostly focus on either specific taxa (Jones *et al.* 2003; Cardillo *et al.* 2004; Fritz *et al.* 2009) or on population declines alone (Purvis *et al.* 2000; Fritz *et al.* 2009; Gilroy *et al.* 2016; Hefley *et al.* 2016; Ceballos *et al.* 2017), and not population change across taxa and geographic regions. Quantifying the variation in trends and fluctuations of populations around the world can provide valuable insight for conservation policy and decision-making by testing the processes that ultimately lead to community reassembly (Batt *et al.* 2017) and by determining the drivers of population change (Ehrlén & Morris 2015; Hefley *et al.* 2016).

Understanding what factors best explain population change and extinction risk across species has been a key focus in population ecology and conservation biology, with possible measures including species' attributes and traits, such as rarity (Gaston &

Blackburn 1995; Owens & Bennett 2000; Purvis *et al.* 2000; Cardillo *et al.* 2004;

O'Grady *et al.* 2004; Olden *et al.* 2007; Collen *et al.* 2009; Ripple *et al.* 2017).

Extinction risk, the likelihood that a species will go extinct within a given time frame

(Mace *et al.* 2008), is widely assumed to increase when species are rare (i.e., they

have small geographic ranges (Jones *et al.* 2003), small population sizes (Green

2003) and are very specialized (Gilroy *et al.* 2016)). Rare species with small

populations are more likely to undergo stochastic fluctuations that could lead to

extinction at any given point as per population dynamics theory (Lande 1993;

Melbourne & Hastings 2008). Population declines can lead to local and eventually

potentially global species extinction (Ceballos *et al.* 2017). Small populations are also

more vulnerable to demographic stochasticity and inbreeding (Kareiva 1990; Hanski

1998). Allee effects, the positive relationship between population growth rate and

population density, further increase the likelihood of declines due to lack of potential

mates and low reproductive output once populations reach a critically low density

(Dennis *et al.* 2016; Sun 2016). Thus, small populations from rare species are

thought to be more likely to experience population declines (McKinney 1997; Purvis

et al. 2000; Manne & Pimm 2001; Hutchings & Reynolds 2004).

Investigating population change over time in relation to rarity can provide insights into

species' extinction risk. Rabinowitz (1981) defined 'seven forms of rarity' based on

the combinations of the extremes of three characteristics: geographic range (small

vs. large), habitat specificity (generalist vs. specialist), and local population size

(small vs. large). Rarity can be either a state in which species naturally occur, or a

state to which they are driven following external perturbations to populations

(Magurran & Henderson 2003). However, previous analyses of the influence of rarity

on population trends and extinction risk have shown mixed results. Geographic range

did not correlate with species' vulnerability to extinction in 45 bird and mammal taxa

from all continents except Antarctica (O'Grady *et al.* 2004) and the minimum viable

population size did not vary with geographic range and population size in 629

invertebrate, 529 vertebrate and 30 plant species (Brook *et al.* 2006). In contrast, in

bats, geographic range was found to be the most important predictor of extinction risk

(Jones *et al.* 2003). Thus, a generalizable relationship between rarity and population

declines and increased extinction risk across taxa and geographic regions remains

unquantified. However, simplifying the concept of rarity so that it can be modelled as

a predictor of population change remains a considerable challenge (Magurran 2013;

Batt *et al.* 2017).

Investigations of potential factors that predispose species to population change

across geographic and taxonomic scales (e.g., from country to biome level, and from

population to taxa level) are particularly valuable for guiding conservation decision-

making (O'Grady *et al.* 2004; Ehrlén & Morris 2015; Hefley *et al.* 2016; Batt *et al.*

2017). Conservation actions are predominantly implemented on smaller, country-

specific, regional or local scales (Mills *et al.* 2010; Guerrero *et al.* 2013), yet policy

may be determined for larger regions or broader taxonomic groups (Agardy 2005;

Pressey *et al.* 2007). Similarly, overall species' Red List status is determined on a

global scale based on multiple criteria (Mace *et al.* 2008), but on a local scale,

populations may vary in the direction and magnitude of experienced change (Ehrlén

& Morris 2015; Lawson *et al.* 2015; Gilroy *et al.* 2016; van Strien *et al.* 2016; Leung *et*

al. 2017). Population time-series data from the Living Planet Database (LPD),

complemented by extensive occurrence data from the Global Biodiversity Information

Facility (GBIF) and information on species habitat preferences and conservation

status from the IUCN Red List, provide a unique opportunity to test which species

attributes explain global and country-specific population trends.

Here, we addressed the research question: Is variation in vertebrate population

change explained by differences in species attributes, such as rarity metrics and

conservation status? We investigated two types of population change – population

trends and population fluctuations. We combined open source data from the LPD,

GBIF and IUCN to examine population change in the UK, a country with high-quality

monitoring and abundance data, and global population change across species' Red

List status, biomes, systems (freshwater, marine, terrestrial) and taxa. We considered

rarity to be the state in which species exist if their geographic range is small vs. large,

their local population size is low vs. high, or they are habitat generalists vs.

specialists. Thus, we quantified rarity through these three individual metrics -

geographic range extent, mean population size, habitat specificity, and additionally

species' Red List status. We tested the following hypotheses: (1) in the UK, rare

species are more likely to be experiencing negative population trends than common

species; (2) on a global scale, declining populations are more likely for species from

more threatened Red List categories, (3) there are geographic and taxonomic

patterns in vertebrate population trends and fluctuations (e.g., hotspots of declines,

increases or fluctuations in abundances in specific regions and in specific taxa) and

(4) rare species or those with a higher Red List status are more likely to experience

greater population fluctuations.

Methods

 Population data

 To quantify vertebrate population change, we extracted the abundance data of all

 populations monitored at more than five different time points from the LPD (LPI

 2016), resulting in 9284 populations from 2084 species for our system, biome, taxa

 and conservation status analysis (global scale) and 508 populations from 237

 species for our rarity analysis (UK scale, but note that habitat specificity data were

 only available for 402 populations from 206 species, Table S1). Monitoring duration

 differed among populations; however, in our analysis, we retained all populations that

 had at least five time points of monitoring data (Figures S1 and S2). Before analysis,

 we scaled the abundance of each population to a common magnitude between zero

 and one, because the populations were monitored using different metrics of

 abundance (e.g. population indices vs. number of individuals). This allowed us to

 explore trends among populations relative to the variation experienced across each

 time series.

 Rarity metrics data

 We defined rarity following a simplified version of the 'seven forms of rarity' model

 (Rabinowitz 1981), and thus consider rarity to be the state in which species exist

 when they have a small geographic range, low population size, or narrow habitat

 specificity. We combined publicly available data from three sources: (1) population

 records for vertebrate species from the LPD (LPI 2016), (2) occurrence data from the

 Global Biodiversity Information Facility (GBIF 2017), and (3) habitat specificity and

 Red List classification data for each species from the International Union for

 Conservation (IUCN 2017). Extracting the necessary data and calculating the rarity

 metrics was challenging and computationally-intensive, thus we did not estimate

rarity metrics for all species globally. Instead, we focused on analyzing rarity effects

for species from the UK, a country with intensive and detailed biodiversity monitoring

of vertebrate species.

To estimate range, we calculated a km^2 square occurrence area for each UK

vertebrate species with available population data in the LPD based on the maximum

and minimum latitudes and longitudes of species' GBIF occurrence data cleaned for

outliers by excluding records at sea for terrestrial species and records on land for

marine species (GBIF 2017, accessed between Feb 2016 and Apr 2017). Latitude-

longitude polygons were transformed to km^2 using the WGS84 Coordinate Reference

System (CRS) from the package *geosphere* (Hijmans 2016). We did not use IUCN

range maps, as they were not available for all of our study species, and previous

studies using GBIF occurrences to estimate range have found a positive correlation

between GBIF-derived and IUCN-derived geographic range (Phillips *et al.* 2017).

We calculated mean population size across the monitoring duration for each

population using the raw population data. To create an index of habitat specificity, we

extracted the number of distinct habitats a species occupies from the online IUCN

profiles for our study species between Jan and Apr 2017 (IUCN 2017). We included

both breeding and non-breeding habitats to determine distinct habitat preferences for

each species (Table S2). We obtained global conservation status data for all study

species through the IUCN Red List classification (IUCN 2017).

Population change modelling

As the first stage of our analysis, we modeled population change to produce a

population trend (slope and variance of population change over time). We used state-

space models that model abundance over time (freely-varying intercept) to calculate

the amount of overall population change experienced by each population (*mu*),

following Dennis *et al.* (2006) and Humbert *et al.* (2009). State-space models account

for process noise and observation error and thus deliver reliable estimates of

population change when working with large datasets where records were collected

using different approaches (Knape *et al.* 2011; Pedersen *et al.* 2011; Leung *et al.*

2017). Not accounting for process and measurement error could lead to over-

estimation of population declines (Rueda-Cediel *et al.* 2018). Positive *mu* values

indicate population increase and negative *mu* values indicate population decline. We

calculated the error around *mu* for each population ((upper 95% confidence limit –

lower 95% confidence limit) / 2), with larger errors indicating populations that have

experienced more fluctuations (variance in abundance metrics) overtime.

Population trends across rarity metrics and species' Red List status in the UK

To test the influence of rarity metrics (geographic range, mean population size and

habitat specificity) and species' Red List status on variation in population trends in

the UK, we conducted a second stage analysis using four Bayesian linear models

(one per metric) based on Markov Chain Monte Carlo Bayesian inference (Hadfield

2010). The response variable was population trend (*mu* values from state-space

models), and fixed effects were geographic range (log transformed), mean population

size (log transformed), habitat specificity (number of distinct habitats occupied) and

Red List status (categorical). We used weakly informative priors for all coefficients

(an inverse Wishart prior for the variances and a normal prior for the fixed effects),

and we refer to predictors as 'significant' when the 95% credible interval for the

corresponding parameter in the fitted models did not overlap zero. We did not include

random effects in the Bayesian models (e.g., a species or region random intercept),

because only a very small proportion of the studied species had more than three

monitored populations and spatial replication was low (Table S1).

Global population trends across species' Red List status

To investigate the relationship between population trends and species' Red List

status on a global scale, we modelled population trends from the first stage of the

analysis (state-space models) as a function of Red List status using a Bayesian linear

model.

Global population trends across latitude, biomes, systems and taxa

To investigate the geographic and taxonomic patterns of population trends, we

estimated population trend from the first stage of the analysis (state-space models)

as a function of (1) latitude and biome (as defined by the 'biome' category in the LPD,

e.g. 'temperate broadleaf forest', Olson & Dinerstein 2002), (2) system (freshwater,

marine, terrestrial), and (3) taxon (*Actinopterygii*, bony fish; *Amphibia*, amphibians;

Aves, birds; *Mammalia*, mammals; *Reptilia*, reptiles). Each categorical model from

this second stage of our analyses was fitted with a zero intercept to allow us to

determine if net population trends differed from zero for each of the categories under

investigation.

Population fluctuations across systems, taxa, Red List status and rarity metrics

To test whether rare species fluctuate more than common species and whether there

is geographic and taxonomic patterning in population fluctuations, we modelled the

error around *mu* (half of the 95% confidence interval) as a response variable across

systems, taxa, Red List status and rarity metrics.

Monitoring duration and units

To assess the influence of monitoring duration on population trends, we used a

Bayesian linear model (Figures S1 and S2). We modelled population trend (*mu*) as a

function of monitoring duration (years) for each population, fitted with a zero

intercept, as when duration is zero, no population change has occurred. The units of

population monitoring across the LPD varied with 1368 different categorical levels

(e.g., population index, nest counts, numbers of individuals). Variation in population

trends were not explained by sampling method across the five most commonly used

units (Figure S3). We estimated population change using state-space models

specifically because of their suitability for analyses of large databases where records

were derived using different methods (Humbert *et al.* 2009; Leung *et al.* 2017).

Comparison of modelling approaches

We conducted the following supplementary analyses: in the second-stage Bayesian

models estimating population trends across systems, biomes, taxa and rarity metrics,

1) we weighed *mu* values by the square of *tau*, the observation error estimate derived

from the state-space models, 2) we used slopes of linear model fits of abundance

(scaled at the population level) instead of the *mu* estimates from state-space models,

3) we modelled the standard error around the slope values of the linear models and

the standard deviation of the raw population data for each time-series as additional

metrics of population fluctuations. All different analytical approaches yielded very

similar results (see main text and Figures S6 and S7, Table S3).

All statistical analyses were conducted using R (version 3.4.2, R Core Team 2017).

Code and data are publicly available on GitHub (link to be added at time of

publication).

Results

Population trends across rarity metrics and species' Red List status in the UK

Across monitored vertebrate populations in the UK, species with smaller ranges,

smaller population sizes, or narrower habitat specificity (i.e. rare species) and

species with more threatened Red List status were not more prone to population

declines than common species (Figure 1a-f, Table S3). Over the monitoring duration

for each population, species classified as least concern and near threatened had a

positive population trend (LC: slope = 0.01, CI = 0.005 to 0.02; NT: slope = 0.03, CI =

0.01 to 0.05), whereas vulnerable, endangered and critically endangered populations

did not experience directional change (Figure 1d, f and Table S3). For example, one

monitored colony for the common species grey seal (*Halichoerus grypus*) in Scotland

increased by 158% over a 17-year period going from 25,824 to 66,522 individuals;

whereas, a population of the vulnerable species Atlantic horse mackerel (*Trachurus*

trachurus) remained stable across a 21-year monitoring period.

Figure 1. Effects of rarity metrics (geographic range, mean population size and habitat specificity, plots a-d) and of Red List status (plots e and f) on population trends of UK species. Results include 508 populations from 237 species in the UK. The *mu* values (plots a-c, e) are state-space model fits of changes

in abundance over the monitoring duration for each population. Effect sizes (plots d

and f) are population trends calculated using state-space and linear models, and

population trends (*mu*) weighted by *tau*, the observation error estimate from the

state-space models. Error bars show confidence intervals around population trend

(*mu,* plots a-c, e) and credible intervals for *mu* across rarity metrics and Red List

status (plots d and f). The numbers on the x axis show the sample size in each

category (plot d, e, f). See Table S3 for full model outputs.

Global analysis of population trends across Red List status

Globally, we found increasing, decreasing and stable populations across all Red List

categories of conservation status. For example, a population of the least concern

species red deer (*Cervus elaphus*) in Canada declined by 68% over 7 years going

from 606 to 194 individuals and a population of the critically endangered Hawksbill

sea turtle (*Eretmochelys imbricate*) from Barbados increased by 269% over 7 years

going from 89 to 328 individuals. Populations from least concern species were more

likely to increase (slope = 0.006, CI = 0.005 to 0.007), whereas populations from

vulnerable and endangered species did not experience net directional change

(Figure 2a). Populations from near threatened and critically endangered species were

more likely to decline (NT: slope = -0.005, CI = -0.01 to -0.001; CR: slope = -0.01, CI

= -0.02 to -0.001). Across all categories, there was no consistent pattern of greater

population declines with increasing Red List status (Table S3). Most of the

populations in the LPD that were monitored at more than five time points were from

species categorized as least concern (Figure 2b).

(a) Distribution of records   (b) Red List status effects

Figure 2. Distribution of population records by Red List status (plot a) and global vertebrate population trends across species' Red List status (plot b). Results include 8064 populations from 1702 species. Effect sizes (plot b) include the population trend (*mu*) calculated using state-space and linear models, and population trend (*mu*) weighted by *tau*, the observation error estimate from the state-space models. The numbers on the x axis show the sample size in each Red List category (plot b). Error bars show 95% credible intervals (plot b). See Table S3 for full model outputs.

Global population trends across latitude and biomes

Global vertebrate population trends did not show distinct geographic patterning, as increasing, stable and declining populations were present on all continents and across all latitudes (Figure 3a and b). Populations in Mediterranean forests, montane grasslands, polar freshwaters, temperate wetlands, tropical forests and tropical coral biomes were more likely to increase, whereas populations from the remaining studied biomes were not more likely to be increasing versus decreasing (Figure 3c and Table S3). The lack of biogeographic patterning in vertebrate population trends was also apparent on a UK scale (Figure S4 and Table S3).

Figure 3. Lack of biogeographic patterning in global vertebrate population

trends (plot a), latitudinal distribution of population trends (plot b), and

population trends across biomes (plot c). Results include 9284 populations from

2084 species. Effect sizes (plot c) include population trend (*mu*) calculated using

state-space and linear models, and population trend (*mu*) weighted by *tau*, the

observation error estimate from the state-space models. The numbers on the x axis

show the sample size for each biome category (plot c). Error bars show 95% credible

intervals (plot c). See Table S3 for full model outputs.

Global population trends across systems and taxa

Globally, monitored vertebrate populations experienced population increases in all

three studied systems (freshwater slope = 0.005, CI = 0.002 to 0.01; marine slope =

0.004, CI = 0.002 to 0.01; terrestrial slope = 0.003, CI = 0.001 to 0.005, Figure 2a-b,

Table S4). In freshwater and terrestrial habitats, there was a bimodal distribution of

population change (Hartigans' dip test, D = 0.04, p < 0.01, Figure S5a). Amphibian

populations were more likely to decline (slope = -0.01, CI = -0.02 to -0.005), whereas

birds, mammals and reptiles were more likely to increase (slope = 0.004, CI = 0.003

to 0.01; slope = 0.01, CI = 0.01 to 0.01; slope = 0.02, CI = 0.01 to 0.02). Bony fish

were not more likely to increase or decline (slope = -0.001, CI = -0.004 to 0.002). The

distribution of *mu* values in bird populations was bimodal (Hartigans' dip test, D =

0.04, p < 0.01, Figure S5b).

Figure 4. Distribution of records across (plot a) freshwater, marine and terrestrial systems and (plot b) bony fish, amphibians, birds, mammals and

reptiles, monitoring duration (plots c and d) and their respective population

trend (*mu*) effect sizes (plots e and f). Results include 9284 populations from 2084

species. Effect sizes are population trend (*mu*) calculated using state-space and

linear models, and population trend (*mu*) weighted by *tau*, the observation error

estimate from the state-space models (plots e and f). The numbers on the x axis

show the sample size for each taxa category (plots e and f). Error bars show 95%

credible intervals (plots e and f). See Table S3 for full model outputs.

Population fluctuations across systems, taxa, Red List status and rarity metrics

Global population fluctuations (variation in population abundance over time) did not

show distinct geographic patterning (Figure 5a-b). Populations that experienced more

fluctuations had a smaller mean population size (slope = -0.008, CI = -0.01 to -0.004)

and were more likely to be generalists in their habitat preferences (slope = 0.005, CI

= 0.001 to 0.01, Figure 5c). On a global scale, population fluctuations were more

common with increasing Red List status (LC: slope = 0.11. CI = 0.11 to 0.12; CR:

slope = 0.17, CI = 0.15 to 0.19, Figure 5d, see Table S3 for all model outputs).

Across the three studied systems, terrestrial populations experienced lower

population fluctuations (slope = 0.10, CI = 0.10 to 0.11, Figure 5e). Across taxa,

fluctuations were most common for amphibian populations (slope = 0.22, CI = 0.20 to

0.23, Figure 5f).

Figure 5. Global population fluctuations in vertebrate species (plot a) across latitude (plot b), rarity metrics (UK scale, plot c), Red List status (global scale, plot d), systems (plot e) and taxa (plot f). Results include 508 populations from

237 species in the UK and 9284 populations from 2084 species globally. The

response variables in the models were: 1) the standard error around the slope

estimates of the linear model fits of abundance versus year, 2) half of the 95%

confidence interval around the *mu* value of population change, 3) half of the 95%

confidence interval around *mu* weighted by *tau*, the observation error estimate from

the state-space models, and 4) the standard deviation of the raw data for each

population time-series. Larger errors indicate more population fluctuations. Effect

size coefficients (c-f) were standardized by dividing the effect size by the standard

deviation of the corresponding input data. Error bars show 95% credible intervals (c-

f). See Table S3 for full model outputs.

Global population trends across monitoring duration

Monitoring duration was weakly positively related to vertebrate population trends,

with slightly more increasing populations found for longer duration studies (Figure S6,

Table S3).

Discussion

Across monitored vertebrate populations in the UK, rarity metrics (geographic range,

population size and habitat specificity) and species' Red List status did not explain

variation in population trends (Figures 1 and 2). There was no consistent pattern of

greater population declines with increasing Red List status in either the UK or globally

(Figures 1e,f and 2). There was no biogeographic patterning of population trends and

fluctuations, as populations increased, remained stable and declined across different

biomes on all continents (Figure 3). Among taxa, birds, mammals and reptiles

increased, bony fish did not experience net directional change and amphibians were

the only taxon with overall declines, as well as the taxon that experienced the most

population fluctuations (Figures 4-5). Fluctuations were more likely for species with

smaller population sizes and broader habitat preferences (Figure 5). Our findings do

not support the application of rarity metrics as generalizable predictors of population

trends in well-monitored vertebrate populations, but rare species were nevertheless

more likely to fluctuate, potentially increasing their extinction risk.

Population change across rarity metrics and species' Red List status

The power of rarity metrics and Red List status to explain population trends could be

mediated by the type of threats to which populations are exposed (Schemske *et al.*

1994), as well as whether species are naturally rare, or have become rare due to

external drivers in recent time (de Lange & Norton 2004; Robbirt *et al.* 2006; Harrison

et al. 2008). Geographic range, population size and habitat specificity may be of

lesser importance in human-dominated landscapes where threats such as habitat

change, pollution and exploitation are more prevalent (Newmark 1996; Purvis *et al.*

2000; O'Grady *et al.* 2004; Brook *et al.* 2006, Sullivan *et al.* 2016 but see Owens &

Bennett 2000 and Julliard *et al.* 2004). For UK vertebrate populations, the lack of

rarity effects on population trends (Figure 1) could be related to urbanization,

intensive land management and to the high investment in conservation (Keith *et al.*

2015), which may be contributing to the observed increases in species classified as

near threatened by the IUCN (Figure 1f). Populations with smaller mean population

size experienced more fluctuations (Figure 5c), which is in line with population

dynamics theory, suggesting that even though rarity metrics did not predict

population trends, they might still be good indicators of extinction risk, as greater

fluctuations could in time lead to extinction (Lande 1993; Melbourne & Hastings

2008).

Global population change across Red List status

Across all Red List status categories, there was no consistent pattern of net declines

with increasing threat (Figure 2). Populations from vulnerable and endangered

species did not experience directional change, suggesting that local scale

conservation actions might have stabilized populations for species which are

declining on a global scale (Saterson *et al.* 2004; Vickery *et al.* 2004; Gillings *et al.*

2005; Hoffmann *et al.* 2010). The threats and mitigation actions to which populations

with more threatened Red List statuses are exposed on a local scale are likely

heterogeneous across different biomes (Pereira *et al.* 2012; Maxwell *et al.* 2016). The

finding that populations from species with higher Red List status were more likely to

fluctuate (Figure 5d) relates to one of the Red List criteria – number of breeding

individuals (Mace *et al.* 2008). Because of Allee effects, extinction risk is higher when

there are very few breeding individuals (Dennis *et al.* 2016; Sun 2016). Thus,

although we did not find evidence to support the hypothesis that species with a

higher Red List status are more likely to be declining, those populations did

experience higher population fluctuations, suggesting that species of concern might

still have a higher risk of extinction.

Global population change across latitude, biomes and systems

Global vertebrate population trends varied among taxa but did not show distinct

geographic patterning (Figures 3-4). Some biomes like Mediterranean forests were

more likely to include increasing populations, whilst populations in other biomes did

not experience directional population change (Figure 3 and Table S3). Similar

heterogeneity in the patterns of population trends across habitats has been

documented in the Netherlands (van Strien *et al.* 2016) and in temperate North

America and Europe (Leung *et al.* 2017) also using the LPD, suggesting that local

scale environmental factors or proximity to anthropogenic drivers of change may be

better predictors of population trends (Brook *et al.* 2006; Fritz *et al.* 2009, but see

Phillips *et al.* 2017). The lack of distinct geographic patterning of population

fluctuations around the world (Figure 5a, b and e) suggests that these well-monitored

vertebrate populations are likely influenced by both stochastic and deterministic

processes (Kareiva 1990; Lundberg *et al.* 2000). Our findings go beyond previous

studies (Ceballos *et al.* 2017; Davidson *et al.* 2017) by quantifying the full spectrum of

population change, including positive, negative and stable trends and fluctuations

(Figures 3-5), which is critical for disentangling the pathways from population change

to biodiversity change (Primack *et al.* 2018; Yoccoz *et al.* 2018).

Global population change across taxa

Taxon-specific factors including reproductive strategy, trophic level, generation time,

life history traits (Purvis *et al.* 2000; Cardillo *et al.* 2004), or sampling effects

(Magurran *et al.* 2010) could explain why some taxa were more likely to increase

(birds, mammals and reptiles), whereas others were more likely to decline

(amphibians) or remain stable (bony fish), (Figure 4 and Table S3). Amphibians were

the only taxa to experience net population declines as well as greater population

fluctuations among monitored vertebrate species in the LPD (Figure 4f and 5f). The

global amphibian decline (Houlahan *et al.* 2000), also demonstrated with this dataset

at the temperate North American and European scale (Leung *et al.* 2017), has been

linked to the spread of a fungal disease (chytrid fungus, *Batrachochytrium*

dendrobatidis), facilitated by global warming (Pounds *et al.* 2006), as well as habitat

loss and Allee effects in small populations (Green 2003).

Study limitations

Our study is limited by three main factors: 1) the gaps in global monitoring of

populations, 2) challenges in estimating geographic range and habitat specificity, and

3) overrepresentation of common species. The LPD under-represents populations

outside of Europe and North America (Pereira & Cooper 2006; Proença *et al.* 2016;

McRae *et al.* 2017) and includes different species abundance measures across

populations that have been monitored for different durations (Figures S1, S2 and S6).

The GBIF database includes sub-optimal representation of some species and

occurrence outliers (Yesson *et al.* 2007; Boakes *et al.* 2010), despite being the

largest international open source database of species locality data (Yesson *et al.*

2007). In addition, we used a static measure of geographic range, which does not

account for changes in species distributions over time. For example, species could

naturally have a small range or the small range size could be due to historic habitat

loss (Magurran & Henderson 2003; Prendergast *et al.* 1993). In addition, UK

populations included in the LPD are predominantly from species with wide

geographic ranges (Table S4). The relationship between population change and

geographic range could differ for highly endemic species with restricted distribution

that are not monitored or included in the LPD dataset.

Conclusion

In summary, rarity metrics did not explain vertebrate population trends in the UK and

we did not find a consistent pattern of net declines with increasing Red List status in

the UK and globally. We found taxonomic but not geographic patterning of population

change around the world. Vertebrate populations experienced different magnitudes

of net change and population fluctuations, which in the future may lead to shifts in

community composition as some species become locally extinct and others expand

into new habitats. Our findings caution the use of rarity metrics for the prediction of

future global population trends, but suggest that such metrics are nevertheless

indicators of population fluctuations, which might ultimately link to increased species

extinction risk.

Acknowledgements

We thank the WWF and the RZSL for compiling and maintaining the Living Planet

Database. We thank Dmitry Schigel and Jan Legind at GBIF for providing access to a

direct download link for the UK species occurrence data. We are grateful to the

Critical Thinking in Ecology and Environmental Sciences 2015-16 tutorial group at the

University of Edinburgh that helped us conceive the idea for the analysis including

Hannah Louise Stevens, Ashlie Nithsdale, Lisa Kopsieker and Claudia Ardrey. We

thank Anne Magurran, Albert Phillimore and Mark Vellend for providing very helpful

feedback on the manuscript. Funding for this research was provided by the NERC

doctoral training partnership grant (NE/L002558/1) at the University of Edinburgh and

the Carnegie Trust for the Universities of Scotland.

References

Agardy, T. (2005). Global marine conservation policy versus site-level

implementation. *Mar. Ecol. Prog. Ser.*, 300, 242–248.

Batt, R.D., Morley, J.W., Selden, R.L., Tingley, M.W. & Pinsky, M.L. (2017). Gradual

changes in range size accompany long-term trends in species richness. *Ecol.*

Lett., 20, 1148–1157.

Boakes, E.H., McGowan, P.J., Fuller, R.A., Chang-qing, D., Clark, N.E., O'Connor,

K., *et al.* (2010). Distorted views of biodiversity: spatial and temporal bias in

species occurrence data. *PLoS Biol.*, 8, e1000385.

Brook, B.W., Traill, L.W. & Bradshaw, C.J.A. (2006). Minimum viable population

sizes and global extinction risk are unrelated: MVP is unrelated to global

extinction risk. *Ecol. Lett.*, 9, 375–382.

Cardillo, M., Mace, G.M., Jones, K.E., Bielby, J., Bininda-Emonds, O.R.P., Sechrest,

W., *et al.* (n.d.). Multiple Causes of High Extinction Risk in Large Mammal

Species. *Science*, 309, 1239–1241.

Cardillo, M., Purvis, A., Sechrest, W., Gittleman, J.L., Bielby, J. & Mace, G.M. (2004).

Human Population Density and Extinction Risk in the World's Carnivores.

PLoS Biol., 2, e197.

Ceballos, G., Ehrlich, P.R. & Dirzo, R. (2017). Biological annihilation via the ongoing

sixth mass extinction signaled by vertebrate population losses and declines.

Proc. Natl. Acad. Sci., 201704949.

Collen, B., Loh, J., Whitmee, S., McRAE, L., Amin, R. & Baillie, J.E.M. (2009).

Monitoring Change in Vertebrate Abundance: the Living Planet Index.

Conserv. Biol., 23, 317–327.

Davidson, A.D., Shoemaker, K.T., Weinstein, B., Costa, G.C., Brooks, T.M.,

Ceballos, G., *et al.* (2017). Geography of current and future global mammal

extinction risk. *PLOS ONE*, 12, e0186934.

Dennis, B., Assas, L., Elaydi, S., Kwessi, E. & Livadiotis, G. (2016). Allee effects and

resilience in stochastic populations. *Theor. Ecol.*, 9, 323–335.

Dennis, B., Ponciano, J.M., Lele, S.R., Taper, M.L. & Staples, D.F. (2006).

Estimating density dependence, process noise, and observation error. *Ecol.*

Monogr., 76, 323–341.

Ehrlén, J. & Morris, W.F. (2015). Predicting changes in the distribution and

abundance of species under environmental change. *Ecol. Lett.*, 18, 303–314.

Fritz, S.A., Bininda-Emonds, O.R.P. & Purvis, A. (2009). Geographical variation in

predictors of mammalian extinction risk: big is bad, but only in the tropics.

Ecol. Lett., 12, 538–549.

Gaston, K. & Fuller, R. (2008). Commonness, population depletion and conservation

biology. *Trends Ecol. Evol.*, 23, 14–19.

Gaston, K.J. & Blackburn, T.M. (1995). Birds, body size and the threat of extinction.

Philos. Trans. R. Soc. Lond. B Biol. Sci., 347, 205–212.

GBIF. (2017). Biodiversity occurrence data. http://www.gbif.org/ Downloaded Feb

2016 - Apr 2017.

Gillings, S., Newson, S.E., Noble, D.G. & Vickery, J.A. (2005). Winter availability of

cereal stubbles attracts declining farmland birds and positively influences

breeding population trends. *Proc. R. Soc. B Biol. Sci.*, 272, 733–739.

Gilroy, J.J., Gill, J.A., Butchart, S.H.M., Jones, V.R. & Franco, A.M.A. (2016).

Migratory diversity predicts population declines in birds. *Ecol. Lett.*, 19, 308–

317.

Green, D.M. (2003). The ecology of extinction: population fluctuation and decline in

amphibians. *Biol. Conserv.*, 111, 331–343.

Guerrero, A.M., McALLISTER, R.R.J., Corcoran, J. & Wilson, K.A. (2013). Scale

Mismatches, Conservation Planning, and the Value of Social-Network

Analyses: Scale Mismatches and Conservation Planning. *Conserv. Biol.*, 27,

35–44.

Gunn, A. (1980). Why should we care about rare species? *Environ. Ethics*, 2, 17–37.

Hadfield, J.D. (2010). MCMC Methods for Multi-Response Generalized Linear Mixed

Models: The MCMCglmm R Package. *J. Stat. Softw.*, 33, 1–22.

Hanski, I. (1998). Metapopulation dynamics. *Nature*, 396, 41.

Harrison, S., Viers, J.H., Thorne, J.H. & Grace, J.B. (2008). Favorable environments

and the persistence of naturally rare species. *Conserv. Lett.*, 1, 65–74.

Hefley, T.J., Hooten, M.B., Drake, J.M., Russell, R.E. & Walsh, D.P. (2016). When

can the cause of a population decline be determined? *Ecol. Lett.*, 19, 1353–

1362.

Hijmans, R.J. (2016). geosphere: Spherical Trigonometry. R package version 1.5-5.

https://CRAN.R-project.org/package=geosphere.

Hoffmann, M., Hilton-Taylor, C., Angulo, A., Bohm, M., Brooks, T.M., Butchart,

S.H.M., *et al.* (2010). The Impact of Conservation on the Status of the

World's Vertebrates. *Science*, 330, 1503–1509.

Houlahan, J.E., Findlay, C.S., Schmidt, B.R., Meyer, A.H. & Kuzmin, S.L. (2000).

Quantitative evidence for global amphibian population declines. *Nature*, 404,

752–755.

Humbert, J.-Y., Scott Mills, L., Horne, J.S. & Dennis, B. (2009). A better way to

estimate population trends. *Oikos*, 118, 1940–1946.

Hutchings, J.A. & Reynolds, J.D. (2004). Marine fish population collapses:

consequences for recovery and extinction risk. *AIBS Bull.*, 54, 297–309.

IUCN. (2017). Conservation status data. http://www.iucnredlist.org/ Downloaded Jan-

April 2017.

Jones, K.E., Purvis, A. & Gittleman, J.L. (2003). Biological Correlates of Extinction

Risk in Bats. *Am. Nat.*, 161, 601–614.

Julliard, R., Jiguet, F. & Couvet, D. (2004). Common birds facing global changes:

what makes a species at risk? *Glob. Change Biol.*, 10, 148–154.

Kareiva, P. (1990). Population dynamics in spatially complex environments: theory

and data. *Phil Trans R Soc Lond B*, 330, 175–190.

Keith, D., Akçakaya, H.R., Butchart, S.H.M., Collen, B., Dulvy, N.K., Holmes, E.E., *et*

al. (2015). Temporal correlations in population trends: Conservation

implications from time-series analysis of diverse animal taxa. *Biol. Conserv.*,

192, 247–257.

Knape, J., Jonzén, N. & Sköld, M. (2011). On observation distributions for state

space models of population survey data: Observation models for population

data. *J. Anim. Ecol.*, 80, 1269–1277.

Lande, R. (1993). Risks of population extinction from demographic and

environmental stochasticity and random catastrophes. *Am. Nat.*, 142, 911–

927.

de Lange, P.J. & Norton, D.A. (2004). The ecology and conservation of *Kunzea*

sinclairii (*Myrtaceae*), a naturally rare plant of rhyolitic rock outcrops. *Biol.*

Conserv., 117, 49–59.

Lawson, C.R., Vindenes, Y., Bailey, L. & van de Pol, M. (2015). Environmental

variation and population responses to global change. *Ecol. Lett.,* 18, 724–

736.

Leung, B., Greenberg, D.A. & Green, D.M. (2017). Trends in mean growth and

stability in temperate vertebrate populations. *Divers. Distrib.*, 23, 1372–1380.

Longton, R.E. & Hedderson, T.A. (2000). What are rare species and why conserve

them? *Lindbergia*, 25, 53–61.

LPI. (2016). Living Planet Index database, 2016. www.livingplanetindex.org/

Downloaded February 2016.

Lundberg, P., Ranta, E., Ripa, J. & Kaitala, V. (2000). Population variability in space

and time. *Trends Ecol. Evol.*, 15, 460–464.

Mace, G.M., Collar, N.J., Gaston, K.J., Hilton-Taylor, C., Akçakaya, H.R., Leader-

Williams, N., *et al.* (2008). Quantification of Extinction Risk: IUCN's System

for Classifying Threatened Species. *Conserv. Biol.*, 22, 1424–1442.

Magurran, A. (2013). *Measuring biological diversity*. John Wiley & Sons.

Magurran, A.E., Baillie, S.R., Buckland, S.T., Dick, J.M., Elston, D.A., Scott, E.M., *et*

al. (2010). Long-term datasets in biodiversity research and monitoring:

assessing change in ecological communities through time. *Trends Ecol.*

Evol., 25, 574–582.

Magurran, A.E. & Henderson, P.A. (2003). Explaining the excess of rare species in

natural species abundance distributions. *Nature*, 422, 714.

Manne, L.L. & Pimm, S.L. (2001). Beyond eight forms of rarity: which species are

threatened and which will be next? *Anim. Conserv.*, 4, 221–229.

Maxwell, S.L., Fuller, R.A., Brooks, T.M. & Watson, J.E. (2016). Biodiversity: The

ravages of guns, nets and bulldozers. *Nature*, 536, 143–145.

McKinney, M.L. (1997). Extinction vulnerability and selectivity: combining ecological

and paleontological views. *Annu. Rev. Ecol. Syst.*, 28, 495–516.

McRae, L., Deinet, S. & Freeman, R. (2017). The Diversity-Weighted Living Planet

Index: Controlling for Taxonomic Bias in a Global Biodiversity Indicator.

PLOS ONE, 12, e0169156.

Melbourne, B.A. & Hastings, A. (2008). Extinction risk depends strongly on factors

contributing to stochasticity. *Nature*, 454, 100–103.

Mills, M., Pressey, R.L., Weeks, R., Foale, S. & Ban, N.C. (2010). A mismatch of

scales: challenges in planning for implementation of marine protected areas

in the Coral Triangle: Mismatch of scales in conservation planning. *Conserv.*

Lett., 3, 291–303.

Newmark, W.D. (1996). Insularization of Tanzanian parks and the local extinction of

large mammals. *Conserv. Biol.*, 10, 1549–1556.

O'Grady, J.J., Reed, D.H., Brook, B.W. & Frankham, R. (2004). What are the best

correlates of predicted extinction risk? *Biol. Conserv.*, 118, 513–520.

Olden, J.D., Hogan, Z.S. & Zanden, M.J.V. (2007). Small fish, big fish, red fish, blue

fish: size-biased extinction risk of the world's freshwater and marine fishes.

Glob. Ecol. Biogeogr., 16, 694–701.

Olson, D.M. & Dinerstein, E. (2002). The Global 200: Priority Ecoregions for Global

Conservation. *Ann. Mo. Bot. Gard.*, 89, 199.

Owens, I.P. & Bennett, P.M. (2000). Ecological basis of extinction risk in birds:

habitat loss versus human persecution and introduced predators. *Proc. Natl.*

Acad. Sci., 97, 12144–12148.

Pedersen, M.W., Berg, C.W., Thygesen, U.H., Nielsen, A. & Madsen, H. (2011).

Estimation methods for nonlinear state-space models in ecology. *Ecol.*

Model., 222, 1394–1400.

Pereira, H.M. & Cooper, D.H. (2006). Towards the global monitoring of biodiversity

change. *Trends Ecol. Evol.*, 21, 123–129.

Pereira, H.M., Navarro, L.M. & Martins, I.S. (2012). Global Biodiversity Change: The

Bad, the Good, and the Unknown. *Annu. Rev. Environ. Resour.*, 37, 25–50.

Phillips, H.R.P., Newbold, T. & Purvis, A. (2017a). Land-use effects on local

biodiversity in tropical forests vary between continents. *Biodivers. Conserv.*,

26, 2251–2270.

Phillips, H.R.P., Newbold, T. & Purvis, A. (2017b). Land-use effects on local

biodiversity in tropical forests vary between continents. *Biodivers. Conserv.*,

26, 2251–2270.

Pigott, C.D. & Walters, S.M. (1977). The scientific basis of practical conservation:

aims and methods of conservation. In: *Proc. R. Soc. Lond. B*. The Royal

Society, pp. 59–68.

Pounds, A.J., Bustamante, M.R., Coloma, L.A., Consuegra, J.A., Fogden, M.P.L.,

Foster, P.N., *et al.* (2006). Widespread amphibian extinctions from epidemic

disease driven by global warming. *Nature*, 439, 161–167.

Povilitis, T. (2001). Toward a robust natural imperative for conservation. *Conserv.*

Biol., 15, 533–535.

Prendergast, J.R., Quinn, R.M., Lawton, J.H., Eversham, B.C. & Gibbons, D.W.

(1993). Rare species, the coincidence of diversity hotspots and conservation

strategies. *Nature*, 365, 335–337.

Pressey, R.L., Cabeza, M., Watts, M.E., Cowling, R.M. & Wilson, K.A. (2007).

Conservation planning in a changing world. *Trends Ecol. Evol.*, 22, 583–592.

Primack, R.B., Miller-Rushing, A.J., Corlett, R.T., Devictor, V., Johns, D.M., Loyola,

R., *et al.* (2018). Biodiversity gains? The debate on changes in local- vs

global-scale species richness. *Biol. Conserv.*, 219, A1–A3.

Proença, V., Martin, L.J., Pereira, H.M., Fernandez, M., McRae, L., Belnap, J., *et al.*

(2016). Global biodiversity monitoring: From data sources to Essential

Biodiversity Variables. *Biol. Conserv.*

Purvis, A., Gittleman, J.L., Cowlishaw, G. & Mace, G.M. (2000). Predicting extinction

risk in declining species. *Proc. R. Soc. B Biol. Sci.*, 267, 1947–1952.

R Core Team. (2017). R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria.

Rabinowitz, D. (1981). Seven forms of rarity. In: *The biological aspects of rare plants*

conservation. pp. 205–217.

Ripple, W.J., Wolf, C., Newsome, T. M., Hoffmann, M., Wirsing, A. J. & McCauley, D.

J. (2017). Extinction risk is most acute for the world's largest and smallest

vertebrates. *114*, 10678–10683.

Robbirt, K.M., Roberts, D.L. & Hawkins, J.A. (2006). Comparing IUCN and

Probabilistic Assessments of Threat: Do IUCN Red List Criteria Conflate

Rarity and Threat? *Biodivers. Conserv.*, 15, 1903–1912.

Rueda-Cediel, P., Anderson, K.E., Regan, T.J. & Regan, H.M. (2018). Effects of

uncertainty and variability on population declines and IUCN Red List

classifications. *Conserv. Biol.*

Saterson, K.A., Christensen, N.L., Jackson, R.B., Kramer, R.A., Pimm, S.L., Smith,

M.D., *et al.* (2004). Disconnects in Evaluating the Relative Effectiveness of

Conservation Strategies. *Conserv. Biol.*, 18, 597–599.

Schemske, D.W., Husband, B.C., Ruckelshaus, M.H., Goodwillie, C., Parker, I.M. &

Bishop, J.G. (1994). Evaluating Approaches to the Conservation of Rare and

Endangered Plants. *Ecology*, 75, 584–606.

Shultz, S., B. Bradbury, R., L. Evans, K., D. Gregory, R. & M. Blackburn, T. (2005).

Brain size and resource specialization predict long-term population trends in

British birds. *Proc. R. Soc. B Biol. Sci.*, 272, 2305–2311.

van Strien, A.J., Meyling, A.W.G., Herder, J.E., Hollander, H., Kalkman, V.J., Poot,

M.J.M., *et al.* (2016). Modest recovery of biodiversity in a western European

country: The Living Planet Index for the Netherlands. *Biol. Conserv.*, 200,

44–50.

Sun, G.-Q. (2016). Mathematical modeling of population dynamics with Allee effect.

Nonlinear Dyn., 85, 1–12.

Vickery, J.A., Bradbury, R.B., Henderson, I.G., Eaton, M.A. & Grice, P.V. (2004). The

role of agri-environment schemes and farm management practices in

reversing the decline of farmland birds in England. *Biol. Conserv.*, 119, 19–

39.

Yesson, C., Brewer, P.W., Sutton, T., Caithness, N., Pahwa, J.S., Burgess, M., *et al.*

(2007). How Global Is the Global Biodiversity Information Facility? *PLoS*

ONE, 2, e1124.

Yoccoz, N.G., Ellingsen, K.E. & Tveraa, T. (2018). Biodiversity may wax or wane

depending on metrics or taxa. *Proc. Natl. Acad. Sci.*, 115, 1681–1683.

 Supplementary Information

 Table S1. Number of species and populations included in analyses.

Scale	Analysis	Number of species	Number of populations	Number of species with >3 populations
UK	Geographic range	167	381	32
	Population size	112	253	19
	Habitat specificity	150	302	29
	Red List status	153	364	31
Global	System	2074	9284	834
	Biome	2074	9284	834
	Taxa			
	- *Actinopterygii*	544	1626	151
	- *Amphibia*	78	193	21
	- *Aves*	968	5852	538
	- *Mammalia*	306	1158	94
	- *Reptilia*	110	322	16
	Red List status	1702	8064	708
	Latitude	2074	9284	834
	Duration	2074	9284	834

759 **Table S2. Key used to determine habitat specificity for 194 species with**

760 **populations in the UK in the LPD.** We extracted the habitats in which each species

761 occurs from their IUCN profiles (http://www.iucnredlist.org/).

Habitat	Considered to be the same as:	Considered to be different to:
Rural park	Suburban park, urban park, rural garden, suburban garden, urban garden	
Lake	Big lake, small lake, pond, pool, dam, oxbow lake, reservoir	
Bog	Swamp, bogland	Lagoon
Coastal cliff	Island cliff	
Shingle beach	Pebble beach, rock beach	Sandy beach
Stream	River	Weir
Fruit Tree Plantation	Fruit Garden, orchard	
Thicket	Copse, grove, small stand	
Forest		Woodland
Glade	Forest Clearing	
Broadleaf	Deciduous	
Urban	Suburban	
River margin	Various types of river margins	
Tidal Creek		Estuary
Harbor	Dock, jetty, pier	
Bush lands	Shrublands	
Irrigation channel	Ditch	
Heath	Moorland	
Sandy beach	Spit, dune	Shingle beach, pebble beach, rock beach
Crag	Rocky outcrop, cliff, rocky slope	
Marsh	Wet meadow	
Islet	Island	

Monitoring duration (years)

764 **Figure S1. Distribution of monitoring duration across (a) freshwater, (b) marine**

765 **and (c) terrestrial systems. Results include 9284 populations from 2084**

766 **species.**

Figure S2. Distribution of monitoring duration across all monitored vertebrate populations (plot a), bony fish (plot b), amphibians (plot c), birds (plot d), mammals (plot e) and reptiles (plot f). See Table S1 for sample sizes of each group.

Figure S3. Population change (*mu* values from state-space models) across the five most commonly used units in the subset of the LPD we analyzed. Numbers on the x axis correspond to sample size in each category.

Figure S4. Geographic distribution of vertebrate population change in the UK, from state-space models (plot a) and linear models (plot b). Results include 508 populations from 237 species in the UK.

783 **Figure. S5. Distribution of population change (*mu*) values from state-space**

784 **models of vertebrate abundance over time across systems (plot a) and across**

785 **taxa (plot b).** See Table S1 for sample sizes.

Figure S6. Vertebrate population trends across monitoring duration. Results are

for 9284 populations from 2084 species. Population trends (*mu*) were estimated for

all populations monitored for more than five time points using state-space models

(plot a) and linear models (plot b). Population fluctuations (plot c) were estimated

using the confidence interval (half of the 95% confidence interval) of the *mu* values

(see also Figure 5). Error bars show 95% confidence intervals.

Figure S7. Geographic distribution of population change, calculated using state-space models (plot a), linear models (plot b), and using the confidence interval (half of the 95% confidence interval) of the *mu* values (plot c). Results include 9284 populations from 2084 species. Larger confidence intervals on (c) correspond to populations that have experienced more fluctuations.

Table S3. Model outputs from UK and global analyses. Sigma is the overall model

residual variance. Net population change is estimated using *mu* values derived from

state-space models of population abundance versus time and slopes of linear models

of population abundance versus time. The weighted *mu* models included *mu* as a

response variable, weighted by *tau*, the observation error estimate derived from the

state-space models. The fluctuation models included half of the 95% confidence

interval around the *mu* value of population change, the standard error around the

slopes of linear models of population abundance versus time, and the standard

deviation of the raw time-series data for each population. Larger confidence intervals

correspond to populations that experienced more fluctuations. The weighted

fluctuation models were weighted by *tau*, the observation error estimate derived from

the state-space models.

Model name	Variable	Posterior mean	Lower 95% CI	Upper 95% CI	Effective sample size	pMCMC	Effect
			UK scale analyses				
Geographic range - mu	(Intercept)	0.02	-0.03	0.06	18,000	0.45	fixed
	log(km2_range)	-0.0004	-0.003	0.002	18,000	0.79	fixed
	sigma	0.003	0.002	0.003	17,265		residual
Geographic range - slope	(Intercept)	0.02	-0.02	0.07	18,937	0.32	fixed
	log(km2_range)	-0.001	-0.004	0.002	19,025	0.52	fixed
	sigma	0.002	0.002	0.003	17,450		residual
Geographic range - weighted	(Intercept)	0.02	-0.03	0.07	18,000	0.47	fixed

	log(km2_range)	-0.0004	-0.003	0.002	18,000	0.76	fixed
	sigma	0.003	0.002	0.003	18,000		residual
Geographic range – fluctuations (SE)	(Intercept)	0.12	-0.05	0.29	18,000	0.17	0.12
	log(km2_range)	0.004	-0.01	0.01	18,000	0.42	0.004
	sigma	0.03	0.03	0.04	18,000		0.03
Geographic range – fluctuations (CI)	(Intercept)	0.25	0.10	0.39	18,000	0.001	fixed
	log(km2_range)	-0.004	-0.01	0.004	18,000	0.37	fixed
	sigma	0.03	0.02	0.03	18,000		residual
Geographic range – fluctuations weighted (CI)	(Intercept)	0.25	0.10	0.40	18,000	0.001	fixed
	log(km2_range)	-0.004	-0.01	0.005	18,000	0.37	fixed
	sigma	0.03	0.02	0.03	18,313		residual
Geographic range – fluctuations (SD)	(Intercept)	0.35	0.23	0.48	18,000	0.0001	0.35
	log(km2_range)	-0.001	-0.01	0.01	18,000	0.84	-0.001
	sigma	0.02	0.02	0.02	18,000		0.02
Mean population size - mu	(Intercept)	0.004	-0.01	0.02	18,549	0.53	fixed
	log(meanpop)	0.001	-0.001	0.002	18,000	0.41	fixed
	sigma	0.003	0.002	0.003	18,000		residual

Mean population size - slope	(Intercept)	0.01	-0.003	0.02	18,000	0.17	fixed
	log(meanpop)	-0.0003	-0.002	0.001	18,000	0.74	fixed
	sigma	0.003	0.002	0.003	18,000		residual
Mean population size - weighted	(Intercept)	0.001	-0.01	0.01	18,000	0.91	fixed
	log(meanpop)	0.001	-0.001	0.003	18,000	0.26	fixed
	sigma	0.003	0.002	0.003	18,000		residual
Mean population size – fluctuations (SE)	(Intercept)	0.34	0.30	0.38	17,499	0.0001	fixed
	log(meanpop)	-0.02	-0.03	-0.02	18,000	0.0001	fixed
	sigma	0.03	0.03	0.04	18,000		residual
Mean population size – fluctuations (CI)	(Intercept)	0.26	0.23	0.30	18,000	0.0001	fixed
	log(meanpop)	-0.01	-0.02	-0.01	18,000	0.0001	fixed
	sigma	0.03	0.02	0.03	18,000		residual
Mean population size – fluctuations weighted (CI)	(Intercept)	0.27	0.23	0.30	17,575	0.0001	fixed
	log(meanpop)	-0.01	-0.02	-0.01	18,000	0.0001	fixed
	sigma	0.03	0.02	0.03	18,000		residual

Mean population size – fluctuations (SD)	(Intercept)	0.48	0.45	0.52	18,000	0.0001	fixed
	log(meanpop)	-0.01	-0.02	-0.01	18,000	0.0001	fixed
	sigma	0.03	0.02	0.03	17,812		residual
Habitat specificity - mu	(Intercept)	0.02	0.01	0.03	18,000	0.002	fixed
	hab_spec	-0.001	-0.003	0.0005	18,000	0.16	fixed
	sigma	0.003	0.003	0.003	18,000		residual
Habitat specificity - slope	(Intercept)	0.01	0.001	0.02	18,000	0.04	fixed
	hab_spec	-0.001	-0.002	0.001	18,172	0.38	fixed
	sigma	0.003	0.002	0.003	18,412		residual
Habitat specificity - weighted	(Intercept)	0.02	0.01	0.03	18,000	0.003	fixed
	hab_spec	-0.001	-0.003	0.0004	18,000	0.15	fixed
	sigma	0.003	0.002	0.003	18,000		residual
Habitat specificity – fluctuations (SE)	(Intercept)	0.18	0.14	0.22	17,681	0.0001	fixed
	hab_spec	0.01	-0.0002	0.01	18,000	0.05	fixed
	sigma	0.04	0.03	0.04	18,510		residual
Habitat specificity – fluctuations (CI)	(Intercept)	0.16	0.13	0.20	18,000	0.0001	fixed

	hab_spec	0.01	0.001	0.01	18,553	0.03	fixed
	sigma	0.03	0.03	0.03	18,000		residual
Habitat specificity – fluctuations weighted (CI)	(Intercept)	0.16	0.13	0.20	18,000	0.0001	fixed
	hab_spec	0.01	0.0004	0.01	18,000	0.04	fixed
	sigma	0.03	0.03	0.03	18,000		residual
Habitat specificity – fluctuations (SD)	(Intercept)	0.36	0.33	0.39	18,000	0.0001	fixed
	hab_spec	0.004	0.0001	0.01	18,000	0.05	fixed
	sigma	0.02	0.02	0.03	18,503		residual
Red List status - mu	Least concern	0.01	0.01	0.02	18,000	0.0001	fixed
	Near threatened	0.03	0.01	0.05	18,000	0.001	fixed
	Vulnerable	-0.01	-0.03	0.01	18,000	0.48	fixed
	Endangered	-0.04	-0.14	0.06	18,000	0.47	fixed
	Critically endangered	0.03	-0.05	0.10	16,574	0.45	fixed
	sigma	0.003	0.002	0.003	18,000		residual
Red List status - slope	Least concern	0.01	0.004	0.01	18,000	0.001	fixed
	Near threatened	0.02	0.01	0.04	18,000	0.005	fixed
	Vulnerable	-0.01	-0.03	0.01	18,000	0.37	fixed
	Endangered	-0.01	-0.10	0.09	18,000	0.91	fixed

	Critically endangered	0.03	-0.04	0.09	18,000	0.43	fixed
	sigma	0.002	0.002	0.003	18,000		residual
Red List status - weighted	Least concern	0.01	0.005	0.02	18,000	0.0002	fixed
	Near threatened	0.04	0.02	0.05	17,614	0.0001	fixed
	Vulnerable	-0.01	-0.03	0.01	18,000	0.46	fixed
	Endangered	-0.04	-0.14	0.06	18,000	0.45	fixed
	Critically endangered	0.03	-0.05	0.11	18,000	0.49	fixed
	sigma	0.003	0.002	0.003	18,000		residual
Red List status – fluctuations (SE)	Least concern	0.19	0.17	0.21	18,523	0.0001	fixed
	Near threatened	0.28	0.22	0.35	17,404	0.0001	fixed
	Vulnerable	0.16	0.09	0.24	18,000	0.0002	fixed
	Endangered	0.18	-0.19	0.54	18,000	0.33	fixed
	Critically endangered	0.09	-0.17	0.34	17,758	0.51	fixed
	sigma	0.03	0.03	0.04	18,000		residual
Red List status – fluctuations (CI)	Least concern	0.18	0.16	0.20	18,000	0.0001	fixed
	Near threatened	0.21	0.16	0.27	18,000	0.0001	fixed
	Vulnerable	0.16	0.09	0.22	18,000	0.0001	fixed
	Endangered	0.41	0.08	0.73	18,000	0.01	fixed

	Critically endangered	0.06	-0.18	0.29	18,000	0.64	fixed
	sigma	0.03	0.02	0.03	18,000		residual
Red List status – fluctuations weighted (CI)	Least concern	0.18	0.16	0.20	18,000	0.0001	fixed
	Near threatened	0.21	0.16	0.27	18,000	0.0001	fixed
	Vulnerable	0.16	0.09	0.23	18,000	0.0001	fixed
	Endangered	0.40	0.07	0.73	19,079	0.02	fixed
	Critically endangered	0.06	-0.18	0.29	18,000	0.63	fixed
	sigma	0.03	0.02	0.03	17,569		residual
Red List status – fluctuations (SD)	Least concern	0.38	0.37	0.40	18,000	0.0001	fixed
	Near threatened	0.37	0.32	0.43	18,000	0.0001	fixed
	Vulnerable	0.27	0.21	0.33	18,000	0.0001	fixed
	Endangered	0.24	-0.06	0.55	18,000	0.12	fixed
	Critically endangered	0.32	0.14	0.51	18,000	0.001	fixed
	sigma	0.02	0.02	0.03	18,043		residual
System - mu	Freshwater	0.01	-0.01	0.02	18,000	0.39	fixed
	Marine	0.004	-0.002	0.01	18,000	0.23	fixed
	Terrestrial	0.03	0.02	0.03	18,000	0.0001	fixed
	sigma	0.003	0.002	0.003	18,000		residual

System - slope	Freshwater	0.01	-0.002	0.02	18,000	0.11	fixed
	Marine	0.004	-0.002	0.01	18,395	0.22	fixed
	Terrestrial	0.02	0.01	0.03	18,000	0.0001	fixed
	sigma	0.002	0.002	0.003	17,258		residual
System - weighted	Freshwater	0.001	-0.01	0.01	18,000	0.86	fixed
	Marine	0.01	-0.002	0.01	18,000	0.15	fixed
	Terrestrial	0.02	0.02	0.03	18,000	0.0001	fixed
	sigma	0.002	0.002	0.003	18,000		residual
Taxa - mu	Actinopterygii	-0.01	-0.02	0.001	18,000	0.09	fixed
	Amphibia	-0.02	-0.05	0.01	18,000	0.15	fixed
	Aves	0.02	0.01	0.02	18,000	0.0001	fixed
	Mammalia	0.02	0.01	0.03	18,000	0.0001	fixed
	Reptilia	-0.02	-0.09	0.05	19,343	0.52	fixed
	sigma	0.003	0.002	0.003	18,000		residual
Taxa - slope	Actinopterygii	-0.01	-0.02	0.0004	18,000	0.07	fixed
	Amphibia	-0.02	-0.05	0.01	18,000	0.13	fixed
	Aves	0.01	0.004	0.02	18,000	0.001	fixed
	Mammalia	0.03	0.02	0.04	18,000	0.0001	fixed
	Reptilia	-0.04	-0.10	0.02	18,000	0.21	fixed
	sigma	0.002	0.002	0.002	18,000		residual
Taxa - weighted	Actinopterygii	-0.01	-0.02	-0.0003	18,000	0.05	fixed

	Amphibia	-0.04	-0.07	-0.002	18,000	0.04	fixed
	Aves	0.01	0.01	0.02	18,000	0.0001	fixed
	Mammalia	0.02	0.01	0.03	18,000	0.0001	fixed
	Reptilia	-0.05	-0.13	0.04	18,862	0.31	fixed
	sigma	0.002	0.002	0.003	18,000		residual
Global scale analyses							
System - mu	Freshwater	0.005	0.002	0.01	18,000	0.0001	fixed
	Marine	0.004	0.002	0.01	18,587	0.001	fixed
	Terrestrial	0.003	0.001	0.005	18,000	0.001	fixed
	sigma	0.003	0.003	0.004	18,000		residual
System - slope	Freshwater	0.005	0.003	0.006	17544	0.0001	fixed
	Marine	0.002	0.001	0.004	18,000	0.04	fixed
	Terrestrial	0.002	0.001	0.003	17,374	0.01	fixed
	sigma	0.002	0.002	0.003	18,175		residual
System - weighted	Freshwater	0.01	0.003	0.01	18,390	0.0001	fixed
	Marine	0.004	0.001	0.01	18,000	0.002	fixed
	Terrestrial	0.004	0.002	0.01	18,000	0.0004	fixed
	sigma	0.003	0.003	0.003	18,000		residual
System – fluctuations (SE)	Freshwater	0.12	0.12	0.13	18,000	0.0001	fixed
	Marine	0.12	0.11	0.12	17,506	0.0001	fixed
	Terrestrial	0.09	0.08	0.09	18,000	0.0001	fixed

	sigma	0.02	0.02	0.02	18,000		residual
System – fluctuations (CI)	Freshwater	0.13	0.13	0.14	18,000	0.0001	fixed
	Marine	0.13	0.13	0.14	18,000	0.0001	fixed
	Terrestrial	0.10	0.10	0.10	18,000	0.0001	fixed
	sigma	0.02	0.02	0.02	18,000		residual
System – fluctuations weighted (CI)	Freshwater	0.13	0.13	0.14	18,000	0.0001	fixed
	Marine	0.14	0.13	0.14	18,550	0.0001	fixed
	Terrestrial	0.10	0.09	0.10	18,590	0.0001	fixed
	sigma	0.02	0.02	0.02	18,000		residual
System – fluctuations (SD)	Freshwater	0.39	0.39	0.40	19,118	0.0001	fixed
	Marine	0.40	0.40	0.41	18,000	0.0001	fixed
	Terrestrial	0.37	0.37	0.38	18,136	0.0001	fixed
	sigma	0.02	0.02	0.02	18,000		residual
Biome - mu	Boreal forests/taiga	-0.001	-0.004	0.002	18,000	0.72	fixed
	Deserts and xeric shrublands	-0.01	-0.03	0.003	18,000	0.10	fixed
	Trop. and subtrop. grasslands savannas and shrublands	-0.01	-0.02	-0.001	17,414	0.03	fixed
	Large lakes	0.003	-0.004	0.01	18,665	0.43	fixed

	Mediterranean forests woodlands and scrub	0.02	0.01	0.02	17,081	0.0001	fixed
	Montane freshwaters	0.01	-0.01	0.04	17,409	0.34	fixed
	Montane grasslands and shrublands	0.02	0.002	0.04	18,000	0.03	fixed
	Polar freshwaters	0.01	0.001	0.01	18,002	0.02	fixed
	Polar seas	-0.002	-0.02	0.02	18,000	0.85	fixed
	Temperate forests	0.005	0.002	0.01	18,000	0.0003	fixed
	Temperate wetlands and rivers	0.005	0.002	0.01	18,000	0.001	fixed
	Temperate grasslands savannas and shrublands	-0.004	-0.01	0.002	16,766	0.22	fixed
	Tropical wetlands and rivers	0.01	-0.003	0.01	18,000	0.22	fixed
	Tropical and subtropical forests	0.02	0.01	0.03	16,634	0.0002	fixed
	Tropical coral	0.02	0.01	0.04	18,000	0.0001	fixed
	Tundra	0.01	-0.001	0.01	19,359	0.10	fixed
	Xeric freshwaters and endorheic basins	-0.002	-0.02	0.01	18,000	0.77	fixed
	sigma	0.003	0.003	0.003	18,000		residual
Biome - slope	Boreal forests/taiga	-0.001	-0.004	0.002	18,686	0.41	fixed

Deserts and xeric shrublands	-0.01	-0.03	0.002	18,000	0.08	fixed
Trop. and subtrop. grasslands savannas and shrublands	-0.001	-0.01	0.01	18,000	0.72	fixed
Large lakes	0.004	-0.003	0.01	18,000	0.28	fixed
Mediterranean forests woodlands and scrub	0.01	0.002	0.01	18,000	0.01	fixed
Montane freshwaters	0.004	-0.02	0.02	18,000	0.66	fixed
Montane grasslands and shrublands	0.01	-0.003	0.03	17,697	0.12	fixed
Polar freshwaters	0.01	0.003	0.01	18,015	0.002	fixed
Polar seas	-0.01	-0.03	0.01	18,000	0.44	fixed
Temperate forests	0.004	0.002	0.01	18,000	0.001	fixed
Temperate wetlands and rivers	0.005	0.002	0.01	18,000	0.0001	fixed
Temperate grasslands savannas and shrublands	-0.004	-0.01	0.001	20,175	0.14	fixed
Tropical wetlands and rivers	0.01	-0.002	0.01	18,000	0.13	fixed
Tropical and subtropical forests	0.01	-0.0005	0.01	17,482	0.06	fixed
Tropical coral	0.02	0.01	0.03	18,000	0.0001	fixed

	Tundra	0.005	-0.002	0.01	18,393	0.17	fixed
	Xeric freshwaters and endorheic basins	-0.01	-0.03	0.0001	18,000	0.04	fixed
	sigma	0.002	0.002	0.002	18,000		residual
Biome - weighted	Boreal forests/taiga	-0.001	-0.004	0.002	18,000	0.56	fixed
	Deserts and xeric shrublands	-0.02	-0.04	-0.003	18,000	0.02	fixed
	Trop. and subtrop. grasslands savannas and shrublands	-0.01	-0.02	-0.0005	18,000	0.04	fixed
	Large lakes	0.002	-0.01	0.01	18,000	0.60	fixed
	Mediterranean forests woodlands and scrub	0.02	0.01	0.03	18,000	0.0001	fixed
	Montane freshwaters	0.02	-0.01	0.05	18,829	0.17	fixed
	Montane grasslands and shrublands	0.03	0.01	0.05	18,000	0.01	fixed
	Polar freshwaters	0.01	0.002	0.02	18,000	0.01	fixed
	Polar seas	-0.002	-0.03	0.02	18,685	0.83	fixed
	Temperate forests	0.01	0.003	0.01	18,000	0.0001	fixed
	Temperate wetlands and rivers	0.01	0.003	0.01	18,000	0.0002	fixed
	Temperate grasslands	-0.004	-0.01	0.003	18,699	0.24	fixed

	savannas and shrublands						
	Tropical wetlands and rivers	0.01	-0.003	0.02	18,000	0.22	fixed
	Tropical and subtropical forests	0.02	0.01	0.03	18,000	0.0001	fixed
	Tropical coral	0.03	0.02	0.04	16,691	0.0001	fixed
	Tundra	0.01	0.001	0.02	18,000	0.04	fixed
	Xeric freshwaters and endorheic basins	-0.003	-0.02	0.01	18,000	0.70	fixed
	sigma	0.003	0.003	0.003	18,000		residual
Biome – fluctuations (SE)	Boreal forests/taiga	0.06	0.06	0.07	18,000	0.0001	fixed
	Deserts and xeric shrublands	0.21	0.17	0.25	19,592	0.0001	fixed
	Trop. and subtrop. grasslands savannas and shrublands	0.11	0.09	0.13	18,342	0.0001	fixed
	Large lakes	0.09	0.07	0.11	17,588	0.0001	fixed
	Mediterranean forests woodlands and scrub	0.16	0.14	0.17	18,000	0.0001	fixed
	Montane freshwaters	0.15	0.10	0.21	18,000	0.0001	fixed
	Montane grasslands and shrublands	0.11	0.07	0.15	18,000	0.0001	fixed

	Polar freshwaters	0.14	0.12	0.15	18,000	0.0001	fixed
	Polar seas	0.08	0.03	0.13	17,192	0.002	fixed
	Temperate forests	0.08	0.08	0.09	17,721	0.0001	fixed
	Temperate wetlands and rivers	0.11	0.11	0.12	18,000	0.0001	fixed
	Temperate grasslands savannas and shrublands	0.07	0.06	0.09	18,000	0.0001	fixed
	Tropical wetlands and rivers	0.21	0.19	0.23	18,385	0.0001	fixed
	Tropical and subtropical forests	0.15	0.13	0.17	17,658	0.0001	fixed
	Tropical coral	0.18	0.16	0.21	18,000	0.0001	fixed
	Tundra	0.13	0.11	0.14	18,000	0.0001	fixed
	Xeric freshwaters and endorheic basins	0.13	0.09	0.16	18,000	0.0001	fixed
	sigma	0.02	0.02	0.02	18,518		residual
Biome – fluctuations (CI)	Boreal forests/taiga	0.07	0.07	0.08	18,000	0.0001	fixed
	Deserts and xeric shrublands	0.22	0.18	0.26	18,000	0.0001	fixed
	Trop. and subtrop. grasslands savannas and shrublands	0.21	0.19	0.23	18,000	0.0001	fixed
	Large lakes	0.11	0.09	0.12	18,725	0.0001	fixed

	Mediterranean forests woodlands and scrub	0.15	0.13	0.16	18,000	0.0001	fixed
	Montane freshwaters	0.14	0.09	0.19	18,000	0.0001	fixed
	Montane grasslands and shrublands	0.17	0.13	0.21	18,000	0.0001	fixed
	Polar freshwaters	0.14	0.12	0.15	18,000	0.0001	fixed
	Polar seas	0.16	0.11	0.20	18,000	0.0001	fixed
	Temperate forests	0.09	0.09	0.10	18,000	0.0001	fixed
	Temperate wetlands and rivers	0.13	0.12	0.13	18,000	0.0001	fixed
	Temperate grasslands savannas and shrublands	0.08	0.07	0.09	17,497	0.0001	fixed
	Tropical wetlands and rivers	0.20	0.19	0.22	18,000	0.0001	fixed
	Tropical and subtropical forests	0.20	0.18	0.21	18,000	0.0001	fixed
	Tropical coral	0.16	0.14	0.19	18,000	0.0001	fixed
	Tundra	0.11	0.09	0.13	18,000	0.0001	fixed
	Xeric freshwaters and endorheic basins	0.18	0.14	0.21	18,000	0.0001	fixed
	sigma	0.02	0.01	0.02	18,440		residual
Biome – fluctuations	Boreal forests/taiga	0.07	0.06	0.08	18,000	0.0001	fixed

weighted (CI)

Deserts and xeric shrublands	0.23	0.19	0.27	18,000	0.0001	fixed
Trop. and subtrop. grasslands savannas and shrublands	0.22	0.20	0.23	18,000	0.0001	fixed
Large lakes	0.11	0.09	0.12	18,000	0.0001	fixed
Mediterranean forests woodlands and scrub	0.15	0.13	0.16	18,000	0.0001	fixed
Montane freshwaters	0.16	0.10	0.21	18,404	0.0001	fixed
Montane grasslands and shrublands	0.17	0.13	0.21	18,000	0.0001	fixed
Polar freshwaters	0.14	0.13	0.15	17,400	0.0001	fixed
Polar seas	0.16	0.11	0.21	18,000	0.0001	fixed
Temperate forests	0.09	0.08	0.10	18,000	0.0001	fixed
Temperate wetlands and rivers	0.12	0.12	0.13	18,000	0.0001	fixed
Temperate grasslands savannas and shrublands	0.08	0.06	0.09	18,000	0.0001	fixed
Tropical wetlands and rivers	0.20	0.18	0.22	18,000	0.0001	fixed

	Tropical and subtropical forests	0.20	0.18	0.22	18,000	0.0001	fixed
	Tropical coral	0.17	0.14	0.19	18,575	0.0001	fixed
	Tundra	0.10	0.09	0.12	17,153	0.0001	fixed
	Xeric freshwaters and endorheic basins	0.18	0.15	0.22	18,000	0.0001	fixed
	sigma	0.02	0.01	0.02	18,683		residual
Biome – fluctuations (SD)	Boreal forests/taiga	0.33	0.32	0.34	18,000	0.0001	fixed
	Deserts and xeric shrublands	0.45	0.41	0.49	18,000	0.0001	fixed
	Trop. and subtrop. grasslands savannas and shrublands	0.47	0.45	0.48	18,000	0.0001	fixed
	Large lakes	0.37	0.35	0.38	18,000	0.0001	fixed
	Mediterranean forests woodlands and scrub	0.44	0.43	0.46	18,000	0.0001	fixed
	Montane freshwaters	0.43	0.37	0.48	18,000	0.0001	fixed
	Montane grasslands and shrublands	0.44	0.40	0.49	18,000	0.0001	fixed
	Polar freshwaters	0.39	0.37	0.40	17,812	0.0001	fixed
	Polar seas	0.39	0.34	0.44	18,000	0.0001	fixed
	Temperate forests	0.36	0.36	0.37	18,000	0.0001	fixed

	Temperate wetlands and rivers	0.39	0.38	0.39	18,000	0.0001	fixed
	Temperate grasslands savannas and shrublands	0.34	0.33	0.36	18,000	0.0001	fixed
	Tropical wetlands and rivers	0.45	0.43	0.47	18,000	0.0001	fixed
	Tropical and subtropical forests	0.48	0.46	0.50	18,000	0.0001	fixed
	Tropical coral	0.47	0.44	0.49	18,391	0.0001	fixed
	Tundra	0.38	0.36	0.39	18,000	0.0001	fixed
	Xeric freshwaters and endorheic basins	0.39	0.36	0.43	18,000	0.0001	fixed
	sigma	0.02	0.02	0.02	18,831		residual
Taxa - mu	Actinopterygii	-0.001	-0.003	0.002	18,000	0.66	fixed
	Amphibia	-0.01	-0.02	-0.005	18,000	0.002	fixed
	Aves	0.004	0.003	0.01	18,000	0.0001	fixed
	Mammalia	0.01	0.01	0.01	18,000	0.0001	fixed
	Reptilia	0.02	0.01	0.02	18,000	0.0001	fixed
	sigma	0.003	0.003	0.004	18,743		residual
Taxa - slope	Actinopterygii	-0.001	-0.004	0.001	18,082	0.33	fixed
	Amphibia	-0.01	-0.02	-0.004	18,000	0.002	fixed
	Aves	0.003	0.002	0.005	18,000	0.0001	fixed
	Mammalia	0.01	0.003	0.01	18,000	0.0001	fixed

	Reptilia	0.01	0.01	0.02	18,000	0.0001	fixed
	sigma	0.002	0.002	0.003	18,000		residual
Taxa - weighted	Actinopterygii	-0.001	-0.004	0.002	18,000	0.67	fixed
	Amphibia	-0.02	-0.03	-0.01	18,000	0.001	fixed
	Aves	0.004	0.003	0.01	18,000	0.0001	fixed
	Mammalia	0.01	0.01	0.02	18,000	0.0001	fixed
	Reptilia	0.02	0.01	0.02	18,000	0.0001	fixed
	sigma	0.003	0.003	0.003	17,290		residual
Taxa – fluctuations (SE)	Actinopterygii	0.14	0.13	0.14	18,000	0.0001	fixed
	Amphibia	0.24	0.23	0.26	18,000	0.0001	fixed
	Aves	0.08	0.07	0.08	18,503	0.0001	fixed
	Mammalia	0.15	0.14	0.16	18,000	0.0001	fixed
	Reptilia	0.19	0.18	0.20	17,583	0.0001	fixed
	sigma	0.02	0.02	0.02	18,000		residual
Taxa – fluctuations (CI)	Actinopterygii	0.15	0.14	0.16	18,000	0.0001	fixed
	Amphibia	0.22	0.20	0.24	18,000	0.0001	fixed
	Aves	0.09	0.09	0.10	17,773	0.0001	fixed
	Mammalia	0.17	0.17	0.18	18,000	0.0001	fixed
	Reptilia	0.17	0.15	0.18	18,000	0.0001	fixed
	sigma	0.01	0.01	0.02	17,064		residual
Taxa – fluctuations	Actinopterygii	0.15	0.15	0.16	16,986	0.0001	fixed

weighted (CI)							
	Amphibia	0.23	0.21	0.25	18,000	0.0001	fixed
	Aves	0.09	0.09	0.09	18,000	0.0001	fixed
	Mammalia	0.18	0.17	0.19	18,000	0.0001	fixed
	Reptilia	0.17	0.16	0.19	18,000	0.0001	fixed
	sigma	0.01	0.01	0.02	18,000		residual
Taxa – fluctuations (SD)	Actinopterygii	0.41	0.40	0.41	18,000	0.0001	fixed
	Amphibia	0.48	0.46	0.50	18,650	0.0001	fixed
	Aves	0.36	0.36	0.36	18,000	0.0001	fixed
	Mammalia	0.45	0.45	0.46	18,000	0.0001	fixed
	Reptilia	0.44	0.43	0.46	17,470	0.0001	fixed
	sigma	0.02	0.02	0.02	18,000		residual
Red List status - mu	Least concern	0.01	0.005	0.01	18,731	0.0001	fixed
	Near threatened	-0.01	-0.01	-0.0003	18,000	0.03	fixed
	Vulnerable	0.002	-0.003	0.01	18,000	0.41	fixed
	Endangered	-0.002	-0.01	0.005	18,000	0.61	fixed
	Critically endangered	-0.01	-0.02	-0.0001	18,000	0.05	fixed
	sigma	0.004	0.003	0.004	18,000		residual
Red List status - slope	Least concern	0.005	0.004	0.01	17,042	0.0001	fixed

	Near threatened	-0.01	-0.01	-0.002	18,000	0.004	fixed
	Vulnerable	0.001	-0.003	0.01	18,374	0.58	fixed
	Endangered	-0.0003	-0.01	0.01	18,000	0.91	fixed
	Critically endangered	-0.003	-0.01	0.005	18,387	0.45	fixed
	sigma	0.003	0.002	0.003	18,000		residual
Red List status - weighted	Least concern	0.01	0.01	0.01	18,000	0.0001	fixed
	Near threatened	-0.01	-0.01	0.0000	18,000	0.05	fixed
	Vulnerable	0.002	-0.003	0.01	18,000	0.39	fixed
	Endangered	-0.003	-0.01	0.01	18,000	0.51	fixed
	Critically endangered	-0.01	-0.02	0.001	18,000	0.09	fixed
	sigma	0.003	0.003	0.004	18,000		residual
Red List status – fluctuations (SE)	Least concern	0.10	0.10	0.10	18,506	0.0001	fixed
	Near threatened	0.13	0.11	0.14	18,000	0.0001	fixed
	Vulnerable	0.11	0.10	0.12	18,000	0.0001	fixed
	Endangered	0.13	0.12	0.15	17,738	0.0001	fixed
	Critically endangered	0.15	0.13	0.17	18,000	0.0001	fixed
	sigma	0.02	0.02	0.02	18,000		residual
Red List status –	Least concern	0.11	0.11	0.12	16,941	0.0001	fixed

fluctuations (CI)							
	Near threatened	0.14	0.13	0.15	18,000	0.0001	fixed
	Vulnerable	0.14	0.13	0.15	17,556	0.0001	fixed
	Endangered	0.16	0.15	0.17	18,000	0.0001	fixed
	Critically endangered	0.17	0.15	0.19	18,000	0.0001	fixed
	sigma	0.02	0.02	0.02	17,523		residual
Red List status – fluctuations weighted (CI)	Least concern	0.11	0.11	0.11	18,000	0.0001	fixed
	Near threatened	0.14	0.13	0.16	18,000	0.0001	fixed
	Vulnerable	0.14	0.13	0.15	18,000	0.0001	fixed
	Endangered	0.16	0.15	0.18	17,504	0.0001	fixed
	Critically endangered	0.17	0.15	0.19	18,000	0.0001	fixed
	sigma	0.02	0.02	0.02	18,000		residual
Red List status – fluctuations (SD)	Least concern	0.38	0.38	0.38	18,000	0.0001	fixed
	Near threatened	0.42	0.41	0.43	18,406	0.0001	fixed
	Vulnerable	0.40	0.39	0.41	17,595	0.0001	fixed
	Endangered	0.44	0.42	0.45	18,000	0.0001	fixed
	Critically endangered	0.47	0.44	0.49	16,838	0.0001	fixed

	sigma	0.02	0.02	0.02	18,000		residual
Latitude - mu	(Intercept)	0.004	0.002	0.01	18,000	0.0001	fixed
	Decimal.Latitude	-0.0000	-0.0000	0.0000	18,000	0.89	fixed
	sigma	0.003	0.003	0.004	18,000		residual
Duration - mu	Duration	0.0000	0.0000	0.0001	18,000	0.04	fixed
Duration - mu	sigma	0.003	0.003	0.004	18,469		residual

Table S4. List of species included in the UK scale analysis of population

change across rarity metrics.

Species name	Number of populations
Acrocephalus schoenobaenus	1
Acrocephalus scirpaceus	1
Agonus cataphractus	1
Alca torda	4
Anarhichas lupus	1
Anas acuta	1
Anas crecca	1
Anas platyrhynchos	2
Anser albifrons	4
Anser fabalis	1
Anthus pratensis	2
Ardea cinerea	1
Arenaria interpres	1
Argentina silus	1
Argentina sphyraena	1
Arnoglossus laterna	1
Asio flammeus	1
Aythya ferina	4
Aythya fuligula	2
Botaurus stellaris	14
Branta bernicla	54
Branta canadensis	1
Branta leucopsis	3
Brosme brosme	1
Bucephala clangula	1
Bufo bufo	1
Burhinus oedicnemus	1
Buteo buteo	1
Calidris alba	1
Calidris alpina	1
Calidris canutus	2
Calidris maritima	2
Callionymus maculatus	1
Capreolus capreolus	3

Carduelis cannabina	2
Cepphus grylle	2
Cervus elaphus	2
Cetorhinus maximus	1
Cettia cetti	1
Charadrius hiaticula	1
Chelidonichthys lucerna	2
Circus aeruginosus	1
Circus cyaneus	1
Clupea harengus	2
Columba oenas	1
Coronella austriaca	1
Corvus corax	1
Corvus corone	1
Corvus monedula	1
Crex crex	1
Cyclopterus lumpus	1
Cygnus columbianus	3
Cygnus cygnus	1
Cygnus olor	1
Delphinus delphis	2
Echiichthys vipera	1
Egretta garzetta	1
Emberiza cirlus	1
Emberiza citrinella	1
Emberiza schoeniclus	3
Eptesicus serotinus	1
Esox lucius	2
Falco peregrinus	1
Falco tinnunculus	1
Fulica atra	1
Fulmarus glacialis	7
Gadus morhua	7
Glyptocephalus cynoglossus	1
Haematopus ostralegus	2
Haliaeetus albicilla	1
Halichoerus grypus	52
Hippoglossus hippoglossus	1
Lagopus lagopus	3

Larus argentatus	1
Larus canus	1
Larus fuscus	1
Larus melanocephalus	1
Lepidorhombus whiffiagonis	2
Lepus timidus	1
Limosa lapponica	2
Limosa limosa	2
Lissotriton vulgaris	5
Lophius budegassa	1
Lophius piscatorius	2
Lullula arborea	1
Melanogrammus aeglefinus	9
Meles meles	1
Mergus serrator	1
Merlangius merlangus	8
Merluccius merluccius	2
Micromesistius poutassou	1
Milvus milvus	3
Molva molva	1
Morus bassanus	4
Muscicapa striata	1
Myotis nattereri	1
Natrix natrix	1
Netta rufina	1
Numenius arquata	2
Nyctalus noctula	1
Oenanthe oenanthe	1
Orcinus orca	1
Oriolus oriolus	1
Oryctolagus cuniculus	6
Oxyura jamaicensis	2
Pandion haliaetus	1
Parus major	2
Passer domesticus	1
Passer montanus	1
Perca fluviatilis	3
Perdix perdix	1
Phalacrocorax aristotelis	8

Phoca vitulina	28
Phrynorhombus norvegicus	1
Pipistrellus pipistrellus	1
Pipistrellus pygmaeus	1
Platichthys flesus	1
Plecotus auritus	1
Plectrophenax nivalis	2
Pleuronectes platessa	6
Pluvialis apricaria	2
Pluvialis squatarola	1
Podiceps cristatus	1
Pollachius pollachius	1
Pollachius virens	3
Prunella modularis	1
Puffinus mauretanicus	2
Pyrrhula pyrrhula	1
Raja brachyura	1
Raja clavata	1
Raja microocellata	1
Raja montagui	1
Rhinolophus ferrumequinum	2
Rhinolophus hipposideros	4
Rissa tridactyla	9
Salmo salar	1
Salmo trutta	1
Scomber scombrus	2
Scyliorhinus canicula	1
Sitta europaea	1
Sprattus sprattus	1
Stenella coeruleoalba	1
Stercorarius parasiticus	3
Sterna dougallii	5
Sterna hirundo	1
Sterna paradisaea	1
Sternula albifrons	1
Streptopelia turtur	1
Strix aluco	1
Sturnus vulgaris	1
Sylvia communis	1

Syngnathus rostellatus	1
Tachybaptus ruficollis	1
Tadorna tadorna	1
Thalasseus sandvicensis	2
Trachurus trachurus	2
Tringa nebularia	1
Tringa totanus	1
Trisopterus esmarkii	2
Trisopterus luscus	1
Trisopterus minutus	1
Triturus cristatus	5
Turdus philomelos	1
Tursiops truncatus	4
Uria aalge	10
Vanellus vanellus	3
Xiphias gladius	1