Rare variant pathogenicity triage and inclusion of synonymous variants

improves analysis of disease associations

Ridge Dershem[1], Raghu P.R. Metpally[1], Kirk Jeffreys[1], Sarathbabu Krishnamurthy[1], Diane T.

Smelser[1], David J. Carey[1], Michal Hershfinkel[2], Janet D. Robishaw[3], Gerda E. Breitwieser[1][§]

[1]Functional and Molecular Genomics, Geisinger Health System, Danville, PA, United States of

America

[2]Faculty of Health Sciences, Ben-Gurion University of the Negev, Israel

[3]Schmidt College of Medicine, Florida Atlantic University, Boca Raton, FL, United States of

America

[§]Corresponding author: gebreitwieser@geisinger.edu

Abstract

Many G protein-coupled receptors (GPCRs) lack common variants that lead to reproducible genome-wide disease associations. Here we used rare variant approaches to assess the disease associations of 85 orphan or understudied GPCRs in an unselected cohort of 51,289 individuals. Rare loss-of-function variants, missense variants predicted to be pathogenic or likely pathogenic, and a subset of rare synonymous variants were used as independent data sets for sequence kernel association testing (SKAT). Strong, phenome-wide disease associations shared by two or more variant categories were found for 39% of the GPCRs. Validating the bioinformatics and SKAT analyses, functional characterization of rare missense and synonymous variants of GPR39, a Family A GPCR, showed altered expression and/or Zn^{2+}-mediated signaling for members of both variant classes. Results support the utility of rare variant analyses for identifying disease associations for genes that lack common variants, while also highlighting the functional importance of rare synonymous variants.

Author summary

Rare variant approaches have emerged as a viable way to identify disease associations for genes without clinically important common variants. Rare synonymous variants are generally considered benign. We demonstrate that rare synonymous variants represent a potentially important dataset for deriving disease associations, here applied to analysis of a set of orphan or understudied GPCRs. Synonymous variants yielded disease associations in common with loss-of-function or missense variants in the same gene. We rationalize their associations with disease by confirming their impact on expression and agonist activation of a representative example, GPR39. This study highlights the importance of rare synonymous variants in human physiology, and argues for their routine inclusion in any comprehensive analysis of genomic variants as potential causes of disease.

Introduction

The superfamily of G protein-coupled receptors (GPCRs) translates extracellular signals from light, metabolites and hormones into cellular changes, which makes them the targets of a significant fraction of drugs currently on the market [1]. Genome-wide association studies (GWAS) on common variants in GPCRs have begun to identify their contributions to various disease processes [2, 3]. However, many GPCRs lack common variants and alternate strategies are needed to understand their roles. Recently, sequence kernel association testing (SKAT) methods have been applied to rare variants in GPCRs to assess disease associations. Initial approaches binned all rare variants in a genomic region or gene [4, 5], but more recent methods group the rare variants most likely to contribute to disease associations, or aggregate variants based on domain or family structures [6, 7].

Large-scale studies of orphan or understudied GPCRs to characterize natural genetic variation in the human population can provide insights into the biological function and/or potential causal contributions to disease processes [8]. The DiscovEHR collaboration represents a tremendous resource [9]; here we use whole exome sequences and clinical information from 51,289 individuals in the DiscovEHR cohort. We identified sequence variants (common variants, mean allele frequency, MAF $\geq$ 1% and rare variants, MAF < 1%) for 85 orphan or understudied GPCRs not amenable to GWAS studies, and performed SKAT analyses. We binned rare variants into functional classes: loss-of-function (LOF, truncation and frameshift) variants; missense variants with predicted pathogenicity; or rare synonymous variants showing altered codon bias. We performed independent SKAT analyses on the various functional classes to determine their disease associations. Remarkably, for those GPCRs with sufficient numbers of patients for

statistical analyses, we found the top disease associations were common to all available variant

classes in the gene.

To assess the validity of the disease association results, we focused on *GPR39*. Among its

particular advantages, *GPR39* contained sufficient numbers of variants in all three functional

classes to be amenable to rare variant approaches. Second, *GPR39* is a small Family A GPCR,

allowing rapid generation of mutants. Third, GPR39 is activated by extracellular Zn^{2+} and is

coupled to inositol phosphate production [10, 11], permitting straightforward functional analyses.

Finally, we have only a rudimentary understanding of the role(s) of GPR39 in human physiology,

despite its broad expression and multiple cellular functions [12, 13]. Our results demonstrate the

validity of combined computational and functional approaches to provide important insights into

the clinical contributions of orphan and understudied GPCRs. Results also focus attention on the

importance of rare synonymous variants for identifying disease associations. The overall strategy

can be readily applied to other classes of genes without clinically important common variants.

Results

Rare variant-based disease associations of orphan / understudied GPCRs

We identified rare variants in whole exome sequences from 51,289 individuals, and to

predict their functional impact, variants were annotated and sorted into three classes in order of

predicted severity, i.e., Loss-of-Function (LOF; premature stop codon, loss of a start or stop codon,

or disruption of a canonical splice site), missense variants (predicted to be likely pathogenic (pLP)

or pathogenic (pP) by RMPath scoring (Supplemental Methods)), and synonymous variants with

significantly different codon frequency relative to the reference codon (termed SYN_ΔCB, i.e.,

synonymous variants with altered codon bias).

Determining disease associations for genes having only low frequency variants requires

binning of variants to increase statistical power. Binning methods have focused on gene or genetic

region, and have recently been expanded to incorporate regulatory regions and/or pathways by

incorporating biological information from curated knowledge databases [14]. For this study, we

were specifically looking for clinically impactful rare coding variants which could be validated by

functional studies of the relevant GPCRs. We used sequence kernel association tests (SKAT) and

binned variants according to functional classes described above, and focused on the disease

associations which were common to more than one class . **Figure 1** illustrates the analysis pipeline

for the entire set of 85 GPCRs. Of the 72 GPCRs with sufficient variants to allow at least 2 variant

classes to be analyzed by SKAT, 28 yielded significant disease associations. Because this was an

exploratory analysis, we used a significance cut-off of 0.05, i.e., P-values < 6.2E-07 after

Bonferonni correction for multiple testing. **Table 1** shows the top disease associations for the 13

GPCRs that had sufficient variants in the LOF class. Supporting validity, associations between

GPR37L1 and epilepsy [15], and *LGR5* and alopecia [16, 17] have been previously identified by

other means. Attesting to the discovery potential, other disease associations were novel, including

GPR161 and hyperhidrosis, *LGR4* and Sicca syndrome, *GPR153* and hyperpotassemia, *GPR84*

and anemia, and *GPR39* and peripheral nerve damage, among others. Notably, the phenome-wide

disease associations for the predicted LOF class showed congruence across the other variant

classes for the subset of GPCRs having sufficient missense or SYN_ΔCB variants for independent

analyses. While predicted LOF variants are easily identified and likely to have the greatest

functional impact, missense variants classed as pLP + pP can also have significant effects on

GPCR function. For the subset of GPCRs without sufficient LOF variants, we found significant

disease associations for the missense and SYN_ΔCB classes of variants, **Table 1**. Some of the

associations validate previous studies, including *GPR183* and disorders of liver [18], and *GPR85*

with acute myocardial infarction [19]. Other associations were novel, and represent potential

targets for future study, including *GPR132* and disturbances in sulphur-bearing amino acid

metabolism, *GPR176* with asthma, *GPR12* with epilepsy, and *GPRC5D* with renal

osteodystrophy. Of note, the SYN_ΔCB variant class had significant disease associations in

common with LOF and/or missense variants, and most consistently had sufficient variants to be

included in the analysis. Parallel SKAT analyses of the three distinct classes of rare variants

produced two important results. First, independent analysis of distinct classes of variants yielded

concordant disease associations, increasing confidence in the results. Second, the most consistent

source of phenome-wide disease associations was SYN_ΔCB. Altogether, these results provide a

strong rationale for including all three functional variant classes in disease association analyses.

Identification of *GPR39* common and rare variants in the 51,289 WES DiscovEHR cohort

122   Disease associations provide a rich source of hypotheses regarding the potential

contributions of GPCRs. The first step in defining a causal role for these rare variants, however,

is functional validation of the SKAT results. Here we focused in particular on rare synonymous

variants with altered codon frequency (SYN_ΔCB), whose functional consequences are not well

understood. Many of the GPCRs have no known agonist, and their dominant signaling pathways

have not been characterized (**Table 1**). We therefore chose to focus on *GPR39*, the Zn^{2+} receptor,

which has been characterized in cellular and knockout mouse models, and for which both agonist

and dominant signaling pathways are known [12, 13].

130   All common and rare variants in *GPR39* were identified in 51,289 individuals in the

DiscovEHR cohort [9] (**Figure 2**). None of the *GPR39* variants were found in the ClinVar

database. Accordingly, all LOF, missense, and synonymous variants were analyzed and classified

as described for **Table 1**. The common variants were used for Phenome-wide Association Studies

(PheWAS), and included a missense variant (2:133174764, Ala50Val) predicted to be benign, a

synonymous variant (2:133174999; Thr128Thr), which changed a common to the rarest codon

(acA (28%) to acG (12%), i.e., a 2.33-fold decrease in frequency), and a non-coding variant

(2:133402607). Notably, no phenome-wide disease associations were detected (Supplemental

Figure S1).

The rare variants used for SKAT, i.e., LOF, missense (pLP + pP), and SYN_ΔCB variants,

were distributed throughout the coding region of *GPR39* (**Figure 3**). Examination of the 39

missense variants that were categorized as pLP and pP confirmed that the RMPath pipeline

identified variants which are likely to impact GPR39 function (**Figure 3**). The three pP variants

(S78L, C108R, R133G) included a cysteine that participates in a disulfide bond at the extracellular

face of GPR39 [20]. The pLP variants included one which contributes to the Zn^{2+} binding site

(H19R) [21], two variants in a disulfide bond cysteine (C191G, C191W) [20], a residue in the

highly conserved NPXXY motif common to Family A GPCRs (N340S) [22], and a variant distal

to a putative palmitoylation site in the proximal carboxyl terminus (R352Q). Other residues in the

pLP category introduce charges or proline residues into transmembrane helices or extracellular

loops (S49N, M87I, T122M, A167S, P179T, R193C, S222Y, T291I, A293P, R302W, R302Q,

A307V, P310A, Y314R, R320Y, S336R, R352Q, R362C, H367R, R386C, and R390C). We also

identified the subset of 21 variants which introduced at least a two-fold change in site-specific

codon frequency (SYN_ΔCB; denoted by arrowheads in **Figure 3**).

Disease associations of *GPR39* rare variant classes

The initial SKAT screen for *GPR39* disease associations compared top associations for the

three independent classes of rare variants. To more fully explore the results for *GPR39* and

increase power to identify all significant disease associations, we combined all individuals

heterozygous for rare LOF, pLP and pP variants (termed LOF/MISS). Analysis was restricted to

only those ICD9 codes with ≥200 individuals independent of genotype [23], and the results are

plotted against ICD9 codes in **Figure 4A**. SKAT analysis was also run on rare SYN_ΔCB variants,

plotted in **Figure 4B**. Both the LOF/MISS and SYN_ΔCB analyses had a significant number of

phenome-wide associations in common. All results for both variant classes that were above the

significance cut-off for multiple testing are listed in **Table 2**. Top associations for *GPR39* included

lesion of the ulnar nerve and benign prostate hyperplasia with lower urinary tract obstruction.

Note that individuals who had both rare LOF/MISS and SYN_ΔCB variants were excluded from

analyses. Also of note in **Table 2**, some significant disease associations were specific to either the

LOF/MISS or SYN_ΔCB variant classes. To assess the importance of rare variant pathogenicity

triage for binning prior to SKAT analysis, we combined all rare variants (LOF, missense, and

synonymous variants; 209 variants, 177 used in analysis) and ran SKAT. The only significant

disease association was upper quadrant abdominal pain (ICD9 789.02; P-value 8.98E-06;

significance cut-off at P < 7.6E-05). We also ran SKAT on those rare missense variants predicted

to be benign (**Figure 2**, pB; 61 variants); no statistically significant disease associations were

obtained. Likewise, we ran SKAT analysis on all rare synonymous variants and found top

associations similar to those obtained for SYN_ΔCB, but with less significant P-values, suggesting

dilution of impactful variants, i.e., lesion of ulnar nerve (P-value 1.08E-07), and benign prostate

hyperplasia with urinary tract obstruction and other lower urinary symptoms (P-value 3.79E-05)

(compare **Tables 2** and **3**). We conclude that pathogenicity triage to refine rare variant binning

was crucial to identifying disease associations for the LOF and missense variants, and binning of

the subset of rare synonymous variants having large changes in local codon frequency bias

strengthened disease associations. Further, refining the rare variant input into the SKAT analyses

yielded disease associations common to multiple independent classes of variants, strengthening

confidence in the associations.

Validation of functional impact of rare *GPR39* variants

To confirm the functional basis for the observed disease associations, we generated the rare

missense and SYN_ΔCB variants in the reference human *GPR39* background. The cDNA

construct included an amino terminal 3X FLAG epitope followed by a minimal bungarotoxin

binding site, for ease of expression analysis and subcellular localization, respectively. Expression

in HEK293 cells of equivalent amounts of cDNA and subsequent western blotting of lysates

showed significant differences in net expression levels of the rare missense and SYN_ΔCB

variants. Ten missense variants had significantly reduced expression and one had significantly

increased expression relative to WT GPR39 (Supplemental **Figure S2**). Rare missense variants

with altered expression were distributed throughout GPR39 structural domains, as color coded in

Figure 5A. Likewise, SYN_ΔCB variant expression was significantly greater than WT GPR39

for three variants, while fourteen variants were expressed significantly less that WT (Supplemental

Figure S2), quantified in **Figure 5B**. Surprisingly, despite the change from rare to common codon

frequency for the majority of SYN_ΔCB variants, only four showed increased expression relative

to WT. Four variants went from common to rare codons, denoted by * on the x-axis in **Figure 5B**.

Contrary to expectations, Q265 was expressed at significantly higher levels than WT (~150%),

despite the large decrease in codon frequency. Overall, results argue that factors in addition to

site-specific codon usage bias determine net expression levels for rare synonymous variants.

GPR39 signaling is complex, with multiple signaling outputs. However, in both

endogenous and heterologous expression systems, GPR39 activates inositol trisphosphate (IP_3)-

dependent release of intracellular Ca^{2+} [10, 11, 24]. The first step in the signaling pathway is G_q-

mediated activation of phospholipase C, which produces diacylglycerol and IP_3. Here we exposed

HEK293 cells transiently expressing WT or variant GPR39 to its agonist, $ZnCl_2$, and used a FRET

assay to quantify accumulation of intracellular inositol monophosphate (IP_1) in the presence of

LiCl. **Figure 6A** illustrates normalized maximal levels of IP_1 for WT and representative missense

variants having decreased (R386C) or increased (W314R) EC_{50} for activation by $ZnCl_2$. **Figure**

6B plots the EC_{50}s for WT and all other missense variants, color-coded for their locations in

GPR39 domains. Four variants, S78L, M87I, T122M and R133G, had no measurable activity,

likely accounted for by low levels of expression (recall **Figure 5A**). Three additional variants,

C108R, R193C and R302Q, had dose-response curves that were significantly right-shifted,

precluding determination of EC_{50} or V_{MAX}, despite expression levels not significantly different

from WT. Likewise, **Figure 6C** illustrates the normalized dose-response relations for WT GPR39

and representative SYN_ΔCB variants which decrease (V354) or increase (I29) the EC_{50} for $ZnCl_2$

activation, and **Figure 6D** plots the EC_{50}s of WT GPR39 and all SYN_ΔCB variants. In contrast

to the missense variants, all SYN_ΔCB variants had $ZnCl_2$-stimulated IP_1 activity. Six variants

had increased EC_{50}s and three variants having significantly reduced EC_{50}s relative to WT. Surface

expression of all rare missense and SYN_ΔCB variants was assessed by BgTx-Alexa594 binding

on non-permeabilized cells (**Figures 6E, 6F**), and generally correlated with the extrapolated V_{max}

values for the IP_1 assays (**Figures 6G, 6H**). Notable exceptions include the missense variants

C108R, R193C and R320Q, which had surface expression comparable to WT but little IP_1 activity.

Likewise, the most striking SYN_ΔCB variant, Q265, had an overall increase in net expression

(**Figure 5B**), increased EC_{50} relative to WT (**Figure 6D**), and increased surface localization and

V_{MAX} relative to WT (**Figure 6F, 6H**). Overall, results argue for a striking similarity in functional

impact on expression and IP_1 signaling for a subset of both rare missense and SYN_ΔCB variants

of GPR39, rationalizing the disease associations identified for both classes of rare variants. The

congruence of top disease associations between rare LOF, pathogenic missense and SYN_ΔCB

variants for a significant subset of GPCRs (recall **Table 1**) argues strongly for the utility of these

analytical approaches for identification of functionally impactful rare variants, and the imperative

to include rare synonymous variants in genomic and functional analyses.

Discussion

The related GWAS and PheWAS approaches, which utilize common variants, have

produced strong, validated targets for drug development [2]. However, many genes do not have

common variants, and additional approaches are required. The advantages of rare coding variant

approaches are two-fold. First, rare variants can have stronger effects on protein function, and

second, assessment of the impact of rare coding variants on protein expression and function is

straightforward. However, due to their low frequency, it is necessary to combine rare coding

variants to increase the statistical power for detecting disease associations. In this study, we used

pathogenicity triage as a basis for rare variant binning, based on the hypothesis that aggregating

variants with similar directions of effect on protein function would amplify disease associations.

We considered, in order of the strength of their potential effects, predicted LOF variants, predicted

likely pathogenic or pathogenic missense variants, and synonymous variants with altered codon

frequency bias. as a test of the importance of rare variant triage on disease associations, we

compared agnostic binning of all rare variants in *GPR39* with the disease associations identified

for the three sorted variant classes. Of note, binning of all rare variants yielded no significant

disease associations, while curated classes of those same rare variants had concordant top

phenome-wide disease associations, obtained with non-overlapping individuals. Likewise, SKAT

analysis of the subset of rare missense variants predicted to be benign yielded no significant disease

associations. Results argue strongly that binning all rare variants in a gene or genetic region dilutes

the effects of impactful variants, and appropriate pathogenicity triage of rare variants is an integral

first step in successful application of disease association analyses.

Over the past 10 years, rare synonymous variants have been shown to impact protein

expression and function with a frequency comparable to non-synonymous variants, although the

mechanism(s) by which they exert their biological function are diverse [25-36]. Rare synonymous

variants have not, however, been systematically utilized for disease association analysis [31, 37-

39]. Here we demonstrate that 39% of understudied GPCRs have impactful rare synonymous

variants which yield disease associations comparable to LOF or missense variants in the same

gene, which may provide impetus for exploratory drug development despite the lack of identified

endogenous agonists. Our data also highlight the necessity for improved approaches for

pathogenicity triage of rare synonymous variants. For *GPR39*, SKAT analysis on all rare

synonymous variants for the most part yielded disease associations similar to those obtained with

the SYN_ΔCB subset, but having reduced significance, arguing for dilution of impactful variants.

However, a few disease associations, including psoriatic arthropathy, showed more significant P-

values in the combined synonymous variant analysis (3.87E-11) versus SYN_ΔCB (1.03E-06),

implying that simply isolating those synonymous variants having altered local codon usage bias

missed some variants with significant functional impacts (see **Tables 2** and **3**). Using both

LOF/missense and synonymous variants as independent data sets in the SKAT analyses increases

confidence in the identified disease associations and provided the impetus for further functional

studies.

A disease association does not prove causation, but implies an impact on protein expression

and/ or function, which must be validated experimentally. As proof-of-concept, *GPR39* was

chosen because its agonist and dominant signaling pathway were known, and rare variant classes

produced strongly congruent disease associations. We identified 168 rare variants in *GPR39* in the

51,289 WES DiscovEHR cohort. Most individuals were heterozygous for the rare variants studied,

and only 1/327 individuals had more than one rare variant. Of the 110 missense variants, the

RMPath pipeline predicted that 28 were likely to be pathogenic (pLP + pP). Among these, rare

variants previously shown to be important for GPR39 function were identified, including the zinc

binding site [21], a critical extracellular disulfide bond [20], the pH sensor [11], and the NPXXY

motif common to all Family A GPCRs [22]. When expressed in HEK293 cells, these variants had

the expected effects on EC_{50}s and maximal activities. Of the 47 synonymous variants, the codon

usage table identified 21 variants having significantly altered codon usage frequency, and many

had significant effects on expression, surface localization, and Zn^{2+}-induced IP_1 production.

Surprisingly, the distributions of rare missense and SYN_ΔCB variants in the GPR39 structural

domains were distinct. The majority of missense variants were localized to the transmembrane

domain (77%), half in the extracellular loops (10/26), one an intracellular loop and the remainder

in the transmembrane helices (9/26). In contrast, ~50% of SYN_ΔCB variants were in the carboxyl

terminus; the remainder were distributed among transmembrane helices (3/20), intracellular (2/20)

and extracellular (4/20) loops. The carboxyl terminal variants in both classes had significant

effects on expression and signaling, primarily reducing V_{MAX}. The carboxyl terminus of GPR39

is relative large (108 amino acids) and has 22 putative phosphorylation sites. GPR39 signaling

outputs are regulated by agonist-toggled binding of protein kinase inhibitor β [40]. In addition,

phosphorylation by Rho kinase mediates desensitization of GPR39 [41]. Thus, carboxyl terminal

missense or synonymous variants may alter secondary structure and/or phosphorylation to impact

signaling. Results focus attention on the carboxyl terminus as a critical domain mediating distinct

aspects of GPR39 signaling. It should be noted that GPR39 signaling is complex, including G_s-

mediated   cAMP   production, G_q-mediated   IP$_3$ accumulation,   and   SRF-SRE-mediated

transcriptional regulation via $G_{12/13}$ [41]. The current study examined only G_q-mediated IP$_3$

generation, leaving open the possible impact of missense and/or synonymous variants on signaling

bias.

Based on ClinVar and related databases, *GPR39* has not been causally linked to disease.

The present study focused on the general patient population, and all patients in the present study

were heterozygous for *GPR39* rare variants. Nevertheless, both LOF/MISS and SYN_ΔCB

variants yielded disease associations by SKAT which suggest a role for GPR39 in nerve function,

benign hyperplasia of the prostate, psoriatic arthropathy, diseases of hair and hair follicles, and

benign essential hypertension. These associations are in agreement with cellular and animal

studies which suggest that GPR39 regulates neuronal transport of potassium and chloride,

attenuating post-synaptic excitability [12, 42-44]. Animal models demonstrate GPR39-mediated

up-regulation of the neuronal K^+/Cl^- cotransporter attenuates seizure activity [44], and knockout

of GPR39 promotes depression [45]. GPR39 is up-regulated by some anti-depressants [46] in

mouse models. One study in humans showed that GPR39 levels were significantly reduced in the

hippocampus and cortex of suicide victims [47]. The highest levels of Zn^{2+} in the body are in the

prostate and seminal fluid [48], and GPR39 is differentially expressed in normal versus malignant

prostate cells, where it regulates cell growth and proliferation [10, 12, 13], potentially contributing

to the association with benign prostate hyperplasia observed in the present work. Finally, the

observation that individuals heterozygous for rare variants in GPR39 have a significant association

with diseases of hair and hair follicles should be considered in light of a recent study that found

that GPR39 marks a class of stem cells in the sebaceous gland and contributes to wound healing

in mice [49]. Overall, the identification of a subset of clinical phenotypes that echo cell culture

and animal model studies supports disparate roles for GPR39 in human physiology, and validates

the SKAT approaches used herein to identify disease associations. Additional phenotypes, not

previously associated with GPR39 including psoriatic arthropathy and macular puckering of the

retina, deserve further attention. In conclusion, we have identified strong disease associations for

GPR39 using either LOF/missense or rare synonymous variants, which confirm and extend cellular

and animals studies. Rare variants from both the missense and synonymous classes altered GPR39

expression and function, providing a rationale for the observed disease associations. The approach

applied to a larger set of orphan or understudied GPCRs also yielded robust associations with

phenome-wide significance, providing potential novel targets for further investigation. The

methods described in this report are easily applied to any set of genes of potential clinical

importance with available genomic and integrated longitudinal electronic health records. Finally,

this study highlights the importance of rare synonymous variants in human physiology, and argues

for their routine inclusion in any comprehensive analysis of genomic variants as potential causes

of disease.

Materials and methods

Whole Exome Sequence (WES) Data

The MyCode® Community Health Initiative recruits Geisinger Health System (Geisinger) patients from a broad range of inpatient and outpatient clinics to integrate genetic information with their clinical EHR data to foster discoveries in health and disease [23]. For this study, we used WES from 51,289 participants [9], which includes patients enrolled through primary care and specialty outpatient clinics. Participants were 59% female, with a median age of 61 years, and predominantly Caucasian (98%) and non-Hispanic/Latino (95%). Patient demographics are in **Supplemental Table 1**. Whole exome sequencing was performed using 75 bp paired-end sequencing on an Illumina v4 HiSeq 2500 to a coverage depth sufficient to provide greater than 20x haploid read depth of over 85% of targeted bases for at least 96% of samples. Sample preparation, sequencing, sequence alignment, variant identification, genotype assignment and quality control steps were carried out as previously described [23].

Analysis of site-specific rare codon changes

Codon usage tables define the relative fraction of different codons used in the genes of an organism [50, 51]. For all rare synonymous codons, we determined the ratio of the genome-wide codon fraction of the *GPR39* reference codon over the synonymous codon variant at that position, e.g., for Ser147Ser (tcG/tcA), 0.15/0.06 = 2.5, i.e., a 2.5-fold change. All synonymous variants with at least a two-fold change in site specific codon usage were used in the SKAT analyses.

Phenome-wide association analysis (PheWAS)

De-identified EHR data was obtained from an approved data broker. All unique ICD9 codes with ≥200 patients (regardless of genotype) with at least 3 independent incidences of the

ICD9 code were extracted from the EHR [52]. Individuals having 1-2 calls were excluded, and

those having no calls of a particular ICD9 code were considered as controls. All non-Europeans

were excluded from the analysis, as was one sample from pairs of closely related individuals up to

first cousins. All models were adjusted for sex, age, age^2 and first 4 principal components. The

PheWAS R package were used for association analyses, which were performed for three GPR39

common variants (2:133174764, Ala50Val; 2:133174999; Thr128Thr; and 2:133402607,

noncoding), and plotting of results was done using GraphPad Prism (V.6).

Sequence Kernel Association Testing (SKAT)

To test for associations with clinical phenotypes, we used the sequence kernel association

test (using default weights; SKAT R package), a gene-based binned statistical test comparing the

burden of rare variants in cases and controls. Cases and controls were defined as for PheWAS

analysis [52]. We defined various variant sets, including (1) only potential loss of function

(splicing, stop gain, stop loss, or frameshift), (2) predicted pathogenic missense (pLP and pP by

RMPath scoring) variants, and (3) synonymous variants having large codon usage frequency

changes (at least a two-fold change in frequency between human reference codon and altered

codon, SYN_ΔCB). For full *GPR39* analysis, we combined categories (1) and (2). Age, age^2, sex,

BMI, and first four principal components were used as co-variates. All non-European samples

were excluded. Statistical significance was determined using the Bonferroni correction, α/m,

where $\alpha=0.05$ and m= # of variant sets and disease codes, as indicated for each analysis [53].

Generation of *GPR39* variants

All missense and SYN_ΔCB variants were generated by gene synthesis (Thermo Fisher Scientific,

Gene Art) based on the NP_001499.1 reference sequence (GeneID 2863, 453 amino acids), and

included tandem tags at the amino terminus to facilitate expression and functional studies. After

the initiation methionine codon, we added a 3X FLAG epitope followed by a minimal

bungarotoxin binding site. The complete inserted sequence is:

GACTACAAAGACCATGACGGTGATTATAAAGATCATGATATCGATTACAAGGATGA

CGATGACAAGGGTTGGAGATACTACGAGAGCTCCCTGGAGCCCTACCCTGACGGT.

Expression of GPR39 variants

WT and variant constructs were transiently transfected into HEK293 cells. Cells were

lysed after 48 hours, and 25 μg of lysates were separated on 4-15% SDS PAGE gels (BioRad),

blotted to nitrocellulose, blocked with 5% milk in TBS-T, and exposed for 1 hour at room

temperature to anti-FLAG monoclonal antibody conjugated to HRP (Sigma). Blots were exposed

to SuperSignal West Pico Chemiluminescence Substrate, (Thermo Fisher), images recorded on a

FUJIFILM LAS-4000mini luminescence analyzer and processed with Image-Gauge version 3.0.

GPR39 expression was quantified using Multi-Gauge software, normalizing to WT expression and

corrected for HEK293 background on the same blot.

Functional analysis of GPR39 variants

WT and variant constructs were transiently transfected into HEK293 cells, cultured for 24

hours, and equivalent numbers of cells were then plated in poly-D-lysine-coated 96 well plates for

a further 24 hours. Replicates were treated with various concentrations of $ZnCl_2$ for 10 min at 37

°C. Levels of IP_1 were determined as recommended by manufacturer (IP-One ELISA kit, Cisbio

USA, Inc.). Plates were read on a POLARStar Omega (BMG Labtech) plate reader. EC_{50} was

determined by fitting normalized dose-response data with the Michaelis-Menten equation, and

V_{MAX} was separately calculated on raw data (GraphPad Prism, V6). Parallel plates were exposed

to Alexa-594-conjugated bungarotoxin (3 μg/ml, 5 min at 37 °C, 5% CO_2) to assess surface

expression of GPR39 variants. After removal of bungarotoxin and washes with PBS, fluorescence

was read on the POLARStar Omega (BMG Labtech) reader. Each variant was assayed in ≥3

independent transfections; WT, standards and solution blanks were included in each assay.

Acknowledgements

415   We are grateful to Geisinger for financial support, Geisinger and the Regeneron Genetics Center

416   for access to the DiscovEHR cohort, and to the patients in the MyCode Biobank.

References

1. Hauser, A. S., Attwood, M. M., Rask-Andersen, M., Schioth, H. B., & Gloriam, D. E. Trends

in GPCR drug discovery: new agents, targets and indications. Nat. Rev. Drug Discov. 2017;

16: 829-842.

2. Roth, B. L., Kroeze, W. K. Integrated approaches for genome-wide interrogation of the

druggable non-olfactory G protein-coupled receptor family. J. Biol. Chem. 2015; 290: 19471-

19477.

3. Kovacs, P., Schonberg, T. The relevance of genomic signatures at adhesion GPCR loci in

humans. Handb. Exp. Pharmacol. 2016; 234: 179-217.

4.   Wu, M. C., Lee, S., Cai, T., Li, Y., Boehnke, M., Lin, X.   Rare-variant association testing for

sequencing data with the sequence kernel association test.   Am. J. Hum. Genet. 2011; 89:   82-

93.

5.   Lee, S., Emond, M. J., Bamshad, M. J., Barnes, K. C., Rieder, M. J., Nickerson, D. A., et al.

Optimal unified approach for rare-variant association testing with application to small-sample

case-control whole-exome sequencing studies.   Am. J. Hum. Genet. 2012; 91:   225-237.

6.   Richardson, T. G., Shihab, H. A., Rivas, M. A., McCarthy, M. I., Campbell, C., Timpson, N.

J., et al.   A protein domain and family based approach to rare variant association analysis.   PLoS

One 2016; 11:   e0153803.

7.   Friedrichs, S., Malzahn, D., Pugh, E. W., Almeida, M., Liu, X. Q., Bailey, J. N.   Filtering

genetic variants and placing informative priors based on putative biological function.   BCM

Genet. 2016; 71 Suppl. 2:   8.

8.   Sriram, K., Insel, P. A.   GPCRs as targets for approved drugs: How many targets and how

many drugs? Mol. Pharmacol. 2018 Jan.3 doi: 10.1124/mol.117.111062.

9. Dewey, F. E., Murray, M. F., Overton, J. D., Habegger, L., Leader, J. B., Fetterolf, S. M., et

al.   Distribution and clinical impact of functional variants in 50,726 whole-exome sequences

from the DiscovEHR study.   Science 2016; 354:   pii aaf6814.

10. Asraf, H., Salomon, S., Nevo, A., Sekler, I., Mayer, D., Hershfinkel, M. The ZnR/GPR39

interacts with the CaSR to enhance signaling in prostate and salivary epithelia.   J. Cell

Physiol. 2014; 229:   868-877.

11. Cohen, L., Asraf, H., Sekler, I., Hershfinkel, M. Extracellular pH regulates zinc signaling via

an Asp reside of the zinc-sensing receptor (ZnR/GPR39).   J. Biol. Chem. 2012; 287:   33339-

33350.

12. Sunuwar, L., Gilad, D., Hershfinkel, M. The zinc sensing receptor, ZnR/GPR39, in health

and disease. Front. Biosci (Landmark Ed) 2017; 22: 1469-1492.

13. Hershfinkel, M. The Zinc Sensing Receptor, ZnR/GPR39, in health and disease. Int. J. Mol.

Sci. 2018; 19: 437-458.

14. Basile, A. O., Byrska-Bishop, M., Wallace, J., Frase, A. T., Ritchie, M. D. Novel features and

enhancements in BioBin, a tool for biologically inspired binning and association analysis of

rare variants. Bioinformatics 2018; 34: 527-529.

15. Giddens, M. M., Wong, J. C., Schroeder, J. P., Farrow, E. G., Smith, B. M., Owino, S., et al.

GPR37L1 modulates seizure susceptibility: Evidence from mouse studies and analyses of a

human GPR37L1 variant. Neurobiology of Disease 2017; 106: 181-190.

16. Hoeck, J. D., Biehs, B., Kurtova, A. V., Kljavin, N. M., de Sousa, E. Melo F., et al. Stem cell

plasticity enables hair regeneration following LGR5+ cell loss. Nat. Cell Biol. 2017; 19: 666-

675.

17. Michel, L., Reygagne, P., Benech, P., Jean-Louis, F., Scalvino, S., Ly Ka So, S., et al. Study

of gene expression alteration in male androgenic alopecia: evidence of predominant molecular

signaling pathways. Brit. J. Dermatol. 2017; 177: 1322-1336.

18. Guillemot-Legris, O., Mutemberezi, V., Cani, P. D., Muccioli, G. G. Obesity is associated

with changes in oxysterol metabolism and levels in mice liver, hypothalamus, adipose tissue

and plasma. Sci. Reports 2016; 6: 19694.

19. Li, Y., Dong, J., Xuan, Y.-H., Liu, S.-S., Luo, J.-Y., Xiao, Y.-J., Tet al. Identification of

microRNA expression in a rat model of post-infarction heart failure. Int. J. Clin. Exp. Pathol.

2017; 10: 4729-4738.

472 20. Storjohann, L., Holst, B., Schwartz, T. W. A second disulfide bridge from the N-terminal

473  domain to extracellular loop 2 dampens receptor activity in GPR39. Biochem. 2008; 47:

474  9198-9207.

475 21. Storjohann, L., Holst, B., Schwartz, T. W. Molecular mechanism of Zn^{2+} agonism in the

476  extracellular domain of GPR39. FEBS Lett. 2008; 582: 2583-2588.

477 22. Yuan, S., Filipek, S., Palxzewski, K., Vogel, H. Activation of G-protein-coupled receptors

478  correlates with the formation of a continuous internal water pathway. Nat. Commun. 2014; 5:

479  4733.

480 23. Carey, D. J., Fetterolf, S. N., Davis, F. D., Faucett, W. A., Kirchner, H. L., Mirshahi, U., et

481  al. The Geisinger MyCode community health initiative: an electronic health record-linked

482  biobank for precision medicine research. Genet. Med. 2016; 18: 906-913.

483 24. Verhulst, P. J., Lintermans, A., Janssen, S., Loeckx, D., Hummelreich, U., Buyse, J., et al.

484  GPR39, a receptor of the ghrelin receptor family, plays a role in the regulation of glucose

485  homeostasis in a mouse model of early onset diet-induced obesity. J. Neuroendocrinol. 2011;

486  23: 490-500.

487 25. Hunt, R. C., Simhadri, V. L., Iandoli, M., Sauna, Z. E., Kimchi-Sarfaty, C. Exposing

488  synonymous mutations. Trends Genet. 2014; 30: 308-321.

489 26. Bali, V., Bebok, Z. Decoding mechanisms by which silent codon changes influence protein

490  biogenesis and function. Int. J. Biochem. Cell. Bol. 2015; 64: 58-74.

491 27. Brar, G. A. Beyond the triplet code: context cues transform translation. Cell 2016; 167: 1681-

492  1692.

493 28. Supek, F., Minana, B., Valcarcel, J., Gabaldon, T., Lehner, B. Synonymous mutations

494  frequently act as driver mutations in human cancers. Cell 2014; 156: 1324-1335.

29. Fernandez-Calero, T., Cabrera-Cabrera, F., Ehrlich, R., Marin, M. Silent polymorphisms: can

the tRNA population explain changes in protein properties? Life 2016; 6:9;

doi:10.3390/life6010009.

30. Jung, H., Lee, D., Park, D., Kim, Y. J., Park, W. Y., Hong, D., Park, P. J., Lee, E. Intron

retention is a widespread mechanism of tumor-suppressor inactivation. Nat. Genet. 2015; 47:

1242-1248.

31. Soussi, T., Taschner, P. E., Samuels, Y. Synonymous somatic variants in human cancer are

not infamous: a plea for full disclosure in databases and publications. Hum. Mutat. 2017; 38:

339-342.

32. Kukreti, R., Tripathi, S., Bhatnagar, P., Gupta, S., Chauhan, C., Kubendran, S., et al.

Association of DRD2 gene variant with schizophrenia. Neurosci. Lett. 2006; 392: 68-71.

33. Grymek, K., Lukasiewica, S., Faron-Goreckaa, A., Tworzydlo, M., Polit, A., Dziedzicka-

Wasylewska, M. Role of silent polymorphrisms within the dopamine D1 receptor associated

with schizophrenia on D1-D2 receptor heterodimerization. Pharmacol. Rep. 2009; 61: 1024-

1033.

34. Duan, J., Wainwright, M.S., Comeron, J.M., Saitou, N., Sanders, A.R., Gelernter, J., et al.

Synonymous mutations in the human dopamine receptor D2 (DRD2) affect mRNA stability

and synthesis of the receptor. Hum. Mol. Genet. 2003; 12: 205-216.

35. Kimchi-Sarfaty, C., Oh, J.M., Kim, I.-W., Sauna, Z E., Calcagno, A. M., Ambudkar, S. V., et

al. A "silent" polymorphism in the MDR1 gene changes substrate specificity. Science 2007;

315: 525-528.

36. Shah, K., Cheng, Y., Hahn, B., Bridges, R., Bradbury, N.A., Mueller, D.M. Synonymous

codon usage affects the expression of wild type and F508del CFTR. J. Mol. Biol. 2015; 427:

1464-1479.

37. Zhang, T., Wu, Y., Lan, Z., Shi, Q., Yang, Y., Guo, J. Syntool: a novel region-based

intolerance score to single nucleotide substitution for synonymous mutations predictions

based on 12,136 individuals. BioMed Res. Int. 2017; 2017: 5096208.

38. Hunt, R. C., Simhadri, V. L., Iandoli, M., Sauna, Z. E., Kimchi-Sarfaty, C. Exposing

synonymous mutations. Trends Genet. 2014; 30: 308-321.

39. Bali, V., Bebok, Z. Decoding mechanisms by which silent codon changes influence protein

biogenesis and function. Int. J. Biochem. Cell. Bol. 2015; 64: 58-74.

40. Kovacs, Z., Schacht, T., Hermann, A. K., Albrecht, P., Lefkimmiatis, K., Methner, A. Protein

kinase inhibitor β enhances the constitutive activity of G-protein-coupled zinc receptor GPR39.

Biochem. J. 2014; 462: 125-132.

41. Shimizu, Y., Koyama, R., Kawamoto, T. Rho kinase-dependent desensitization of GPR39; a

unique mechanism of GPCR downregulation. Biochem. Pharmacol. 2017; 140: 105-114.

42. Chorin, E., Vinograd, O., Fleidervish, I., Gilad, D., Hermann, S., Sekler, I., et al.

Upregulation of KCC2 activity by zinc-mediated neurotransmission via the mZnR/GPR39

receptor. J. Neurosci. 2011; 31: 12916-12926.

43. Saadi, R. A., He, K., Hartnett, K. A., Kandler, K, Hershfinkel, M., Aizenman, E. SNARE-

dependent upregulation of potassium chloride co-transporter 2 activity after metabotropic zinc

receptor activation in rat cortical neurons in vitro. Neurosci. 2012; 210: 38-46.

44. Gilad, D., Shorer, S., Ketzef, M., Friedman, A., Sekler, I., Aizenman, E., et al. Homeostatic

regulation of KCC2 activity by the zinc receptor mZnR/GPR39 during seizures. Neurobiol.

Dis. 2015; 81: 4-13.

45. Mlyniec, K., Budziszewska, B., Holst, B., Ostachowicz, B., Nowak, G. GPR39 (zinc

receptor) knockout mice exhibit depression-like behavior and CREB/BDNF down-regulation

in the hippocampus. Int. J. Neuropsychopharmacol. 2014; 18: pii pyu002.

46. Mlyniec, K., Nowak, G. Up-regulation of the GPR39 Zn^{2+}-sensing receptor and

CREB/BDNF/TrkB pathway after chronic but not acute antidepressant treatment in the frontal

cortex of zinc-deficient mice. Pharmacol. Rep. 2015; 67: 1135-1140.

47. Mlyniec, K., Doboszewska, U., Szewczyk, B., Sowa-Kucma, M., Misztak, P., Piekoszewski,

W., et al. The involvement of the GPR39-Zn^{2+}-sensing receptor in the pathophysiology of

depression. Studies in rodent models and suicide victims. Neuropharmacol. 2014; 79: 290-

297.

48. Costello, L. C., Franklin, R. B. A comprehensive review of the role of zinc in normal prostate

function and metabolism; and its implications in prostate cancer. Arch. Biochem. Biophys.

2016; 611: 100-112.

49. Zhao, H., Qiao, J., Zhang, S., Zhang, H., Lei, H., Lei, X., et al. GPR39 marks specific cells

within the sebaceous gland and contributes to skin wound healing. Sci. Rep. 2015; 4: 7913.

50. Nakamura, Y., Gojobori, T., Ikemura, T. Codon usage tabulated from international DNA

sequence databases: status for the year 2000. Nucleic Acids Res. 2000; 28: 292.

51. Athey, J., Alexaki, A., Osipova, E., Rostovtsev, A., Santana-Quintero, L. V., Katneni, U., et

al. A new and updated resource for codon usage tables. BMC Bioinformatics 2017; 18: 391.

52. Verma, A., Bradford, Y., Dudek, S., Lucas, A.M., Verma, S.S., et al. A simulation study

investigating power estimates in phenome-wide association studies. BCM Bioinformatics

2018; 19: 120.

53. Verma, A., Lucas, A., Verma, S.S., Zhang, Y., Josyula, N., et al. PheWAS and Beyond: The

landscape of associations with medical diagnoses and clinical measures across 38,662

individuals from Geisinger. Am. J. Hum. Genet. 2018; 102: 592-608.

Table 1. Orphan GPCRs with overlapping disease associations for multiple classes of rare variants. All orphan GPCRs with sufficient numbers of rare LOF, MISS or SYN_ΔCB variants in at least two categories were analyzed by SKAT. Absence of results in a category (marked by X) indicates insufficient number of variants and/or individuals having the variant for well-powered SKAT. A Bonferroni correction for multiple testing (123 sets, 656 ICD9 codes) was used to establish an exploratory significance cut-off of 0.05 (adjusted P-value threshold 6.2E-7). NOS, not otherwise specified.

GPCR	ICD9	DESCRIPTION	LOF	MISS	SYN_ΔCB
LGR4	710.2	Sicca syndrome	1.58E-31	1.28E-13	2.31E-14
GPR150	372.14	Other chronic allergic conjunctivitis	2.59E-31	1.45E-32	5.48E-07
GPR153	276.7	Hyperpotassemia	2.32E-30	8.73E-16	1.14E-22
GPR149	374.3	Ptosis of eyelid	5.89E-30	4.62E-08	1.53E-07
GPR161	780.8	Generalized hyperhidrosis	3.78E-26	X	9.03E-27
GPR37L1	345.1	Generalized convulsive epilepsy, w/o intractable epilepsy	2.04E-18	1.92E-19	4.95E-20
GPR84	280.8	Other specified iron deficiency anemias	2.31E-15	2.99E-08	2.07E-15
GPR39	**354.2**	**Lesion of ulnar nerve**	**8.04E-15**	**1.62E-09**	**8.69E-15**
LGR6	583.81	Nephritis and nephropathy, not specified as acute or chronic	4.52E-11	X	1.31E-19
LGR5	704	Alopecia NOS	5.73E-13	X	1.36E-12
GPR1	287.5	Thrombocytopenia NOS	1.5E-09	1.26E-07	1.5E-09
GPR18	583.81	Nephritis and nephropathy, not specified as acute or chronic	2.61E-08	1.01E-08	3.77E-09
GPR179	270.4	Disturbances of sulphur-bearing amino acid metabolism	1.38E-07	X	1.05E-07
GPR176	493.91	Asthma, unspecified type, with status asthmaticus	X	1.35E-33	6.03E-26
GPR132	216.3	Disturbances of sulphur-bearing amino acid metabolism	X	4.13E-30	9.61E-24
GPR85	410.1	Acute myocardial infarction, of other anterior wall	X	2.06E-26	1.33E-35
GPR183	573.8	Other specified disorders of liver	X	1.34E-19	1.5E-19
GPR12	345.1	Generalized convulsive epilepsy, w/o intractable epilepsy	X	5.64E-14	5.48E-14
GPR135	350.1	Trigeminal neuralgia	X	1.91E-12	1.99E-12
GPR26	372.14	Other chronic allergic conjunctivitis	X	4.56E-12	4.35E-12
MRGPRE	401.1	Benign essential hypertension	X	2.17E-10	8.61E-08
GPR15	455.6	Hemorrhoids NOS	X	1.61E-10	1.97E-10
GPR83	250.8	Diabetes mellitus type II (non-insulin-dependent)	X	1.29E-09	5.71E-12
GPRC5D	588	Renal osteodystrophy	X	7.5E-09	2.0E-20
GPCR5B	381.81	Dysfunction of Eustachian tube	X	1.4E-07	1.23E-07
GPR141	786.07	Wheezing	X	3.17E-09	2.76E-10
GPR6	214.1	Lipoma of other skin and subcutaneous tissue	X	1.82E-07	2.51E-09
GPR31	41.02	Group B Streptococcus infection	X	2.04E-07	2.04E-07

Table 2. SKAT results for rare GPR39 variants in 51,289 DiscovEHR cohort. All LOF plus pathogenic missense (pLP + pP) were binned and analyzed by SKAT (53 markers; 221 individuals). The SYN_ΔCB were separately binned and analyzed (22 markers; 198 individuals). Listed are all significant associations when either/both variant categories had a P-value below the exploratory threshold of 0.05 (adjusted P-value threshold of 3.8E-5; 2 sets, 656 codes). Associations that were below the adjusted P-value threshold for only one variant category are also listed. NOS, not otherwise specified.

CODE	DESCRIPTION	LOF + pLP + pP	SYN_ΔCB
354.2	Lesion of ulnar nerve	6.2E-08	8.69E-15
995.3	Allergy NOS	6.67E-06	7.22E-14
600.21	Benign prostate hyperplasia w urinary obstruction	1.24E-07	2.57E-09
362.56	Macular puckering of retina	X	7.65E-07
786.6	Swelling, mass or lump in chest	3.83E-05	2.4E-07
704.8	Other specified diseases of hair and hair follicles	X	6.72E-07
696	Psoriatic arthropathy	3.24E-13	1.03E-06
727.05	Other tenosynovitis of hand or wrist	X	1.78E-06
372.3	Conjunctivitis NOS	1.68E-05	2.75E-06
998.59	Other postoperative infection	X	5.0E-06
401.1	Benign essential hypertension	X	9.55E-06
70.54	Chronic hepatitis C w/o mention of hepatic coma	X	1.09E-05
655.83	Other known or suspected fetal abnormality	9.71E-06	1.41E-05
591	Hydronephrosis	X	2.06E-05
110.4	Dermatophytosis of foot	1.62E-06	X
726.2	Other affections of shoulder region NOS	4.54-E06	X
429.2	Cardiovascular disease (ASCVD)	2.26E-05	X
707.19	Ulcer of lower limb	2.46E-05	X

Table 3. SKAT analysis of all rare synonymous GPR39 variants in 51,289 DiscovEHR cohort. All rare synonymous variants were binned and analyzed by SKAT (46 markers; 940 individuals). Shown are all associations which were below the exploratory threshold of 0.05 (adjusted P-value 7.62E-05; 1 set, 656 codes). NOS, not otherwise specified.

CODE	DESCRIPTION	P-value
696	Psoriatic arthropathy	3.87E-11
354.2	Lesion of ulnar nerve	1.08E-07
110.4	Dermatophytosis of foot	8.34-E07
726.2	Other affectations of shoulder region, NOS	2.24E-06
995.3	Allergy, NOS	3.51E-06
655.83	Other known or suspected fetal abnormality	4.38E-06
707.19	Ulcer of lower limb	1.06E-05
786.6	Swelling, mass or lump in chest	2.06E-05
429.2	Cardiovascular disease (ASCVD)	3.03E-05
649.13	Obesity complicating pregnancy, childbirth, antepartum	3.58E-05
600.21	Benign prostate hyperplasia with lower urinary tract symptoms	3.79E-05
440.9	Atherosclerosis NOS	6.31E-05
649	Tobacco use disorder complicating pregnancy, childbirth	6.8E-05
368.2	Diplopia	7.46E-05

Figure Legends

Figure 1. Analysis flow for orphan/understudied GPCR SKAT analysis. Rare variants from 85 GPCRs were identified in 51,289 WES dataset and sorted into LOF, missense and synonymous categories. Missense variants were further triaged to identify those predicted to be likely pathogenic or pathogenic by RMPath analysis (Supplemental Methods), and the subset of synonymous variants with large changes in codon usage bias were identified. Those GPCRs with at least 2 categories of variant having $\geq$ 7 distinct members were further analyzed (72 of 85 GPCRs). SKAT analysis was run independently on 123 sets of variants, against 656 disease codes having $\geq$ 200 individuals with $\geq$3 independent encounters in their EHR.

Figure 2. Pathogenicity triage of 159 coding variants in *GPR39* identified in 51,289 WES. Common variants with MAF$\geq$1% were categorized as likely benign (MAF between 1-5%) and benign (MAF$\geq$5%). All variants having MAF<1% sorted to synonymous, loss of function (LOF; frameshift or premature stop codon), and missense. Missense variants were thus Variants of Unknown Significance (VUS) and subjected to bioinformatics pathogenicity triage using RMPath pipeline, and classified as predicted benign, pB, likely benign, pLB, likely pathogenic, pLP or pathogenic pP.

Figure 3. Snake plot of GPR39 topology indicating rare variants identified in 51,289 WES. The GPR39 sequence was color-coded to indicate reference codon usage frequency. For all amino acids having multiple potential codons, red indicates the most common codon, with progression from common to rare colored as gradations to light pink (rare). Arrowheads indicate the locations of rare synonymous variants exhibiting the largest changes in codon bias, either from common to

rare or rare to common. Asterisks indicate rare LOF variants (truncations or frameshifts) and

arrows indicate locations of rare missense variants combined for SKAT analysis.

Figure 4. Manhattan plots of SKAT results for *GPR39* rare variants. A. Rare LOF and MISS

variants were combined (39 total variants) and analysis run using SKAT. Top phenome-wide

associations are labelled. B. Rare SYN_ΔCB variants (21 total variants) were analyzed with

SKAT; top phenome-wide associations are labelled. Full details of associations common to both

variant categories are presented in **Table 2**. For both **A.** & **B.**, the significance threshold was set

at 0.05 (adjusted P-value 3.8E-05, 2 sets, 656 codes).

Figure 5. Expression of rare missense and synonymous GPR39 variants. A. Quantitation of

expression levels of *GPR39* rare variants relative to WT. All blots had WT and HEK293 only

lanes for background subtraction and normalization. Data presented as % of WT expression $\pm$

S.D. (n $\geq$ 3 independent transfections). Representative blots in **Figure S2**. **B.** Quantitation of

expression levels of GPR39 rare SYN_ΔCB variants relative to WT. Details same as in **A**.

Statistical analysis for both figures was two-tailed t-test relative to WT, with *p<0.05, ** p<0.01,

***p<0.001 and ****p<0.0001.

Figure 6. Functional analysis of rare missense and synonymous GPR39 variants. A. IP1

responses to various concentration of $ZnCl_2$ for all variants that expressed $\geq$ 50% of WT.

Representative curves and fits for WT, W314R and R386C are illustrated. Plotted are mean $\pm$ S.D.

of three independent experiments, lines show fits to Michaelis-Menten equation. **B.** EC_{50}s for all

dose-response relations (as in **A**), plotted as mean $\pm$ S.E. for n=3 independent experiments. **C.** IP1

responses of representative SYN_ΔCB variants showing altered $EC_{50}s$ compared to WT, I29 and

V354. Details as in **A**. **D.** $EC_{50}s$ for all dose-response relations (as in **A**) for SYN_ΔCB variants.

E.-F. Bungarotoxin binding (see Methods) was used to assess surface localization of rare missense

(**E**) and SYN_ΔCB (**F**) variants. **G.-H.** Extrapolated $V_{MAX}s$ of Michaelis-Menten fits to IP_1 dose-

response relations (see Methods). For all plots, $*p<0.05$, $**p<0.01$, $***p<0.001$, and

$****p<0.0001$, by two-tailed t-test relative to WT, assuming unequal variances.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Supplemental Methods.

In-silico Radical Mutations Pathogenicity (RMPath) prediction pipeline.

Rare variants are categorized into two groups: (1) nonsense and frame-shift variants, and (2)

missense variants, and were scored by summing numerical values derived from predictions.

(1) Scoring of nonsense and frame-shift variants.

SnpEff fields: LOF[*].PERC ≥ 0.5 or NMD[*].NUMTR > 3 are considered potential Loss of

Function (LOF) variants.

(2) Scoring of missense variants.

1. SIFT scoring: D (damaging) = 2; All others = 0.

2. PolyPhen2 scoring: (i) HDIV: Probably damaging (D) = 2; Possibly damaging (P) = 1; all other

results = 0. (ii) HVAR: Probably damaging (D) = 2; Possibly damaging (P) = 1; all other results =

0.

3. MutationTaster scoring: Disease causing (D) = 2; all other results = 0.

4. MutationAssessor scoring: High (H) = 2; Medium (M) = 1; all other results = 0.

5. FATHMM scoring: D (damaging) = 2; all other results = 0.

6. LRT: D(deleterious) → score 2; All others → score 0.

7. PROVEAN scoring: D(damaging) = 2; all other results = 0.

8. GRANTHAM scoring: ≥140 = 4; ≥120 = 3; ≥90 = 2; ≥70 = 1; all other results = 0.

9. Genomic Evolutionary Rate Profiling GERP++ rejected substitutions (RS) scoring: ≥6 = 4; ≥5

= 3; ≥3.5 = 2; ≥2 = 1; all other results = 0.

Scores from all tools were summed to generate a cumulative score (= RMPath score), with a

maximum possible value of 24. Variants with RMPath scores ≤8 were predicted to be Benign

(pB), those with scores in the range from 6 to 11 were predicted to be Likely Benign (pLB); those

with scores between 11 and 17 were predicted to be Likely Pathogenic (pLP); and those with scores

≥17 were predicted to be Pathogenic (pP). For clinical purposes, all variants are considered

Variants of Unknown Significance (VUS) until family co-segregation and/or functional analyses

definitively establish a variant as pathogenic.

Supplemental Table 1. Demographics of the DiscovEHR cohort used in the present study.

Abbreviations, EHR, electronic health record, BMI, body mass index.

Basic demographics	DiscoveEHR Sequenced Patients
Total patients (N)	51,289
Female, N (%)	30,290 (59%)
Median age, yrs	61 (range 48-72)
Median BMI, kg/m^2	30 (range 26-36)
Median years of EHR data	14 (range 11-17)
Median medication orders/patient	129 (range 37-221)

Supplemental Figure S1. Manhattan plots of results of PheWAS analyses of common

GPR39 variants. No association results exceeded the minimal P-value threshold of 0.05,

uncorrected for multiple testing.

Supplemental Figure S2. Representative western blots of rare missense and synonymous

variants identified in the 51,289 WES cohort. Western blots of HEK293 cells lysates expressing

WT or rare variant missense or SYN_ΔCB cDNAs were run as described in Methods. **A.** Missense

variants run in order from amino to carboxyl terminal location; each gel also had WT and

untransfected HEK samples for normalization; GAPDH was used as loading control. **B.**

Expression of rare synonymous variants was more variable, and rare variants were sorted for high

expression (top row) or low expression (bottom row) for better quantitation; WT and HEK were

run on each gel for normalization; GAPDH was used as loading control.