Mesolimbic dopamine projections mediate cue-motivated reward seeking but not

reward retrieval

Briac Halbout[1*], Andrew T. Marshall[1], Ali Azimi[1], Mimi Liljeholm[2], Stephen V. Mahler[3], Kate M. Wassum[4], Sean B. Ostlund[1*]

1. *Department of Anesthesiology and Perioperative Care and Irvine Center for Addiction Neuroscience, University of California, Irvine, CA, USA*
2. *Department of Cognitive Sciences, University of California, Irvine, CA, USA*
3. *Department of Neurobiology & Behavior and Irvine Center for Addiction Neuroscience, University of California, Irvine, CA, USA*
4. *Department of Psychology and Brain Research Institute University of California, Los Angeles, CA, USA.*

Corresponding authors (*):
Briac Halbout
University of California, Irvine
837 Health Science Road, 2216 GNRF
Irvine, CA 92697
Email: halboutb@uci.edu
Tel: 949-824-9826

Sean Ostlund
University of California, Irvine
837 Health Science Road, 3111GNRF
Irvine, CA 92697
Email: sostlund@uci.edu
Tel: 949-824-1347

Running Title: VTA-dopamine neurons mediate cue-triggered reward seeking

Abstract 150

Article: 7173
    Introduction: 794
    Materials and Methods: 2558
    Results: 2255
    Discussion: 1544
Figures count: 5
References count: 70
Supplementary Materials: 1

Abstract

Efficient foraging requires an ability to coordinate discrete reward-seeking and reward-retrieval behaviors. We used pathway-specific chemogenetic inhibition to investigate how mesolimbic and mesocortical dopamine circuits contribute to the expression and modulation of reward seeking and retrieval. Inhibiting ventral tegmental area dopamine neurons disrupted the tendency for reward-paired cues to motivate reward seeking, but spared their ability to increase attempts to retrieve reward. Similar effects were produced by inhibiting dopamine inputs to nucleus accumbens, but not medial prefrontal cortex. Inhibiting dopamine neurons spared the suppressive effect of reward devaluation on reward seeking, an assay of goal-directed behavior. Attempts to retrieve reward persisted after devaluation, indicating they were habitually performed as part of a fixed action sequence. Our findings show that complete bouts of reward seeking and retrieval are behaviorally and neurally dissociable from bouts of reward seeking without retrieval. This dichotomy may prove useful for uncovering mechanisms of maladaptive behavior.

Foraging and other reward-motivated behaviors tend to unfold as a sequence of

actions, beginning with a *reward-seeking* phase and ending with an attempt to retrieve

and consume any rewards produced by this activity. Coordinating the discrete reward-

seeking and reward-retrieval behaviors that make up these action sequences is

important for efficient foraging. When rewards are sparse or otherwise difficult to obtain,

attempts to retrieve them are often unnecessary and should therefore be withheld to

conserve energy and minimize opportunity costs (Stephens and Krebs, 1986; Niv et al.,

2007). Consistent with this, studies on self-paced instrumental behavior show that the

ability to efficiently pattern reward-seeking and -retrieval responses based on task

demands (e.g., reinforcement schedule) can strongly impact the rate at which rewards

are obtained (Ostlund et al., 2012; Wassum et al., 2012; Matamales et al., 2017).

However, such behaviors must remain sensitive to changes in internal and external

states. For instance, environmental cues that signal reward availability increase

attempts to seek out (Estes, 1948; Corbit and Balleine, 2016) and retrieve reward

(Marshall and Ostlund, 2018). While the ability to develop and modify action sequences

is normally adaptive, this process may become dysregulated in certain conditions, such

as obsessive-compulsive disorder (Joel and Avisar, 2001; Korff and Harvey, 2006;

Frederick and Cocuzzo, 2017) and drug addiction (Tiffany, 1990; Graybiel, 2008;

Volkow et al., 2013), leading to maladaptive behaviors. Despite this, the behavioral and

neural mechanisms responsible for regulating reward seeking and retrieval are not well

understood.

Previous studies strongly implicate dopamine in learning new action sequences

(Graybiel, 1998; Jin and Costa, 2015). While other findings suggest that dopamine is

not as important for the expression of well-established action sequences (Levesque et

al., 2007; Wassum et al., 2012), it remains possible that dopamine contributes to action

sequence performance when changes in task conditions prompt a reorganization of

reward seeking and retrieval. For instance, previous studies indicate that the tendency

for reward-paired cues to motivate reward-*seeking* behavior critically depends on

dopamine signaling (Dickinson et al., 2000; Ostlund and Maidment, 2011; Wassum et

al., 2011), particularly in the nucleus accumbens (NAc)(Wyvell and Berridge, 2000; Lex

and Hauber, 2008; Wassum et al., 2013; Ostlund et al., 2014; Aitken et al., 2016).

Interestingly, we recently found that such cues do not simply provoke reward-seeking

behavior (e.g., lever pressing), they also increase the likelihood that such behavior will

be followed by an attempt to retrieve reward (e.g., food-cup approach)(Marshall and

Ostlund, 2018). Although this finding suggests that reward-paired cues preferentially

motivate complete bouts of reward seeking and retrieval, it has yet to be established if

this modulation of action sequence performance depends on dopamine.

Dopamine may also contribute to regulating attempts to seek out and retrieve a

reward when the value of that reward changes. Self-paced, instrumental reward-seeking

actions are normally performed in a goal-directed manner, such that they are sensitive

to changes in reward value (Balleine and Dickinson, 1998). However, they can develop

into inflexible stimulus-response habits with extended training (Dickinson, 1985;

Dickinson et al., 1995). In contrast, it is not well understood how changes in reward

value modulate attempts to retrieve rewards produced through instrumental reward-

seeking behavior. For example, it has been suggested that rats' tendency to approach

the food cup after lever pressing may represent a discrete goal-directed action – one

that is selected independently of the initial decision to press the lever (Rescorla, 1964).

Alternatively, rats may concatenate the press-approach sequence to form an *action*

chunk, which can then be selected and deployed as a single unit of behavior (Lashley,

1951; Graybiel, 1998; Jin and Costa, 2015). Action chunks are thought to represent a

special form of habit, or behavioral chain, in which each element of the chain

automatically elicits the next response. This allows for efficient action sequencing but

comes with a decrease in behavioral flexibility. Once an action chunk has been initiated,

it should be automatically completed without further consideration of reward value

(Dezfouli et al., 2014; Smith and Graybiel, 2016).

In the current study, we applied a chemogenetic approach to investigate the role

of the mesocorticolimbic dopamine system in action sequence performance in rats. We

used a combination of well-established behavioral assays and novel microstructural

analyses to selectively probe the influence of reward-paired cues and expected reward

value on the regulation of reward-seeking and -retrieval responses. We found that

inhibiting dopamine neurons in the ventral tegmental area (VTA) or their inputs to the

NAc, but not the medial prefrontal cortex (mPFC), reversibly disrupted cue-motivated

reward seeking, but spared the tendency for reward-paired cues to trigger complete

bouts of seeking and retrieval. These dopamine manipulations had no impact on rats'

tendency to adjust their reward-seeking behavior in response to reward devaluation.

Importantly, attempts to retrieve reward were not suppressed by reward devaluation,

suggesting that this behavior was the product of action chunking.

Materials and Methods

Animals: In total, 89 male and female Long-Evans Tyrosine hydroxylase

(Th):Cre+ rats (hemizygous Cre+) (Witten et al., 2011; Mahler et al., 2019) and wildtype

(WT) littermates were used for this study. Subjects were at least 3 months of age at the

start of the experiment and were single- or paired-housed in standard Plexiglas cages

on a 12h/12h light/dark cycle. Animals were maintained at ~85% of their free-feeding

weight during behavioral procedures. All experimental procedures that involved rats

were approved by the UC Irvine Institutional Animal Care and Use Committee and were

in accordance with the National Research Council Guide for the Care and Use of

Laboratory Animals.

Apparatus: Behavioral procedures took place in sound- and light-attenuated Med

Associates chambers (St Albans, VT, USA; ENV-007). Individual chambers were

equipped with two retractable levers (Med Associates; ENV-112CM) positioned to the

left and right of recessed food cup. Grain-based dustless precision pellets (45 mg,

BioServ, Frenchtown, NJ, USA) were delivered into the cup using a pellet dispenser

(Med Associates; ENV-203M-45). Sucrose solution (20% wt/vol) was delivered into the

cup with a syringe pump (Med Associates; PHM-100). A photobeam detector (Med

Associates; ENV-254-CB) positioned across the magazine entrance was used to record

food-cup approaches. Chambers were illuminated by a houselight during all sessions.

Surgery: Th:Cre+ rats were anesthetized using isoflurane and placed in a

stereotaxic frame for microinjections of a Cre-dependent (DIO) serotype 2 adeno-

associated virus (AAV) vectors to induce dopamine neuron-specific expression of the

inhibitory designer receptor exclusively activated by designer drug (DREADD) hM4Di

fused to mCherry (AAV-hSyn-DIO-hM4Di-mCherry), or mCherry alone (AAV-hSyn-DIO-

mCherry) (University of North Carolina Chapel Hill vector Core, Chapel Hill, NC, USA /

Addgene, Cambridge, MA, USA; Experiment 2 was replicated with both sources)

(Witten et al., 2011; Mahler et al., 2019). The AAV was injected bilaterally into the VTA

(-5.5 mm AP, +/- 0.8 mm ML, -8.15 mm DV; 1uL / side). Experiment 3 rats were

bilaterally implanted with guide cannulae (22 gauge, Plastic One) 1 mm dorsal to NAc

(+1.3 AP, +/-1.8 ML, -6.2 DV) or mPFC (+3.00 AP, +/-0.5 ML, -3.0 DV) for subsequent

clozapine-n-oxide (CNO) microinjections. Animals were randomly assigned to virus

(hM4Di or mCherry) and cannula location (NAc or mPFC) groups. Animals were allowed

at least 5 days of recovery before undergoing food restriction and behavioral training.

Testing occurred at least 25 days after surgery to allow adequate time for viral

expression of hM4Di throughout dopamine neurons, including in terminals within the

NAc and mPFC.

Experiment 1: Effects of response-contingent feedback about reward delivery on reward

retrieval

Instrumental learning. WT rats (n = 9) underwent 2 d of magazine training. In

each session, 40 pellets were delivered into the food cup on a random 90-sec intertrial

interval (ITI). Rats then received 9 d of instrumental lever-press training. In each

session, rats had continuous access to the right lever, which could be pressed to deliver

food pellets into the food cup. The schedule of reinforcement was adjusted over days

from continuous reinforcement (CRF) to increasing random intervals (RI), such that

reinforcement only became available once a randomly determined interval had elapsed

since the last reinforcer delivery. Rats received one day each of CRF, RI-15s, and RI-

30s training, before undergoing 6 days of training with RI-60s. Each session was

terminated after 30 min or after 20 rewards deliveries.

Varying response-contingent feedback. Following training, rats were given a

series of tests to assess the influence of response-contingent feedback about reward

delivery on instrumental reward-seeking (lever presses) and reward-retrieval responses

(press-contingent food-cup approach). Rats were given 3 tests (30-min each,

pseudorandom order over days) during which lever pressing caused: 1) activation of the

pellet dispenser to deliver a pellet into the food cup (RI-60s schedule; *Food and Cues*

Test), 2) activation of the pellet dispenser to deliver a pellet into an external cup not

accessible to the rats, producing associated sound and tactile cues but no reward (also

RI-60s schedule; *Cues Only Test*), or 3) no dispenser activation (i.e., extinction; *No*

Food or Cues Test).

Experiments 2 and 3: Role of mesocorticolimbic dopamine in cue-motivated reward

seeking and retrieval:

Pavlovian conditioning. Th:Cre+ rats (n = 60) underwent 2 d of magazine training,

as in Experiment 1 (40 pellets on 90s random ITI). Rats then received 8 daily Pavlovian

conditioning sessions. Each session consisted of a series of 6 presentations of a two-min

audio cue (CS+; either a pulsating 2 kHz pure tone (0.1s on and 0.1s off) or white noise;

80dB), with trials separated by a 5-min variable ITI (range 4-6-min between CS onsets).

During each CS+ trial, pellets were delivered on a 30-s random time schedule, resulting

in an average of 4 pellets per trial. Rats were separately habituated to an unpaired

auditory stimulus (CS-; alternative audio stimulus; 2-min duration). CS- exposure

procedures differed slightly across experiments. For Experiment 2, which assessed the

effects of system-wide dopamine neurons inhibition, rats received a final Pavlovian

conditioning session consisting of four trials with the CS+ (reinforced, as described above)

followed by four trials with the CS- (nonreinforced), separated by a 5-min variable ITI. In

Experiment 3, which assessed the effects of local inhibition of dopamine terminals in NAc

or mPFC, rats were given 2 days of CS- only exposure (8 nonreinforced trials per session,

5-min variable ITI) following initial CS+ training. Conditioning was measured by

comparing the rate of food-cup approach between the CS onset and the first pellet

delivery (to exclude unconditioned behavior) to the rate of approach during the pre-CS

period.

Instrumental training. Following Pavlovian conditioning, rats were given 9 d of

instrumental training, as in Experiment 1, with one day each of CRF, RI-15s, RI-30s,

and 6 days of RI-60s. Sessions ended after 30 min or 20 rewards were earned.

Pavlovian-to-instrumental transfer (PIT) test. After the last instrumental training

session, rats were given a session of Pavlovian (CS+) training, identical to initial

training. They were then given a 30-min extinction session, during which lever presses

were recorded but had no consequence (i.e., no food or cues). On the next day, rats

were given a PIT test, during which the lever was continuously available but produced

no rewards. Following 8 min of extinction, the CS+ and CS− were each presented 4

times (2 min per trial) in pseudorandom order and separated by a 3-min fixed ITI. Before

each new round of testing, rats were given two sessions of instrumental retraining (RI-

60s), one session of CS+ retraining, and one 30-min extinction session, as described

above. Test procedures differed slightly between Experiments 2 and 3.

Experiment 2: Th:Cre+ rats expressing hM4Di (n = 18) or mCherry only (n = 14)

in VTA dopamine neurons were used to assess the effects of system-wide inhibition of

the mesocorticolimbic dopamine system on PIT performance. These groups were run

together and received CNO (5mg/kg, i.p.) or vehicle (5% DMSO in saline) injections 30

min prior to testing. They underwent a second test following retraining (described

above), prior to which the alternative drug pretreatment was administered.

Experiment 3: In Experiment 3A, Th:Cre+ rats expressing hM4Di in VTA

dopamine neurons were used to assess the impact of locally inhibiting dopaminergic

terminals in the NAc (n = 7) or mPFC (n = 9) on PIT performance. Because

microinjection procedures produced additional variability in task performance, rats in

this experiment underwent a total of 4 tests. Rats received either CNO microinfusions

(1mM, 0.5uL/side or 0.3uL/side, for NAc and mPFC respectively) or vehicle (DMSO 5%

in aCSF) 5 min before the start of each test and were given two rounds of testing each

with CNO and vehicle (test order counterbalanced across other experimental

conditions). To determine if the effects of CNO microinjections depended on hM4Di

expression, a separate control study (Experiment 3B) was run using Th:Cre+ rats

expressing mCherry only in VTA dopamine neurons. Experiments 3A and 3B were run

and analyzed separately.

Experiment 4: *Role of mesocorticolimbic dopamine in goal-directed action selection*:

Instrumental Training. Th:Cre+ rats expressing hM4Di (n = 11) or mCherry only

(n = 9) in VTA dopamine neurons began with 2 d of magazine training, during which

they received 20 grain-pellets and 20 liquid sucrose rewards (0.1 mL of 20% sucrose

solution, wt/vol) in random order according to a common 30s random ITI. This was

followed by 11 d of instrumental training with two distinct action–outcome contingencies

(e.g., left-lever press → grain; right-lever press→ sucrose). The reinforcement schedule

that was gradually shifted over days with 2d of CRF to increasingly effortful random ratio

(RR) schedules, with 3 d of RR-5, 3 d of RR-10, and 3d of RR-20 reinforcement. The

left and right lever-press responses were trained in separate sessions, at least 2 h

apart, on each day. Action-outcome contingencies were counterbalanced across

subjects. Sessions were terminated after 30 min elapsed or 20 pellets were earned.

Devaluation Testing. To selectively devalue one of the food rewards prior to

testing, rats were satiated on grain pellets or sucrose solution by providing them with 90

min of unrestricted access to that food in the home cage. After 60 min of feeding, rats

received CNO (5mg/kg, i.p.) or vehicle injections. After an additional 30 min of feeding,

rats were placed in the chamber for a test in which they had continuous access to both

levers. The test began with a 5-min nonreinforced phase (no food or cues), which was

immediately followed by a 15-min reinforced phase, during which each action was

reinforced with its respective reward (CRF for the first 5 rewards, then RR-20 for the

remainder of the session). Rats were given a total of 4 devaluation tests, 2 after CNO

and 2 after vehicle, alternating the identity of the devalued reward across the 2 tests in

each drug condition (test order counterbalanced across training and drug conditions).

Histology: Rats were deeply anesthetized with a lethal dose of pentobarbital and

perfused with 1x PBS followed by 4% paraformaldehyde. Brains were postfixed in 4%

paraformaldehyde, cryoprotected in 20% sucrose and sliced at 40 μm on a cryostat. To

visualize hM4Di expression, we performed immunohistochemistry for Th and mCherry

tag. Tissue was first incubated in 3% normal donkey serum PBS plus Triton X-100

(PBST; 2 h) and then in primary antibodies in PBST at 4°C for 48 hours using rabbit

anti-DsRed (mCherry tag; 1:500; Clontech; 632496), and mouse anti-Th (1:1,000,

Immunostar; 22941) antibodies. Sections were incubated for 4 h at room temperature in

fluorescent conjugated secondary antibodies (Alexa Fluor 488 goat anti-mouse (Th;

1:500; Invitrogen; A10667) and Alexa Fluor 594 goat anti-rabbit (DsRed; 1:500;

Invitrogen; A11037)).

Drugs: CNO was obtained from NIMH (Experiments 2 and 4) or Sigma-Aldrich (St.

Louis, MO, USA; Experiment 3), and dissolved in 5% DMSO in saline, or aCSF for

microinjection.

Behavioral measures:

Reward-*seeking* actions were quantified as the total number (frequency) of lever

presses performed per unit time. Based on microstructural analyses described below,

lever presses that were followed by a food-cup approach (≤ 2.5 sec) were distinguished

from presses that were not followed by an approach. The proportion of presses that

were followed by an approach response served as our primary measure of press-

contingent reward retrieval. We also analyzed bouts of noncontingent food-cup

approach (occurring > 2.5 sec after the most recent press or approach), which served

as a measure of spontaneous or cue-evoked reward retrieval.

Statistical analysis:

Data were analyzed using general(ized) linear mixed-effects models (Pinheiro

and Bates, 2000), which allows for simultaneous parameter estimation as a function of

condition (fixed effects) and the individual rat (random effects) (Pinheiro and Bates,

2000; Bolker et al., 2008; Boisgontier and Cheval, 2016). Analyses on count data (e.g.,

response frequency) incorporated a Poisson response distribution and a log link

function (Coxe et al., 2009). Fixed-effects structures included an overall intercept and

the full factorial of all primary manipulations (Experiment 2: Group, Drug, CS Type, CS

Period; Experiment 3: Site, Drug, CS Type, CS Period; Experiment 4: Group, Drug,

Lever), and the random-effects structures included by-subjects uncorrelated intercepts

adjusted for the within-subjects manipulations (i.e., Experiments 2 and 3: Drug, CS

Type, and CS Period; Experiment 4: Drug, Lever). "CS Type" refers to the distinction

between the CS+ and CS-, while "CS Period" refers to the distinction between the 120-s

CS duration and the 120-s period preceding its onset. Proportion data were square-root

transformed prior to analysis to correct positive skew, but are plotted in non-transformed

space for ease of interpretation. These data were collapsed across pre-CS+ and pre-

CS- periods, such that the factor "CS Period" had 3 levels (CS+, CS-, and Pre-CS). The

fixed- and random-effects structures of this analysis was identical to the frequency

analysis above with the exception that CS Type was not included in the analysis, and

the random-effects structure only included by-subjects intercepts.

All statistical analyses were conducted using the Statistics and Machine Learning

Toolbox in MATLAB (The MathWorks; Natick, MA, USA). The alpha level for all tests

was .05. As all predictors were categorical in the mixed-effects analysis, effect size was

represented by the unstandardized regression coefficient (Baguley, 2009), reported as *b*

in model output tables. Mixed-effects models provide *t*-values to reflect the statistical

significance of the coefficient relative to the population mean (i.e., simple effects).

These simple effects are indicative of main effects and interactions when a factor has

only two levels. For factors with at least 3 levels, *F*-tests were conducted to reveal the

overall significance of the effect or interaction(s) involving this factor. The source of

significant interactions was determined by secondary mixed-effects models identical to

those described above but split by the relevant factor of interest. For analyses in which

a significant main effect had more than two levels, post-hoc tests of main effects

employed MATLAB's *coefTest* function, and interactions were reported in-text as the

results of ANOVA *F*-tests (i.e., whether the coefficients for each fixed effect were

significantly different from 0).

When analyzing data from PIT experiments, the ability of the CS+ to selectively

increase performance of a response (relative to the CS-) over baseline (pre-CS) levels

was indicated by a significant CS Type * CS Period interaction. We were particularly

interested in treatment-induced alterations in the expression of this effect, as indicated

by significant 3-way and 4-way interactions involving this CS Type * CS Period term, in

combination with Drug and/or Group factors. We were also interested in potential main

effects of Drug and/or Group factors, reflecting broad, cue-independent behavioral

effects. While statistical output tables include a summary of all fixed effects included in

the model, only these theoretically interesting findings are discussed in the main text.

Lower level interactions involving only CS Type *or* CS Period, but not their combination,

are provided in the output tables but are not discussed in the main text given that they

may be the product of incidental or spurious behavioral differences across cue

conditions.

PIT Scores (CS+ – pre-CS+) were calculated for more focused analysis of CS+

elicited lever pressing. One-sample t-tests were used to assess the effect of CNO for

each group. Because inhibiting VTA dopamine neurons or their NAc terminals

predominantly disrupted the ability of the CS+ to elicit lever presses that were not

followed by an approach response, we also assessed if differences across rats in their

tendency to exhibit such behavior in the Vehicle Test (PIT score; presses without

approach) correlated with differences in their sensitivity to the response-suppressive

effect of CNO on CS+ elicited lever pressing (CNO – Vehicle; PIT score, all presses).

Results

Effects of response-contingent feedback about reward delivery on reward

retrieval

We first characterized the relationship between reward-seeking and -retrieval

responses when rewards are sparse (Figure 1A). Rats were trained to lever press on a

RI-60s schedule, such that this action was often nonreinforced and only occasionally

earned food pellet delivery into a recessed food cup. Not surprisingly, we found that the

probability of food-cup approach was elevated for several seconds after performance of

the lever-press action (Figures 1B and 1C). This timeframe for press-contingent food-

cup approach behavior is consistent with previous reports (Nicola, 2010; Marshall and

Ostlund, 2018), and was relatively consistent across the current experiments (see

Figure 3-Supplement 1). We therefore used a cutoff value of 2.5 sec to identify reward-

retrieval attempts. To control for reward-retrieval *opportunities*, which were contingent

on lever pressing, our analysis focuses on a normalized measure – the proportion of

lever presses that were followed by food-cup approach.

We found that rats were much more likely to approach the food cup after

reinforced presses than after nonreinforced presses ($t(8) = 19.33$, $p < .001$), suggesting

they could detect when pellets were delivered based on sound and tactile cues

produced by the dispenser. This was confirmed in subsequent tests, during which lever

pressing produced either 1) pellet dispenser cues and actual pellet delivery (Food and

Cues), 2) pellet dispenser cues only (Cues Only), or 3) no pellet dispenser cues or

pellet delivery (No Food or Cues). Here too, we found that food-cup approaches were

more likely after reinforced than nonreinforced lever presses, regardless of whether

pellet dispenser cues were presented alone or together with actual food delivery (Figure

1F; $ts(8) \geq 13.74$, $ps < .001$; the overall frequency of lever pressing (Figure 1D) and the

frequency of complete bouts of presses that were followed by an approach (Figure 1E)

are presented for comparison). Although pellet dispenser cues were clearly an effective

trigger for rats to shift from the lever to the food cup, they also made these shifts

spontaneously, indicating that they had developed the tendency to perform the

complete press-approach action sequence. These unprompted approaches occurred

after a relatively small subpopulation of nonreinforced lever presses, which is consistent

with our previous data (Marshall and Ostlund, 2018).

Figure 1. Microstructural organization of instrumental behavior. **A.** Hungry rats were trained to perform a self-paced "reward seeking" task, in which pressing a lever was intermittently reinforced with food pellets (RI-60s schedule). Press-contingent food-cup approaches were taken as a measure of attempted "reward retrieval". **B.** Probability of food-cup approaches as a function of time surrounding reinforced (purple) and nonreinforced (gray) lever presses. **C.** Representative pattern of food-cup approach behavior for an individual rat surrounding reinforced and nonreinforced lever presses. Individual reinforced trials are separately presented across the y-axis aligned at the point at which the lever became activated (i.e., primed for reinforcement). **D, E.** Effects of manipulating instrumental reinforcement contingency on the organization of reward-seeking and -retrieval responses. Total lever presses (D) or presses followed by an approach (E) during tests in which lever pressing was intermittently reinforced (RI-60s) either with food pellets and associated cues (Food and Cues) or with pellet dispenser cues but no actual food delivery (Cues Only). Rats were also tested without any reinforcement (No Food or Cues). **F.** The proportion of lever presses that were followed by food-cup approach was higher for reinforced presses than for nonreinforced presses, regardless of whether pressing was reinforced with Food and Cues, or Cues Only. Rats also continued to sporadically check the food cup after nonreinforced lever presses, albeit at a much lower level than after reinforced presses.

Inhibiting dopamine neurons during Pavlovian-to-instrumental transfer

preferentially disrupts cue-motivated reward seeking, but not reward retrieval

Our previous findings suggest that reward-predictive cues *both* invigorate

reward-seeking behavior (i.e., the PIT effect) and increase the likelihood that such

actions will be followed by an attempt to retrieve reward from the food cup (Marshall and

Ostlund, 2018). Experiment 2 investigated the contributions of the mesocorticolimbic

dopamine system to these distinct behavioral effects of reward-paired cues.

Rats with dopamine neuron-specific expression of the inhibitory DREADD hM4Di

or mCherry in the VTA (Figure 2) were trained on a PIT task (Figure 3A) consisting of a

Pavlovian conditioning phase, in which two different auditory cues were paired (CS+) or

unpaired (CS-) with food pellets, and a separate instrumental training phase, in which

rats were trained to lever press for pellets. During PIT testing, we noncontingently

presented the CS+ and CS- while rats were free to lever press and check the food cup

without response-contingent food or cue delivery.

Figure 2. DREADD expression in Th:Cre+ rats. **A.** Th:Cre+ rats received bilateral injections of AAV-hSyn-DIO-hM4Di-mCherry or AAV-hSyn-DIO-mCherry in the VTA. **B.** Representative expression of the mCherry-tagged inhibitory DREADD hM4Di (red) in VTA Th positive neurons (green) of Th:Cre+ rats, as well as in neuronal terminals (**C.**) projecting to the nucleus accumbens (NAc) and medial prefrontal cortex (mPFC). Scale bar is 500um.

We found that rats selectively increased their lever press performance during

CS+ presentations, relative to the CS- and pre-CS response rates (Figure 3B; CS

Period * CS Type interaction, $p < .001$; see Supplementary File Table 1 for full

generalized linear mixed-effects model output). This effect was significantly attenuated

by CNO in a group-specific manner (Group * Drug * CS Period * CS Type interaction, p

= .002). Analysis of data from CS+ trials (only) found that CNO selectively suppressed

cue-induced lever pressing in hM4Di relative to mCherry rats (Drug * Group * CS Period

interaction, p = .013). Further analysis found that the mCherry group displayed a

pronounced increase in lever pressing during CS+ trials (CS Period * CS Type

interaction, $p <$ .001), and this effect was not altered by CNO (Drug * CS Period * CS

Type interaction, p = .780). In contrast, CNO pretreatment significantly disrupted

expression of CS+ induced lever pressing in the hM4Di group (Drug * CS Period * CS

Type interaction, $p <$ .001). hM4Di rats showed a CS+ specific elevation in lever

pressing when pretreated with vehicle (CS Period * CS Type interaction, $p <$ .001) but

not CNO (CS Period * CS Type interaction, p = .684). While these findings indicate that

CNO selectively disrupted the response-invigorating influence of the CS+ by inhibiting

VTA dopamine neurons in hM4Di rats, there was also some indication that CNO may

have produced a nonspecific, group-independent, suppression of PIT performance

(Drug x CS Period x CS Type, p = .007). We therefore conducted a more focused

analysis of CS+ induced changes in lever-press performance (PIT score: CS+ - pre-

CS+; Figure 3C), which confirmed that CNO significantly suppressed this behavioral

effect in the hM4Di group (t(17) = -3.83, $p <$ .001), but not in the mCherry group (t(13)

= -1.21, p = .249). This is in line with recent findings that similar CNO treatment does

not significantly alter PIT performance in DREADD-free rats (Collins et al., *in press*).

We also investigated if VTA dopamine neuron inhibition impacts the tendency for

the CS+ to increase attempts to retrieve reward after performing the reward-seeking

response (Figures 3D and 3E; see Figure 3-Supplement 1 for illustration of the

probability of food-cup approach surrounding lever presses during nonreinforced PIT

trials). We found that the CS+ ($p < .001$) but not the CS- ($p = .501$) increased the

proportion of lever presses that were followed by a food-cup approach, even though no

rewards were actually delivered at test (see Supplementary File Table 2 for full

generalized linear mixed-effects model output; see Figure 3-Supplements 2 and 3 for

analysis of total press-contingent and noncontingent approaches, respectively).

Importantly, CNO did not alter this response to the CS+ in a group-specific manner

(Group * Drug * CS+ Period, $p = .835$), indicating that VTA dopamine neuron function is

not required for this behavior. However, CNO did induce some nonspecific, group-

independent alterations in the proportion of presses that were followed by food-cup

approach, lowering the overall likelihood of this behavior (Drug effect, $p = .019$), but

enhancing the effect of the CS+ (Drug * CS+ Period, $p < .037$).

Figure 3. Chemogenetic inhibition of dopamine neurons on Pavlovian to instrumental transfer (PIT) performance. **A.** Experimental design: Following viral vectors injections and recovery, rats received Pavlovian training, during which they learned to associate an auditory cue (CS+) with food pellet delivery. During instrumental conditioning, rats performed the same lever-press task used in Experiment 1. Lever pressing was extinguished (Ext) before rats were submitted to a PIT test, which included separate noncontingent presentations of the CS+ and an unpaired control cue (CS-). **B.** Chemogenetic inhibition of VTA dopamine neurons disrupted cue-motivated reward seeking. Total lever presses during PIT trials for rats expressing the inhibitory DREADD hM4Di or mCherry following vehicle (left) or CNO (5mg/kg, right) treatment prior to test. Presses during pre-CS (gray) and CS periods (red) are plotted separately. **C.** PIT expression is specifically impaired in hM4Di expressing Th:Cre+ rats. PIT scores (total presses: CS+ - pre-CS+) show that the CS+ increased lever pressing after vehicle treatment for both groups, but that CNO suppressed this effect in the hM4Di group but not the mCherry group. **p<0.01. **D.** The CS+ increased the proportion of lever presses that were followed by a food-cup approach during PIT testing. Inhibiting VTA dopamine neurons did not disrupt expression of this effect. Instead, rats in both groups showed a modest increase in their likelihood of checking the food cup after lever pressing when treated with CNO. **E.** Representative organization of the effects of the CS+ and CS- on attempts to seek out and retrieve reward during PIT. Data show lever presses and food-cup approaches (press-contingent or noncontingent) for two control rats (Th:Cre+ rats expressing mCherry and receiving vehicle).

Pathway-specific inhibition of dopamine projections to NAc, but not mPFC,

disrupts cue-motivated reward seeking but not retrieval

As previously reported (Mahler et al., 2019), hM4Di expression in VTA dopamine

neurons resulted in transport of DREADDs to axonal terminals in the NAc and mPFC

(Figure 2). We took advantage of this to investigate the roles of these two pathways in

PIT performance, again distinguishing between the influence of reward-paired cues on

reward seeking and reward retrieval. Guide cannulae were aimed at the NAc or mPFC

in rats expressing hM4Di in VTA dopamine neurons (Experiment 3A; Figure 4A and

Figure 4-Supplement 1). These rats underwent training and testing for PIT (Figure 4B),

as described above, but were pretreated with intra-NAc or mPFC injections of CNO

(1mM) or vehicle to achieve local inhibition of neurotransmitter release (Mahler et al.,

2014; Stachniak et al., 2014; Lichtenberg et al., 2017), an approach previously shown to

be effective in inhibiting dopamine release (Mahler et al., 2019). Figure 4C shows that,

in hM4Di-expressing rats, the CS+ specific increase in lever pressing (CS Period * CS

Type interaction, $p < .001$) was disrupted by CNO in a manner that depended on

microinjection site (Drug * CS Period * CS Type * Site interaction, $p = .003$;

Supplementary File Table 3 for full generalized linear mixed-effects model output). After

intracranial vehicle injections, rats showed a CS+ specific elevation in pressing (CS

Period * CS Type interaction, $p < .001$), which did not differ significantly across vehicle

injection sites (CS Period * CS Type * Site interaction, $p = .151$). Unlike with systemic

CNO, the CS+ remained effective in increasing lever pressing after CNO microinjection

into the mPFC (CS Type * CS Period interaction, $p < .001$) and NAc (CS Type * CS

Period interaction, $p < .001$). However, this effect was significantly attenuated when

CNO was injected into the NAc versus the mPFC (CS Period * CS Type * Site

interaction, p = .012; analysis of CNO data only). A more focused analysis of CS+

elicited lever pressing (Fig 4D; PIT score) confirmed that CNO disrupted this effect in

the NAc group ($t(6)$ = -2.49, p = .047), but not in the mPFC group ($t(8)$ = 0.34, p = .746).

The disruptive effect of intra-NAc CNO administration on PIT performance did not

systematically vary as a function of injection site (data not presented), which is not

surprising given previous findings that this effect is modulated by dopamine signaling in

both the core and shell of the NAc (Lex and Hauber, 2008; Pecina and Berridge, 2013).

We also conducted a separate experiment (Experiment 3B) with rats expressing

the mCherry reporter (only) in VTA dopamine neurons to determine if this behavioral

effects of CNO microinfusion was hM4Di-dependent. While there was evidence that

CNO may have produced some nonspecific response suppression when injected into

the mPFC but not the NAc (Drug * Site * CS Period * CS Type, p = .068), this drug

treatment did not significantly disrupt expression of CS+ elicited lever pressing for either

injection site (p's > .165; Figure 4-Supplement 2).

Figure 4. Pathway specific chemogenetic inhibition of dopamine on PIT performance. **A.** Th:Cre+ rats initially received VTA AAV-hSyn-DIO-hM4Di-mCherry injections and were implanted with guide cannulas aimed at the medial prefrontal cortex (mPFC) or nucleus accumbens (NAc) for microinjection of CNO (1mM) or vehicle to inhibit dopamine terminals at test. **B.** Following surgery, rats underwent training and testing for PIT, as described above. We analyzed the microstructural organization of behavior (Lever presses: seeking, and presses followed by a food-cup approach: retrieval) at test. **C.** Pathway specific inhibition of dopamine terminals in the NAc but not the mPFC disrupted cue-motivated reward seeking. Total lever presses during PIT trials for rats expressing the inhibitory DREADD hM4Di and receiving CNO or vehicle microinfusions in either the mPFC or NAc prior to test. Presses during pre-CS (gray) and CS periods (red) are plotted separately. **D.** PIT expression was specifically impaired following NAc CNO treatment. PIT scores (total presses: CS+ - pre-CS+) show that the CS+ increased lever pressing following vehicle treatment in both groups, but that CNO suppressed this effect when injected into the NAc but not the mPFC. *p<0.05. **E.** The CS+ increased the proportion of lever presses that were followed by a food-cup approach during PIT testing. This effect did not significantly vary as a function of drug treatment or group. **F.** Scatter plots show the relationship between individual differences in the effect of the CS+ on lever presses that were not followed by food-cup approach in the vehicle condition (Δ Presses without approach) and the suppressive effect of CNO on CS+ evoked lever pressing (PIT Score for vehicle test - PIT Score for CNO test). Data points are from individual rats receiving intra-mPFC (left panel) or intra-NAc (right panel) microinjections.

As in the previous experiment, we found that the CS+ ($p < .001$) increased the

proportion of lever presses that were followed by an attempt to retrieve reward from the

food cup (Figure 4E; Supplementary File Table 4 for full generalized linear mixed-effects

model output; see Figure 4–Supplements 3 and 4 for analysis of total press-contingent

and noncontingent approaches, respectively). CNO seemed to generally reduce the

likelihood that lever pressing would be followed by food-cup approach, though this effect

did not reach statistical significance (Drug effect, $p = .057$). If anything, intra-NAc

injections of CNO tended to *enhance* the effect of the CS+ on this approach response,

though this effect also failed to reach significance (Drug * Site * CS+ Period, $p = .093$).

The above findings indicate that VTA dopamine circuitry supports the

motivational influence of the CS+ on reward seeking but does not mediate that cue's

ability to promote reward retrieval. We wondered if this might account for variability in

the partial, response-suppressive effect of NAc dopamine terminal inhibition.

Specifically, we hypothesized that rats inclined to respond to the CS+ by engaging in

discrete bouts of lever pressing, without attempting to retrieve reward, would be

particularly sensitive to inhibition of NAc dopamine inputs. Consistent with this, we

found that for the NAc group, individual differences in the effect of the CS+ on lever

presses without subsequent food cup approach (during the vehicle test) were negatively

correlated with CNO-induced suppression of CS+ evoked lever pressing (all presses; r

$= -0.81$, $p = .027$; Figure 4F). No such relationship was found for the mPFC group (r

$= -0.19$, $p = .618$), which did not show sensitivity to dopamine terminal inhibition. Similar

analysis of data from Experiment 2 also found no correlation between these measures

(Figure 4-Supplement 5), which may not be surprising given that systemic inhibition of

VTA dopamine neurons led to a more robust and consistent suppression of CS+ evoked

lever pressing (Figure 3B).

Altogether, these findings demonstrate that the mesolimbic dopamine system

selectively mediates cue-motivated reward seeking, and suggest that dopamine inputs

to the NAc are particularly important for individuals that tend to respond to such cues

with discrete bouts of reward seeking without subsequent reward retrieval.

Inhibiting dopamine neurons spares the sensitivity of reward-seeking actions to

reward devaluation

It is unclear from the above findings if rats' tendency to approach the food cup

after lever pressing reflects a discrete goal-directed action or if this response tends to

be performed habitually, as part of a fixed press-approach action chunk. We conducted

a reward devaluation experiment to probe this issue and investigate the role of VTA

dopamine neurons in goal-directed action selection. Rats expressing mCherry or hM4Di

in VTA dopamine neurons were trained on two distinct instrumental action-outcome

contingencies, after which they underwent reward devaluation testing after pretreatment

with CNO (5mg/kg) or vehicle (Figure 5A). Rats performed significantly fewer presses

on the devalued lever than on the valued lever (Figure 5B; Lever effect, $p < .001$;

Supplementary File Table 5 for full generalized linear mixed-effects model output). CNO

treatment did not significantly alter the effect of reward devaluation on lever pressing in

either hM4Di or mCherry rats (Drug * Lever, $p = .146$; Group * Drug * Lever interaction,

$p = .591$), indicating that VTA dopamine neuron function is not required for this aspect of

goal-directed action selection. Inhibiting VTA dopamine neurons also failed to disrupt

sensitivity to devaluation during reinforced testing (see Figure 5-Supplement 1).

VTA dopamine neuron inhibition did not significantly alter the overall likelihood of

press-contingent approach behavior or its sensitivity to reward devaluation (Figure 5C;

ps ≥ .109; see Supplementary File Table 6 for full generalized linear mixed-effects

model output). Interestingly, we found that the proportion of presses that were followed

by a food-cup approach was actually *greater* for the devalued lever than for the valued

lever (Lever effect, p = .040). This effect was driven by the fact that lever presses that

were not followed by approach were more strongly suppressed by reward devaluation

than presses that were directly followed by an approach (Press Type * Lever interaction,

p < .001; see Figure 5D and Supplementary File Table 7 for full generalized linear

mixed-effects model output).

Figure 5. Chemogenetic inhibition of dopamine neurons on reward devaluation performance. **A.** Th:Cre+ rats received VTA injections of AAV-hSyn-DIO-hM4Di-mCherry or AAV-hSyn-DIO-mCherry. Following recovery, rats were trained on two distinct lever-press actions for two different rewards (Instrumental Learning). Rats then underwent reward-specific devaluation testing following treatment with CNO (5mg/kg) or vehicle. **B.** Chemogenetic VTA dopamine inhibition did not alter the impact of reward devaluation on reward seeking. Total lever presses on the valued (red bars) and devalued (gray) levers in hM4Di or mCherry expressing Th:Cre+ rats, following CNO (5 mg/kg) or vehicle treatments. **C.** Proportion of valued (blue) and devalued (gray) lever-press actions that were followed by a food-cup approach. Rats were more likely to attempt to retrieve reward after performing the devalued lever-press action. This effect was not altered by VTA dopamine neuron inhibition. **D.** Lever presses performed without a subsequent food-cup approach response (red) were more sensitive to reward devaluation than presses that were not followed by an approach (blue).

____

Discussion

We investigated the role of mesocorticolimbic dopamine circuitry in regulating

reward-seeking (lever pressing) and reward-retrieval responses (press-contingent food-

cup approach). Consistent with a recent study (Marshall and Ostlund, 2018), we found

that noncontingent CS+ presentations increased reward seeking, generally, but also

increased the likelihood that rats would attempt to retrieve reward after performing such

actions. These behaviors were differentially mediated by the mesolimbic dopamine

system. Specifically, chemogenetic inhibition of VTA dopamine neurons or their inputs

to NAc, but not mPFC, disrupted the excitatory influence of the CS+ on reward seeking,

but spared that cue's ability to increase attempts to retrieve reward. These behaviors

were also differentially sensitive to reward devaluation, which suppressed reward

seeking but actually increased the likelihood that rats would attempt to retrieve reward.

VTA dopamine neurons inhibition did not impact the influence of reward devaluation on

either component of behavior.

We found that attempts to retrieve reward by transitioning from the lever to the

food cup were executed in a habitual manner, without consideration of reward value,

consistent with action chunking (Dezfouli et al., 2014; Smith and Graybiel, 2016).

However, task performance was not limited to these press-approach action chunks.

When rats pressed the lever but were not reinforced (with food or cues), they would

occasionally check the food cup but often omitted this response. This sporadic pattern

of reward retrieval is adaptive given that strict press-approach action sequencing is

unnecessary under such conditions, when rewards are sparse and uncertain. Instead,

rats seemed to vacillate between two different strategies when initiating the lever-press

response, performing it as part of a complete action chunk (press-approach) or as a

discrete action (press only). These distinct patterns of reward seeking appeared to be

differentially sensitive to reward devaluation. While rats were generally less likely to

lever press for the devalued reward than for the valued reward, press-approach action

chunks tended to be less sensitive to reward devaluation than presses that were not

followed by approach. Because of this differential sensitivity to reward devaluation, the

proportion of all lever presses followed by an attempt to retrieve reward was actually

greater for devalued action than for the valued action. Such findings supports the

connection between action chunking and habitual behavior (Graybiel, 2008; Dezfouli et

al., 2014; Smith and Graybiel, 2016), and suggest that moment-to-moment control over

self-paced, reward-seeking behavior may shift back and forth between habit and goal-

directed systems.

PIT testing revealed that the CS+ generally increased lever pressing, but

disproportionately increased the performance of press-approach action chunks, at least

relative to their otherwise low frequency of occurring in the absence of the CS+. This

finding further bolsters the connection between action chunking and habitual control

given previous reports that habitual reward-seeking actions are particularly sensitive to

the motivational effects of reward-paired cues (Holland, 2004; Wiltgen et al., 2012).

However, while press-approach action chunks were elevated during the CS+, they still

accounted for only a minority (between 30% and 50%) of lever presses that were

performed during these trials. Most lever presses evoked by the CS+ were *not* followed

by a food-cup approach, and it was this component of the PIT effect that was selectively

disrupted by chemogenetic inhibition of VTA dopamine neurons or their inputs to NAc.

The ability of the CS+ to promote press-approach chunks was, in contrast, completely

spared by these manipulations. Consistent with this, we found that the response-

suppressive effect of NAc dopamine terminal inhibition varied across rats based on the

way they normally responded to the CS+. Rats that responded to that cue with a large

increase in discrete lever presses (i.e., *without* subsequent food-cup approach) showed

the greatest suppression. We suggest that this may reflect differences across rats in

their sensitivity to the dopamine-dependent motivational effects of reward-paired cues.

Previous studies have found that dopamine receptor antagonists either

selectively suppress lever pressing without affecting concomitant food-cup approach

(Nelson and Killcross, 2013), or suppress both types behavior to a similar extent

(Wassum et al., 2011; Ostlund et al., 2012). Even this latter finding is consistent with

dopamine contributing more to reward seeking than reward retrieval, since a reduction

in reward seeking creates fewer opportunities to retrieve reward. Interpreting these

findings is problematic, however, because such studies typically have not applied

microstructural analyses, like those used here, to distinguish between press-contingent

and noncontingent food-cup approaches. One exception is a study by Nicola (2010)

showing that blocking dopamine receptors in the NAc attenuates cue-triggered lever

pressing without impacting the latency of subsequent food-cup approach behavior.

Building on such findings, the current study used the PIT paradigm to show that the

mesolimbic dopamine system specifically mediates the motivational influence of reward-

paired cues on reward seeking but not their dissociable ability to increase the likelihood

that such actions will be followed by an attempt to retrieve reward.

Our previous studies monitoring mesolimbic dopamine release during PIT

performance are also interesting to consider together with the current findings. For

instance, we found that CS+ evoked phasic dopamine release in the NAc correlates

with that cue's effect on lever pressing (Wassum et al., 2013; Ostlund et al., 2014) but

not food-cup approaches (Aitken et al., 2016). We also found that individual CS+

evoked lever presses are temporally correlated with transient bouts of phasic dopamine

release (Ostlund et al., 2014). The current findings suggest that this relationship

between NAc dopamine release and cue-motivated reward seeking may be stronger for

discrete presses that are performed without a subsequent food-cup approach than for

complete press-approach chunks. This question remains to be investigated, and would

help resolve whether the mesolimbic dopamine system is involved in modulating reward

seeking, generally, or whether its activity becomes uncoupled from the execution of

action chunks, which may become differentially associated with nigrostriatal dopamine

system activity (Jin and Costa, 2010).

While dopamine is known to play a crucial role in forming new action chunks

(Graybiel, 1998; Jin and Costa, 2015), its role in the expression of previously learned

action chunks is less clear. Our findings indicate that VTA dopamine circuitry does not

play a necessary role in the execution of press-approach action chunks, regardless of

whether they are self-initiated or are prompted by a reward-paired cue. This is generally

compatible with previous findings. For instance, dopamine receptor blockade

suppresses action sequence performance early but not late in training (Levesque et al.,

2007; Wassum et al., 2012). Moreover, the phasic NAc dopamine release that normally

precedes action sequence performance tends to become attenuated as rats acquire

efficient task performance, presumably through action chunking (Cacciapaglia et al.,

2012; Wassum et al., 2012; Klanker et al., 2015; Collins et al., 2016). That said, the

mesolimbic dopamine system continues to contribute to action sequence tasks that

require considerable effort, such as the execution of a long series of lever presses

(Fischbach-Weiss et al., 2017).

Inhibiting VTA dopamine neurons did not impact rats' sensitivity to reward

devaluation, which is consistent with other findings in the literature (Dickinson et al.,

2000; Lex and Hauber, 2009; Lex and Hauber, 2010; Wassum et al., 2011). Such

findings are interesting given that regions innervated by this dopamine system, including

the NAc and mPFC, are known to make important contributions to goal-directed

decision making (Bradfield and Balleine, 2017; Sharpe et al., 2019). Of course,

dopamine likely contributes to goal-directed decision making in more demanding tasks

that require greater cognitive resources (Floresco, 2013; Cools, 2015; Westbrook and

Braver, 2016).

It is also notable that inhibiting mPFC dopamine terminals had no detectable

effects on expression of PIT, since food-paired cues are known to elicit dopamine

release (Bassareo and Di Chiara, 1997; Feenstra et al., 1999) and neural activity

(Homayoun and Moghaddam, 2009) in the mPFC. It is possible that the dissociable

effects of NAc versus mPFC dopamine terminal inhibition reported here may relate to

inherent differences between the mesolimbic and mesocortical dopamine systems,

which include regional differences in release kinetics and in the density of dopamine

terminals or receptors (Lammel et al., 2008; Vander Weele et al., 2018; Mahler et al.,

2019). However, previous lesion studies suggest that the mPFC may not be an

essential component of the circuitry that mediates PIT performance (Cardinal et al.,

2003; Corbit and Balleine, 2003), which is more in line with the current results.

Our findings may also have implications for understanding the role of dopamine

in pathologies of behavioral control such as obsessive-compulsive disorder (OCD). In

the signal attenuation model of OCD (Joel and Avisar, 2001), rats learn that response-

contingent cues no longer signal that an instrumental reward-seeking action will

produce reward. In this case, the logical organization of reward-seeking and -retrieval

actions disintegrates, such that rats exhibit persistent reward seeking, typically without

attempting to collect reward from the food cup. It was previously reported that blocking

D1-dopamine receptors disrupts expression of these incomplete bouts of compulsive-

like reward seeking, without affecting the production of complete bouts of reward

seeking and retrieval, which continue to be performed on some test trials (Joel and

Doljansky, 2003). Considered in this light, our findings suggest that the mesolimbic

dopamine system may mediate the tendency for reward-paired cues to promote this

potentially compulsive component of cue-motivated reward seeking. This link deserves

further research, and may facilitate research to advance understanding and treatment of

compulsive disorders like OCD and addiction (Joel et al., 2008; Robinson et al., 2014).

Funding and Disclosure

Our work is supported by National Institutes of Health (NIH) grants MH106972, DK098709, AG045380, and DA029035. The authors declare no competing interests.

Acknowledgment

The authors acknowledge the assistance of Christy N. Munson in the acquisition of behavioral data.

References

Aitken TJ, Greenfield VY, Wassum KM (2016) Nucleus accumbens core dopamine signaling tracks the need-based motivational value of food-paired cues. Journal of neurochemistry 136:1026-1036.

Baguley T (2009) Standardized or simple effect size: what should be reported? British Journal of Psychology 100:603-617.

Balleine BW, Dickinson A (1998) Goal-directed instrumental action: contingency and incentive learning and their cortical substrates. Neuropharmacology 37:407-419.

Bassareo V, Di Chiara G (1997) Differential influence of associative and nonassociative learning mechanisms on the responsiveness of prefrontal and accumbal dopamine transmission to food stimuli in rats fed ad libitum. J Neurosci 17:851-861.

Boisgontier MP, Cheval B (2016) The anova to mixed model transition. Neuroscience & Biobehavioral Reviews 68:1004-1005.

Bolker BM, Brooks ME, Clark CJ, Geange SW, Poulsen JR, Stevens MHH, White J-SS (2008) Generalized linear mixed models: a practical guide for ecology and evolution. Trends in Ecology & Evolution 24:127-135.

Bradfield L, Balleine B (2017) The Learning and Motivational Processes Controlling Goal-Directed Action and Their Neural Bases. In: Decision Neuroscience, pp 71-80: Elsevier.

Cacciapaglia F, Saddoris MP, Wightman RM, Carelli RM (2012) Differential dopamine release dynamics in the nucleus accumbens core and shell track distinct aspects of goal-directed behavior for sucrose. Neuropharmacology 62:2050-2056.

Cardinal RN, Parkinson JA, Marbini HD, Toner AJ, Bussey TJ, Robbins TW, Everitt BJ (2003) Role of the anterior cingulate cortex in the control over behavior by Pavlovian conditioned stimuli in rats. Behav Neurosci 117:566-587.

Collins AL, Greenfield VY, Bye JK, Linker KE, Wang AS, Wassum KM (2016) Dynamic mesolimbic dopamine signaling during action sequence learning and expectation violation. Scientific reports 6.

Collins AL, Aitken TJ, Huang IW, Shieh C, Greenfield VY, Monbouquette HG, Ostlund SB, Wassum KM (*in press*) Nucleus accumbens cholinergic interneurons oppose cue-motivated behavior. Biological Psychiatry.

Cools R (2015) The cost of dopamine for dynamic cognitive control. Current Opinion in Behavioral Sciences 4:152-159.

Corbit LH, Balleine BW (2003) The role of prelimbic cortex in instrumental conditioning. Behav Brain Res 146:145-157.

Corbit LH, Balleine BW (2016) Learning and Motivational Processes Contributing to Pavlovian-Instrumental Transfer and Their Neural Bases: Dopamine and Beyond. Curr Top Behav Neurosci 27:259-289.

Coxe S, West SG, Aiken LS (2009) The analysis of count data: A gentle introduction to Poisson regression and its alternatives. Journal of Personality Assessment 91:121-136.

Dezfouli A, Lingawi NW, Balleine BW (2014) Habits as action sequences: hierarchical action control and changes in outcome value. Philosophical Transactions of the Royal Society B: Biological Sciences 369:20130482.

Dickinson A (1985) Actions and habits: the development of behavioural autonomy. Philosophical Transactions of the Royal Society of London Series B, Biological Sciences:67-78.

Dickinson A, Smith J, Mirenowicz J (2000) Dissociation of Pavlovian and instrumental incentive learning under dopamine antagonists. Behavioral neuroscience 114:468.

Dickinson A, Balleine B, Watt A, Gonzales F, Boakes R (1995) Overtraining and the motivational control of instrumental action. Anim Learn Behav 22:197-206.

Estes WK (1948) Discriminative conditioning. II. Effects of a Pavlovian conditioned stimulus upon a subsequently established operant response. Journal of experimental psychology 38:173.

Feenstra MG, Teske G, Botterblom MH, De Bruin JP (1999) Dopamine and noradrenaline release in the prefrontal cortex of rats during classical aversive and appetitive conditioning to a contextual stimulus: interference by novelty effects. Neurosci Lett 272:179-182.

Fischbach-Weiss S, Reese RM, Janak PH (2017) Inhibiting Mesolimbic Dopamine Neurons Reduces the Initiation and Maintenance of Instrumental Responding. Neuroscience.

Floresco SB (2013) Prefrontal dopamine and behavioral flexibility: shifting from an "inverted-U" toward a family of functions. Front Neurosci 7:62.

Frederick MJ, Cocuzzo SE (2017) Contrafreeloading in Rats Is Adaptive and Flexible: Support for an Animal Model of Compulsive Checking. Evol Psychol 15:1474704917735937.

Graybiel AM (1998) The basal ganglia and chunking of action repertoires. Neurobiology of learning and memory 70:119-136.

Graybiel AM (2008) Habits, Rituals, and the Evaluative Brain. Annual Review of Neuroscience 31:359-387.

Holland PC (2004) Relations between Pavlovian-instrumental transfer and reinforcer devaluation. Journal of Experimental Psychology: Animal Behavior Processes 30:104.

Homayoun H, Moghaddam B (2009) Differential representation of Pavlovian–instrumental transfer by prefrontal cortex subregions and striatum. European Journal of Neuroscience 29:1461-1476.

Jin X, Costa RM (2010) Start/stop signals emerge in nigrostriatal circuits during sequence learning. Nature 466:457.

Jin X, Costa RM (2015) Shaping action sequences in basal ganglia circuits. Current Opinion in Neurobiology 33:188-196.

Joel D, Avisar A (2001) Excessive lever pressing following post-training signal attenuation in rats: a possible animal model of obsessive compulsive disorder? Behavioural brain research 123:77-87.

Joel D, Doljansky J (2003) Selective Alleviation of Compulsive Lever-Pressing in Rats by D 1, but not D 2, Blockade: Possible Implications for the Involvement of D 1 Receptors in Obsessive–Compulsive Disorder. Neuropsychopharmacology 28:77.

Joel D, Stein DJ, Schreiber R (2008) CHAPTER 6 - Animal Models of Obsessive–Compulsive Disorder: From Bench to Bedside via Endophenotypes and

Biomarkers A2 - McArthur, Robert A. In: Animal and Translational Models for CNS Drug Discovery (Borsini F, ed), pp 133-164. San Diego: Academic Press.

Klanker M, Sandberg T, Joosten R, Willuhn I, Feenstra M, Denys D (2015) Phasic dopamine release induced by positive feedback predicts individual differences in reversal learning. Neurobiology of Learning and Memory 125:135-145.

Korff S, Harvey BH (2006) Animal models of obsessive-compulsive disorder: rationale to understanding psychobiology and pharmacology. The Psychiatric clinics of North America 29:371-390.

Lammel S, Hetzel A, Häckel O, Jones I, Liss B, Roeper J (2008) Unique properties of mesoprefrontal neurons within a dual mesocorticolimbic dopamine system. Neuron 57:760-773.

Lashley KS (1951) The problem of serial order in behavior: Bobbs-Merrill.

Levesque M, Bedard MA, Courtemanche R, Tremblay PL, Scherzer P, Blanchet PJ (2007) Raclopride-induced motor consolidation impairment in primates: role of the dopamine type-2 receptor in movement chunking into integrated sequences. Experimental Brain Research 182:499-508.

Lex A, Hauber W (2008) Dopamine D1 and D2 receptors in the nucleus accumbens core and shell mediate Pavlovian-instrumental transfer. Learn Mem 15:483-491.

Lex B, Hauber W (2009) The role of dopamine in the prelimbic cortex and the dorsomedial striatum in instrumental conditioning. Cerebral cortex 20:873-883.

Lex B, Hauber W (2010) The role of nucleus accumbens dopamine in outcome encoding in instrumental and Pavlovian conditioning. Neurobiol Learn Mem 93:283-290.

Lichtenberg NT, Pennington ZT, Holley SM, Greenfield VY, Cepeda C, Levine MS, Wassum KM (2017) Basolateral Amygdala to Orbitofrontal Cortex Projections Enable Cue-Triggered Reward Expectations. The Journal of Neuroscience 37:8374-8384.

Mahler SV, Vazey EM, Beckley JT, Keistler CR, McGlinchey EM, Kaufling J, Wilson SP, Deisseroth K, Woodward JJ, Aston-Jones G (2014) Designer receptors show role for ventral pallidum input to ventral tegmental area in cocaine seeking. Nat Neurosci 17:577-585.

Mahler SV, Brodnik ZD, Cox BM, Buchta WC, Bentzley BS, Quintanilla J, Cope ZA, Lin EC, Riedy MD, Scofield MD (2019) Chemogenetic Manipulations of Ventral Tegmental Area Dopamine Neurons Reveal Multifaceted Roles in Cocaine Abuse. Journal of Neuroscience 39:503-518.

Marshall AT, Ostlund SB (2018) Repeated cocaine exposure dysregulates cognitive control over cue-evoked reward-seeking behavior during Pavlovian-to-instrumental transfer. Learning & Memory 25:399-409.

Matamales M, Skrbis Z, Bailey MR, Balsam PD, Balleine BW, Gotz J, Bertran-Gonzalez J (2017) A corticostriatal deficit promotes temporal distortion of automatic action in ageing. Elife 6.

Nelson AJD, Killcross S (2013) Accelerated habit formation following amphetamine exposure is reversed by D1, but enhanced by D2, receptor antagonists. Frontiers in neuroscience 7:76.

Nicola SM (2010) The flexible approach hypothesis: unification of effort and cue-responding hypotheses for the role of nucleus accumbens dopamine in the activation of reward-seeking behavior. J Neurosci 30:16585-16600.

Niv Y, Daw ND, Joel D, Dayan P (2007) Tonic dopamine: opportunity costs and the control of response vigor. Psychopharmacology 191:507-520.

Ostlund SB, Maidment NT (2011) Dopamine Receptor Blockade Attenuates the General Incentive Motivational Effects of Noncontingently Delivered Rewards and Reward-Paired Cues Without Affecting Their Ability to Bias Action Selection. Neuropsychopharmacology 37:508.

Ostlund SB, Kosheleff AR, Maidment NT (2012) Relative Response Cost Determines the Sensitivity of Instrumental Reward Seeking to Dopamine Receptor Blockade. Neuropsychopharmacology 37:2653.

Ostlund SB, LeBlanc KH, Kosheleff AR, Wassum KM, Maidment NT (2014) Phasic mesolimbic dopamine signaling encodes the facilitation of incentive motivation produced by repeated cocaine exposure. Neuropsychopharmacology 39:2441-2449.

Pecina S, Berridge KC (2013) Dopamine or opioid stimulation of nucleus accumbens similarly amplify cue-triggered 'wanting' for reward: entire core and medial shell mapped as substrates for PIT enhancement. Eur J Neurosci 37:1529-1540.

Pinheiro J, Bates D (2000) Mixed-effects models in S and S-Plus. New York: Springer.

Rescorla RA (1964) Relation of bar-presses to magazine approaches. Psychological Reports 14:943-948.

Robinson TE, Yager LM, Cogan ES, Saunders BT (2014) On the motivational properties of reward cues: individual differences. Neuropharmacology 76:450-459.

Sharpe MJ, Stalnaker T, Schuck NW, Killcross S, Schoenbaum G, Niv Y (2019) An Integrated Model of Action Selection: Distinct Modes of Cortical Control of Striatal Decision Making. Annual Review of Psychology 70:null.

Smith KS, Graybiel AM (2016) Habit formation. Dialogues in clinical neuroscience 18:33.

Stachniak TJ, Ghosh A, Sternson SM (2014) Chemogenetic synaptic silencing of neural circuits localizes a hypothalamus-->midbrain pathway for feeding behavior. Neuron 82:797-808.

Stephens DW, Krebs JR (1986) Foraging theory: Princeton University Press.

Tiffany ST (1990) A cognitive model of drug urges and drug-use behavior: role of automatic and nonautomatic processes. Psychological review 97:147.

Vander Weele CM, Siciliano CA, Tye KM (2018) Dopamine tunes prefrontal outputs to orchestrate aversive processing. Brain research.

Volkow ND, Wang GJ, Tomasi D, Baler RD (2013) Unbalanced neuronal circuits in addiction. Curr Opin Neurobiol 23:639-648.

Wassum KM, Ostlund SB, Maidment NT (2012) Phasic mesolimbic dopamine signaling precedes and predicts performance of a self-initiated action sequence task. Biological psychiatry 71:846-854.

Wassum KM, Ostlund SB, Balleine BW, Maidment NT (2011) Differential dependence of Pavlovian incentive motivation and instrumental incentive learning processes on dopamine signaling. Learning & memory 18:475-483.

Wassum KM, Ostlund SB, Loewinger GC, Maidment NT (2013) Phasic mesolimbic dopamine release tracks reward seeking during expression of Pavlovian-to-instrumental transfer. Biol Psychiatry 73:747-755.

Westbrook A, Braver TS (2016) Dopamine Does Double Duty in Motivating Cognitive Effort. Neuron 89:695-710.

Wiltgen BJ, Sinclair C, Lane C, Barrows F, Molina M, Chabanon-Hicks C (2012) The effect of ratio and interval training on Pavlovian-instrumental transfer in mice. PloS one 7:e48227.

Witten IB, Steinberg EE, Lee SY, Davidson TJ, Zalocusky KA, Brodsky M, Yizhar O, Cho SL, Gong S, Ramakrishnan C, Stuber GD, Tye KM, Janak PH, Deisseroth K (2011) Recombinase-driver rat lines: tools, techniques, and optogenetic application to dopamine-mediated reinforcement. Neuron 72:721-733.

Wyvell CL, Berridge KC (2000) Intra-accumbens amphetamine increases the conditioned incentive salience of sucrose reward: enhancement of reward "wanting" without enhanced "liking" or response reinforcement. J Neurosci 20:8122-8130.

Figure 3-Supplement 1. Probability of food-cup approaches as a function of time surrounding individual lever-press responses during PIT testing, plotted separately for CS+ (blue), CS- (red), and pre-CS (yellow) periods. This analysis was restricted to vehicle tests completed by rats in Experiments 2 and 3A. Shewhart process control chart analyses were used to determine the times when food-cup approach behavior was elevated with respect to constant background rates. This value approximated 2.5 s following each lever press, as shown by the shaded box.

Figure 3-Supplement 2. Frequency of lever presses that were followed by a food-cup approach during PIT testing by rats expressing the inhibitory DREADD hM4Di or mCherry following vehicle or CNO (5mg/kg) treatment in Experiment 2. Data from pre-CS (gray) and CS (blue) periods are plotted separately. Error bars represent ±1 standard error of the estimated marginal means from the corresponding fitted generalized linear mixed-effects model. We found that the CS+ but not the CS- strongly increased the frequency of these press-approach sequences (CS Period * CS Type interaction, $t(240) = 7.87$, $p <$ .001), which did not vary as a function of Group or Drug treatment (interactions involving these factors, ps ≥ .18). However, while CNO administration did not disrupt the ability of the CS+ to elicit these press-approach sequences, it did result in a general, hM4Di-independent reduction in the frequency of these sequences (Drug effect, $t(240) = -4.52$, $p <$ .001; Drug * Group interaction, $t(240) = -1.19$, $p = .234$).

Figure 3-Supplement 3. Noncontingent (press-independent) food-cup approaches during PIT testing by rats expressing the inhibitory DREADD hM4Di or mCherry following vehicle or CNO (5mg/kg) treatment in Experiment 2. Data from pre-CS (gray) and CS (green) periods are plotted separately. Error bars represent ±1 standard error of the estimated marginal means from the corresponding fitted generalized linear mixed-effects model. This behavior was selectively increased by the CS+ relative to the CS- (CS Period * CS Type interaction, $t(240) = 11.72$, $p < .001$). CNO produced a nonspecific (hM4Di-independent) suppression in this approach response (Drug effect, $t(240) = -1.99$, $p = .047$), which was more pronounced for the mCherry group (Group * Drug * CS Type * CS Period interaction, $t(240) = 2.73$, $p = .0067$).

Figure 4-Supplement 1. Cannulae placements for Experiment 3A hM4Di expressing rats. Individual placements in nucleus accumbens (NAc) and medial prefrontal cortex (mPFC).

Figure 4-Supplement 2. Frequency of lever presses during PIT testing by rats expressing mCherry following microinjection of vehicle (**A**) or CNO (**B**) into the mPFC or NAc in Experiment 3B. Data from pre-CS (gray) and CS (red) periods are plotted separately. Error bars represent ±1 standard error of the estimated marginal means from the corresponding fitted generalized linear mixed-effects model. The CS+ induced a cue-specific increase in lever pressing (CS Type * CS Period interaction, $t(176) = 4.51$, $p <$ .001). Rats in the NAc group exhibited a marginally weaker CS+ specific increase in lever pressing than rats in the mPFC group (Site * CS Type * CS Period interaction, $t(176) = -$ 1.89, $p = .060$). CNO appeared to slightly attenuate CS+ elicited lever pressing, particularly when injected into the mPFC, though this effect was not significant (effect was not significant (Drug * CS Period * CS Type interaction, $t(176) = -0.0022$, $p = .998$; Site * Drug * CS Period * CS Type interaction, $t(176) = 1.83$, $p = .068$). **C.** A focused analysis of CS+ elicited lever pressing (PIT Score; ± 1 between-subjects SEM) confirmed that CNO injections did not significantly disrupt this effect in either the mPFC ($t(5) = 1.63$, $p =$ .165) or NAc ($t(5) = 0.33$, $p = .753$) group. These results indicate that the tendency for intra-NAc CNO injections to disrupt CS+ elicited lever pressing in hM4Di expressing rats in Experiment 3A (see Figure 4, main text) was due to dopamine terminal inhibition and not a nonspecific CNO effect.

Figure 4-Supplement 3. Frequency of presses that were followed by a food-cup approach during PIT testing by rats expressing the inhibitory DREADD hM4Di following microinjection of CNO or vehicle into the mPFC or NAc in Experiment 3A. Data from pre-CS (gray) and CS (blue) periods are plotted separately. Error bars represent ±1 standard error of the estimated marginal means from the corresponding fitted generalized linear mixed-effects model. The cue-induced increase in the frequency of these press-approach sequences was specific to the CS+ (CS Type * CS Period interaction, $t(240) = 7.34$, $p <$ .001), which did not vary as a function of drug and/or injection site ($ps > .689$), nor were there any main effects of these treatment variables (Drug effect, $t(240) = 1.05$, $p = .294$; Site effect, $t(240) = -0.26$, $p = .798$; Drug * Site interaction, $t(240) = 1.28$, $p = .203$).

Figure 4-Supplement 4. Noncontingent (press-independent) food-cup approaches during PIT testing in rats expressing the inhibitory DREADD hM4Di following microinjection of CNO or vehicle into the mPFC or NAc in Experiment 3A. Data from pre-CS (gray) and CS (green) periods are plotted separately. Error bars represent ±1 standard error of the estimated marginal means from the corresponding fitted generalized linear mixed-effects model. Noncontingent approach behavior was selectively elicited by the CS+ but not the CS- (CS Period * CS Type interaction, $t(240)$ = 12.98, p < .001). CNO administration led to a very modest but reliable *enhancement* in the tendency for the CS+ to increase this behavior over baseline levels (Drug * CS Type * CS Period interaction, $t(240)$ = 2.98, p = .0032), which did not depend on injection site (Drug * Site * CS Type * CS Period interaction, $t(240)$ = -1.45, p = .147). CNO did not have any general (cue-independent) effects on this response (Drug effect, $t(240)$ = -0.41, p = .680; Site effect, $t(240)$ = 0.75, p = .456; Drug * Site interaction, $t(240)$ = -0.96, p = .338).

Figure 4-Supplement 5. Scatter plots show the relationship between individual

differences in the effect of the CS+ on lever presses that were not followed by food-cup approach in the vehicle condition (Δ Presses without approach) and the suppressive effect of CNO on CS+ evoked lever pressing (PIT Score for vehicle test - PIT Score for CNO test). Data points are from individual rats expressing mCherry (left panel) or hM4Di (right panel).

Figure 5-Supplement 1. Data from the reinforced phase of reward devaluation testing for rats expressing the inhibitory DREADD hM4Di or mCherry following vehicle or CNO treatment in Experiment 4. (**A**) Significantly fewer lever presses were performed on the devalued lever compared to the valued lever, $t(148) = -5.55$, $p < .001$, which did not interact with Group or Drug, $ps ≥ .095$. (**B**) Similarly, the frequency of presses that were followed by a food-cup approach was lower on the devalued versus the valued lever, $t(148) = -5.46$, $p < .001$, which also did not depend on Group or Drug, $ps ≥ .128$. In A-B, error bars represent ±1 standard error of the estimated marginal means from the corresponding fitted generalized linear mixed-effects model. (**C**) The proportion of lever presses that were followed by a food-cup approach was significantly higher for the devalued versus the valued lever, $t(131) = 4.11$, $p < .001$, which also did not depend on Group or Drug conditions, $ps ≥ .679$.