Neural correlates of learning pure tones versus natural sounds in the auditory cortex

Ido Maor[1,2], Ravid Shwartz-Ziv[2], Libi Feigin[2], Yishai Elyada[2], Haim Sompolinsky[2,3], and Adi

Mizrahi[1,2]*

[1]Department of Neurobiology, [2]The Edmond and Lily Safra Center for Brain Sciences, [3]Racah

Institute of Physics, The Hebrew University of Jerusalem, Jerusalem, Israel.

* Corresponding author:

Adi Mizrahi: Mizrahi.adi@mail.huji.ac.il

Acknowledgements: We thank Israel Nelken and Livia de Hoz for help in setting up the initial

behavioral experiments. We thank Yoni Cohen, David Pash and Joseph Jubran for writing the

software code of the Educage. We thank Yonatan Loewenstein, Merav Ahissar, and the Mizrahi

lab members for comments on the manuscript. This work was supported by an ERC consolidators

grant to A.M (#616063) and by Gatsby Charitable Foundation.

Author contributions: I.M. and A.M. designed the experiments. R.S.Z and H.S analyzed data.

L.F. conducted experiments. Y.E. constructed the vocalization stimuli and software for their

delivery. I.M. conducted experiments and analyzed the data. I.M. H.S. and A.M. wrote the paper.

Conflict of interest: The authors declare no competing financial interests.

ABSTRACT

Auditory perceptual learning of pure tones causes tonotopic map expansion in the primary

auditory cortex (A1), but the function this plasticity sub-serves is unclear. We developed an

automated training platform called the 'Educage', which was used to train mice on a go/no-go

auditory discrimination task to their perceptual limits, for difficult discriminations among pure

tones or natural sounds. Spiking responses of excitatory and inhibitory L2/3 neurons in mouse A1

revealed learning-induced overrepresentation of the learned frequencies, in accordance with

previous literature. Using a novel computational model to study auditory tuning curves we show

that overrepresentation does not necessarily improve discrimination performance of the network

to the learned tones. In contrast, perceptual learning of natural sounds induced 'sparsening' of the

neural response, and consequently improving discrimination of these complex sounds. The

signature of plasticity in A1 highlights its central role in coding natural sounds as compared to

pure tones.

INTRODUCTION

Learning is accompanied by plastic changes in brain circuits. This plasticity is often viewed as

substrate for improving computations that sub-serve learning and behavior. A well-studied

example of learning-induced plasticity is following perceptual learning where cortical

representations change towards the learned stimuli (Roelfsema and Holtmaat 2018, Gilbert,

Sigman, and Crist 2001). Whether such changes improve discrimination is still debated and the

mechanisms leading to these changes are still largely unknown. Furthermore, learning paradigms

in animal models are often limited to simplified, unnatural stimuli that deviate further from real

learning experience and that brain circuits actually compute.

46   Perceptual learning is an implicit form of lifelong learning during which perceptual

performance improves with practice (Gibson 1969). Perceptual learning spans a wide range of

sensory modalities and tasks like learning and improving reading skills, acquiring language, or

the learning to discriminate different shades of color and flavors. Extensive psychophysical

research on perceptual learning tasks led to a general agreement on some attributes of this type of

learning (Hawkey, Amitay, and Moore 2004). For example, perceptual learning has been shown

to be task specific, poorly generalized to other senses or tasks. It is also largely agreed upon that

gradual training is essential for improvement (Ahissar and Hochstein 2004, Ball and Sekuler

1987, Berardi and Fiorentini 1987, Ericsson 2006, Irvine et al. 2000, Karni and Sagi 1991, Kurt

and Ehret 2010, Lawrence 1952, Ramachandran and Braddick 1973, Wright and Fitzgerald

2001). Given the specificity observed at the behavioral level, functional correlates of perceptual

learning are thought to involve neural circuits as early as primary sensory regions (Schoups et al.

2001, Gilbert, Sigman, and Crist 2001). In auditory learning paradigms, changes are already

observed at the level of primary auditory cortex (Weinberger 2004). Learning to discriminate

among tones results in tonotopic map plasticity towards the trained stimulus (Bieszczad and

Weinberger 2010, Polley, Steinberg, and Merzenich 2006, Recanzone, Schreiner, and Merzenich

1993, Rutkowski and Weinberger 2005); reviewed in (Irvine 2017). Notably, however, not all

studies could replicate the learning induced changes in the tonotopic map (Brown, Irvine, and

Park 2004). Furthermore, artificially induced map plasticity was shown to be unnecessary for

better discrimination performances *per se*, (Talwar and Gerstein 2001, Reed et al. 2011). Our

understanding of the mechanisms underlying auditory cortex plasticity remains rudimentary, let

alone for more natural stimuli.

68   To gain understanding of learning-induced plasticity at single neuron resolution, animal

models have proven very useful. Mice, for example, offer the advantage of a rich genetic

experimental toolkit to study neurons and circuits with high efficiency and specificity (Luo,

Callaway, and Svoboda 2018). Historically, the weak aspect of using mice as a model was its

limited behavioral repertoire to learn difficult tasks. However, in the past decade technical

difficulties to train mice to their limits have been steadily improving with increasing number of

software and hardware tools to probe mouse behavior in high resolution (Egnor and Branson

2016, Krakauer et al. 2017, Aoki et al. 2017, Murphy et al. 2016, Erskine et al. 2018). Here, we

developed our own experimental system for training groups of mice on an auditory perceptual

task – an automatic system called the 'Educage'. The Educage is a simple affordable system that

allows efficient training of several mice simultaneously to discriminate among pure tones or

complex sounds.

A1 is well known for its tonotopic map plasticity following simple forms of learning in

other animal models (Irvine 2017). An additional interest in primary auditory cortex is its

increasing recognition as a brain region involved in coding complex sounds (Bizley and Cohen

2013, Kuchibhotla and Bathellier 2018). We thus asked what are the similarities and differences

in the plastic changes single neurons undergo following training to discriminate pure tones or

natural stimuli. We describe distinct changes in the long-term stimulus representations by L2/3

neurons of mice following perceptual learning and assess how these contribute to information

processing by local circuits. Using two photon targeted electrophysiology, we also describe how

L2/3 parvalbumin-positive neurons change with respect to their excitatory counterparts. Our work

provides a behavioral, physiological and computational foundation to questions of auditory-

driven plasticity in mice.

METHODS

<u>Animals</u>

A total of n=88, 7-8 week-old female mice were used in this work as follows. 44 mice were

C57BL/6 mice and 44 mice were PV–Cre; Ai9 double-heterozygous (PV × Ai9;(Hippenmeyer et

al. 2005, Madisen et al. 2010)). All experiments were approved by the Hebrew University's

IACUC.

<u>Behavioral setup</u>

The 'Educage' is a small chamber (10x10x10 cm), opening on one end into a standard animal

home cage where mice can enter and exit freely (Fig. 1a and Supplementary figure 1a). On the

other end, the chamber contains the hardware that drives the system, hardware for identifying

mice and measuring behavioral performance. Specifically, at the port entrance there is a coil radio

antenna (ANTC40 which connected to LID665 stationary decoder; Dorset) followed by infra-red

diodes used to identify mice individually and monitor their presence in the port. This port is the

only access to water for the mice. Water is delivered *via* a solenoid valve (VDW; SMC) allowing

precise control of the water volume provided on each visit. Water is delivered via a water spout,

which is also a lickometer (based on 1 microampere current). An additional solenoid valve is

positioned next to the water spout in order to deliver a mild air puff as a negative reinforcement,

if necessary. For sound stimulation, we positioned a speaker (ES1; TDT), at the center of the top

wall of the chamber. Sound was delivered to the speaker at 125 kHz sampling rate *via* a driver

and a programmable attenuator (ED1, PA5; TDT). For high speed data acquisition, reliable

control, on-board sound delivery, and future flexibility, the system was designed via a field

programmable gate array (FPGA) module and a real-time operating system (RTOS) module

(CompactRIO controller; National Instruments). A custom-made A/D converter was connected to

the CompactRIO controller, mediated signal from infra-red diodes and lickometer and controlled

the valves. A custom code was written in Labview to allow an executable user-friendly interface

with numerous options for user-input and flexibility for designing custom protocols. All software

and hardware design are freely available for download at

https://github.com/MizrahiTeam/Educage.

<u>Training paradigm</u>

Prior to the training, each mouse was implanted, under light and very short period of Isoflurane

anesthesia, with Radio Frequency Identification (RFID; Trovan) chip on the back of its head.

RFID chips allow identification of mice individually, which is then used by the system to control

the precise stimulus delivery and track behavioral performance, on a per-mouse basis. Food and

water were provided *ad libitum*. While access to water was only in the Educage, mice could

engage the water port without restriction. Thus, mice were never deprived of food nor water. At

the beginning of each experiment, RFID-tagged mice were placed in a large home-cage that was

connected to the Educage. Before training, we removed the standard water bottle from the home

cage. Mice were free to explore the Educage and drink the water at the behavioral port. Every

time a mouse entered the behavioral port it was identified individually by the antenna and the

infra-red beam and a trial was initiated. Before learning, any entry to the port immediately

resulted in a drop of water, but no sound was played. Following 24hrs of exploration and

drinking, mice were introduced for the first time to the 'target' stimulus – a series of six 10 kHz

pure tones (100 ms duration, 300 ms interval; 3 ms on and off linear ramps; 62 dB SPL) played

every time the mouse crossed the IR beam. To be rewarded with water, mice were now required

to lick the spout during the 900 ms response window. This stage of operant conditioning lasted

for 2-3 days until mice performed at >70% hit rates.

We then switched the system to the first level of discrimination when mice learned to

identify a non-target stimulus (a 7.1 kHz pure tone series) from the already known target stimulus

(separated by 49%/octave). Thus, on each trial one of two possible stimuli were played for 2.1

seconds- either a target tone or a non-target tone. Mice were free to lick the water spout anytime,

but for the purpose of evaluating mouse decisions we defined a response window and counted the

licking responses only within 900 ms after the sound terminated. Target tones were played at 70%

probability and non-target tones at 30% probability, in a pseudorandom sequence. A 'lick'

response to a target was rewarded with a water drop (15 microliter) and considered as a 'Hit' trial.

A 'no lick' response to a target sound was considered as a 'Miss' trial. A lick response to the non-

target was considered a 'False Alarm' (FA) trial, which was negatively reinforced by a mild air

puff (60 PSI; 600ms) followed by a 9 second 'timeout'. A 'no lick' response to the non-target was

considered a correct rejection (CR) and was not rewarded.

Once mice learned the easy discrimination, we switched the system to the second

discrimination stage. Here, we increased task difficulty by changing the non-target tone to 8.333

kHz, thus decreasing the inter tone distance to 26%/octave. Then, at the following stage, the inter

tone distance was further decreased to 14%/octave and then down to 6%/octave. This last

transition was often done in a gradual manner (»12%/octave»10%/octave»

8%/octave»6%/octave). In some of the animals (n=25), we trained mice to their JND and then

changed the task back to an easier level. For some of the mice (n=13) we played 'catch trials'

during the first and second sessions of the 14%/octave discrimination stages. In catch trials,

different tones spanning the frequency range of the whole training (7-10 kHz) were presented to

the animals in low probability (6% of the total number of sounds), and were neither negatively

nor positively reinforced.

For the vocalizations task, we used playback of pups' wriggling calls (WC) as the target

stimulus. These vocalizations were recorded with a one-quarter inch microphone (Brüel & Kjær)

from P4–P5 PV × Ai9 pups (n = 3), were sampled at 500 kHz and identified offline (Digidata

1322A; Molecular Devices). As the non-target stimulus, we used manipulations of the WC.

During the first stage of the operant learning, mice learned to discriminate between WC and a

fully reversed version of this call. Then, the second manipulation on the non-target stimulus was a

gradual change of the frequency modulation (FM) of each syllable in the call while leaving the

temporal structure of the call intact. To manipulate the syllable FM we used a dynamic linear FM

ramp. This operation multiplies each sampling interval within the syllable, by a dynamic speeding

factor, which changed according to the relative distance from the start and end of the syllable, and

generated a new waveform by interpolation from the original waveform. For example, for a 0.6

speeding factor, the beginning of each syllable was slower by a factor of 0.4 while the end of each

syllable accelerated by a factor of 0.4. The range of sound modulation used here was 0.66-0.9. A

value of 0.66 is away from the WC, 0.9 similar to the WC and 1 exactly the same as the WC. The

basic task design for the non-target sound was as follows: Reverse» 0.66 »0.81» 0.9.

<u>Surgical procedure</u>

Mice were anesthetized with an intraperitoneal injection of ketamine and medetomidine (0.80 and

0.65 mg/kg, respectively) and a subcutaneous injection of Carprofen (0.004 mg/g). Additionally,

dextrose–saline was injected to prevent dehydration. Experiments lasted up to 8 h. The depth of

anesthesia was assessed by monitoring the pinch withdrawal reflex, and ketamine/medetomidine

was added to maintain it. The animal's rectal temperature was monitored continuously and

maintained at 36 ± 1°C. For imaging and recording, a custom-made metal pin was glued to the

skull using dental cement and connected to a custom stage to allow precise positioning of the

head relative to the speaker (facing the right ear). The muscle overlying the left auditory cortex

was removed, and a craniotomy ($\sim 2 \times 2$ mm) was performed over A1 (coordinates, 2.3 mm

posterior and 4.2 mm lateral to bregma) as described previously(Cohen, Rothschild, and Mizrahi

2011, Stiebler et al. 1997).

Imaging and electrophysiology

Cell-attached recordings were obtained using targeted patch-clamp recording by a previously

described procedure (Cohen and Mizrahi 2015, Maor, Shalev, and Mizrahi 2016, Margrie et al.

2003). For visualization, the electrode was filled with a green fluorescent dye (Alexa Flour-488;

50 μM). Imaging of A1 was performed using an Ultima two-photon microscope from Prairie

Technologies equipped with a 16X water-immersion objective lens (0.8 numerical aperture;

CF175; Nikon). Two-photon excitation of the electrode and somata was used at 930 nm (DeepSee

femtosecond laser; Spectraphysics). The recording depths of cell somata were restricted to

subpial depths of 180–420 μm, documented by the multiphoton imaging. Spike waveform

analysis was performed on all recorded cells, verifying that tdTomato+ cells in L2/3 had

faster/narrower spikes relative to tdTomato-negative (tdTomato−) cells (See also(Cohen et al.

2013)).

Auditory stimuli

The auditory protocol comprised 18/24 pure tones (100 ms duration, 3 ms on and off linear

ramps) logarithmically spaced between 3-40 kHz and presented at four sound pressure levels (72–

42 dB SPLs). Each stimulus/intensity combination was presented 10/12 times at a rate of 1.4 Hz.

The vocalizations protocol comprised the playback of pups' wriggling calls (WC) and 3

additional frequency modulated (FM) calls, presented at 62 dB SPL for 16 repetitions.

Behavioral data analysis

To evaluate behavioral performance we calculated, for different time bins (normally 20 trials),

Hit and FA rates, which are the probability to lick in response to the target and non-target tones,

respectively. In order to compensate for the individual bias, we used a measure of discriminability

from signal detection theory – d-prime (d'). D' is defined as the difference between the normal

inverse cumulative distribution of the 'hit' and FA rates (Nevin 1969). D' for each discrimination

stage was calculated based on trials from the last 33% of the indicated stage. Psychometric curves

were extracted based on mouse performance in response to the catch trials. By fitting a sigmoidal

function to these curves we calculated decision boundaries as the inflection point of each curve.

Detection time was calculated for each mouse individually, by determining the time in which lick

patterns in the correct reject *vs.* the hit trials diverged (i.e. the time when significance levels

crossed $P<0.001$ in a two-sample t-test).

<u>Data analysis - electrophysiology</u>

Data analysis and statistics were performed using custom-written code in MATLAB

(MathWorks). Spikes were extracted from raw voltage traces by thresholding. Spike times were

then assigned to the local peaks of suprathreshold segments and rounded to the nearest

millisecond. For each cell, we obtained peri-stimulus time histogram (PSTH) and determined the

response window as the 100 ms following stimulus onset which evoked the maximal response

integral. Only neurons that had tone-evoked response (determined by a two sample t-test) were

included in our dataset. Based on this response window, we extracted the cell's tuning curve and

frequency-response area (FRA). Evoked firing rate was calculated as the average response to all

frequencies that evoked a significant response. Firing rate in the training band was calculated as

the response to frequencies inside the training band (7-10 kHz), averaged across all intensities.

Best frequency (BF) of each cell was determined as the tone frequency that elicited the strongest

response averaged across all intensities. The selectivity of the cell is the % of all frequency-

intensity combinations that evoked significant response (determined by a two sample t-test).

Pairwise signal correlations (r_{sc}) were calculated as Pearson correlation between FRA's matrices

of neighboring cells. The spontaneous firing rate of the cell was calculated based on the 100 ms

preceding each stimulus presentation. Response latency is the time point after stimulus onset at

which average spike count reached maximum.

<u>Statistical model based on the Independent Basis Functions (IBF) Method</u>

Since the measured responses before and after learning are not from the same cells, we cannot

estimate the changes of individual tuning curves due to learning. Instead, we must rely on

estimated learning-induced changes in the *ensemble* of single-neuron responses. Our goal,

therefore, was to build a statistical model of single-neuron tuning curves before and after

learning. This model is different before and after learning, and was used to estimate the learning-

induced changes in the population of A1 responses. Furthermore, we use this model as a

generative model that allowed us to generate a large number of 'model neurons' with statistically

similar response properties as the measured ones.

In principle, one could use a parametric model, by fitting each observed tuning curve to a

specific shape of functions (e.g., Gaussian tuning curves). However, since the tuning curves of

neurons to tone frequencies do not have symmetric 'gaussian' shapes, and some are bimodal,

fitting them to a parametric model has not been successful. Instead, we chose to model each

single neuron response as a weighted sum of a small set of orthogonal basis functions,

$$r_i(f) = \sum_{l=1}^{K} a_l^i g_l(f)$$

Here, $r_i(f)$ is the firing rate (i.e., the trial-averaged spike count) of the i-th neuron in response to

the stimulus with frequency f; K is the number of orthogonal basis functions denoted by $g_l(f)$

(dependencies in f are in log scale). In order to determine the basis functions and the coefficients,

a_l^i we performed Singular Value Decomposition (SVD) of the matrix of the measured neuronal

firing rates for the 18 values of f. Our model (1) uses a subset of the K modes with the largest

singular values (the determination of K is described below). The SVD yields the coefficients

$a_{l,data}^i$ for the *N observed* neurons and (2) smoothes the resultant SVD f-dependent vectors using

a simple 'moving average' technique to generate the basis functions, $g_l(f)$ (3) Importantly, to use

the SVD as a generative model, for each l we compute the histogram of the N $a_{l,data}^i$

coefficients. To generate 'new neurons', we sample each coefficient independently from the

corresponding histogram. In other words, we approximate the joint distribution of the coefficients

by a factorized distribution. This allowed us to explore the effect of changing the number of

neurons that downstream decoders use in order to perform the perceptual task.

Model (1) describes the variability of the population responses to the stimulus, in terms of tuning

curves of the trial averages firing rates. Additional variability in the data is the single trial spike

count. We model these as independent Poisson random variables with means given by $r_i(f)$.

Since neurons are not simultaneously recorded, we do not include noise correlations in the model.

We performed this procedure for the *naïve* and *expert* measured responses separately, so that both

the basis functions and the coefficient histograms are evaluated for the two conditions separately.

Note we do not make Gaussian assumptions about the coefficient histograms. In fact, the

observed histograms are in general far from Gaussian.

The choice of number of basis functions

Due to a limited number of trials that we sampled for each neuron, taking a large value of K can

result in over-fitting the model to the noise caused by the finite number of trials. To estimate the

optimal number of basis functions, we evaluated the percentage of response firing rate variability

of the population (i.e, the fraction of the sum of the squared SVD eigenvalues) as a function of K.

We also evaluated the parameters of model (1) based on a subset of trials and checked how well it

accounts for the observed tuning curves that are calculated from the test trials. We took K that

produces the smallest test error and saturated the fractional variance.

We used Model (1) with the above choice of K in order to evaluate the discrimination

ability of the population of A1 neurons, by creating *an ensemble* of single-neuron responses for

the naïve and expert conditions. To generate the model neurons, we sampled the coefficients of

the basis function independently from the corresponding histogram of the measured neurons, and

used these neurons for the calculations depicted below.

Fisher Information

We calculated the Fisher Information (FI) for each condition (naïve vs. expert) using our model

(1). FI measure bounds the mean squared error of an (unbiased) estimator of the stimulus from

the noisy single trial neuronal responses. When the neuronal population is large (and they are

noise- independent) FI also determines the discriminability d' of a maximum likelihood

discriminator between two nearby values of the stimulus (Seung and Sompolinsky 1993). Under

the above Poisson assumption, the FI for the *i*-th neuron is equal to $I_i = \frac{r_i'(f)^2}{r_i(f)}$, where $r_i'(f)$ is

the derivative of the firing rate with respect to the stimulus value *f*. The total FI is the sum of the

FIs of individual neurons (Seung and Sompolinsky 1993). This has been evaluated in both naïve

and expert conditions. Note that the FI are functions of the stimulus value *f*, around which the

discrimination task is performed.

Discrimination by linear readout

We applied a linear decoder to assess the ability to discriminate between nearby stimuli on the

basis of the neuronal population responses. We trained a support vector machine (SVM) with a

linear kernel, which finds an 'optimal' linear classifier that discriminate between two nearby

frequencies on the basis of single-trial vectors of spike counts generated with our generative

model (1) and Poisson variability. We then evaluated the probability of classification errors to test

trials, in both naïve and expert conditions. Since our training set is not linearly separable, we

used SVM with slack variables (Vapnik 1998), which incorporates a 'soft' cost for classification

errors. Each classification was iterated maximum 500 times (or until converged). In each

iteration, 16 trials were used for training the classifier and 4 trials were used to test the decoder

accuracy.

<u>Data analysis – vocalization responses</u>

Similarity of response to different vocalizations was calculated as Pearson correlation between

the PSTH's of the different stimuli. Classification of vocalization identity based on population

activity was determined using the Support Vector Machine (SVM) decoder with a linear kernel

and slack variables. The decoder was tested for its accuracy to differentiate between responses to

two different vocalizations. The input to the SVM consisted of the spike count of each neuron in

the syllable response window. The same number of neurons (37) was used in both groups to

avoid biases. We then evaluated the probability of classification errors to test trials, using leave-

one-out cross validation. Each classification was iterated 1000 times. In each iteration, 15 trials

were used for training the classifier and one trial was used to test decoder accuracy. The number

of syllables utilized in the decoder were increased cumulatively.

RESULTS

Behavior – Discrimination of pure tones

To study perceptual learning in mice, we developed a behavioral platform named the 'Educage'

(all software and hardware design are freely available for download at

https://github.com/MizrahiTeam/Educage). The Educage is an automated home-cage training

apparatus designed to be used simultaneously with several mice. One advantage of the Educage

over other procedures is that human interference is brought to minimum and training efficiency

increases. The 'Educage' is a small modular chamber (10x10X10 cm), opening on one end into a

standard animal home cage where mice can enter and exit freely (Fig. 1a). On its other end, the

chamber contains the hardware that drives the system, hardware for identifying mice and

measuring behavioral performance (Fig. 1a, Fig. S1a, and Methods). Mice were free to engage

the behavioral port at their own will, where they consume all of their water intake. Following

habituation, mice were trained on a go/no-go auditory discrimination task to lick in response to a

target tone (a series of 10 kHz pure tones) and withhold licking in response to the non-target tone.

A 'lick' response to a target was rewarded with a water drop (15 microliter) and considered as a

'Hit' trial. A 'no lick' response to a target sound was considered as a 'Miss' trial. A lick response

to the non-target was considered a 'False Alarm' (FA) trial, which was negatively reinforced by a

mild air puff (60 PSI; 600ms) followed by a 9 second 'timeout'. A 'no lick' response to the non-

target was considered a correct rejection (CR) and was not rewarded (Fig. 1a). On average, mice

performed 327 ± 71 trials per day, mainly during dark hours (Fig. S1b).

The initial level of learning was to identify a non-target stimulus (a series of 7.1 kHz pure

tones) from the 10 kHz target stimulus. These stimuli are separated by 49% of an octave and are

perceptually easily separated by mice. Despite the simplicity of the task, behavioral performance

varied widely between mice (Fig. 1b, Fig. S1c). On average, it took mice 54 ± 38 trials to cross our

criterion of learning, which was set arbitrarily at d'=1 (Fig. 1b; dotted line), and gradually

increased to plateau at d'=2.35 ± 0.64 (Fig. 1d). To extend the task to more challenging levels, we

gradually increased task difficulty by changing the non-target tone closer to the target tone. The

target tone remained constant at 10 kHz throughout the experiment and only the non-target

stimulus changed. The lowest distance used between target and non-target was 3%/octave (9.6

kHz vs 10 kHz). A representative example from one mouse's performance in the Educage

throughout a complete experiment is shown in figure 1c. The just noticeable difference (JND) for

each mouse was determined when mice could no longer discriminate (e.g. the JND of the mouse

shown in figure 1c was determined between 6-10%/octave). The range of JND's was 3-

14%/octave and averaged 8.6±4.7%/octave. Most mice improved their performance with training

(Fig. 1e, f), showing improved perceptual abilities along the task. The duration to reach JND

varied as well, ranging 3069±1099 trials. Detection times, defined as the time in which lick

patterns in the correct reject trials diverged from the lick patterns of the hit trials, increased

monotonically by ~177 ms per each step of task difficulty demonstrating the increased perceptual

load during the harder tasks (Fig. S1d).

To show that gradual training is necessary for perceptual learning we trained groups of

littermate mice on different protocols simultaneously. In one group of mice we used a standard

protocol and the animals were trained on the gradually increasing task difficulty described above.

Simultaneously, in the second group of mice – termed 'easy only' – animals were trained

continuously on the easy task. Although both groups of mice trained together, only the mice that

underwent gradual training were able to perform the hard task (Fig. 1g). Taking together, these

data demonstrate the efficiency of the Educage to train groups of mice to become experts in

discriminating between a narrowband of frequencies in a relatively short time and with minimal

human intervention.

Representation of pure tones in A1 following perceptual learning

To evaluate cortical plasticity following perceptual learning, we compared how pure tones are

represented in A1 of naïve and expert mice. We used *in vivo* loose patch recording of L2/3

neurons in anesthetized mice to record tone-evoked spiking activity in response to 3-40kHz pure

tones (Fig. 2a,b; Table 1). In naïve mice, responses were highly heterogeneous, with best

frequencies covering the whole frequency range (Fig. 2c,d; n=105 neurons, n=22 mice, red). In

expert mice, best frequencies of tuning curves were biased towards the frequencies that were

presented during learning (Fig. 2c,d; n=107 neurons, n=21 mice, blue). These data show, as

expected from previous literature, that learned frequencies in A1 become overrepresented at least

as measured by the neuron's best frequency (BF). We next showed that this overrepresentation

was specific to the learned tone by training mice on 4 kHz as the target tone and recorded neurons

in a similar manner to the abovementioned experiment. Indeed, L2/3 neurons in A1 of mice

training on 4kHz showed BF shifts towards 4 kHz (Fig. S2a). Temporal responses to the trained

frequencies were only slightly different between naive and expert mice. Specifically, average

spiking responses were slightly but significantly faster and stronger in experts (Fig. S2b, Table 1).

395    We recorded the responses to pure tones at different intensities and constructed frequency

396 response areas (Fig. S2c). Pairwise signal correlation of neighboring neurons, calculated from

397 these frequency response areas were high in naïve mice (0.2±0.28) but even higher in experts

398 (0.3±0.24; Fig. 2E). Thus, the basal level of functional heterogeneity in A1 (Maor, Shalev, and

399 Mizrahi 2016, Rothschild, Nelken, and Mizrahi 2010, Bandyopadhyay, Shamma, and Kanold

400 2010) is reduced following learning. This learning-induced increase in functional homogeneity of

401 the local circuit, emphasizes the kind of shift that local circuits undergo. Since neurons in expert

402 mice did not have wider response areas (Fig. 2F, Table 1), our data suggests that neurons shifted

403 their response properties towards the learned tones at the expense of frequencies outside the

404 training band. These results are largely consistent with previous studies in monkeys, cats, and rats

405 (reviewed in (Irvine 2017)), extending this phenomenon of learning-induced changes in tuning

406 curves to the mouse, to L2/3 neurons and to local circuits.

408 *A generative model of A1 population responses to pure tones*

409 To what extent does overrepresentation of a learned stimulus sub-serve better discrimination by

410 the neural population? To answer this question, we built a statistical model of tuning curves of

411 neurons in A1 using six basis functions (Fig. S3), that correspond to the six largest Singular

412 Value Decomposition (SVD) vectors of the population responses (Independent Basis Functions

413 (IBF) method; see Methods). In figure 3a we show two representative examples of tuning curves

414 and their reconstruction by our model. In contrast to Gaussian fits used previously (Vapnik

415 1998), our model captures the salient features of the shape of the auditory tuning curves

416 (asymmetry, multimodality), yet also smoothed the *raw* response vectors to reduce overfitting due

417 to finite sampling.

418    In order to choose the appropriate number of basis functions, we determined the minimal

419 number of basis functions that achieves good performance in reconstructing test single trial

420 responses. In figure 3b we show the fraction of explained variance as a function of the number of

421 basis functions, K. For both the naïve and expert groups the explained variance reaches above

422 96% after five basis functions. Figure 3c shows the Mean Square Error (MSE) on the unseen

423 trials as a function of K in data from both the naïve and expert animals. The MSE exhibits a broad

424 minimum for K in a range between 6-13. Interestingly, both naïve and experts achieve roughly

425 the same MSE values, although for experts the MSE values at both low and large values of K are

426 considerably larger than that of the naïve, presumably due to the smaller number of cells in the

expert conditions. Taken together, we conclude that for these conditions six basis functions are

the appropriate number and used this value for our calculations.

Based on the IBF method described above, we generated a population of tuning curves

(500 'new neurons'), and estimated their total Fisher Information (FI; see Methods). Figure 3d

shows the FI as a function of the stimulus *f* for both the naïve and expert conditions. Surprisingly,

the FI of the neurons from expert animals was enhanced relative to the naïve group, but only for

stimuli at both flanks of the training band. Importantly, the FI within the band of the trained

frequencies remained unchanged (Fig. 3d, within the black lines). The same result holds true for a

performance of a support vector machine (SVM) classifier. Using SVM to separate any two

frequencies that are 0.2198 octave apart, discriminability (d') values derived from the classifier's

error show similar results as the FI (Fig. 3e; (Seung and Sompolinsky 1993)). The value of d' is

larger in the expert groups as compared to the naïve but only outside the training band, whereas

within the band discriminability is not improved (or even slightly compromised).

The results shown in figure 3 do not change qualitatively if we use our SVD model for

the recorded neurons, as opposed to newly modelled neurons. One advantage of having a

generative model for the population responses is that we can generate an unlimited number of

trials and tuning curves. We took advantage of this to explore whether the results of the FI and

SVM change with population size. To answer this question, we evaluated the mean

discrimination performance (over test neurons) as a function of the number of sampled cells, *N*,

which increases as expected. Consistent with the results of the SVM, the performances in the

naïve and expert groups are similar with slight tendency for a higher accuracy in the naive

population at large *Ns* (Fig. 3f*)*. In contrast, for frequencies near the training band, the accuracy is

substantially larger in the expert than in the naïve group for virtually all *N* (Fig. 3g). Thus, it

seems that learning induced changes in tuning curves that do not improve discriminability of the

learned stimuli.

Perceptual learning of natural sounds

Natural sounds are characterized by rich spectro-temporal structures with frequency and

amplitude modulations over time (Mizrahi, Shalev, and Nelken 2014). Discrimination of such

complex stimuli could be different from that of pure tones. Thus, we next designed a task similar

to that with the pure tones but using mouse vocalizations as the training stimuli. We used

playback of pups' wriggling calls (WC) as the target stimulus (Fig. 4a - top). As the non-target

stimuli, we used frequency modulations of the WC; a manipulation that allowed us to morph one

460 stimulus to another by a continuous metric (Fig. 4a). The range of sound modulation used here

461 was indexed as a "speeding factor" (see Methods for details). In short, a modulation factor of 0.66

462 affected the original WC more than a modulation factor of 0.9 did, and is therefore easier to

463 discriminate (Fig. 4b). To reach perceptual limits we trained mice gradually, starting with an easy

464 version of the task (WC vs a temporally reversed version of the WC) and then gradually to

465 modulated calls starting at 0.66 modulation. Once mice reached >80% hit rates we changed the

466 non-target stimulus to more difficult stimuli until mice could no longer discriminate (Fig. 4c).

467 Mice (n=11) learned the easy task, i.e., discriminating WC from a 0.66 modulated call, with

468 average d' values of 2.5±0.4 (Fig.4d). On average, mice could only barely discriminate between a

469 WC and its 0.9 modulation (d' at 0.9 was 1±0.8; Fig. 4d). While these discrimination values were

470 comparable to the performance of pure tones, detection times were substantially higher (Fig.

471 S4a). For similar d' values, discriminating between the vocalizations took 300-1000ms longer as

472 compared to the pure tone tasks (Fig.4e). In addition, learning curves were slower for the

473 vocalization task as compared to the pure tones task. The average number of trials to reach d'=1

474 for vocalizations was 195 trials, more than three times longer as compared with pure tones

475 (compare Fig. S4b and 1b, respectively). These differences may arise from the difference in the

476 delay, inter-syllable interval and temporal modulation of each stimulus type, which we did not

477 further explore. Taken together, these behavioral results demonstrate a gradual increase in

478 perceptual difficulty using a manipulation of a natural sound.

480 *Sparser response in A1 following perceptual learning of natural sounds*

481 To study the neural correlates in A1 that support natural sound discrimination we recorded L2/3

482 neurons in response to the learned stimuli (Fig. 5a), expecting increased representation of these

483 particular stimuli. Surprisingly, and in contrast to the results from pure tone discrimination, we

484 did not find an increase in the representation of the learned stimuli. The fraction of cells

485 responding to the trained vocalization remained constant (Fig. S5a) as well as the evoked firing

486 rate for the preferred vocalization or the preferred syllable within a vocalization (Table 2).

487 Instead, representation in expert mice trained on vocalizations was sparser. Specifically,

488 responses were more selective in their response within the call, i.e. each cell responded to less

489 syllables and each syllable was represented by fewer cells, Table 2; Fig. S5b). Sparseness was

490 also evident from the average population response to the vocalization (Fig. 5b). Nearly all

491 syllables had weaker responses, three of which were statistically weaker (Fig. 5b, gray bars).

492 Notably, the smaller fraction of responses was not just an apparent sparseness due to increase in

493   trial to trial variability, as reliability of responses by neurons in the expert group remained similar

494   to reliability of responses in the naïve group (Table 2).

495   We next asked whether the 'sparsening' described above bears more information to the

learned stimulus. We analyzed population responses by including all neurons from all mice as if

they are a single population (naïve: n=41 neurons from 8 mice; expert: n=37 neurons from 6

mice). We calculated the Pearson correlation of the population response to all responses in a

pairwise manner (Fig. 6a). As compared to naïve mice, the absolute levels of correlations in

expert mice were significantly lower for nearly all pairs of comparisons (Fig. 6a, asterisks). As

expected, weaker modulations of the call and, hence, high similarity among stimuli, were

expressed as higher correlations in the neuronal responses (Fig. 6b). The pairs of stimuli which

mice successfully discriminated in the behavior (0.66 *vs.* the original WC and 0.81 *vs.* the original

WC), had significantly lower correlation in the expert mice (Fig. 6b, rank sum test, p<0.05).

Responses to the more similar stimuli that were near perceptual thresholds (i.e. 0.9 *vs.* the original

WC) were lower in expert mice, but not significantly (Fig. 6b, rank sum test, p>0.05). This

reduced Pearson correlation suggested that plasticity in A1 supports better discrimination among

the learned natural stimuli. Indeed, a SVM decoder performed consistently better in expert mice,

discriminating more accurately the original WC from the manipulated ones (Fig. 6c). As

expected, the decoder performance monotonically increased when utilizing the responses to more

syllables in the call. However, in the expert mice, performances reached a plateau already half

way through the call, suggesting that neuronal responses to the late part of the call carried no

additional information useful for discrimination. Similarly, the correlation of the population

responses along the call, shows that responses were separated already following the first syllable,

but that the lowest level of the correlation was in the 5th - 7th syllables range, which then rapidly

recovered by the end of the call (Fig. S5c). These findings are also consistent with the behavioral

performance of the mice as decisions are made within the first 1.5 seconds of the trial (or the first

7 syllables of the sentence). Specifically, the head of the mouse is often retracted by the time the

late syllables are played (Fig. 4e). Taken together, 'sparser' responses improve neural

discrimination of learned natural sounds.

Learning induced plasticity of Parvalbumin neurons

The mechanisms responsible for the learning-induced changes following learning are currently

unknown. We used mouse genetics and two-photon targeted patch to ask whether local inhibitory

neurons could contribute to the observed plasticity we describe above. To this end, we focused

only on parvalbumin inhibitory (PV$^+$) interneurons as recent evidence indicates their role in a

variety of learning related plasticity processes (Letzkus et al. 2011, Wolff et al. 2014, Kaplan et

al. 2016, Lagler et al. 2016, Lee et al. 2017, Goel et al. 2017). Here we probe the role of

inhibition in perceptual learning by measuring learning induce plasticity in the response

properties of the PV$^+$ neurons. We trained PV$^+$-Cre x Ai9 mice in the Educage and then patched

single neurons under visual guidance (Fig. 7a; (Cohen and Mizrahi 2015, Maor, Shalev, and

Mizrahi 2016). In order to increase the sample of PV$^+$ neurons we used targeted patch and often

patched both PV$^+$ and PV$^-$ neurons in the same mice. PV$^-$ neurons were used as proxy for

excitatory neurons (these neurons were also included in the analysis shown in Fig. 2-6). All the

TdTomato$^+$ neurons that we patched were also verified as having a fast spike shape (Fig. 7a), a

well-established electrophysiological signature of PV$^+$ cells. PV$^+$ neurons had response properties

different from PV$^-$ neurons in accordance with our previous work (Maor, Shalev, and Mizrahi

2016). For example, PV$^+$ neurons responses were stronger and faster to both pure tones and

natural sounds (Fig. 7b-c and Table 3; compare with Fig. 2b and 5a, see also (Maor, Shalev, and

Mizrahi 2016)).

Following pure tone learning, PV$^+$ neurons also changed their response profile. On

average, the BF of PV$^+$ neurons shifted towards the learned frequencies, similar to what we

described for PV$^-$ neurons (Fig. 8a,b). The shift in tuning curves of PV$^+$ neurons was also

accompanied by a significant widening of their receptive fields (Fig. S6a). When we compared

the BF's of PV$^-$ and PV$^+$ neurons within the same brain (within 250 microns of each other), we

found that excitatory and inhibitory neurons became more functionally homogeneous as

compared to naïve mice (Fig. 8c). As both neuronal groups show similar trends in the shift of

their preferred frequencies we rule out a simple scenario whereas parvalbumin neurons increase

their responses in the sidebands of the learning frequency. In other words, plasticity does not

seem to be induced by lateral inhibition *via* parvalbumin neurons, but rather maintain a strict

balance between excitation and inhibition, regardless of whether they are naïves or experts (Wehr

and Zador 2003, Zhou et al. 2014). Note that the peak of the PV$^+$ population response and their

BF distribution was on the outskirts of the training band, rather than within it (compare Fig. 8a,b

with Fig. 2c,d), and concomitantly, the slope of the population responses at the trained frequency

band increased due to learning (Fig. 8a). To assess the computational effect of the plasticity in

the inhibitory neurons' responses we have applied on their responses the same d' and Fisher

Information calculated for the PV$^-$ neurons (Fig. 8d,e). Overall, the discrimination performance of

the two cell populations (when equalized in size) is similar. However, the PV$^+$ population shows a

significant learning related increase in tone discrimination performance by these cells within the

training band (Fig. 8d,e) in contrast to the results for PV$^-$ neurons (Fig. 3d and e). This result is

consistent with the above-mentioned increase in their response slopes in the trained frequency.

Following natural sound learning, we found no significant changes in basic response

properties to any parameter in the vocalizations by the PV$^+$ neurons (Fig. S6d, Table 3). The

relationship between the responses of PV$^+$ and their PV$^-$ neighbors remained constant as reflected

in the similar slopes of the functions describing PV$^-$ firing versus PV$^+$ firing (Fig. S6c). Note that

this common trend of plasticity (i.e. lower responses after learning) is also reflected in the

population PSTHs. The PSTHs in expert mice of both PV$^-$ and PV$^+$ neurons tended to have lower

firing rates than in naïve mice (Fig. S6b). This result suggests that the excitation-inhibition

balance, as reflected in the responses of PV$^-$ versus PV$^+$, remains. In PV$^+$ neurons, the temporal

correlation along the call as well as the decoding performance from these neurons showed

changes that are qualitatively similar to their PV$^-$ counterparts, but the data across the population

was noisier (Fig. 8f-g; Fig. S6d) perhaps due to the smaller sample of the PV$^+$ dataset.

DISCUSSION

Plasticity in frequency tuning following perceptual learning

Shifts in the average stimulus representation towards the learned stimuli is not a new

phenomenon. Similar findings were observed in numerous studies, multiple brain areas, animal

models and sensory systems, including in auditory cortex (Buonomano and Merzenich 1998,

Karni and Sagi 1991, Weinberger 2004). In fact, the model of learning-induced plasticity in A1,

also known as tonotopic map expansion, is an exemplar in neuroscience (Bakin and Weinberger

1990, Polley, Steinberg, and Merzenich 2006, Recanzone, Schreiner, and Merzenich 1993,

Rutkowski and Weinberger 2005); but see (Brown, Irvine, and Park 2004, Crist, Wu, and Gilbert

2001, Ghose, Yang, and Maunsell 2002, Kato, Gillet, and Isaacson 2015). Indeed, we also found

that the number of neurons tuned to the learned frequency band increased in expert mice after

perceptual learning (Fig. 2). Thus, our results support the observations of tuning curve plasticity

in a primary sensory cortex, showing this here specifically for L2/3 neurons and extending it to

A1 in mice.

Since we sampled only a small number of neurons in areas smaller than 250μm³, our

observations cannot be inferred as direct evidence for tonotopic map expansion. Rather, our data

emphasizes that plastic-shifts occur in local circuits (Fig. 2e). Given that neurons in A1 are

functionally heterogeneous within local circuits (Bandyopadhyay, Shamma, and Kanold 2010,

Maor, Shalev, and Mizrahi 2016, Rothschild, Nelken, and Mizrahi 2010), any area in A1 that

represents a range of frequencies prior to learning could become more frequency-tuned once

learned. Such a mechanism allows a wide range of modifications within local circuits to enable

increased representation of the learned stimuli without necessarily perturbing gross tonotopic

order. One advantage of local circuit heterogeneity is that it allows circuits to maintain a dynamic

balance between plasticity and stability (Mermillod, Bugaiska, and Bonin 2013). Only mapping

the full extent of auditory cortex, at both global and local scale and preferably over time in the

same neurons (Rothschild and Mizrahi 2015), will enable to reveal the precise type of changes the

cortex undergoes and how it exploits its variability to sub-serve perceptual learning.

Quantifying overrepresentation across a population of neurons and interpreting these in

the context of the learned perceptual tasks have been the subject of extensive research (Bao

2015). It has commonly been assumed that the learning-induced changes in tuning properties

improves the accuracy of coding of the trained stimuli. In particular, for discrimination tasks

between two stimuli differing by a small change in a one-dimensional stimulus parameter,

perceptual learning theory predicts that sharpening of the slope of the tuning curves with respect

to this parameter improves the discriminability of the stimuli (Seung and Sompolinsky 1993).

However, the observed increased representation of the BFs towards the training band in expert

mice may not increase over all tuning slopes and may even decrease them, especially since the

slopes tend to be small at the best frequencies. A closer look at the tuning curves in A1 shows that

they are often irregular with multiple slopes and peaks (i.e. not having simple unimodal Gaussian

shapes). Furthermore, the learning induced changes in the ensemble of tuning curves are not

limited to shifting the BF, hence a more quantitative approach was required to assess the

consequences of the observed learning induced plasticity on discrimination accuracy. This

motivated us to develop a method that takes into account the irregularities in tuning curves of A1

neurons. To this end, we have used a new, SVD based, generative model (Fig. 3), allowing us to

assess the combined effects of changes in BFs as well other changes in the shapes of the tuning

curves. Surprisingly, both Fisher Information analysis as well as estimated classification errors of

an optimal linear classifier, show that learning induced changes in tuning curves does not

improve tone discriminability at trained values. This conclusion is consistent with previous work

on the effect of exposure to tones during development. Early life experience during development

which induces similar changes to those observed during learning has been argued to decrease tone

discriminability for similar reasons (Han et al. 2007). However, in that work, the functional effect

of tuning curve changes was consistent with an observed impaired behavioral performance,

suggesting that plasticity in A1 sub-serves discrimination behavior. In contrast, the stable (or

even reduced) accuracy in the coding of the trained frequency we observed occurs despite the

improved behavioral performance after training. What may this finding underlie?

The changes that we recorded in A1 following pure tone learning do not improve their

contribution to discrimination of the task stimuli, and are thus orthogonal to direct behavioral

performance. One interpretation of this finding is that learning induced plasticity in other brain

regions, either in downstream (or parallel) auditory areas or in task related 'readout' areas enable

improved behavioral performance. Notably, even the representation in naïve mice would be

sufficient for reliable decoding at behavioral resolution – utilizing a population of only two-three

hundred A1 neurons (Fig. 3g). Thus, changes in the tuning of A1 neurons are not necessary for

discriminating the pure-tone task. Further, we suggest that the observed learning-induced changes

in the tuning of pure tones in A1 are the result of unsupervised Hebbian learning induced by over

exposure to the trained tones during the training period; similar to the reported results in early

overexposure (Han et al. 2007). Unsupervised learning signals are not driven by task related

reward *per se* and may increase representation rather than discriminability. Increased

representation of trained stimuli may lead to improved discriminability to untrained tones, as

observed experimentally. Importantly, we find that for natural sounds, learning-induced changes

in A1 responses improved the discriminability of the trained stimuli relative to the naïve

responses. This suggests that A1's primary function is in processing and coding complex auditory

stimuli like e.g. natural sounds, rather than pure tones.

Coding of Natural sounds in A1

Unlike pure tones, natural sounds are composed of rich spectro-temporal energies and include a

set of sound features such as amplitude modulations, frequency modulations, harmonics, and

noise. Although auditory cortex is organized tonotopically, it may not be essential for processing

simple sounds, as these stimuli are accurately represented in earlier stages in the auditory

hierarchy, as early as the brainstem (Mizrahi, Shalev, and Nelken 2014, Nelken 2004). This is

consistent with experiments showing that auditory cortex is necessary for associative fear

conditioning with complex sounds (Letzkus et al. 2011) but not with pure tones (Weible et al.

2014, LeDoux 2000, Ohl et al. 1999).

A multitude of studies have shown that single neurons and populations in A1 respond to

sounds in a non-linear fashion (Angeloni and Geffen 2018, Harper et al. 2016, Bathellier,

Ushakova, and Rumpel 2012). For example, neurons in A1 can be selective to harmonic content

that are prevalent in vocalizations and other natural stimuli (Feng and Wang 2017). Some

neurons in A1 show strong correlations to global stimulus statistics (Theunissen and Elie 2014).

Furthermore, neurons in A1 are sensitive to the fine-grained spectrotemporal environments of the

sounds, expressed as strong gain modulation to local sound statistics (Williamson et al. 2016), as

well as to sound contrast and noise (Rabinowitz et al. 2013, Rabinowitz et al. 2011). All of these

features (harmonics, globally and locally rich statistics, and noise) are well represented in the

wriggling calls we played during learning. Which of these particular sensitivities changes after

perceptual learning is not yet known but one expression of this plasticity can be the increased

sparse sound representation we found here (Fig. 5, S5). Sparseness can take different forms

(Barth and Poulet 2012). Here, sparseness is expressed mainly as a reduced number of neurons in

the network that respond to any of the 12 syllables played (Fig. S5). Such increase in selectivity

to the syllables could arise from disparate mechanisms, and changes in the structure of local

inhibition was one suspect that we tested (Froemke 2015).

Inhibitory plasticity follows excitatory plasticity

Cortical inhibitory neurons are central players in many forms of learning (Hennequin, Agnes, and

Vogels 2017, Kullmann et al. 2012, Sprekeler 2017). Inhibitory interneurons have been

implicated as important for experience dependent plasticity in the developing auditory system

(Hensch 2005), during fear learning in adulthood (Courtin et al. 2014, Letzkus et al. 2011), and

following injury (Resnik and Polley 2017). Surprisingly, however, and despite the numerous

studies on parvalbumin neurons, we could not find any references in the literature of recordings

from parvalbumin neurons after perceptual learning. Two simple (non mutually exclusive)

hypotheses are naively expected. One is that plasticity in inhibitory neurons are negative mirrors

of the plasticity in excitatory neurons. This would predict that inhibitory neurons would increase

their responses to the stimuli for which responses of excitatory neurons are downregulated (as

found in the plasticity of somatostatin expressing (SOM) neurons following passive sound

exposure (Kato et al. 2015), or in multisensory plasticity in mothers (Cohen and Mizrahi, 2015).

The second is that inhibitory neurons enhance their response to 'lateral' stimuli thus enhancing

selectivity to the trained stimulus, as suggested by the pattern of maternal related plasticity to pup

calls (Galindo-Leon, Lin, and Liu 2009). To the best of our knowledge, our study thus provides a

first test of these hypotheses in the context of perceptual learning. We found no evidence for

these scenarios. Instead, a common motif in the local circuit was that parvalbumin neurons

changed in a similar manner to their excitatory counterparts.

The cortex hosts several types of inhibitory cells (Hattori et al. 2017, Zeng and Sanes

2017), presumably serving distinct roles. While PV neurons are considered a rather homogeneous

pool of neurons based on molecular signature, their role in coding sounds is not. This is evident in

recordings from pyramidal neurons in A1 while optogenetically inhibiting the PV$^+$ neurons

resulted in mixed effects (Seybold et al. 2015, Phillips and Hasenstaub 2016). In the visual

cortex, PV$^+$ cells' activity (and presumably its plasticity) was correlated with stimulus-specific

response potentiation but not in ocular dominance plasticity (Kaplan et al. 2016), again,

suggesting that PV$^+$ neurons are not necessarily involved in all forms of experience-dependent

plasticity. Inhibitory neurons have been suggested to play a key role in enhancing the detection of

behaviorally significant vocalization by lateral inhibition (Galindo-Leon, Lin, and Liu 2009). But

recent imaging data argue that somatostatin interneurons rather than PV$^+$ interneurons govern

lateral inhibition in A1 (Kato, Asinof, and Isaacson 2017). Our results are also consistent with the

observation that in contrast to the SOM neurons, the changes in responses following sound

exposure are similar in PV$^+$ and pyramidal neurons (Kato et al 2015). To what extent

somatostatin or other interneurons subtypes contribute to excitatory plasticity after perceptual

learning remains to be studied.

Simplifying training with the Educage

Training rodents on perceptual learning paradigms has traditionally been performed manually

over many weeks, sometimes months, of training (Polley, Steinberg, and Merzenich 2006, Kurt

and Ehret 2010). Behavior is confounded by all sorts of variability, even in the controlled

conditions of a laboratory and within inbred mice of the same age and sex (Renart and Machens

2014). The Educage is a new tool for standardizing cognitive tasks, as we demonstrated here for

perceptual learning. The Educage allows training of larger animal datasets, effortlessly collecting

thousands of trials per day, and effectively increasing statistical power for assessing inter-animal

variability. Since mice train simultaneously, it potentially reduces stress-associated with factors

like social deprivation. Other behavioral tools have been introduced in recent years. Each system

has its own strengths or weaknesses in cost, design and operation load (e.g. see a new system for

olfactory perceptual learning (Erskine et al. 2018), and new systems with options for combining

physiology and imaging (Aoki et al. 2017)). The simple design and operation of the 'Educage'

together with its affordable hardware and software makes it yet another valuable tool for

simplifying training.

a

Figure 1- Perceptual learning in the 'Educage'

A. Left: Schematic design of the 'Educage' system and its components. **Right:** Schematic

representation of the go/no-go auditory discrimination task. CR- correct rejection, FA- False

Alarm. **B.** d' values of three representative mice and the population average± s.e.m for the first

stage of discrimination. Learning criterion is represented as dashed line (d'=1). **C.** Lick responses

to the target tone (solid line) and non-target tone (dashed line) of one representative mouse along

different discrimination stages. **D.** population average d' values for the different discrimination

stages. N=39 mice (mean ± s.e.m). Shades denote the level of difficulty (from 49% to 4-10%

octave apart). **E.** Individual d' values at the end of each level as a function of d' in the beginning

of that level. Shades denote the level of difficulty. Learning criterion represented as dashed lines

(d'=1). **F.** Normalized psychometric curves of five mice calculated from the first (light curves)

and the second (dark curves) 14%/octave session. Light and dark arrows indicate average

decision boundaries in the first and second sessions respectively (Mann-Whitney U-test on

criteria: p=0.03). **G.** d' values in easy (49%/octave) and more difficult (10% /octave)

discrimination stages of individual mice from the 'Easy only' group (filled circles) and from the

perceptual learning group (blank circles). d' values are significantly different between groups

only for the hard discrimination level (Mann-Whitney U-test: ***p<0.001).

Figure 2- Learning induces over representation of the learned stimuli

A. Schematic representation of the experimental setup and a sample of the loose patch recording
showing a representative cell from naïve (left) and expert (right) mice (gray markers indicate tone
stimuli). **B.** Raster plots and peri-stimulus time histograms (PSTH) in response to pure tones of
the cells shown. Gray bars indicate the time of stimulus presentation (100 ms). Color bars and
arrows indicate the training frequency band (7.1-10 kHz) and BF, respectively. **C.** Population
average of normalized response tuning curves of 105 neurons from naïve mice (red) and 107

neurons from expert mice (blue; mean ± s.e.m). Gray area indicates the training frequency band.

Asterisks correspond to frequencies with significant response difference (Mann-Whitney U-test:

*p<0.05). **D.** Best frequency (BF) of individual neurons from naïve mice (red markers) and expert

mice (blue markers; Mann-Whitney U-test: **p<0.01). **E.** Pairwise signal correlations (r_{sc}) values

between all neighboring neuronal pairs in naïve (red) and expert (blue) mice. Neurons in expert

mice have higher r_{sc} (Mann-Whitney U-test: p<0.001). **F.** Cumulative distribution of response

selectivity in naïve (red) and expert (blue) mice. Response selectivity was determined as the % of

all frequency-intensity combinations that evoked a significant response. Distributions are not

significantly different (Kolmogorov Smirnov test; p=0.542).

Figure 3- Plasticity in A1 does not improve the discrimination of the learned tones

A. Two representative examples of tuning curves of neurons recorded in A1. Average spike rates

are shown in blue and the SVD approximation of the particular curves are shown in orange. Note

that although the SVD approximation is smooth, it captures the irregular dynamics (i.e. non-

Gaussian) of the tuning curves. **B,C.** The explained variance and error as a measure of the

number of SVD vectors used in the model. **D.** Fisher Information calculated form the tuning

curves of the both populations along the frequency dimension. Note the increased FI for the

expert neurons in the flanks of the training band but not within it (gray band). **E.** Discriminability

(d') of SVM decoder along the range of frequencies. Pairwise comparison along the continuum

are performed for frequencies 0.2198 octave apart. In accordance with 'd' the decoder does not

perform better in the training band (gray shade). **F,G.** Classification performance of the decoder

as a function of the number of neurons in the model. In the training band (F), the performance is

similar for both naïve and expert mice. Outside the training band (0.4396 octave apart; G),

performance improved rapidly in the expert mice.

Figure 4- Perceptual learning of vocalizations

A. Spectrograms of the wriggling call (WC, 'target' stimulus, top panel) and the manipulated

WC's ('non-target' stimuli, bottom panels). **B.** Enlargement of the spectrogram's 4th syllable of

the target WC and the manipulated calls. Top, a large manipulation (speeding factor-0.66), which

is perceptually easy to discriminate from the WC. Bottom, a minor manipulation (speeding

factor-0.9), which is perceptually closer to the WC. **C.** Lick responses to the target tone (solid

line) and non-target tone (dashed line), binned over 50 trials, of one representative mouse along

the different discrimination stages. The task of the first stage was to discriminate between WC vs

reversed playback of the call ('Reverse'). The following stages are different degrees of call

modulation. Titles correspond to the speeding factors used for the non-target stimulus. **D.**

Population average d' values for the different discrimination levels. N=9 mice (mean ± s.e.m).

Shades denote the level of difficulty. **E.** Comparison between detection times during the easy and

difficult stages of pure tone (blank circles) and vocalizations (filled circles) discrimination tasks.

Detection times are significantly different between all groups (Mann-Whitney U-test:

***p<0.001).

a

b

Figure 5- Learning complex sounds induces 'sparsening'

A. Representative examples of raster plots in response to 4 modulated wriggling calls from naïve
(top) and expert (bottom) mice. Red lines indicate spikes in response windows which are
significantly above baseline. The stimulus power spectrum shown in light pink in the background.
B. Average normalized PSTHs calculated from all neurons in response to the original WC. Data
is shown overlaid for naïve (red) and expert (blue) mice. In expert mice, only syllables 1, 7 and
11 evoked significantly weaker responses as compared to naïve mice (dark gray bars, Mann-
Whitney U-test; p=0.03, 0.001, 0.03).

Figure 6- Decorrelation improves coding

A. Matrices describing the average response similarity of individual neurons between all combinations of stimuli in naïve (left) and expert (right) mice. Each pixel indicates the average Pearson correlation value calculated from all syllables' evoked spike rate from all neurons to two different calls. Neurons from expert mice have lower correlation between responses to different modulated calls (asterisks indicate significant differences between naïve and expert groups; Mann-Whitney U-test, P<0.05). **B.** Pearson correlation between responses to modulated WCs and responses to the original WC in naïve (red) and expert (blue) mice (mean± s.e.m). Correlations are significantly different in the 0.66 and 0.81 modulation (Mann-Whitney U-test: * p<0.05). **C.** Classification performance of a Support Vector Machine (SVM) decoder. The decoder was tested for its accuracy to differentiate between the modulated WC stimuli against the original WC. The performance of the decoder is shown for neurons from the naïve (red) and expert (blue) groups. Decoder performance is plotted separately for the three different pairs of stimuli. Each point in each graph shows the number of syllables the decoder was trained on and allowed to use. Error bars are SEM for 1000 repetitions of leave-one-out cross validation.

Figure 7- Response properties of PV⁺ neurons

A. Left: Schematic representation of the experimental setup for two-photon targeted patch and average spike waveform of 203 PV⁺ neurons. **Right**: Representative two-photon micrograph (projection image of 120 microns) of tdTomato+ cells (red) and the recording electrode (Alexa Fluor-488, green). **B.** Raster plots and peri-stimulus time histograms (PSTH) in response to pure tones of a representative PV⁺ neuron from naïve (left) and expert (right) mice. Gray bars indicate

828   the time of stimulus presentation (100 ms). Color bars and arrows indicate the training frequency

829   band (7.1-10 kHz) and BF, respectively. **C.** Representative examples of raster plots from PV$^+$

830   neurons in response to 4 modulated wriggling calls from naïve (left) and expert (right) mice. Red

831   lines indicate spikes in response windows which are significantly above baseline. Stimulus power

832   spectrum shown in light pink in the background.

Figure 8- Plasticity of PV⁺ neurons after learning

A. Population average of normalized response tuning curves of n=120 PV⁺ neurons from naïve

mice (red) and 83 PV⁺ neurons from expert mice (blue; mean ± s.e.m). Gray area indicates the

training frequency band. Asterisks correspond to frequencies with significant response difference

(Mann-Whitney U-test: *p<0.05). **B.** Best frequency (BF) of individual PV⁺ neurons from naïve

mice (red) and expert mice (blue; Mann-Whitney U-test: **p<0.01) **C.** Average distance in BF

between PV$^-$ and PV$^+$ neurons from the same penetration sites. Distances were significantly

smaller in expert mice (Mann-Whitney U-test: ***p<0.001). **D.** Fisher Information calculated

form the tuning curves of PV$^+$ neurons **E.** Discriminability (d') of SVM decoder along the range

of frequencies. **F**. Matrices describing the average response similarity of individual PV$^+$ neurons

between all combinations of different stimuli in naïve (left) and expert (middle) mice. Each pixel

indicates the average Pearson correlation value calculated from all syllables evoked spike rate

from all neurons to two different calls. There was no significant difference between correlations

of responses to different modulated calls in naïve and expert mice (Mann-Whitney U-test,

P>0.05). **G.** Pearson correlation between responses to modulated WCs versus the responses to

the original WCs in naïve (red) and expert (blue) mice (mean± s.e.m).). Correlations are not

significantly different for all comparisons (Mann-Whitney U-test: p=0.4, 0.5, 0.5).

TABLES

Group	Animals (n)	Cells (n)	BF (kHz)	Spontaneous spike rate (Hz)	Evoked firing rate (Hz)	FR in T.B (Hz)	Response latency (ms)	Selectivity (% of all stimuli)
Naïve	22	105	11.3 ±8.4	0.45±0.58	15.5 ±5.3	3.4 ±5	35.2±8.7	9.7±13.9
Expert	21	107	8.6 ±4	0.59±0.65	13.2 ±3.4	4.2 ±4	32.6±9.7	8.5±9.9
Mann-Whitney U-test			0.004	0.06	0.01	0.02	0.007	0.58

857   **Table 1- Learning induced physiological changes**

858   A summary table of the complete dataset of recordings from excitatory neurons in naïve and

859   expert mice after perceptual learning of pure tones. Columns show different parameters of the

860   dataset or property tested. The third row shows the statistical p value between naïve and experts

861   using a Mann-Whitney U-test.

Group	Animals (n)	Cells (n)	Spontaneous firing rate (Hz)	Evoked firing rate (Hz)	Response latency (ms)	Response selectivity (% of evoked syllables)	Fano-Factor
Naïve	8	41	1.8±2.9	7.6±5.9	36±11	25.4±21.2	1 ±0.45
Expert	6	37	0.77±0.7	6.6±3.6	43±13	15.3 ±12.2	1.1 ±0.35
Mann-Whitney U-test			P=0.1	P=0.48	P=0.02	p=0.046	P=0.12

864   **Table 2- Vocalization learning induced physiological changes**

865   A summary table of the complete dataset of recordings from excitatory neurons in naïve and

866   expert mice after perceptual learning of vocalizations. Columns show different parameters of the

867   dataset or property tested. The third row shows the statistical p value between naïve and experts

868   using a Mann-Whitney U-test.

Group	Animals (n)	Cells (n)	BF (kHz)	Spontaneous firing rate (Hz)	Evoked firing rate (Hz)	FR in training band (Hz)	Response latency (ms)	Response selectivity (% of evoked stimuli)
Naïve PV[+] Pure tones	22	120	16.2 ±10.4	2.2±2.1	21±9	9.2±9	31.1±6.7	17±17
Expert PV[+] Pure tones	21	83	11.8 ±8.1	1.7±1.8	22±10	11.9±10	33.2±9.3	25±18
Mann-Whitney U-test			0.006	0.05	0.7	0.02	0.2	<0.001
Naïve PV[+] Vocalizations	6	17		4±3.7	14.3 ±10.8		31.5±8.6	27±14.9
Expert PV[+] Vocalizations	5	18		3.4±3.9	14.9 ±12.9		32±8	25.6±15.6
Mann-Whitney U-test				0.4	0.97		1	0.72

Table 3- Physiological changes in PV[+] neurons

A summary table of the complete dataset of recordings from PV neurons in naïve and expert mice

after perceptual learning of pure tones (rows 1-2) and vocalizations (rows 4-5). Columns show

different parameters of the dataset or property tested. The white color row shows the statistical p

value between naïve and experts using a Mann-Whitney U-test for each group separately.

REFERENCES

Ahissar, Merav, and Shaul Hochstein. 2004. "The reverse hierarchy theory of visual perceptual
learning." *Trends in cognitive sciences* 8 (10):457-464.
Angeloni, C, and MN Geffen. 2018. "Contextual modulation of sound processing in the auditory
cortex." *Current opinion in neurobiology* 49:8-15.
Aoki, Ryo, Tadashi Tsubota, Yuki Goya, and Andrea Benucci. 2017. "An automated platform for
high-throughput mouse behavior and physiology with voluntary head-fixation." *Nature*
communications 8 (1):1196.
Bakin, Jonathan S, and Norman M Weinberger. 1990. "Classical conditioning induces CS-specific
receptive field plasticity in the auditory cortex of the guinea pig." *Brain research* 536
(1):271-286.
Ball, Karlene, and Robert Sekuler. 1987. "Direction-specific improvement in motion
discrimination." *Vision research* 27 (6):953-965.
Bandyopadhyay, Sharba, Shihab A Shamma, and Patrick O Kanold. 2010. "Dichotomy of
functional organization in the mouse auditory cortex." *Nature neuroscience* 13 (3):361-
368.
Bao, Shaowen. 2015. "Perceptual learning in the developing auditory cortex." *European Journal*
of Neuroscience 41 (5):718-724.
Barth, Alison L, and James FA Poulet. 2012. "Experimental evidence for sparse firing in the
neocortex." *Trends in neurosciences* 35 (6):345-355.
Bathellier, Brice, Lyubov Ushakova, and Simon Rumpel. 2012. "Discrete neocortical dynamics
predict behavioral categorization of sounds." *Neuron* 76 (2):435-449.
Berardi, N, and A Fiorentini. 1987. "Interhemispheric transfer of visual information in humans:
spatial characteristics." *The Journal of physiology* 384 (1):633-647.
Bieszczad, Kasia M, and Norman M Weinberger. 2010. "Representational gain in cortical area
underlies increase of memory strength." *Proceedings of the National Academy of*
Sciences 107 (8):3793-3798.
Bizley, Jennifer K, and Yale E Cohen. 2013. "The what, where and how of auditory-object
perception." *Nature Reviews Neuroscience* 14 (10):693-707.
Brown, Mel, Dexter RF Irvine, and Valerie N Park. 2004. "Perceptual learning on an auditory
frequency discrimination task by cats: association with changes in primary auditory
cortex." *Cerebral cortex* 14 (9):952-965.
Buonomano, Dean V, and Michael M Merzenich. 1998. "Cortical plasticity: from synapses to
maps." *Annual review of neuroscience* 21 (1):149-186.
Cohen, Lior, Noa Koffman, Hanoch Meiri, Yosef Yarom, Ilan Lampl, and Adi Mizrahi. 2013. "Time-
lapse electrical recordings of single neurons from the mouse neocortex." *Proceedings of*
the National Academy of Sciences:201214434.
Cohen, Lior, and Adi Mizrahi. 2015. "Plasticity during motherhood: changes in excitatory and
inhibitory layer 2/3 neurons in auditory cortex." *Journal of Neuroscience* 35 (4):1806-
1815.
Cohen, Lior, Gideon Rothschild, and Adi Mizrahi. 2011. "Multisensory integration of natural
odors and sounds in the auditory cortex." *Neuron* 72 (2):357-369.
Courtin, Julien, Fabrice Chaudun, Robert R Rozeske, Nikolaos Karalis, Cecilia Gonzalez-Campo,
Hélène Wurtz, Azzedine Abdi, Jerome Baufreton, Thomas CM Bienvenu, and Cyril Herry.

2014. "Prefrontal parvalbumin interneurons shape neuronal activity to drive fear
expression." *Nature* 505 (7481):92-96.
Crist, Roy E, Li Wu, and Charles D Gilbert. 2001. "Learning to see: experience and attention in
primary visual cortex." *Nature neuroscience* 4 (5):519.
Egnor, S. E., and K. Branson. 2016. "Computational Analysis of Behavior." *Annu Rev Neurosci*
39:217-36. doi: 10.1146/annurev-neuro-070815-013845.
Ericsson, K Anders. 2006. "The influence of experience and deliberate practice on the
development of superior expert performance." *The Cambridge handbook of expertise
and expert performance* 38:685-705.
Erskine, Andrew, Thorsten Bus, Jan T Herb, and Andreas T Schaefer. 2018. "AutonoMouse: High
throughput automated operant conditioning shows progressive behavioural impairment
with graded olfactory bulb lesions." *bioRxiv*:291815.
Feng, Lei, and Xiaoqin Wang. 2017. "Harmonic template neurons in primate auditory cortex
underlying complex sound processing." *Proceedings of the National Academy of
Sciences* 114 (5):E840-E848.
Froemke, Robert C. 2015. "Plasticity of cortical excitatory-inhibitory balance." *Annual review of
neuroscience* 38:195-219.
Galindo-Leon, Edgar E, Frank G Lin, and Robert C Liu. 2009. "Inhibitory plasticity in a lateral band
improves cortical detection of natural vocalizations." *Neuron* 62 (5):705-716.
Ghose, Geoffrey M, Tianming Yang, and John HR Maunsell. 2002. "Physiological correlates of
perceptual learning in monkey V1 and V2." *Journal of neurophysiology* 87 (4):1867-
1888.
Gibson, Eleanor Jack. 1969. "Principles of perceptual learning and development."
Gilbert, Charles D, Mariano Sigman, and Roy E Crist. 2001. "The neural basis of perceptual
learning." *Neuron* 31 (5):681-697.
Goel, Anubhuti, Daniel Cantu, Janna Guilfoyle, Gunvant R Chaudhari, Aditi Newadkar, Barbara
Todisco, Diego de Alba, Nazim Kourdougli, Lauren M Schmitt, and Ernest Pedapati. 2017.
"Impaired perceptual learning in Fragile X syndrome is mediated by parvalbumin neuron
dysfunction in V1 and is reversible." *bioRxiv*:217414.
Han, Yoon K, Hania Köver, Michele N Insanally, John H Semerdjian, and Shaowen Bao. 2007.
"Early experience impairs perceptual discrimination." *Nature neuroscience* 10 (9):1191.
Harper, Nicol S, Oliver Schoppe, Ben DB Willmore, Zhanfeng Cui, Jan WH Schnupp, and Andrew J
King. 2016. "Network receptive field modeling reveals extensive integration and multi-
feature selectivity in auditory cortical neurons." *PLoS computational biology* 12
(11):e1005113.
Hattori, Ryoma, Kishore V Kuchibhotla, Robert C Froemke, and Takaki Komiyama. 2017.
"Functions and dysfunctions of neocortical inhibitory neuron subtypes." *Nature
neuroscience* 20 (9):1199.
Hawkey, David JC, Sygal Amitay, and David R Moore. 2004. "Early and rapid perceptual
learning." *Nature neuroscience* 7 (10):1055.
Hennequin, Guillaume, Everton J Agnes, and Tim P Vogels. 2017. "Inhibitory plasticity: balance,
control, and codependence." *Annual review of neuroscience* 40:557-579.
Hensch, Takao K. 2005. "Critical period plasticity in local cortical circuits." *Nature Reviews
Neuroscience* 6 (11):877.
Hippenmeyer, Simon, Eline Vrieseling, Markus Sigrist, Thomas Portmann, Celia Laengle, David R
Ladle, and Silvia Arber. 2005. "A developmental switch in the response of DRG neurons
to ETS transcription factor signaling." *PLoS biology* 3 (5):e159.
Irvine, Dexter RF. 2017. "Plasticity in the auditory system." *Hearing research*.

Irvine, Dexter RF, Russell L Martin, Ester Klimkeit, and Rachel Smith. 2000. "Specificity of
perceptual learning in a frequency discrimination task." *The Journal of the Acoustical
Society of America* 108 (6):2964-2968.
Kaplan, Eitan S, Sam F Cooke, Robert W Komorowski, Alexander A Chubykin, Aurore Thomazeau,
Lena A Khibnik, Jeffrey P Gavornik, and Mark F Bear. 2016. "Contrasting roles for
parvalbumin-expressing inhibitory neurons in two forms of adult visual cortical
plasticity." *Elife* 5:e11450.
Karni, Avi, and Dov Sagi. 1991. "Where practice makes perfect in texture discrimination:
evidence for primary visual cortex plasticity." *Proceedings of the National Academy of
Sciences* 88 (11):4966-4970.
Kato, Hiroyuki K, Samuel K Asinof, and Jeffry S Isaacson. 2017. "Network-level control of
frequency tuning in auditory cortex." *Neuron* 95 (2):412-423. e4.
Kato, Hiroyuki K, Shea N Gillet, and Jeffry S Isaacson. 2015. "Flexible sensory representations in
auditory cortex driven by behavioral relevance." *Neuron* 88 (5):1027-1039.
Krakauer, John W, Asif A Ghazanfar, Alex Gomez-Marin, Malcolm A MacIver, and David Poeppel.
2017. "Neuroscience needs behavior: correcting a reductionist bias." *Neuron* 93 (3):480-
490.
Kuchibhotla, Kishore, and Brice Bathellier. 2018. "Neural encoding of sensory and behavioral
complexity in the auditory cortex." *Current opinion in neurobiology* 52:65-71.
Kullmann, Dimitri M, Alexandre W Moreau, Yamina Bakiri, and Elizabeth Nicholson. 2012.
"Plasticity of inhibition." *Neuron* 75 (6):951-962.
Kurt, Simone, and Günter Ehret. 2010. "Auditory discrimination learning and knowledge transfer
in mice depends on task difficulty." *Proceedings of the National Academy of Sciences*
107 (18):8481-8485.
Lagler, Michael, A Tugrul Ozdemir, Sabria Lagoun, Hugo Malagon-Vina, Zsolt Borhegyi, Romana
Hauer, Anna Jelem, and Thomas Klausberger. 2016. "Divisions of identified parvalbumin-
expressing basket cells during working memory-guided decision making." *Neuron* 91
(6):1390-1401.
Lawrence, Douglas H. 1952. "The transfer of a discrimination along a continuum." *Journal of
Comparative and Physiological Psychology* 45 (6):511.
LeDoux, Joseph E. 2000. "Emotion circuits in the brain." *Annual review of neuroscience* 23
(1):155-184.
Lee, Kwang, Sandra M Holley, Justin L Shobe, Natalie C Chong, Carlos Cepeda, Michael S Levine,
and Sotiris C Masmanidis. 2017. "Parvalbumin interneurons modulate striatal output
and enhance performance during associative learning." *Neuron* 93 (6):1451-1463. e4.
Letzkus, Johannes J, Steffen BE Wolff, Elisabeth MM Meyer, Philip Tovote, Julien Courtin, Cyril
Herry, and Andreas Lüthi. 2011. "A disinhibitory microcircuit for associative fear learning
in the auditory cortex." *Nature* 480 (7377):331.
Luo, Liqun, Edward M Callaway, and Karel Svoboda. 2018. "Genetic dissection of neural circuits:
a decade of progress." *Neuron* 98 (2):256-281.
Madisen, Linda, Theresa A Zwingman, Susan M Sunkin, Seung Wook Oh, Hatim A Zariwala, Hong
Gu, Lydia L Ng, Richard D Palmiter, Michael J Hawrylycz, and Allan R Jones. 2010. "A
robust and high-throughput Cre reporting and characterization system for the whole
mouse brain." *Nature neuroscience* 13 (1):133-140.
Maor, Ido, Amos Shalev, and Adi Mizrahi. 2016. "Distinct spatiotemporal response properties of
excitatory versus inhibitory neurons in the mouse auditory cortex." *Cerebral Cortex* 26
(11):4242-4252.

Margrie, Troy W, Axel H Meyer, Antonio Caputi, Hannah Monyer, Mazahir T Hasan, Andreas T
Schaefer, Winfried Denk, and Michael Brecht. 2003. "Targeted whole-cell recordings in
the mammalian brain in vivo." *Neuron* 39 (6):911-918.
Mermillod, Martial, Aurélia Bugaiska, and Patrick Bonin. 2013. "The stability-plasticity dilemma:
Investigating the continuum from catastrophic forgetting to age-limited learning
effects." *Frontiers in psychology* 4:504.
Mizrahi, Adi, Amos Shalev, and Israel Nelken. 2014. "Single neuron and population coding of
natural sounds in auditory cortex." *Current opinion in neurobiology* 24:103-110.
Murphy, Timothy H, Jamie D Boyd, Federico Bolaños, Matthieu P Vanni, Gergely Silasi, Dirk
Haupt, and Jeff M LeDue. 2016. "High-throughput automated home-cage mesoscopic
functional imaging of mouse cortex." *Nature communications* 7:11611.
Nelken, Israel. 2004. "Processing of complex stimuli and natural scenes in the auditory cortex."
Current opinion in neurobiology 14 (4):474-480.
Nevin, John A. 1969. "Signal detection theory and operant behavior: A review of david m. green
and john a. swets' signal detection theory and psychophysics. 1." *Journal of the
Experimental Analysis of Behavior* 12 (3):475-480.
Ohl, Frank W, Wolfram Wetzel, Thomas Wagner, Alexander Rech, and Henning Scheich. 1999.
"Bilateral ablation of auditory cortex in Mongolian gerbil affects discrimination of
frequency modulated tones but not of pure tones." *Learning & Memory* 6 (4):347-362.
Phillips, Elizabeth AK, and Andrea R Hasenstaub. 2016. "Asymmetric effects of activating and
inactivating cortical interneurons." *Elife* 5:e18383.
Polley, Daniel B, Elizabeth E Steinberg, and Michael M Merzenich. 2006. "Perceptual learning
directs auditory cortical map reorganization through top-down influences." *Journal of
neuroscience* 26 (18):4970-4982.
Rabinowitz, Neil C, Ben DB Willmore, Andrew J King, and Jan WH Schnupp. 2013. "Constructing
noise-invariant representations of sound in the auditory pathway." *PLoS biology* 11
(11):e1001710.
Rabinowitz, Neil C, Ben DB Willmore, Jan WH Schnupp, and Andrew J King. 2011. "Contrast gain
control in auditory cortex." *Neuron* 70 (6):1178-1191.
Ramachandran, VS, and Oliver Braddick. 1973. "Orientation-specific learning in stereopsis."
Perception 2 (3):371-376.
Recanzone, Gregg H, Christoph E Schreiner, and Michael M Merzenich. 1993. "Plasticity in the
frequency representation of primary auditory cortex following discrimination training in
adult owl monkeys." *Journal of Neuroscience* 13 (1):87-103.
Reed, Amanda, Jonathan Riley, Ryan Carraway, Andres Carrasco, Claudia Perez, Vikram
Jakkamsetti, and Michael P Kilgard. 2011. "Cortical map plasticity improves learning but
is not necessary for improved performance." *Neuron* 70 (1):121-131.
Renart, Alfonso, and Christian K Machens. 2014. "Variability in neural activity and behavior."
Current opinion in neurobiology 25:211-220.
Resnik, Jennifer, and Daniel B Polley. 2017. "Fast-spiking GABA circuit dynamics in the auditory
cortex predict recovery of sensory processing following peripheral nerve damage." *eLife*
6:e21452.
Roelfsema, Pieter R, and Anthony Holtmaat. 2018. "Control of synaptic plasticity in deep cortical
networks." *Nature Reviews Neuroscience* 19 (3):166.
Rothschild, Gideon, and Adi Mizrahi. 2015. "Global order and local disorder in brain maps."
Annual review of neuroscience 38.

Rothschild, Gideon, Israel Nelken, and Adi Mizrahi. 2010. "Functional organization and
population dynamics in the mouse primary auditory cortex." *Nature neuroscience* 13
(3):353-360.
Rutkowski, Richard G, and Norman M Weinberger. 2005. "Encoding of learned importance of
sound by magnitude of representational area in primary auditory cortex." *Proceedings*
of the National Academy of Sciences of the United States of America 102 (38):13664-
13669.
Schoups, Aniek, Rufin Vogels, Ning Qian, and Guy Orban. 2001. "Practising orientation
identification improves orientation coding in V1 neurons." *Nature* 412 (6846):549.
Seung, H Sebastian, and Haim Sompolinsky. 1993. "Simple models for reading neuronal
population codes." *Proceedings of the National Academy of Sciences* 90 (22):10749-
10753.
Seybold, Bryan A, Elizabeth AK Phillips, Christoph E Schreiner, and Andrea R Hasenstaub. 2015.
"Inhibitory actions unified by network integration." *Neuron* 87 (6):1181-1192.
Sprekeler, Henning. 2017. "Functional consequences of inhibitory plasticity: homeostasis, the
excitation-inhibition balance and beyond." *Current Opinion in Neurobiology* 43:198-203.
Stiebler, I, R Neulist, I Fichtel, and G Ehret. 1997. "The auditory cortex of the house mouse: left-
right differences, tonotopic organization and quantitative analysis of frequency
representation." *Journal of Comparative Physiology A* 181 (6):559-571.
Talwar, Sanjiv K, and George L Gerstein. 2001. "Reorganization in awake rat auditory cortex by
local microstimulation and its effect on frequency-discrimination behavior." *Journal of*
neurophysiology 86 (4):1555-1572.
Theunissen, Frédéric E, and Julie E Elie. 2014. "Neural processing of natural sounds." *Nature*
Reviews Neuroscience 15 (6):355.
Vapnik, Vladimir. 1998. *Statistical learning theory. 1998*. Vol. 3: Wiley, New York.
Wehr, Michael, and Anthony M Zador. 2003. "Balanced inhibition underlies tuning and sharpens
spike timing in auditory cortex." *Nature* 426 (6965):442.
Weible, Aldis P, Christine Liu, Cristopher M Niell, and Michael Wehr. 2014. "Auditory cortex is
required for fear potentiation of gap detection." *Journal of Neuroscience* 34 (46):15437-
15445.
Weinberger, Norman M. 2004. "Specific long-term memory traces in primary auditory cortex."
Nature Reviews Neuroscience 5 (4):279.
Williamson, Ross S, Misha B Ahrens, Jennifer F Linden, and Maneesh Sahani. 2016. "Input-
specific gain modulation by local sensory context shapes cortical and thalamic responses
to complex sounds." *Neuron* 91 (2):467-481.
Wolff, Steffen BE, Jan Gründemann, Philip Tovote, Sabine Krabbe, Gilad A Jacobson, Christian
Müller, Cyril Herry, Ingrid Ehrlich, Rainer W Friedrich, and Johannes J Letzkus. 2014.
"Amygdala interneuron subtypes control fear learning through disinhibition." *Nature*
509 (7501):453.
Wright, Beverly A, and Matthew B Fitzgerald. 2001. "Different patterns of human discrimination
learning for two interaural cues to sound-source location." *Proceedings of the National*
Academy of Sciences 98 (21):12307-12312.
Zeng, Hongkui, and Joshua R Sanes. 2017. "Neuronal cell-type classification: challenges,
opportunities and the path forward." *Nature Reviews Neuroscience* 18 (9):530-546.
Zhou, Mu, Feixue Liang, Xiaorui R Xiong, Lu Li, Haifu Li, Zhongju Xiao, Huizhong W Tao, and Li I
Zhang. 2014. "Scaling down of balanced excitation and inhibition by active behavioral
states in auditory cortex." *Nature neuroscience* 17 (6):841-850.