Neural correlates of learning pure tones versus natural sounds in the auditory cortex

Ido Maor[1,2], Ravid Shwartz-Ziv[2], Libi Feigin[2], Yishai Elyada[2], Haim Sompolinsky[2,3], and Adi
Mizrahi[1,2]*

[1]Department of Neurobiology, [2]The Edmond and Lily Safra Center for Brain Sciences, [3]Racah
Institute of Physics, The Hebrew University of Jerusalem, Jerusalem, Israel.

* Corresponding author:

Adi Mizrahi: Mizrahi.adi@mail.huji.ac.il

Acknowledgements: We thank Israel Nelken and Livia de Hoz for help in setting up the initial
behavioral experiments. We thank Yoni Cohen, David Pash and Joseph Jubran for writing the
software code of the Educage. We thank Yonatan Loewenstein, Merav Ahissar, and the Mizrahi
lab members for comments on the manuscript. This work was supported by an ERC consolidators
grant to A.M (#616063) and by Gatsby Charitable Foundation.

Author contributions: I.M. and A.M. designed the experiments. R.S.Z and H.S analyzed data.
L.F. conducted experiments. Y.E. constructed the vocalization stimuli and software for their
delivery. I.M. conducted experiments and analyzed the data. I.M. H.S. and A.M. wrote the paper.

Conflict of interest: The authors declare no competing financial interests.

ABSTRACT

Auditory perceptual learning of pure tones causes tonotopic map expansion in the primary

auditory cortex (A1), but the function this plasticity sub-serves is unclear. We developed an

automated training platform called the 'Educage', which was used to train mice on a go/no-go

auditory discrimination task to their perceptual limits, for difficult discriminations among pure

tones or natural sounds. Spiking responses of excitatory and inhibitory L2/3 neurons in mouse A1

revealed learning-induced overrepresentation of the learned frequencies, in accordance with

previous literature. Using a novel computational model to study auditory tuning curves we show

that overrepresentation does not necessarily improve discrimination performance of the network

to the learned tones. In contrast, perceptual learning of natural sounds induced 'sparsening' and

decorrelation of the neural response, and consequently improving discrimination of these

complex sounds. The signature of plasticity in A1 highlights its central role in coding natural

sounds as compared to pure tones.

INTRODUCTION

Learning is accompanied by plastic changes in brain circuits. This plasticity is often viewed as

substrate for improving computations that sub-serve learning and behavior. A well-studied

example of learning-induced plasticity is following perceptual learning where cortical

representations change towards the learned stimuli [1,2]. Whether such changes improve

discrimination is still debated and the mechanisms leading to these changes are still largely

unknown. Furthermore, learning paradigms in animal models are often limited to simplified,

unnatural stimuli that deviate further from real learning experience and that brain circuits actually

compute.

Perceptual learning is an implicit form of lifelong learning during which perceptual

performance improves with practice [3]. Perceptual learning spans a wide range of sensory

modalities and tasks like learning and improving reading skills, acquiring language, or the

learning to discriminate different shades of color and flavors. Extensive psychophysical research

on perceptual learning tasks led to a general agreement on some attributes of this type of learning

[4]. For example, perceptual learning has been shown to be task specific, poorly generalized to

other senses or tasks. It is also largely agreed upon that gradual training is essential for

improvement [5-14]. Given the specificity observed at the behavioral level, functional correlates of

perceptual learning are thought to involve neural circuits as early as primary sensory regions [2,15].

In auditory learning paradigms, changes are already observed at the level of primary auditory

cortex [16]. Learning to discriminate among tones results in tonotopic map plasticity towards the

trained stimulus [17-20]; reviewed in [21]. Notably, however, not all studies could replicate the learning

induced changes in the tonotopic map [22]. Furthermore, artificially induced map plasticity was

shown to be unnecessary for better discrimination performances *per se*, [23,24]. Our understanding

of the mechanisms underlying auditory cortex plasticity remains rudimentary, let alone for more

natural stimuli.

To gain understanding of learning-induced plasticity at single neuron resolution, animal

models have proven very useful. Mice, for example, offer the advantage of a rich genetic

experimental toolkit to study neurons and circuits with high efficiency and specificity [25].

Historically, the weak aspect of using mice as a model was its limited behavioral repertoire to

learn difficult tasks. However, in the past decade technical difficulties to train mice to their limits

have been steadily improving with increasing number of software and hardware tools to probe

mouse behavior in high resolution [26-33]. Here, we developed our own experimental system for

training groups of mice on an auditory perceptual task – an automatic system called the

'Educage'. The Educage is a simple affordable system that allows efficient training of several

mice simultaneously to discriminate among pure tones or complex sounds.

A1 is well known for its tonotopic map plasticity following simple forms of learning in

other animal models [21]. An additional interest in primary auditory cortex is its increasing

recognition as a brain region involved in coding complex sounds [34,35]. We thus asked what are the

similarities and differences in the plastic changes single neurons undergo following training to

discriminate pure tones or natural stimuli. We describe distinct changes in the long-term stimulus

representations by L2/3 neurons of mice following perceptual learning and assess how these

contribute to information processing by local circuits. Using two photon targeted

electrophysiology, we also describe how L2/3 parvalbumin-positive neurons change with respect

to their excitatory counterparts. Our work provides a behavioral, physiological and computational

foundation to questions of auditory-driven plasticity in mice.

METHODS

<u>Animals</u>

A total of n=88, 10-11 week-old female mice were used in this work as follows. 44 mice were

C57BL/6 mice and 44 mice were PV–Cre; Ai9 double-heterozygous (PV × Ai9;[36,37]). All

experiments were approved by the Hebrew University's IACUC.

<u>Behavioral setup</u>

The 'Educage' is a small chamber (10x10x10 cm), opening on one end into a standard animal

home cage where mice can enter and exit freely (Fig. 1a and Supplementary figure 1a). On the

other end, the chamber contains the hardware that drives the system, hardware for identifying

mice and measuring behavioral performance. Specifically, at the port entrance there is a coil radio

antenna (ANTC40 which connected to LID665 stationary decoder; Dorset) followed by infra-red

diodes used to identify mice individually and monitor their presence in the port. This port is the

only access to water for the mice. Water is delivered *via* a solenoid valve (VDW; SMC) allowing

precise control of the water volume provided on each visit. Water is delivered via a water spout,

which is also a lickometer (based on 1 microampere current). An additional solenoid valve is

positioned next to the water spout in order to deliver a mild air puff as a negative reinforcement,

if necessary. For sound stimulation, we positioned a speaker (ES1; TDT), at the center of the top

wall of the chamber. Sound was delivered to the speaker at 125 kHz sampling rate *via* a driver

and a programmable attenuator (ED1, PA5; TDT). For high speed data acquisition, reliable

control, on-board sound delivery, and future flexibility, the system was designed via a field

programmable gate array (FPGA) module and a real-time operating system (RTOS) module

(CompactRIO controller; National Instruments). A custom-made A/D converter was connected to

the CompactRIO controller, mediated signal from infra-red diodes and lickometer and controlled

the valves. A custom code was written in Labview to allow an executable user-friendly interface

with numerous options for user-input and flexibility for designing custom protocols. All software

and hardware design are freely available for download at

https://github.com/MizrahiTeam/Educage.

<u>Training paradigm</u>

Prior to the training, each mouse was implanted, under light and very short period of Isoflurane

anesthesia, with Radio Frequency Identification (RFID; Trovan) chip on the back of its head.

RFID chips allow identification of mice individually, which is then used by the system to control

the precise stimulus delivery and track behavioral performance, on a per-mouse basis. Food and

water were provided *ad libitum*. While access to water was only in the Educage, mice could

engage the water port without restriction. Thus, mice were never deprived of food nor water. At

the beginning of each experiment, RFID-tagged mice were placed in a large home-cage that was

connected to the Educage. Before training, we removed the standard water bottle from the home

cage. Mice were free to explore the Educage and drink the water at the behavioral port. Every

time a mouse entered the behavioral port it was identified individually by the antenna and the

infra-red beam and a trial was initiated. Before learning, any entry to the port immediately

resulted in a drop of water, but no sound was played. Following 24hrs of exploration and

drinking, mice were introduced for the first time to the 'target' stimulus – a series of six 10 kHz

pure tones (100 ms duration, 300 ms interval; 3 ms on and off linear ramps; 62 dB SPL) played

every time the mouse crossed the IR beam. To be rewarded with water, mice were now required

to lick the spout during the 900 ms response window. This stage of operant conditioning lasted

for 2-3 days until mice performed at >70% hit rates.

We then switched the system to the first level of discrimination when mice learned to

identify a non-target stimulus (a 7.1 kHz pure tone series) from the already known target stimulus

(separated by 49%/octave). Thus, on each trial one of two possible stimuli were played for 2.1

seconds- either a target tone or a non-target tone. Mice were free to lick the water spout anytime,

but for the purpose of evaluating mouse decisions we defined a response window and counted the

licking responses only within 900 ms after the sound terminated. Target tones were played at 70%

probability and non-target tones at 30% probability, in a pseudorandom sequence. A 'lick'

response to a target was rewarded with a water drop (15 microliter) and considered as a 'Hit' trial.

A 'no lick' response to a target sound was considered as a 'Miss' trial. A lick response to the non-

target was considered a 'False Alarm' (FA) trial, which was negatively reinforced by a mild air

puff (60 PSI; 600ms) followed by a 9 second 'timeout'. A 'no lick' response to the non-target was

considered a correct rejection (CR) and was not rewarded.

Once mice learned the easy discrimination, we switched the system to the second

discrimination stage. Here, we increased task difficulty by changing the non-target tone to 8.333

kHz, thus decreasing the inter tone distance to 26%/octave. Then, at the following stage, the inter

tone distance was further decreased to 14%/octave and then down to 6%/octave. This last

transition was often done in a gradual manner (»12%/octave»10%/octave»

8%/octave»6%/octave). In some of the animals (n=25), we trained mice to their JND and then

changed the task back to an easier level. For some of the mice (n=13) we played 'catch trials'

during the first and second sessions of the 14%/octave discrimination stages. In catch trials,

different tones spanning the frequency range of the whole training (7-10 kHz) were presented to

the animals in low probability (6% of the total number of sounds), and were neither negatively

nor positively reinforced.

For the vocalizations task, we used playback of pups' wriggling calls (WC) as the target

stimulus. These vocalizations were recorded with a one-quarter inch microphone (Brüel & Kjær)

from P4–P5 PV × Ai9 pups (n = 3), were sampled at 500 kHz and identified offline (Digidata

1322A; Molecular Devices). As the non-target stimulus, we used manipulations of the WC.

During the first stage of the operant learning, mice learned to discriminate between WC and a

fully reversed version of this call. Then, the second manipulation on the non-target stimulus was a

gradual change of the frequency modulation (FM) of each syllable in the call while leaving the

temporal structure of the call intact. To manipulate the syllable FM we used a dynamic linear FM

ramp. This operation multiplies each sampling interval within the syllable, by a dynamic speeding

factor, which changed according to the relative distance from the start and end of the syllable, and

generated a new waveform by interpolation from the original waveform. For example, for a 0.6

speeding factor, the beginning of each syllable was slower by a factor of 0.4 while the end of each

syllable accelerated by a factor of 0.4. The range of sound modulation used here was 0.66-0.9. A

value of 0.66 is away from the WC, 0.9 similar to the WC and 1 exactly the same as the WC. The

basic task design for the non-target sound was as follows: Reverse» 0.66 »0.81» 0.9.

<u>Surgical procedure</u>

Mice were anesthetized with an intraperitoneal injection of ketamine and medetomidine (0.80 and

0.65 mg/kg, respectively) and a subcutaneous injection of Carprofen (0.004 mg/g). Additionally,

dextrose–saline was injected to prevent dehydration. Experiments lasted up to 8 h. The depth of

anesthesia was assessed by monitoring the pinch withdrawal reflex, and ketamine/medetomidine

was added to maintain it. The animal's rectal temperature was monitored continuously and

maintained at $36 \pm 1°C$. For imaging and recording, a custom-made metal pin was glued to the

skull using dental cement and connected to a custom stage to allow precise positioning of the

head relative to the speaker (facing the right ear). The muscle overlying the left auditory cortex

was removed, and a craniotomy ($\sim 2 \times 2$ mm) was performed over A1 (coordinates, 2.3 mm

posterior and 4.2 mm lateral to bregma) as described previously [38-40].

Imaging and electrophysiology

Cell-attached recordings were obtained using targeted patch-clamp recording by a previously

described procedure [40-42]. For visualization, the electrode was filled with a green fluorescent dye

(Alexa Flour-488; 50 μM). Imaging of A1 was performed using an Ultima two-photon

microscope from Prairie Technologies equipped with a 16X water-immersion objective lens (0.8

numerical aperture; CF175; Nikon). Two-photon excitation of the electrode and somata was used

at 930 nm (DeepSee femtosecond laser; Spectraphysics). The recording depths of cell somata

were restricted to subpial depths of 180–420 μm, documented by the multiphoton imaging. Spike

waveform analysis was performed on all recorded cells, verifying that tdTomato+ cells in L2/3

had faster/narrower spikes relative to tdTomato-negative (tdTomato−) cells (See also[43]).

Auditory stimuli

The auditory protocol comprised 18/24 pure tones (100 ms duration, 3 ms on and off linear

ramps) logarithmically spaced between 3-40 kHz and presented at four sound pressure levels (72–

42 dB SPLs). Each stimulus/intensity combination was presented 10/12 times at a rate of 1.4 Hz.

The vocalizations protocol comprised the playback of pups' wriggling calls (WC) and 3

additional frequency modulated (FM) calls, presented at 62 dB SPL for 16 repetitions.

Behavioral data analysis

To evaluate behavioral performance we calculated, for different time bins (normally 20 trials),

Hit and FA rates, which are the probability to lick in response to the target and non-target tones,

respectively. In order to compensate for the individual bias, we used a measure of discriminability

from signal detection theory – d-prime (d'). D' is defined as the difference between the normal

inverse cumulative distribution of the 'hit' and FA rates [44]. D' for each discrimination stage was

calculated based on trials from the last 33% of the indicated stage. Psychometric curves were

extracted based on mouse performance in response to the catch trials. By fitting a sigmoidal

function to these curves we calculated decision boundaries as the inflection point of each curve.

Detection time was calculated for each mouse individually, by determining the time in which lick

patterns in the correct reject *vs.* the hit trials diverged (i.e. the time when significance levels

crossed P<0.001 in a two-sample t-test).

<u>Data analysis - electrophysiology</u>

Data analysis and statistics were performed using custom-written code in MATLAB

(MathWorks). Spikes were extracted from raw voltage traces by thresholding. Spike times were

then assigned to the local peaks of suprathreshold segments and rounded to the nearest

millisecond. For each cell, we obtained peri-stimulus time histogram (PSTH) and determined the

response window as the 100 ms following stimulus onset which evoked the maximal response

integral. Only neurons that had tone-evoked response (determined by a two sample t-test) were

included in our dataset. Based on this response window, we extracted the cell's tuning curve and

frequency-response area (FRA). Evoked firing rate was calculated as the average response to all

frequencies that evoked a significant response. Firing rate in the training band was calculated as

the response to frequencies inside the training band (7-10 kHz), averaged across all intensities.

Best frequency (BF) of each cell was determined as the tone frequency that elicited the strongest

response averaged across all intensities. The selectivity of the cell is the % of all frequency-

intensity combinations that evoked significant response (determined by a two sample t-test).

Pairwise signal correlations (r_{sc}) were calculated as Pearson correlation between FRA's matrices

of neighboring cells. The spontaneous firing rate of the cell was calculated based on the 100 ms

preceding each stimulus presentation. Response latency is the time point after stimulus onset at

which average spike count reached maximum.

<u>Statistical model based on the Independent Basis Functions (IBF) Method</u>

Since the measured responses before and after learning are not from the same cells, we cannot

estimate the changes of individual tuning curves due to learning. Instead, we must rely on

estimated learning-induced changes in the *ensemble* of single-neuron responses. Our goal,

therefore, was to build a statistical model of single-neuron tuning curves before and after

learning. This model is different before and after learning, and was used to estimate the learning-

induced changes in the population of A1 responses. Furthermore, we use this model as a

generative model that allowed us to generate a large number of 'model neurons' with statistically

similar response properties as the measured ones.

In principle, one could use a parametric model, by fitting each observed tuning curve to a

specific shape of functions (e.g., Gaussian tuning curves). However, since the tuning curves of

neurons to tone frequencies do not have symmetric 'gaussian' shapes, and some are bimodal,

fitting them to a parametric model has not been successful. Instead, we chose to model each

single neuron response as a weighted sum of a small set of orthogonal basis functions,

$$r_i(f) = \sum_{l=1}^{K} a_l^i g_l(f)$$

Here, $r_i(f)$ is the firing rate (i.e., the trial-averaged spike count) of the i-th neuron in response to

the stimulus with frequency f; K is the number of orthogonal basis functions denoted by $g_l(f)$

(dependencies in f are in log scale). In order to determine the basis functions and the coefficients,

a_l^i we performed Singular Value Decomposition (SVD) of the matrix of the measured neuronal

firing rates for the 18 values of f. Our model (1) uses a subset of the K modes with the largest

singular values (the determination of K is described below). The SVD yields the coefficients

$a_{l,data}^i$ for the *N observed* neurons and (2) smoothes the resultant SVD f-dependent vectors using

a simple 'moving average' technique to generate the basis functions, $g_l(f)$ (3) Importantly, to use

the SVD as a generative model, for each l we compute the histogram of the $N\ a_{l,data}^i$

coefficients. To generate 'new neurons', we sample each coefficient independently from the

corresponding histogram. In other words, we approximate the joint distribution of the coefficients

by a factorized distribution. This allowed us to explore the effect of changing the number of

neurons that downstream decoders use in order to perform the perceptual task.

Model (1) describes the variability of the population responses to the stimulus, in terms of tuning

curves of the trial averages firing rates. Additional variability in the data is the single trial spike

count. We model these as independent Poisson random variables with means given by $r_i(f)$.

Since neurons are not simultaneously recorded, we do not include noise correlations in the model.

We performed this procedure for the *naïve* and *expert* measured responses separately, so that both

the basis functions and the coefficient histograms are evaluated for the two conditions separately.

Note we do not make Gaussian assumptions about the coefficient histograms. In fact, the

observed histograms are in general far from Gaussian.

<u>The choice of number of basis functions</u>

Due to a limited number of trials that we sampled for each neuron, taking a large value of *K* can

result in over-fitting the model to the noise caused by the finite number of trials. To estimate the

optimal number of basis functions, we evaluated the percentage of response firing rate variability

of the population (i.e, the fraction of the sum of the squared SVD eigenvalues) as a function of *K*.

We also evaluated the parameters of model (1) based on a subset of trials and checked how well it

accounts for the observed tuning curves that are calculated from the test trials. We took *K* that

produces the smallest test error and saturated the fractional variance.

276   We used Model (1) with the above choice of *K* in order to evaluate the discrimination

ability of the population of A1 neurons, by creating *an ensemble* of single-neuron responses for

the naïve and expert conditions. To generate the model neurons, we sampled the coefficients of

the basis function independently from the corresponding histogram of the measured neurons, and

used these neurons for the calculations depicted below.

<u>Fisher Information</u>

We calculated the Fisher Information (FI) for each condition (naïve vs. expert) using our model

(1). FI measure bounds the mean squared error of an (unbiased) estimator of the stimulus from

the noisy single trial neuronal responses. When the neuronal population is large (and they are

noise- independent) FI also determines the discriminability d' of a maximum likelihood

discriminator between two nearby values of the stimulus [45]. Under the above Poisson assumption,

the FI for the *i*-th neuron is equal to $I_i = \frac{r_i'(f)^2}{r_i(f)}$, where $r_i'(f)$ is the derivative of the firing rate

with respect to the stimulus value *f*. The total FI is the sum of the FIs of individual neurons [45].

This has been evaluated in both naïve and expert conditions. Note that the FI are functions of the

stimulus value *f*, around which the discrimination task is performed.

<u>Discrimination by linear readout</u>

We applied a linear decoder to assess the ability to discriminate between nearby stimuli on the

basis of the neuronal population responses. We trained a support vector machine (SVM) with a

linear kernel, which finds an 'optimal' linear classifier that discriminate between two nearby

frequencies on the basis of single-trial vectors of spike counts generated with our generative

model (1) and Poisson variability. We then evaluated the probability of classification errors to test

trials, in both naïve and expert conditions. Since our training set is not linearly separable, we

used SVM with slack variables [46], which incorporates a 'soft' cost for classification errors. Each

classification was iterated maximum 500 times (or until converged). In each iteration, 16 trials

were used for training the classifier and 4 trials were used to test the decoder accuracy.

<u>Data analysis – vocalization responses</u>

Similarity of response to different vocalizations was calculated as Pearson correlation between

the PSTH's of the different stimuli. To quantify life time sparseness we used the following

measure $S= (1 − [(\Sigma r_i /n)2/\Sigma(ri\ 2/n)])/[1 − (1/n)]$, where r_i is the response to the ith syllable in the

original vocalization (averaged across trials) and n is the number of syllables. Values of S near

0% indicate a dense code, and values near 100% indicate a sparse code [47]. Population sparseness

was calculated as 100- the percent of cells that evoked a significant response to each syllable in

the call [48]. Classification of vocalization identity based on population activity was determined

using the Support Vector Machine (SVM) decoder with a linear kernel and slack variables. The

decoder was tested for its accuracy to differentiate between responses to two different

vocalizations. The input to the SVM consisted of the spike count of each neuron in the syllable

response window. The same number of neurons (37) was used in both groups to avoid biases. We

then evaluated the probability of classification errors to test trials, using leave-one-out cross

validation. Each classification was iterated 1000 times. In each iteration, 15 trials were used for

training the classifier and one trial was used to test decoder accuracy. The number of syllables

utilized in the decoder were increased cumulatively.

RESULTS

Behavior – Discrimination of pure tones

To study perceptual learning in mice, we developed a behavioral platform named the 'Educage'

(all software and hardware design are freely available for download at

https://github.com/MizrahiTeam/Educage). The Educage is an automated home-cage training

apparatus designed to be used simultaneously with several mice. One advantage of the Educage

over other procedures is that human interference is brought to minimum and training efficiency

increases. The 'Educage' is a small modular chamber (10x10X10 cm), opening on one end into a

standard animal home cage where mice can enter and exit freely (Fig. 1a). On its other end, the

chamber contains the hardware that drives the system, hardware for identifying mice and

measuring behavioral performance (Fig. 1a, Fig. S1a, and Methods). Mice were free to engage

the behavioral port at their own will, where they consume all of their water intake. Following

habituation, mice were trained on a go/no-go auditory discrimination task to lick in response to a

target tone (a series of 10 kHz pure tones) and withhold licking in response to the non-target tone.

A 'lick' response to a target was rewarded with a water drop (15 microliter) and considered as a

'Hit' trial. A 'no lick' response to a target sound was considered as a 'Miss' trial. A lick response

to the non-target was considered a 'False Alarm' (FA) trial, which was negatively reinforced by a

mild air puff (60 PSI; 600ms) followed by a 9 second 'timeout'. A 'no lick' response to the non-

target was considered a correct rejection (CR) and was not rewarded (Fig. 1a). On average, mice

performed 327 ± 71 trials per day, mainly during dark hours (Fig. S1b).

The initial level of learning was to identify a non-target stimulus (a series of 7.1 kHz pure

tones) from the 10 kHz target stimulus. These stimuli are separated by 49% of an octave and are

perceptually easily separated by mice. Despite the simplicity of the task, behavioral performance

varied widely between mice (Fig. 1b, Fig. S1c). On average, it took mice 54 ± 38 trials to cross our

criterion of learning, which was set arbitrarily at d'=1 (Fig. 1b; dotted line), and gradually

increased to plateau at d'=2.35 ± 0.64 (Fig. 1d). To extend the task to more challenging levels, we

gradually increased task difficulty by changing the non-target tone closer to the target tone. The

target tone remained constant at 10 kHz throughout the experiment and only the non-target

stimulus changed. The lowest distance used between target and non-target was 3%/octave (9.6

kHz vs 10 kHz). A representative example from one mouse's performance in the Educage

throughout a complete experiment is shown in figure 1c. The just noticeable difference (JND) for

each mouse was determined when mice could no longer discriminate (e.g. the JND of the mouse

shown in figure 1c was determined between 6-10%/octave). The range of JND's was 3-

14%/octave and averaged 8.6±4.7%/octave. These values of JND are typical for frequency

discrimination in mice [31,49]. Most mice improved their performance with training (Fig. 1e, f),

showing improved perceptual abilities along the task. The duration to reach JND varied as well,

ranging 3069±1099 trials (14±3 days). Detection times, defined as the time in which lick patterns

in the correct reject trials diverged from the lick patterns of the hit trials, increased monotonically

by ~177 ms per each step of task difficulty demonstrating the increased perceptual load during

the harder tasks (Fig. S1d).

To show that gradual training is necessary for perceptual learning [5,8,11] we trained groups

of littermate mice on different protocols simultaneously. In one group of mice we used a standard

protocol and the animals were trained on the gradually increasing task difficulty described above.

Simultaneously, in the second group of mice – termed 'easy only' – animals were trained

continuously on the easy task. Although both groups of mice trained together, only the mice that

underwent gradual training were able to perform the hard task (Fig. 1g). Taking together, these

data demonstrate the efficiency of the Educage to train groups of mice to become experts in

discriminating between a narrowband of frequencies in a relatively short time and with minimal

human intervention.

Representation of pure tones in A1 following perceptual learning

To evaluate cortical plasticity following perceptual learning, we compared how pure tones are

represented in A1 of expert mice, trained to discriminate between narrowband frequencies, and

age matched naive mice who were never introduced to these tones. We used *in vivo* loose patch

recording of L2/3 neurons in anesthetized mice to record tone-evoked spiking activity in response

to 3-40kHz pure tones (Fig. 2a,b; Table 1). We targeted our recording electrode to the center of

A1 based on previously validated stereotactic coordinates [40]. Loose patch recording enables

superb spatial resolution, is not biased to specific cell types and has superb signal to noise ratio

for spike detection. However, one caveat of this technique is a potential bias of recording sites

along the tonotopic axis. In order to overcome it we measured from neurons in a large number of

animals, such that possible biases are averaged out. In naïve mice, responses were highly

heterogeneous, with best frequencies covering the whole frequency range (Fig. 2c,d; n=105

neurons, n=22 mice, red). In expert mice, best frequencies of tuning curves were biased towards

the frequencies that were presented during learning (Fig. 2c,d; n=107 neurons, n=21 mice, blue).

These data show, as expected from previous literature, that learned frequencies in A1 become

overrepresented at least as measured by the neuron's best frequency (BF). We next showed that

this overrepresentation was specific to the learned tone by training mice on 4 kHz as the target

tone and recorded neurons in a similar manner to the abovementioned experiment. Indeed, L2/3

neurons in A1 of mice training on 4kHz showed BF shifts towards 4 kHz (Fig. S2a). Temporal

responses to the trained frequencies were only slightly different between naive and expert mice.

Specifically, average spiking responses were slightly but significantly faster and stronger in

experts (Fig. S2b, Table 1).

We recorded the responses to pure tones at different intensities and constructed frequency

response areas (Fig. S2c). Pairwise signal correlation of neighboring neurons, calculated from

these frequency response areas were high in naïve mice (0.2±0.28) but even higher in experts

(0.3±0.24; Fig. 2E). Notably, the increased signal correlation was not an artifact of differences in

response properties between naïve and expert group (Fig. S2d) but reflected true similarity in

receptive fields (Fig. S2e). Thus, the basal level of functional heterogeneity in A1 [40,50,51] is

reduced following learning. This learning-induced increase in functional homogeneity of the local

circuit, emphasizes the kind of shift that local circuits undergo. Since neurons in expert mice did

not have wider response areas (Fig. 2F, Table 1), our data suggests that neurons shifted their

response properties towards the learned tones at the expense of frequencies outside the training

band. These results are largely consistent with previous studies in monkeys, cats, and rats

(reviewed in [21]), extending this phenomenon of learning-induced changes in tuning curves to the

mouse, to L2/3 neurons and to local circuits.

A generative model of A1 population responses to pure tones

To what extent does overrepresentation of a learned stimulus sub-serve better discrimination by

the neural population? To answer this question, we built a statistical model of tuning curves of

neurons in A1 using six basis functions (Fig. S3), that correspond to the six largest Singular

Value Decomposition (SVD) vectors of the population responses (Independent Basis Functions

(IBF) method; see Methods). In figure 3a we show two representative examples of tuning curves

and their reconstruction by our model. In contrast to Gaussian fits used previously[46,52], our model

captures the salient features of the shape of the auditory tuning curves (asymmetry,

multimodality), yet also smoothed the *raw* response vectors to reduce overfitting due to finite

sampling.

In order to choose the appropriate number of basis functions, we determined the minimal

number of basis functions that achieves good performance in reconstructing test single trial

responses. In figure 3b we show the fraction of explained variance as a function of the number of

basis functions, K. For both the naïve and expert groups the explained variance reaches above

96% after five basis functions. Figure 3c shows the Mean Square Error (MSE) on the unseen

trials as a function of K in data from both the naïve and expert animals. The MSE exhibits a broad

minimum for K in a range between 6-13. Interestingly, both naïve and experts achieve roughly

the same MSE values, although for experts the MSE values at both low and large values of K are

considerably larger than that of the naïve, presumably due to the smaller number of cells in the

expert conditions. Taken together, we conclude that for these conditions six basis functions are

the appropriate number and used this value for our calculations.

Based on the IBF method described above, we generated a population of tuning curves

(500 'new neurons'), and estimated their total Fisher Information (FI; see Methods). Figure 3d

shows the FI as a function of the stimulus f for both the naïve and expert conditions. Surprisingly,

the FI of the neurons from expert animals was enhanced relative to the naïve group, but only for

stimuli at both flanks of the training band. Importantly, the FI within the band of the trained

frequencies remained unchanged (Fig. 3d, within the black lines). The same result holds true for a

performance of a support vector machine (SVM) classifier. Using SVM to separate any two

frequencies that are 0.2198 octave apart, discriminability (d') values derived from the classifier's

error show similar results as the FI (Fig. 3e; [45]). The value of d' is larger in the expert groups as

compared to the naïve but only outside the training band, whereas within the band

discriminability is not improved (or even slightly compromised).

The results shown in figure 3 do not change qualitatively if we use our SVD model for

the recorded neurons, as opposed to newly modelled neurons. One advantage of having a

generative model for the population responses is that we can generate an unlimited number of

trials and tuning curves. We took advantage of this to explore whether the results of the FI and

SVM change with population size. To answer this question, we evaluated the mean

discrimination performance (over test neurons) as a function of the number of sampled cells, N,

which increases as expected. Consistent with the results of the SVM, the performances in the

naïve and expert groups are similar with slight tendency for a higher accuracy in the naive

population at large Ns (Fig. 3f). In contrast, for frequencies near the training band, the accuracy is

substantially larger in the expert than in the naïve group for virtually all N (Fig. 3g). Thus, it

seems that learning induced changes in tuning curves that do not improve discriminability of the

learned stimuli.

Perceptual learning of natural sounds

Natural sounds are characterized by rich spectro-temporal structures with frequency and

amplitude modulations over time [53]. Discrimination of such complex stimuli could be different

from that of pure tones. Thus, we next designed a task similar to that with the pure tones but

using mouse vocalizations as the training stimuli. We used playback of pups' wriggling calls

(WC) as the target stimulus (Fig. 4a - top). As the non-target stimuli, we used frequency

modulations of the WC; a manipulation that allowed us to morph one stimulus to another by a

continuous metric (Fig. 4a). The range of sound modulation used here was indexed as a "speeding

factor" (see Methods for details). In short, a modulation factor of 0.66 affected the original WC

more than a modulation factor of 0.9 did, and is therefore easier to discriminate (Fig. 4b). To

reach perceptual limits we trained mice gradually, starting with an easy version of the task (WC

vs a temporally reversed version of the WC) and then gradually to modulated calls starting at 0.66

modulation. Once mice reached >80% hit rates we changed the non-target stimulus to more

difficult stimuli until mice could no longer discriminate (Fig. 4c). Mice (n=11) learned the easy

task, i.e., discriminating WC from a 0.66 modulated call, with average d' values of 2.5±0.4

(Fig.4d). On average, mice could only barely discriminate between a WC and its 0.9 modulation

(d' at 0.9 was 1±0.8; Fig. 4d). While these discrimination values were comparable to the

performance of pure tones, detection times were substantially higher (Fig. S4a). For similar d'

values, discriminating between the vocalizations took 300-1000ms longer as compared to the

pure tone tasks (Fig.4e). In addition, learning curves were slower for the vocalization task as

compared to the pure tones task. The average number of trials to reach d'=1 for vocalizations was

195 trials, more than three times longer as compared with pure tones (compare Fig. S4b and 1b,

respectively). These differences may arise from the difference in the delay, inter-syllable interval

and temporal modulation of each stimulus type, which we did not further explore. Taken together,

these behavioral results demonstrate a gradual increase in perceptual difficulty using a

manipulation of a natural sound.

Sparser response in A1 following perceptual learning of natural sounds

To study the neural correlates in A1 that support natural sound discrimination we recorded L2/3

neurons in response to the learned stimuli (Fig. 5a), expecting increased representation of these

particular stimuli. Surprisingly, and in contrast to the results from pure tone discrimination, we

did not find an increase in the representation of the learned stimuli. The fraction of cells

responding to the trained vocalization remained constant (Fig. S5a) as well as the evoked firing

rate for the preferred vocalization or the preferred syllable within a vocalization (Table 2).

Instead, representation in expert mice trained on vocalizations was sparser. Specifically, we

measured the 'lifetime sparseness' [47,48] of each neuron and found that neurons of expert mice had

higher sparseness and were more selective in their response within the call, regardless of their

evoked firing rate (Fig. 5b; Naïve: 47±14%; Expert: 63±15%; p=0.001). Sparseness was also

evident from the average population response to the vocalization (Fig. 5c). Nearly all syllables

had weaker responses, three of which were statistically weaker (Fig. 5c, gray bars). Moreover, the

population sparseness[48] for almost all syllables in the call were higher in expert animal (Fig. 5d;

Naïve: 55±16%; Expert: 70±15%; p=0.026). Notably, the smaller fraction of responses was not

just an apparent sparseness due to increase in trial to trial variability, as reliability of responses by

neurons in the expert group remained similar to reliability of responses in the naïve group (Table

2). Thus, 'sparsening' of A1 responses is a main feature of plasticity following learning to

discriminate complex sounds.

We next asked whether the 'sparsening' described above bears more information to the

learned stimulus. We analyzed population responses by including all neurons from all mice as if

they are a single population (naïve: n=41 neurons from 8 mice; expert: n=37 neurons from 6

mice). We calculated the Pearson correlation of the population response to all responses in a

pairwise manner (Fig. 6a). As compared to naïve mice, the absolute levels of correlations in

expert mice were significantly lower for nearly all pairs of comparisons (Fig. 6a, asterisks). As

expected, weaker modulations of the call and, hence, high similarity among stimuli, were

expressed as higher correlations in the neuronal responses (Fig. 6b). The pairs of stimuli which

mice successfully discriminated in the behavior (0.66 *vs.* the original WC and 0.81 *vs.* the original

WC), had significantly lower correlation in the expert mice (Fig. 6b, rank sum test, p<0.05).

Responses to the more similar stimuli that were near perceptual thresholds (i.e. 0.9 *vs.* the original

WC) were lower in expert mice, but not significantly (Fig. 6b, rank sum test, p>0.05). This

reduced Pearson correlation suggested that plasticity in A1 supports better discrimination among

the learned natural stimuli. Indeed, a SVM decoder performed consistently better in expert mice,

discriminating more accurately the original WC from the manipulated ones (Fig. 6c). As

expected, the decoder performance monotonically increased when utilizing the responses to more

syllables in the call. However, in the expert mice, performances reached a plateau already half

way through the call, suggesting that neuronal responses to the late part of the call carried no

additional information useful for discrimination. Similarly, the correlation of the population

responses along the call, shows that responses were separated already following the first syllable,

but that the lowest level of the correlation was in the 5^{th} - 7^{th} syllables range, which then rapidly

recovered by the end of the call (Fig. S5c). These findings are also consistent with the behavioral

performance of the mice as decisions are made within the first 1.5 seconds of the trial (or the first

7 syllables of the sentence). Specifically, the head of the mouse is often retracted by the time the

late syllables are played (Fig. 4e). Taken together, 'sparser' responses improve neural

discrimination of learned natural sounds.

Learning induced plasticity of Parvalbumin neurons

The mechanisms responsible for the learning-induced changes following learning are currently

unknown. We used mouse genetics and two-photon targeted patch to ask whether local inhibitory

neurons could contribute to the observed plasticity we describe above. To this end, we focused

only on parvalbumin inhibitory (PV[+]) interneurons as recent evidence indicates their role in a

variety of learning related plasticity processes [54-59]. Here we probe the role of inhibition in

perceptual learning by measuring learning induce plasticity in the response properties of the PV[+]

neurons. We trained PV[+]-Cre x Ai9 mice in the Educage and then patched single neurons under

visual guidance (Fig. 7a; [40,41]. In order to increase the sample of PV[+] neurons we used targeted

patch and often patched both PV[+] and PV[-] neurons in the same mice. PV[-] neurons were used as

proxy for excitatory neurons (these neurons were also included in the analysis shown in Fig. 2-6).

All the TdTomato[+] neurons that we patched were also verified as having a fast spike shape (Fig.

7a), a well-established electrophysiological signature of PV[+] cells. PV[+] neurons had response

properties different from PV[-] neurons in accordance with our previous work [40]. For example, PV[+]

neurons responses were stronger and faster to both pure tones and natural sounds (Fig. 7b-c and

Table 3; compare with Fig. 2b and 5a, see also [40]).

541   Following pure tone learning, PV[+] neurons also changed their response profile. On

average, the BF of PV[+] neurons shifted towards the learned frequencies, similar to what we

described for PV[-] neurons (Fig. 8a,b). This result is consistent with recent evidence from the

visual cortex showing increased selectivity to trained stimuli following learning [60].The shift in

tuning curves of PV[+] neurons was also accompanied by a significant widening of their receptive

fields (Fig. S6a). When we compared the BF's of PV[-] and PV[+] neurons within the same brain

(within 250 microns of each other), we found that excitatory and inhibitory neurons became more

functionally homogeneous as compared to naïve mice (Fig. 8c). As both neuronal groups show

similar trends in the shift of their preferred frequencies we rule out a simple scenario whereas

parvalbumin neurons increase their responses in the sidebands of the learning frequency. In other

words, plasticity does not seem to be induced by lateral inhibition *via* parvalbumin neurons, but

rather maintain a strict balance between excitation and inhibition, regardless of whether they are

naïves or experts [61,62]. Note that the peak of the PV$^+$ population response and their BF distribution

was on the outskirts of the training band, rather than within it (compare Fig. 8a,b with Fig. 2c,d),

and concomitantly, the slope of the population responses at the trained frequency band increased

due to learning (Fig. 8a). To assess the computational effect of the plasticity in the inhibitory

neurons' responses we have applied on their responses the same d' and Fisher Information

calculated for the PV$^-$ neurons (Fig. 8d,e). Overall, the discrimination performance of the two cell

populations (when equalized in size) is similar. However, the PV$^+$ population shows a significant

learning related increase in tone discrimination performance by these cells within the training

band (Fig. 8d,e) in contrast to the results for PV$^-$ neurons (Fig. 3d and e). This result is consistent

with the above-mentioned increase in their response slopes in the trained frequency.

Following natural sound learning, we found no significant changes in basic response

properties of the PV$^+$ neurons (Table 3), nor in the degree of sparseness of their representation of

the learned vocalizations (Fig. S6b, .The relationship between the responses of PV$^+$ and their PV$^-$

neighbors remained constant as reflected in the similar slopes of the functions describing PV$^-$

firing versus PV$^+$ firing (Fig. S6c). This result suggests that the excitation-inhibition balance, as

reflected in the responses of PV$^-$ versus PV$^+$, remains. In PV$^+$ neurons, the temporal correlation

along the call as well as the decoding performance from these neurons showed changes that are

qualitatively similar to their PV$^-$ counterparts, but the data across the population was noisier (Fig.

8f-g; Fig. S6d) perhaps due to the smaller sample of the PV$^+$ dataset.

DISCUSSION

Plasticity in frequency tuning following perceptual learning

Shifts in the average stimulus representation towards the learned stimuli is not a new

phenomenon. Similar findings were observed in numerous studies, multiple brain areas, animal

models and sensory systems, including in auditory cortex [10,16,63]. In fact, the model of learning-

induced plasticity in A1, also known as tonotopic map expansion, is an exemplar in neuroscience

[18-20,64]; but see [22,65-67]. Indeed, we also found that the number of neurons tuned to the learned

frequency band increased in expert mice after perceptual learning (Fig. 2). Thus, our results

support the observations of tuning curve plasticity in a primary sensory cortex, showing this here

specifically for L2/3 neurons and extending it to A1 in mice.

583   Since we sampled only a small number of neurons in areas smaller than $250\mu m^3$, our

observations cannot be inferred as direct evidence for tonotopic map expansion. Rather, our data

emphasizes that plastic-shifts occur in local circuits (Fig. 2e). Given that neurons in A1 are

functionally heterogeneous within local circuits [40,50,51], any area in A1 that represents a range of

frequencies prior to learning could become more frequency-tuned once learned. Such a

mechanism allows a wide range of modifications within local circuits to enable increased

representation of the learned stimuli without necessarily perturbing gross tonotopic order. One

advantage of local circuit heterogeneity is that it allows circuits to maintain a dynamic balance

between plasticity and stability [68]. Only mapping the full extent of auditory cortex, at both global

and local scale and preferably over time in the same neurons [69], will enable to reveal the precise

type of changes the cortex undergoes and how it exploits its variability to sub-serve perceptual

learning.

595   Quantifying overrepresentation across a population of neurons and interpreting these in

the context of the learned perceptual tasks have been the subject of extensive research [70]. It has

commonly been assumed that the learning-induced changes in tuning properties improves the

accuracy of coding of the trained stimuli. In particular, for discrimination tasks between two

stimuli differing by a small change in a one-dimensional stimulus parameter, perceptual learning

theory predicts that sharpening of the slope of the tuning curves with respect to this parameter

improves the discriminability of the stimuli [45]. However, the observed increased representation of

the BFs towards the training band in expert mice may not increase over all tuning slopes and may

even decrease them, especially since the slopes tend to be small at the best frequencies. A closer

look at the tuning curves in A1 shows that they are often irregular with multiple slopes and peaks

(i.e. not having simple unimodal Gaussian shapes). Furthermore, the learning induced changes in

the ensemble of tuning curves are not limited to shifting the BF, hence a more quantitative

approach was required to assess the consequences of the observed learning induced plasticity on

discrimination accuracy. This motivated us to develop a method that takes into account the

irregularities in tuning curves of A1 neurons. To this end, we have used a new, SVD based,

generative model (Fig. 3), allowing us to assess the combined effects of changes in BFs as well

other changes in the shapes of the tuning curves. Surprisingly, both Fisher Information analysis as

well as estimated classification errors of an optimal linear classifier, show that learning induced

changes in tuning curves does not improve tone discriminability at trained values. This

conclusion is consistent with previous work on the effect of exposure to tones during

development. Early life experience during development which induces similar changes to those

observed during learning has been argued to decrease tone discriminability for similar reasons [71].

However, in that work, the functional effect of tuning curve changes was consistent with an

observed impaired behavioral performance, suggesting that plasticity in A1 sub-serves

discrimination behavior. In contrast, the stable (or even reduced) accuracy in the coding of the

trained frequency we observed occurs despite the improved behavioral performance after training.

What may this finding underlie?

The changes that we recorded in A1 following pure tone learning do not improve their

contribution to discrimination of the task stimuli, and are thus orthogonal to direct behavioral

performance. One interpretation of this finding is that learning induced plasticity in other brain

regions, either in downstream (or parallel) auditory areas or in task related 'readout' areas enable

improved behavioral performance. Notably, even the representation in naïve mice would be

sufficient for reliable decoding at behavioral resolution – utilizing a population of only two-three

hundred A1 neurons (Fig. 3g). Thus, changes in the tuning of A1 neurons are not necessary for

discriminating the pure-tone task. Further, we suggest that the observed learning-induced changes

in the tuning of pure tones in A1 are the result of unsupervised Hebbian learning induced by over

exposure to the trained tones during the training period; similar to the reported results in early

overexposure [71]. Unsupervised learning signals are not driven by task related reward *per se* and

may increase representation rather than discriminability. Increased representation of trained

stimuli may lead to improved discriminability to untrained tones, as observed experimentally.

Importantly, we find that for natural sounds, learning-induced changes in A1 responses improved

the discriminability of the trained stimuli relative to the naïve responses. This suggests that A1's

primary function is in processing and coding complex auditory stimuli like e.g. natural sounds,

rather than pure tones.

Coding of Natural sounds in A1

Unlike pure tones, natural sounds are composed of rich spectro-temporal energies and include a

set of sound features such as amplitude modulations, frequency modulations, harmonics, and

noise. Although auditory cortex is organized tonotopically, it may not be essential for processing

simple sounds, as these stimuli are accurately represented in earlier stages in the auditory

hierarchy, as early as the brainstem [53,72]. This is consistent with experiments showing that

auditory cortex is necessary for associative fear conditioning with complex sounds [54] but not with

pure tones [73-75].

A multitude of studies have shown that single neurons and populations in A1 respond to

sounds in a non-linear fashion [76-78]. For example, neurons in A1 can be selective to harmonic

content that are prevalent in vocalizations and other natural stimuli [79]. Some neurons in A1 show

strong correlations to global stimulus statistics [80]. Furthermore, neurons in A1 are sensitive to the

fine-grained spectrotemporal environments of the sounds, expressed as strong gain modulation to

local sound statistics [81], as well as to sound contrast and noise [82,83]. All of these features

(harmonics, globally and locally rich statistics, and noise) are well represented in the wriggling

calls we played during learning. Which of these particular sensitivities changes after perceptual

learning is not yet known but one expression of this plasticity can be the increased sparse sound

representation we found here (Fig. 5, S5). Sparseness can take different forms [84]. Here,

sparseness, expressed both in the temporal pattern of individual neurons and at the level of the

population, is mainly a result of reduced number of neurons in the network that respond to any of

the 12 syllables played (Fig. S5). Such increase in selectivity to the syllables could arise from

disparate mechanisms, and changes in the structure of local inhibition was one suspect that we

tested [85].

Inhibitory plasticity follows excitatory plasticity

Cortical inhibitory neurons are central players in many forms of learning [86-88]. Inhibitory

interneurons have been implicated as important for experience dependent plasticity in the

developing auditory system [89], during fear learning in adulthood [54,90], and following injury [91].

Surprisingly, however, and despite the numerous studies on parvalbumin neurons, we could not

find any references in the literature of recordings from parvalbumin neurons after auditory

perceptual learning. Two simple (non mutually exclusive) hypotheses are naively expected. One

is that plasticity in inhibitory neurons are negative mirrors of the plasticity in excitatory neurons.

This would predict that inhibitory neurons would increase their responses to the stimuli for which

responses of excitatory neurons are downregulated (as found in the plasticity of somatostatin

expressing (SOM) neurons following passive sound exposure (Kato et al. 2015), or in

multisensory plasticity in mothers (Cohen and Mizrahi, 2015). The second is that inhibitory

neurons enhance their response to 'lateral' stimuli thus enhancing selectivity to the trained

stimulus, as suggested by the pattern of maternal related plasticity to pup calls (Galindo-Leon,

Lin, and Liu 2009). To the best of our knowledge, our study thus provides a first test of these

hypotheses in the context of perceptual learning. We found no evidence for these scenarios.

Instead, a common motif in the local circuit was that parvalbumin neurons changed in a similar

manner to their excitatory counterparts. These results are in line with the observation that PV

neurons in V1 following visual discrimination task become as selective as their Pyramidal

neighbors [60].

The cortex hosts several types of inhibitory cells [92,93], presumably serving distinct roles.

While PV neurons are considered a rather homogeneous pool of neurons based on molecular

signature, their role in coding sounds is not. This is evident in recordings from pyramidal neurons

in A1 while optogenetically inhibiting the PV$^+$ neurons resulted in mixed effects [94,95]. In the

visual cortex, PV$^+$ cells' activity (and presumably its plasticity) was correlated with stimulus-

specific response potentiation but not in ocular dominance plasticity [56], again, suggesting that PV$^+$

neurons are not necessarily involved in all forms of experience-dependent plasticity. Inhibitory

neurons have been suggested to play a key role in enhancing the detection of behaviorally

significant vocalization by lateral inhibition [96]. But recent imaging data argue that somatostatin

interneurons rather than PV$^+$ interneurons govern lateral inhibition in A1 [97]. Our results are also

consistent with the observation that in contrast to the SOM neurons, the changes in responses

following sound exposure are similar in PV$^+$ and pyramidal neurons [60,67]. To what extent

somatostatin or other interneurons subtypes contribute to excitatory plasticity after auditory

perceptual learning remains to be studied.

Simplifying training with the Educage

Training rodents on perceptual learning paradigms has traditionally been performed manually

over many weeks, sometimes months, of training [11,18]. Behavior is confounded by all sorts of

variability, even in the controlled conditions of a laboratory and within inbred mice of the same

age and sex [98]. The Educage is a new tool for standardizing cognitive tasks, as we demonstrated

here for perceptual learning. The Educage allows training of larger animal datasets, effortlessly

collecting thousands of trials per day, and effectively increasing statistical power for assessing

inter-animal variability. Since mice train simultaneously, it potentially reduces stress-associated

with factors like social deprivation. Other behavioral tools have been introduced in recent years

[31,32,99,100]. Each system has its own strengths or weaknesses in cost, design and operation load

[33](e.g. see a new system for olfactory perceptual learning [30], and new systems with options for

combining physiology and imaging [28]). The simple design and operation of the 'Educage'

together with its affordable hardware and software makes it yet another valuable tool for

simplifying training.

a

Figure 1- Perceptual learning in the 'Educage'

A. Left: Schematic design of the 'Educage' system and its components. **Right:** Schematic representation of the go/no-go auditory discrimination task. CR- correct rejection, FA- False Alarm. **B.** d' values of three representative mice and the population average± s.e.m for the first stage of discrimination. Learning criterion is represented as dashed line (d'=1). **C.** Lick responses to the target tone (solid line) and non-target tone (dashed line) of one representative mouse along

different discrimination stages. **D.** population average d' values for the different discrimination

stages. N=39 mice (mean ± s.e.m). Shades denote the level of difficulty (from 49% to 4-10%

octave apart). **E.** Individual d' values at the end of each level as a function of d' in the beginning

of that level. Shades denote the level of difficulty. Learning criterion represented as dashed lines

(d'=1). **F.** Normalized psychometric curves of five mice calculated from the first (light curves)

and the second (dark curves) 14%/octave session. Light and dark arrows indicate average

decision boundaries in the first and second sessions respectively (Mann-Whitney U-test on

criteria: p=0.03). **G.** d' values in easy (49%/octave) and more difficult (10% /octave)

discrimination stages of individual mice from the 'Easy only' group (filled circles) and from the

perceptual learning group (blank circles). d' values are significantly different between groups

only for the hard discrimination level (Mann-Whitney U-test: ***p<0.001).

Figure 2- Learning induces over representation of the learned stimuli

A. Schematic representation of the experimental setup and a sample of the loose patch recording

showing a representative cell from naïve (left) and expert (right) mice (gray markers indicate tone

stimuli). **B.** Raster plots and peri-stimulus time histograms (PSTH) in response to pure tones of

the cells shown. Gray bars indicate the time of stimulus presentation (100 ms). Color bars and

arrows indicate the training frequency band (7.1-10 kHz) and BF, respectively. **C.** Population

average of normalized response tuning curves of 105 neurons from naïve mice (red) and 107

neurons from expert mice (blue; mean ± s.e.m). Gray area indicates the training frequency band.

Asterisks correspond to frequencies with significant response difference (Mann-Whitney U-test:

*p<0.05). **D.** Best frequency (BF) of individual neurons from naïve mice (red markers) and expert

mice (blue markers; Mann-Whitney U-test: **p<0.01). **E.** Pairwise signal correlations (r_{sc}) values

between all neighboring neuronal pairs in naïve (red) and expert (blue) mice. Neurons in expert

mice have higher r_{sc} (Mann-Whitney U-test: *** p<0.001). **F.** Cumulative distribution of

response selectivity in naïve (red) and expert (blue) mice. Response selectivity was determined as

the % of all frequency-intensity combinations that evoked a significant response. Distributions

are not significantly different (Kolmogorov Smirnov test; p=0.542).

Figure 3- Plasticity in A1 does not improve the discrimination of the learned tones

A. Two representative examples of tuning curves of neurons recorded in A1. Average spike rates
are shown in blue and the SVD approximation of the particular curves are shown in orange. Note
that although the SVD approximation is smooth, it captures the irregular dynamics (i.e. non-

Gaussian) of the tuning curves. **B,C.** The explained variance and error as a measure of the

number of SVD vectors used in the model. **D.** Fisher Information calculated form the tuning

curves of the both populations along the frequency dimension. Note the increased FI for the

expert neurons in the flanks of the training band but not within it (gray band). **E.** Discriminability

(d') of SVM decoder along the range of frequencies. Pairwise comparison along the continuum

are performed for frequencies 0.2198 octave apart. In accordance with 'd' the decoder does not

perform better in the training band (gray shade). **F,G.** Classification performance of the decoder

as a function of the number of neurons in the model. In the training band (F), the performance is

similar for both naïve and expert mice. Outside the training band (0.4396 octave apart; G),

performance improved rapidly in the expert mice.

Figure 4- Perceptual learning of vocalizations

A. Spectrograms of the wriggling call (WC, 'target' stimulus, top panel) and the manipulated

WC's ('non-target' stimuli, bottom panels). **B.** Enlargement of the spectrogram's 4th syllable of

the target WC and the manipulated calls. Top, a large manipulation (speeding factor-0.66), which

is perceptually easy to discriminate from the WC. Bottom, a minor manipulation (speeding

factor-0.9), which is perceptually closer to the WC. **C.** Lick responses to the target tone (solid

line) and non-target tone (dashed line), binned over 50 trials, of one representative mouse along

the different discrimination stages. The task of the first stage was to discriminate between WC vs

reversed playback of the call ('Reverse'). The following stages are different degrees of call

modulation. Titles correspond to the speeding factors used for the non-target stimulus. **D.**

Population average d' values for the different discrimination levels. N=9 mice (mean ± s.e.m).

Shades denote the level of difficulty. **E.** Comparison between detection times during the easy and

difficult stages of pure tone (blank circles) and vocalizations (filled circles) discrimination tasks.

Detection times are significantly different between all groups (Mann-Whitney U-test:

***p<0.001).

Figure 5- Learning complex sounds induces 'sparsening'

A. Representative examples of raster plots in response to 4 modulated wriggling calls from naïve (top) and expert (bottom) mice. Red lines indicate spikes in response windows which are significantly above baseline. The stimulus power spectrum shown in light pink in the background. **B. Top:** Scatter plot of 'life time sparseness' (along the vocalization, see methods) and evoked firing rate for individual neurons from naïve (red circles) and expert (blue circles) mice. The middle range of firing rate distribution (0.5 SD bellow and above the median) is indicated as a blue rectangle. **Bottom:** Life time sparseness for all cells from the middle range of the distribution show significant difference between the groups (Mann-Whitney U-test; p=0.001). **C.** Average normalized PSTHs calculated from all neurons in response to the original WC. Data is

shown overlaid for naïve (red) and expert (blue) mice. In expert mice, only syllables 1, 7 and 11

evoked significantly weaker responses as compared to naïve mice (dark gray bars, Mann-Whitney

U-test; p=0.03, 0.001, 0.03). **D.** Population sparseness (% of cells that evoked a significant

evoked response to a given syllable) of all neurons in naïve (red) and expert (blue) for the

different syllables in the call.

Figure 6- Decorrelation improves coding

A. Matrices describing the average response similarity of individual neurons between all

combinations of stimuli in naïve (left) and expert (right) mice. Each pixel indicates the average

Pearson correlation value calculated from all syllables' evoked spike rate from all neurons to two

different calls. Neurons from expert mice have lower correlation between responses to different

modulated calls (asterisks indicate significant differences between naïve and expert groups;

Mann-Whitney U-test, P<0.05). **B.** Pearson correlation between responses to modulated WCs and

responses to the original WC in naïve (red) and expert (blue) mice (mean± s.e.m). Correlations

are significantly different in the 0.66 and 0.81 modulation (Mann-Whitney U-test: * p<0.05). **C.**

Classification performance of a Support Vector Machine (SVM) decoder. The decoder was tested

for its accuracy to differentiate between the modulated WC stimuli against the original WC. The

performance of the decoder is shown for neurons from the naïve (red) and expert (blue) groups.

Decoder performance is plotted separately for the three different pairs of stimuli. Each point in

each graph shows the number of syllables the decoder was trained on and allowed to use. Error

bars are SEM for 1000 repetitions of leave-one-out cross validation.

Figure 7- Response properties of PV⁺ neurons

A. Left: Schematic representation of the experimental setup for two-photon targeted patch and average spike waveform of 203 PV⁺ neurons. **Right**: Representative two-photon micrograph (projection image of 120 microns) of tdTomato+ cells (red) and the recording electrode (Alexa

Fluor-488, green). **B.** Raster plots and peri-stimulus time histograms (PSTH) in response to pure

tones of a representative PV$^+$ neuron from naïve (left) and expert (right) mice. Gray bars indicate

the time of stimulus presentation (100 ms). Color bars and arrows indicate the training frequency

band (7.1-10 kHz) and BF, respectively. **C.** Representative examples of raster plots from PV$^+$

neurons in response to 4 modulated wriggling calls from naïve (left) and expert (right) mice. Red

lines indicate spikes in response windows which are significantly above baseline. Stimulus power

spectrum shown in light pink in the background.

Figure 8- Plasticity of PV⁺ neurons after learning

A. Population average of normalized response tuning curves of n=120 PV⁺ neurons from naïve

mice (red) and 83 PV⁺ neurons from expert mice (blue; mean ± s.e.m). Gray area indicates the

training frequency band. Asterisks correspond to frequencies with significant response difference

(Mann-Whitney U-test: *p<0.05). **B.** Best frequency (BF) of individual PV⁺ neurons from naïve

mice (red) and expert mice (blue; Mann-Whitney U-test: **p<0.01) **C.** Average distance in BF

between PV⁻ and PV⁺ neurons from the same penetration sites. Distances were significantly

smaller in expert mice (Mann-Whitney U-test: ***p<0.001). **D.** Fisher Information calculated

form the tuning curves of PV⁺ neurons **E.** Discriminability (d') of SVM decoder along the range

of frequencies. **F**. Matrices describing the average response similarity of individual PV⁺ neurons

between all combinations of different stimuli in naïve (left) and expert (middle) mice. Each pixel

indicates the average Pearson correlation value calculated from all syllables evoked spike rate

from all neurons to two different calls. There was no significant difference between correlations

of responses to different modulated calls in naïve and expert mice (Mann-Whitney U-test,

P>0.05). **G.** Pearson correlation between responses to modulated WCs versus the responses to

the original WCs in naïve (red) and expert (blue) mice (mean± s.e.m).). Correlations are not

significantly different for all comparisons (Mann-Whitney U-test: p=0.4, 0.5, 0.5).

TABLES

Group	Animals (n)	Cells (n)	BF (kHz)	Spontaneous spike rate (Hz)	Evoked firing rate (Hz)	FR in T.B (Hz)	Response latency (ms)	Selectivity (% of all stimuli)
Naïve	22	105	11.3 ±8.4	0.45±0.58	15.5 ±5.3	3.4 ±5	35.2±8.7	9.7±13.9
Expert	21	107	8.6 ±4	0.59±0.65	13.2 ±3.4	4.2 ±4	32.6±9.7	8.5±9.9
Mann-Whitney U-test			0.004	0.06	0.01	0.02	0.007	0.58

854 **Table 1- Learning induced physiological changes**

855 A summary table of the complete dataset of recordings from excitatory neurons in naïve and
856 expert mice after perceptual learning of pure tones. Columns show different parameters of the
857 dataset or property tested. The third row shows the statistical p value between naïve and experts
858 using a Mann-Whitney U-test.

Group	Animals (n)	Cells (n)	Spontaneous firing rate (Hz)	Evoked firing rate (Hz)	Response latency (ms)	Response selectivity (% of evoked syllables)	Fano-Factor
Naïve	8	41	1.8±2.9	7.6±5.9	36±11	25.4±21.2	1 ±0.45
Expert	6	37	0.77±0.7	6.6±3.6	43±13	15.3 ±12.2	1.1 ±0.35
Mann-Whitney U-test			P=0.1	P=0.48	P=0.02	p=0.046	P=0.12

861 **Table 2- Vocalization learning induced physiological changes**

862 A summary table of the complete dataset of recordings from excitatory neurons in naïve and
863 expert mice after perceptual learning of vocalizations. Columns show different parameters of the
864 dataset or property tested. The third row shows the statistical p value between naïve and experts
865 using a Mann-Whitney U-test.

Group	Animals (n)	Cells (n)	BF (kHz)	Spontaneous firing rate (Hz)	Evoked firing rate (Hz)	FR in training band (Hz)	Response latency (ms)	Response selectivity (% of evoked stimuli)
Naïve PV$^+$ Pure tones	22	120	16.2 ±10.4	2.2±2.1	21±9	9.2±9	31.1±6.7	17±17
Expert PV$^+$ Pure tones	21	83	11.8 ±8.1	1.7±1.8	22±10	11.9±10	33.2±9.3	25±18
Mann-Whitney U-test			0.006	0.05	0.7	0.02	0.2	<0.001
Naïve PV$^+$ Vocalizations	6	17		4±3.7	14.3 ±10.8		31.5±8.6	27±14.9
Expert PV$^+$ Vocalizations	5	18		3.4±3.9	14.9 ±12.9		32±8	25.6±15.6
Mann-Whitney U-test				0.4	0.97		1	0.72

Table 3- Physiological changes in PV$^+$ neurons

A summary table of the complete dataset of recordings from PV neurons in naïve and expert mice

after perceptual learning of pure tones (rows 1-2) and vocalizations (rows 4-5). Columns show

different parameters of the dataset or property tested. The white color row shows the statistical p

value between naïve and experts using a Mann-Whitney U-test for each group separately.

SUPPLEMENTARY FIGURES

Supplementary figure 1-

A. Photograph of the Educage platform (top view). 1. Home cage; 2. Connecting tunnel; 3. Auditory

speaker; 4. IR diodes monitoring the behavioral port; 5. Water spout and licko-meter; 6. RFID antenna;

7. Connection to D/A converter; 8. Electrically conductive grid floor (for lickometer circuit). **B.**

Histogram of daily hourly distribution of voluntary training. Gray area represents dark hours. N=39

mice (mean ± s.e.m). **C.** Representative examples of the performance of three mice (the same mice

shown in 'Figure 1B') during the first stage of discrimination. The graph shows % of lick responses to

the target tone ('Hit rate'-solid line) and to the non-target tone ('FA rate'- dashed line). **D.** Population

averages of detection times for the different discrimination levels. N=12 mice (mean ± s.e.m). Shades

denote the level of difficulty (from 49% to 10% octave apart).

Supplementary figure 2 -

A. The fraction of neurons with BF that resides within a 4kHz frequency band in three different

experimental groups. Naïve, (n=22 mice) - red; mice trained on a 10kHz (n=21 mice), and 4kHz

(n=4 mice) as the target tone - blue. **B.** Population average of normalized PSTHs from naïve (red)

and expert mice trained (blue; mean ± s.e.m). The PSTHs shown were collapsed for all intensities

and for all frequencies inside the training band, binned at 1 ms time bins. Inset: zoom-in on the

first 50 ms from stimulus onset. **C.** Representative examples of frequency response areas (FRA's)

and pairwise signal correlations (r_{sc}) of neighboring neurons from the same local circuits in naïve

(top) and expert (bottom) mice. **D.** Left, Scatter plot of pairwise r_{sc} and pair average of evoked

firing rate or response Fano-factor for naive (red circles) and expert (blue circles). The range

common to the naïve and expert (0.5 SD bellow and above the median) is indicated as blue

rectangle. Right, All pairwise r_{sc} values from the common range show consistent differences

between groups, independent of these parameters (Mann-Whitney U-test: *** p<0.05). **E.** Scatter

plot of pairwise r_{sc} and pair best frequency difference showing strong dependency (Naïve: slope=-

0.3; p=$2E^{-28}$; R^2= 0.42; Expert: slope=-0.27; p=$5.8E^{-17}$; R^2=0.22).

Supplementary figure 3 -

A-F. 6 basis functions coming out of the SVD analysis for naïve (red) and expert (blue) neurons

Supplementary figure 4 -

A. Population average of detection times for the different discrimination stages. N=9 mice (mean

± s.e.m). Shades denote the level of difficulty. **B.** d' values (mean± s.e.m) for the first stage of

discrimination (WC vs Reverse). The learning criterion is represented as dashed line (d'=1).

Supplementary Figure 5 -

A. Fraction of cells with significant response to frequencies within the training band (left) and to

the original WC in naïve (red) and expert (blue) mice (mean ± s.e.m). Fraction of cells

responding to the trained vocalization remained constant (Mann-Whitney U-test: p=0.09) **B.**

Correlation matrices for different syllables individually. Each pixel indicates the average Pearson

correlation value from two different calls calculated from the evoked spike rate from all neurons

in response to one syllable. Neurons from expert mice have lower correlation between responses

to different modulated calls, and specifically in syllables 5-7.

Supplementary Figure 6-

A. Cumulative distribution of response selectivity of PV[+] neurons in naïve (red) and expert (blue)

mice. Response selectivity was determined as the % of all frequency-intensity combinations that

evoked significant response. PV[+] neurons in expert were less selective (Kolmogorov Smirnov

test; p=0.0015). **B.** Life time sparseness for all cells from the middle range of the firing rate

distribution show no significant difference between naïve (red circles) and expert (blue circles)

groups (Mann-Whitney U-test; p=0.46).   **C.** Scatter plot showing linear dependency of average

syllable evoked firing rate of PV[-] and PV[+] neurons in naïve (red; R^2=0.5; slope=0.37; p=0.008)

and expert (blue red; R^2=0.7; slope=0.31; p<0.001) mice. Each marker corresponds to the

average response for one syllable in the sentence. **D.** Classification performance of a Support

Vector Machine (SVM) decoder based on PV[+] responses.

REFERENCES

1   Roelfsema, P. R. & Holtmaat, A. Control of synaptic plasticity in deep cortical networks.
Nature Reviews Neuroscience **19**, 166 (2018).
2   Gilbert, C. D., Sigman, M. & Crist, R. E. The neural basis of perceptual learning. *Neuron*
31, 681-697 (2001).
3   Gibson, E. J. Principles of perceptual learning and development. (1969).
4   Hawkey, D. J., Amitay, S. & Moore, D. R. Early and rapid perceptual learning. *Nature*
neuroscience **7**, 1055 (2004).
5   Ahissar, M. & Hochstein, S. The reverse hierarchy theory of visual perceptual learning.
Trends in cognitive sciences **8**, 457-464 (2004).
6   Ball, K. & Sekuler, R. Direction-specific improvement in motion discrimination. *Vision*
research **27**, 953-965 (1987).
7   Berardi, N. & Fiorentini, A. Interhemispheric transfer of visual information in humans:
spatial characteristics. *The Journal of physiology* **384**, 633-647 (1987).
8   Ericsson, K. A. The influence of experience and deliberate practice on the development
of superior expert performance. *The Cambridge handbook of expertise and expert*
performance **38**, 685-705 (2006).
9   Irvine, D. R., Martin, R. L., Klimkeit, E. & Smith, R. Specificity of perceptual learning in a
frequency discrimination task. *The Journal of the Acoustical Society of America* **108**,
2964-2968 (2000).
10   Karni, A. & Sagi, D. Where practice makes perfect in texture discrimination: evidence for
primary visual cortex plasticity. *Proceedings of the National Academy of Sciences* **88**,
4966-4970 (1991).
11   Kurt, S. & Ehret, G. Auditory discrimination learning and knowledge transfer in mice
depends on task difficulty. *Proceedings of the National Academy of Sciences* **107**, 8481-
8485 (2010).
12   Lawrence, D. H. The transfer of a discrimination along a continuum. *Journal of*
Comparative and Physiological Psychology **45**, 511 (1952).
13   Ramachandran, V. & Braddick, O. Orientation-specific learning in stereopsis. *Perception*
2, 371-376 (1973).
14   Wright, B. A. & Fitzgerald, M. B. Different patterns of human discrimination learning for
two interaural cues to sound-source location. *Proceedings of the National Academy of*
Sciences **98**, 12307-12312 (2001).
15   Schoups, A., Vogels, R., Qian, N. & Orban, G. Practising orientation identification
improves orientation coding in V1 neurons. *Nature* **412**, 549 (2001).
16   Weinberger, N. M. Specific long-term memory traces in primary auditory cortex. *Nature*
Reviews Neuroscience **5**, 279-290 (2004).
17   Bieszczad, K. M. & Weinberger, N. M. Representational gain in cortical area underlies
increase of memory strength. *Proceedings of the National Academy of Sciences* **107**,
3793-3798 (2010).
18   Polley, D. B., Steinberg, E. E. & Merzenich, M. M. Perceptual learning directs auditory
cortical map reorganization through top-down influences. *Journal of neuroscience* **26**,
4970-4982 (2006).

19   Recanzone, G. H., Schreiner, C. E. & Merzenich, M. M. Plasticity in the frequency
representation of primary auditory cortex following discrimination training in adult owl
monkeys. *Journal of Neuroscience* **13**, 87-103 (1993).
20   Rutkowski, R. G. & Weinberger, N. M. Encoding of learned importance of sound by
magnitude of representational area in primary auditory cortex. *Proceedings of the
National Academy of Sciences of the United States of America* **102**, 13664-13669 (2005).
21   Irvine, D. R. Plasticity in the auditory system. *Hearing research* (2017).
22   Brown, M., Irvine, D. R. & Park, V. N. Perceptual learning on an auditory frequency
discrimination task by cats: association with changes in primary auditory cortex.
Cerebral cortex **14**, 952-965 (2004).
23   Talwar, S. K. & Gerstein, G. L. Reorganization in awake rat auditory cortex by local
microstimulation and its effect on frequency-discrimination behavior. *Journal of
neurophysiology* **86**, 1555-1572 (2001).
24   Reed, A. *et al.* Cortical map plasticity improves learning but is not necessary for
improved performance. *Neuron* **70**, 121-131 (2011).
25   Luo, L., Callaway, E. M. & Svoboda, K. Genetic dissection of neural circuits: a decade of
progress. *Neuron* **98**, 256-281 (2018).
26   Egnor, S. E. & Branson, K. Computational Analysis of Behavior. *Annu Rev Neurosci* **39**,
217-236, doi:10.1146/annurev-neuro-070815-013845 (2016).
27   Krakauer, J. W., Ghazanfar, A. A., Gomez-Marin, A., MacIver, M. A. & Poeppel, D.
Neuroscience needs behavior: correcting a reductionist Bias. *Neuron* **93**, 480-490 (2017).
28   Aoki, R., Tsubota, T., Goya, Y. & Benucci, A. An automated platform for high-throughput
mouse behavior and physiology with voluntary head-fixation. *Nature communications* **8**,
1196 (2017).
29   Murphy, T. H. *et al.* High-throughput automated home-cage mesoscopic functional
imaging of mouse cortex. *Nature communications* **7**, 11611 (2016).
30   Erskine, A., Bus, T., Herb, J. T. & Schaefer, A. T. AutonoMouse: High throughput
automated operant conditioning shows progressive behavioural impairment with
graded olfactory bulb lesions. *bioRxiv*, 291815 (2018).
31   de Hoz, L. & Nelken, I. Frequency tuning in the behaving mouse: different bandwidths
for discrimination and generalization. *PloS one* **9**, e91676 (2014).
32   Cruces-Solís, H. *et al.* Auditory midbrain coding of statistical learning that results from
discontinuous sensory stimulation. *PLoS biology* **16**, e2005114 (2018).
33   Francis, N. A. & Kanold, P. O. Automated operant conditioning in the mouse home cage.
Frontiers in neural circuits **11**, 10 (2017).
34   Bizley, J. K. & Cohen, Y. E. The what, where and how of auditory-object perception.
Nature Reviews Neuroscience **14**, 693-707 (2013).
35   Kuchibhotla, K. & Bathellier, B. Neural encoding of sensory and behavioral complexity in
the auditory cortex. *Current opinion in neurobiology* **52**, 65-71 (2018).
36   Hippenmeyer, S. *et al.* A developmental switch in the response of DRG neurons to ETS
transcription factor signaling. *PLoS biology* **3**, e159 (2005).
37   Madisen, L. *et al.* A robust and high-throughput Cre reporting and characterization
system for the whole mouse brain. *Nature neuroscience* **13**, 133-140 (2010).
38   Cohen, L., Rothschild, G. & Mizrahi, A. Multisensory integration of natural odors and
sounds in the auditory cortex. *Neuron* **72**, 357-369 (2011).
39   Stiebler, I., Neulist, R., Fichtel, I. & Ehret, G. The auditory cortex of the house mouse:
left-right differences, tonotopic organization and quantitative analysis of frequency
representation. *Journal of Comparative Physiology A* **181**, 559-571 (1997).

| 1036 | 40 | Maor, I., Shalev, A. & Mizrahi, A. Distinct spatiotemporal response properties of excitatory versus inhibitory neurons in the mouse auditory cortex. *Cerebral Cortex* **26**, 4242-4252 (2016). |

40   Maor, I., Shalev, A. & Mizrahi, A. Distinct spatiotemporal response properties of
excitatory versus inhibitory neurons in the mouse auditory cortex. *Cerebral Cortex* **26**,
4242-4252 (2016).
41   Cohen, L. & Mizrahi, A. Plasticity during motherhood: changes in excitatory and
inhibitory layer 2/3 neurons in auditory cortex. *Journal of Neuroscience* **35**, 1806-1815
(2015).
42   Margrie, T. W. *et al.* Targeted whole-cell recordings in the mammalian brain in vivo.
Neuron **39**, 911-918 (2003).
43   Bizley, J. K. & Cohen, Y. E. The what, where and how of auditory-object perception.
Nature Reviews. Neuroscience **14**, 693 (2013).
44   Nevin, J. A. SIGNAL DETECTION THEORY AND OPERANT BEHAVIOR: A Review of David M.
Green and John A. Swets' Signal Detection Theory and Psychophysics. *Journal of the
Experimental Analysis of Behavior* **12**, 475-480 (1969).
45   Seung, H. S. & Sompolinsky, H. Simple models for reading neuronal population codes.
Proceedings of the National Academy of Sciences **90**, 10749-10753 (1993).
46   Vapnik, V. *Statistical learning theory. 1998*. Vol. 3 (Wiley, New York, 1998).
47   Vinje, W. E. & Gallant, J. L. Sparse coding and decorrelation in primary visual cortex
during natural vision. *Science* **287**, 1273-1276 (2000).
48   Willmore, B. & Tolhurst, D. J. Characterizing the sparseness of neural codes. *Network:
Computation in Neural Systems* **12**, 255-270 (2001).
49   Ehret, G. Frequency and intensity difference limens and nonlinearities in the ear of the
housemouse (Mus musculus). *Journal of Comparative Physiology A: Neuroethology,
Sensory, Neural, and Behavioral Physiology* **102**, 321-336 (1975).
50   Rothschild, G., Nelken, I. & Mizrahi, A. Functional organization and population dynamics
in the mouse primary auditory cortex. *Nature neuroscience* **13**, 353-360 (2010).
51   Bandyopadhyay, S., Shamma, S. A. & Kanold, P. O. Dichotomy of functional organization
in the mouse auditory cortex. *Nature neuroscience* **13**, 361-368 (2010).
52   Briguglio, J. J., Aizenberg, M., Balasubramanian, V. & Geffen, M. N. Cortical neural
activity predicts sensory acuity under optogenetic manipulation. *Journal of
Neuroscience*, 2457-2417 (2018).
53   Mizrahi, A., Shalev, A. & Nelken, I. Single neuron and population coding of natural
sounds in auditory cortex. *Current opinion in neurobiology* **24**, 103-110 (2014).
54   Letzkus, J. J. *et al.* A disinhibitory microcircuit for associative fear learning in the auditory
cortex. *Nature* **480**, 331 (2011).
55   Wolff, S. B. *et al.* Amygdala interneuron subtypes control fear learning through
disinhibition. *Nature* **509**, 453 (2014).
56   Kaplan, E. S. *et al.* Contrasting roles for parvalbumin-expressing inhibitory neurons in
two forms of adult visual cortical plasticity. *Elife* **5**, e11450 (2016).
57   Lagler, M. *et al.* Divisions of identified parvalbumin-expressing basket cells during
working memory-guided decision making. *Neuron* **91**, 1390-1401 (2016).
58   Lee, K. *et al.* Parvalbumin interneurons modulate striatal output and enhance
performance during associative learning. *Neuron* **93**, 1451-1463. e1454 (2017).
59   Goel, A. *et al.* Impaired perceptual learning in Fragile X syndrome is mediated by
parvalbumin neuron dysfunction in V1 and is reversible. *bioRxiv*, 217414 (2017).
60   Khan, A. G. *et al.* Distinct learning-induced changes in stimulus selectivity and
interactions of GABAergic interneuron classes in visual cortex. *Nature neuroscience*, 1
(2018).

61   Wehr, M. & Zador, A. M. Balanced inhibition underlies tuning and sharpens spike timing
in auditory cortex. *Nature* **426**, 442 (2003).
62   Zhou, M. *et al.* Scaling down of balanced excitation and inhibition by active behavioral
states in auditory cortex. *Nature neuroscience* **17**, 841-850 (2014).
63   Buonomano, D. V. & Merzenich, M. M. Cortical plasticity: from synapses to maps.
Annual review of neuroscience **21**, 149-186 (1998).
64   Bakin, J. S. & Weinberger, N. M. Classical conditioning induces CS-specific receptive field
plasticity in the auditory cortex of the guinea pig. *Brain research* **536**, 271-286 (1990).
65   Crist, R. E., Wu, L. & Gilbert, C. D. Learning to see: experience and attention in primary
visual cortex. *Nature neuroscience* **4**, 519 (2001).
66   Ghose, G. M., Yang, T. & Maunsell, J. H. Physiological correlates of perceptual learning in
monkey V1 and V2. *Journal of neurophysiology* **87**, 1867-1888 (2002).
67   Kato, H. K., Gillet, S. N. & Isaacson, J. S. Flexible sensory representations in auditory
cortex driven by behavioral relevance. *Neuron* **88**, 1027-1039 (2015).
68   Mermillod, M., Bugaiska, A. & Bonin, P. The stability-plasticity dilemma: Investigating
the continuum from catastrophic forgetting to age-limited learning effects. *Frontiers in*
psychology **4**, 504 (2013).
69   Rothschild, G. & Mizrahi, A. Global order and local disorder in brain maps. *Annual review*
of neuroscience **38** (2015).
70   Bao, S. Perceptual learning in the developing auditory cortex. *European Journal of*
Neuroscience **41**, 718-724 (2015).
71   Han, Y. K., Köver, H., Insanally, M. N., Semerdjian, J. H. & Bao, S. Early experience
impairs perceptual discrimination. *Nature neuroscience* **10**, 1191 (2007).
72   Nelken, I. Processing of complex stimuli and natural scenes in the auditory cortex.
Current opinion in neurobiology **14**, 474-480 (2004).
73   Weible, A. P., Liu, C., Niell, C. M. & Wehr, M. Auditory cortex is required for fear
potentiation of gap detection. *Journal of Neuroscience* **34**, 15437-15445 (2014).
74   LeDoux, J. E. Emotion circuits in the brain. *Annual review of neuroscience* **23**, 155-184
(2000).
75   Ohl, F. W., Wetzel, W., Wagner, T., Rech, A. & Scheich, H. Bilateral ablation of auditory
cortex in Mongolian gerbil affects discrimination of frequency modulated tones but not
of pure tones. *Learning & Memory* **6**, 347-362 (1999).
76   Angeloni, C. & Geffen, M. Contextual modulation of sound processing in the auditory
cortex. *Current opinion in neurobiology* **49**, 8-15 (2018).
77   Harper, N. S. *et al.* Network receptive field modeling reveals extensive integration and
multi-feature selectivity in auditory cortical neurons. *PLoS computational biology* **12**,
e1005113 (2016).
78   Bathellier, B., Ushakova, L. & Rumpel, S. Discrete neocortical dynamics predict
behavioral categorization of sounds. *Neuron* **76**, 435-449 (2012).
79   Feng, L. & Wang, X. Harmonic template neurons in primate auditory cortex underlying
complex sound processing. *Proceedings of the National Academy of Sciences* **114**, E840-
E848 (2017).
80   Theunissen, F. E. & Elie, J. E. Neural processing of natural sounds. *Nature Reviews*
Neuroscience **15**, 355 (2014).
81   Williamson, R. S., Ahrens, M. B., Linden, J. F. & Sahani, M. Input-specific gain modulation
by local sensory context shapes cortical and thalamic responses to complex sounds.
Neuron **91**, 467-481 (2016).

82   Rabinowitz, N. C., Willmore, B. D., King, A. J. & Schnupp, J. W. Constructing noise-
invariant representations of sound in the auditory pathway. *PLoS biology* **11**, e1001710
(2013).
83   Rabinowitz, N. C., Willmore, B. D., Schnupp, J. W. & King, A. J. Contrast gain control in
auditory cortex. *Neuron* **70**, 1178-1191 (2011).
84   Barth, A. L. & Poulet, J. F. Experimental evidence for sparse firing in the neocortex.
Trends in neurosciences **35**, 345-355 (2012).
85   Froemke, R. C. Plasticity of cortical excitatory-inhibitory balance. *Annual review of*
neuroscience **38**, 195-219 (2015).
86   Hennequin, G., Agnes, E. J. & Vogels, T. P. Inhibitory Plasticity: Balance, Control, and
Codependence. *Annual Review of Neuroscience* **40** (2017).
87   Kullmann, D. M., Moreau, A. W., Bakiri, Y. & Nicholson, E. Plasticity of inhibition. *Neuron*
75, 951-962 (2012).
88   Sprekeler, H. Functional consequences of inhibitory plasticity: homeostasis, the
excitation-inhibition balance and beyond. *Current Opinion in Neurobiology* **43**, 198-203
(2017).
89   Hensch, T. K. Critical period plasticity in local cortical circuits. *Nature Reviews*
Neuroscience **6**, 877-888 (2005).
90   Courtin, J. *et al.* Prefrontal parvalbumin interneurons shape neuronal activity to drive
fear expression. *Nature* **505**, 92-96 (2014).
91   Resnik, J. & Polley, D. B. Fast-spiking GABA circuit dynamics in the auditory cortex
predict recovery of sensory processing following peripheral nerve damage. *eLife* **6**,
e21452 (2017).
92   Hattori, R., Kuchibhotla, K. V., Froemke, R. C. & Komiyama, T. Functions and
dysfunctions of neocortical inhibitory neuron subtypes. *Nature neuroscience* **20**, 1199
(2017).
93   Zeng, H. & Sanes, J. R. Neuronal cell-type classification: challenges, opportunities and
the path forward. *Nature Reviews Neuroscience* **18**, 530-546 (2017).
94   Seybold, B. A., Phillips, E. A., Schreiner, C. E. & Hasenstaub, A. R. Inhibitory actions
unified by network integration. *Neuron* **87**, 1181-1192 (2015).
95   Phillips, E. A. & Hasenstaub, A. R. Asymmetric effects of activating and inactivating
cortical interneurons. *Elife* **5**, e18383 (2016).
96   Galindo-Leon, E. E., Lin, F. G. & Liu, R. C. Inhibitory plasticity in a lateral band improves
cortical detection of natural vocalizations. *Neuron* **62**, 705-716 (2009).
97   Kato, H. K., Asinof, S. K. & Isaacson, J. S. Network-Level Control of Frequency Tuning in
Auditory Cortex. *Neuron* (2017).
98   Renart, A. & Machens, C. K. Variability in neural activity and behavior. *Current opinion in*
neurobiology **25**, 211-220 (2014).
99   Poddar, R., Kawai, R. & Ölveczky, B. P. A fully automated high-throughput training
system for rodents. *PLoS One* **8**, e83171 (2013).
100   Murphy, T. H. *et al.* High-throughput automated home-cage mesoscopic functional
imaging of mouse cortex. *Nature communications* **7** (2016).

1172