Transcription factor TAp73 and microRNA-449 complement each other to support

multiciliogenesis

Running title: TAp73 and miR449 cooperate in multiciliogenesis

Merit Wildung[1,*], Tilman Uli Esser[1,*], Katie Baker Grausam[2,3], Cornelia Wiedwald[1], Larisa
Volceanov-Hahn[1], Dietmar Riedel[4], Sabine Beuermann[1], Li Li[2], Jessica Zylla[2], Ann-Kathrin
Guenther[5], Magdalena Wienken[6], Evrim Ercetin[1], Zhiyuan Han[7], Felix Bremmer[8], Orr Shomroni[9],
Stefan Andreas[1], Haotian Zhao[2,3,7,#] and Muriel Lizé[1,#]

* = equal contribution; # = corresponding authors

1) Molecular & Experimental Pneumology Group, Clinic for Cardiology and Pneumology,
12   University Medical Center Goettingen, Germany
13   2) Cancer Biology and Immunotherapeutics Group, Sanford Research, Sioux Falls, South
14   Dakota, USA
15   3) Division of Basic Biomedical Sciences, University of South Dakota, Sanford School of
16   Medicine, Vermillion, South Dakota
17   4) Electron Microscopy, Max-Planck-Institute for Biophysical Chemistry, Goettingen,
18   Germany
19   5) Department of Genes and Behavior, MPI for Biophysical Chemistry, Goettingen, Germany
20   6) Institute of Molecular Oncology, University Medical Center Goettingen, Germany
21   7) Department of Biomedical Sciences, New York Institute of Technology College of
22   Osteopathic Medicine, Old Westbury, New York, USA
23   8) Institute of Pathology, University Medical Center Goettingen, Goettingen, Germany
24   9) Microarray and Deep-Sequencing Core Facility, University Medical Center Goettingen,
25   Germany

Corresponding authors

Muriel Lizé, University Medical Center Goettingen, Clinic for Cardiology and Pneumology, section

Molecular & Experimental Pneumology, email: mlize@gwdg.de

Haotian Zhao, Department of Biomedical Sciences, New York Institute of Technology College of

Osteopathic Medicine, Riland Building Room 027, Old Westbury, New York 11568-8000.

Telephone: (516) 686-1271; Fax: (516) 686-1454; email: hzhao10@nyit.edu

Abstract

Motile cilia serve vital functions in development, homeostasis and regeneration. We recently demonstrated that TAp73 is an essential transcriptional regulator of respiratory multiciliogenesis. Here, we show that TAp73 is expressed in multiciliated cells (MCCs) of diverse tissues. Analysis of *TAp73* mutant animals revealed that TAp73 regulates *Foxj1*, *Rfx2, Rfx3*, axonemal dyneins *Dnali1* and *Dnai1*, plays a pivotal role in the generation of MCCs in male and female reproductive ducts, and contributes to fertility. However, the function of MCCs in the brain appears to be preserved despite the loss of *TAp73*, and robust activity of cilia-related networks is maintained in the absence of *TAp73*. Notably, *TAp73* loss leads to distinct changes in ciliogenic microRNAs: *miR34bc* expression is reduced, whereas the *miR449* cluster is induced in diverse multiciliated epithelia. Among different MCCs, choroid plexus (CP) epithelial cells in the brain display prominent *miR449* expression, whereas brain ventricles exhibit significant increase in *miR449* levels along with an increase in the activity of ciliogenic E2F4/MCIDAS circuit in *TAp73* mutant animals. Conversely, E2F4 induces robust transcriptional response from *miR449* genomic regions. To address whether increased *miR449* levels in the brain maintain the multiciliogenesis program in the absence of *TAp73*, we deleted both *TAp73* and *miR449* in mice. Although loss of *miR449* alone led to a mild ciliary defect in the CP, more pronounced ciliary defects and hydrocephalus were observed in the brain lacking both *TAp73* and *miR449*. In contrast, *miR449* loss in other MCCs failed to enhance ciliary defects associated with *TAp73* loss. Together, our study shows that, in addition to the airways, TAp73 is essential for generation of MCCs in male and female reproductive ducts, whereas *miR449* and TAp73 complement each other to support multiciliogenesis and CP development in the brain.

Introduction

Cilia are hair-like appendages protruding from the cell membrane into the surrounding environment. Solitary immotile primary cilia are a common organelle in most mammalian cells, whereas motile cilia are restricted to a subset of cell types. This subset includes multiciliated cells (MCCs) lining brain ventricles, tracheal, and bronchial epithelium as well as the epithelium of male efferent ducts (EDs) and fallopian tubes (FTs) in females [1].

Multiciliogenesis requires precise regulation of the production, transport and assembly of a large number of different structural components, a process critically dependent on a hierarchical network of transcriptional and post-transcriptional regulators [2]. Geminin Coiled-Coil Domain Containing 1 (GEMC) [3–5] and multiciliate differentiation and DNA synthesis associated cell cycle protein (MCIDAS or Multicilin) [6–8], members of the Geminin family, are early regulators of the MCC fate, downstream of the NOTCH pathway. MCC differentiation is also regulated by post-transcriptional mechanisms including microRNAs (miRNAs). *miR-34/449* constitute a conserved family that encodes six homologous miRNAs (*miR34a, 34b, 34c, 449a, 449b,* and *449c*) from three genomic loci in vertebrates. Inhibition of the NOTCH pathway e.g. by *miR449* is required for multiciliogenesis through de-repression of the transcriptional network of *GEMC1*, *MCIDAS*, E2F transcription factors (*E2F4*, *E2F5*), forkhead box J1 (*FOXJ1*), and v-myb avian myeloblastosis viral oncogene homolog (*MYB)* [9–11]. Disturbance of the molecular circuit leads to defective multiciliogenesis and ciliopathies in the airways, reproductive tracts, and the brain [1].

Transformation related protein 73 (*Trp73*) is a member of the p53 family with distinct isoforms generated from two alternative promoters: isoforms containing the N-terminal transactivation domain (TAp73), and N-terminally truncated dominant-negative isoforms (ΔNp73). Recently, we and others showed that TAp73 is essential for airway multiciliogenesis [12,13]. Gene expression analysis and chromatin immunoprecipitation (ChIP) identified TAp73 as a critical regulator of multiciliogenesis: TAp73 acts downstream of E2F4/MCIDAS and regulates the expression of *FOXJ1*, *RFX2,* and *RFX3* in pulmonary tissues [12,14–17].

The FT of female reproductive tract consists of MCCs that possess hundreds of motile cilia beating

in a wave-like manner which, along with musculature contraction, moves the oocyte or zygote

towards the uterus [18–20]. Defects in ciliary functions may lead to ectopic pregnancies or infertility

[19,21]. In the male reproductive tract, MCCs in the EDs are involved in the transport of

spermatozoa from testis to epididymis (Epi), their maturation and concentration [22–25].

MCCs in the brain can be found in a single layer of ependymal cells facing the ventricles and

choroid plexus (CP). The CP epithelium, a specialized secretory epithelium that secretes

cerebrospinal fluid, arises from monociliated progenitors in the roof plate around embryonic day

(E) 12 and undergoes multi-ciliate differentiation to form multiple primary cilia [26,27]. Ependymal

cells in mice are specified around day E16 and form multiple motile cilia on the apical surface after

birth to facilitate cerebrospinal fluid movement [28,29]. Defects in the ependymal and CP lineages

are implicated in aging, hydrocephalus, and brain tumors [30,31].

In this study, we detected robust TAp73 expression in MCCs in diverse tissues. In reproductive

ducts, *TAp73* loss leads to a profound reduction of multiciliogenesis and suppression of *TAp73*-

dependent transcriptional network activity. However, MCCs in the brain maintain robust

multiciliogenesis activity despite *TAp73* loss. Molecular studies revealed alterations in *miR-34/449*

family members in diverse MCCs from *TAp73* mutant mice: decreased levels of TAp73 target

miR34bc concurrent with increased expression of *miR449*. In the brain, *miR449* is highly

expressed in the CP and experiences significant upregulation following *TAp73* deletion. In

addition, brain ventricles but no other multiciliated tissues from *TAp73* mutant animals exhibit

increased expression of E2F4, which in turn is capable of eliciting robust transcriptional response

from *miR449* genomic loci, suggesting that *miR449* plays a crucial role in brain multiciliogenesis

in collaboration with *TAp73*. Indeed, *miR449* loss alone results in ciliary reduction in the CP,

whereas loss of both *TAp73* and *miR449* leads to a dramatic reduction of multiciliogenesis in the

CP and severe hydrocephalus. Therefore, the molecular network governing MCC fate is subjected

to tissue-specific feedback modulation by transcriptional and post-transcriptional mechanisms.

Materials and Methods

Animals

TAp73 mutant mice with a targeted deletion of exons 2 and 3 of the *Trp73* gene were a generous

gift from Dr. Tak Mak (Princess Margaret Cancer Centre, *Toronto, Canada*) [32]. *miR449* mutants

were previously described [33]. Both strains were maintained in C57Bl/6 background (n8) at the

animal facility of the European Neuroscience Institute Goettingen, Germany in full compliance with

institutional guidelines. The study was approved by the Animal Care Committee of the University

Medical Centre Goettingen and the authorities of Lower-Saxony under the number 16/2069.

Human samples

Human epididymis samples were procured with informed consent from two patients (42 and 41

years of age, respectively). All experimental procedures were approved and performed in

accordance with the requirements set forth by Ethics Committee of the University Medical Centre

Goettingen (application number: 18/2/16).

Histology and immunostaining

Paraformaldehyde-fixed, paraffin-embedded tissues were treated with heat-induced epitope

retrieval using Rodent Decloaker (RD913 L, Biocare Medical, *Pacheco, CA, USA*). For

immunohistochemistry, endogenous peroxidase activity was quenched with 3% H_2O_2 for 10 min.

Tissue sections were blocked with 10% fetal calf serum (FCS) in phosphate-buffered saline (PBS)

with 0.1% Triton X-100, and subsequently incubated with primary antibodies (List of antibodies is

provided in **Supplementary Table 1**). Biotinylated secondary antibodies were applied for 1 h at

room temperature (List of antibodies is provided in **Supplementary Table 2**), after which avidin

enzyme complex and substrate/chromogen were used for color development (Vector laboratories,

Burlingame, CA, USA). Stained tissue sections were counterstained with hematoxylin. For

immunofluorescence, sections were stained with fluorescently labeled secondary antibodies (List

of antibodies is provided in **Supplementary Table 2**) for 1 h at room temperature. Nuclei were

counterstained with 4`, 6-Diamidin-2-phenylindol (DAPI). Histology of tissue sections was

assessed by using hematoxylin (Merck, *Darmstadt, Germany*) and eosin (Carl Roth, *Karlsruhe,*

Germany) staining.

Electron microscopy

Transmission electron microscopy (TEM) was performed as previously described [12]. Briefly,

murine tissue samples were fixed by immersion using 2% glutaraldehyde in 0.1 M cacodylate

buffer (Science Services, *München, Germany*) at pH 7.4 overnight at 4°C. Post-fixation was

performed using 1% osmium tetroxide diluted in 0.1 M cacodylate buffer. After pre-embedding

staining with 1% uranyl acetate, tissue samples were dehydrated and embedded in Agar 100

(Plano, *Wetzler, Germany*). Thin tissue sections (100 nm) were examined using a Philips CM 120

BioTwin transmission electron microscope (Philips Inc., *Eindhoven, The Netherlands*) and images

were taken with a TemCam F416 CMOS camera (TVIPS, *Gauting, Germany*).

Quantification of cilia markers

145 Cilia were quantified using the *ImageJ* software [34]. Briefly, the region of interest was selected

146 and a threshold was set to exclude unspecific background signals. The *Analyze Particles* tool was

147 used to measure the area of the ciliary staining. Values were normalized to the length of the

148 epithelia measured.

Western blot

Samples were homogenized in RIPA buffer (20 mM TrisHCl pH 7.5, 150 mM NaCl, 9.5 mM EDTA,

1% Triton X100, 0.1% SDS, 1% sodium deoxycholate) supplemented with urea (2.7 M) and

protease inhibitors (Complete Mini EDTA-free, Roche, *Basel, Switzerland*). Equal amounts of

protein extracts were separated by SDS-polyacrylamide gels prior to transfer onto a nitrocellulose

membrane and incubated with primary antibodies (List of antibodies is provided in

Supplementary Table 1). The membrane was washed and incubated for 1 h with horse radish

peroxidase (HRP)-conjugated secondary antibodies (List of antibodies is provided in

Supplementary Table 2) followed by chemiluminescence detection. β-ACTIN or heat shock

cognate 71 kDa protein (HSC70) were used as protein loading controls.

RNA extraction, quantitative PCR, small RNA sequencing, and RNAscope

Tissue samples were snap-frozen in liquid nitrogen and total RNA was isolated by Extrazol (7BioScience, *Hartheim, Germany*)/Chloroform extraction followed by 80% ethanol precipitation at -20°C. For cDNA synthesis, 1 μg of total RNA was incubated with the M-MuLV reverse transcriptase and a mix of random nonameric and polyA tail primers at 42°C for 1 h in a total volume of 50 μl. All reactions were set up in triplicate with self-made SYBR Green quantitative PCR (qPCR) Mix (Tris-HCl [75 mM], $(NH_4)_2SO_4$ [20 mM], Tween-20 [0.01% v/v], $MgCl_2$ [3 mM], Triton X-100 [0.25% v/v], SYBR Green I (1:40 000), dNTPs [0.2 mM] and Taq-polymerase [20 U/ml]) using 250 nM of each gene-specific primer (List of primers is provided in **Supplementary Table 3**). Standard curve method was used to assess relative transcript content. Transcript of interests were normalized to the reference transcript of ribosomal phosphoprotein P0 (Rplp0, or *36b4*) and normalized to the mean value of control samples. The results for each sample were obtained by averaging transcript levels of technical triplicates. No RT controls and dilution curves as well as melting curves and gel electrophoresis assessment of amplicons were performed for all primer combinations. For *miR449a*, *miR34b*, and *miR34c* quantification, TaqMan MicroRNA Assay (Applied Biosystems, Thermo Fisher Scientific, *Waltham, MA, USA*) was performed according to the manufacturer's instructions with U6 snRNA as internal control.

Copy number in RNA samples was determined by qPCR using a murine TAp73 plasmid (MC219984, Origene, *Rockville, USA*) with a known copy number as standard curve. Copy number of the TAp73 plasmid was determined using the following formula: number of copies = (plasmid amount [ng] * $6.022x10^{23}$ [molecules/mole]) / (plasmid length [bp] * $1x10^9$ [ng/g] * 650 [g/mol])

The libraries for small RNA samples were prepared using TruSeq Small RNA Library Prep Kit - Set A (24 rxns) (Set A: indexes 1-12; Cat N°: RS-200-001, Illumina, *San Diego, CA, USA*) using 1 μg of total RNA according to manufacturer's recommendations. Samples were sequenced on the Illumina HiSeq 4000 using a 50 bp single-end approach. Mapping, prediction of novel miRNAs,

quality control, and differential expression (DE) analysis were carried out using Oasis2.0 (*Oasis:*

online analysis of small RNA deep sequencing data) [35]. In brief, FASTQ files were trimmed with

cutadapt 1.7.1 [36] removing Truseq adapter sequences (TGGAATTCTCGGGTGCCAAGG)

followed by removing sequences smaller than 15 or larger than 32 nucleotides. Trimmed FASTQ

sequences were aligned to mouse small RNAs using STAR version 2.4.1d [37] with a mismatch

of 5% of the sequence length and by utilizing the following databases: Mirbase version 21 for

miRNAs; piRNAbank V.2 for piwiRNA; and Ensembl v84 for small nuclear RNA, small nucleolar

RNA, and ribosomal RNA. Counts per small RNA were calculated using featureCounts v1.4.6 [38].

Novel miRNAs were searched for using miRDeep2 version 2.0.0.5 [39]. Differential expression of

small RNA was determined by DESeq2 [40], where small RNAs were considered differentially

expressed with an adjusted p-value <0.05 and absolute log2 fold-change >1. The results of the

DE analysis can be found in **Supplementary Table S6**, and the small RNA-seq data sets can be

found in Gene Expression Omnibus (GEO) with accession number **GSE108385**.

TAp73 (probe no. 475741), *Mcidas* (probe no. 510401-C2), *Hes1* (probe no. 417701), and *Hes5*

(probe no. 400991-C2) were visualized using RNAscope 2.5 HD Duplex Reagent Kit (#322430,

Advanced Cell Diagnostics, *Hayward, CA, USA*) according to manufacturer's instructions.

Chromatin immuno-precipitation (ChIP)

Chromatin was harvested from Saos2 cells transiently overexpressing TAp73α, TAp73β, and the

control vector pcDNA3.1. Saos2 cells were routinely tested negative for Mycoplasma. ChIP and

qPCR was performed as previously described using gene specific primers (sequence information

is provided in **Supplementary Table 4**) [12]. Enrichment levels were determined as the number

of PCR products for each gene relative to total input.

Luciferase assay

Luciferase assay was performed as previously described [12]. Briefly, Saos2 cells were

transfected with pcDNA3.1 empty vector, or pcDNA3.1 vector carrying *E2F4* or *MCIDAS*, or both

E2F4 and *MCIDAS* vectors. Moreover, a firefly luciferase reporter construct containing the putative

three wild type E2F-binding sequences of *miR449* genomic region (wild type, or "WT"), or the

same sequences lacking the strongest predicted E2F-binding motif (mutant, or "Mut") were

transfected (sequence information is provided **Supplementary Table 5**). In addition, a Renilla TK

luciferase vector was co-transfected. At 24 h after transfection, cells were harvested and the

luciferase activities were measured using the dual luciferase assay. Firefly luciferase activities

were determined relative to those of Renilla TK luciferase vector and normalized to the mean

value of samples from the control vector. Luciferase assays were performed as technical triplicates

on every biological replicate.

Video microscopy

Murine fallopian tube and testis connected to the epididymis were dissected and transferred to

Dulbecco's Modified Eagle's Medium (DMEM, Gibco, Thermo Fisher Scientific, *Waltham, MA,*

USA). To image spermatozoa, the epididymis was separated from testis and vas deferens and an

incision was made at distal end to release the spermatozoa. Spermatozoa as well as the peristaltic

contraction of the fallopian tube were imaged with an inverse microscope.

Imaging of cilia-generated bead-flow and cilia beating in the brain ventricular system

Mouse brains were dissected and transferred to DMEM 21063 (Gibco, Thermo Fisher Scientific).

Coronal slices containing the lateral ventricle, ventral third ventricle, aqueduct, and fourth ventricle

were prepared by using a coronal adult brain matrix (ASI Instruments, *Warren, MI, USA*). The

ventral third ventricle was processed further as previously described [41]. Tissue explant was

placed in DMEM containing fluorescent latex beads (Fluoresbrite Multifluorescent 1.0 micron

Microspheres, Polysciences, *Warrington, PA, USA*). Movement of fluorescent beads along the

ventricular wall and within ventricular lumen was observed by fluorescence microscopy using a

DMR (Leica, *Wetzlar, Germany*) upright microscope with an epifluorescence lamp. Ciliary beating

was observed by differential interference contrast microscopy using the same set-up. Bead

movement was recorded using a high-speed camera (Cascade II-512, Photometrics, *Tucson, AZ,*

USA) operated by MultiRecorder Software (developed by Johannes Schröder_Schetlig) and

analyzed using ImageJ software [34].

Statistical Analysis

One-tailed, unpaired Student's *t*-test assuming normal distribution and equal variances was used

to calculate statistical significance for pairwise comparisons. Luciferase assay statistics were

assessed using one-way ANOVA assuming normal distribution followed by Dunnett's multiple

comparison tests. The following indications of significance were used: *$*P<0.05$, $**P <0.01$, $***P$

<0.001. N values represent biological replicates. Error bars indicate standard error of the mean

(SEM).

Results

TAp73 is expressed in diverse multiciliated epithelia

We and others previously showed that TAp73 expressed in respiratory epithelia controls

multiciliogenesis [12,13]. However, little is known about the expression and function of TAp73 in

other MCCs. To address this, we performed immunostaining and *in situ* hybridization and

demonstrated that in addition to the testis [42,43], TAp73 is expressed in EDs, FTs, and

ependymal and CP epithelial cells in the brain (**Fig. 1a-f; Supplementary Fig. 1**). qPCR and

western blot analyses showed that, among different multiciliated epithelia, FTs and EDs exhibit

higher levels of TAp73 expression than testis or brain (**Fig. 1g-i**). Taken together, these results

demonstrate robust TAp73 expression in different MCC types.

TAp73 **is crucial for the molecular circuit of multiciliogenesis in efferent ducts**

Loss of *TAp73* leads to male infertility that has been attributed to defective germ cell maintenance

during spermatogenesis [42,43]. Interestingly, we detected spermatozoa in testis from *TAp73* KO

mice, although at a markedly reduced levels (**Supplementary Fig. 2a, b**). Despite normal

morphology and mobility of these cells, no mature spermatozoa were detected in the epididymis

(Epi) of these mice (**Fig. 2a; Supplementary Video 1a-d**), suggesting that additional defects may

contribute to infertility. The multiciliated epithelium of the EDs contributes to gamete transport by facilitating testicular fluid circulation, fluid reabsorption, and spermatozoa concentration [22,24,25], all essential aspects of male fertility [9,44,45]. Indeed, though no gross morphological difference was observed in EDs between control and *TAp73* KO animals (**Fig. 2a**), immunofluorescent staining of the cilia components acetylated alpha-tubulin (Ac-α-TUB) and dynein axonemal intermediate chain 1 (DNAI1) showed a dramatic reduction in the number and length of cilia in the EDs from *TAp73* KO mice (**Fig. 2b, c)**. In contrast to the abundant long cilia of WT cells, mutant MCCs generated far fewer cilia as observed by transmission electron microscopy (TEM) (**Fig. 2d; Supplementary Fig. 2c**), resembling the loss of airway cilia in these animals [12]. Consistent with its role as transcriptional regulator, ChIP followed by qPCR revealed significant enrichment of TAp73 in genomic loci of *FOXJ1* [12] and dynein axonemal light intermediate chain 1 (*DNALI1)* and *DNAI1,* both encoding axonemal dyneins (**Fig. 2e; Supplementary Fig. 3**). Accordingly, expression of *Dnali1*, *Foxj1*, *Rfx2,* and *Rfx3* was reduced or almost completely lost in male reproductive ducts from *TAp73* KO animals (**Fig. 2f, g; Supplementary Fig. 2d**). Together, our data indicate that TAp73 directs *Dnali1* and *Dnai1* in addition to known critical nodes including *Foxj1*, *Rfx2,* and *Rfx3* to mediate multiciliogenesis in EDs (**Fig. 7a, b**). Thus, these additional defects in the multiciliated epithelium of the EDs may contribute to male infertility in *TAp73* KO mice.

TAp73-driven transcriptional network regulates multiciliogenesis in fallopian tubes

Though infertility in *TAp73* KO females is thought to arise from defects of oocyte development and release from the ovary [32,46], it remains unclear whether *TAp73* loss affects the multiciliated epithelium of the FT, thereby possibly influencing ova transport. Despite normal tubal morphology, analysis of Ac-α-TUB and DNAI1 expression showed reduced cilia coverage of the oviduct epithelium (**Fig. 3a-c)**. Consistently, TEM demonstrated reduced cilia and mislocated basal bodies in FTs from *TAp73* KO mice (**Fig. 3d; Supplementary Fig. 4a**). Transcript levels of *Dnali1*, *Foxj1*, and *Rfx2,* but not *Rfx3* were reduced in *TAp73* KO FTs (**Fig. 3e**), which was accompanied by declined protein expression of FOXJ1, DNAI1, DNALI1 (all expressed in the human FTs, **Supplementary Fig. 4b**), and gamma-tubulin (γ-TUB, basal body marker) (**Fig. 3f**), though to a lesser degree when compared to the decrease in multiciliogenesis activity observed in *TAp73*-deficient EDs. Further, smooth muscle contraction pattern in FTs is similar between control and *TAp73* KO animals (**Supplementary Video 2a, b**). Taken together, our data indicate that *TAp73* loss leads to reduced multiciliogenesis in the oviducts (**Fig. 7a, c**).

Ciliary function in the brain is intact in the absence of *TAp73*

Given TAp73 expression in ependymal and CP epithelial cells, we further evaluated TAp73 expression during embryonic brain development. Immunofluorescent studies showed that proliferative progenitors (KI-67$^+$) are present in hindbrain roof plate at day E14.5, whereas post-mitotic cells expressing aquaporin 1 (AQP1) [31,47] are detected in CP epithelium (KI-67$^-$/AQP1$^+$) (**Fig. 4a**). Notably, a portion of the roof plate exists between the progenitors and CP epithelium that remains undifferentiated after cell cycle exit (KI-67$^-$/AQP1$^-$) (**Fig. 4a**). In contrast to progenitors with a solitary primary cilium, the "transition" zone is comprised of MCCs that exhibit TAp73 expression (**Fig. 4b**).

The expression of TAp73 in ependymal and CP epithelial cells, along with recent studies demonstrating the role of E2F4/MCIDAS in multiciliogenesis of ependymal cells [7,48,49], led us

to examine the role of TAp73 in MCCs in the brain. Immunostainings confirmed the loss of TAp73

expression in ependymal cells and the CP from *TAp73* KO mice (**Fig. 4c**), whereas morphological

analysis revealed no apparent defect in these cells (**Supplementary Fig. 5a**). We performed

immunostainings for the cilia markers ADP-ribosylation factor-like 13b (ARL13B) [50], Ac-α-TUB,

and DNAI1 in the 4th and lateral ventricles. In contrast to FTs and EDs, MCCs in ependyma and

CP from *TAp73* KO animals are similar to those of WT mice (**Fig. 4d, e; Supplementary Fig. 5b-**

d). No significant difference was observed in the expression of markers for epithelial differentiation

of CP between control and *TAp73* KO animals (**Supplementary Fig. 6a-d**). RT-qPCR analysis

demonstrated similar expression levels of *Dnali1* and *Foxj1,* whereas increased *Rfx2* and *Rfx3*

mRNA levels were observed in brain ventricles from *TAp73* KO mice (**Fig. 4f**). Consistently, ciliary

beating and bead flow in the cerebrospinal fluid appeared unaffected by *TAp73* loss (**Fig. 4g;**

Supplementary Video 3a, b). Taken together, these results indicate that, unlike EDs, FTs, and

the airways [12], the differentiation and function of MCCs in the brain remain intact despite *TAp73*

loss.

TAp73 regulates *miR-34/449* family members in diverse MCCs

Functional MCCs in the brain from *TAp73* KO mice suggest that other ciliogenic factors may

rescue brain multiciliogenesis in the absence of *TAp73*. TAp73 influences post-transcriptional

mechanisms *via* regulation of miRNAs [12]. Analysis of small RNA species from brain ventricles

in *TAp73* KO mice revealed reduced *miR34bc* levels, along with a strong induction of the *miR449*

cluster that works together with *miR34bc* to regulate multiciliogenesis in different tissues across

species (**Fig. 5a, c; Supplementary Table 6**) [11,33,51–53]. In the brain, *miR449* is

predominantly detected in the CP [54], where its expression undergoes >10 fold increase upon

TAp73 loss, whereas *miR34bc* levels strongly decline (**Fig. 5b, c**). Although *miR34b* levels were

down-regulated as well in trachea from *TAp73* KO (**Supplementary Fig. 7a)**, *miR449* induction

was less pronounced and more variable in FTs and EDs (**Fig. 5d**). Altogether, these results reveal

a conserved reaction from the *miR-34/449* family following *TAp73* loss in diverse multiciliated

epithelia.

In an effort to understand *miR449* upregulation we analyzed potential changes in the RB-E2F

pathway known to regulate *miR449* levels [55,56]. However, expression of *E2f1*, *E2f3*, *Cdkn1a,*

and *Cdkn1b* in brain ventricles were comparable between WT and mutant animals,

(**Supplementary Fig. 8a, b**), indicative of a RB-E2F pathway unaffected by *TAp73* loss in brain

MCCs. Interestingly, transcript and protein levels of the other E2F family member E2F4, which is

a potent inducer of multiciliogenesis [6–8,48,57,58], were markedly increased in *TAp73* KO

ventricles, despite only a mild increase of its cofactor *Mcidas* (**Fig. 5e, f**). In contrast, E2F4 levels

in FTs and EDs were unaltered and even downregulated in tracheae (**Supplementary Fig. 7b-e**).

Therefore, increased E2F4 levels concurrent with a *miR449* increase are restricted to the brain in

TAp73 KO mice.

To assess potential E2F4 contribution to *miR449* elevation, we used the genomic region of

miR449 containing three putative E2F binding sites in a reporter-based assay. Indeed, E2F4 in

combination with MCIDAS elicited a strong transcriptional response from the *miR449* locus, a

reaction almost abolished by mutating the strongest out of three E2F consensus motifs (**Fig. 5g;**

Supplementary Table 5). Together, these results indicate that increased E2F4/MCIDAS activity

may stimulate *miR449* expression in *TAp73* KO brains.

TAp73 collaborates with *miR449* in brain multiciliogenesis

Our data suggest that *miR449* upregulation may compensate at least partially for *TAp73* loss to

maintain brain multiciliogenesis. To address this, we generated mice with a deletion of the *miR449*

cluster in addition to *TAp73*. Strikingly, *TAp73$^{-/-}$;miR449$^{-/-}$* (*TAp73xmiR449* KO) mice developed

severe hydrocephalus (**Fig. 6a; Supplementary Fig. 9a**). Since defective ependymal and CP cilia

contribute to the development of hydrocephalus [59–61], we next assessed ciliation in the

ventricles of *TAp73xmiR449* KO mice. Analysis of the expression of ARL13B in CP epithelium

revealed a decrease in cilia number and length in the absence of *miR449*, whereas a more

pronounced reduction in cilia was observed in *TAp73xmiR449* KO mice (**Fig. 6b, c;**

Supplementary Fig. 9b). TEM studies also revealed mildly disorganized apical docking of basal

bodies in ependymal cells in *TAp73* KO and *TAp73xmiR449* KO mice (**Fig. 6d; Supplementary**

Fig. 9c); however, Ac-α-TUB content was similar in ependymal cells among WT and

TAp73xmiR449 KO animals (**Supplementary Fig. 9d**). Consistently, ciliary beating and bead flow

over ventricles appeared unaffected in *TAp73xmiR449* KO animals (**Supplementary Fig. 9e;**

Supplementary Video 3a, c). Furthermore, expression of cytokeratins, AQP1, and OTX2 in CP

epithelial cells was similar among WT, *miR449* KO, and *TAp73xmiR449* KO animals

(**Supplementary Fig. 10a-c**). Despite the role of Notch signaling in CP development and

tumorigenesis [31,62], RNAscope studies revealed similar expression of NOTCH targets *Hes1*

and *Hes5* in the roof plate of WT, *miR449* KO, and *TAp73xmiR449* KO embryos at day E14.5

(**Supplementary Fig. 11)**. In summary, additional loss of *miR449* in *TAp73* KO mice strongly

impairs ciliogenesis in the CP, but only slightly affects ependymal cilia, which is consistent with its

prominent expression in the CP [54] (**Fig. 5b**). Thus, our data indicate that *miR449* collaborates

with *TAp73* to drive multiciliogenesis in the brain.

As *miR449* was induced upon *TAp73* deletion in further multiciliated tissues, we analyzed tracheae

and EDs in *TAp73xmiR449* KO mice. Immunostainings and TEM consistently revealed a dramatic

decrease in cilia coverage and an increase in defective basal body docking in trachea from

TAp73xmiR449 KO animals compared to WT animals (**Fig. 6e-g; Supplementary Fig. 12a**), a

phenotype bearing resemblance to our previous findings in the airways of *TAp73* KO animals [12].

Likewise, loss of *miR449* did not further enhance MCC reduction in *TAp73*-deficient EDs

(**Supplementary Fig. 12b, c**). Thus, additional deletion of the *miR449* cluster fails to exacerbate

ciliary defects in trachea and EDs in the absence of *TAp73*.

Overall, our data indicate that TAp73 utilizes the unique topology of its transcriptional circuit to

communicate with the *miR-34/449* family and other crucial regulators of motile multiciliogenesis

e.g. *E2F4/MCIDAS* to regulate brain multiciliogenesis (**Fig. 7a, d, e**).

Discussion

TAp73 activates a plethora of ciliogenic effectors to drive multiciliogenesis in the airways [12,13].

The current study examines the role of TAp73-driven molecular circuit in MCCs of reproductive

tracts and the brain. Our results revealed a profound reduction of cilia in EDs and FTs from *TAp73*-

deficient mice, as well as diminished *Foxj1*, *Rfx2, and Rfx3* expression. These molecular and

cellular changes in MCCs are reminiscent of our previous findings in respiratory epithelia of these

mice, suggesting that male and female infertility associated with *TAp73* loss could be in part

related to the observed cilia loss. The expression of the axonemal dyneins *Dnai1* and *Dnali1*, both

of which exhibit TAp73 binding in their genomic loci, was also significantly reduced in EDs and

FTs from mutant animals, indicating that they are part of the TAp73-directed multiciliogenesis

program in reproductive tracts.

Consistent with previous reports, we found partial degradation of the germinal epithelium and

reduced sperm cell production in *TAp73*KO mice [42,43]. The EDs are comprised of MCCs, which

are required for fluid circulation and reabsorption, thereby facilitating the transport of spermatozoa

to their storage and maturation in the epididymis [22,24,25]. Despite the presence of flagellated

spermatozoa in testis, lack of spermatozoa in epididymis of *TAp73* KO mice indicates that

defective multiciliogenesis may contribute to male sterility. Indeed, disruption of transcriptional

regulators of multiciliogenesis has been shown to cause infertility in mice and humans [3,63],

whereas fertility issues have been reported in female primary ciliary dyskinesia patients [20,21].

Importantly, *TAp73* is downregulated as women age [64], and certain single nucleotide

polymorphisms in *TP73* are associated with female patients over 35 years of age seeking *in vitro*

fertilization [65,66]. Hence, the integrity of MCCs is critical for reproductive health. Further studies

using tissue-specific deletion of *TAp73* in MCCs of EDs and oviducts are necessary to delineate

its role in reproductive motile cilia maintenance and fertility.

In the brain, *TAp73* expression is initiated at the onset of multiciliated differentiation of ependymal

and CP epithelial cells. However, our data indicate that *TAp73* is dispensable for the generation

of cilia in the brain, although it is plausible that *TAp73* loss results in more subtle defects such as

polarity and cilia orientation [67,68]. In contrast to the dynamic TAp73-dependent program in the

airways and reproductive tracts, expression of *Foxj1, Rfx2*, *and Rfx3* in the brain remains mostly

unaltered in the absence of *TAp73*, suggesting that other effectors maintain the activity of the

molecular circuit to support MCC differentiation.

Previous studies revealed robust expression of *GemC1* and *E2f/Mcidas*, all of which are capable

of transcriptional activation of *Foxj1*, *TAp73* itself, and many other ciliogenic effectors e.g. *Rfx2*

and *Rfx3* in MCCs of the brain [4,6,8,48,69]. Indeed, E2F4/MCIDAS expression is upregulated in

the brain but not in other multiciliated tissues upon *TAp73* loss, and therefore may facilitate brain

multiciliogenesis. In agreement, loss of either *Mcidas* or *GemC1*, both transcriptional activators of

TAp73, leads to defect in MCC differentiation and hydrocephalus [3,6].

Although it is less clear how *TAp73* loss results in enhanced E2F/MCIDAS activity in the brain, a

quick look downstream of TAp73 provides some clues: reduced expression of the TAp73 target

miR34bc is concurrent with an induction of *miR449* in the absence of *TAp73*. Interestingly,

expression of Cdkn1a/p21, Cdkn1b/p27, E2f1, and E2f3 in brain ventricles remain unchanged

following *TAp73* loss, suggesting that *TAp73* loss regulates E2F and *miR449* activity

independently of the conserved RB-E2F1 axis. *miR449* induction is commonly observed in *miR34-*

deficient MCCs, whereas ablation of the entire *miR-34/449* family severely impairs

multiciliogenesis in diverse tissues [33,70]. *miR449* is known to inhibit the NOTCH pathway to

relieve the suppression of MCC fate determination; however, NOTCH pathway activity in the CP

remains unchanged after *miR449* loss. Given the diverse targets of the *miR-34/449* family, it is

plausible that *miR449* may indirectly increase E2F/MCIDAS activity in MCCs of the brain

independent of NOTCH inhibition. Conversely, transcriptional activation of *miR449* by

E2F/MCIDAS complexes may complete the feedback loop to keep the molecular circuit fully

engaged in the absence of *TAp73*.

This interpretation posits that the crosstalk between *miR449* and E2F/MCIDAS serves as a crucial

backup circuit for TAp73-driven multiciliogenesis network in the brain. Indeed, combined deletion

of *TAp73* and *miR449* results in disruption of multiciliogenesis in the brain and hydrocephalus,

defects distinct from those associated with complete loss of the *miR-34/449* family [33,71]. In

TAp73-deficient MCCs outside the brain that exhibit less prominent increase in *miR449* and no

increase in E2F4 levels, further deletion of *miR449* cluster fails to exacerbate multiciliogenesis

defects caused by *TAp73* loss, indicating that TAp73 functions at least partially through *miR449*

to support MCCs in the brain. Recent studies also demonstrated the role of *TAp73*-driven *miR34a*

expression in neuronal development [72]. Therefore, interaction of TAp73 with *miR-34/449* family

members is crucial for normal brain functions.

Nonetheless, detailed studies are necessary to clarify the interaction between *miR449* and

E2F4/MCIDAS pathway in MCCs in the brain, but also to address *miR449* regulation in *TAp73*-

deficient MCCs outside the brain.

Unlike *TAp73* mutant animals, *p73* KO mice lacking both *TAp73* and *ΔNp73* exhibit

hydrocephalus, defective ependymal cell maturation and aqueduct stenosis, suggesting a

potential role for *ΔNp73* in ependymal cells [73,74]. Given the abnormal apical localization of basal

bodies in ependymal cells along with ciliary defects in the CP from *TAp73xmiR449* KO mice, it is

conceivable that *ΔNp73* may regulate *miR449* expression indirectly in these cells. In support of

this notion, *miR449* is highly expressed in the CP whereby its loss alone leads to ciliary defects,

whereas *ΔNp73* deletion also results in defects in the CP [75]. Further analysis of the

multiciliogenesis network and *miR449* expression in MCCs in the brains of *p73* KO and *ΔNp73*

KO animals are necessary to resolve these questions.

Acknowledgments

We thank Tak Mak for providing *TAp73* KO mice, Gerd Hasenfuß for support, Matthias Dobbelstein for hosting and Karola Metze and Verena Siol for assistance. M.L. is supported by Deutsche Forschungsgemeinschaft (DFG Li 2405); H.Z. by New York Institute of Technology, Sanford Research, Matthew Larson Foundation, Institutional Development Award from the National Institute of General Medical Sciences under grant numbers 5P20GM103548, 1P20GM103620-01A1, and National Cancer Institute (R01CA220551); F.B. by Wilhelm-Sander-Stiftung (2016.041.1); A.K.G by the Max Planck Society. We thank Heymut Omran's group for introduction to cilia microscopy and Travis Stracker for disclosure of non-published data.

Author Contributions

Me.W. and T.E. characterized cilia defects and gene expression and generated figures. Me.W. and Ma.W. validated TAp73 targets by WB and ChIP. E.E. and F.B. contributed IF analysis of human epididymis. E.E. performed cilia quantification on tracheae. D.R. performed electron microscopy analysis. C.W. maintained mice, performed RNA isolation and qPCRs. L.V-H. and S.B. contributed to Western blot analysis of different tissues. Z. H. performed RNAscope analysis for *TAp73* on diverse tissues. K.B.G., J.Z., L.L., and H.Z. contributed brain analyses. A-K.G analyzed *ex vivo* ciliary beating. O.S. analyzed small RNA sequencing data. S.A. contributed to interpretation and supported the group. M.L. developed the project, interpreted the data, designed and coordinated the experiments to complete this study. Me.W., T.E., H.Z., and M.L. were major contributors to manuscript preparation.

Conflict of Interest

The authors declare that they have no conflict of interest.

483   **Electronic supplementary material.** The online version of this article contains supplementary

484   material, which is available to authorized users.

485   **References**

486   1.   Spassky N, Meunier A. The development and functions of multiciliated epithelia. Nat. Rev.
487   Mol. Cell Biol. 2017;18:423–36.

488   2.   Choksi SP, Lauter G, Swoboda P, Roy S. Switching on cilia: transcriptional networks
489   regulating ciliogenesis. Dev. Camb. Engl. 2014;141:1427–41.

490   3.   Terré B, Piergiovanni G, Segura-Bayona S, Gil-Gómez G, Youssef SA, Attolini CS-O, et al.
491   GEMC1 is a critical regulator of multiciliated cell differentiation. EMBO J. 2016;35:942–60.

492   4.   Arbi M, Pefani D-E, Kyrousi C, Lalioti M-E, Kalogeropoulou A, Papanastasiou AD, et al.
493   GemC1 controls multiciliogenesis in the airway epithelium. EMBO Rep. 2016;17:400–13.

494   5.   Zhou F, Narasimhan V, Shboul M, Chong YL, Reversade B, Roy S. Gmnc Is a Master
495   Regulator of the Multiciliated Cell Differentiation Program. Curr. Biol. 2015;25:3267–73.

496   6.   Boon M, Wallmeier J, Ma L, Loges NT, Jaspers M, Olbrich H, et al. MCIDAS mutations
497   result in a mucociliary clearance disorder with reduced generation of multiple motile cilia.
498   Nat. Commun. 2014;5:4418.

499   7.   Ma L, Quigley I, Omran H, Kintner C. Multicilin drives centriole biogenesis via E2f proteins.
500   Genes Dev. 2014;28:1461–71.

501   8.   Stubbs JL, Vladar EK, Axelrod JD, Kintner C. Multicilin promotes centriole assembly and
502   ciliogenesis during multiciliate cell differentiation. Nat. Cell Biol. 2012;14:140–7.

503   9.   Danielian PS, Hess RA, Lees JA. E2f4 and E2f5 are essential for the development of the
504   male reproductive system. Cell Cycle Georget. Tex 2016;15:250–60.

505   10.   Danielian PS, Bender Kim CF, Caron AM, Vasile E, Bronson RT, Lees JA. E2f4 is required
506   for normal development of the airway epithelium. Dev. Biol. 2007;305:564–76.

507   11.   Marcet B, Chevalier B, Luxardi G, Coraux C, Zaragosi L-E, Cibois M, et al. Control of
508   vertebrate multiciliogenesis by miR-449 through direct repression of the Delta/Notch
509   pathway. Nat. Cell Biol. 2011;13:693–9.

510   12.   Nemajerova A, Kramer D, Siller SS, Herr C, Shomroni O, Pena T, et al. TAp73 is a central
511   transcriptional regulator of airway multiciliogenesis. Genes Dev. 2016;30:1300–12.

512   13.   Marshall CB, Mays DJ, Beeler JS, Rosenbluth JM, Boyd KL, Santos Guasch GL, et al. p73
513   Is Required for Multiciliogenesis and Regulates the Foxj1-Associated Gene Network. Cell
514   Rep. 2016;14:2289–300.

515   14.   Blatt EN, Yan XH, Wuerffel MK, Hamilos DL, Brody SL. Forkhead transcription factor HFH-
516   4 expression is temporally related to ciliogenesis. Am. J. Respir. Cell Mol. Biol.
517   1999;21:168–76.

518   15.   Brody SL, Yan XH, Wuerffel MK, Song SK, Shapiro SD. Ciliogenesis and left-right axis
519   defects in forkhead factor HFH-4-null mice. Am. J. Respir. Cell Mol. Biol. 2000;23:45–51.

16. Chen J, Knowles HJ, Hebert JL, Hackett BP. Mutation of the mouse hepatocyte nuclear
factor/forkhead homologue 4 gene results in an absence of cilia and random left-right
asymmetry. J. Clin. Invest. 1998;102:1077–82.

17. Yu X, Ng CP, Habacher H, Roy S. Foxj1 transcription factors are master regulators of the
motile ciliogenic program. Nat. Genet. 2008;40:1445–53.

18. Crow J, Amso NN, Lewin J, Shaw RW. Morphology and ultrastructure of fallopian tube
epithelium at different stages of the menstrual cycle and menopause. Hum. Reprod. Oxf.
Engl. 1994;9:2224–33.

19. Lyons RA, Saridogan E, Djahanbakhch O. The reproductive significance of human
Fallopian tube cilia. Hum. Reprod. Update 2006;12:363–72.

20. Raidt J, Werner C, Menchen T, Dougherty GW, Olbrich H, Loges NT, et al. Ciliary function
and motor protein composition of human fallopian tubes. Hum. Reprod. Oxf. Engl.
2015;30:2871–80.

21. Vanaken GJ, Bassinet L, Boon M, Mani R, Honoré I, Papon J-F, et al. Infertility in an adult
cohort with primary ciliary dyskinesia: phenotype–gene association. Eur. Respir. J.
2017;50:1700314.

22. Ilio KY, Hess RA. Structure and function of the ductuli efferentes: a review. Microsc. Res.
Tech. 1994;29:432–67.

23. Lambot M-AH, Mendive F, Laurent P, Van Schoore G, Noël J-C, Vanderhaeghen P, et al.
Three-dimensional reconstruction of efferent ducts in wild-type and Lgr4 knock-out mice.
Anat. Rec. Hoboken NJ 2007 2009;292:595–603.

24. Hess RA. The Efferent Ductules: Structure and Functions [Internet]. In: Robaire B, Hinton
BT, editors. The Epididymis: From Molecules to Clinical Practice. Boston, MA: Springer US;
2002 [cited 2017 Jan 19]. page 49–80.Available from: http://link.springer.com/10.1007/978-
1-4615-0679-9_4

25. Hess RA. Small tubules, surprising discoveries: from efferent ductules in the turkey to the
discovery that estrogen receptor alpha is essential for fertility in the male. Anim. Reprod.
2015;12:7–23.

26. Lun MP, Johnson MB, Broadbelt KG, Watanabe M, Kang Y -j., Chau KF, et al. Spatially
Heterogeneous Choroid Plexus Transcriptomes Encode Positional Identity and Contribute
to Regional CSF Production. J. Neurosci. 2015;35:4903–16.

27. Silva-Vargas V, Maldonado-Soto A, Mizrak D, Codega P, Doetsch F. Age-Dependent
Niche Signals from the Choroid Plexus Regulate Adult Neural Stem Cells. Cell Stem Cell
2016;19:643–52.

28. Spassky N, Merkle FT, Flames N, Tramontin AD, García-Verdugo JM, Alvarez-Buylla A.
Adult ependymal cells are postmitotic and are derived from radial glial cells during
embryogenesis. J. Neurosci. Off. J. Soc. Neurosci. 2005;25:10–8.

29. Lun MP, Monuki ES, Lehtinen MK. Development and functions of the choroid plexus-
cerebrospinal fluid system. Nat. Rev. Neurosci. 2015;16:445–57.

30. Del Bigio MR. Ependymal cells: biology and pathology. Acta Neuropathol. (Berl.)
2010;119:55–73.

31. Li L, Grausam KB, Wang J, Lun MP, Ohli J, Lidov HGW, et al. Sonic Hedgehog promotes
proliferation of Notch-dependent monociliated choroid plexus tumour cells. Nat. Cell Biol.
2016;18:418–30.

32. Tomasini R, Tsuchihara K, Wilhelm M, Fujitani M, Rufini A, Cheung CC, et al. TAp73
knockout shows genomic instability with infertility and tumor suppressor functions. Genes
Dev. 2008;22:2677–91.

33. Song R, Walentek P, Sponer N, Klimke A, Lee JS, Dixon G, et al. miR-34/449 miRNAs are
required for motile ciliogenesis by repressing cp110. Nature 2014;510:115–20.

34. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an
open-source platform for biological-image analysis. Nat. Methods 2012;9:676–82.

35. Capece V, Garcia Vizcaino JC, Vidal R, Rahman R-U, Pena Centeno T, Shomroni O, et al.
Oasis: online analysis of small RNA deep sequencing data. Bioinformatics 2015;31:2205–
7.

36. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal 2011;17:10.

37. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast
universal RNA-seq aligner. Bioinforma. Oxf. Engl. 2013;29:15–21.

38. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinforma. Oxf. Engl. 2014;30:923–30.

39. Friedländer MR, Mackowiak SD, Li N, Chen W, Rajewsky N. miRDeep2 accurately
identifies known and hundreds of novel microRNA genes in seven animal clades. Nucleic
Acids Res. 2012;40:37–52.

40. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. Genome Biol. 2014;15:550.

41. Faubel R, Westendorf C, Bodenschatz E, Eichele G. Cilia-based flow network in the brain
ventricles. Science 2016;353:176–8.

42. Holembowski L, Kramer D, Riedel D, Sordella R, Nemajerova A, Dobbelstein M, et al.
TAp73 is essential for germ cell adhesion and maturation in testis. J. Cell Biol.
2014;204:1173–90.

43. Inoue S, Tomasini R, Rufini A, Elia AJ, Agostini M, Amelio I, et al. TAp73 is required for
spermatogenesis and the maintenance of male fertility. Proc. Natl. Acad. Sci.
2014;111:1843–8.

44. Dacheux J-L, Dacheux F. New insights into epididymal function in relation to sperm
maturation. Reproduction 2013;147:R27–42.

45.   Mendive F, Laurent P, Van Schoore G, Skarnes W, Pochet R, Vassart G. Defective
596   postnatal development of the male reproductive tract in LGR4 knockout mice. Dev. Biol.
597   2006;290:421–34.

598   46.   Santos Guasch GL, Beeler JS, Marshall CB, Shaver TM, Sheng Q, Johnson KN, et al. p73
599   Is Required for Ovarian Follicle Development and Regulates a Gene Network Involved in
600   Cell-to-Cell Adhesion. iScience 2018;8:236–49.

601   47.   Huang X, Ketova T, Fleming JT, Wang H, Dey SK, Litingtung Y, et al. Sonic hedgehog
602   signaling regulates a novel epithelial progenitor domain of the hindbrain choroid plexus.
603   Dev. Camb. Engl. 2009;136:2535–43.

604   48.   Kyrousi C, Lalioti M-E, Skavatsou E, Lygerou Z, Taraviras S. Mcidas and GemC1/Lynkeas
605   specify embryonic radial glial cells. Neurogenesis 2016;3:e1172747.

606   49.   Mori M, Hazan R, Danielian PS, Mahoney JE, Li H, Lu J, et al. Cytoplasmic E2f4 forms
607   organizing centres for initiation of centriole amplification during multiciliogenesis. Nat.
608   Commun. 2017;8:15857.

609   50.   Caspary T, Larkins CE, Anderson KV. The Graded Response to Sonic Hedgehog Depends
610   on Cilia Architecture. Dev. Cell 2007;12:767–78.

611   51.   Lizé M, Herr C, Klimke A, Bals R, Dobbelstein M. MicroRNA-449a levels increase by
612   several orders of magnitude during mucociliary differentiation of airway epithelia. Cell Cycle
613   Georget. Tex 2010;9:4579–83.

614   52.   Marcet B, Chevalier B, Coraux C, Kodjabachian L, Barbry P. MicroRNA-based silencing of
615   Delta/Notch signaling promotes multiple cilia formation. Cell Cycle 2011;10:2858–64.

616   53.   Otto T, Candido SV, Pilarz MS, Sicinska E, Bronson RT, Bowden M, et al. Cell cycle-
617   targeting microRNAs promote differentiation by enforcing cell-cycle exit. Proc. Natl. Acad.
618   Sci. U. S. A. 2017;114:10660–5.

619   54.   Redshaw N, Wheeler G, Hajihosseini MK, Dalmay T. microRNA-449 is a putative regulator
620   of choroid plexus development and function. Brain Res. 2009;1250:20–6.

621   55.   Yang X, Feng M, Jiang X, Wu Z, Li Z, Aau M, et al. miR-449a and miR-449b are direct
622   transcriptional targets of E2F1 and negatively regulate pRb-E2F1 activity through a
623   feedback loop by targeting CDK6 and CDC25A. Genes Dev. 2009;23:2388–93.

624   56.   Lizé M, Pilarski S, Dobbelstein M. E2F1-inducible microRNA 449a/b suppresses cell
625   proliferation and promotes apoptosis. Cell Death Differ. 2010;17:452–8.

626   57.   Kyrousi C, Arbi M, Pilz G-A, Pefani D-E, Lalioti M-E, Ninkovic J, et al. Mcidas and GemC1
627   are key regulators for the generation of multiciliated ependymal cells in the adult
628   neurogenic niche. Development 2015;142:3661–74.

629   58.   Kim S, Ma L, Shokhirev MN, Quigley I, Kintner C. Multicilin and activated E2f4 induce
630   multiciliated cell differentiation in primary fibroblasts. Sci. Rep. 2018;8:12369.

631   59.   Ibanez-Tallon I. Dysfunction of axonemal dynein heavy chain Mdnah5 inhibits ependymal
632   flow and reveals a novel mechanism for hydrocephalus formation. Hum. Mol. Genet.
633   2004;13:2133–41.

634   60.   Banizs B, Pike MM, Millican CL, Ferguson WB, Komlosi P, Sheetz J, et al. Dysfunctional
635   cilia lead to altered ependyma and choroid plexus function, and result in the formation of
636   hydrocephalus. Dev. Camb. Engl. 2005;132:5329–39.

637   61.   Banizs B, Komlosi P, Bevensee MO, Schwiebert EM, Bell PD, Yoder BK. Altered pH(i)
638   regulation and Na(+)/HCO3(-) transporter activity in choroid plexus of cilia-defective
639   Tg737(orpk) mutant mouse. Am. J. Physiol. Cell Physiol. 2007;292:C1409-1416.

640   62.   Bill BR, Balciunas D, McCarra JA, Young ED, Xiong T, Spahn AM, et al. Development and
641   Notch Signaling Requirements of the Zebrafish Choroid Plexus. PLoS ONE 2008;3:e3114.

642   63.   Amirav I, Wallmeier J, Loges NT, Menchen T, Pennekamp P, Mussaffi H, et al. Systematic
643   Analysis of CCNO Variants in a Defined Population: Implications for Clinical Phenotype and
644   Differential Diagnosis. Hum. Mutat. 2016;37:396–405.

645   64.   Guglielmino MR, Santonocito M, Vento M, Ragusa M, Barbagallo D, Borzì P, et al. TAp73
646   is downregulated in oocytes from women of advanced reproductive age. Cell Cycle
647   Georget. Tex 2011;10:3253–6.

648   65.   Hu W, Zheng T, Wang J. Regulation of Fertility by the p53 Family Members. Genes Cancer
649   2011;2:420–30.

650   66.   Feng Z, Zhang C, Kang H-J, Sun Y, Wang H, Naqvi A, et al. Regulation of female
651   reproduction by p53 and its family members. FASEB J. Off. Publ. Fed. Am. Soc. Exp. Biol.
652   2011;25:2245–55.

653   67.   Fujitani M, Sato R, Yamashita T. Loss of p73 in ependymal cells during the perinatal period
654   leads to aqueductal stenosis. Sci. Rep. 2017;7:12007.

655   68.   Gonzalez-Cano L, Fuertes-Alvarez S, Robledinos-Anton N, Bizy A, Villena-Cortes A,
656   Fariñas I, et al. p73 is required for ependymal cell maturation and neurogenic SVZ
657   cytoarchitecture. Dev. Neurobiol. 2016;76:730–47.

658   69.   Pefani D-E, Dimaki M, Spella M, Karantzelis N, Mitsiki E, Kyrousi C, et al. Idas, a Novel
659   Phylogenetically Conserved Geminin-related Protein, Binds to Geminin and Is Required for
660   Cell Cycle Progression. J. Biol. Chem. 2011;286:23234–46.

661   70.   Bao J, Li D, Wang L, Wu J, Hu Y, Wang Z, et al. MicroRNA-449 and MicroRNA-34b/c
662   Function Redundantly in Murine Testes by Targeting E2F Transcription Factor-
663   Retinoblastoma Protein (E2F-pRb) Pathway. J. Biol. Chem. 2012;287:21686–98.

664   71.   Fededa JP, Esk C, Mierzwa B, Stanyte R, Yuan S, Zheng H, et al. MicroRNA-34/449
665   controls mitotic spindle orientation during mammalian cortex development. EMBO J.
666   2016;35:2386–98.

667   72.   Agostini M, Tucci P, Killick R, Candi E, Sayan BS, Rivetti di Val Cervo P, et al. Neuronal
668   differentiation by TAp73 is mediated by microRNA-34a regulation of synaptic protein
669   targets. Proc. Natl. Acad. Sci. U. S. A. 2011;108:21093–8.

73. Medina-Bolívar C, González-Arnay E, Talos F, González-Gómez M, Moll UM, Meyer G.
Cortical hypoplasia and ventriculomegaly of p73-deficient mice: Developmental and adult
analysis: p73 in developing and adult cortex. J. Comp. Neurol. 2014;522:2663–79.

74. Yang A, Walker N, Bronson R, Kaghad M, Oosterwegel M, Bonnin J, et al. p73-deficient
mice have neurological, pheromonal and inflammatory defects but lack spontaneous
tumours. Nature 2000;404:99–103.

75. Tissir F, Ravni A, Achouri Y, Riethmacher D, Meyer G, Goffinet AM. DeltaNp73 regulates
neuronal survival in vivo. Proc. Natl. Acad. Sci. U. S. A. 2009;106:16871–6.

76. Koeppel M, van Heeringen SJ, Kramer D, Smeenk L, Janssen-Megens E, Hartmann M, et
al. Crosstalk between c-Jun and TAp73alpha/beta contributes to the apoptosis-survival
balance. Nucleic Acids Res. 2011;39:6069–85.

77. Diez-Roux G, Banfi S, Sultan M, Geffers L, Anand S, Rozado D, et al. A High-Resolution
Anatomical Atlas of the Transcriptome in the Mouse Embryo. PLoS Biol. 2011;9:e1000582.

 Figure Legends

Fig. 1. TAp73 is expressed in diverse multiciliated epithelia. (**a**) Schematic illustration of

efferent ducts (EDs, red arrows) that connect testis and epididymis (Epi). Blue dotted lines indicate

regions used for histological, protein, and RNA analyses. (**b**) Representative images of the

expression of P73 (green) and the axonemal marker acetylated-alpha tubulin (Ac-α-TUB, red) in

the human ED. White bracket circles delineate P73 nuclear staining. DAPI staining (blue) marks

nuclei. (**c**) Schematic illustration of the fallopian tube (FT) that connects ovary and uterus. Blue

dotted line illustrates the region used for immunofluorescence analysis including fimbriae (red

arrow). (**d**) Expression of P73 in human FT. Upper panel depicts a magnification of the boxed

region in the lower panel. Arrowheads mark P73$^+$ cells. Images were retrieved from Human Protein

Atlas (https://www.proteinatlas.org/ENSG00000078900-TP73/tissue/fallopian+tube). (**e**)

Schematic illustration of murine brain ventricles (red arrows). Blue dotted lines indicate the position

of coronal brain slices used in protein and RNA analyses. (**f**) Expression of TAp73 in lateral and

4th ventricles of wild type (WT) adult mice. Red dotted lines demarcate ependymal cells lining brain

ventricles. Notice that both ependymal and choroid plexus (CP) epithelial cells express TAp73. (**g**)

Quantitative PCR analysis of *TAp73* expression in the testis, EDs, FTs, and brain ventricles from

WT adult mice. Expression levels are shown in copy number. Data from a single experiment are

shown (testis, *n*=3; FT and ventricle, *n*=4; ED, *n*=6). (**h**) Western blot analysis of the expression

of TAp73 and ΔNp73 in testis, Epi, FT, and brain ventricle from WT adult mice. Heat shock cognate

71 kDa protein (HSC70) serves as a loading control. Data are representative of two independent

experiments. (**i**) Quantitation of the signal intensity of TAp73 bands relative to that of HSC70 (**h**)

is shown. All data are presented as mean ± SEM with *P<0.05.

Fig. 2. TAp73 controls multiciliogenesis in the male reproductive tract. (**a**) Representative images of hematoxylin and eosin (H&E) staining of Epi sections from WT and *TAp73* knockout (KO) animals. Bracket lines demarcate the border of EDs and Epi. Notice the lack of mature spermatozoa in cauda Epi from *TAp73* KO mice (arrowheads). (**b**) Representative images of the expression of Ac-α-TUB (green) and axonemal dynein DNAI1 (red) in EDs from WT and *TAp73* KO mice. DAPI staining (blue) labels nuclei. Boxed regions are magnified in the bottom panel. Note that *TAp73* KO mice have less cilia that also exhibit reduced length (white bars). (**c**) Quantitation of Ac-α-TUB and DNAI1 signals normalized to epithelial length. Data from a single experiment are shown (WT, n=6 images from 3 animals; *TAp73* KO, n=11 images from 4 animals). (**d**) Representative photomicrographs of transmission electron microscopy (TEM) in EDs from WT and *TAp73* KO mice. Dotted lines mark apical region of the cells. Notice the abundant cilia (white arrows) and clustered basal bodies (white arrowhead) docked to the apical surface of WT cells, whereas mutant cells exhibit fewer cilia (red arrow). Interspersed microvilli are marked with asterisks. (**e**) Chromatin immunoprecipitation was performed for Saos2 cells transfected with TAp73α, TAp73β, and empty vector. Binding of TAp73α and TAp73β to genomic regions of *FOXJ1*, axonemal dyneins *DNALI1* and *DNAI1* was evaluated by quantitative PCR and compared to vector control (n=3 for each antibody/gene pair, except for DNALI1 [n=4], genomic regions examined are illustrated in **Supplementary Fig. 3**; [76]). (**f**) Semi-quantitative PCR analysis of *Dnali1, Foxj1*, *Rfx2, and Rfx3* expression in EDs from WT and *TAp73* KO mice. Data from a single experiment are shown (WT, n=4 for *Dnali1, Foxj1*, and *Rfx3, n*=3 for Rfx2; *TAp73* KO, n=3). (**g**) Immunoblot analysis of the expression of TAp73, ΔNp73, DNAI1, and DNALI1 in Epi from WT and *TAp73* KO animals. HSC70 serves as a loading control. Representative result of three independent experiments is displayed. All data are presented as mean ± SEM and relative to the WT group with *P<0.05, ** P <0.01, *** P <0.001.

Fig. 3. TAp73 controls multiciliogenesis in the oviducts. (**a**) Representative H&E staining of FTs from WT and *TAp73* KO animals. (**b**) The expression of Ac-α-TUB (green) and DNAI1 (red) in FTs from WT and *TAp73* KO mice. DAPI staining (blue) labels nuclei. Boxed regions are magnified in the bottom panel. In contrast to multiciliated epithelia in WT mice, *TAp73* KO mice exhibit FT segments devoid of cilia (arrowheads). (**c**) Quantitation of Ac-α-TUB and DNAI1 signals normalized to epithelial length. Data from a single experiment are shown (WT, n=6 images from 4 mice; *TAp73* KO, n=6 images from 3 mice). (**d**) Representative TEM photomicrographs of FTs from WT and *TAp73* KO animals. Dotted lines mark apical region of the cells. Notice the presence of abundant cilia (black arrows) and basal bodies (white arrowhead) docked at the apical surface of WT cells, whereas mutant cells display fewer cilia (red arrows). Interspersed microvilli are marked with asterisks. (**e**) Semi-quantitative PCR analysis of *Dnali1, Foxj1, Rfx2,* and *Rfx3* expression in oviducts from WT and *TAp73* KO mice. Data from a single experiment are shown (n=3). (**f**) Immunoblot analysis of the expression of TAp73, ΔNp73, FOXJ1, DNAI1, DNALI1, and gamma tubulin (γ-TUB) in oviducts from WT and *TAp73* KO animals. HSC70 serves as a loading control. Data are representative of three independent experiments. All data are presented as mean ± SEM and relative to the WT group with *P<0.05, ** P<0.01.

Fig. 4. *TAp73* is dispensable for brain multiciliogenesis. (**a**) The expression of KI-67, Aquaporin 1 (AQP1, green), and ADP-ribosylation factor-like 13b (ARL13B, red) in WT hindbrain roof plate/CP at embryonic (E) day 14.5. Notice that KI-67$^+$ roof plate progenitors, and AQP1$^+$ CP epithelial cells are spatially separated. ARL13B labels monociliated roof pate progenitors and multiciliated CP epithelial cells. White lines demarcate roof pate epithelium (KI-67$^+$/AQP1$^-$, arrows), CP epithelium (KI-67$^-$/AQP1$^+$, asterisks), and "transition zone" (KI-67$^-$/AQP1$^-$, arrowheads) in which MCCs appear. Dotted lines mark apical cell surface with cilia. DAPI staining (blue) labels nuclei. (**b**) Expression of TAp73 (green, red), AQP1 (green), and ARL13B (red) in WT hindbrain roof plate/CP at E14.5. Dotted lines mark apical cell surface of roof plate (TAp73$^-$,

arrow) and transition zone (TAp73$^+$, arrowhead). White lines mark transition zone (TAp73$^+$/AQP1$^-$

, arrowhead) and CP epithelium (TAp73$^+$/AQP1$^+$, asterisk). DAPI staining (blue) labels nuclei. (**c**)

Representative images of TAp73 expression of ependymal and CP epithelial cells in hindbrain

and lateral ventricle from WT and *TAp73* KO. Red dotted lines mark ventricles lined with

ependymal cells. Note that p73 expression is lost in *TAp73* KO mice. (**d**) Expression of the cilia

marker ARL13B (red) in CP epithelial cells from WT and *TAp73* KO. White arrowheads mark cilia

on cell surface. DAPI staining (blue) labels nuclei. Quantitation of average cilia length is shown in

the lower panel. Data from a single experiment are shown (WT, *n*=12 cells [hindbrain] and 9 cells

[lateral ventricle] from 2 mice; *TAp73* KO, *n*=17 cells [hindbrain] and 15 cells [lateral ventricle] from

3 mice). (**e**) Immunoblot analysis of Ac-α-TUB in brain ventricles from WT and *TAp73* KO animals.

β-ACTIN serves as a loading control. Data are representative of two independent experiments. (**f**)

Semi-quantitative PCR of *Dnali1, Foxj1, Rfx2,* and *Rfx3* in brain ventricles from WT and *TAp73*

KO. Data from a single experiment are shown (WT, *n*=3; *TAp73* KO, *n*=4). (**g**) Movement of

fluorescent beads along the ventricular system in WT and *TAp73* KO mice. Images of maximum

intensity projections of representative movies of the lateral and the ventral 3rd ventricles are shown

(WT, *n*=2; *TAp73* KO, *n*=3; *TAp73* heterozygous, *n*=1). Red arrows mark the direction of bead

flow. Bracket lines depict ependymal layer lining the lateral ventricles. Refer to **Supplementary**

Video S3a, b for examples of recording of ciliary beating. All data are presented as mean ± SEM

and relative to the WT group with **P<0.05, ** $P<0.01$.

Fig. 5. *TAp73* loss leads to changes in *miR-34/449* family and E2F4/MCIDAS circuit in the

brain. (**a**) Hierarchical clustering of differentially expressed miRNAs in brain ventricles from WT

and *TAp73* KO mice (WT, *n*=3; *TAp73* KO, *n*=4, one-way ANOVA, FDR < 0.05, fold change is

shown). log$_2$ values for miRNAs are plotted on the right. (**b**) *In situ* hybridization analysis of the

expression of *miR449* in WT roof plate/CP at E14.5 (http://www.eurexpress.org/ee/) [77]). Semi-

quantitative PCR analysis of *miR449a*, *miR34b,* and *miR34c* in brain ventricles (**c**), *miR449a*

expression in EDs, FTs, and trachea (**d**), and *E2f4* and *Mcidas* levels in brain ventricles (**e**) from

WT and *TAp73* KO mice. Data from a single experiment are shown (WT: ED, n=3; FT, n=7;

trachea, n=4; ventricle, n=3. *TAp73* KO: ED, n=4; FT, n=8; trachea, n=4; ventricle, n=4). (**f**)

Immunoblot of E2F4 in WT and *TAp73* KO ventricles. β-ACTIN serves as a loading control.

Representative result of three independent experiments. (**g**) Luciferase assay of *miR449*

regulatory regions containing E2F binding motifs. Three consensus E2F binding sites in *miR449*

locus (http://jaspar.binf.ku.dk/) were placed in front of a luciferase cassette. A deletion mutant

(Mut) that lacks the strongest consensus site was also created (**Supplementary Table 5**). WT or

Mut luciferase vector was then co-transfected with empty vector (control), or vectors expressing

E2F4, MCIDAS, or both. Fold changes in luciferase activities relative to those of control vector are

shown. Date from 5 independent experiments are shown. All data are presented as mean ± SEM

and relative to the WT group with *P<0.05, ** P <0.01.

Fig. 6 TAp73 functions through *miR449* in brain multiciliogenesis. (**a**) Coronal brain slices

from WT, *TAp73* KO, *miR449* KO and *TAp73xmiR449* KO mice. Note that *TAp73xmiR449* KO

mice display enlarged lateral ventricles. (**b**) ARL13B (red) expression in CP epithelial cells of the

4[th] and lateral ventricles from WT, *miR449* KO, and *TAp73xmiR449* KO animals. White

arrowheads mark cilia on cell surface. DAPI staining (blue) labels nuclei. (**c**) Quantitation of

average cilia length of CP epithelial cells shown in (**b**). Data from a single experiment are shown

(WT, n=4 cells [hindbrain, lateral ventricle] from 2 mice; *miR449* KO, n=18 cells [hindbrain] and

14 cells [lateral ventricle] from 4 mice; *TAp73xmiR449* KO, n=8 cells [hindbrain, lateral ventricle]

from 3 mice). (**d**) Representative TEM photomicrographs of ependymal cells in WT, *TAp73* KO,

and *TAp73x<i>miR449</i>* KO mice. Dotted lines mark apical region of the cells. Notice that WT cells

possess cilia (white arrow) and basal bodies (white arrowhead) docked to the apical surface,

whereas mutant cells have a similar number of cilia (red arrows) but disorganized basal bodies

(red arrowheads) located further away from the apical surface. Interspersed microvilli are marked

with asterisks. (**e**) Representative staining of Ac-α-TUB (green) in tracheae from WT and

TAp73xmiR449 KO mice. DAPI staining (blue) labels nuclei. Note that mutants harbor less and

shorter cilia (arrowhead) than WTs. (**f**) Quantitation of Ac-α-TUB signals normalized to epithelial

length is shown. Data from a single experiment are shown (*n*=4 samples/genotype). (**g**)

Representative TEM photomicrographs of tracheae from WT and *TAp73xmiR449* KO mice. Dotted

lines mark apical region of the cells. Notice the abundant cilia (black arrows) and clustered basal

bodies (white arrowhead) docked to apical surface in WT cells, whereas mutant cells exhibit fewer

cilia (red arrow). Interspersed microvilli are marked with asterisks. All data are presented as mean

± SEM and relative to the WT group with ** P <0.01, *** P <0.001.

Fig. 7. Schematic diagram of the molecular circuits of TAp73-driven multiciliogenesis in

diverse tissues. (**a**) TAp73-dependent transcriptional network, including dyneins, *miR34bc*,

Foxj1, Rfx2, and Rfx3 factors, critically regulates multiciliogenesis in various ciliated epithelia

downstream of *E2f4*/*Mcidas*. In the EDs (**b**) and FTs (**c**) *TAp73* KO impairs multiciliogenesis

concurrent with male and female fertility. (**d**) TAp73 is not essential for multiciliogenesis in the

brain; however, *TAp73* loss leads to upregulation of pro-ciliogenic *E2f4* and its target *miR449*. (**e**)

Further removal of *miR449* in *TAp73* KO animals leads to reduced number and length of CP cilia

and severe hydrocephalus, indicating that *miR449* and TAp73 complement each other to support

brain ciliogenesis.

Wildung et al. Figure 1

a WT   *TAp73* KO

ED   Epi   100 µm

Cauda Epi   Cauda Epi   100 µm

Epididymis

b WT   *TAp73* KO

Ac-α-TUB

DNAI1

Merge   100 µm

Merge   100 µm

Efferent duct

c Efferent duct

☐ WT ■ *TAp73* KO

Relative Cilia marker / epithelial length [AU]

Ac-α-TUB   DNAI1

*** **

d WT   *TAp73* KO

2 µm

Efferent duct

e Saos2 cells

■ *Empty* ☐ *TAp73α* ■ *TAp73β*

% of input

FOXJ1 down   *FOXJ1* end   *DNALI1*   *DNAI1*   Avg. IgG

* **   * ***   * **   p=0.06 ***

f Epididymis

☐ WT ■ *TAp73 KO*

Relative gene expression

Dnali1   *Foxj1*   *Rfx2*   *Rfx3*

**   *   *   **

g Epididymis

WT   *TAp73* KO   kDA

TAp73   100 / 70

ΔNp73   70

DNAI1   100

DNALI1   35

HSC70   70

Wildung et al. Figure 3

Wildung et al. Figure 4

Wildung et al. Figure 6

a

WT   *TAp73* KO   *miR449* KO   *TAp73xmiR449* KO

b

4th ventricle   Lateral ventricle

ARL13B

WT

miR449 KO

TAp73xmiR449 KO

5 μm

Choroid plexus

c

4th ventricle   Lateral ventricle

Relative cilia length

***

WT   *miR449* KO   *TAp73xmiR449* KO

d

WT   *TAp73* KO   *TAp73xmiR449* KO

2 μm

Ependyma

e

WT   *TAp73xmiR449* KO

Ac-α-TUB

100 μm

Trachea

f

Relative Ac-α-TUB / epithelial length [AU]

**

WT   *TAp73xmiR449* KO

g

WT   *TAp73xmiR449* KO

2 μm   2 μm

Trachea

a Wild type

Multiciliated tissues

b *TAp73* KO ED ♂

c *TAp73* KO FT ♀

Reproductive ducts

d *TAp73* KO

e *TAp73xmiR449* KO

Brain ventricles/Choroid plexus

Supplementary information

Transcription factor TAp73 and microRNA-449 complement each other to support multiciliogenesis

Running title: TAp73 and miR449 cooperate in multiciliogenesis

Merit Wildung[1,*], Tilman Uli Esser[1,*], Katie Baker Grausam[2,3], Cornelia Wiedwald[1], Larisa Volceanov-Hahn[1], Dietmar Riedel[4], Sabine Beuermann[1], Li Li[2], Jessica Zylla[2], Ann-Kathrin Guenther[5], Magdalena Wienken[6], Evrim Ercetin[1], Zhiyuan Han[7], Felix Bremmer[8], Orr Shomroni[9], Stefan Andreas[1], Haotian Zhao[2,3,7,#] and Muriel Lizé[1,#]

* = equal contribution; # = corresponding authors

Content:

Supplementary Figures

Supplementary Fig. 1. TAp73 is expressed in diverse multiciliated tissues. RNAscope analysis of *TAp73* (blue) and *Mcidas* (red) expression in testis, efferent ducts, fallopian tube, trachea, lung, ependymal and choroid plexus epithelial cells in hindbrain and lateral ventricle from adult wild type (WT) and *TAp73* knockout (KO) mice (3 months of age). Arrowheads mark cells in boxed regions that are also shown in higher magnification. Notice that *TAp73* is absent in *TAp73* KO tissues.

Supplementary Fig. 2. Loss of _TAp73_ impairs multiciliogenesis in the male reproductive duct. (**a**) Representative images of the expression of acetylated alpha-tubulin (Ac-α-TUB, green) in testes from WT and _TAp73_ KO mice. DAPI staining (blue) labels nuclei. Notice, although at a reduced level, spermatozoa (Ac-α-TUB$^+$) are present in mutant testis (arrowheads). (**b**) Expression of TAp73 in testes from WT and _TAp73_ KO mice at 3 months of age, respectively. Notice that TAp73 is expressed in testes from WT mice, but absent in mutant testes. (**c**) Representative photomicrographs of transmission electron microscopy (TEM) of efferent ducts from WT and _TAp73_ KO mice. Dotted lines mark apical region of the cells. Notice the abundant cilia (white arrow) and clustered basal bodies (white arrowheads) on WT cells, whereas mutant cells exhibit disorganized basal bodies (red arrowheads) located away from apical surface. Interspersed microvilli are marked with asterisks. (**d**) Immunoblot analysis of the expression of DNAI1 and DNALI1 in testes from WT and _TAp73_ KO animals. β-ACTIN serves as a loading control. DNAI1 and DNALI1 levels are reduced in mutant testis compared to WT animals. Representative results of three independent experiments are shown.

Supplementary Fig. 3. TAp73 is associated with ciliary genes. TAp73 binding at *FOXJ1*, *DNALI1,* and *DNAI1* genomic loci is shown in results from ChIP-seq [1], Geo accession no. **GSE15780**). Boxed regions mark genomic loci enriched with TAp73 binding and validated by ChIP-qPCR (**Fig. 2e**).

a

WT TAp73 KO

Fallopian tube

b

DNALI1 FOXJ1 DNAI1

Fallopian tube

Supplementary Fig. 4. TAp73 KO mice show defective motile cilia in the fallopian tube.
(**a**) Representative photomicrographs of transmission electron microscopy (TEM) of fallopian tubes from WT and *TAp73* KO mice. Dotted lines mark apical region of the cells. Notice the presence of abundant cilia (white arrow) and clustered basal bodies (white arrowhead) docked to apical surface of the WT cell, whereas the mutant cell displays fewer cilia (red arrow) and disorganized basal bodies (red arrowheads) located away from apical surface. Interspersed microvilli are marked with asterisks. (**b**) Expression of DNALI1, FOXJ1, and DNAI1 in human FT epithelia. Images were retrieved from the Human Protein Atlas (DNALI1: http://www.proteinatlas.org/ENSG00000163879-DNALI1/tissue/fallopian+tube, FOXJ1: http://www.proteinatlas.org/ENSG00000129654-FOXJ1/tissue/fallopian+tube, DNAI1: http://www.proteinatlas.org/ENSG00000122735-DNAI1/tissue/fallopian+tube).

Supplementary Fig. 5. *TAp73* **loss does not affect multiciliogenesis in the brain.** (**a**) Representative images of H&E staining of choroid plexus (CP) in the 4[th] and lateral ventricle from WT and *TAp73* KO animals. (**b**) Representative images of the expression of ARL13B in CP of the 4[th] and lateral ventricle from WT and *TAp73* KO animals. Arrowheads mark cilia on CP epithelial cells. Red dotted lines delineate the boundary of lateral ventricles lined with ependymal cells. Expression of Ac-α-TUB (green) and DNAI1 (red) in ependymal cells of lateral ventricle (**c**) and the 4[th] ventricle (**d**) from WT and *TAp73* KO animals. Boxed regions are magnified in bottom panels. DAPI staining (blue) labels nuclei.

Supplementary Fig. 6. *TAp73* **loss does not affect epithelial differentiation of choroid plexus cells.** Representative images of the expression of cytokeratins (**a**), aquaporin 1 (AQP1, **b**), transthyretin (TTR, **c**), and orthodenticle homeobox 2 (OTX2, **d**) in CP epithelium of the 4th and lateral ventricles from WT and *TAp73* KO animals.

Supplementary Fig. 7. Analysis of gene expression in multiciliated tissues of *TAp73* KO animals. (**a**) Semi-quantitative PCR analysis of *miR34b* expression in trachea from WT and *TAp73* KO mice. Data from a single experiment are shown (WT: *n*=3; *TAp73* KO: *n*=4). (**b**) Semi-quantitative PCR analysis of *E2f4* expression in brain ventricles, efferent ducts (ED), fallopian tubes (FT), and tracheae from WT and *TAp73* KO mice. Data from a single experiment are shown (WT: ED, *n*=5; FT, *n*=7; trachea, *n*=4; *TAp73* KO: ED, *n*=4; FT, *n*=8; trachea, *n*=4). Immunoblot analysis of the expression of E2F4 in FTs (**c**), EDs (**d**), and tracheae (**e**) from WT and *TAp73* KO animals. β-ACTIN or HSC70 serve as loading controls. Quantitation of the signal intensity of E2F4 bands normalized to that of the loading control is shown below each immunoblot (n.s. non-significant). Representative results from three independent experiments are shown. All data are presented as mean ± SEM and relative to the WT group with *P<0.05, ** P <0.01.

Supplementary Fig. 8. pRb/E2F pathway activity is not deregulated in *TAp73* KO ventricles. (**a**) Semi-quantitative PCR analysis of *Cdkn1a*, *Cdkn1b*, *E2f1*, and *E2f3* in brain ventricles from WT and *TAp73* KO animals. Data from a single experiment are shown (*n*=4). (**b**) Immunoblot analysis of P21 expression in brain ventricles from WT and *TAp73* KO animals. Data are representative of three independent experiments. Quantitation of the signal intensity of P21 bands relative to that of β-ACTIN is shown (*n*=3). All data are presented as mean ± SEM and relative to the WT group.

a

WT   *TAp73* KO   *miR449* KO   *TAp73xmiR449* KO

Lateral ventricle

1 mm

1 mm

b

4th ventricle   Lateral ventricle

WT

miR449 KO

TAp73xmiR449 KO

20 µm

ARL13B

d

Ac-α-TUB

WT

TAp73xmiR449 KO

100 µm   100 µm

Lateral ventricle

c

WT   *TAp73* KO   *TAp73xmiR449* KO

Ependyma

2 µm

e

4th ventricle   Aqueduct   Lateral ventricle

WT

TAp73xmiR449 KO

200 µm

Supplementary Fig. 9. *miR449* **collaborates with** *TAp73* **in brain multiciliogenesis.** (**a**) Representative images of H&E staining of brain sections from WT, *TAp73* KO, *miR449* KO, and *TAp73xmiR449* KO animals. Notice that *TAp73xmiR449* KO mice display enlarged lateral ventricles (arrows). (**b**) Representative images of the expression of ARL13B in cilia (arrowheads) of CP in the 4th and lateral ventricle from WT, *TAp73* KO, *miR449* KO, and *TAp73xmiR449* KO animals. (**c**) Representative photomicrographs of transmission electron microscopy (TEM) of ependymal cells from WT, *TAp73* KO, and *TAp73xmiR449* KO mice. Dotted lines mark apical region of the cells. Notice that the WT and *TAp73* KO cell possesses abundant cilia (arrows) and basal bodies (arrowhead) docked to the apical surface, whereas the *TAp73xmiR449* KO cell has disorganized basal bodies (red arrowheads) located further away from the apical surface. Interspersed microvilli are marked with asterisks. (**d**) Expression of Ac-α-TUB (green) in ependymal cells of the lateral ventricle from WT and *TAp73xmiR449* KO mice. Boxed regions are magnified on the right. DAPI staining (blue) labels nuclei. (**e**) Analysis of the movement of fluorescent beads along the ventricular system from WT and *TAp73xmiR449* KO mice. Images of maximum intensity projections of representative movies of the 4th and lateral ventricle, and aqueduct are shown (*n*=2). Red arrows mark the direction of bead flow. Bracket lines delineate ependymal layer lining the ventricles. Refer to **Supplementary Video 3c** for examples of recording of ciliary beating.

Supplementary Fig. 10. Combined loss of *TAp73* and *miR449* does not affect epithelial differentiation of choroid plexus cells. Representative images of the expression of cytokeratins (**a**), aquaporin 1 (AQP1, **b**), and orthodenticle homeobox 2 (OTX2, **c**) in CP epithelium of the 4th and lateral ventricles from WT, *miR449* KO, and *TAp73xmiR449* KO mice.

Supplementary Fig. 11. NOTCH signaling is unaltered in *miR449* KO and *TAp73xmiR449* KO brains. RNAscope analysis of the expression of NOTCH targets *Hes1* (blue) and *Hes5* (red) in roof plate of hindbrain and lateral ventricles from WT, *miR449* KO, and *TAp73xmiR449* KO mice.

Supplementary Fig. 12. Additional loss of *miR449* does not exacerbate ciliary defects in the airways and efferent ducts in *TAp73* KO animals. (**a**) Representative photomicrographs of transmission electron microscopy (TEM) of trachea from WT and *TAp73xmiR449* KO mice. Dotted lines mark apical region of the cells. Notice the abundant cilia (white arrow) and clustered basal bodies (white arrowhead) docked to apical surface in the WT cell, whereas the mutant cell exhibits disorganized basal bodies (red arrowheads) located away from apical surface. Interspersed microvilli are marked with asterisks. (**b**) Representative images of the expression of Ac-α-TUB (green) in ED from WT and *TAp73xmiR449* KO mice. DAPI staining (blue) labels nuclei. Notice that mutant cells have less and shorter cilia (white bars) compared to WT mice. (**c**) Quantitation of Ac-α-TUB signals normalized to epithelial length (*n*=7 images from 4 WT mice; *n*=6 images from 3 *TAp73xmiR449* KO mice). Data are presented as mean ± SEM and relative to the WT group with *P<0.05.

Supplementary Tables

Supplementary Table 1. Primary antibody information.

Antibodies	Dilution (Application)	Company	Catalog # [clone]
Mouse monoclonal anti-Ac-α-TUB	1:1 000 (IF)	Merck (*Darmstadt, Germany*)	T6793 [6-11B-1]
Mouse monoclonal anti-ARL13B	1:500 (IF)	UC Davis/NIH NeuroMab Facility (*Davis, CA, USA*)	75-287 [N295B/66]
Rabbit polyclonal anti-ARL13B	1:500 (IHC)	Proteintech (*Rosemont, IL, USA*)	17711-1-AP
Mouse monoclonal anti-AQP1	1:1 000 (IHC, IF)	Abcam (*Cambridge, UK*)	ab9566 [1/22]
Rabbit polyclonal anti-AQP1	1:1 000 (IF)	Merck	AB2219
Rabbit polyclonal anti-β-ACTIN	1:10 000 (WB)	Abcam	ab8227
Rabbit polyclonal anti-Cytokeratins	1:100 (IHC)	Agilent Technologies (*Santa Clara, CA, USA*)	Z0622
Rabbit polyclonal anti-DNAI1	1:500 (IF); 1:700 (WB)	Merck	HPA021649
Goat polyclonal anti-DNALI1	1:300 (WB)	Santa Cruz Biotechnology (*Dallas, TX, USA*)	sc-160296
Mouse monoclonal anti-E2F4	1:300 (WB)	Santa Cruz Biotechnology	sc-6851 [D-3]
Rabbit polyclonal anti-FOXJ1	1:500 (WB)	Merck	HPA005714
Mouse monoclonal anti-HSC70	1:20 000 (WB)	Santa Cruz Biotechnology	sc-7298 [B-6]
Rabbit monoclonal anti-KI-67	1:100 (IF)	Abcam	ab16667 [SP6]
Rabbit polyclonal anti-OTX2	1:500 (IHC)	Merck	AB9566
Rabbit monoclonal anti-p21	1:300 (WB)	Abcam	ab188224 [EPR18021]
Rabbit monoclonal anti-P73	1:100 (IF, IHC); 1:300 (WB)	Abcam	ab40658 [EP436Y]
Rabbit polyclonal anti-TTR	1:100 (IHC)	Proteintech	1189-1-AP
Rabbit polyclonal anti-γ-Tubulin	1:400 (WB)	Merck	T5192

IF= Immunofluorescence, IHC= Immunohistochemistry, WB= Western blot

Supplementary Table 2. Secondary antibody information.

Antibodies	Dilution (Application)	Company	Catalog #
Alexa Fluor 488 donkey anti-mouse	1:500 (IF)	Invitrogen, Thermo Fisher Scientific (*Waltham, MA, USA*)	A21202
Alexa Fluor 594 goat anti-rabbit	1:500 (IF)	Invitrogen, Thermo Fisher Scientific	A11012
Peroxidase-conjugated donkey anti-mouse	1:10 000 (WB)	Jackson ImmunoResearch (*West Grove, PA, USA*)	715-036-150
Peroxidase-conjugated donkey anti-goat	1:10 000 (WB)	Jackson ImmunoResearch	705-036-147
Peroxidase-conjugated donkey anti-rabbit	1:10 000 (WB)	Jackson ImmunoResearch	711-036-152
Biotin-SP-conjugated AffiniPure Goat anti-Rabbit IgG	1:1 000 (IHC)	Jackson ImmunoResearch	111-065-144
Biotin-SP-conjugated AffiniPure Donkey anti-Mouse	1:1 000 (IHC)	Jackson ImmunoResearch	111-065-144

IF= Immunofluorescence, IHC= Immunohistochemistry, WB= Western blot

Supplementary Table 3. Sequence information for primers used in semi-quantitative PCR.

Gene	Accession number	Ampli con (bp)	Exons	Forward primer (5´- 3´)	Reverse primer (5´- 3´)
TAp73	NM_011642	163	Ex2-Ex3	AGCAGAATGAGCGGCAGCGTT	TGTTGGACTCCTCGCTGCCTGA
Foxj1	NM_008240	200	Ex2-Ex3	CCATGCAGACCCCACCTGGCA	GGGCAAAGGCAGGGTGGATGT
Dnali1	NM_175223	213	Ex4-Ex5	TTTGGCATGAGGAAGGCACT	CTGGTTGGTCCGTTTCAGGA
Mcidas	NM_001037914	137	Ex7	AACAACGAAAAGGAGCCTGGA	GCCGCTTAGGGTCACGATTG
E2f4	NM_148952	199	Ex7-Ex9	GCACTGGACACTCGGCCT	TGCACTCTCTCGTGGGGTCG
E2f1	NM_007891.5	151	Ex3-Ex4	AACTGGGCAGCTGAGGTGC	CAAGCCGCTTACCAATCCC
E2f3	NM_001359994.1	73	Ex3	AAACGCGGTATGATACGTCCC	CCATCAGGAGACTGGCTCAG
Cdkn1a	NM_007669.5	126	Ex2-Ex3	GTGGCCTTGTCGCTGTCTT	GCGCTTGGAGTGATAGAAATCTG
Cdkn1b	NM_009875.4	130	Ex1-Ex2	AGTGTCCAGGGATGAGGAAGCGAC	TTCTTGGGCGTCTGCTCCACAGTG
Rfx2	NM_027787	74	Ex5-Ex6	GACGGCACAAGACACTCTCTG	AGAGTCTCAATCGCCATTTCAAG
Rfx3	NM_001360357	107	Ex3	ATGCAGACTTCAGAGACGGGT	ACTGGCACTTGCTGTACCAC
36b4	NM_007475	155	Ex6-Ex7	GCAGATCGGGTACCCAAC	CAGCAGCCGCAAATGCAG

Supplementary Table 4. Sequence information for primers used in ChIP-qPCR.

Gene	Forward primer (5´- 3´)	Reverse primer (5´- 3´)
FOXJ1 down	CAGCATGCCCAGAAGCTTTG	TCAGGGGCTGCATTCTTCC
FOXJ1 end	AGGGCACACTTAGCCTTTG	AGGAGACAAAGGGAGGAGG
DNAI1	CCCAAGCGGGGTAATCTCT	CTTGAGGTTGTGGGACTTCAC
DNALI1	CACGCCCGGCAAATTTCTG	CAAGGTGGGCAGATCATGTG

Supplementary Table 5. Luciferase constructs for E2F4/MCIDAS luciferase reporter assay.

RS: Restriction site

Construct name	DNA sequence of inserts The strongest E2F binding site is depicted in grey and the removed sequence in **bold underlined**. Two consensus sequences with lower score were retained in the mutant (==pink== and ==red==).	Vector	5' RS	3' RS
miR449/**Cdc20b** WT E2F binding site = **WT**	GCCAGAAAGCTGAGCACACTGGGGACT CCGTGATAAAGGGG==GAGAGGAAGAT==AT TGAGGGTTGAGGAAGAGGTCTG**GCGGG AAAT**==GACAGGGAACC==AGATGGGCTGTG CAGCCTTAGCTGCCCATCTGAGCTGCC AAGAGAGCCGAGTTGTGCCATATGGCA GGAG	pGL4.23	NheI	EcoRV
miR449/**Cdc20b** Mut E2F binding site = **Mut**	GCCAGAAAGCTGAGCACACTGGGGACT CCGTGATAAAGGGG==GAGAGGAAGAT==AT TGAGGGTTGAGGAAGAGGTCTG==GACAG GGAACC==AGATGGGCTGTGCAGCCTTAG CTGCCCATCTGAGCTGCCAAGAGAGCC GAGTTGTGCCATATGGCAGGAG	pGL4.23	NheI	EcoRV

Supplementary Table 6. Summary of small RNA-seq data from WT (*n*=3) and *TAp73* KO (*n*=4) brains. GEO accession number: **GSE108385.**

- small RNA-seq read counts from WT and TAp73 KO ventricles
- small RNA-seq differential gene expression results from WT vs. TAp73 KO ventricles.

Supplementary Video legends

Supplementary Video 1. Movement of spermatozoa from *TAp73*[+/-] (**a**, **b**) and *TAp73* KO mice (**c**, **d**).

Supplementary Video 2. Smooth muscle contraction in fallopian tubes from WT (**a**) and *TAp73* KO (**b**) mice.

Supplementary Video 3. Ciliary beating in WT (**a**, 3[rd] ventricle), *TAp73* KO (**b**, lateral ventricle), and *TAp73xmiR449* KO (**c**, lateral ventricle) mice.

References

1. Koeppel M, van Heeringen SJ, Kramer D, Smeenk L, Janssen-Megens E, Hartmann M, et al. Crosstalk between c-Jun and TAp73alpha/beta contributes to the apoptosis-survival balance. Nucleic Acids Res. 2011;39:6069–85.