Structural and functional studies of the RBPJ-SHARP complex reveal conserved corepressor binding site

Zhenyu Yuan[1*], Bradley D. VanderWielen[1*], Benedetto Daniele Giaimo[2], Leiling Pan[3], Courtney E. Collins[1], Aleksandra Turkiewicz[2], Kerstin Hein[2], Franz Oswald[3], Tilman Borggrefe[2] & Rhett A. Kovall[1]

[1]*Department of Molecular Genetics, Biochemistry and Microbiology, University of Cincinnati College of Medicine, Cincinnati, Ohio, USA,* [2]*Institute of Biochemistry, University of Giessen, Giessen, Germany,* [3]*Department of Internal Medicine I, Center for Internal Medicine, University Medical Center Ulm, 89081 Ulm, Germany. Correspondence should be addressed to RAK, E-mail: kovallra@ucmail.uc.edu*

These authors contributed equally to this work.

Abstract

The Notch pathway is a conserved signaling mechanism that is essential for cell fate decisions during pre and postnatal development. Dysregulated signaling underlies the pathophysiology of numerous human diseases, most notably T-cell acute lymphoblastic leukemia. Receptor-ligand interactions result in changes in gene expression, which are regulated by the transcription factor CSL. CSL forms a complex with the intracellular domain of the Notch receptor and the transcriptional coactivator Mastermind, which is required to activate transcription of all Notch target genes. CSL can also function as repressor by interacting with corepressor proteins, *e.g.* SHARP in mammals and Hairless in *Drosophila melanogaster*; however, its role as a transcriptional repressor is not well understood. Here we determine the high-resolution structure of RBPJ, the mouse CSL ortholog, bound to the corepressor SHARP and DNA, which reveals a new mode of corepressor binding to CSL and an interesting example for how ligand binding sites evolve in proteins. Based on the structure, we designed and tested a number of mutants in biophysical, biochemical, and cellular assays to characterize the role of RBPJ as a repressor of Notch target genes. Our cellular studies clearly demonstrate that RBPJ mutants that are deficient for binding SHARP are incapable of repressing transcription from genes responsive to Notch signaling. Altogether, our structure-function studies of the RBPJ-SHARP corepressor

complex bound to DNA provide significant insights into the repressor function of RBPJ and

identify a new binding pocket on RBPJ that could be targeted for therapeutic benefit.

Introduction

The Notch pathway is a highly conserved cell-to-cell signaling mechanism that is indispensable

for cell fate decisions during prenatal development and postnatal tissue homeostasis.[1,2]

Dysregulated signaling underlies the pathogenesis of many human diseases, including certain

types of cancer, congenital defects, and cardiovascular disease[3]. Given its association with

human disease, there have been extensive efforts to identify reagents that pharmaceutically

modulate the Notch pathway with most efforts focused on modalities that curtail overactive

Notch signaling[4,5]. However, there is a need to identify components in the Notch pathway that

when pharmacologically targeted result in upregulated signaling to treat diseases associated

with insufficient Notch signaling[3], such as myeloid leukemia, head and neck squamous cell

carcinoma, bicuspid aortic valve disease, and Alagille syndrome.

Notch signaling is initiated when Notch receptors on one cell interact with a DSL (Delta,

Serrate, Lag-2) ligand on a neighboring cell (Figure 1A)[6]. Extracellular Notch-DSL interactions

result in proteolytic cleavage of the receptor, which frees the intracellular domain of Notch

(NICD) from the cell membrane, allowing NICD to migrate into the nucleus. Nuclear NICD

directly binds the transcription factor CSL [CBF1/RBPJ, Su(H), Lag-1] and the CSL-NICD

complex recruits a member of the Mastermind (MAM) family of transcriptional coactivators

[Mastermind-like (MAML1-3) in mammals], resulting in the transcriptional activation of Notch

target genes (Figure 1A)[7]. CSL can also function as a repressor by interacting with corepressor

proteins, such as SHARP [SMRT/HDAC1-associated repressor protein, *also known as* MINT

(Msx2-interacting nuclear target) or SPEN (split ends)][8,9], Hairless in *Drosophila melanogaster*[10],

FHL1 (Four and a half LIM domains 1, *also known as* KyoT2)[11], L3MBTL3 (lethal 3 malignant

brain tumor-like 3)[12], and RITA1 (RBPJ-interacting and tubulin associated)[13,14]. Corepressors

are part of large higher order transcriptional repression complexes that contain histone-

modifying activity, *e.g.* histone deacetylase or histone demethylase, which convert chromatin

into a transcriptionally repressed state (Figure 1A)[7].

Crystal structures of Notch transcription complexes have revealed that all CSL proteins

contain a conserved structural core composed of three domains termed NTD (N-terminal

domain), BTD (β-trefoil domain), and CTD (C-terminal domain) (Figure 1B,C)[15-17]. The NTD and

CTD are immunoglobulin (Ig) domains that are structurally similar to the Rel homology region

(RHR) of transcription factors, such as NF-κB and NFAT (Nuclear factor of activated T-cells),

whereas the fold of the BTD is related to cytokine and growth factor structures, such as

Interleukin1 and FGF (Fibroblast Growth Factor)[17]. The NTD and BTD form an electropositive

surface that recognizes nucleotides in the major and minor grooves of DNA, respectively[17]. In

the transcriptionally active CSL-NICD-MAM ternary complex bound to DNA structure (Figure

1B,C), the RAM and ANK domains of NICD bind the BTD and CTD domains of CSL,

respectively, and MAM forms an elongated bent helical structure that binds the CTD-ANK

interface and the NTD of CSL[15,16]. In addition to the CSL-NICD-MAM activator complex, several

CSL-corepressor structures have been determined, including the *Drosophila* corepressor

Hairless bound to Su(H) (the fly CSL ortholog)[18], as well as FHL1 and RITA1 bound to RBPJ

(mouse/human CSL ortholog)[13,19]. These studies reveal that the binding of Hairless to the CTD

of Su(H) produces a large conformational change in CTD[18]; whereas, FHL1 and RITA1 have

regions similar to the RAM domain of NICD and bind the BTD of RBPJ in a structurally similar

manner[13,19].

SHARP is a large multidomain transcriptional coregulator protein that has folded

functional domains separated by long regions that are intrinsically disordered. Notably, SHARP

has no sequence similarity with any of the previously identified corepressors that bind CSL

(Figure 1C). SHARP was originally identified in yeast two-hybrid screens for factors that interact

with SMRT/NCoR and the transcription factor MSX2[20,21]. SHARP has traditionally been thought

of as a corepressor, because it binds SMRT/NCoR through its C-terminal SPOC domain and

represses transcription[9,20,22,23]; however, it has also been shown that through its SPOC domain

SHARP can also recruit the KMT2D coactivator complex to Notch target genes[24]. More recently,

it has been demonstrated that the N-terminal RRM (RNA recognition motif) domains of SHARP

bind the lncRNA Xist and this interaction is important for X chromosome inactivation[25,26].

Previously, we defined a region in SHARP that binds RBPJ, termed RBPID (RBPJ-interacting

domain, Figure 1C) and showed that RBPJ and SHARP form a high affinity complex[9,27].

Here we determine the X-ray structure of the RBPJ-SHARP corepressor complex bound

to DNA. We identify structure-based mutants that are essential for RBPJ-mediated repression

and characterize these mutants both *in vitro* and in cellular assays. Taken together, our studies

reveal the conserved interface of the RBPJ/SHARP corepressor complex, which provides

molecular insights into RBPJ repressor function and identify a potential site on RBPJ that could

be pharmacologically targeted to upregulate Notch signaling.

Results

Structure determination of the RBPJ/SHARP/DNA complex

In order to determine the X-ray structure of the RBPJ/SHARP corepressor complex bound to

DNA, we purified recombinant RBPJ and SHARP proteins from bacteria, corresponding to the

structural core of RBPJ (residues 53-474) and the RBPJ-interacting domain (RBPID) of SHARP

(residues 2776-2820), formed complexes with a 13mer oligomeric DNA duplex with single-

stranded overhangs, containing a single RBPJ binding site, and screened the

RBPJ/SHARP/DNA complex for crystallization conditions. While we were able to isolate

crystallization conditions for the complex, despite extensive optimization efforts, we were unable

to produce diffraction quality crystals amenable for structural analysis. Therefore, we produced

an MBP (maltose binding protein) fusion protein with the RBPID of SHARP (MBP-SHARP), *i.e.*

fixed-arm carrier approach[28], in which the MBP moiety also has surface entropy reduction

mutations engineered into it, in order to identify new crystallization conditions for the

RBPJ/MBP-SHARP/DNA complex. This strategy successfully led to crystals that nominally

diffract to 2.8 Å resolution and belong to the space group $P2_1$ (a=54.5 Å, b=231.6 Å, c=90.3 Å,

and β=99.88°) (Table 1). We demonstrated that our MBP-SHARP construct binds RBPJ

similarly as the native SHARP construct, albeit with somewhat weaker affinity (Figure S1B). This

is likely due to the close crystal contacts between MBP and RBPJ required for crystallization of

the complex. The RBPJ/MBP-SHARP/DNA complex structure was solved by molecular

replacement and refined to a final R factor and free R factor of 19.4% and 22.8%, respectively

(Table 1).

There are two RBPJ-MBP-SHARP/DNA complexes in the asymmetric unit (AU) of the

crystals (Figure S1A) that overall are very structurally similar (RMSD 2.28 for 819 cα atoms),

with even higher structural correspondence when either MBP (RMSD 0.07 for 297 cα atoms) or

RBPJ (RMSD 0.66 for 388 cα atoms) are aligned individually (Figure S1C-E). The largest

structural difference between the two complexes in the AU is a poorly ordered linker region in

SHARP (Figure S1F), whereby the different conformations are likely influenced by different

environments within the crystals. For clarity in subsequent figures, we do not show the MBP

moiety and all structural comparisons are performed with chains G & H of RBPJ and SHARP,

respectively, as these proteins chains have overall lower temperature factors.

Bipartite interaction of SHARP with RBPJ

As shown in Figure 2A-B, SHARP interacts with two distinct surfaces on RBPJ, contacting the CTD and the BTD of RBPJ, which is consistent with previous binding studies[27]. Starting at the N-terminus of its RBPID, SHARP forms a β-hairpin motif that binds between the two β-sheets that compose the Ig domain of the CTD. The second β-strand of SHARP, residues 2788-2794, pairs with β-strand βg of the CTD, extending this from a three- to five-stranded β-sheet (Figure 2C). The binding mode of SHARP is structurally analogous to how the *Drosophila melanogaster* corepressor Hairless interacts with the CTD of Su(H) (*the fly CSL ortholog*)[18], which is discussed further below. The SHARP/CTD interaction is followed by a short, approximately six residues, linker region that is poorly ordered and makes no contacts with RBPJ. The C-terminal portion of the SHARP RBPID binds in an extended fashion across the BTD in a manner that is structurally similar to the RAM domain of NICD[15,29,30], and the corepressors FHL1 and RITA1 (Figure 2A-B and described below)[13,19]. The complex between SHARP and RBPJ is largely driven by hydrophobic interactions between nonpolar side chains on SHARP, and the CTD and BTD of RBPJ (Figure 2B); with SHARP residue L2791, as well as V2789 and Y2793, anchoring its interaction with the CTD; and I2811 anchoring the interaction between SHARP and the BTD. Additionally, there are key ionic interactions that appear to play auxiliary roles in complex formation, including salt bridges between E2786 of SHARP and R438 of the CTD (Figure 2C), and K2807 of SHARP and the BTD residue E259 (Figure 2B). It should also be mentioned that when complexed with SHARP, RBPJ maintains similar nonspecific and specific contacts with DNA, suggesting that SHARP binding does not affect the affinity of RBPJ for DNA, which is also consistent with previous binding studies[27].

As shown in Figure 2C, SHARP binding to RBPJ induces a structural change in CTD that results in translation of β-strands βf and βg outward, modestly expanding the CTD by as much as ~4 Å. A significant structural rearrangement is also observed for the loop that connects strands βe with βf, which strikingly repositions W441 from a buried to a solvent exposed

155 conformation when comparing unbound RBPJ with the RBPJ/SHARP complex (Figure 2C). The

156 repositioning of W441 allows for E2786 of SHARP to form a salt bridge with R438 of RBPJ. This

157 structural rearrangement has not been observed for any other CSL structure determined to

158 date.

Structural comparison of coregulator binding sites on RBPJ

The intracellular domain of Notch (NICD) together with Mastermind (MAM) and RBPJ form a

transcriptionally active ternary complex[6]. NICD interacts with high affinity to RBPJ[30-33], binding

the BTD and CTD of RBPJ via its RAM and ANK domains, respectively[15,16]. Figure 3A

compares the interfaces that SHARP and NICD/MAM use to interact with RBPJ, illustrating the

overlap of these binding sites. There is partial overlap of the SHARP and ANK/MAM binding

sites on the CTD of RBPJ, whereas there is completely overlapping binding of SHARP and the

RAM domain of NICD on the BTD of RBPJ. While SHARP and ANK/MAM use different

strategies and target different key residues to interact with the CTD, clearly binding of SHARP

and ANK/MAM to the CTD is mutually exclusive (Figure 3A). Despite the absence of sequence

similarity between SHARP and the RAM domain of NICD, unexpectedly, SHARP interacts with

the BTD in a manner that is structurally virtually identical to how RAM, as well as the

corepressors FHL1 and RITA1, interacts with BTD (Figure 3B). As shown in Figure 3B, RAM

and other coregulators that bind BTD have a characteristic hydrophobic tetrapeptide sequence

(-ϕWϕP-) that is essential for binding BTD and serves as the linchpin for high affinity

interactions. SHARP is lacking this and other conserved elements that contribute to interactions

with BTD[31]. However, a structural alignment of SHARP with RAM and other BTD-binders

reveals some sequence conservation of hydrophobic residues that play important roles in

SHARP-RBPJ complex formation, including I2804, A2806, I2808, I2811, and P2812 (Figure

3B).

The corepressor Hairless is the major antagonist of Notch signaling in *Drosophila* and

directly binds the CTD of Su(H) (fly *CSL* ortholog) with high affinity in order to repress Notch

target gene transcription in flies[18,34,35]. While there is no sequence similarity between SHARP

and Hairless, these two corepressors have evolved to bind the same CTD interface on RBPJ

and Su(H) (Figure 3C). However, there are major structural differences in how SHARP and

Hairless form complexes with the CTD. In contrast to SHARP, which only induces a modest

conformational change in CTD, Hairless binding dramatically opens up the Ig domain of CTD,

interacting with residues that form the hydrophobic core of CTD[18]. This allows Hairless to bind

exclusively to the CTD with high affinity[34].

The CTD is structurally similar to the RHR-C domains of transcription factors, such as

NF-κB and NFAT[17]. Figure 4A shows the overlay of the CTD from RBPJ with the RHR-C

domain from NFAT. Importantly, the CTD deviates from canonical RHR-C domains by the

absence of β-strand a′ (βa′), which lies between the two β−sheets that compose the Ig domain

of a canonical RHR-C domain. Strikingly, SHARP binding to the CTD serves as a structural

surrogate for βa′, occupying this region in the complex structure (Figure 4A), which has

interesting implications for how binding sites have evolved on RBPJ and will be discussed

further below (Figure 4B).

Binding analysis of RBPJ and SHARP mutants

To gain further insights into RBPJ/SHARP complex formation, SHARP function, and to validate

our structural studies, we used a combination of assays, both *in vitro* and in cells, to analyze

structure-based mutants of RBPJ and SHARP. As shown in Figure 5 and Tables 2 and 3, we

used isothermal titration calorimetry (ITC) to quantitate binding between SHARP and RBPJ

mutants. SHARP binds RBPJ with high affinity (K_d~5nM, Figure 5A and Table 2), wherein

binding is enthalpically driven and entropically unfavorable, in accordance with the RBPID of

SHARP being an intrinsically disordered region[27]. Consistent with its side chain burying ~84 Å^2

in the SHARP-CTD complex, mutation of L2791 to alanine (L2791A) in SHARP resulted in

greater than 350 fold decrease in binding (ΔΔG°=3.4kcal/mol, Figure 5C and Table 2). The

SHARP mutants V2789A and Y2793A (Figure 5B,D and Table 2) also significantly reduced

binding to RBPJ by >10 fold and >20 fold, respectively, in agreement with these residues also

burying considerable amounts of surface area at the SHARP-CTD interface (V2789=54 Å^2 and

Y2793=118 Å^2). The side chain of SHARP residue I2811 is buried at the BTD-SHARP interface,

lying within a small hydrophobic pocket and burying a substantial 160 Å^2. Mutation of I2811 to

alanine (I2811A) results in more than a 60 fold reduction in binding (Figure 5F and Table 2).

Mutation of K2807, which makes electrostatic interactions with E259 of the BTD, to alanine

results in a much more modest effect on binding (~6 fold, Figure 5E and Table 2). Consistent

with SHARP binding independently to the BTD and CTD, which results in an avidity effect[27],

single alanine mutants are unable to completely abrogate binding. However, the SHARP double

mutant L2791A/I2811A, which targets mutations to key residues that interact with the BTD and

CTD, results in a complete loss of binding in our ITC studies (Table 2).

Similarly, we performed ITC binding studies with structure-based RBPJ mutants and

native SHARP. We performed differential scanning fluorimetry (DSF) to confirm that our RBPJ

mutants are correctly folded (Figure S2A); EMSA to show that our RBPJ mutants bind DNA

similarly to wild-type (WT) (Figure S2B); immunofluorescence microscopy to demonstrate that

our RBPJ mutants properly localize to the nucleus (Figure S2C); and coimmunoprecipitation

(CoIP) from cells to show that our RBPJ mutants still bind NICD (Figure S2D). However, we

were unable to purify and test by ITC some RBPJ mutants that target key residues in CTD-

SHARP complex, *e.g.* F468A, as these residues are buried within the hydrophobic core of the

CTD and are required for proper folding of RBPJ. Nonetheless, as shown in Figure 5I and Table

3, the mutation of L388, which buries ~103 Å^2 in the complex, to alanine results in an

approximately 70 fold reduction in binding. Whereas the CTD mutants L386A, L397A, and

L466A had only a minor effect on SHARP binding (Table 3), consistent with these residues

burying much less surface area at the CTD-SHARP interface (L386=39 $Å^2$, L397=3 $Å^2$, and

L466=44 $Å^2$). Because SHARP binds the BTD of RBPJ in a structurally similar manner to the

RAM domain of NICD, and the corepressors FHL1 and RITA1, we used a set of alanine mutants

(F261A, V263A, A284V, and Q333A) that we have previously characterized for RAM, FHL1,

L3MBTL3, and RITA1 binding to RBPJ (Figure 5G-H and Table 3)[12,13,19,36]. The BTD mutants

F261A and A284V, which are more centrally located in the SHARP-BTD interface, have a

stronger effect, significantly reducing binding by ~45 fold and ~50 fold, respectively (Figure 5G-

H and Table 3). V263A and Q333A, which significantly reduced RAM binding to RBPJ[36], but

target more peripheral interactions in the BTD-SHARP complex, only modestly affect binding

(Table 3). Interestingly, similar binding trends were observed when these BTD mutants were

previously tested for interactions with FHL1 and RITA1[13,19].

We also tested our SHARP mutants, and a subset of our RBPJ mutants, in cells using

CoIP of exogenously expressed RBPJ and SHARP RBPID proteins, and a mammalian two-

hybrid assay (Figure 6). Overall, we observed excellent correspondence between our ITC

binding studies and our cellular assays. As shown in Figure 6A, the CoIP of SHARP mutants

L2791A and I2811A with RBPJ is greatly reduced compared to wild-type SHARP, and the CoIP

of the double mutant L2791A/I2811A is scarcely detectable on the Western blot (Figure 6A, lane

9); whereas, the CoIP of SHARP mutants V2789A, Y2793A, K2807A with RBPJ is only

modestly affected (Figure 6A, lane 4-6), consistent with our ITC binding studies and the RBPJ-

SHARP structure. Similarly, the CoIP of the BTD mutant F261A and the CTD mutant L388A with

the RBPID of SHARP are greatly reduced compared to wild-type (Figure 6B, lanes 4-5), but the

double mutant F261A/L388A was unable to CoIP SHARP (Figure 6B, lane 6). In our mammalian

two-hybrid assays (Figure 6C-E), in which RBPJ and SHARP are fused to VP16 and Gal4,

respectively, SHARP mutants L2791A and I2811A have the strongest effect on reporter

expression, whereas V2798A, Y2793A, and K2807A only modestly affect reporter expression

(Figure 6C). As expected from the complex structure, the SHARP double mutant L279A/I2811A,

which does not bind RBPJ in our ITC assays, displayed no more than 1% expression from the

reporter compared to wild-type (Figure 6C). The RBPJ mutants F261A and L388A were

severely compromised for reporter activity (~1%), whereas no reporter activity was observed for

the F261A/L388A double mutant. As a control, all RBPJ mutants fused to the VP16

transactivation domain activated a luciferase reporter containing 12 CSL binding sites (Figure

6E) and bound DNA in an EMSA (Figure S2B) similar to wild-type.

RBPJ/SHARP interaction is required to repress Notch target genes

In order to investigate the contribution of the RBPJ/SHARP interaction in the regulation of Notch

target genes in cells, we analyzed the Notch target genes *Hes1* and *Hey1* in a mature T-cell line

(MT) that lacks Notch activity. First, to interfere with RBPJ/SHARP-mediated repression in MT

cells, we expressed the RBPID of SHARP as a GFP fusion protein either as wild-type (GFP-

SHARP/RBPID^{WT}) or the RBPJ-interacting defective L2791A/I2811A mutant (GFP-

SHARP/RBPID$^{LI/AA}$) (Figure 7A). Expression of GFP-SHARP/RBPID^{WT} leads to the upregulation,

i.e. derepression, of *Hes1* and *Hey1*, whereas GFP-SHARP/RBPID$^{LI/AA}$ has little to no effect on

Hes1 and Hey1 expression (Figure 7A). This suggests that GFP-SHARP/RBPID^{WT}, but not

GFP-SHARP/RBPID$^{LI/AA}$, effectively outcompetes endogenous SHARP for binding to RBPJ.

Next, we used CRISPR/Cas9 technology to deplete RBPJ (Figure 7B). Consistent with RBPJ

repressor function, we observed robust upregulation of *Hes1* and *Hey1* in the absence of RBPJ

(Figure 7B). Correspondingly, shRNA-mediated knockdown of RBPJ also resulted in the

upregulation of *Hes1* and *Hey1* (Figure 7C). Importantly, reintroduction of wild-type RBPJ

(RBPJ^{WT}) in the CRISPR/Cas9-mediated RBPJ depleted background rescues the repression of

Hes1 and *Hey1* Notch target gene expression, whereas the SHARP-interacting defective RBPJ

mutant F261A/L388A (RBPJ$^{FL/AA}$) does not (Figure 7D). Altogether, our data demonstrates that

the RBPJ/SHARP interaction is strongly required to repress transcription of Notch target genes

in cells.

Discussion

Notch is a highly conserved signaling pathway that ultimately results in changes in gene

expression, a process that is controlled by the transcription factor CSL (Figure 1A)[1,2]. CSL forms

an activation complex with the intracellular domain of the Notch receptor (NICD) and the

coactivator Mastermind (MAM) to activate transcription from Notch target genes (Figure 1B)[15,16].

This activity of CSL is required at all Notch target genes and is conserved in all metazoans. CSL

can also function as a transcriptional repressor, by interacting with corepressors such as

SHARP[8,9,27], L3MBTL3[12], FHL1[11,19], and RITA1[13,14] in mammals, and Hairless in

Drosophila[18,34,35]; however, its role as a repressor, in particular in mammals, is not well

understood. In order to begin to address this gap in our understanding, here we determine the

X-ray structure of the RBPJ-SHARP corepressor complex bound to DNA and use a multitude of

in vitro and cellular assays to characterize structure based RBPJ and SHARP mutants in order

to better understand CSL-corepressor function.

Together with previous studies, the complex structure provides a detailed mechanism for

how the corepressor SHARP interacts with the transcription factor RBPJ to repress

transcription. Prior to binding RBPJ, the RBPID of SHARP is intrinsically disordered[27], which

then forms a bipartite interaction with RBPJ. As shown in Figure 2, the N-terminal portion of the

RBPID assumes a β-hairpin motif that interacts with the CTD of RBPJ, resulting in a modest

conformational change in the CTD; the SHARP-CTD interaction is followed by a poorly

structured linker and then an extended region that binds across a nonpolar surface on the BTD.

Thus, SHARP forms a high affinity complex with RBPJ by interacting with two distant binding

surfaces on RBPJ, which individually are low affinity, but when tethered by the linker region

result in an avidity effect[27]. RBPJ-SHARP complex formation localizes the RRM and SPOC

domains of SHARP to Notch target genes (Figure 1C). In the case of SPOC, this recruits other

factors involved in transcriptional repression, such as NCoR/SMRT[20,24], CtIP (C-terminal binding

protein interacting protein)/CtBP (C-terminal binding protein)[23], and ETO (eight-twenty-one)[22],

as well as the KMT2D coactivator complex[24]. Whether the RRM domains of SHARP recruit Xist

lncRNA or other lncRNAs, such as SRA (steroid receptor coactivator)[37], to Notch target genes

remains to be determined

Analogous to SHARP, NICD also binds the BTD and CTD of RBPJ (Figure 1B and 3A);

however, the two complexes with RBPJ are mutually exclusive (Figure 3A) and the distribution

of binding energies for surfaces on RBPJ is distinct. In mammals, the RAM domain of NICD

forms a high affinity interaction with the BTD[30-33], comparable to the affinity of SHARP for

RBPJ[27], and is structurally very similar to how SHARP interacts with the BTD despite no

sequence similarity between the two proteins (Figure 3B). The ANK domain of NICD, however,

binds the CTD of RBPJ in a manner that is structurally very different than SHARP and the ANK-

CTD interaction is extremely weak, barely discernible in some binding assays[27,30,32,33]. MAM is

then required to stabilize the RBPJ-NICD-MAM ternary complex. The functional consequences

of these structural and binding differences between SHARP and NICD are currently unclear, but

it may be important for the dynamic association/dissociation of RBPJ-coregulator complexes

and the lifetimes of these complexes in the nucleus. Certainly, single molecule studies of wild-

type and mutant SHARP molecules will be important for addressing the dynamics of RBPJ-

coregulator complexes in live mammalian cells.

Our cellular studies clearly demonstrate that CRISPR/Cas9-mediated depletion of RBPJ

in MT cells results in derepression of the well-established Notch target genes *Hes1 and Hey1*

(Figure 7)[38]. Moreover, reconstitution of repression by wild-type RBPJ, but not a SHARP-binding

mutant, strongly suggests that SHARP is the primary corepressor that mediates repression via

RBPJ at Notch target genes in lymphocytes, and likely other cells and tissues, which is

consistent with previous *in vivo* studies of SHARP in mice[8,39,40]. These data are also consistent

with research in other experimental systems that show that loss of CSL results in transcriptional

derepression at some, but not all, Notch target genes[41,42]. Why RBPJ-SHARP corepressor

complexes are recruited to some Notch targets, but not others, remains an open question.

Clearly, the use of RBPJ mutants defined here that affect SHARP interactions, but leave NICD

interactions largely intact, will be instrumental in addressing this and other questions regarding

the function of RBPJ as a transcriptional repressor.

Classical models of transcriptional regulation in the Notch pathway posit that CSL is

constitutively bound to DNA, and that corepressors are replaced by NICD when signaling is

activated[43,44]. Given the comparable high affinities of both SHARP and NICD for RBPJ[27,30], as

well as the more modest affinity of CSL for DNA (~0.5-1.0 µM)[45,46], it is difficult to reconcile how

cofactor exchange on DNA, *i.e.* NICD replacing SHARP on RBPJ, might work at the molecular

level. The affinities rather suggest that preformed RBPJ-corepressor complexes and RBPJ-

NICD-MAM coactivator complexes are dynamically associating/dissociating with sites on the

DNA. This scenario is supported by recent data by Bray and colleagues investigating the DNA-

binding dynamics of Su(H) complexes with NICD and Hairless in Notch-ON and Notch-OFF

states in the salivary gland of *Drosophila melanogaster*[47].

In contrast to the CSL-NICD-MAM activator complex, CSL-corepressor interactions

appear to be less conserved across disparate organisms. For example, while SHARP, also

known as SPEN in *Drosophila* and DIN-1 in *C. elegans*, is conserved from nematodes to flies to

mammals[48], the RBPID of SHARP is only conserved in vertebrates[27]. Similarly, the corepressor

Hairless, which is the major antagonist of Notch signaling in *Drosophila*, is not conserved

outside of insects and crustaceans[49]. However, while the corepressors that bind CSL are not

strictly conserved across disparate organisms, interestingly, the corepressor binding sites on

CSL are conserved. For example, SHARP, as well as the corepressors RITA1, FHL1, and likely

L3MBTL3, binds to the BTD of RBPJ in a structurally similar manner (Figure 3B); and strikingly,

despite no sequence conservation between SHARP and Hairless, both proteins bind to the

same surface of the CTD of CSL (Figure 3C). Thus, it seems likely that other species-specific

transcriptional corepressors that bind CSL will be identified, but they will likely utilize the

aforementioned conserved binding surfaces on the BTD and CTD to interact with CSL.

In a broader context, CSL provides an interesting example for how ligand binding sites

evolve in proteins. CSL is composed of two types of domains – the NTD and CTD are

immunoglobulin domains that are structurally related to the rel homology region (RHR) of

proteins, such as NF-κB and NFAT, and the BTD contains a β-trefoil fold that is structurally

related to growth factors such as interleukins and FGF[17]. As shown in Figure 4A, the canonical

RHR-C fold contains a β-strand, βa', that lies in between β-strands βa and βg; however, the

CTD of CSL is missing βa' typically found in RHR-C domains. The ligand, in this case the

corepressor SHARP, serves as a structural surrogate binding precisely in the region where the

missing βa' strand would lie. Strikingly, this is the second example in CSL proteins where this

type of ligand evolution has occurred. The first example, as shown in Figure 4B, was uncovered

following the structure determination of the CSL-NICD-MAM activation complex[15], whereby it

was shown that the BTD of CSL is atypical, as it deviates from the canonical β-trefoil fold such

that it is missing two of the 12 strands of the consensus fold[17]. In this instance, the ligand, the

RAM domain of NICD, again serves as a structural surrogate, binding across the BTD exactly

where the two missing β-strands would normally lie.

Finally, because of its association with human disease, there have been wide-ranging

efforts to pharmacologically target the Notch pathway, but the majority of this work has focused

on reagents that blunt overactive Notch signaling, *e.g.* in T-cell acute lymphoblastic leukemia[4,5].

However, there are numerous human diseases, including certain types of cancer,

cardiovascular defects, and congenital syndromes that are associated with insufficient Notch

signaling[3]. Our structure-function studies of the RBPJ-SHARP corepressor complex provide a

roadmap for developing reagents that lead to the derepression of Notch target genes, which

have the potential to become new treatment options for diseases connected to deficiencies in

Notch signaling.

Methods

Protein expression and purification

The cloning, expression, and purification of *Mus musculus* core RBPJ, residues 53–474, and

SMT3-SHARP, residues 2776-2833, which corresponds to the RBPJ-interacting region, were

described previously[27]. An MBP-SHARP fusion protein was used to crystallize the

RBPJ/SHARP/DNA complex. SHARP, corresponding to residues 2776-2820, was cloned into

pMALX-E, which encodes maltose binding protein with the following surface entropy reduction

mutations D82A, K83A, E172A, N173A, and K239A to aide in crystallization. The MBP-SHARP

fusion construct was overexpressed in Tuner *E. coli* and cells were lysed by sonication. The

lysate was incubated with amylose resin and eluted with 10 mM maltose. The MBP-SHARP

fusion protein was further purified by size exclusion chromatography.

Crystallization and Data Collection

RBPJ/MBP-SHARP/DNA complexes were setup in a 1:1.1:1.1 molar ratio and screened for

crystallization conditions using an Art Robbins Phoenix Crystallization Robot at 4°C. The

RBPJ/MBP-SHARP/DNA complex crystallized in 100mM Bis-Tris pH 6.6, 100mM NaCl, 40%

PEG 400, and 200mM NDSB-256. Crystals were harvested, flash frozen in liquid nitrogen,

diffraction data were collected at the Advanced Photon Source, beamline 21-ID-F (LS-CAT).

The RBPJ/MBP-SHARP/DNA crystals nominally diffract to 2.8Å and belong to the spacegroup

P2$_1$ with unit cell dimensions a=54.5 Å, b=231.6 Å, c=90.3 Å, and β=99.88°.

Structure Determination, Model Building, and Refinement

Molecular replacement with Phaser[50] was used to determine the RBPJ/MBP-SHARP/DNA

complex structure using the following search models DNA (3BRG), CSL (3IAG), and MBP

(3OB4). Two RBPJ/MBP-SHARP/DNA complexes were identified in the asymmetric unit. Phenix

was used for the initial stages of refinement[51]. Manual model building was performed with

COOT[52]. TLS parameters were generated and the model was subsequently refined using

BUSTER[53] and validated with MolProbity[54]. The final RBPJ/MBP-SHARP/DNA model was

refined to an R$_{work}$ = 18% and R$_{free}$ = 23% with good overall geometry (see Table 1). PyMOL

(The PyMOL Molecular Graphics System, Version 2.0 Schrödinger, LLC) was used to generate

figures and perform structural overlays. The PISA server was used to analyze protein

interfaces[55].

Coordinates and structure factors have been deposited into the Protein Data Bank (XYZ1)

Isothermal titration calorimetry (ITC)

ITC experiments were performed using a Microcal VP-ITC micocalorimeter. For all binding

reactions, SMT3-SHARP (2776-2883) at ~100µM was placed in the syringe concentrations and

RBPJ (53-474) at ~10 µM was placed in the cell. Titrations consisted of an initial 1µl injection

followed by 39 7µl injections. ITC binding experiments were performed in 50 mM sodium

phosphate pH 6.5, 150 mM NaCl at 25°C. Samples were buffer matched using size-exclusion

chromatography or dialysis. The raw data were analyzed using ORIGIN and fit to a one site

binding model.

Cell culture and preparation of cell extracts

Mouse hybridoma mature T (MT) cell line was grown in Iscove's Modified Dulbecco Medium

(IMDM, Gibco) supplemented with 2% FCS, 0.3 mg/l peptone, 5 mg/l insulin, nonessential

aminoacids and penicillin/streptomycin. Cells were grown at 37°C with 5% CO_2. HeLa (ATCC:

CCL2), HEK-293 (ATCC: CRL1573), 293T and *Phoenix*[TM] packaging cells (*Orbigen*, Inc., San

Diego, CA, USA) were cultivated in Dulbecco's Modified Eagle Medium (DMEM, Gibco)

supplemented with 10% fetal calf serum (FCS) and penicillin/streptomycin.

DNA transfection

HEK-293 and HeLa cells were transfected using the Nanofectin transfection reagent (PAA)

according to the manufacturer's instructions.

Coimmunoprecipitation experiments

HEK-293 cells were transfected with the indicated constructs for expression of GFP- and Flag-

tagged wild-type and mutant proteins. 24 hours after transfection cells were lysed with 600 µl

CHAPS lysis buffer [10 mM 3-[(3-Cholamidopropyl)dimethylammonio]-1-propanesulfonate

hydrate (CHAPS, Merck), 50 mM Tris-HCl (pH 7.8), 150 mM NaCl, 5 mM NaF, 1 mM

Dithiothreitol (DTT, Merck), 0.5 mM Phenylmethanesulfonyl fluoride **(**PMSF, Merck) and 40 µl/ml

"Complete Mix" protease inhibitor cocktail (Roche)]. The extracts were incubated with 40 µl

agarose-conjugated anti-Flag antibody (M2, Sigma) at 4°C overnight. Precipitates were washed

6 to 8 times with CHAPS lysis buffer and finally resuspended in SDS-polyacrylamide gel loading

buffer. For Western blotting the proteins were resolved in SDS-polyacrylamide gels and

transferred electrophoretically at room temperature to PVDF membranes (Millipore) for 1 h at 50

mA using a Tris-glycine buffer system. The membranes were pre-blocked for 1 h in a solution of

3% milk powder in PBS-T (0.1% Tween 20 in PBS) before adding antibodies. The following

antibodies were used: anti-GFP (7.1/13.1, mouse monoclonal IgG, secondary antibody

peroxidase conjugated sheep anti-mouse IgG, NA931V, GE healthcare) or anti-Flag (M5,

Sigma; secondary antibody, NA931V, GE healthcare).

In vitro protein translation

The *in vitro* protein translations were performed using the TNT-assay (#L4610) from Promega

according to manufacturer´s instructions. Prior to EMSAs the *in vitro* translations of RBPJ (wt)

and mutant proteins were monitored by western blotting using an anti-Flag antibody (M5,

Sigma).

Electro Mobility Shift Assay (EMSA)

Reticulocyte lysates from *in vitro* translations were used for electromobility shift assays (EMSAs)

in a binding buffer consisting of 10 mM Tris-HCl (pH 7.5), 100 mM NaCl, 0.1 mM EDTA, 0.5 mM

DTT, and 4% glycerol. For binding reaction, 2 µg poly(dI-dC) (GE healthcare) and approximately

0.5 ng of ^{32}P-labelled oligonucleotides were added. The sequence of the double-stranded

oligonucleotide FO-233 (Supplementary Table S1) corresponds to the two RBP-J-binding sites

within the EBV TP-1 promoter. Super shifting of complexes was achieved by adding 1 µg of

anti-Flag (M5, Sigma) antibody. The reaction products were separated using 5% polyacrylamide

gels with 1x Tris-glycine-EDTA at room temperature. Gels were dried and exposed to X-ray

films (Kodak).

Fluorescence microscopy

HeLa cells were cultured on glass cover slips in a 25-well plate (Bibby Sterilin Ltd) at a density

of 10^5 cells per cm^2. After 16 h cells were transfected with 400 ng of expression plasmids using

the Nanofectin transfection reagent (see above). 24 h after transfection cells were rinsed with

PBS, fixed with 4% paraformaldehyde (PFA, Merck) in PBS (pH = 7.5). Specimens were

embedded in "ProLong© Gold antifade" reagent (Invitrogen) supplemented with 2-(4-

carbamimidoylphenyl)-1H-indol-6-carboximidamide (DAPI) and stored at 4 °C overnight.

Pictures were taken using a fluorescence microscope (IX71, Olympus) equipped with a digital

camera (C4742, Hamamatsu), and a 100-W mercury lamp (HBO 103W/2, Osram). The

following filter sets were used: Green, (EGFP) ex: HQ470/40, em: HQ525/50, blue (DAPI)

D360/50, em: D460/50.

Luciferase assay

HeLa cells were seeded in 48-well plates at a density of 20 x 10^4 cells. Transfection was

performed with Nanofectin reagent (see above) using 1 µg of reporter plasmid alone or together

with various amounts of expression plasmid (given in the corresponding figure legends). After

24 h luciferase activity was determined from at least six independent experiments with 20 µl of

cleared lysate in an LB 9501 luminometer (Berthold) by using the luciferase assay system from

Promega.

Infection of hybridoma mature T-cell line

5 x 10^6 *Phoenix*[TM] cells were seeded and 24 h later they were transfected with the plasmid DNA

of choice. Briefly, 20 μg of DNA were mixed with 860 μl of H_2O and 120 μl of 2 M $CaCl_2$ and

mixed by vortexing. The DNA solution was transferred dropwise to 1 ml of 2 x HBS buffer (50

mM HEPES pH 7.05, 10 mM KCl, 12 mM Glucose, 280 mM NaCl, 1.5 mM $NaHPO_4$) while

vortexing and the solution was incubated 20 min at room temperature. In the meantime, 25 μM

Chloroquine solution (Sigma-Aldrich) was added to the *Phoenix*[TM] cells (1 μl/ml) and the cells

were incubated for 10 min. The DNA solution was added to the cells and 12 h later the medium

was replaced. After 24 h of incubation, the medium containing the retroviral suspension was

filtered and 2 mg/ml Polybrene (Sigma-Aldrich) solution was added (1 μl/ml). Fresh medium was

added to the *Phoenix*[TM] cells that were maintained in culture for further infections. The retroviral

solution was used for spin infection of MT cells by centrifuging 45 min at 1800 rpm at 37°C. In

total, four spin infections were performed over two days. Positively infected cells were selected

with puromycine (Serva) or blasticidin (Gibco) and, eventually, GFP positivity was analyzed

using a BD FACS Calibur.

Generation of CRISPR/Cas9 depleted MT cells

CRISPR/Cas9 *Rbpj* depleted MT cells were generated as follows: 3 x 10^6 293T cells were

seeded and, after 24 h, transfected with 2.5 μg psPAX, 1 μg pMD2G and 3.33 μg of the desired

lentiCRISPR v2 vector using Lipofectamine 2000 Transfection Reagent (Invitrogen 11668-019)

accordingly to manifacturer's instructions. After at least 6 h of incubation at 37°C the medium

was replaced with fresh one and 48 h post-transfection the supernatant was filtered,

supplemented with polybrene and used for infection of MT cells. Positively infected cells were

selected with puromycin and dilutions were performed to establish single cell clones. Individual

clones were screened by Western blotting versus RBPJ.

shRNA knockdown

For the knockdown in MT cells, the pLKO.1 TRC1 shRNA library (SIGMA-ALDRICH) was used.

Transfection of 293T cells and infection and selection of MT cells was performed as previously

described. Sequence of the hairpins used in this study is indicated in Table S1.

Constructs

The expression plasmid pcDNA3-Flag-mNotch-1-IC (Flag-NICD) and the luciferase reporter

construct pGa981/6 (12 x CSL-RE-LUC) were previously described[14]. The Gal4-reporter

plasmid pFR-Luc (5 x Gal4-RE-LUC) was previously described[9]. The pMSCV-FLAG-hRBP-J

IRES Blasticidin was kindly provided by Dr. R. Liefke. All oligonucleotides used in this study are

listed in Table S1. PCR products were cloned in the pSC-A-amp/kan (Agilent Technologies

240205-5), digested with the selected restriction enzymes (New England Biolabs) and cloned

into the destination vectors accordingly to Table S2. All plasmids were analyzed by sequencing.

The pcDNA 3.1 Flag2 (Invitrogen) was commercially acquired while the pMY BioTip60 IRES-

GFP was previously described.

An engineered CRISPR/Cas9 resistant mouse RBPJ cDNA was synthetized at

GENEART/Life Technologies and inserted into the pcDNA3.1 Flag2 via NotI digestion. The

RBPJ mutants R218H, F261A, L388A and the F261A/L388A double mutant was generated by

site directed mutagenesis using the QuikChange II XL Site-Directed Mutagenesis Kit (Agilent

Technologies 200521-5) accordingly to manufacturer's instructions with primers listed in Table

S1 and using the pcDNA3.1 Flag-mRBPJ wt CRISPR/Cas9 resistant plasmid as template. The

mRBPJ wt and RBPJ F261A/L388A CRISPR/Cas9 resistant cDNAs were subcloned into the

pMY-Bio-IRES Blasticidin. The mouse RBPJ-VP16 expression plasmids (pcDNA3.1-Flag-2-

mRBPJ-VP16 wt, F261A, L388A and F261A/L388A) were constructed as follows: The stop

codon was deleted by a mRBP-J specific PCR fragment (mRBP_VP16_UP, mRBP-TAA_DO)

resulting in the pcDNA3.1-Flag-2-mRBPJΔstop constructs. A VP16 specific PCR-fragment

(VP16_XhoI_UP, VP16_XbaI_DO) was inserted into the corresponding sites of pcDNA3.1Flag-

2-mRBPJΔstop constructs resulting in the pcDNA3.1-Flag2-mRBPJ-VP16 plasmids. The Gal-4-

Mint (2776-2833) and the EGFP-Mint (2776-2833) constructs were generated by PCR assisted

cloning (Gal-Mint_F, Gal-Mint_R) into the BamHI/XbaI sites of pFa-CMV (Stratagene) and

pEGFP-C1 (Clontech), respectively. The mutated constructs (V2785A, Y2793A, K2807A,

I2811A, L2791A and I2811A/L2791A) were obtained by site directed mutagenesis.

The lentiCRISPR v2 was a gift from Dr. F. Zhang (Addgene plasmid # 52961). The

CRISPR/Cas9 guides were designed using the online tool available at http://crispr.mit.edu/. The

desired 5' overhangs were added and oligos were phosphorylated, annealed and ligated into

the lentiCRISPRv2 predigested with BsmBI.

RNA extraction, RT-PCR and qPCR

Total RNA was purified using Trizol reagent (Ambion, 15596018) accordingly to manufacturer's

instructions. 1 μg of RNA was reverse transcribed in cDNA using random hexamers and M-

MuLV reverse transcriptase (NEB). qPCRs were assembled with Absolute QPCR ROX Mix

(Thermo Scientific, AB-1139), gene-specific oligonucleotides and double-dye probes (see Table

S1) and analyzed using the StepOne Plus Real Time PCR system (Applied Biosystem). Data

were normalized to the housekeeping gene *glucuronidase* β (*GusB*).

Preparation of protein extracts and Western Blotting from MT cells

Whole Cell Extract (WCE) was prepared as follows. Briefly, cells were washed twice in PBS,

lysed in WCE buffer (20 mM Tris-HCl pH 8.0, 150 mM NaCl, 1% NP-40, 10% glycerol, 0.5 mM

Na_3VO_4, 10 mM NaF, 1 mM PMSF, 1 x Protease inhibitor cocktail mix) and incubated 20 min on

ice. After centrifuging 15 min at 13200 rpm at 4°C, the supernatant was recovered.

The Nuclear Extract (NE) from MT cells overexpressing the SHARP constructs was

prepared as follows. Briefly, cells were washed with PBS and resuspended in Buffer A (20 mM

Hepes pH 7.9 / 20 mM NaCl / 5 mM $MgCl_2$ / 10% glycerol / 0.2 mM PMSF) at the concentration

of 1 x 10^6 cells/ml. The cell suspension was incubated 20 min on ice and mixed by vortexing.

After 5 min centrifugation at 4000 rpm at 4°C, the pellet was washed twice in PBS and

resuspended in Buffer C (20 mM Hepes pH 7.9 / 300 mM NaCl / 0.2% NP-40 / 25% glycerol / 1

mM $MgCl_2$/ 0.2 mM PMSF / 1 x Protease inhibitor mix / 0.3 mM DTT) at the concentration of 1 x

10^6 nuclei/100 μl. After 20 min of incubation on ice, the nuclei suspension was centrifuged 5 min

at 13200 rpm at 4°C and the supernatant was collected.

The NE from MT cells overexpressing the RBPJ constructs was prepared as follows.

Briefly, 10 x 10^6 cells were washed with PBS and resuspended in 200 μl of Buffer 1 (10 mM

Hepes pH 7.9 / 10 mM KCl / 0.1 mM EDTA / 0.1 mM EGTA / 1 mM ßME, supplemented with

PMSF). The cell suspension was incubated 10 min on ice, 5 μl of 10% NP-40 were added and

mixed by vortexing. After 10 sec of centrifugation at 13000 rpm at 4°C, the nuclei pellet was

washed twice in 500 μl of Buffer 1 and resuspended in 100 μl of Buffer 2 (20 mM Hepes pH 7.9

/ 400 mM NaCl / 1 mM EDTA / 1 mM EGTA / 1 mM ßME, supplemented with PMSF). After 20

min of incubation on ice, the nuclei suspension was centrifuged 10 min at 13000 rpm at 4°C and

the supernatant was collected for further analysis.

Protein concentration was measured by Bradford assay (Sigma-Aldrich) and samples

were boiled after adding SDS-polyacrylamide gel loading buffer. Samples were resolved by

SDS-Page and analyzed by Western blotting using antibodies against GAPDH (abcam,

ab8245), GFP (Roche, 11814460001) or TBP (Santa Cruz, sc-273). Briefly, membranes were

blocked in 5% milk, 1x TBS, 0.1% Tween-20 (TBS-T) and primary antibodies were diluted in 5%

milk, TBS-T. After incubation over night at 4°C, membranes were washed in TBS-T, secondary

antibodies against mouse (Cell Signaling, #7076S) or rabbit (Cell Signaling, #7074S) were

diluted in 5% milk TBS-T and finally membranes were washed in TBS-T.

In the case of the RBPJ Western blotting the procedure was as follows. Briefly,

membranes were blocked in 5% milk, 1x TBS and the RBPJ antibody (Cosmo Bio Co. LTD,

2ZRBP2) was diluted 1:1000 in 5% BSA, 1x TBS, 0.3% NP40. After incubation over night at

4°C, membranes were washed three times 15 min each in 1x TBS / 0.5 M NaCl / 0.5% Triton X-

100 and the secondary antibody against rat (Jackson ImmunoResearch, 112-035-072) was

diluted 1:5000 in 5% BSA, 1x TBS, 0.3% NP-40. Membranes were washed three times 15 min

each in 1x TBS / 0.5 M NaCl / 0.5% Triton X-100. All membranes were finally incubated with

ECL solution and chemiluminescence was detected with a light sensitive film.

Acknowledgements

We thank Roswitha Rittelmann, Sabine Schirmer, and Thomas Schmidt-Wöll for excellent

technical assistance. This work was supported by National Institutes of Health Grants

5R01CA178974 (RAK). It was also supported by the collaborative research grant TRR81 and

the grant (BO 1639/9-1) by the DFG (German Research Foundation) and the Excellence Cluster

for Cardio Pulmonary System (ECCPS) in Giessen to TB and the University Medical Center

Giessen and Marburg (UKGM to BG). This work was further supported by German Research

Foundation (DFG) Grant SFB1074/A3 and by the BMBF (Federal Ministry of Education and

Research, research nucleus SyStAR) to FO. LP is supported by the DFG (GRK 2253 – HEIST).

This research used resources of the Advanced Photon Source, a U.S. Department of Energy

(DOE) Office of Science User Facility operated for the DOE Office of Science by Argonne

National Laboratory under Contract No. DE-AC02-06CH11357.

638 **Table 1: X-ray Data collection and refinement statistics.**

Data Collection Statistics

Resolution (Å)	50.0 – 2.8 (2.85 – 2.78)
Space Group	P2$_1$
Wavelength (Å)	0.97872
Unit Cell a, b, c (Å)	54.5, 231.6, 90.3
Unit Cell α, β, γ (°)	90.00, 99.9, 90.00
R$_{merge}$	0.084 (0.48)
I/σI	14.9 (2.3)
Completeness (%)	99.3 (97.6)
Redundancy	3.7 (3.4)

Refinement Statistics

R$_{work}$/R$_{free}$ (%)	19.4 / 22.8
Number of reflections	54,503
Number of atoms	13,392
Complexes/asymmetric unit	2
Wilson B/Mean B value (Å^2)	72.68 / 77.80
RMSD Bond Lengths (Å)	0.008
RMSD Bond Angles (°)	0.96
Ramachandran (favored/outliers)	96.34% / 0.37%

639 Highest resolution shell shown in parentheses.

Table 2: Calorimetric binding data for SHARP mutants and native RBPJ

SHARP	K (M^{-1})	K_d (uM)	$\Delta G°$ (kcal/mol)	$\Delta H°$ (kcal/mol)	$-T\Delta S°$ (kcal/mol)	$\Delta\Delta G°$ (kcal/mol)
WT	$2.0 \pm 0.5 \times 10^8$	0.005	-11.3 ± 0.1	-13.5 ± 0.4	2.2 ± 0.3	--
V2789A	$1.9 \pm 1.3 \times 10^7$	0.067	-9.9 ± 0.4	-11.0 ± 0.9	1.1 ± 1.2	1.4
L2791A	$5.8 \pm 1.0 \times 10^5$	1.75	-7.9 ± 0.1	-7.8 ± 0.8	-0.1 ± 0.7	3.4
Y2793A	$8.7 \pm 2.5 \times 10^6$	0.122	-9.4 ± 0.2	-9.0 ± 0.2	-0.4 ± 0.4	1.9
K2807A	$3.6 \pm 0.8 \times 10^7$	0.029	-10.3 ± 0.2	-13.3 ± 0.1	3.0 ± 0.1	1.0
I2811A	$3.3 \pm 0.6 \times 10^6$	0.316	-8.9 ± 0.1	-12.3 ± 0.2	3.4 ± 0.2	2.4
L2791A/I2811A	NBD	---	---	---	---	---

All experiments were performed at 25°C. NBD represents no binding detected.

Table 3: Calorimetric binding data for native SHARP and RBPJ mutants

	RBPJ	K (M^{-1})	K_d (uM)	$\Delta G°$ (kcal/mol)	$\Delta H°$ (kcal/mol)	$-T\Delta S°$ (kcal/mol)	$\Delta\Delta G°$ (kcal/mol)
	WT	$2.0 \pm 0.5 \times 10^8$	0.005	-11.3 ± 0.1	-13.5 ± 0.4	2.2 ± 0.3	--
BTD mutants	F261A	$4.5 \pm 0.9 \times 10^6$	0.23	-9.1 ± 0.1	-9.9 ± 0.5	0.8 ± 0.4	2.2
	V263A	$3.2 \pm 0.4 \times 10^8$	0.003	-11.6 ± 0.1	-18.3 ± 0.8	6.7 ± 0.9	-0.3
	A284V	$4.0 \pm 1.0 \times 10^6$	0.27	-9.0 ± 0.2	-12.9 ± 0.6	3.9 ± 0.7	2.3
	Q333A	$2.6 \pm 0.3 \times 10^8$	0.004	-11.4 ± 0.1	-16.8 ± 0.7	5.4 ± 0.7	-0.1
CTD mutants	L386A	$7.5 \pm 3.5 \times 10^7$	0.016	-10.7 ± 0.3	-13.2 ± 0.4	2.5 ± 0.7	0.6
	L388A	$2.6 \pm 0.5 \times 10^6$	0.391	-8.7 ± 0.1	-7.5 ± 0.6	-1.2 ± 0.7	2.6
	L397A	$4.2 \pm 0.8 \times 10^7$	0.025	-10.4 ± 0.1	-11.3 ± 0.2	0.9 ± 0.3	0.9
	L466A	$9.6 \pm 2.4 \times 10^7$	0.011	-10.9 ± 0.1	-13.3 ± 0.2	2.4 ± 0.2	0.4
BTD/CTD	F261A/L388A	NBD	---	---	---	---	---

All experiments were performed at 25°C. NBD represents no binding detected.

Table S1. Oligonucleotides used for plasmid construction, EMSAs, shRNAs, CRISPR/Cas9, and qPCR experiments.

Constructs		
Blasticidin NotI fw	5'-GCG GCC GCC CCC CCC CCC TAA CGT TA-3'	
Blasticidin SalI rev	5'-GTC GAC TTA GCC CTC CCA CAC ATA ACC AGA GG-3'	
Puro_EcoRI_F	5´-CTA TGA ATT CGG TGT GGA AAG TCC CCA GG-3´	
Puro_HindIII_R	5´-TTC AAA GCT TTC AGG CAC CGG GCT TGC G-3´	
mRBP_VP16_F	5´-GCC AGC CAG TCC AGG TTC C-3´	
mRBP-TAA_R	5´-ATC TCG AGG GAC ACC ACG TTG CTG TG G-3´	
VP16_XhoI_F	5´-ATC TCG AGA GCT TGG CCC CCC CG-3´	
VP16_XbaI_R	5´-CAT CTA GAT TAC CAC CGT ACT CGT CAA TTC CA AG-3´	
Gal_Mint_F	5´-ATG GAT CCG GCG CGG GCC TGC G-3´	
Gal_Mint_R	5´-CCT CTA GAC TAG AGC GGC CG-3´	
shRNA		
Rbpj_sh1	5´-GCA GAC TCA TTG GGC TAC ATT-3´	
Rbpj_sh4	5´-CCC TGT GCG TTT ATT GGA ATA-3´	
CRISPR/Cas9 guides		
sgRbpj_oligo_3	5´-GCT TGA CAG TCT GCC CGT AA-3´	
sgRbpj_oligo_4	5´-CGA CGA CGA GTC GGA AGG AG-3´	
Site directed mutagenesis		
mRBPJ_F261A_F	5´-GAA AGT GAG GGG GAA GAG GCC ACA GTT AGA GAT GGC TAC-3´	
mRBPJ_F261A_R	5´-GTA GCC ATC TCT AAC TGT GGC CTC TTC CCC CTC ACT TTC-3´	
mRBPJ_L388A_F	5´-CTG TCG TAG AAA GTC TTC AGG CGA ATG GCG GCG GGG ACG TAG-3´	
mRBPJ_L388A_R	5´-CTA CGT CCC CGC CGC CAT TCG CCT GAA GAC TTT CTA CGA CA-3´	
mRBPJ_R218H_F	5'-CGA AGG TGG CAC TGT TCA ATC ACC TTC GGT CCC AGA CAG-3'	
mRBPJ_R218H_R	5'-CTG TCT GGG ACC GAA GGT GAT TGA ACA GTG CCA CCT TCG-3'	
Mint-I2811A-F	5´-GAT CAG CCA GGC CCC TCC AG-3´	
Mint-I2811A-R	5´-CTG GAG GGG CCT GGC TGA TC-3´	
Mint-V2789A-F	5´- catcagaaggggtcgcgctgctgagctactc -3´	
Mint-V2789A-R	5´- gagtagctcagcagcgcgaccccttctgatg -3´	
Mint-Y2793A-F	5´- gtcgtgctgctgagcgcctcgggacagaagac -3´	
Mint-Y2793A-R	5´- gtcttctgtcccgaggcgctcagcagcacgac -3´	
Mint-K2807A-F	5´- agggatctggctgatcgcagcactgatccgctgg -3´	
Mint-K2807A-R	5´- ccagcggatcagtgctgcgatcagccagatccct -3´	
Mint-L2791A-F	5´- gtcccgagtagctcgccagcacgacccctt -3´	
Mint-L2791A-R	5´- aaggggtcgtgctggcgagctactcgggac -3´	

Mint-I2811A-F	5´- gcgctggctggaggggcctggctgatcttagc -3´	
Mint-I2811A-R	5´- gctaagatcagccaggcccctccagccagcgc -3´	
real time PCR		
	Mus musculus	*Probe*
GusB_F	5´-TGT GGG CAT TGT GCT ACC T-3´	25
GusB_R	5´-ATT TTT GTC CCG GCG AAC-3´	
Hes1_F	5´-ACA CCG GAC AAA CCA AAG AC-3´	99
Hes1_R	5´-CGC CTC TTC TCC ATG ATA GG-3´	
Hes1_2_F	5´-TGC CAG CTG ATA TAA TGG AGA A-3´	20
Hes1_2_R	5´-CCA TGA TAG GCT TTG ATG ACT TT-3´	
Tbp_F	5´-GGG GAG CTG TGA TGT GAA GT-3´	97
Tbp_R	5´-CCA GGA AAT AAT TCT GGC TCA T-3´	
Hey1_F	5´-CAT GAA GAG AGC TCA CCC AGA-3´	17
Hey1_R	5´-CGC CGA ACT CAA GTT TCC-3´	
EMSA		
FO233_F	5´-CCT GGA ACT ATT <u>TTC CCA CGG</u> TGC CCT TCC GCC CA TTT <u>TCC CAC</u> GAG TCG-3´ (RBP-J binding sites underlinded)	
FO233_R	5´-CTC GCG ACT C<u>GT GGG AAA</u> ATG GGC GGA AGG GCA C C <u>GTG GGA AA</u>A TAG TTC-3´ (RBP-J binding sites underlinded)	

Table S2. Constructs cloned by PCR.

Plasmid	Template	Primer Forward	Primer Reverse	Enzymes	Ligation
pMIGR1 pSV40-Puro	pLXSPHAMel18 ATG	Puro EcoRI F	Puro HindIII R	EcoRI/ HindIII	pMIGRI IRES-GFP
pcDNA3.1.Flag-2-mRBPJ	pcDNA3.1.Flag-2-mRBPJ	mRBP_VP16_F	mRBP-TAA_R	PshAI/XhoI	pcDNA3.1.Flag-2-mRBPJΔstop
pcDNA3.1.Flag-2-mRBPJΔstop	pFA-CMV	VP16_XhoI_F	VP16_XbaI_R	XhoI/XbaI	pcDNA3.1-Flag-2-mRBPJ-VP16
pFA-CMV	pcDNA3.1_HisC_MINT_2776-2833	Gal-Mint_F	Gal-Mint_R	BamHI/XbaI	pFA-CMV-Mint-2776-2833(wt)
pMY-Bio-IRES-Blasticidin	pMSCV-FLAG-hRBP-J Ires Blasticidin	Blasticidin NotI fw	Blasticidin SalI rev	NotI/SalI	pMY BioTip60 IRES-GFP

 Figure Legends

 Figure 1. <u>Overview of Notch signaling</u>**.** (A) Notch signaling occurs between neighboring cells, in

 which extracellular Notch-DSL interactions initiate signaling, resulting in the cleavage of Notch

 and release of its intracellular domain (NICD) from the cell membrane that subsequently transits

 to the nucleus. In the absence of Notch signaling, CSL can bind corepressors (CoR) to repress

 transcription from Notch target genes. Upon pathway activation, NICD and Mastermind (MAM)

 form a ternary complex with CSL that activates transcription from Notch target genes. (B)

 Structure of the CSL-NICD-MAM ternary complex bound to DNA (PDB ID: 2FO1). The structural

 core of CSL is composed of three domains: NTD (N-terminal domain), BTD (β-trefoil domain),

 and CTD (C-terminal domain), which are colored cyan, green, and orange, respectively. A β-

 strand that makes hydrogen-bonding interactions with all three domains is colored magenta.

 The RAM and ANK domains of NICD are colored yellow and blue, respectively. MAM and DNA

 are colored red and light pink/blue, respectively. (C) Domain schematics of CSL, NICD, and

 MAM are colored similarly to the structure. SHARP is a multidomain transcriptional coregulator

 that contains N-terminal RRM (RNA Recognition Motif) domains, multiple NLS (Nuclear

 Localization Sequence), a RID (Receptor Interaction Domain), a RBPID (RBPJ Interacting

 Domain), and a C-terminal SPOC (Spen Paralog and Ortholog C-terminal) domain.

 Figure 2. <u>X-ray structure of the RBPJ-SHARP corepressor complex bound to DNA</u>. (A) Ribbon

 diagram of RBPJ-SHARP-DNA complex with the NTD, BTD, and CTD of RBPJ colored cyan,

 green, and orange, respectively. SHARP is colored purple, and the DNA is colored light pink

 and light blue. Magnified views of the interaction of SHARP with the CTD (top) and BTD

 (bottom) of RBPJ. RBPJ is represented as a molecular surface with CTD and BTD residues that

 contact SHARP colored orange and green, respectively. The ribbon diagram of SHARP is

 colored purple with the side chains of SHARP that contact either the BTD or CTD shown as

sticks. SHARP residues that were mutated and tested for activity in this study are labeled. (B)

Open book representation of RBPJ-SHARP interfaces. (Left) Ribbon diagram with transparent

surface of the CTD-SHARP interface, and (right), ribbon diagram with transparent surface of the

BTD-SHARP interface. Side chains in RBPJ and SHARP that contribute to the interface are

shown. Key residues at the interface that were mutated and tested for activity in this study are

labeled. (C) Conformational changes in the CTD of RBPJ as a result of SHARP binding. (Left)

Magnified view of the CTD-SHARP complex with labeled secondary structural elements of the

CTD. The R438-E2786 salt bridge is also shown. (Middle) Structural overlay of RBPJ (bound)

from RBPJ-SHARP complex with an unbound structure of RBPJ (PDBID: 3IAG). The RBPJ

bound and unbound structures are colored orange and yellow, respectively. RBPJ residue

W441, which undergoes a large conformational change in the bound structure to accommodate

the R438-E2786 salt bridge, is shown in both structures. (Right) Ribbon diagram of the unbound

RBPJ structure 3IAG.

Figure 3. Comparison of coregulator binding sites on RBPJ. (A) Both SHARP and NICD bind

the BTD and CTD of RBPJ. (Left) RBPJ/SHARP/DNA complex structure with RBPJ represented

as a gray molecular surface, SHARP as a ribbon diagram colored purple, and the DNA as CPK

colored light blue and pink. The RBPJ residues that contact SHARP in the BTD and CTD are

colored green and orange, respectively. (Right) LAG-1/NICD/MAM/DNA complex structure

(PDBID: 2FO1) with LAG-1 (*C. elegans* CSL ortholog) represented as a gray molecular surface.

NICD is represented as a ribbon diagram with its RAM and ANK domains colored yellow and

blue, respectively. MAM is depicted as a ribbon diagram and colored red. The DNA is

represented in CPK and colored light blue and pink. RBPJ residues that contact RAM in the

BTD are colored green and residues that contact ANK in the CTD are colored orange. (B)

Structural alignment of SHARP and other coregulators that bind the BTD of CSL. (Left) The

BTD of RBPJ is represented as a green molecular surface. SHARP, the RAM domain from LIN-

12, RITA1 and FHL1 are colored purple, yellow, light pink, and gray, respectively. The

hydrophobic tetrapeptide (φWφP) is labeled. (Right) Sequence alignment of coregulators that

bind the BTD of CSL, including the RAM domains of Notch from mammals, *D. melanogaster*,

and *C. elegans*, and the corepressors RITA1, FHL1, and SHARP. The conserved φWφP, which

is absent in SHARP, is boxed in green. Other conserved regions in RAM that contribute to BTD

binding are highlighted, including the basic region (blue), the -HG- motif (orange), and the -GF-

motif (magenta). Structurally similar residues in SHARP that align with other BTD binders are

boxed and highlighted in gray. (C) Structural similarity of RBPJ-SHARP and Su(H)-Hairless

corepressor complexes. Despite no sequence similarity, both SHARP and Hairless bind the

CTD in a structurally analogous manner. RBPJ and Su(H) are represented as a gray molecular

surface with the SHARP and Hairless binding sites colored orange. SHARP and Hairless are

shown as ribbon diagrams and colored purple and yellow, respectively. The DNA is shown as

CPK spheres and colored light blue and light pink.

Figure 4. <u>Evolution of coregulator binding sites on CSL.</u> (A) Structural comparison of the CTD

of RBPJ with the RHR-C domain of NFAT. The ribbon diagram of the RBPJ-SHARP complex is

colored similarly as in Figure 2, whereas NFAT is colored green. The NFAT β-strand βa', which

is absent in the CTD of RBPJ, but occupied by SHARP in the complex structure, is labeled. (B)

Structural alignment of the BTD from RBPJ with a canonical β-trefoil fold from the ryanodine

receptor (RYR). A canonical BTD is composed of 12 β-strands, in which four β-strands are

arranged in a pseudo-threefold symmetrical arrangement. The atypical BTD of CSL is missing

two of the 12 β-strands that compose a canonical BTD. The RAM domain of NICD acts as a

structural surrogate, binding the region left absent by the missing β-strands. The BTD of CSL

and RYR are colored green and light pink, respectively. The two β-strands that are missing in

the CSL BTD fold are colored light blue and highlighted with red arrowheads. The RAM domain

of NICD is colored yellow.

Figure 5. ITC binding analysis of RBPJ and SHARP structure based mutants. Figure shows

representative thermograms (raw heat signal and nonlinear least squares fit to the integrated

data) for RBPJ binding to SHARP. Each experiment was performed at 25°C, with 40 titrations of

7 µl injections spaced 120 s apart. The dissociation constants (K_d) shown for each experiment

are indicated in Tables 2 and 3. (A) Native RBPJ and SHARP form a high affinity 1:1 complex

with a 5 nM dissociation constant. The SHARP mutants L2791A (C), Y2793A (D), and I2811A

(F) have the strongest effect on RBPJ-SHARP complex formation, whereas SHARP mutants

V2789A (B) and K2807A (E) only modestly affect RBPJ binding. The RBPJ BTD mutants F261A

(G) and A284V (H), and the CTD mutant L388A (I), strongly affect binding to SHARP.

Figure 6. Characterization of RBPJ and SHARP structure based mutants in cells. (A)

Coimmunoprecipitation (CoIP) of RBPJ with wild-type and mutant SHARP constructs. HEK293

cells were cotransfected with GFP-SHARP (2766-2833), which corresponds to the RBPJ-

interaction-domain of SHARP (RBPID), and Flag-RBPJ. Expression of GFP-SHARP (middle

panel) and Flag-RBPJ (bottom panel) proteins was verified by Western blotting. CoIPs were

performed 24h after transfection. The asterisk denotes the heavy chain of anti-Flag antibody

used for immunoprecipitation. (B) CoIP of SHARP with wild-type and mutant RBPJ constructs.

HEK293 cells were cotransfected with GFP-RBPJ and Flag-SHARP (RBPID). Expression of

Flag-SHARP (middle panel) and GFP-RBPJ (bottom panel) proteins was verified by Western

blotting. CoIPs were performed 24h after transfection. The asterisk denotes the heavy chain of

anti-Flag antibody used for immunoprecipitation. (C-D) Analysis of structure based SHARP and

RBPJ mutants using a mammalian two-hybrid assay. HeLa cells were cotransfected with the

indicated Gal4-SHARP (300ng) and RBPJ-VP16 (150ng) constructs together with the pFR-Luc

reporter (500ng), which contains five Gal4 DNA binding sites upstream of the luciferase gene.

Relative luciferase activity was determined after cotransfection of the pFR-Luc reporter

construct alone. Mean values and standard deviations are from at least six independent

experiments. (E) Control experiments showing RBPJ-VP16 mutants are competent to bind and

activate reporter similar to wild-type. HeLa cells were cotransfected with RBPJ-VP16 constructs

(150ng) and the 12xCSL-RE luciferase reporter (500ng), which contains 12 CSL DNA binding

sites upstream of the luciferase gene. Relative luciferase activity was determined after

cotransfection of the reporter construct alone. Mean values and standard deviations are from

four independent experiments [***P < 0.001, (ns) not significant, unpaired student´s *t*-test].

Figure 7. The RBPJ-SHARP interaction is required for repression of Notch target genes in cells.

(A) The wild-type SHARP RBPID (RBPID^{WT}), but not the RBPJ-interacting defective SHARP

RBPID (RBPID$^{LI/AA}$) mutant, causes upregulation of Notch target genes in mouse mature T (MT)

cells by outcompeting endogenous SHARP for RBPJ binding. MT cells were infected with

plasmids encoding GFP-tagged SHARP (2776-2833), either wild-type (GFP-SHARP/RBPID^{WT},

black bars) or the RBPJ-interacting defective mutant L2791A/I2811A (GFP-SHARP/RBPID$^{LI/AA}$,

gray bars), or an empty vector control (Control, white bars). (Left) Total RNA from MT cells

infected with plasmids encoding GFP-SHARP/RBPID^{WT}, GFP-SHARP/RBPID$^{LI/AA}$, or an empty

vector control, were reverse transcribed into cDNA and analyzed by qPCR using primers

specific for *Tbp*, *Hes1* or *Hey1*. Data were normalized to the housekeeping gene *GusB*

(glucuronidase β). Data shown represent the mean ± SD of triplicate experiments ([**] P < 0.01,

[***] P < 0.001, unpaired Student's *t*-test). (Right) Nuclear extracts (NE) were prepared from MT

cells infected with GFP-SHARP/RBPID^{WT}, GFP-SHARP/RBPID$^{LI/AA}$, or an empty vector control

and analyzed by Western blotting using a GFP antibody and with TBP used as a loading

control. (B) RBPJ is required to repress Notch target genes *Hes1* and *Hey1* in MT cells as

revealed by CRISPR/Cas9 depletion of RBPJ. (Left) Total RNA from wild-type (*Control*) or

RBPJ depleted (clones *sgRbpj 2-12* and *sgRbpj 2-14*) MT cells was reverse transcribed into

cDNA and analyzed by qPCR using primers specific for *Tbp*, *Hes1* or *Hey1*. Data were

normalized to the housekeeping gene *GusB* (*glucuronidase* β). Shown is the mean ± SD of

triplicate experiments ([*] P < 0.05, [**] P < 0.01, [***] P < 0.001, unpaired Student's *t*-test).

(Right) Whole cell extracts (WCE) were prepared from wild-type (*Control*) or RBPJ depleted

(clones *sgRbpj 2-12* and *sgRbpj 2-14*) MT cells and analyzed by Western blotting using an anti-

RBPJ antibody. GAPDH was used as loading control. (C) *Hes1* and *Hey1* Notch target genes

are upregulated upon shRNA-mediated *Rbpj* knockdown, but not with the *SCR* control shRNA.

(Left) Total RNA from MT cells infected with shRNAs targeting *Rbpj* (*Rbpj sh1* or *Rbpj sh4*) or

scrambled shRNA control (SCR) was reverse transcribed into cDNA and analyzed by qPCR

using primers specific for *Tbp*, *Hes1* or *Hey1*. Data were normalized to the housekeeping gene

GusB (*glucuronidase* β). Shown is the mean ± SD of quadruplicate experiments ([**] P < 0.01,

[***] P < 0.001, unpaired Student's *t*-test). (Right) WCE was prepared from MT cells infected

with shRNAs targeting *Rbpj* (*Rbpj sh1* or *Rbpj sh4*) or a scrambled shRNA control (*SCR*) and

analyzed by Western blotting using a RBPJ antibody. GAPDH was used as loading control. (D)

Expression of wild-type RBPJ (RBPJ^{WT}), but not the SHARP-interacting defective mutant

F261A/L388A (RBPJ$^{FL/AA}$), in the RBPJ-depleted background rescues the repression of Notch

target genes. (Left) Total RNA from *sgRbpj 2-12* MT cells infected with empty vector (Control),

RBPJ^{WT} or RBPJ$^{FL/AA}$ was reverse transcribed into cDNA and analyzed by qPCR using primers

specific for *Tbp*, *Hes1* or *Hey1*. Data were normalized to the housekeeping gene *GusB*

(*glucuronidase* β). Shown is the mean ± SD of three independent experiments measured twice

each ([*] P < 0.05, [**] P < 0.01, [***] P < 0.001, unpaired Student's *t*-test). (Right) NEs were

prepared from *sgRbpj 2-12* MT cells infected with empty vector (Control), RBPJ^{WT} or RBPJ$^{FL/AA}$

and analyzed by Western blotting using anti-RBPJ antibodies. Histone H3 was used as a

loading control.

Figure S1. <u>Composition and analysis of the asymmetric unit of the RBPJ/MBP-SHARP/DNA</u> <u>complex crystals</u>. (A) The asymmetric unit of the RBPJ/MBP-SHARP/DNA complex crystals contains two RBPJ/MBP-SHARP/DNA complexes. RBPJ, MBP, and SHARP are shown as ribbon diagrams and colored green, cyan, and purple, respectively. The DNA is colored gray. (B) ITC thermogram showing the affinity of MBP-SHARP for RBPJ. (C-D) Structural overlays of the two complexes contained within the asymmetric unit of the crystals. (C) Structural alignment overall all cα atoms. (D) Resulting structural overlay when only RBPJ cα atoms are aligned. (E) Resulting structural overlay when only MBP cα atoms are aligned. (F) Resulting structural overlay when only SHARP cα atoms are aligned. Red arrows highlight the large conformational differences in the poorly ordered linkers of SHARP that lie between the regions that bind the BTD and CTD of RBPJ.

Figure S2. <u>Control studies of the structure based RBPJ mutants.</u> (A) Differential scanning fluorimetry (DSF) was used to analyze the melting temperatures (Tm) of wild-type and mutant RBPJ proteins. Figure shows melt curve and derivative plot. Tm determined for wild-type and mutant RBPJ proteins is the mean ± SD of experiments performed in triplicate.   (B) Upper panel, analysis of DNA binding by EMSA of wild-type RBPJ and select RBPJ mutants. Oligomeric duplex DNA probe used in EMSA 5´-CCTGGAACTATT**TTCCCACG**GTGCCCTTCCGCCCATT**TTCCCACG**AGTCG-3´ with RBPJ binding sites in bold and underlined. The DNA interaction bands are labeled A and B. The asterisk denotes a nonspecific background band. Addition of α-Flag antibodies lead to supershifted Flag-RBPJ/DNA complexes (complexes a and b) in the EMSA, demonstrating specificity. Lower panel,N-terminally tagged Flag-RBPJ fusion proteins were translated *in vitro* with the TNT T7 coupled Reticulocyte Lysate System Kit (Promega) and relative protein levels

were analyzed by Western blotting. Flag-RBPJ/DNA complexes, labeled a and b, were

supershifted in the EMSA using α-Flag antibodies, demonstrating specificity. (C) Subcellular

localization of RBPJ proteins after expression in HeLa cells. HeLa cells were transiently

transfected with 0.3 µg of the respective plasmids GFP-RBPJ: WT, F261A, L388A, and

F261A/L388A. After 24 h cells were fixed and the images were taken 24 h after transfection with

DAPI staining under the fluorescence microscope using a 63x lens. The superimposition (right

column) of the DAPI signal (center) with the GFP signal (left) illustrates the predominantly

nuclear localization of the RBPJ proteins. (D) NICD binding of wild-type and mutant RBPJ

proteins from cells. HEK293 cells were transfected with the plasmids GFP-RBPJ^{WT}, GFP-

RBPJ^{F261A} GFP-RBPJ^{L388A}, or GFP-RBPJ$^{F261A/L388A}$ plasmids together with with Flag-NICD (lanes

2-6). Immunoprecipitation was performed with the α-Flag antibody agarose beads and detected

by Western blotting using an α-GFP antibody. The expression of the Flag-tagged NICD protein

(lower blot) could be detected using an α-Flag antibody. The asterisk in the upper blot marks the

heavy chain of the antibody used for immunoprecipitation.

References

1   Kovall, R. A., Gebelein, B., Sprinzak, D. & Kopan, R. The Canonical Notch Signaling Pathway:
Structural and Biochemical Insights into Shape, Sugar, and Force. *Dev Cell* **41**, 228-241,
doi:10.1016/j.devcel.2017.04.001 (2017).
2   Bray, S. J. Notch signalling in context. *Nat Rev Mol Cell Biol* **17**, 722-735,
doi:10.1038/nrm.2016.94 (2016).
3   Siebel, C. & Lendahl, U. Notch Signaling in Development, Tissue Homeostasis, and Disease.
Physiological reviews **97**, 1235-1294, doi:10.1152/physrev.00005.2017 (2017).
4   Braune, E. B. & Lendahl, U. Notch -- a goldilocks signaling pathway in disease and cancer
therapy. *Discovery medicine* **21**, 189-196 (2016).
5   Ntziachristos, P., Lim, J. S., Sage, J. & Aifantis, I. From fly wings to targeted cancer therapies: a
centennial for notch signaling. *Cancer cell* **25**, 318-334, doi:10.1016/j.ccr.2014.02.018 (2014).
6   Kovall, R. A. & Blacklow, S. C. Mechanistic insights into Notch receptor signaling from structural
and biochemical studies. *Curr Top Dev Biol* **92**, 31-71, doi:S0070-2153(10)92002-4 [pii]

10.1016/S0070-2153(10)92002-4 (2010).
7   Borggrefe, T. & Oswald, F. The Notch signaling pathway: transcriptional regulation at Notch
target genes. *Cell Mol Life Sci* **66**, 1631-1646 (2009).

8   Kuroda, K. *et al.* Regulation of marginal zone B cell development by MINT, a suppressor of
Notch/RBP-J signaling pathway. *Immunity* **18**, 301-312 (2003).
9   Oswald, F. *et al.* SHARP is a novel component of the Notch/RBP-Jkappa signalling pathway. *Embo*
J **21**, 5417-5426 (2002).
10   Maier, D. Hairless: the ignored antagonist of the Notch signalling pathway. *Hereditas* **143**, 212-
221, doi:10.1111/j.2007.0018-0661.01971.x (2006).
11   Taniguchi, Y., Furukawa, T., Tun, T., Han, H. & Honjo, T. LIM protein KyoT2 negatively regulates
transcription by association with the RBP-J DNA-binding protein. *Mol Cell Biol* **18**, 644-654
(1998).
12   Xu, T. *et al.* RBPJ/CBF1 interacts with L3MBTL3/MBT1 to promote repression of Notch signaling
via histone demethylase KDM1A/LSD1. *EMBO J* **36**, 3232-3249, doi:10.15252/embj.201796525
(2017).
13   Tabaja, N., Yuan, Z., Oswald, F. & Kovall, R. A. Structure-function analysis of RBP-J-interacting
and tubulin-associated (RITA) reveals regions critical for repression of Notch target genes. *J Biol*
Chem **292**, 10549-10563, doi:10.1074/jbc.M117.791707 (2017).
14   Wacker, S. A. *et al.* RITA, a novel modulator of Notch signalling, acts via nuclear export of RBP-J.
Embo J **30**, 43-56, doi:emboj2010289 [pii]

10.1038/emboj.2010.289 (2011).
15   Wilson, J. J. & Kovall, R. A. Crystal structure of the CSL-Notch-Mastermind ternary complex
bound to DNA. *Cell* **124**, 985-996, doi:10.1016/j.cell.2006.01.035 (2006).
16   Nam, Y., Sliz, P., Song, L., Aster, J. C. & Blacklow, S. C. Structural basis for cooperativity in
recruitment of MAML coactivators to Notch transcription complexes. *Cell* **124**, 973-983,
doi:10.1016/j.cell.2005.12.037 (2006).
17   Kovall, R. A. & Hendrickson, W. A. Crystal structure of the nuclear effector of Notch signaling,
CSL, bound to DNA. *EMBO J* **23**, 3441-3451, doi:10.1038/sj.emboj.7600349 (2004).
18   Yuan, Z. *et al.* Structure and Function of the Su(H)-Hairless Repressor Complex, the Major
Antagonist of Notch Signaling in Drosophila melanogaster. *PLoS Biol* **14**, e1002509,
doi:10.1371/journal.pbio.1002509 (2016).
19   Collins, K. J., Yuan, Z. & Kovall, R. A. Structure and function of the CSL-KyoT2 corepressor
complex: a negative regulator of Notch signaling. *Structure* **22**, 70-81,
doi:10.1016/j.str.2013.10.010 (2014).
20   Shi, Y. *et al.* Sharp, an inducible cofactor that integrates nuclear receptor repression and
activation. *Genes Dev* **15**, 1140-1151, doi:10.1101/gad.871201 (2001).
21   Newberry, E. P., Latifi, T. & Towler, D. A. The RRM domain of MINT, a novel Msx2 binding
protein, recognizes and regulates the rat osteocalcin promoter. *Biochemistry* **38**, 10678-10690,
doi:10.1021/bi990967j (1999).
22   Salat, D., Liefke, R., Wiedenmann, J., Borggrefe, T. & Oswald, F. ETO, but not leukemogenic
fusion protein AML1/ETO, augments RBP-Jkappa/SHARP-mediated repression of notch target
genes. *Mol Cell Biol* **28**, 3502-3512, doi:10.1128/MCB.01966-07 (2008).
23   Oswald, F. *et al.* RBP-Jkappa/SHARP recruits CtIP/CtBP corepressors to silence Notch target
genes. *Mol Cell Biol* **25**, 10379-10390, doi:10.1128/MCB.25.23.10379-10390.2005 (2005).
24   Oswald, F. *et al.* A phospho-dependent mechanism involving NCoR and KMT2D controls a
permissive chromatin state at Notch target genes. *Nucleic Acids Res* **44**, 4703-4720 (2016).
25   McHugh, C. A. *et al.* The Xist lncRNA interacts directly with SHARP to silence transcription
through HDAC3. *Nature* **521**, 232-236, doi:10.1038/nature14443 (2015).
26   Chu, C. *et al.* Systematic discovery of Xist RNA binding proteins. *Cell* **161**, 404-416,
doi:10.1016/j.cell.2015.03.025 (2015).

27   VanderWielen, B. D., Yuan, Z., Friedmann, D. R. & Kovall, R. A. Transcriptional repression in the
Notch pathway: thermodynamic characterization of CSL-MINT (Msx2-interacting nuclear target
protein) complexes. *J Biol Chem* **286**, 14892-14902, doi:10.1074/jbc.M110.181156 (2011).
28   Moon, A. F., Mueller, G. A., Zhong, X. & Pedersen, L. C. A synergistic approach to protein
crystallization: combination of a fixed-arm carrier with surface entropy reduction. *Protein Sci* **19**,
901-913, doi:10.1002/pro.368 (2010).
29   Choi, S. H. *et al.* Conformational locking upon cooperative assembly of notch transcription
complexes. *Structure* **20**, 340-349, doi:10.1016/j.str.2011.12.011 (2012).
30   Friedmann, D. R., Wilson, J. J. & Kovall, R. A. RAM-induced allostery facilitates assembly of a
notch pathway active transcription complex. *J Biol Chem* **283**, 14781-14791, doi:M709501200
[pii]

10.1074/jbc.M709501200 (2008).
31   Johnson, S. E., Ilagan, M. X., Kopan, R. & Barrick, D. Thermodynamic analysis of the CSL x Notch
interaction: distribution of binding energy of the Notch RAM region to the CSL beta-trefoil
domain and the mode of competition with the viral transactivator EBNA2. *J Biol Chem* **285**,
6681-6692, doi:10.1074/jbc.M109.019968 (2010).
32   Del Bianco, C., Aster, J. C. & Blacklow, S. C. Mutational and energetic studies of Notch 1
transcription complexes. *J Mol Biol* **376**, 131-140, doi:S0022-2836(07)01560-4 [pii]

10.1016/j.jmb.2007.11.061 (2008).
33   Lubman, O. Y., Ilagan, M. X., Kopan, R. & Barrick, D. Quantitative dissection of the Notch:CSL
interaction: insights into the Notch-mediated transcriptional switch. *J Mol Biol* **365**, 577-589,
doi:10.1016/j.jmb.2006.09.071 (2007).
34   Maier, D. *et al.* Structural and functional analysis of the repressor complex in the Notch signaling
pathway of Drosophila melanogaster. *Mol Biol Cell* **22**, 3242-3252, doi:mbc.E11-05-0420 [pii]

10.1091/mbc.E11-05-0420 (2011).
35   Kurth, P., Preiss, A., Kovall, R. A. & Maier, D. Molecular analysis of the notch repressor-complex
in Drosophila: characterization of potential hairless binding sites on suppressor of hairless. *PLoS*
One **6**, e27986, doi:10.1371/journal.pone.0027986

PONE-D-11-18755 [pii] (2011).
36   Yuan, Z., Friedmann, D. R., VanderWielen, B. D., Collins, K. J. & Kovall, R. A. Characterization of
CSL (CBF-1, Su(H), Lag-1) mutants reveals differences in signaling mediated by Notch1 and
Notch2. *J Biol Chem* **287**, 34904-34916, doi:M112.403287 [pii]

10.1074/jbc.M112.403287 (2012).
37   Jung, C., Mittler, G., Oswald, F. & Borggrefe, T. RNA helicase Ddx5 and the noncoding RNA SRA
act as coactivators in the Notch signaling pathway. *Biochimica et biophysica acta* **1833**, 1180-
1189, doi:10.1016/j.bbamcr.2013.01.032 (2013).
38   Bray, S. & Bernard, F. Notch targets and their regulation. *Curr Top Dev Biol* **92**, 253-275,
doi:S0070-2153(10)92008-5 [pii]

10.1016/S0070-2153(10)92008-5 (2010).
39   Surendran, K. *et al.* The contribution of Notch1 to nephron segmentation in the developing
kidney is revealed in a sensitized Notch2 background and can be augmented by reducing Mint
dosage. *Developmental biology* **337**, 386-395, doi:10.1016/j.ydbio.2009.11.017 (2010).

40 Tsuji, M., Shinkura, R., Kuroda, K., Yabe, D. & Honjo, T. Msx2-interacting nuclear target protein
(Mint) deficiency reveals negative regulation of early thymocyte differentiation by Notch/RBP-J
signaling. *Proc Natl Acad Sci USA* **104**, 1610-1615, doi:10.1073/pnas.0610520104 (2007).
41 Chan, S. K. K. *et al.* Role of co-repressor genomic landscapes in shaping the Notch response. *PLoS*
genetics **13**, e1007096, doi:10.1371/journal.pgen.1007096 (2017).
42 Castel, D. *et al.* Dynamic binding of RBPJ is determined by Notch signaling status. *Genes Dev* **27**,
1059-1071, doi:10.1101/gad.211912.112 (2013).
43 Kao, H. Y. *et al.* A histone deacetylase corepressor complex regulates the Notch signal
transduction pathway. *Genes Dev* **12**, 2269-2277 (1998).
44 Hsieh, J. J. & Hayward, S. D. Masking of the CBF1/RBPJ kappa transcriptional repression domain
by Epstein-Barr virus EBNA2. *Science* **268**, 560-563 (1995).
45 Torella, R. *et al.* A combination of computational and experimental approaches identifies DNA
sequence constraints associated with target site binding specificity of the transcription factor
CSL. *Nucleic Acids Res* **42**, 10550-10563, doi:10.1093/nar/gku730 (2014).
46 Friedmann, D. R. & Kovall, R. A. Thermodynamic and structural insights into CSL-DNA complexes.
Protein Sci **19**, 34-46, doi:10.1002/pro.280 (2010).
47 Gomez-Lamarca, M. J. *et al.* Activation of the Notch Signaling Pathway In Vivo Elicits Changes in
CSL Nuclear Dynamics. *Dev Cell* **44**, 611-623 e617, doi:10.1016/j.devcel.2018.01.020 (2018).
48 Ariyoshi, M. & Schwabe, J. W. A conserved structural motif reveals the essential transcriptional
repression function of Spen proteins and their role in developmental signaling. *Genes Dev* **17**,
1909-1920 (2003).
49 Zehender, A. *et al.* Conservation of the Notch antagonist Hairless in arthropods: functional
analysis of the crustacean Daphnia pulex Hairless gene. *Development genes and evolution* **227**,
339-353, doi:10.1007/s00427-017-0593-4 (2017).
50 McCoy, A. J. *et al.* Phaser crystallographic software. *Journal of applied crystallography* **40**, 658-
674, doi:10.1107/S0021889807021206 (2007).
51 Adams, P. D. *et al.* PHENIX: a comprehensive Python-based system for macromolecular structure
solution. *Acta crystallographica. Section D, Biological crystallography* **66**, 213-221,
doi:10.1107/S0907444909052925 (2010).
52 Emsley, P. & Cowtan, K. Coot: model-building tools for molecular graphics. *Acta*
crystallographica. Section D, Biological crystallography **60**, 2126-2132 (2004).
53 Smart, O. S. *et al.* Exploiting structure similarity in refinement: automated NCS and target-
structure restraints in BUSTER. *Acta crystallographica. Section D, Biological crystallography* **68**,
368-380, doi:10.1107/S0907444911056058 (2012).
54 Davis, I. W. *et al.* MolProbity: all-atom contacts and structure validation for proteins and nucleic
acids. *Nucleic Acids Res* **35**, W375-383 (2007).
55 Krissinel, E. & Henrick, K. Inference of macromolecular assemblies from crystalline state. *J Mol*
Biol **372**, 774-797 (2007).

1014

1015

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7