Quantifying cell densities and biovolumes of phytoplankton communities and

functional groups using scanning flow cytometry, machine learning and

unsupervised clustering

Mridul K. Thomas[1]*, [#a], Simone Fontana[1], [#b], Marta Reyes[1] & Francesco Pomati[1]

[1] Department of Aquatic Ecology, Eawag: Swiss Federal Institute of Aquatic Science

and Technology, Dübendorf, Switzerland

[#a] Current address: Centre for Ocean Life, DTU Aqua, Technical University of

Denmark, Lyngby, Denmark

[#b] Current address: Biodiversity and Conservation Biology, Swiss Federal Research

Institute WSL, Birmensdorf, Switzerland

11
12
13 *Corresponding author.
14 Email: mrit@dtu.dk

15
16 Running head: Quantifying phytoplankton using SFCM

17
18 Key words: traits, phytoplankton, cell size, flow cytometry, automated monitoring

Abstract

Scanning flow cytometry (SFCM) is characterized by the measurement of time-resolved pulses of fluorescence and scattering, enabling the high-throughput quantification of phytoplankton morphology and pigmentation. Quantifying variation at the single cell and colony level improves our ability to understand dynamics in natural communities. Automated high-frequency monitoring of these communities is presently limited by the absence of repeatable, rapid protocols to analyse SFCM datasets, where images of individual particles are not available. Here we demonstrate a repeatable, semi-automated method to (1) rapidly clean SFCM data from a phytoplankton community by removing signals that do not belong to live phytoplankton cells, (2) classify individual cells into trait clusters that correspond to functional groups, and (3) quantify the biovolumes of individual cells, the total biovolumes of the whole community and the total biovolumes of the major functional groups. Our method involves the development of training datasets using lab cultures, the use of an unsupervised clustering algorithm to identify trait clusters, and machine learning tools (random forests) to (1) evaluate variable importance, (2) classify data points, and (3) estimate biovolumes of individual cells. We provide example datasets and code for our analytical approach that can be adapted for analysis of datasets from other flow cytometers or scanning flow cytometers.

Introduction

Flow cytometry (FCM) has enabled the monitoring of natural microbial communities by capturing point estimates of cellular characteristics and images, respectively ([1–7]. Developed more recently, scanning flow cytometry (SFCM) records time-resolved pulses of fluorescence and scattering for every cell [4,8–10]. The fluorescence and scattering pulses are summarized using parameters that characterise changes in morphology and pigmentation over the length of a cell. This vast amount of individual-level information can be used to quantify the distributions of important cellular *traits* within communities. High-throughput quantification of traits governing organism-environment interactions [11] would increase our ability to understand ecological and evolutionary changes in microbial communities. However, the utility of traditional FCM and scanning flow cytometry (SFCM) in monitoring natural communities has been limited by a lack of protocols that allow us to automate 1) cleaning of SFCM data by removing signals that do not belong to living cells, 2) classifying individual cells into functional groups, and 3) quantifying the biovolumes of individual cells, and the total biovolumes of the whole community and major functional groups (we follow convention by referring to cyanobacteria, diatoms, etc. as 'functional groups', but these groups are defined taxonomically). The first two goals are a challenge in FCM and SFCM analyses because validating results from an analysis is difficult in the absence of images (which are available only in imaging flow cytometers, IFCM, for which high-throughput analytical methods are available e.g. [6,12]). This paper demonstrates a protocol to achieve these three goals in FCM or SFCM datasets.

Cleaning datasets (i.e. remove signals that are not from live cells) and identifying functional groups in FCM has traditionally been done manually. This has involved

visually identifying clusters of points with similar trait values using small numbers of

bivariate plots. These manually-identified clusters are used to separate

measurements of live cells from other signals ('gating'), and to identify distinct cell

types for further analyses. However, a manual sample-by-sample approach is not

practical when large numbers of samples are measured, such as in high-frequency

monitoring efforts or large experiments. Additionally, the large number of variables of

unknown importance captured by SFCM makes the choice of variables for plotting

difficult, and the results sensitive to these choices. Previous efforts to avoid these

problem in FCM and SFCM have involved (i) focusing on very small numbers of

variables, (ii) the use of linear thresholds for separation of signal types, which

neglects the possibility that thresholds may be nonlinear, in single or multiple

dimensions, or more recently (iii) the use of a variety of clustering algorithms to

identify groups of points that are similar to each other in multiple dimensions

[9,10,13–17]. The use of clustering algorithms reduces subjectivity associated with

cluster identification and makes analyses repeatable. We have adapted this

approach to scale easily across large SFCM datasets using limited computing

resources.

Estimating the biovolume of cells and colonies is an important goal in the

phytoplankton, which are responsible for nearly half of global primary production

[18]. Cell biovolume is also a major determinant of how individuals interact with their

abiotic and biotic environment. Called a 'master trait', biovolume influences nutrient-

uptake rates, nutrient quotas, predator avoidance, and low-light performance [19–

21]. Additionally, distributions of biovolume can - in combination with cell density

(typically estimated by flow cytometers) - provide us with accurate estimates of total

phytoplankton community biovolume (highly correlated with biomass), the most

important algal parameter characterizing water quality. Despite its importance,

quantifying biovolumes using FCM and SFCM remains a challenge because of the

complex shapes that phytoplankton take [22]. Present approaches to estimating

biovolume involve the use of a single scattering channel and are based on calibrated

relationships with beads [9,23]; they therefore assume that phytoplankton cells

possess similar scattering properties to these beads. However, scattering properties

vary between phytoplankton taxa based on differences in cell wall, vacuoles, and

internal cellular structure, necessitating the development of better methods for

biovolume estimation.

Identifying the functional groups that phytoplankton belong to is fundamental to

understanding ecological patterns because members of these groups share

similarities in biochemistry, edibility, toxicity and roles in biogeochemical cycles

[21,24]. Although we follow convention by referring to 'functional groups', these

groups are defined taxonomically (e.g. cyanobacteria, diatoms, chrysophytes, green

algae, etc.), but at different levels in the taxonomic hierarchy (phylum Cyanobacteria

as opposed to class Bacillariophyceae). The term 'functional group' retains its

usefulness because the groups are paraphyletic: some members of the

clades/taxonomic groups are no longer phytoplanktonic (e.g. plants exist within the

clade of the green algae). Identification of functional groups by SFCM is possible

because groups differ in their autofluorescent photosynthetic pigments [24], leading

to differences in cell absorption and emission profiles [25].

The protocol that we present here (outlined in Fig. 1) quantifies the cell densities and

biovolumes of a phytoplankton communities and their major constituent functional

groups. It relies on three tools: (i) multiple training datasets that we generated, (ii) an

unsupervised clustering algorithm (flowPeaks [26]), and (iii) a machine learning

125   algorithm (random forests [27]). These tools are used for three distinct roles: (1)

126   identifying important variables for clustering, (2) classifying points into clusters, and

127   (3) estimating cell biovolumes.

Our advance consists in coupling these together in a manner that allows for

processing and analysis of large FCM and SFCM datasets without the use of

specialised computing resources such as high performance computing facilities.

Unsupervised clustering algorithms identify groups of points that are similar to each

other in multiple dimensions [28,29]. Here we use one such algorithm (flowPeaks

[26]) that has previously been shown to provide reasonable results when applied to

phytoplankton populations [17]. Random forests are a machine learning tool

composed of ensembles of decision trees that can be used to identify features that

separate groups in high-dimensional data. Each component decision tree uses a

random subset of 66% of the dataset, and a randomly selected subset of variables

that are assessed for correlation strength at each node. By aggregating predictions

from all decision trees, random forests limit the overfitting problems associated with

the use of individual decision trees [27]. No assumptions of linearity are required,

with the algorithm capable of identifying curved boundaries in multiple dimensions.

Additionally, they can be used to rank the importance of variables by assessing the

differences in error rate (or tree node purity [27]) when individual variables are

permuted across all component decision trees. While a number of machine learning

approaches could be applied to FCM and SFCM data, random forests have the

advantage of being flexible, fast, and robust, negating the need for specialized

computing resources.

We rely on training datasets based on lab cultures to (1) identify the traits that are

best at distinguishing live cells from other signals, (2) distinguish functional groups

from each other, and (3) quantify how all measured SFCM traits relate to cell

biovolume measured by microscopy. Briefly, we used the most important variables

identified by a random forest (applied to the lab culture training dataset) in

combination with the flowPeaks algorithm [26] to clean SFCM data and to identify

functional groups of cells in the cleaned data. The data we used to test this protocol

were collected from a natural lake community across several months of automated

monitoring. We used two additional random forests trained on clustered data to

classify all points from 191 files containing >20 million data points into phytoplankton

cells and other particles, and subsequently classify the cells into clusters

corresponding to functional groups. We used a fourth random forest trained on our

lab culture training dataset to estimate the biovolume of every phytoplankton cell

based on all the measured SFCM parameters.

We describe this procedure in detail below and validate it using 191 microscopy

measurements corresponding to the same depths and times of the SFCM

measurements (Fig. 1). We include the SFCM dataset we used to evaluate this

method on the data repository Zenodo, accessible at

https://doi.org/10.5281/zenodo.977772 [30]. We also include code for the entire

analysis on Github, accessible at https://doi.org/10.5281/zenodo.999747 [31],

implemented in the R software environment [32]. Associated microscopy data used

in this paper may be found in the supporting information.

Methods

Overview of approach:

Our approach is shown in flow chart form in Fig. 1. It involves 10 steps:

(1) Generation of a training dataset using lab cultures with known species identity,

functional group identity, and mean cell biovolume (Table 1).

(2) Identifying the parameters that most accurately distinguish between live cells and

other signals (a combination of bacteria, detritus, and electronic noise), using the

training dataset.

(3) Identifying clusters of similar points in a subset of the field dataset based on the

parameters identified in step 2.

(4) Classifying points from the complete field dataset into these clusters, and

removing all points except for those from live cells (data cleaning).

(5) Identifying the parameters that most accurately distinguish between different

functional groups, using the training dataset.

(6) Identifying clusters of similar points (corresponding to different functional groups)

in a subset of the cleaned field dataset based on the parameters identified in step 5.

(7) Classifying all cells from the cleaned field dataset into these clusters.

(8) Training a machine learning algorithm (a random forest) to estimate the

biovolumes of individual cells based on all measured SFCM parameters, using the

training dataset.

(9) Using the trained random forest from step 8 to estimate the biovolumes of every

cell in the cleaned field dataset based on all measured SFCM parameters.

(10) Estimating cell density and biovolume of the total phytoplankton community and

of the major clusters (corresponding to functional groups), using the assigned

clusters for each individual cell (from step 7) as well as their biovolume estimates

(from step 9) and cell density estimates.

Table 1. *Cultures used in biovolume estimation. Mean biovolume values were estimated by microscopy as part of this study.*

Species	Functional group	Coloniality	Mean biovolume (μm^3)
Synechococcus sp.	Cyanobacterium	Unicellular	13.4
Microcystis aeruginosa	Cyanobacterium	Unicellular (in culture)	17.2
Chlorella sp.	Green alga	Unicellular	28.7
Ankistrodesmus bibraianus	Green alga	Unicellular	31.6
Chroomonas sp.	Cryptophyte	Unicellular	60.8
Synura sp.	Chrysophyte	Unicellular	152.7
Cyclotella cf. glomerata	Diatom	Unicellular	226.2
Cryptomonas sp*.*	Cryptophyte	Unicellular	896.8
Asterionella formosa	Diatom	6 cells per colony, with rare exceptions	1964.8
Anabaena flos-aquae	Cyanobacterium	Long filaments with highly variable numbers of cells.	15478.7

Procedures:

1. Sampling campaign

Between August 17th and October 26th 2014, we collected SFCM measurements at 6

depths (1.0, 2.5, 4.0, 5.5, 7.0 and 8.5 m) every 4 hours in Greifensee, a eutrophic

lake in Switzerland (47.35°N, 8.68°E), using the automated monitoring station

Aquaprobe [4]. We also collected weekly microscopy samples at every depth within

1 hour of an SFCM measurement, obtaining a set of paired SFCM and microscopy

measurements. A small number of SFCM samples were lost, leaving us with 191

paired samples [30].

Over this time period, we saw environmental changes of approximately 15°C in

water temperature, and variation of an order of magnitude in dissolved phosphorus

concentration and N:P ratio.

The Office of Waste, Water, Energy and Air (AWEL) of Canton Zürich provided

permission for in situ monitoring.

2. Microscopy protocol

2.1. Microscopy sample collection and counting

Microscopy samples were collected manually using a Niskin bottle. The Niskin bottle

collects a vertical column of water 46 cms in length, which we centred at each of the

6 depths sampled by SFCM (therefore, the microscopy samples average over a

water sample 46 cms in length as opposed to 1 cm in length as in the case of

SFCM). 300 mL of water was fixed using Lugol's solution and stored in a brown

glass bottle in the dark till measurement.

Cells were counted using a Utermöhl counting chamber (3 mL) under 200X and

400X magnification [33]. 40 fields were counted and identified to a species level in

most cases, and to the genus level in the remaining cases. Every sample was

measured under both magnifications, with different species counted in each case

based on their size.

2.2. Microscopy biovolume estimation

The total biovolume of the whole community and of the individual functional groups

was calculated by multiplying the density of individual species by their mean

biovolume, and summing appropriately. Mean biovolume values of each species

were taken from an existing database of measurements made using standard

protocols on individual cells from the same lake, Greifensee (Buergi unpublished).

Local environmental conditions were therefore accounted for in these

measurements, but intraspecific variation and changes through time were not

considered. However, as intraspecific variation is considerably lower than

interspecific variation (which can be >7 orders of magnitude, [34]), we believe this is

likely to be a small source of error.

3. SFCM protocol

3.1. Instrument description

SFCM measurement was performed using the CytoSense

(http://www.cytobuoy.com), which is designed to characterize the scattering and

fluorescence of individual particles; here we use it to study phytoplankton cells. The

instrument measures particles across a large proportion of the phytoplankton length

range, between approximately 2 µm and 1 mm in length. Although this instrument is

capable of taking images, it is unable to resolve small cells clearly (approximately

<10 µm). As these small cells are abundant in natural systems such as freshwater

lakes, it necessitated the development of the method we present here.

Particles cross two coherent 15mW solid-state lasers (488 nm and 642 nm). Cells

absorb and scatter these wavelengths, and also fluoresce at longer wavelengths

determined by their specific pigment composition. Scattering (forward and sideward;

the latter is typically referred to as 'side-scatter' in the flow cytometry literature) is

measured, as well as fluorescence at each of three channels hereafter referred to as

Red (668 – 734nm), Orange (601 – 667nm) and Yellow (536 – 601nm). These

channels are listed below, followed by a more detailed description:

Red fluorescence Channel Laser 1: 488: 701/33

Red fluorescence Channel Laser 2: 642: 701/33

Orange fluorescence Channel: 488: 634/33

Yellow fluorescence Channel: 488: 569/33

Fluorescence in the Yellow and Orange channels is exclusively stimulated by the

488nm laser, while fluorescence in the Red channel is stimulated by both lasers.

Therefore, the Red channel is electronically deconvoluted into Red 1 (from the

488nm laser) and Red 2 (a weaker signal from the 642nm laser). These wavelength

bands are not highly specific in terms of the pigments they target, but generally

target chlorophyll-a (Red 1), phycocyanin (Red 1 and Red 2), phycoerythrin

(Orange), and carotenoids and decaying pigments (Yellow).

Hereafter, names of parameters representing the different fluorescence channels are

preceded by '*FL*', the Red 1 channel is referred to as '*FL.Red*' and the Red 2

channel as '*X2.FL.Red*'. Names of parameters representing the scattering channels

are preceded by *'FWS*' (forward scatter; typically abbreviated FSC in the flow

cytometry literature) and '*SWS*' (sideward scatter or side-scatter' typically

abbreviated SSC in the flow cytometry literature).

The signal produced by each particle is a time series of measurements for each

channel, which describe the variation in scattering and fluorescence over the length

of the particle. This high-resolution time series describes a pulse for each channel (4

fluorescence and 2 scattering). These pulses may be highly irregular in shape (Fig.

S1) and are therefore characterized using a number of parameters (see Table S1 for

parameter descriptions).

3.2. Instrument configuration

286 Internal flow rates were set at 2 µL.s^{-1}. A trigger threshold of 99.7 mV on the

287 sideward scatter channel was enforced; particles whose scattering did not rise above

288 this level were therefore not recorded. Field samples measurements were terminated

289 when 500 µL was measured or 9 minutes elapsed, whichever was earlier.

3.3. Field sampling procedure

Every four hours, the sampling tube was automatically deployed to each of the 6

depths by the automated station (described in [4]). Water samples were pumped to a

250 mL sampling chamber at the surface through a tube with a 1-cm diameter

opening, making these highly depth-specific measurements. The sampling chamber

was flushed with water from the sampling depth four to five times over 2 minutes

before the CytoSense collected a water sample of up to 500 µL for measurement.

3.4 Generation of training datasets using lab cultures

Our training dataset for the random forest served 3 purposes: (1) to enable the

identification of parameters that most strongly distinguish between live cells and

other signals, (2) to enable the identification of parameters that most strongly

distinguish between cells belonging to different functional groups, and (3) to train the

algorithm to estimate cell biovolume based on all the measured CytoSense

parameters.

We measured ten laboratory cultures belonging to multiple functional groups (Table

1). In clonal lab cultures, manual identification of live cells and other signals is

straightforward (examples in Fig. S2). We generated a dataset containing

approximately 1,200 measurements of live cells (manually identified as in Fig. S2)

and 8,400 measurements of other signals, equally sampled from all species'

measurements. The proportion was chosen to approximately mirror the low

proportion of live cells expected from field measurements.

We note that overall performance will likely improve if the training dataset is

improved. This may be done by increasing the number of species, number of

functional groups, range of sizes, and range of culture conditions.

3.5 Data cleaning

3.5.1. Identifying parameters that distinguish live phytoplankton cells from other

signals

The CytoSense measures 94 parameters for every particle. However, due to the

'curse of dimensionality' problem for clustering algorithms, we chose to reduce the

number of variables examined. We used a machine learning algorithm, a random

forest [27], to identify the variables that most strongly distinguished between live

cells and other signals. Most importantly for this purpose, random forests can be

used to generate a ranking of variable importance based on the change in

classification error rate when a particular variable is permuted across individual

decision trees. For this and all subsequent uses of random forests, we used the R

package *randomForest* [35].

Before analysis, we log-transformed all variables whose minimum value was > 0,

because most variables exhibited highly skewed distributions that could prove

challenging for subsequent analyses. Then, to avoid problems associated with

multicollinearity, we removed one member of all pairs of variables in the laboratory

dataset whose correlation coefficient was > 0.8 or < -0.8. We were left with 51

variables. We trained a random forest with 10,001 trees on this dataset and found

that the 10 most important variables enabled a clear distinction between live cells

and other signals (Fig. 2). In descending order of importance, they were:

FL.Red.Range, *X2.FL.Red.Range*, *X2.FL.Red.Gradient, FL.Red.Number.of.cells,*

X2.FL.Red.Last, FWS.Range, *FL.Red.Fill.factor, X2. FL.Red.Fill.factor,*

FL.Yellow.Range, and *FL.Orange.Range*. We note that the specific number of

variables used is a feature of the dataset being considered, and readers are advised

to explore using differing numbers of variables to identify the optimal number for their

data.

3.5.2 Generating a random subset of the raw field data

The complete dataset of 191 measurements contained >20 million particles and

could not be clustered simultaneously using standard computing resources.

Furthermore, clustering each measurement separately would have led to the

identification of different clusters as the community changed, limiting the possibility to

make comparisons between measurements. We therefore generated a file

containing an equal number of randomly selected data points from every measured

sample, to distinguish between living phytoplankton cells and other signals. The data

subset contained approximately 100,000 points which was both computationally

tractable, and large enough to enable the identification of distinct clusters containing

relatively low numbers of points that may be missed with smaller file sizes.

Fig. 2. *Importance of different traits in distinguishing between particle types, as determined by random forests applied to laboratory training data (Fig. S2). Variable importance is assessed using a Gini coefficient that measures the change in classification error when a variable is omitted [27]. The 30 most important variables are shown here. a) Variables that most strongly distinguish between live cells and other signals (electronic noise, detritus and bacterial cells). The top ten were used for unsupervised clustering of the raw field data. a) Variables that most strongly distinguish between major functional groups. The top eight were used for unsupervised clustering of the cleaned field data.*

3.5.3 Unsupervised clustering of the subset of raw data

We selected the 10 variables identified in section 3.5.1, log-transformed them and
used the *flowPeaks* algorithm [26] to identify clusters of similar data points.
flowPeaks (implemented in the R statistical environment [32]) uses a k-means-based
algorithm to first cluster the entire dataset into large numbers of small clusters, and
then merges clusters based on density gradients. The smoothness of the density
function is determined by tuneable parameters; we used values of 0.25 for *tol*, 0.05
for *h0*, and 2 for *h* [26] as these provided clusters that matched our expectations
based on visual examination of 3D plots. However, the best parameter values may
differ between datasets.

The algorithm identified 8 clusters, which we visually inspected using 3D plots (Fig.
S3 and animated versions of S3a and S3b in the supporting information, Table S2).
From these plots, we identified seven clusters corresponding to live phytoplankton
cells, which together constituted 5% of the data. The remaining cluster corresponded
to signals we were uninterested in, and cannot distinguish between: a combination of
bacteria, dead phytoplankton, detritus, and electronic noise. Phytoplankton clusters
were primarily characterized by high pigment fluorescence, particularly
FL.Red.Range (a proxy for chlorophyll-a) (Table S2) and high forward scatter (FWS).

3.5.4 Using random forests to classify raw data points and clean dataset

We used a second random forest [27,35] to classify all points in every sample into
one of the eight clusters identified in the previous step using *flowPeaks* (code
available at [31]). We trained the random forest with 1,001 trees on the clustered
100,000 point data subset, allowing it to identify nonlinear boundaries between the
eight clusters. The out-of-bag classification error rate of this random forest was <2%,

indicating acceptably accurate performance in this classification step. We then used

this random forest to classify all points in every sample into one of the eight clusters

identified earlier. Points identified as belonging to one of the seven phytoplankton

clusters were retained and the remainder discarded. The retained points from each

sample were saved in separate files for further analysis.

3.6 Functional group identification

We identified the functional groups that individual cells belonged to by essentially

repeating the data cleaning procedure (section 3.5) on the cleaned data. There were

minor differences in procedure that we note below.

3.6.1. Identifying parameters that distinguish between phytoplankton functional

groups

For this step, we used the lab culture training dataset containing 1,200

measurements of live phytoplankton cells only, belonging to five different functional

groups – chrysophytes, cryptophytes, cyanobacteria, green algae, and diatoms

(Table 1).

We then used a random forest to identify the variables that most strongly

distinguished between these functional groups. As earlier, we log-transformed all

variables whose minimum value was > 0 and removed one member of all pairs of

variables in the laboratory dataset whose correlation coefficient was > 0.8 or < -0.8.

We were left with 43 variables. We trained a random forest with 10,001 trees [35]

and found that 8 variables enabled a clear distinction between cells belonging to

different functional groups (Fig. 2). In descending order of importance, they were:

FL.Red.Range, Red1Red2.ratio, FWS.Length, FL.Orange.Range, X2.FL.Red.Range,

FL.Red.First, X2.FL.Red.Gradient and FWS.Fill.factor.

3.6.2 Generating a random subset of the cleaned field data

To identify phytoplankton cells belonging to different functional groups, we generated

a file containing an equal number of randomly selected data points from every

cleaned data file, for a total of approximately 100,000 points.

3.6.3 Unsupervised clustering of the subset of cleaned data

We selected the 8 variables identified in section 3.6.1, log-transformed them and

applied *flowPeaks* [26] to identify clusters of similar data points. We used the same

parameter values as earlier ($tol = 0.25$, $h0 = 0.05$, and $h = 2$). The algorithm

identified 12 clusters, which we visually inspected using 3D plots (Figs. S4 and

animated versions of S4a and S4b in the supporting information, Table S3). We

limited our subsequent analyses to the four clusters that formed >5% of the dataset

(totalling >90% of the data), but note here that (i) functional groups may comprise

multiple clusters and (ii) rarer functional groups are unassigned. The latter this is not

a limitation of the method; a detailed laboratory comparison of the trait signatures of

all functional groups would permit the assignment of every cluster to a functional

group.

3.6.4 Assignment of clusters to specific functional groups

Based on the fluorescence and scattering signals of these clusters and their relative

abundance in the microscopy data, we assigned the four clusters to four functional

groups. We note that our microscopy data showed that diatoms, dinoflagellates,

euglenophytes, and desmids were almost entirely absent from the lake during the

periods we sampled, leaving us with four well-represented functional groups –

chrysophytes, cryptophytes, cyanobacteria, and green algae.

Cyanobacteria are characterised by a low *Red1Red2.ratio* value (indicative of high

phycocyanin content), which was characteristic of Cluster 2 (Tables S3, S4). Our

assignment of the remaining clusters is necessarily more speculative and would be

improved by a broader sampling of the phenotypic diversity of these groups using lab

cultures. Cluster 1, the most abundant group in the SFCM data, was designated as

chrysophytes, the most abundant eukaryotic group in the microscopy counts. Cluster

3 was characterized by a low *X2.FL.Red.Gradient* and *FL.Red.First* and an

intermediate *Red1Red2.ratio*, indicative of cryptophytes. Cluster 4 was characterized

by relatively high *Red1Red2.ratio* and *FL.Red.Range*, suggesting that they represent

green algae. Note that in most cases other than *Red1Red2.ratio*, we did not have an

a priori expectation relating to the importance of these variables in differentiating

between functional groups. Therefore, we do not speculate on the reasons behind

their importance here.

3.6.5 Using random forests to classify cleaned data into trait clusters

We trained a random forest with 1,001 trees on the 100,000 point cleaned data

subset (code available at https://doi.org/10.5281/zenodo.999747), allowing it to

identify nonlinear boundaries between the clusters identified in the previous stage.

Out-of-bag error rates for this random forest were <2%, again suggesting acceptably

low levels of classification error. We then used this random forest to classify all cells

in the cleaned data files into one of the 12 trait clusters identified earlier. We

recorded the cell densities of every trait cluster in every sample.

3.7. Biovolume estimation

3.7.1 Estimating the biovolumes of single cells and colonies

We trained a random forest with 10,001 trees on our lab culture training dataset

(Table 1) to estimate the biovolumes of individual cells using all measured SFCM

parameters. We used all the parameters and a training dataset with taxa from all the

major functional groups in order to train a random forest that accounts for taxonomic

differences in scattering and pigmentation properties. This approach was

substantiated by our finding that *Red1Red2.ratio* (indicative of whether a cell was a

cyanobacterium or a eukaryote) was the most important predictor of biovolume (Fig.

S5) and not scattering, which has previously been used in biovolume estimation of

phytoplankton cells by SFCM [9,10]. Furthermore, the trained random forest had a

model R^2 (based on out-of-bag estimates) of 0.93 as opposed to a linear model

based on the best single scattering variable (FWS.Length), which had an R^2 of 0.50.

We used this trained random forest to predict the biovolume of every individual cell in

our cleaned field data based on all their SFCM parameters.

3.7.2 Estimating the density of the whole community and major functional groups

We first quantified the cell density of each community by multiplying the particle

concentration (estimated by the CytoSense) with the proportion of particles that were

cells in the sample. Subsequently, we estimated the cell density of each cluster

(functional group) by multiplying the cell density of the community by the proportion

of cells in the sample belonging to the cluster.

3.7.3 Estimating the total biovolume of the whole community and major functional

groups

We estimated total biovolume of each community by multiplying the cell density

estimated in section 3.7.2 with the mean biovolume of all cells in the community.

Similarly, we estimated the biovolume of each cluster (functional group) by

multiplying the estimated cell density of each cluster by the mean biovolume of all

cells belonging to that cluster.

Results and assessment

We assessed the accuracy of our data cleaning, functional group identification, and

biovolume estimation procedures by using field data to compare SFCM estimates

with microscopy estimates of (1) cell density at the whole-community level, (2) total

biovolume at the whole-community level, (3) cell density of the major functional

groups, and (4) total biovolume of the major functional groups.

1) Cell density of the phytoplankton community

SFCM estimates of whole-community cell density were strongly positively correlated

with microscopy estimates ($r = 0.73$, $p < 0.001$, Fig. 3a). The slope of the regression

between the two estimates is indistinguishable from 1 (mean = 0.99, CIs: 0.86, 1.13)

and the intercept is indistinguishable from 0 (mean = -0.13, CIs: -0.65, 0.38).

Quantitative estimates of cell density were extremely similar in both cases. SFCM

estimates were marginally lower on average, possibly due to (i) calibration of the

SFCM inlet pump, leading to lower flow rates and consequently cell density

estimates, (ii) difficulties in evaluating whether a cell was alive or dead in fixed

microscopy samples, leading to overestimation of the microscopy cell density, or (iii)

break-up of colonial organisms during preservation for microscopy. Whatever the

underlying reason, the apparent difference is small, and variation in SFCM cell

Fig. 3. *Estimates of whole-community cell density and biovolume by SFCM are strongly correlated with those by microscopy (on the log scale), and are quantitatively accurate. Each point represents a single depth-specific sample measured using both methods. The black line is the 1:1 line and the red line represents the regression relationship. a) Cell density estimates are strongly correlated (r = 0.73, p < 0.001). b) Whole-community total biovolume estimates are strongly correlated (r = 0.67, p < 0.001). The slope of the regression is indistinguishable from 1 (mean = 0.90, CIs: 0.76, 1.05) and the intercept is indistinguishable from 0 (mean = -0.40, CIs: -0.54, 1.33).*

density estimates accurately captures variation in cell density by microscopy

measurements.

2) Total biovolume of the phytoplankton community

SFCM estimates of total community biovolume are strongly correlated with

microscopy estimates ($r = 0.67$, $p < 0.001$, Fig. 3b). The slope of the regression is

indistinguishable from 1 (mean = 0.90, CIs: 0.76, 1.05) and the intercept is

indistinguishable from 0 (mean = -0.40, CIs: -0.54, 1.33). Quantitative estimates of

total biovolume were similar in absolute magnitude over an approximately 30-fold

range in biovolume, but appear to be slightly underestimated. This underestimation

is partly driven by differences in cell density estimates (Fig. 3a), because density

estimates are involved in total biovolume calculation. Differences in total biovolume

are also influenced by our microscopy biovolume estimation procedure: estimates

were generated by referring to a database of mean cell biovolumes and were not

measured in the individual samples. Therefore, errors in our cell size database are

propagated through to our estimates of total community biovolume by microscopy.

However, intraspecific variation in cell biovolume is substantially lower than

interspecific variation (which varies by >7 orders of magnitude [34]), and so we

expect that this is at most a small source of error.

3) Cell density of the major functional groups

SFCM estimates of the cell density of major functional groups were moderately to

strongly correlated with microscopy estimates (r ranges from 0.44 to 0.76, $p < 0.001$

in all cases, Fig. 4). Chrysophytes were the most abundant group overall and also

show the strongest correlation and quantitative agreement between estimates.

Cryptophytes and green algae show weaker correlations with slopes that were <1,

Fig. 4. *Estimates of cell density of four major functional groups by SFCM and microscopy are moderately to strongly correlated on the log scale (r between 0.44 and 0.76, p < 0.001 in all cases). Each point represents a single depth-specific sample measured using both methods. The black line is the 1:1 line and the red line represents the regression relationship (with confidence bands). In a small number of microscopy measurements, no cyanobacteria were found. These points are plotted with high transparency assuming that the true concentration was half the detection limit of 28 cells per mL. A small number of points were excluded from the regressions due to their being excessively influential, but the effect of this exclusion on slopes and correlation coefficients is modest.*

but also exhibited relatively low variation over the sampling period, making it

challenging to compare accurately. Cyanobacterial density estimates also showed a

strong positive relationship, but this relationship was influenced by a small number of

microscopy measurements in which no cyanobacterial cells were identified. These

samples appeared to have been substantially overestimated by SFCM or

underestimated by microscopy. This may partly be explained by the lower detection

limit and consequently lower sampling variability of SFCM measurements when

compared to microscopy measurements, but may also be a result of cell degradation

in samples which were not properly fixed. Though the quantitative estimates are

substantially better if we exclude these points, the relationship remains reasonably

strong even if we include them by assuming a cell density of 50% of the detection

limit.

4) Total biovolume of the major functional groups

SFCM estimates of the total biovolume of major functional groups were moderately

to strongly correlated with microscopy estimates (*r* ranges from 0.41 to 0.75*, p <*

0.001 in all cases, Fig. 5). As with cell density, chrysophytes showed not just a

strong correlation between estimates, but also highly similar quantitative estimates.

Cryptophytes and green algae exhibited a relatively small degree of variation and

differed in absolute value, but were positively correlated in both cases. As in Fig. 4,

cyanobacterial estimates were reasonably accurate, but a small number of

measurements showed no cells in microscopy measurements even though the

SFCM estimates were moderately high. Notably, slopes and correlation coefficients

are relatively similar in both cell density and total biovolume. As cell density is used

to estimate total biovolume, this suggests that the discrepancy is largely caused by

Fig. 5. *Estimates of total biovolume of four major functional groups by SFCM and microscopy are moderately well-correlated on the log scale (r between 0.41 and 0.75, p < 0.001 in all cases). Each point represents a single depth-specific sample measured using both methods. The black line is the 1:1 line and the red line represents the regression relationship (with confidence bands). In a small number of microscopy measurements, no cyanobacteria were found. These points are plotted with high transparency assuming that the true concentration was the minimum cyanobacterial biovolume detected. A small number of points were excluded from the regressions due to their being excessively influential, but the effect of this exclusion on slopes and correlation coefficients is modest.*

these estimates, and that mean cell biovolume of each group is being estimated

accurately. We have demonstrated that this method produces estimates of cell

density and biovolume that are reasonably well-correlated with estimates by

microscopy, especially at the whole community level. These correlations may have

been even stronger if microscopy samples could be collected at the same depth

resolution as our SFCM measurements (we sampled a 46 cm-wide water column for

microscopy and a 1 cm-wide water column for SFCM). And although we ignored less

abundant clusters in this analysis because of limitations in the number of lab-

measured taxa, our method can easily be expanded upon to include finer-scale

variation and a wider range of functional groups.

Our protocol (Fig. 1) is composed of three parts that can each be improved upon

with more data or improved algorithms, while following essentially the same protocol.

The strong agreement between SFCM results and microscopy results from a

complex natural community supports our claim. It improves on existing methods by

(i) reducing the subjectivity associated with manual cluster designation, and (ii)

increasing the scalability of FCM and SFCM analyses by enabling the study of

datasets of arbitrarily large size using basic computing facilities. This scalability is

achieved by coupling unsupervised clustering with machine learning techniques

(which have frequently been used independently previously) and reducing the size of

the major tasks through the use of training datasets and data subsets.

However, there are important limitations to our method:

1) The generation of training datasets is necessary and can be time-consuming.

Training datasets must contain representatives of all groups present in the system

under study (with the possible exception of taxa that are rare both in terms of

abundance and biovolume contribution). Moreover, this training exercise will need to be done independently for every instrument and potentially for changes in important instrument settings (such as photomultiplier tube gain) as well. It is possible that a training dataset also includes measurements of taxa grown under a variety of environmental conditions would improve performance by accounting environment-dependent plasticity. Currently, our method might lead to biased estimates of biovolume due to plastic changes in fluorescence traits (such as due to decreases in light with depth) that our random forest does not account for. Functional groups that are similar to each other in traits may also not be distinguished well (i.e. there may be misclassification) if this plasticity leads to changes in their trait distributions. Additionally, functional groups are diverse and our training dataset contains relatively few species from each cluster, potentially biasing biovolume estimates. However, this can be solved by including more representative species from each group in the training dataset.

2) Many clustering algorithms have difficulties in identifying clusters that are highly nonlinear in multiple dimensions. Unsupervised clustering is an active area of research and future advances in density-based clustering methods and neural network-based methods may alleviate some of these problems [36]. However, every such algorithm has its own weaknesses and such improvements may come at the cost of other features (e.g. computational speed).

3) By clustering datasets with equal numbers of data points randomly selected from every time point, we might miss out on clusters that are present for very short periods of the time series. These clusters are poorly represented even if abundant for short time periods. This limitation is even more relevant if our approach is used to study datasets collected across space instead of time; clusters present at just one or

two of a large number of sites may be difficult to detect. Addressing this issue will

require careful sampling, but may also be aided by advances in clustering methods.

A sequential clustering approach that re-clusters the data after the most abundant

cluster is removed may also be useful in identifying these cases.

4) Our assignment of clusters to functional groups is at present provisional, except in

the case of cyanobacteria, which are very strongly differentiated by their ratio of Red

1 to Red 2 fluorescence. This may be improved on by using instruments configured

with more laser and detectors than the one used here, enabling the more precise

targeting of pigments characterising different functional groups. In the absence of

this, the problem may be mitigated by using larger training datasets with more

representative species from each functional group.

Future work may therefore most profitably expand on our work by using a much

broader training dataset to train machine learning algorithms. Capturing the entire

range of functional groups and a broad sampling of the phenotypic variation within

these groups would enable the classification of cells into functional groups (and

potentially lower level taxonomic categories) without the need for unsupervised

clustering.

Discussion

Scanning flow cytometry has the potential to enable the high-frequency monitoring of

natural phytoplankton communities in real time, improving our ability to understand

their ecology [4,9]. For some ecological questions, such as those relating to the

influence of short-term environmental changes on populations and communities,

they are (along with IFCM) the only feasible measurement tool at present. Tools to

automate the analysis of FCM and SFCM data will greatly increase the utility of

these instruments, increasing our ability to understand the drivers of high-frequency

dynamics in natural communities.

A very large number of unsupervised clustering algorithms exist with differing

strengths and weaknesses [28,29,37,38], and we did not attempt to compare the

efficacy of different algorithms here. Instead, we use an algorithm (flowPeaks [26])

that has previously been applied to phytoplankton populations [17] and provides

rapid, reasonably accurate results using relatively limited computational resources.

More sophisticated algorithms that are capable of identifying points that do not below

to any clear cluster ('noise') such as DBSCAN [39] may improve on this, but they

generally require computationally expensive calculations, including the distance

between all pairs of points to be clustered. Our approach is flexible and alternative

algorithms can be substituted here if found to be more suitable.

A more general problem that our approach solves is that unsupervised clustering of

high-dimensional SFCM data suffers from the 'curse of dimensionality'. Output is

less reliable as dimensionality increases, and visualisation of clusters also becomes

difficult beyond three dimensions [40]. Therefore, there is a strong need to reduce

the dimensionality of the problem for both reliable clustering and for visualisation.

This may be done by two methods: (i) we could use a dimension-reduction algorithm

such as principal components analysis, or (ii) we may limit the number of dimensions

that we choose to cluster. We followed the second approach here because the

former suffers from the weakness that the major principal components may not be

meaningful in distinguishing between the signals we are interested in; in other words,

there may be more variation in unimportant dimensions than in important ones.

Although machine learning techniques have previously been applied to FCM data

[12,41–43], our work differs in using it for tasks beyond classification. We have also

used machine learning to predict biovolumes of individual cells, and to identify the

phenotypic features that most reliably distinguish between particle types (Figs. 1, 2).

When used in combination with unsupervised clustering, it constitutes a powerful and

flexible tool in processing large FCM and SFCM datasets. In principle, a machine

learning algorithm trained on lab cultures may be able to identify trait clusters in field

data. However, in practice we need to couple this with unsupervised clustering

because trait cluster boundaries differ between lab and field communities. This is

because (i) natural communities contain species that may not be in the training

dataset, and (ii) changes in environmental conditions may lead to plastic changes in

fluorescence and scattering, shifting the boundaries between clusters.

Our method provides a repeatable, semi-automated protocol with which to process

large FCM and SFCM datasets using readily available computing resources and free

software tools (the R software environment and packages *randomForest* and

flowPeaks). Our work integrates existing tools - lab measurements, machine

learning, and unsupervised classification - from multiple fields to improve the speed,

repeatability, and robustness of SFCM analysis. We believe that this protocol will

greatly increase the ease and speed with which large SFCM datasets can be

processed and analysed. In the future, automated (or semi-automated) analytical

tools such as these may enable the monitoring of water bodies for harmful algal

bloom development in real time, through classification of cells based on previously

developed training datasets. Cyanobacteria, which are responsible for freshwater

harmful algal blooms, possess fluorescence traits were especially easy to distinguish

with our instrument configuration, making the identification and quantification of this

taxon simple. Examination of a broader range of cyanobacteria may allow us to

identify features that distinguish between potentially toxic cyanobacterial taxa and

taxa that do not produce toxins, greatly increasing water quality monitoring efforts.

However, this protocol will also improve our ability to study fundamental aspects of

the physiology, ecology and evolution of phytoplankton communities, by enabling

their monitoring on time-scales orders of magnitude lower than is presently feasible

using microscopy methods.

Acknowledgements:

This work was supported by the Swiss National Science Foundation project

CRSII2_147654, "In-situ Sensing Tools for Understanding Rapid Microscale

Plankton Dynamics". We would like to thank M. Stravs and L. Nizzetto for providing

phytoplankton cultures, Esther Keller for microscopy analysis and Hannele Penson

for assistance with field work, and Michael Kehoe for advice related to machine

learning techniques.

Statement of author contributions

712 MKT and FP conceived of the method with input from SF. FP, SF, MR & MKT

713 collected the field data. MR maintained lab cultures and collected lab SFCM and

714 microscopy measurements. MKT wrote the manuscript with substantial input from

715 FP, SF and MR.

References:

1. Hammes F, Egli T. Cytometric methods for measuring bacteria in water:
Advantages, pitfalls and applications. Anal Bioanal Chem. 2010;397: 1083–
1095. doi:10.1007/s00216-010-3646-3

2. Sosik HM, Olson RJ, Armbrust EV. Flow cytometry in phytoplankton research.

In: Suggest DJ, Prášil O, Borowitzka MA, editors. Chlorophyll a Fluorescence
in Aquatic Sciences: Methods and Applications. Dordrecht: Springer
Netherlands; 2010. pp. 171–185. doi:10.1007/978-90-481-9268-7

3.   Wang Y, Hammes F, De Roy K, Verstraete W, Boon N. Past, present and
future applications of flow cytometry in aquatic microbiology. Trends
Biotechnol. 2010;28: 416–424. doi:10.1016/j.tibtech.2010.04.006

4.   Pomati F, Jokela J, Simona M, Veronesi M, Ibelings BW. An automated
platform for phytoplankton ecology and aquatic ecosystem monitoring. Environ
Sci Technol. 2011;45: 9658–65. doi:10.1021/es201934n

5.   Arnoldini M, Heck T, Blanco-Fernández A, Hammes F. Monitoring of dynamic
microbiological processes using real-time flow cytometry. PLoS One. 2013;8:
e80117. doi:10.1371/journal.pone.0080117

6.   Hunter-Cevera KR, Neubert MG, Solow AR, Olson RJ, Shalapyonok A, Sosik
HM. Diel size distributions reveal seasonal growth dynamics of a coastal
phytoplankter. PNAS. 2014;111: 9852–9857. doi:10.1073/pnas.1321421111

7.   Besmer MD, Weissbrodt DG, Kratochvil BE, Sigrist JA, Weyland MS, Hammes
F. The feasibility of automated online flow cytometry for in-situ monitoring of
microbial dynamics in aquatic ecosystems. Front Microbiol. 2014;5: 1–12.
doi:10.3389/fmicb.2014.00265

8.   Dubelaar GBJ, Gerritzen PL. A Step Forward towards Using Flow Cytometry in
Operational Oceanography. Sci Mar. 2000;64: 255–265.

9.   Pomati F, Kraft NJB, Posch T, Eugster B, Jokela J, Ibelings BW. Individual cell
based traits obtained by scanning flow-cytometry show selection by biotic and
abiotic environmental factors during a phytoplankton spring bloom. PLoS One.
2013;8: e71677. doi:10.1371/journal.pone.0071677

10.   Pomati F, Nizzetto L. Assessing triclosan-induced ecological and trans-
generational effects in natural phytoplankton communities: a trait-based field
method. Ecotoxicology. 2013;22: 779–94. doi:10.1007/s10646-013-1068-7

11.   Dubelaar GBJ, Jonker RR. Flow cytometry as a tool for the study of
phytoplankton. Sci Mar. 2000;64: 135–156. doi:10.3989/scimar.2000.64n2135

12.   Sosik HM, Olson RJ. Automated taxonomic classification of phytoplankton
sampled with imaging in-flow cytometry. Limnol Oceanogr Methods. 2007;5:
204–216. doi:10.4319/lom.2007.5.204

13.   Lo K, Brinkman RR, Gottardo R. Automated gating of flow cytometry data via
robust model-based clustering. Cytom Part A. 2008;73: 321–332.
doi:10.1002/cyto.a.20531

14.   Aghaeepour N, Nikolic R, Hoos HH, Brinkman RR. Rapid cell population
identification in flow cytometry data. Cytom Part A. 2011;79A: 6–13.
doi:10.1002/cyto.a.21007

15.   Malkassian A, Nerini D, Van Dijk MA, Thyssen M, Mante C, Gregori G.
Functional analysis and classification of phytoplankton based on data from an

automated flow cytometer. Cytom Part A. 2011;79 A: 263–275. doi:10.1002/cyto.a.21035

16. Glüge S, Pomati F, Albert C, Kauf P, Ott T. The challenge of clustering flow cytometry data from phytoplankton in lakes. In: Mladenov V, Ivanov PC, editors. Nonlinear Dynamics of Electronic Systems: 22nd International Conference, NDES 2014, Albena, Bulgaria, July 4-6, 2014 Proceedings. Springer International Publishing Switzerland; 2014. pp. 379–386.

17. Hyrkas J, Clayton S, Ribalet F, Halperin D, Virginia Armbrust E, Howe B. Scalable clustering algorithms for continuous environmental flow cytometry. Bioinformatics. 2015;32: 417–423. doi:10.1093/bioinformatics/btv594

18. Field CB, Behrenfeld MJ, Randerson JT, Falkowski PG. Primary production of the biosphere: Integrating terrestrial and oceanic components. Science. 1998;281: 237–240.

19. Litchman E, Klausmeier CA. Trait-based community ecology of phytoplankton. Annu Rev Ecol Evol Syst. 2008;39: 615–639. doi:10.1146/annurev.ecolsys.39.110707.173549

20. Edwards KF, Thomas MK, Klausmeier CA, Litchman E. Allometric scaling and taxonomic variation in nutrient utilization traits and maximum growth rate of phytoplankton. Limnol Oceanogr. 2012;57: 554–566. doi:10.4319/lo.2012.57.2.0554

21. Edwards KF, Thomas MK, Klausmeier CA, Litchman E. Light and growth in marine phytoplankton: allometric, taxonomic, and environmental variation. Limnol Oceanogr. 2015;60: 540–552. doi:10.1002/lno.10033

22. Sun J, Liu D. Geometric models for calculating cell biovolume and surface area for phytoplankton. J Plankton Res. 2003;25: 1331–1346. doi:10.1093/plankt/fbg096

23. Foladori P, Quaranta A, Ziglio G. Use of silica microspheres having refractive index similar to bacteria for conversion of flow cytometric forward light scatter into biovolume. Water Res. 2008;42: 3757–3766. doi:10.1016/j.watres.2008.06.026

24. Reynolds CS. The ecology of phytoplankton. Cambridge University Press; 2006.

25. Steinberg CEW, Schäfer H, Siedler M, Beisker W. Ataxonomic assessment of phytoplankton integrity by means of flow cytometry. In: Seiler JP, Kroftová O, Eybl V, editors. Toxicology - From Cells to Man. Springer Berlin Heidelberg; 1996. pp. 417–434.

26. Ge Y, Sealfon SC. Flowpeaks: A fast unsupervised clustering for flow cytometry data via K-means and density peak finding. Bioinformatics. 2012;28: 2052–2058. doi:10.1093/bioinformatics/bts300

27. Breiman L. Random Forests. Mach Learn. 1999;45: 1–35. doi:10.1023/A:1010933404324

804   28.   Jain AK, Dubes RC. Algorithms for clustering data. Prentice-Hall, Inc.; 1988.

805   29.   Jain AK, Murty MN, Flynn PJ. Data clustering: A review. ACM Comput Surv.
806   1999;31: 264–323. doi:10.1145/331499.331504

807   30.   Thomas MK, Fontana S, Reyes M, Pomati F. Dataset: Quantifying cell
808   densities and biovolumes of phytoplankton communities and functional groups
809   using scanning flow cytometry, machine learning and unsupervised clustering.
810   2017. doi:10.5281/zenodo.977772. https://doi.org/10.5281/zenodo.977772

811   31.   Thomas MK. R Code: Cleaning and clustering SFCM data. 2017.
812   doi:10.5281/zenodo.999747. https://doi.org/10.5281/zenodo.999747

813   32.   R Core Team. R: A language and environment for statistical computing.
814   Vienna, Austria: R Foundation for Statistical Computing; 2017.

815   33.   Utermöhl H von. Neue Wege in der quantitativen Erfassung des Planktons.
816   (Mit besondere Beriicksichtigung des Ultraplanktons). Verhandlungen der Int
817   Vereinigung für Theor und Angew Limnol. 1931;5: 567–596.

818   34.   Litchman E, Klausmeier CA, Schofield OME, Falkowski PG. The role of
819   functional traits and trade-offs in structuring phytoplankton communities:
820   Scaling from cellular to ecosystem level. Ecol Lett. 2007;10: 1170–81.
821   doi:10.1111/j.1461-0248.2007.01117.x

822   35.   Liaw A, Wiener M. Classification and Regression by randomForest. R News.
823   2002;2: 18–22.

824   36.   Stoop R, Kanders K, Lorimer T, Held J, Albert C. Big data naturally rescaled.
825   Chaos, Solitons and Fractals. 2016;90: 81–90.

826   37.   Parsons L, Parsons L, Haque E, Haque E, Liu H, Liu H. Subspace clustering
827   for high dimensional data: A review. ACM SIGKDD Explor Newsl. 2004;6: 90–
828   105. doi:10.1145/1007730.1007731

829   38.   Aghaeepour N, Finak G, Dougall D, Khodabakhshi AH, Mah P, Obermoser G,
830   et al. Critical assessment of automated flow cytometry data analysis
831   techniques. Nat Methods. 2013;10: 228–238. doi:10.1038/nmeth.2365

832   39.   Ester M, Kriegel HP, J S, Xu X. A density-based algorithm for discovering
833   clusters in large spatial databases with noise. KDD. 1996;96.

834   40.   Newell EW, Cheng Y. Mass cytometry: Blessed with the curse of
835   dimensionality. Nat Immunol. Nature Publishing Group; 2016;17: 890–5.
836   doi:10.1038/ni.3485

837   41.   Balfoort HW, Snoek J, Smiths JRM, Breedveld LW, Hofstraat JW, Ringelberg
838   J. Automatic identification of algae: Neural network analysis of flow cytometric
839   data. J Plankton Res. 1992;14: 575–589. doi:10.1093/plankt/14.4.575

840   42.   Boddy L, Morris CW, Wilkins MF, Al-Haddad L, Tarran G a, Jonker RR, et al.
841   Identification of 72 phytoplankton species by radial basis function neural
842   network analysis of flow cytometric data. Mar Ecol Prog Ser. 2000;195: 47–59.
843   doi:10.3354/meps195047

844   43.   Boddy L, Wilkins MF, Morris CW. Pattern recognition in flow cytometry.
845   Cytometry. 2001;44: 195–209.