Comparison of ultracentrifugation and a commercial kit for isolation of exosomes derived from glioblastoma and breast cancer cells

Frøydis Sved Skottvoll[a]*, Henriette Engen Berg[a]*, Kamilla Bjørseth[a], Kaja Lund[b], Norbert Roos[c], Sara Bekhradnia[a], Bernd Thiede[c], Cecilie Sandberg[d], Einar Osland Vik-Mo[d,e], Hanne Roberg-Larsen[a], Bo Nyström[a], Elsa Lundanes[a], Steven Ray Wilson[a†]

[a] Department of Chemistry, University of Oslo, Post Box 1033, Blindern, NO-0315 Oslo, Norway

[b] Department of Microbiology, Unit Cell Signaling, Oslo University Hospital, Gaustadalleen 34, NO-0372 Oslo, Norway

[c] Department of Biosciences, University of Oslo, Post Box 1066, Blindern, NO-0316 Oslo, Norway

[d] Vilhelm Magnus Laboratory of Neurosurgical Research, Institute for Surgical Research and Department of Neurosurgery, Oslo University Hospital, 4950 Nydalen, NO-0424 Oslo, Norway

[e] Institute of Clinical Medicine, Faculty of Medicine, University of Oslo, Post Box 1171, Blindern, 0318 Oslo, Norway

Corresponding author: stevenw@kjemi.uio.no, +47 97010953. https://orcid.org/0000-0002-9755-1188

Abstract

Exosomes are a potentially rich source of biomarkers, but their isolation and characterization can

be challenging. For isolation of exosomes, differential ultracentrifugation (a traditional approach)

and an isolation kit from a major vendor (Total Exosome Isolation Reagent from Thermo Fisher

Scientific) were compared. "Case study" exosomes were isolated from cell culture media of two

different cell sources, namely patient-derived cells from glioblastoma multiforme and the breast

cancer cell line MDA-MB-231. For both isolation methods, transmission electron microscopy

and dynamic light scattering indicated the presence of exosomes. The kit- and UC isolates

contained similar amounts of protein measured by the bicinchoninic acid (BCA) assay with

absorbance at 562 nm. Using western blot, positive exosome markers were identified in all

isolates. Potential biomarkers for both diseases were also identified in the isolates using LC-

MS/MS. However, WB and LC-MS/MS also revealed negative exosome markers regarding both

isolation approaches. The two isolation methods had an overall similar performance, but we

hesitate to use the term "exosome isolation" as impurities may be present with both isolation

methods. LC-MS/MS can detect disease biomarkers in exosomes and is also highly useful for

critical assessment of exosome enrichments.

Key words: Exosomes; Ultracentrifugation; Proteomics; Glioblastoma; Breast cancer; LC-

MS/MS

1 Introduction

Exosomes are extracellular vesicles (EVs) with membrane-surrounded bodies which are secreted from cells to the extracellular environment as a part of the endocytic pathway [1]. Exosomes are formed by invagination of an endosome membrane to create intraluminal vesicles inside the endosome, i.e. multivesicular bodies (MVBs), and are secreted when the endosomes fuse with the plasma membrane [2]. Exosomes commonly contain proteins originating from the cellular cytosol and the plasma membrane, nucleic acids (e.g. DNA, mRNA, microRNA and non-coding RNA), lipids and metabolites [3-5,1,6-8], and are believed to take part in e.g. cell-cell communication, transfer of proteins/nucleic acids, coagulation and antigen presentation [6,9].

Cancer cells have been found to release more exosomes than stromal cells [10,11] and exosomes are associated with metastasis and tumor progression [7,12,13]. Hence, cancer exosomes may be a source of biomarkers for diagnosing cancers such as breast cancer (BC) and glioblastoma multiforme (GBM) when e.g. isolated from body fluids. BC is the predominant type of female cancer [14], with recurrent metastatic disease being responsible for the majority of BC-caused deaths [15]. GBM is the most frequent and malignant form of brain cancer [16-18]. The diagnosis of both BC and GBM rely on highly invasive patient tissue biopsies at relatively late stages [16,19,20]. Thus, a non-invasive disease monitoring is desirable for both BC and GBM, and can be achieved by measuring biomarkers in accessible body fluids, such as blood (liquid biopsy), for early diagnosis and prognosis assessment [16,21-23]. Hence, the isolation of exosomes for cancer biomarker discovery has emerged as an alternative to invasive methodologies [24-31,23].

Isolation of exosomes is predominantly performed from body fluids (e.g. blood, urine, and saliva) or cell culture media by centrifugation-based methods, e.g. sucrose density gradient centrifugation or ultracentrifugation (UC) [32,33]. However, common drawbacks of using UC-

based exosome isolation methods are the large amounts of starting material needed, low yield,

and poor reproducibility [34,35]. Moreover, there is a great need for exosome isolation protocols

tailored towards smaller starting volumes for e.g. miniaturized cell culture models like organoids

and "organ on a chip" [36,37]. Other exosome isolation protocols and principles have been

developed to overcome the drawbacks of UC based methods. Among these, filtration,

immunoaffinity capturing, size exclusion chromatography, flow field-flow fractionation and also

acoustic trapping have been attempted [34,38-42,8,43,44]. In addition, different commercial

exosome isolation kits are available (e.g. ExoQuick[TM] from Systems Biosciences, and Total

Exosome Isolation[TM] from Thermo Fisher), enabling simple isolation of exosomes from small

starting volumes from a wide range of matrices. The exosome isolation kits are known to be

based on exosome precipitation at low-speed centrifugation after sample incubation with water-

excluding polymers such as polyethylene glycol (PEG) [45].

We have compared two exosome isolation methods, namely UC and a commercial kit for

precipitation of exosomes. The methods were evaluated using the following characterization

techniques: WB, transmission electron microscopy (TEM), dynamic light scattering (DLS),

quantitative total protein analysis using UV-Vis spectrophotometry and LC-MS/MS. "Case

study" exosomes were isolated from cell culture media from free-floating patient-derived primary

cell cultures from GBM biopsies (T1018) and a serum cultivated, adherently growing BC cell

line (MDA-MB-231).

2 Materials and Methods

2.1 MDA MB-231 cell culturing

The BC cell line was purchased from American Type Culture Collection (ATCC, Sesto San Giovanni, Milan, Italy) and is derived from a triple-negative human metastatic breast carcinoma. The cells were maintained in Rosewell Park Memorial Institute (RPMI) 1640 growth medium depleted of phenol red (Sigma-Aldrich, St. Louis, MO, USA) supplemented with 10 % exosome-depleted fetal bovine serum (FBS) (System Biosciences, Palo Alto, CA, USA) and 1 % penicillin/streptomycin (Sigma- Aldrich). The cells were incubated in a humidifying atmosphere at 5 % CO_2 and at 37 $^{\circ}$C. Prior to exosome isolation, 1-2.3 million cells (in T75-T175 culturing flasks) were incubated for 6-7 days (always using a passage lower than 12). The incubated cell culture medium was centrifuged at 906 $\times$ g (30 minutes at 23 °C). See also **Supplementary 1 (S1).**

2.2 Glioblastoma cell culturing

The GBM cells (T1018) were derived from biopsies from a primary GBM tumor, obtained after informed consent through a biobank approved by the Regional Ethical Authorities operated at Oslo University Hospital (2016/1791). The cells were maintained in Dulbecco's modified eagle medium with nutrient mixture F-12 (DMEM/F12, Gibco, Thermo Fisher Scientific, Waltham, MA, USA), supplemented with HEPES buffer (10 mM) and penicillin/streptomycin (100 U/mL) from Lonza (Basel, Switzerland), B27 without vitamin A (1/50) from Thermo Fisher Scientific, epidermal growth factor (20 ng/mL) and basic fibroblast growth factor (10 ng/ mL) from R&D Systems (Minneapolis, MN, USA) and heparin (2.5 µg/mL) obtained from LEO Pharma AS

(Ballerup, Denmark). Under these culturing conditions, cells express stem cell markers *in vitro*,

differentiate upon removal of growth factors and give rise to diffusely infiltrative tumors upon

xenografting [46]. The cells were incubated in a humidifying atmosphere at 5 % CO_2 and 37 °C

in T25 flasks (Thermo Fisher Scientific). Prior to exosome isolation, the incubated cell culture

medium was centrifuged twice at 453 × *g* and 1811 × *g* for 5 minutes each. The cell pellets were

harvested for WB analysis. See also **S1.**

2.3   Exosome isolation by ultracentrifugation

For the BC and GBM cells, 9-12 mL and 60 mL cell culture media were used for centrifugation,

respectively. Cell culture media were first centrifuged at 1811 × *g* (5 minutes at 20 °C). The

supernatants were then centrifuged at 20 000 × *g* (20 minutes at 4 °C) with an Allegra 25R

centrifuge (with TA-14-50 rotor) from Beckman Coulter (Brea, CA, USA) and the supernatants

were transferred to polycarbonate ultracentrifugation tubes (Beckman Coulter) and diluted with

PBS (~60 mL in each). The tubes were centrifuged twice at 100 000 × *g* (90 minutes at 4 °C)

with an L-80 ultracentrifuge (45 Ti rotor) from Beckman Coulter. The supernatants were

removed (leaving suspension 1 cm above the pellets) and the pellets were suspended with PBS

between the centrifugations. Upon centrifugation, the supernatants were discarded and the

exosome pellets (UC isolates) were suspended in either PBS (3 mL for DLS- and 50-100 μL for

TEM analysis) or the preferred lysis buffer.

2.4   Exosome isolation by isolation kit

The isolation of exosomes with the kit was performed with the Total Exosome Isolation Reagent

(from cell culture media) from Thermo Fisher Scientific (catalog no. 4478359). The isolation was

performed according to the protocol of the supplier [47]. Starting volumes ranged from 0.5 mL to

9 mL cell culture medium for the BC cells and 5 mL to 6 mL for the GBM cells. The samples

were centrifuged with the Allegra 25R centrifuge, and the exosome pellets (kit isolates) were

suspended as with UC.

2.5   Protein extraction

Cell and exosome protein extracts were made by lysis with RIPA- or Nonidet™ P40 (NP40)

buffer (both from Thermo Fisher Scientific) containing protease inhibitors (Protease Inhibitor

Cocktail Tablets, Roche, Basel, Switzerland) and phosphatase inhibitors (PhosStop Tablets,

Sigma-Aldrich). See also **S2.**

2.6   UV-Vis spectrophotometry

The protein amount was measured using Pierce™ BCA protein Assay Kit (Thermo Fisher

Scientific), by measuring the absorbance at 562 nm. See also **S3.**

2.7   Western blotting

For information about WB antibodies, procedures and equipment, see **S4.**

2.8   Immunogold labeling and transmission electron microscopy

Samples were visualized with a JEM-1400Plus transmission electron microscope from JEOL

(Tokyo, Japan) and images were recorded at 80 kV. See also **S5.**

2.9 Dynamic light scattering

The DLS experiments were conducted with the aid of an ALV/CGS-8F multi-detector version

compact goniometer system, with 8 fiber-optical detection units, from ALV-GmbH, Langen,

Germany. See **S6** for more details.

2.10 LC-MS/MS analysis

LC-MS/MS was performed using Q-Exactive mass spectrometers (Thermo) coupled with liquid

nano chromatography. Samples were prepared by in-solution and in-gel protease digestion. See

S7-9 for additional information related to LC-MS/MS analysis**.**

3 Results and Discussion

3.1 Similar content of protein measured in kit- and UC isolates

The protein amount per million cells (hereafter referred to as protein amount) in the BC- (**Figure**

1A) and GBM- (**Figure 1B**) isolates was measured using UV-Vis spectrophotometry. The

measurements for kit isolates were 15-28 times higher than for UC isolates. A higher protein

amount in exosomes isolated by the kit compared to that by UC was also observed in a study by

Van Deun et al. who compared UC to the same isolation kit used in the present study for MCF7

derived exosomes [48]. However, we observed that the measured absorbance in the kit blanks

was high in comparison to UC blanks, where the absorbance was below the limit of

quantification. The high absorbance from the kit blanks was further assessed to establish possible

UV-absorbents or scattering components in the kit reagent. However, no absorbance was

measured in the kit reagent using the same protocol (i.e. absorbance at 562 nm after BCA-

reaction) as for the isolates and blanks, and NMR spectroscopy showed sharp peaks implying an

absence of relaxation-perturbing components, e.g. particles (results not shown). The high

absorbance in the kit blanks might therefore indicate co-precipitation of proteins or other UV-

absorbing compounds from the blank media. When correcting for the blank (subtracting the

protein amount measured in blank samples from the protein amount in exosome isolates), the

measured protein content for exosomes isolated by the kit and UC was similar.

3.2 TEM and DLS detected vesicles in the expected size range for exosomes

Morphological analysis of the exosome samples was performed using TEM. In addition, the

hydrodynamic particle size distribution was measured using DLS. Clusters of vesicles were

observed in the micrographs of the samples isolated with both kit and UC (**Figure 2, AI and**

AIII). Vesicle structures similar to that described in literature were observed [49,50,6].

Regarding GBM exosomes: With TEM, the UC isolates presented somewhat more distinct

double membranes compared to the kit isolates. The blank samples for both isolation methods did

not display membrane structures (**Figure 2, AII** and **AIV**). The DLS-analysis of the GBM

isolates exhibited particles of similar sizes of 51 and 73 nm (mean) with both isolation methods

(**Figure 2B**). Thus, both isolation methods gave rise to comparable exosome populations.

Regarding BC exosomes: Clusters of vesicles were also in here observed in the micrographs of

the samples isolated with both kit and UC (**Figure 2, CI and CIII**). Blank isolates displayed

contaminations (**Figure 2, CII and CIV**), e.g. exosome-resembling vesicles were found in the

UC blank using TEM (red dashed circles), and the kit blank displayed 67 nm (mean)

contaminations when using DLS (**Figure 2D**). The DLS analysis also presented two distinct

particle diameters in kit isolates (28 and 95 nm, mean values) while only one particle diameter

was present in UC isolates (137 nm, mean value), indicating some differences in the mean

particle sizes isolated with the two isolation methods. However, the sizes observed with DLS

correlates well with that found in other studies (30–250 nm) [51,52,13,53,54,48,55]. Overall, the

isolates showed structures resembling those of EVs, but some blanks were not entirely devoid of

vesicles or particles.

3.3. Western blot analyses indicated the presence of exosomes for all samples.

WB was performed using antibodies for a selection of positive exosome markers, namely the

tetraspanins CD81, CD9 and CD63, TSG101 and flotillin-1. Calnexin was selected as a negative

marker for purity evaluation as recommended by the International Society of Extracellular

vesicles (ISEV) [56]. This protein is located at the endoplasmic reticulum (ER) and is assumed to

signalize ER-contamination. For the GBM cells and exosomes, positive and negative exosome

markers were detected in isolates from both the kit and UC (**Figure 3**). For the BC cells and

exosomes, positive markers TSG101, flotillin-1 and CD9 (barely visible in the UC isolates) were

detected using both isolation methods, and calnexin was not detected. The positive markers

overall demonstrate the presence of exosomes in the isolates obtained using both methods, but the

GBM samples could contain impurities.

220
3.4 LC-MS/MS studies reveal impurities, and biomarkers

The absence of calnexin (see above) in BC exosomes from both isolation methods indicates that

the isolates are not contaminated with the ER. However, general proteins related to e.g. the

nucleus, Golgi apparatus, mitochondrion, and ER were identified in the BC exosomes using LC-

MS/MS and gene ontology (GO) annotations (**Figure 4**). Hence, untargeted LC-MS/MS

suggested the presence of impurities also in the BC samples. Proteins related to the nucleosome,

Golgi apparatus, mitochondrion, and ER were also identified by GO-annotation in the GBM

isolates.

LC-MS/MS could also identify a number of positive markers (see **Figure 5** for examples).

However, there was expectedly not a complete overlap with those observed with WB, as e.g.

sensitivity can vary between WB and untargeted LC-MS/MS. In-house prepared nanoLC

columns packed with core shell particles provided high-resolution separations (**Figure 5**, and see

reference [57] for packing procedure). Examples of potential biomarkers for GBM, e.g. heat

shock proteins 70 kDa and 90 kDa [58-60], chondroitin sulfate proteoglycan 4 [58,61], CD44

[62,58,61] and CD276 [63] were identified using LC-MS/MS. Examples of LC-MS/MS-detected

biomarkers related to triple negative breast cancer were e.g. heat shock 90 kDa α and β protein

[64], calmodulin and epithermal growth factor receptor [65] (see **Supplemental Proteins**). When

comparing cell sources, the number of identified proteins was lower in GBM isolates than BC

isolates, but the number of identified proteins for GBM isolates was comparable to another LC-

MS/MS study on GBM exosomes [66].

4 Conclusions

Regarding our glioblastoma/breast cancer "case study" samples, the UC/kit isolation methods

overall were approximately equal in quality. Kit isolation however has an advantage of requiring

less starting material compared to conventional UC equipment. Untargeted LC-MS/MS revealed

a number of biomarkers related to the diseases, supporting the concept of exosomes being an

interesting matrix towards diagnostics. In addition to exosomes, our analyses suggest the

presence of cellular contaminations and other vesicles. Hence, the "isolations" should perhaps be

considered "enrichments". Considering that the methods do not fully provide isolations, we

welcome alternative approaches to preparing and analyzing these important extracellular vesicles.

Acknowledgments

This work was supported by the Department of Chemistry, University of Oslo, Norway. We

would like to acknowledge DIATECH@UiO, since parts of this work have been carried out

within this strategic research initiative at the Faculty of Mathematics and Natural Sciences,

University of Oslo. This work has also been supported by the UiO:Life Science funded

convergence environment "Organ on a chip and nano-devices". Conflict of Interest: The authors

declare that they have no conflict of interest.

Figure Captions

Figure 1 Measured relative protein amount pr. million cells in exosome samples from GBM- and BC cells isolated by kit and UC (n ≥ 2). A) The measured relative protein amount (%) for the BC exosome isolates. **B**) The measured relative protein amount (%) for the GBM exosome isolates. Each replicate is depicted as circles, and the median depicted as a line. The X-mark shows the measured relative protein amount in the blank sample (isolated cell culture medium). The protein amounts were measured by UV-Vis spectrophotometry (absorption at $\lambda = 562$ nm) after reaction with BCA kit reagents.

Figure 2 Transmission electron micrographs and hydrodynamic particle size (nm) distribution by DLS analysis of exosomes isolated by kit and UC from GBM- and BC cells.

Images were taken with a magnification of 400 000, and the dashed areas were additionally zoomed. **A**) Micrographs of GBM exosome isolates (not CD9-labelled). **I** depict the micrograph from a kit isolate, **II** the kit blank, **III** a UC isolate, and **IV** the UC blank. **B**) DLS analysis of GBM exosomes isolated by kit and UC (n = 1). No particles were detected in the UC blank (n = 1). DLS analysis of the kit blank was not performed. **C**) Micrographs of BC exosome isolates (successfully CD9-labelled). **I** depict the micrograph from a kit isolate, **II** the kit blank, **III** a UC isolate, and **IV** the UC blank. **D**) DLS analysis of BC exosomes isolated by kit (n = 2) and UC (n = 3), including the kit blank (n = 1). No particles were detected in the UC blank.

Figure 3 Western blot of common protein exosome markers. The protein markers CD81, CD9, CD63, TSG101, flotillin-1 (positive markers, +) and calnexin (negative marker, -) were targeted in cell lysates and exosomes isolated by kit and UC (n ≥ 2). Monoclonal mouse antibodies were used for CD81, CD9, CD63, flotillin-1 and calnexin, while a polyclonal rabbit antibody was used for TSG101. For the BC exosomes, 15 µg protein was loaded for kit isolates and 3 µg for UC isolates. For the GBM exosomes, ~14 µg was loaded for kit isolates and ~8 µg for UC isolates. Uncropped western blots are presented in **Supplemental Western Blots**.

Figure 4 Chromatograms and MS/MS spectrums from LC-MS/MS analysis of GBM- and BC exosome peptides. **A**) Chromatogram with corresponding MS/MS spectrum for the CD9 signature peptide KDVLETFTVK (*m/z*=393.89, z=3) in BC exosomes isolated by UC. **C**) Chromatogram with corresponding MS/MS spectrum for the calnexin signature peptide AEEDEILNR (*m/z*=544.77, z=2) from GBM exosomes isolated by UC. An in-house packed 50 µm x 150 mm column with 80 Å Accucore particles with C_{18} stationary phase was used for separation. A 50 µm x ~3 mm in-house packed pre-column with the same column material was

used for trapping. The elution was performed with a linear gradient of 3-15 % MP B in 120

minutes. See **Section 2.11.1** for more LC-MS/MS parameters.

Figure 5 GO annotation of proteins in BC exosomes to different cellular locations. The

identified proteins classified by their cellular location (GO annotations) grouped based on their

positive/ negative relevance towards exosomes. The annotated proteins (% of total proteins) and

their cellular location, with proteins annotated from the kit isolates are shown in red (from 749

DAVID ID's), while proteins annotated from the UC isolates are shown in blue (from 615

DAVID ID's).

Figure 6 Venn diagram presenting the number of proteins identified by LC-MS/MS in

exosomes isolated by kit and UC from GBM- and BC cell culture medium. The numbers are

the total number of unique proteins identified when trypsin, keratin related proteins and the

proteins identified in blank isolates were disregarded. One signature peptide was selected as

requirement for positive identifications during database search. Equal amounts of protein were

injected for both kit- and UC isolates (~ 1.5 µg protein for GBM isolates (n = 6) and ~2-5 µg

protein for BC exosomes (n=3)). A list of all proteins identified is presented in **Supplemental**

Proteins.

References

317   1. Théry C, Zitvogel L, Amigorena S (2002) Exosomes: composition, biogenesis and function.
318   Nature Reviews Immunology 2:569. doi:10.1038/nri855
319   2. Lener T, Gimona M, Aigner L, Borger V, Buzas E, Camussi G, Chaput N, Chatterjee D, Court FA,
320   del Portillo HA, O'Driscoll L, Fais S, Falcon-Perez JM, Felderhoff-Mueser U, Fraile L, Gho YS,

Gorgens A, Gupta RC, Hendrix A, Hermann DM, Hill AF, Hochberg F, Horn PA, de Kleijn D, Kordelas L, Kramer BW, Kramer-Albers EM, Laner-Plamberger S, Laitinen S, Leonardi T, Lorenowicz MJ, Lim SK, Lotvall J, Maguire CA, Marcilla A, Nazarenko I, Ochiya T, Patel T, Pedersen S, Pocsfalvi G, Pluchino S, Quesenberry P, Reischl IG, Rivera FJ, Sanzenbacher R, Schallmoser K, Slaper-Cortenbach I, Strunk D, Tonn T, Vader P, van Balkom BWM, Wauben M, El Andaloussi S, Thery C, Rohde E, Giebel B (2015) Applying extracellular vesicles based therapeutics in clinical trials - an ISEV position paper. J Extracell Vesicles 4:31. doi:10.3402/jev.v4.30087

3. Zhao H, Yang L, Baddour J, Achreja A, Bernard V, Moss T, Marini JC, Tudawe T, Seviour EG, San Lucas FA, Alvarez H, Gupta S, Maiti SN, Cooper L, Peehl D, Ram PT, Maitra A, Nagrath D (2016) Tumor microenvironment derived exosomes pleiotropically modulate cancer cell metabolism. eLife 5:e10250. doi:10.7554/eLife.10250

4. Skotland T, Sandvig K, Llorente A (2017) Lipids in exosomes: Current knowledge and the way forward. Progress in Lipid Research 66:30-41. doi:https://doi.org/10.1016/j.plipres.2017.03.001

5. Beach A, Zhang H-G, Ratajczak MZ, Kakar SS (2014) Exosomes: an overview of biogenesis, composition and role in ovarian cancer. Journal of Ovarian Research 7:14-14. doi:10.1186/1757-2215-7-14

6. Li P, Kaslan M, Lee SH, Yao J, Gao Z (2017) Progress in Exosome Isolation Techniques. Theranostics 7 (3):789-804. doi:10.7150/thno.18133

7. Becker A, Thakur BK, Weiss JM, Kim HS, Peinado H, Lyden D (2016) Extracellular Vesicles in Cancer: Cell-to-Cell Mediators of Metastasis. Cancer Cell 30 (6):836-848. doi:https://doi.org/10.1016/j.ccell.2016.10.009

8. Pocsfalvi G, Stanly C, Fiume I, Vekey K (2016) Chromatography and its hyphenation to mass spectrometry for extracellular vesicle analysis. J Chromatogr A 1439:26-41. doi:10.1016/j.chroma.2016.01.017

9. Farahani M, Rubbi C, Liu L, Slupsky JR, Kalakonda N (2015) CLL exosomes modulate the transcriptome and behaviour of recipient stromal cells and are selectively enriched in miR-202-3p. PloS one 10 (10):e0141429

10. Logozzi M, De Milito A, Lugini L, Borghi M, Calabro L, Spada M, Perdicchio M, Marino ML, Federici C, Iessi E, Brambilla D, Venturi G, Lozupone F, Santinami M, Huber V, Maio M, Rivoltini L, Fais S (2009) High Levels of Exosomes Expressing CD63 and Caveolin-1 in Plasma of Melanoma Patients. Plos One 4 (4):10. doi:10.1371/journal.pone.0005219

11. Parolini I, Federici C, Raggi C, Lugini L, Palleschi S, De Milito A, Coscia C, Iessi E, Logozzi M, Molinari A, Colone M, Tatti M, Sargiacomo M, Fais S (2009) Microenvironmental pH Is a Key Factor for Exosome Traffic in Tumor Cells. J Biol Chem 284 (49):34211-34222. doi:10.1074/jbc.M109.041152

12. Peinado H, Alečković M, Lavotshkin S, Matei I, Costa-Silva B, Moreno-Bueno G, Hergueta-Redondo M, Williams C, García-Santos G, Ghajar CM (2012) Melanoma exosomes educate bone marrow progenitor cells toward a pro-metastatic phenotype through MET. Nature medicine 18 (6):883-891

13. Harris DA, Patel SH, Gucek M, Hendrix A, Westbroek W, Taraska JW (2015) Exosomes released from breast cancer carcinomas stimulate cell movement. PloS one 10 (3):e0117495

14. DeSantis CE, Lin CC, Mariotto AB, Siegel RL, Stein KD, Kramer JL, Alteri R, Robbins AS, Jemal A
(2014) Cancer treatment and survivorship statistics, 2014. CA: A Cancer Journal for Clinicians 64
(4):252-271. doi:10.3322/caac.21235
15. Redig AJ, McAllister SS (2013) Breast cancer as a systemic disease: a view of metastasis.
Journal of Internal Medicine 274 (2):113-126. doi:10.1111/joim.12084
16. Touat M, Duran-Pena A, Alentorn A, Lacroix L, Massard C, Idbaih A (2015) Emerging
circulating biomarkers in glioblastoma: promises and challenges. Expert Rev Mol Diagn 15
(10):1311-1323. doi:10.1586/14737159.2015.1087315
17. Molina JR, Hayashi Y, Stephens C, Georgescu MM (2010) Invasive Glioblastoma Cells Acquire
Stemness and Increased Akt Activation. Neoplasia 12 (6):453-U437. doi:10.1593/neo.10126
18. Preusser M, de Ribaupierre S, Wohrer A, Erridge SC, Hegi M, Weller M, Stupp R (2011)
Current Concepts and Management of Glioblastoma. Ann Neurol 70 (1):9-21.
doi:10.1002/ana.22425
19. Jan Chrastina ZN, Milan Brazdil, Marketa Hermanova (2015) Glioblastoma multiforme in a
patient with isolated hemimegalencephaly. Journal of Neurological Surgery Reports 76 (1):160-
163. doi:ht tp://dx.doi.org/
10.1055/s -0035-1554929.
20. Shao HL, Chung J, Lee K, Balaj L, Min C, Carter BS, Hochberg FH, Breakefield XO, Lee H,
Weissleder R (2015) Chip-based analysis of exosomal mRNA mediating drug resistance in
glioblastoma. Nat Commun 6:9. doi:10.1038/ncomms7999
21. Best MG, Sol N, Zijl S, Reijneveld JC, Wesseling P, Wurdinger T (2015) Liquid biopsies in
patients with diffuse glioma. Acta Neuropathol 129 (6):849-865. doi:10.1007/s00401-015-1399-
y
22. Cristofanilli M, Budd GT, Ellis MJ, Stopeck A, Matera J, Miller MC, Reuben JM, Doyle GV,
Allard WJ, Terstappen LW (2004) Circulating tumor cells, disease progression, and survival in
metastatic breast cancer. New England Journal of Medicine 351 (8):781-791
23. He M, Zeng Y (2016) Microfluidic Exosome Analysis toward Liquid Biopsy for Cancer. Journal
of laboratory automation 21 (4):599-608. doi:10.1177/2211068216651035
24. Melo SA, Luecke LB, Kahlert C, Fernandez AF, Gammon ST, Kaye J, LeBleu VS, Mittendorf EA,
Weitz J, Rahbari N (2015) Glypican-1 identifies cancer exosomes and detects early pancreatic
cancer. Nature 523 (7559):177-182
25. Taylor DD, Gercel-Taylor C (2008) MicroRNA signatures of tumor-derived exosomes as
diagnostic biomarkers of ovarian cancer. Gynecologic Oncology 110 (1):13-21
26. Roberg-Larsen H, Lund K, Seterdal KE, Solheim S, Vehus T, Solberg N, Krauss S, Lundanes E,
Wilson SR (2017) Mass spectrometric detection of 27-hydroxycholesterol in breast cancer
exosomes. The Journal of steroid biochemistry and molecular biology 169:22-28
27. Skog J, Würdinger T, Van Rijn S, Meijer DH, Gainche L, Curry WT, Carter BS, Krichevsky AM,
Breakefield XO (2008) Glioblastoma microvesicles transport RNA and proteins that promote
tumour growth and provide diagnostic biomarkers. Nature cell biology 10 (12):1470-1476
28. Hochberg FH, Atai NA, Gonda D, Hughes MS, Mawejje B, Balaj L, Carter RS (2014) Glioma
diagnostics and biomarkers: an ongoing challenge in the field of medicine and science. Expert
Rev Mol Diagn 14 (4):439-452. doi:10.1586/14737159.2014.905202

29. Saadatpour L, Fadaee E, Fadaei S, Mansour RN, Mohammadi M, Mousavi SM, Goodarzi M,
Verdi J, Mirzaei H (2016) Glioblastoma: exosome and microRNA as novel diagnosis biomarkers.
Cancer Gene Ther 23 (12):415-418. doi:10.1038/cgt.2016.48
30. Mondal A, Singh DK, Panda S, Shiras A (2017) Extracellular Vesicles As Modulators of Tumor
Microenvironment and Disease Progression in Glioma. Front Oncol 7:8.
doi:10.3389/fonc.2017.00144
31. Keller S, Ridinger J, Rupp A-K, Janssen JW, Altevogt P (2011) Body fluid derived exosomes as
a novel template for clinical diagnostics. Journal of translational medicine 9 (1):86
32. Théry C, Amigorena S, Raposo G, Clayton A (2006) Isolation and characterization of
exosomes from cell culture supernatants and biological fluids. Current Protocols in Cell
Biology:3.22. 21-23.22. 29
33. Cantin R, Diou J, Bélanger D, Tremblay AM, Gilbert C (2008) Discrimination between
exosomes and HIV-1: purification of both vesicles from cell-free supernatants. Journal of
Immunological Methods 338 (1):21-30
34. Greening DW, Xu R, Ji H, Tauro BJ, Simpson RJ (2015) A protocol for exosome isolation and
characterization: evaluation of ultracentrifugation, density-gradient separation, and
immunoaffinity capture methods. Proteomic Profiling: Methods and Protocols:179-209
35. Cvjetkovic A, Lötvall J, Lässer C (2014) The influence of rotor type and centrifugation time on
the yield and purity of extracellular vesicles. Journal of Extracellular Vesicles 3:DOI:
10.3402/jev.v3403.23111. doi:10.3402/jev.v3.23111
36. Tauro BJ, Greening DW, Mathias RA, Mathivanan S, Ji H, Simpson RJ (2013) Two distinct
populations of exosomes are released from LIM1863 colon carcinoma cell-derived organoids.
Molecular & Cellular Proteomics 12 (3):587-598
37. Bhatia SN, Ingber DE (2014) Microfluidic organs-on-chips. Nature Biotechnology 32 (8):760-
772
38. Böing AN, Van Der Pol E, Grootemaat AE, Coumans FA, Sturk A, Nieuwland R (2014) Single-
step isolation of extracellular vesicles by size-exclusion chromatography. Journal of Extracellular
Vesicles 3:DOI: 10.3402/jev.v3403.23430
39. Gámez-Valero A, Monguió-Tortajada M, Carreras-Planella L (2016) Size-Exclusion
Chromatography-based isolation minimally alters Extracellular Vesicles' characteristics
compared to precipitating agents. Scientific Reports 6:DOI: 10.1038/srep33641.
doi:10.1038/srep33641
40. Grant R, Ansa-Addo E, Stratton D, Antwi-Baffour S, Jorfi S, Kholia S, Krige L, Lange S, Inal J
(2011) A filtration-based protocol to isolate human plasma membrane-derived vesicles and
exosomes from blood plasma. Journal of Immunological Methods 371 (1):143-151
41. Clayton A, Court J, Navabi H, Adams M, Mason MD, Hobot JA, Newman GR, Jasani B (2001)
Analysis of antigen presenting cell derived exosomes, based on immuno-magnetic isolation and
flow cytometry. Journal of Immunological Methods 247 (1):163-174
42. Yang JS, Lee JC, Byeon SK, Rha KH, Moon MH (2017) Size dependent lipidomic analysis of
urinary exosomes from patients with prostate cancer by flow field-flow fractionation and
nanoflow liquid chromatography-tandem mass spectrometry. Analytical Chemistry 89:2488-
2496

43. Abramowicz A, Widlak P, Pietrowska M (2016) Proteomic analysis of exosomal cargo: the
challenge of high purity vesicle isolation. Mol Biosyst 12 (5):1407-1419.
doi:10.1039/c6mb00082g
44. Ku A, Lim HC, Evander M, Lilja H, Laurell T, Scheding S, Ceder Y (2018) Rapid Acoustic
Enrichment of Extracellular Vesicles from Biological Fluids. Analytical Chemistry
45. Rider MA, Hurwitz SN, Meckes Jr DG (2016) ExtraPEG: a polyethylene glycol-based method
for enrichment of extracellular vesicles. Scientific reports 6:23978
46. Vik-Mo EO, Sandberg C, Olstorn H, Varghese M, Brandal P, Ramm-Pettersen J, Murrell W,
Langmoen IA (2010) Brain tumor stem cells maintain overall phenotype and tumorigenicity after
in vitro culturing in serum-free conditions. Neuro-Oncology 12 (12):1220-1230.
doi:10.1093/neuonc/noq102
47. ThermoFisher Total Exosome Isolation Reagent (from cell culture media).
https://www.thermofisher.com/order/catalog/product/4478359?SID=srch-srp-4478359.
48. Van Deun J, Mestdagh P, Sormunen R, Cocquyt V, Vermaelen K, Vandesompele J, Bracke M,
De Wever O, Hendrix A (2014) The impact of disparate isolation methods for extracellular
vesicles on downstream RNA profiling. J Extracell Vesicles 3 (1):24858. doi:10.3402/jev.v3.24858
49. Pan B-T, Teng K, Wu C, Adam M, Johnstone RM (1985) Electron microscopic evidence for
externalization of the transferrin receptor in vesicular form in sheep reticulocytes. The Journal
of Cell Biology 101 (3):942-948. doi:10.1083/jcb.101.3.942
50. Pan B-T, Johnstone RM (1983) Fate of the transferrin receptor during maturation of sheep
reticulocytes in vitro: selective externalization of the receptor. Cell 33 (3):967-978
51. Lobb RJ, Becker M, Wen SW, Wong CSF, Wiegmans AP, Leimgruber A, Moller A (2015)
Optimized exosome isolation protocol for cell culture supernatant and human plasma. J
Extracell Vesicles 4:11. doi:10.3402/jev.v4.27031
52. Jenjaroenpun P, Kremenska Y, Nair VM, Kremenskoy M, Joseph B, Kurochkin IV (2013)
Characterization of RNA in exosomes secreted by human breast cancer cell lines using next-
generation sequencing. PeerJ 1:e201. doi:10.7717/peerj.201
53. Palazzolo G, Albanese NN, Di Cara G, Gygax D, Vittorelli ML, Pucci-Minafra I (2012)
Proteomic analysis of exosome-like vesicles derived from breast cancer cells. Anticancer
Research 32 (3):847-860
54. ThermoFisherScientific (2016) Boldly go above and beyond.
https://tools.thermofisher.com/content/sfs/brochures/Exosomes-brochure.pdf. Accessed
COL12585 0916
55. Liu F, Vermesh O, Mani V, Ge TJ, Madsen SJ, Sabour A, Hsu E-C, Gowrishankar G, Kanada M,
Jokerst JV, Sierra RG, Chang E, Lau K, Sridhar K, Bermudez A, Pitteri SJ, Stoyanova T, Sinclair R,
Nair VS, Gambhir SS, Demirci U (2017) The Exosome Total Isolation Chip. ACS Nano 11
(11):10712-10723. doi:10.1021/acsnano.7b04878
56. Lötvall J, Hill AF, Hochberg F, Buzás EI, Di Vizio D, Gardiner C, Gho YS, Kurochkin IV,
Mathivanan S, Quesenberry P (2014) Minimal experimental requirements for definition of
extracellular vesicles and their functions: a position statement from the International Society for
Extracellular Vesicles. Journal of Extracellular Vesicles 3:26913-23078
57. Berg HS, Seterdal KE, Smetop T, Rozenvalds R, Brandtzaeg OK, Vehus T, Lundanes E, Wilson
SR (2017) Self-packed core shell nano liquid chromatography columns and silica-based
monolithic trap columns for targeted proteomics. Journal of Chromatography A 1498:111-119

58. Redzic JS, Ung TH, Graner MW (2014) Glioblastoma extracellular vesicles: reservoirs of
potential biomarkers. Pharmacogenomics and personalized medicine 7:65-77.
doi:10.2147/pgpm.s39768
59. Thuringer D, Hammann A, Benikhlef N, Fourmaux E, Bouchot A, Wettstein G, Solary E,
Garrido C (2011) Transactivation of the epidermal growth factor receptor by heat shock protein
90 via Toll-like receptor 4 contributes to the migration of glioblastoma cells. The Journal of
biological chemistry 286 (5):3418-3428. doi:10.1074/jbc.M110.154823
60. Graner MW, Cumming RI, Bigner DD (2007) The Heat Shock Response and Chaperones/Heat
Shock Proteins in Brain Tumors: Surface Expression, Release, and Possible Immune
Consequences. The Journal of Neuroscience 27 (42):11214-11227. doi:10.1523/jneurosci.3588-
07.2007
61. Wade A, Robinson AE, Engler JR, Petritsch C, James CD, Phillips JJ (2013) Proteoglycans and
their roles in brain cancer. The FEBS journal 280 (10):2399-2417. doi:10.1111/febs.12109
62. Mooney KL, Choy W, Sidhu S, Pelargos P, Bui TT, Voth B, Barnette N, Yang I (2016) The role
of CD44 in glioblastoma multiforme. Journal of Clinical Neuroscience 34:1-5.
doi:https://doi.org/10.1016/j.jocn.2016.05.012
63. Kraan J, van den Broek P, Verhoef C, Grunhagen DJ, Taal W, Gratama JW, Sleijfer S (2014)
Endothelial CD276 (B7-H3) expression is increased in human malignancies and distinguishes
between normal and tumour-derived circulating endothelial cells. British journal of cancer 111
(1):149-156. doi:10.1038/bjc.2014.286
64. Beliakoff J, Whitesell L (2004) Hsp90: an emerging target for breast cancer therapy. Anti-
Cancer Drugs 15 (7):651-662
65. Price JT, Tiganis T, Agarwal A, Djakiew D, Thompson EW (1999) Epidermal growth factor
promotes MDA-MB-231 breast cancer cell migration through a phosphatidylinositol 3′-kinase
and phospholipase C-dependent mechanism. Cancer Research 59 (21):5475-5478
66. de Vrij J, Maas SLN, Kwappenberg KMC, Schnoor R, Kleijn A, Dekker L, Luider TM, de Witte
LD, Litjens M, van Strien ME, Hol EM, Kroonen J, Robe PA, Lamfers ML, Schilham MW, Broekman
MLD (2015) Glioblastoma-derived extracellular vesicles modify the phenotype of monocytic
cells. International Journal of Cancer 137 (7):1630-1642. doi:10.1002/ijc.29521