Critical comparison of ultracentrifugation and a commercial kit for isolation of

exosomes derived from glioblastoma and breast cancer cell lines

Frøydis Sved Skottvoll[a]*****, Henriette Engen Berg[a]*****, Kamilla Bjørseth[a], Kaja Lund[b], Norbert

Roos[c], Sara Bekhradnia[a], Bernd Thiede[c], Hanne Roberg-Larsen[a], Bo Nyström[a], Elsa

Lundanes[a], Steven Ray Wilson[a]**

[a]Department of Chemistry, University of Oslo, Post Box 1033, Blindern, NO-0315 Oslo, Norway

[b]Department of Microbiology, Unit Cell Signaling, Oslo University Hostpital, Gaustadalleen 34,

NO-0372 Oslo, Norway

[c]Department of Biosciences, University of Oslo, Post Box 1066, Blindern, NO-0316 Oslo, Norway

* These authors contributed equally to the work.

** stevenw@kjemi.uio.no, +47 97010953

Abbreviations: ABC, ammonium bicarbonate; ACN, acetonitrile; AGC, automatic gain

control; BC, breast cancer; BSA, bovine serum albumin; dd, data-dependent; DLS, dynamic

light scattering; DTT, dithiothreitol; ECL, enhanced chemiluminescence; ER, endoplasmic

reticulum; EVs, extracellular vesicles; FA, formic acid; FBS, fetal bovine serum; FSG, fish

serum gelatine; GBM, glioblastoma multiforme; GO, gene ontology; IAM, 2-iodoacetamide;

ISEV, The International Society of Extracellular Vesicles; LC-MS/MS, liquid

chromatography tandem mass spectrometry; MP, mobile phase; MS, mass spectrometry;

MVBs, multivesicular bodies; nuclear magnetic resonance, NMR; PBS, phosphate buffered

saline; RPMI, Rosewell Park Memorial Institute; RT, room temperature; S/N, signal to noise

ratio; TEM, transmission electron microscopy; TFA, trifluoroacetic acid; UC,

ultracentrifugation; WB, Western blot

Key words: *Exosomes, Ultracentrifugation, LC-MS, Proteomics, Glioblastoma, Breast*

cancer

ABSTRACT

Exosomes are small extracellular vesicles (around 30-100 nm in diameter) that are secreted from cells and can be found in most body fluids. Exosomes can be a vital source of biomarkers as they contain various substances (e.g. lipids, RNAs, metabolites and proteins) that can reflect the cell of origin (e.g. cancer cells). For isolating exosomes present in biological matrices, ultracentrifugation (UC)-based procedures are most common. Other approaches exist, including commercial kits developed for easy and low sample volume isolation. In this study, the performance of differential UC and an isolation kit from a major vendor (Total Exosome Isolation Reagent from Thermo Fisher) were compared. Exosomes were isolated from cell culture media of two different cell lines (patient derived cells from glioblastoma multiforme and the breast cancer cell line MDA-MB-231). Transmission electron microscopy (TEM), dynamic light scattering (DLS) and Western blot (WB) indicated the presence of exosomes for both isolation methods. The isolation kit (using <10 % of the sample volume than for UC) and UC contained similar amounts of protein determined by bicinchoninic acid assay (BCA) with absorbance at 562 nm. Using WB, positive exosome markers were identified in all samples. Using mass spectrometry (MS)-based proteomics, additional exosome markers were identified. Regarding glioblastoma exosomes, the number of proteins identified with MS was higher for the UC samples when injecting equal protein amounts, opposite to the breast cancer exosomes. However, negative markers could also be found using MS. Thus, we are cautious to use the term "exosome isolation" as impurities may be present regarding both procedures. Notably, known potential biomarkers for both diseases were detected in the samples using MS. In our opinion, the two isolation methods had rather similar performance, although with some minor differences based on cell of origin.

1 Introduction

Exosomes are extracellular vesicles (EVs) with small membrane-bound bodies of 30-100 nm sizes which are secreted from cells to the extracellular environment as a part of the endocytic pathway [1]. Exosomes are formed by invagination of an endosome membrane to create intraluminal vesicles inside the endosome (multivesicular bodies (MVBs)) and are secreted when the endosomes fuse with the plasma membrane [2]. Exosomes commonly contain proteins originating from the cellular cytosol and the plasma membrane (for both common proteins and proteins specific to the cell type), nucleic acids (e.g. DNA, mRNA, microRNA and non-coding RNA), lipids and metabolites [1, 3-8], and are believed to take part in e.g. cell-cell communication, transfer of proteins/nucleic acids, coagulation and antigen presentation [6, 9].

Cancer cells have been found to release more exosomes than stromal cells [10, 11] and are linked to metastasis and tumor progression [7, 12, 13]. Hence, cancer exosomes, when e.g. isolated from body fluids, may be a source of biomarkers for diagnosing cancers such as breast cancer (BC) and glioblastoma multiforme (GBM). BC is the most dominant type of cancer in women [14], with recurrent metastatic disease being responsible for the majority of BC deaths [15]. GBM is the most frequent occurring form of malignant glioma (tumor originating in glial cells in the brain) in adults and is a highly aggressive astrocytoma [16-18]. The diagnosis of both BC and GBM rely on highly invasive patient tissue biopsies at relatively late stages [16, 19, 20]. Thus, a non-invasive disease monitoring e.g. for early diagnosis and prognosis assessment using biomarkers in more accessible body fluids such as blood (liquid biopsy) is desirable for both GBM and BC [16, 21, 22]. Hence, the isolation of exosomes for cancer biomarker discovery has emerged [23-30].

The isolation of exosomes is mostly performed from body fluids (e.g. blood, urine, and saliva) or cell culture media by centrifugation-based methods (e.g. sucrose gradient ultracentrifugation and ultracentrifugation, UC) [31, 32]. In addition, other isolation protocols and principles have been developed to overcome the drawbacks of ultracentrifugation such as large amounts of starting volume needed and poor reproducibility [8, 33-39]. Moreover, there is a great need for exosome isolation protocols to tailor towards the isolation of smaller starting volumes ($< \mu L$) for e.g. miniaturized cell culture models like organoids and "organ on a chip" [40, 41]. However, there is a lack of consensus between the laboratories as the methods for rigorous isolation are still largely empirical.

The protein content of exosomes has been previously extensively characterized using Western

blotting, but also by liquid chromatography tandem mass spectrometry (LC-MS/MS) [42-44].

Tetraspanins (e.g. CD9, CD63 and CD81) have been used as positive exosome protein

markers for targeted analysis as they are particularly known to be enriched in exosomes

compared to cells [1, 45-48]. These positive exosome markers generally take part in exosome

biogenesis, and are hence expected to be present in, but are not specific to, exosomes [49]. A

large range of exosome markers (both positive and negative) for exosome characterization are

presented by The International Society of Extracellular Vesicles (ISEV) [45]. In addition,

larger protein databases covering occurring proteins in exosomes and other extracellular

vesicles are available [50-52]. ISEV also recommends using other characterization methods in

addition to WB and LC-MS/MS, to study the heterogeneity and morphology of vesicles

present in the isolated sample.

In the present study, we have compared two exosome isolation methods (UC and a Total

Exosome Isolation kit from Thermo Fisher). The methods were evaluated using

characterization techniques recommended by ISEV: WB, transmission electron microscopy

(TEM), dynamic light scattering (DLS), quantitative total protein analysis using UV-Vis

spectrophotometry and LC-MS/MS for untargeted proteomic analysis. Exosomes were

isolated from cell culture media from patient-derived GBM cells (T1018) and one BC cell line

(MDA-MB-231). The impact of different isolation techniques for these cell lines have not yet

been studied, and studies of exosome presence and purity are also limited for these cell lines.

We also present a critical view of the characterization methods used to detect exosomes and

evaluate the purity of the exosomes samples.

2   Material and Methods

Unless otherwise stated, water (commonly type 1 water purified by a Direct-Q® water

purification system from Millipore (Billerica, MA, USA)) was used as solvent. For detailed

information about chemicals, solutions and experimental methods used, see **Supplementary**

Materials 1 (**SM-1**).

2.1   MDA MB-231 cell culturing

The BC cell line was purchased from American Type Culture Collection (ATCC, Sesto San

Giovanni, Milan, Italy) and is a triple-negative human metastatic breast carcinoma. The cells

were maintained in Rosewell Park Memorial Institute (RPMI) 1640 growth medium depleted

of phenol red (Sigma-Aldrich, St.Louis, MO, USA) supplemented with 10% exosome-

depleted fetal bovine serum (FBS) (System Biosciences, Palo Alto, CA, USA) and 1 %

penicillin/ streptomycin (Sigma- Aldrich). The cells were incubated in a humidifying

atmosphere at 5% CO_2 and at 37 $^\circ$C. Prior to exosome isolation, 1-2.3 million cells (in T75-

T175 culturing flasks) were incubated for 6-7 days (always using a passage lower than 12).

The incubated cell culture medium was spun down at a RCF of 906 g (30 minutes at 23 °C).

2.2   Glioblastoma cell culturing

The glioblastoma (GBM) cell line (T1018) was obtained from the Langmoen Lab at Oslo

University Hospital, Norway. The cell line is derived from primary glioblastoma multiforme

tumours. The cells were maintained in DMEM/F12 medium (Gibco, Thermo Fisher

Scientific, Waltham, MA, USA). The medium was supplied with HEPES buffer (10mM) and

penicillin/ streptomycin from Lonza (Basel, Switzerland), B27 w/o Vit A from Thermo

Fisher, epidermal growth factor (20 ng/mL) and basic fibroblast growth factor (10 ng/ mL)

from R&D Systems (Minneapolis, MN, USA) and heparin (2,5 µg/mL) obtained from LEO

Pharma AS (Ballerup, Denmark). The cells were incubated in a humidifying atmosphere at 5

% CO_2 and 37 °C. Prior to exosome isolation, the incubated cell culture media was

centrifuges twice at a RCF of 453 g and 1811 g for 5 minutes each.

2.3   Exosome isolation by ultracentrifugation

For the BC and GM cells, 9-12 mL and 60 mL cull culture media were used for

centrifugation. Cell culture media was thawed on ice and centrifuged at a RCF of 1811 g (5

minutes). The supernatant was centrifuged at a RCF of 20 000 g (20 minutes at 20 °C) with an

Allegra 25R centrifuge (with TA-14-50 rotor) from Beckman Coulter (Brea, CA, USA) and

the supernatant was transferred to polycarbonate ultracentrifugation tubes (Beckman Coulter)

and diluted with phosphate buffered saline (PBS) until the tubes were full. The tubes were

centrifuged twice at a RCF of 100 000 g (90 minutes at 4 °C) with an L-80 ultracentrifuge (45

Ti rotor) from Beckman Coulter. The supernatant was removed (leaving suspension 1 cm

above the pellet) and the pellet was suspended with PBS between the centrifugations. Upon

centrifugation, the supernatant was discarded and the pellet was suspended in either PBS (3

mL for DLS- and 50-100 µL for TEM analysis) or the preferred lysis buffer (**Section 0 and**

0).

 2.4   Exosome isolation by isolation kit

The isolation of exosomes with the use of the Total Exosome Isolation Reagent (from cell

culture media) was from Thermo Fisher (catalog nr. 4478359). The isolation was performed

according to the protocol of the supplier [53]. Starting volumes were ranging from 0.5-9 mL

cell culture media for the BC cells and 5-6 mL for the GBM cells. The samples were

centrifuged with the Allegra 25R centrifuge, and the pellet was suspended as with

ultracentrifugation (**Section 0**).

2.5   Protein extraction and measurement with UV-Vis spectrophotometry

Cell and exosome protein extracts were made by adding RIPA buffer (Thermo Fisher)

containing protease inhibitors (Protease Inhibitor Coctail Tablets, Roche, Basel, Switzerland)

and phosphatase inhibitors (PhosStop Tablets, Sigma-Aldrich). The BC exosomes were

extracted in 50 µL of the RIPA solution, and the BC cells in the 100 µL RIPA solution. The

GBM exosomes were extracted in 300 µL of the RIPA solution, and the cells in 1 mL RIPA

solution. The extracts were incubated while rotating for 30 minutes (4 ℃), and then snap

frozen (at -80 ℃). Extracted samples were thawed and centrifuged at a RCF of 20 570 g (30

minutes at 2℃) using a Heraeus Fresco 21 centrifuge or Eppendorf Centrifuge 5424R. The

pellet was discarded. The protein amount was determined using Pierce™ BCA protein Assay

Kit (Thermo Fisher), by measuring the absorbance at 562 nm. The BC protein measurements

were performed using a Wallac Victor2 1420 multilabel counter and Wallac 1420

Workstation software (version 3.00), both from Perkin Elmer (Waltham, MA, USA). The

GBM protein measurements were performed on a Nanodrop 2000 Spectrophotometer

(Thermo Fisher).

2.6   Western blotting

The samples containing 3-15 µg protein were diluted giving equal volume of water before

adding 5x loading buffer (see **SM-1** for solution preparation) to yield 1x. Prior to

electrophoresis, the samples were boiled for 3 minutes at 92 ℃. The samples with added

PageRuler™ Prestained protein ladder (Thermo Fisher) were loaded on to 4-12 % Bis-Tris

gels (Thermo Fisher) and run at 70-75 V for 1 hour. The voltage was increased to 110 V after

the first hour if the bromophenol blue lane was horizontally distributed. MOPS or MES was

used as SDS running buffers (Thermo Fisher). Proteins were transferred to a 45 µm

nitrocellulose membrane (Thermo Fisher) by semi-dry electroblotting (1 W, at 4 ℃ overnight)

with a transfer chamber from Bio-Rad (Hercules, CA, USA). The transferred proteins were

blocked with a blocking solution consisting of (5:95, *w/v*) non-fat dry milk (PanReac
AppliChem ITW reagents, Darmstadt, Germany) in tween-20/ TBS (0.05:99.95, *v/v*)
(Medicago, Uppsala, Sweden) for 1 hour on a mixing plate. The proteins were then stained
with primary antibodies (at 4 °C overnight) in blocking solution and with secondary
antibodies for 2 hours at room temperature with rocking in blocking solution. The antibodies
CD9 (10626D), CD63 (10628D) and CD81 (MA5-13548 and 10630D) were purchased from
Thermo Fisher. The antibody TSG101 (T5701) was purchased from Sigma-Aldrich, and anti-
flotillin-1 (610821) and anti-calnexin (610523) antibodies were purchased from BD
Biosciences (San Jose, CA, USA). The secondary antibodies (sc-2954 and sc-2955) were
purchased from Santa Cruz (Dallas, TX, USA). Next, the membranes were further washed
with tween-20 in TBS (0.05:99.95, *v/v*) for 30 minutes on a mixing plate. Protein bands were
visualized using an ECL- prime from GE Healthcare (Buckinghamshire, UK) and
Transparency films from Nobo (integrated part of ACCO Brands Corporation, Lake Zurich,
IL, USA). The bands were developed in a ChemidocTM touch imaging system (Bio-Rad).
The antibody actin (A2066, from Sigma) was used as a positive control (results not shown).
209
2.7   Immunogold labelling and transmission electron microscopy
One drop of 5-50 μL of the exosome samples was placed on clean Parafilm, and the formvar
coated copper grid (100 square mesh) was carefully placed to float on the drop with the
coated side facing the suspension. The material was adsorbed for 5-20 minutes, before rinsing
on two large drops of PBS for 5 minutes followed by incubation on a drop (8 μL) of the
primary anti-CD9 (PA5-11559) from Thermo Fisher (diluted 1+ 9 with fish serum gelatine
(FSG) in PBS (1:99, *v/v*)) for 20 minutes. Next the grids were again washed on two large
drops of PBS for 5 minutes and incubated on a drop (5 μL) of rabbit anti-mouse antibody
(Z0259, Dako Glostrup, Denmark) (diluted 1:200 with the FSG in PBS solution) for 25
minutes, before repeating the washing on two drops of PBS for 5 minutes. Prior to gold
labelling, the grid was incubated on one drop of the diluted protein A-gold solution (1:50 in
the FSG in PBS solution) (10 nm particle size) for 20 minutes, from Cell Microscopy Core
(CMC, University Medical Center Utrecht, Utrecht, The Netherlands). A final wash on 5
drops of PBS (2 minutes) followed by 5 drops of water (3 minutes) was performed before
negative staining with uranyl acetate/water (4:96, *w/v*) for 2 minutes. Excess fluid was
removed, and the grids were stored in a storage box at room temperature (RT) until use. The
samples were visualized with a JEM-1400Plus transmission electron microscope from JEOL
(Tokyo, Japan) and images were recorded at 80 kV.

2.8 Dynamic light scattering

The dynamic light scattering (DLS) experiments were conducted with the aid of an
ALV/CGS-8F multi-detector version compact goniometer system, with 8 fiber-optical
detection units, from ALV-GmbH., Langen, Germany. The beam from a Uniphase cylindrical
22 mW HeNe-laser, operating at a wavelength of 632.8 nm with vertically polarized light,
was focused on the sample cell (10-mm NMR tubes, Wilmad Glass Co., of highest quality)
through a temperature-controlled cylindrical quartz container (with 2 plane-parallel windows),
vat (the temperature constancy being controlled to within $\pm$ 0.01 ^{o}C with a heating/cooling
circulator), which is filled with a refractive index matching liquid (*cis*-decalin). The polymer
solutions were filtered in an atmosphere of filtered air through a 5 µm filter (Millipore)
directly into precleaned NMR tubes. The measurements were carried out at 25 ^{o}C. The
measurements revealed two relaxation modes, one fast and one slow mode. This suggests that
there is a coexistence between single entities and aggregates in the solution. The analyses of
the correlation function data are presented in **SM-1** (**SM-1.6 DLS**).

2.9 Protein digestion

2.9.1 In-gel digestion

Prior to in-gel digestion, the samples were prepared and run using the same procedures as
with gel electrophoresis (**Section 0**). Bovine serum albumin (BSA) was prepared in parallel.
The gel was covered by a fixation buffer (water/ methanol/ acetic acid, 40:50:10, *v/v/v*)
overnight (18 hours at 4°C), stained with Coomassie brilliant blue for 4 hours at RT and
destained overnight with water. The gel was cut to yield four fractions from each gel lane;
approximately 0-25 kDa, 25-70 kDa, 70-130 kDa (70-250 for BC samples) and 130-up kDa
(250-up kDa for BC samples). The lane containing BSA was cut in the mass range of 55-70
kDa. Each fraction was transferred to Protein LoBind tubes. The samples were further
reduced, alkylated and digested using the protocol of Shevchenko et al. [54], with trypsin
from Promega Biotech AB (Nacka, Sweden). The digested samples were concentrated to
dryness and kept at -20 °C until further use (not more than 7 days).

2.9.2 In-solution digestion with peptide desalting

The samples were concentrated, and the pellet was dissolved with 25 µL 6 M urea in 100 mM
ammonium bicarbonate (ABC). Then, the samples were reduced with 9.5 mM dithiothreitol

(DTT, 30 minutes at 30 °C) and alkylated with 25 mM 2-iodoacetamide (IAM, 60 minutes, at

RT and in the dark). The reduction with DTT was repeated for the BC samples, with 28 mM

DTT (30 minutes at 30 °C). The proteins in the BC exosomes were initially digested by

adding 0.1 µg Lys-C (120 minutes at 37 °C), before the samples were diluted to a final

concentration of 19 mM ABC. The trypsin digestion of the samples was performed with 1 µg

trypsin (16 hours at 37 °C) and the protease activity was terminated with formic acid,

FA/water (0.9:99.1, *v/v*). The desalting and concentrating of the samples were performed

using ZipTip® (silica particles with C_{18} resin) from Millipore. The ZipTip was wetted with

neat acetonitrile (ACN) and equilibrated with trifluoroacetic acid, TFA/water (0.1:99.9, *v/v*).

The peptide samples were then pipetted through the ZipTip. Washing the ZipTip was

performed with water/ methanol/ TFA (94.9:5:0.1, *v/v/v*) and the peptides were eluted with 5

µL water/ACN/TFA (29.9:70:0.1, *v/v/v*). The filtrate was concentrated to dryness at 30°C and

frozen at -20°C until further use (not more than 30 days).

2.10 LC-MS/MS analysis

Unless otherwise stated, the dilutions during sample preparations were performed using water

(HiPerSolv Chromanorm®) from VWR. A Q-Exactive™ equipped with a nanoFlex

nanospray ion source from Thermo Fisher was used for the LC-MS/MS analysis at the two

laboratories.

2.10.1 LC-MS/MS analysis at laboratory 1

The precolumn (50 µm ID x 20-50 mm) and analytical column (50 µm ID x 150 mm) were

packed with C_{18}- Accucore particles (2.6 µm beads, 80 Å pore size) from Thermo Fisher,

using the developed method as described in our previous study [55]. An EASY-nLC 1000

pump (with autosampler) from Thermo Fisher was applied throughout the experiments. The

solvents used for peptide separation was of FA/ water (0.1:99.9, *v/v*) (mobile phase A, MP A)

and of ACN/ FA (0.1:99.9, *v/v*) (mobile phase B, MP B). Prior to LC-MS analyses, the

samples were thawed and dissolved in 10-15 µL FA/ water (0.1:99.9, *v/v*). 10 µL of the

sample was injected with a flowrate of 20 µL/min. Trapping of the analytes in the pre-column

was performed with 100 % MP A at a maximum flow rate restricted not to reach above 500

bar (12 µL). A 120- minute linear gradient elution from 3-15 % MP B with a flow rate of 130

nL/min was set for the exosome samples, starting at 3 % MP B for 3 minutes before the

percentage of MP B was increased to 15 % in 120 minutes. The MP B was increased to 50 %

for 5 minutes before the percentage of MP B increased to 80 % for 2 minutes (flow rate was

also increased to 173 nL/ min) and kept at 80 % MP B for another 15 minutes. The eluting

peptides were ionized at 1.8 kV at 250 °C for the BC exosome samples and at 275 °C for the

GBM exosome samples. The MS was operated in data-dependent (dd) positive mode to

automatically switch between MS and MS/MS acquisition. Survey full scan MS spectra (with

a mass filter of *m/z* 350 to 1850) were acquired with a resolution of 70 000, automatic gain

control (AGC) of $1x10^6$ and a maximum injection time of 120 ms. When dd/MS/MS, the

resolving power was set to 17 500, the AGC to $1 x10^5$ and the maximum injection time to 60

ms. Charges of 1, 7 or $\geq$ 8 were excluded and dynamic exclusion was set to 70.0 seconds. The

method allowed sequential isolation of up to the ten most intense ions depending on signal

intensity (intensity threshold 2.0 $x10^4$), with isolation window of *m/z* 1.8.

2.10.2 LC-MS/MS analysis at laboratory 2

The analytical column applied was an Acclaim PepMap 100 column (C18, 3 µm beads, 100

Å, 75 µm ID) of 500 mm bed length and an Ultimate 3000 nano ultra-HPLC system from

Dionex (Sunnyvale, CA, USA) was connected to the LC-MS system. The MP A was set to

FA/water (0.1:99.9, *v/v*) and the MP B was set to be ACN/ FA/ water (90:0.1:9.9, *v/v/v*). A

207- minute linear gradient elution from 4-35 % MP B with a flow rate of 300 nL/ min was

set for the exosome samples. The percentage of MP B was increased to 50 % in 20 minutes

and 80 % MP B in 2 minutes. The MS was operated in data-dependent (dd) positive mode.

Survey full scan MS spectra (with a mass filter of *m/z* 400 to 1700) were acquired with a

resolution of 70 000, automatic gain control (AGC) of $3x10^6$ and maximum injection time of

100 milliseconds. When in dd/MS/MS the resolving power was set to 35 000 and the

maximum injection time to 120 milliseconds. The dynamic exclusion was set to 60.0 seconds.

The method allowed sequential isolation of up to the ten most intense ions depending on

signal intensity (intensity threshold 1.7x10^4) isolation window was *m/z* 2 without offset.

2.10.3 Data processing and protein identification

Proteome Discoverer (version 1.4.0.228) was used to identify the peptides and proteins. The

proteins were identified using both SEQUEST and MASCOT algorithms. All searches were

performed setting the digestion enzyme to trypsin with maximum one missed cleavage,

fragment ion mass tolerance of 0.10 Da and a precursor mass tolerance of 10.0 ppm. Signal to

noise (S/N) threshold was set to 1.5 and the minimum ion count to 1. Carbamidomethylation

of cysteine was specified as static modification. Oxidation of methionine, acetylation of the

N-terminus and deamidation of glutamine and asparagine were specified as dynamic

modifications. The proteins were identified with high peptide confidence filter. Proteins
identified as keratin or trypsin were removed from the list, in addition to proteins found in the
blank samples.

3   Results and Discussion

For comparison of the two exosome isolation methods (UC and Total Exosome Isolation kit)
using the cell lines T1018 (GBM) and MDA-MB-231 (BC), common characterization
methods were investigated and evaluated for their ability to prove the presence of exosomes
and/or determine the purity. The standard techniques TEM for morphological analysis and
WB for exosome marker analysis were performed in addition to total protein determination,
DLS and LC-MS/MS. Comprehensive proteome analysis using LC-MS/MS was applied to
complement WB for detecting exosome protein markers (hereby referred to as exosome
markers) and inclusion of additional positive and negative markers for a more complete
investigation.

3.1   Similar protein yields in kit and UC samples

The protein amount in the BC (**Figure 1A**) and GBM (**Figure 1B**) exosomes per million cells
was measured using UV-Vis spectrophotometry (i.e. absorbance at 562 nm). The total protein
amount measured for exosomes isolated by the kit was 15-28 times higher than for exosomes
isolated by UC. Elevated protein amounts in exosomes isolated by kit compared to UC was
also observed in a previous study by Van Deun et al. (UC compared to an identical isolation
kit for MCF7 derived exosomes) [56]. However, increased absorbance (indicating possible
protein contaminations) was measured in the kit blanks (i.e. isolation of cell culture medium
grown w/o cells with isolation kit), in comparison to UC blank (i.e. cell culture medium
grown w/o cells isolated by UC) where the absorbance was below the limit of quantification.
When correcting for the blank (subtracting the protein amount measured in blank samples
from the protein amount in exosome samples), the measured protein amount for exosomes
isolated by the kit and UC was similar.

Figure 1: Measured protein amount pr. million cells in exosome samples from GBM- and BC cells isolated by kit and UC (n ≥ 2). A) Measured absolute protein amount (µg) for the BC exosome sample. **B**) The measured relative protein amount (due to low regression coefficients, %) for the GBM exosome samples. Each replicate is depicted as circles, and the median depicted as a line. The X-mark shows the measured protein amount in the blank sample (isolated cell culture medium). The protein amounts were measured by UV-Vis spectroscopy (absorption at λ= 562 nm) after reaction with BCA kit reagents.

3.2   TEM and DLS detected vesicles in the expected size range for exosomes

Morphological analysis of the exosome samples was performed using TEM and immunogold

labelling of CD9. In addition, the hydrodynamic particle size distribution was measured using

DLS analysis. Clusters of vesicles in the samples isolated with both kit and UC were observed

in the micrographs (**Figure 2**). Vesicle structures similar to that described in literature were

observed [6, 57, 58]. The DLS experiments disclosed the coexistence of two populations of

moieties, single entities and clusters, both with a narrow size distribution.

3.2.1   GBM exosomes

No CD9-labelling was observed for the vesicle structures observed in the GBM exosomes

(**Figure 2A-I** and **2A-III**) and could not confirm the presence of a membrane enclosing the

vesicles. The UC sample presented more distinct double membranes in the expected size

range for exosomes compared to the kit sample. The blank samples for both isolation methods

did not display membrane structures and seemed pure (**Figure 2A-II** and **2A-IV**). The high

378   purity was further confirmed by DLS analysis of the UC blank (**Figure 2B**). The DLS-
379   analysis of the exosome samples exhibited particles of similar sizes of 51 and 73 nm (mean)
380   with both isolation methods (**Figure 2B**). Thus, both isolation methods isolated comparable
381   exosome populations.
382
3.2.2   BC exosomes
Several of the BC vesicle structures were CD9-labelled (**Figure 2B-I** and **B-III**). CD9-
labelled vesicles have also been observed in a previous study of the same cell line [59].
Notably, the blank samples displayed contamination (**Figure 2C-II and C-IV**), e.g. exosome-
resembling vesicles were found in the UC blank (red dashed circles). However, no
contaminations were found in the UC blank using DLS, while the kit blank displayed 67 nm
(mean) contaminations (**Figure 2D**). The DLS analysis also presented two distinct particle
diameters in exosome samples from kit (28 and 95 nm, mean values) while only one was
presented using UC (137 nm, mean value), indicating differences in the particle sizes isolated
with the two isolation methods.
393
The sizes observed with DLS correlates well with other studies (30– 250 nm), also displaying
a heterogeneity of particle sizes in exosome samples [13, 56, 60-64]. In conclusion, the
isolated samples did show structures resembling to those of EVs, but some blank samples
were not entirely pure from vesicles or particles. Observations made with TEM are not
necessarily detectable with DLS because TEM measure on dry samples, whereas DLS probes
on solution or suspension of particles. In addition, the micrographs taken with TEM display a
narrow section of the grid, which again represent a small part of the isolated sample.
401

Figure 2: Transmission electron micrographs and hydrodynamic particle size (nm) distribution by DLS analysis of exosome samples from GBM- and BC cells isolated by kit and UC. The TEM-samples were immunogold labelled with anti-CD9, with gold particles depicted as 10 nm black dots. Images were taken with a magnification of 400 000, and the dashed areas were additionally zoomed. **A**) Micrographs of CD9-labelled GBM exosome samples. **I** depict the micrograph from a kit sample, **II** the kit blank, **III** the UC sample, and **IV** the UC blank. **B**) DLS analysis of GBM exosome samples isolated by kit and UC (n = 1). No particles were detected in the blank sample isolated with UC (n = 1). DLS analysis of the kit blank was not performed. **C**) Micrographs of CD9-labelled BC exosome samples. **I** depicts the micrograph from the kit sample, **II** the kit blank, **III** a UC sample, and **IV** the UC blank. **D**) DLS analysis of BC exosome samples isolated by kit (n = 2) and UC (n = 3), including the kit blank (n = 1). No particles were detected in the blank sample isolated with UC.

3.3 Western blot analyses indicated the presence of exosomes, but detected impurities in the GBM exosome samples

According to ISEV, at least three markers should be included for characterization of exosomes, originating from both transmembrane proteins (e.g. tetraspanins), cytosolic proteins (e.g. TSG101 or annexins) and negative markers (e.g. calnexin) [45]. In the present study, WB was performed using antibodies for a selection of positive exosome markers (the tetraspanins CD81, CD9 and CD63, TSG101 and flotillin-1). Calnexin was selected as a negative marker for purity evaluation as recommended by ISEV. This protein is located at the endoplasmic reticulum (ER) and have been absent in exosome samples in some studies [45, 60]. Hence, the presence of calnexin is assumed to signalize ER-contamination (which enables possibilities of contaminations from other cellular organelles).

3.3.1 GBM exosomes

For the GBM cells and exosomes, all exosome markers (both positive and negative) were found in samples from both kit- and UC isolation except for CD81, which was only found in UC samples (**Figure 3**). The WB-bands were more apparent for most positive markers for exosomes isolated by UC (even with higher protein amount loaded for the kit samples), which is also in accordance with Van Deun's study [56].

3.3.2 BC exosomes

For the BC exosomes, inconsistency for several positive exosome markers were observed between the kit and UC samples (**Figure 3**). The positive marker CD63 was only detected in kit samples, while CD81 was only detected in samples from UC (similar to GBM exosomes). The proteins TSG101, flotillin-1 and CD9 (barely visible in the UC samples) were detected for both isolation methods. Although, a study by Harris et al. did not detect TSG101 using WB on BC exosomes isolated by UC [13]. The proteins CD9, CD81 and flotillin-1 were detected in other WB-studies on UC samples from the same cell line [23, 65, 66].

The reason for the absence of some tetraspanins in kit and others in UC samples could be due
to protein concentrations below detection limits or poor antibody quality. Several antibodies
for CD63 and CD81 (from different companies) were tested on the BC exosomes before a
signal was obtained, which could indicate poor antibody quality. On the other hand, the WB
was performed under reducing conditions. When the epitope binds to cysteine-conserved
protein domains (i.e. tetraspanins), performing WB under non-reducing conditions is more
commonly selected. The stronger signals for the kit samples from BC could be due to the
higher loaded protein amount. Nevertheless, the presence of positive markers indicates the
presence of exosomes in the isolates obtained using both methods. The absence of calnexin in
BC exosomes from both isolation methods indicate that the samples do not contain
contamination from the ER. However, only one negative marker is insufficient to exclude cell

Figure 3: Western blot of common protein organelle impurities. Further
450 **exosome markers.** The protein markers CD81, investigation by implementing more
451 CD9, CD63, TSG101, flotillin-1 (positive markers, comprehensive methods like LC-
+) and calnexin (negative marker, -) were targeted 452 MS/MS was considered to be
in cell lysates and exosome samples isolated with beneficial.
kit and by UC (n ≥ 2). For the BC exosomes, 15 μg 453
protein was loaded for kit samples and 3 μg for UC 454
samples. For the GBM exosomes, ~14 μg was
loaded for kit samples and ~8 μg for UC samples. 455 **3.4   LC-MS/MS studies**
Uncropped Western blots are presented in 456 **confirmed and complemented the**
Supplementary Materials 2 (SM-2). 457 **WB study**

LC-MS/MS was performed to confirm the WB observations, using in-house packed nano
separation columns [55]. One positive marker (annexin A2) and one negative marker
(serine/threonine-protein kinase 26) was added to complement the WB study. The detection
of the selected exosome markers using LC-MS/MS is presented in **Table 1**, with
chromatograms and MS/MS spectra of a CD9 signature peptide (BC, **Figure 4A**) and
calnexin signature peptide (GBM, **Figure 4B**). Similar to the WB analysis, the same
tetraspanins including calnexin observed in kit and UC samples were detected for the GBM
exosomes using LC-MS/MS. However, the tetraspanins detected from kit samples were only
found in one replicate, indicating low concentrations and high detection uncertainty. Flotillin-
1 was also detected, but only for UC samples with LC-MS/MS. TSG101 was identified in
neither kit nor UC (GBM exosomes) using LC-MS/MS in contradiction to WB.

Table 1: A selection of common protein markers (from LC-MS analyses) with a specific
exosome related function. The exosome markers were identified in exosome samples from

 MDA-MB-231 and glioblastoma isolated with kit and by UC. In addition, proteins identified in blank samples are shown.

Protein marker	Exosome related function	GBM				BC			
		UC	UC blank	Kit	Kit blank	UC	UC blank	Kit	Kit blank
CD81	Membrane protein	x	-	-	-	x	-	-	-
CD9	Membrane protein	x	-	x	-	x	x	x	-
CD63	Membrane protein	x	-	x	-	x	-	-	-
TSG101	Exosome biogenesis	-	-	-	-	x	-	-	-
Annexin A2	Exosome biogenesis	x	-	x	-	x	-	x	-
Flotillin-1	Exosome biogenesis	-	-	-	-	x	-	-	-
Calnexin	Negative	x	-	x	-	-	-	-	-
Serine/ threonine-protein kinase 26	Negative	-	-	-	-	-	-	-	-
Actin	General marker	x	x	x	x	x	-	x	x

For the BC exosomes, the LC-MS/MS analysis is contradictory to the trend observed in the
WB analysis where kit samples provided more pronounced signals for positive markers than
the UC samples. Using LC-MS/MS, several positive markers were absent in the kit samples
when injecting similar amounts of protein as with UC samples. The reason for the difference
to WB could hence be partially due to higher protein amount loaded onto the gel for kit
samples. Calnexin was, similar to the WB, not detected in the BC samples. Annexin A2
(positive marker) was found in all samples and serine/threonine-protein kinase 26 (second
negative marker) was not detected in any samples.

Other negative markers from peroxisomes (PMP70), mitochondria (prohibitin-1, hexokinase-
2 and mitochondrial phosphoenolpyruvate carboxykinase [GTP]), Golgi apparatus (GM130,
translocation protein SEC62, translocation protein SEC63 and protein disulfide-isomerase
TMX3), nucleus (Bcl-2-associated transcription factor 1 and c) and ER (calreticulin), as used
in other studies [56, 67], were not detected in any BC samples in our study (only translocation

protein SEC63 and protein disulfide-isomerase TMX3 detected in GBM samples). The
absence of several negative markers in the BC exosomes supports the low cell organelle
contamination observed with WB and LC-MS/MS.

Figure 4: Chromatograms and MS/MS spectrums from LC-MS/MS analysis of GBM-
and BC exosome peptides. A) Chromatogram with corresponding MS/MS spectrum for the
CD9 signature peptide KDVLETFTVK (*m/z*=393.89, z=3) in BC exosomes isolated by
UC. **C)** Chromatogram with corresponding MS/MS spectrum for the calnexin signature
peptide AEEDEILNR (*m/z*=544.77, z=2) from GBM exosomes isolated by UC.

3.5   GO annotations reveal proteins annotated to cell organelles in BC exosomes

However, the possibility of contaminants being present in BC samples cannot be excluded.

Other general proteins related to e.g. the nucleus, Golgi apparatus, mitochondrion and ER was

indeed identified in the BC exosomes using LC-MS/MS (**Figure 5**). The proteins identified in

BC exosomes were classified based on their gene ontology (GO) annotations to different

cellular localizations, where one protein can be annotated to several cellular localizations. Out

of the 668 proteins identified from the UC and 814 from the kit, 615 and 749 DAVID ID`s

were annotated to selected cellular localizations. From them, both isolation methods

generated samples enriched in exosome related proteins. Several proteins were annotated to

the cellular organelles mitochondria (11-13 %), ER (6-8 %) and Golgi apparatus (7 %).

Interestingly, 35-42 % of the proteins were also annotated to the nucleus (e.g. histones),

which seems to imply impure exosome samples [45]. On the other hand, a high percentage

(20-40 %) of other proteins related to the nucleus has also been found in exosome samples

from other studies [68, 69]. The presence of cell organelle annotated proteins could point

toward cellular impurities in the samples, which would not have been discovered by targeted

protein characterization methods (e.g. WB). However, there is not sufficient knowledge on

whether cell organelle proteins derive exclusively from cell impurities, or if they occur
naturally in EVs. To summarize, from our point of view, complete information about
exosome purity cannot be obtained by any of the common characterization techniques used
today.

Figure 5*: **GO annotation of proteins in BC exosomes to different cellular locations.** The
identified proteins classified by their cellular location (GO annotations) grouped based on
their positive/ negative relevance towards exosomes. The annotated proteins (% of total
proteins) and their cellular location, with proteins annotated from the kit samples are shown in
red (from 749 DAVID ID's), while proteins annotated from the UC samples are shown in
blue (from 615 DAVID ID's).

3.6   The isolation method and cell line influence the number of cancer related proteins
identified

The total number of proteins identified in the GBM and BC exosome samples using LC-
MS/MS (performed in two different laboratories) is presented in the Venn diagrams in **Figure**
6 (see **SM-3** for a list of all identified proteins). For the GBM exosomes, the number of
identified proteins reflects the findings in both WB and LC-MS/MS exosome marker
investigations. UC-isolated exosomes provided more unique proteins than the kit-isolated
exosomes (75 % higher number of identified proteins than the kit). An increased number of
potential biomarkers for GBM (e.g. heat shock proteins 70 kDa and 90 kDa [70-72],
chondroitin sulfate proteoglycan 4 [70, 73], CD44 [70, 73, 74] and CD276 [75]) were also
identified in the UC samples compared to the kit samples using LC-MS/MS. The

identification of relevant biomarkers is of great interest for further studies on exosomes.
However, the identified biomarkers cannot exclusively be related to exosomes due to the
presence of negative exosomes markers indicating cellular contaminations.

For the BC exosomes, the tables are turned, and the kit-isolated exosomes provided 12 %
higher number of identified proteins than UC-isolated exosomes. However, there was no
correlation between the injected protein amount or the starting volume used for isolation, and
the number of identified proteins in kit- vs UC samples for BC exosomes (result not shown).
Thus, the reason for why the protein number is changed between the two cell lines and
isolation methods is unknown. The number of identified biomarkers related to the triple
negative breast cancer (e.g. histone H4 [76], heat shock 90 kDa α and β protein [77],
calmodulin and epithermal growth factor receptor [78]) in the UC samples compared to kit
samples were comparable.

When comparing cell lines, the number of identified proteins was lower in GBM exosomes
than BC exosomes, but the number for GBM exosomes is comparable to another LC-MS/MS
study on GBM exosomes [79].

Figure 6: Venn Diagram presenting the number of proteins identified by LC-MS/MS in
GBM- and BC exosome samples isolated with kit and by UC. The numbers are the total
number of unique proteins identified when trypsin, keratin related proteins and the proteins
identified in blank samples were disregarded. One signature peptide was selected as
requirement for positive identifications during database search. Equal amounts of protein
were injected for both kit and UC (~ 1.5 µg protein for GBM exosomes (n= 6) and ~2- 5 µg
protein for BC exosomes (n= 3)). A list of all proteins identified is presented in
Supplementary Materials 3 (SM-3).

3.7 Choosing the proper exosome isolation method is not straight forward

A complete comparison of the characteristics of exosome isolation methods is given in **Table 2**. For all exosome samples, the kit and UC samples display similarities and differences.

3.7.1 GBM exosomes

For the GBM exosomes, one of the positive markers detected in the UC samples (CD81) was not found in kit samples by WB. In TEM, double membrane structures were more eminent in the UC samples, but the existence of double membranes cannot be excluded by looking at the micrographs from the kit samples. The largest differences between kit and UC for the GBM exosomes were found in the LC-MS/MS studies recoding positive markers and general protein number. All tetraspanins investigated were detected in the UC samples in several replicates. For the kit, CD81 was not found, and the detected tetraspanins (CD63 and CD9) were only found in one replicate each. A larger number of proteins and biomarker candidates were also identified in the UC samples compared to kit samples. However, the negative marker calnexin was detected in more replicates for the UC than the kit using LC-MS/MS. In total, from **Table 2**, UC appear to be the method of choice for isolation of GBM (T1018) exosomes.

3.7.2 BC exosomes

For the BC exosomes, there was a slight difference in favor of the kit method regarding positive markers in WB and the number of identified proteins (LC-MS/MS). However, using LC-MS/MS, more relevant protein markers were found in the UC samples in contrary to the WB study. In addition, TEM presented slightly more defined double membrane structures with more labeling using UC. However, the micrograph displays an extremely small part of the whole sample. The isolation methods also performed similarly regarding biomarker identifications. Based on the above, there is no obvious reason for choosing one method over the other. However, tendencies towards different characteristics of the isolated exosomes from kit and UC have been observed.

Table 2: Comparison of characteristics of exosome isolated from GBM and BC cell culture medium. In the table, UC and kit was compared for their ability to isolate pure exosomes based on the characterization techniques used. Increasing number of + signs indicate positive relation regarding indications of exosomes and/or high purity of exosomes.

Characterization method	GBM		BC	
	UC	Kit	UC	Kit

Protein amount measured (corrected for blank)	+	++	+	++
TEM	++	+	+++	++
DLS	+++	+++	+	++
WB (positive markers)	++	+	+	++
WB (negative marker)	+	+	+++	+++
LC-MS/MS (positive markers)	++	+	+++	+
LC-MS/MS (negative markers)	+	++	+++	+++
LC-MS/MS number of proteins identified	+++	+	+	++
LC-MS/MS relevant biomarkers	++	+	++	++

4   Conclusions

The observations in our study support the perception of exosome isolation being highly dependent on the isolation protocol used, general differences in the behavior of cells/exosomes between cell lines and the characterization methods and conditions applied. The knowledge about potentially natural occurring proteins in exosomes is insufficient and a selection of expected negative markers cannot exclusively be used to determine the purity of exosome samples. Hence, the application area (e.g. proving presence of exosomes, determine exosome purity, biomarker discovery, or lower the dynamic range of proteins in body fluids like blood) and sample volume available for the exosome isolation should be stronger determent factors when selecting the proper isolation method. Thus, the characterization methods used in this study are not able to distinguish exosomes from cellular contaminations and other vesicles. Untargeted proteome analyses using LC-MS/MS did provide more extensive and versatile information on the protein content of the samples than most commonly used targeted WB of few proteins. Consequently, we suggest that LC-MS/MS should be implemented to a higher extend regarding exosome characterization. Considering our findings, it is important to state that the term "exosome enrichment" is more appropriate than "exosome isolation".

5   References

[1] C. Théry, L. Zitvogel, S. Amigorena, Exosomes: composition, biogenesis and function, Nature Reviews Immunology 2 (2002) 569.

[2] T. Lener, M. Gimona, L. Aigner, V. Borger, E. Buzas, G. Camussi, N. Chaput, D. Chatterjee, F.A. Court, H.A. del Portillo, L. O'Driscoll, S. Fais, J.M. Falcon-Perez, U. Felderhoff-Mueser, L. Fraile, Y.S. Gho, A. Gorgens, R.C. Gupta, A. Hendrix, D.M. Hermann, A.F. Hill, F. Hochberg, P.A. Horn, D. de Kleijn, L. Kordelas, B.W. Kramer, E.M. Kramer-Albers, S. Laner-Plamberger, S. Laitinen, T. Leonardi, M.J. Lorenowicz, S.K. Lim, J. Lotvall, C.A. Maguire, A. Marcilla, I.

Nazarenko, T. Ochiya, T. Patel, S. Pedersen, G. Pocsfalvi, S. Pluchino, P. Quesenberry, I.G.
Reischl, F.J. Rivera, R. Sanzenbacher, K. Schallmoser, I. Slaper-Cortenbach, D. Strunk, T. Tonn,
P. Vader, B.W.M. van Balkom, M. Wauben, S. El Andaloussi, C. Thery, E. Rohde, B. Giebel,
Applying extracellular vesicles based therapeutics in clinical trials - an ISEV position paper, J.
Extracell. Vesicles 4 (2015) 31.

[3] H. Zhao, L. Yang, J. Baddour, A. Achreja, V. Bernard, T. Moss, J.C. Marini, T. Tudawe, E.G.
Seviour, F.A. San Lucas, H. Alvarez, S. Gupta, S.N. Maiti, L. Cooper, D. Peehl, P.T. Ram, A.
Maitra, D. Nagrath, Tumor microenvironment derived exosomes pleiotropically modulate
cancer cell metabolism, eLife 5 (2016) e10250.

[4] T. Skotland, K. Sandvig, A. Llorente, Lipids in exosomes: Current knowledge and the way
forward, Progress in Lipid Research 66 (2017) 30-41.

[5] A. Beach, H.-G. Zhang, M.Z. Ratajczak, S.S. Kakar, Exosomes: an overview of biogenesis,
composition and role in ovarian cancer, Journal of Ovarian Research 7 (2014) 14-14.

[6] P. Li, M. Kaslan, S.H. Lee, J. Yao, Z. Gao, Progress in Exosome Isolation Techniques,
Theranostics 7(3) (2017) 789-804.

[7] A. Becker, B.K. Thakur, J.M. Weiss, H.S. Kim, H. Peinado, D. Lyden, Extracellular Vesicles in
Cancer: Cell-to-Cell Mediators of Metastasis, Cancer Cell 30(6) (2016) 836-848.

[8] G. Pocsfalvi, C. Stanly, I. Fiume, K. Vekey, Chromatography and its hyphenation to mass
spectrometry for extracellular vesicle analysis, J. Chromatogr. A 1439 (2016) 26-41.

[9] M. Farahani, C. Rubbi, L. Liu, J.R. Slupsky, N. Kalakonda, CLL exosomes modulate the
transcriptome and behaviour of recipient stromal cells and are selectively enriched in miR-
202-3p, PloS one 10(10) (2015) e0141429.

[10] M. Logozzi, A. De Milito, L. Lugini, M. Borghi, L. Calabro, M. Spada, M. Perdicchio, M.L.
Marino, C. Federici, E. Iessi, D. Brambilla, G. Venturi, F. Lozupone, M. Santinami, V. Huber, M.
Maio, L. Rivoltini, S. Fais, High Levels of Exosomes Expressing CD63 and Caveolin-1 in Plasma
of Melanoma Patients, Plos One 4(4) (2009) 10.

[11] I. Parolini, C. Federici, C. Raggi, L. Lugini, S. Palleschi, A. De Milito, C. Coscia, E. Iessi, M.
Logozzi, A. Molinari, M. Colone, M. Tatti, M. Sargiacomo, S. Fais, Microenvironmental pH Is a
Key Factor for Exosome Traffic in Tumor Cells, J. Biol. Chem. 284(49) (2009) 34211-34222.

[12] H. Peinado, M. Alečković, S. Lavotshkin, I. Matei, B. Costa-Silva, G. Moreno-Bueno, M.
Hergueta-Redondo, C. Williams, G. García-Santos, C.M. Ghajar, Melanoma exosomes
educate bone marrow progenitor cells toward a pro-metastatic phenotype through MET,
Nature medicine 18(6) (2012) 883-891.

[13] D.A. Harris, S.H. Patel, M. Gucek, A. Hendrix, W. Westbroek, J.W. Taraska, Exosomes
released from breast cancer carcinomas stimulate cell movement, PloS one 10(3) (2015)
e0117495.

[14] C.E. DeSantis, C.C. Lin, A.B. Mariotto, R.L. Siegel, K.D. Stein, J.L. Kramer, R. Alteri, A.S.
Robbins, A. Jemal, Cancer treatment and survivorship statistics, 2014, CA: A Cancer Journal
for Clinicians 64(4) (2014) 252-271.

[15] A.J. Redig, S.S. McAllister, Breast cancer as a systemic disease: a view of metastasis,
Journal of Internal Medicine 274(2) (2013) 113-126.

[16] M. Touat, A. Duran-Pena, A. Alentorn, L. Lacroix, C. Massard, A. Idbaih, Emerging
circulating biomarkers in glioblastoma: promises and challenges, Expert Rev. Mol. Diagn.
15(10) (2015) 1311-1323.

[17] J.R. Molina, Y. Hayashi, C. Stephens, M.M. Georgescu, Invasive Glioblastoma Cells
Acquire Stemness and Increased Akt Activation, Neoplasia 12(6) (2010) 453-U37.

[18] M. Preusser, S. de Ribaupierre, A. Wohrer, S.C. Erridge, M. Hegi, M. Weller, R. Stupp,
Current Concepts and Management of Glioblastoma, Ann. Neurol. 70(1) (2011) 9-21.

[19] Z.N. Jan Chrastina, Milan Brazdil, Marketa Hermanova, Glioblastoma multiforme in a
patient with isolated hemimegalencephaly, Journal of Neurological Surgery Reports 76(1)
(2015) 160-163.

[20] H.L. Shao, J. Chung, K. Lee, L. Balaj, C. Min, B.S. Carter, F.H. Hochberg, X.O. Breakefield,
H. Lee, R. Weissleder, Chip-based analysis of exosomal mRNA mediating drug resistance in
glioblastoma, Nat. Commun. 6 (2015) 9.

[21] M.G. Best, N. Sol, S. Zijl, J.C. Reijneveld, P. Wesseling, T. Wurdinger, Liquid biopsies in
patients with diffuse glioma, Acta Neuropathol. 129(6) (2015) 849-865.

[22] M. Cristofanilli, G.T. Budd, M.J. Ellis, A. Stopeck, J. Matera, M.C. Miller, J.M. Reuben,
G.V. Doyle, W.J. Allard, L.W. Terstappen, Circulating tumor cells, disease progression, and
survival in metastatic breast cancer, New England Journal of Medicine 351(8) (2004) 781-
791.

[23] S.A. Melo, L.B. Luecke, C. Kahlert, A.F. Fernandez, S.T. Gammon, J. Kaye, V.S. LeBleu, E.A.
Mittendorf, J. Weitz, N. Rahbari, Glypican-1 identifies cancer exosomes and detects early
pancreatic cancer., Nature 523(7559) (2015) 177-182.

[24] D.D. Taylor, C. Gercel-Taylor, MicroRNA signatures of tumor-derived exosomes as
diagnostic biomarkers of ovarian cancer, Gynecologic Oncology 110(1) (2008) 13-21.

[25] H. Roberg-Larsen, K. Lund, K.E. Seterdal, S. Solheim, T. Vehus, N. Solberg, S. Krauss, E.
Lundanes, S.R. Wilson, Mass spectrometric detection of 27-hydroxycholesterol in breast
cancer exosomes, The Journal of steroid biochemistry and molecular biology 169 (2017) 22-
28.

[26] J. Skog, T. Würdinger, S. Van Rijn, D.H. Meijer, L. Gainche, W.T. Curry, B.S. Carter, A.M.
Krichevsky, X.O. Breakefield, Glioblastoma microvesicles transport RNA and proteins that
promote tumour growth and provide diagnostic biomarkers, Nature cell biology 10(12)
(2008) 1470-1476.

[27] F.H. Hochberg, N.A. Atai, D. Gonda, M.S. Hughes, B. Mawejje, L. Balaj, R.S. Carter,
Glioma diagnostics and biomarkers: an ongoing challenge in the field of medicine and
science, Expert Rev. Mol. Diagn. 14(4) (2014) 439-452.

[28] L. Saadatpour, E. Fadaee, S. Fadaei, R.N. Mansour, M. Mohammadi, S.M. Mousavi, M.
Goodarzi, J. Verdi, H. Mirzaei, Glioblastoma: exosome and microRNA as novel diagnosis
biomarkers, Cancer Gene Ther. 23(12) (2016) 415-418.

[29] A. Mondal, D.K. Singh, S. Panda, A. Shiras, Extracellular Vesicles As Modulators of Tumor
Microenvironment and Disease Progression in Glioma, Front. Oncol. 7 (2017) 8.

[30] S. Keller, J. Ridinger, A.-K. Rupp, J.W. Janssen, P. Altevogt, Body fluid derived exosomes
as a novel template for clinical diagnostics, Journal of translational medicine 9(1) (2011) 86.

[31] C. Théry, S. Amigorena, G. Raposo, A. Clayton, Isolation and characterization of
exosomes from cell culture supernatants and biological fluids, Current Protocols in Cell
Biology (2006) 3.22. 1-3.22. 29.

[32] R. Cantin, J. Diou, D. Bélanger, A.M. Tremblay, C. Gilbert, Discrimination between
exosomes and HIV-1: purification of both vesicles from cell-free supernatants, Journal of
Immunological Methods 338(1) (2008) 21-30.

[33] D.W. Greening, R. Xu, H. Ji, B.J. Tauro, R.J. Simpson, A protocol for exosome isolation
and characterization: evaluation of ultracentrifugation, density-gradient separation, and
immunoaffinity capture methods, Proteomic Profiling: Methods and Protocols (2015) 179-
209.

[34] M.A. Rider, S.N. Hurwitz, D.G. Meckes Jr, ExtraPEG: a polyethylene glycol-based method
for enrichment of extracellular vesicles, Scientific reports 6 (2016) 23978.
[35] A.N. Böing, E. Van Der Pol, A.E. Grootemaat, F.A. Coumans, A. Sturk, R. Nieuwland,
Single-step isolation of extracellular vesicles by size-exclusion chromatography, Journal of
Extracellular Vesicles 3 (2014) DOI: 10.3402/jev.v3.23430.
[36] A. Gámez-Valero, M. Monguió-Tortajada, L. Carreras-Planella, Size-Exclusion
Chromatography-based isolation minimally alters Extracellular Vesicles' characteristics
compared to precipitating agents, Scientific Reports 6 (2016) DOI: 10.1038/srep33641.
[37] R. Grant, E. Ansa-Addo, D. Stratton, S. Antwi-Baffour, S. Jorfi, S. Kholia, L. Krige, S. Lange,
J. Inal, A filtration-based protocol to isolate human plasma membrane-derived vesicles and
exosomes from blood plasma, Journal of Immunological Methods 371(1) (2011) 143-151.
[38] A. Clayton, J. Court, H. Navabi, M. Adams, M.D. Mason, J.A. Hobot, G.R. Newman, B.
Jasani, Analysis of antigen presenting cell derived exosomes, based on immuno-magnetic
isolation and flow cytometry, Journal of Immunological Methods 247(1) (2001) 163-174.
[39] J.S. Yang, J.C. Lee, S.K. Byeon, K.H. Rha, M.H. Moon, Size dependent lipidomic analysis of
urinary exosomes from patients with prostate cancer by flow field-flow fractionation and
nanoflow liquid chromatography-tandem mass spectrometry, Analytical Chemistry 89 (2017)
2488-2496.
[40] B.J. Tauro, D.W. Greening, R.A. Mathias, S. Mathivanan, H. Ji, R.J. Simpson, Two distinct
populations of exosomes are released from LIM1863 colon carcinoma cell-derived
organoids, Molecular & Cellular Proteomics 12(3) (2013) 587-598.
[41] S.N. Bhatia, D.E. Ingber, Microfluidic organs-on-chips, Nature Biotechnology 32(8)
(2014) 760-772.
[42] A. Abramowicz, P. Widlak, M. Pietrowska, Proteomic analysis of exosomal cargo: the
challenge of high purity vesicle isolation, Mol. Biosyst. 12(5) (2016) 1407-1419.
[43] M. He, Y. Zeng, Microfluidic Exosome Analysis toward Liquid Biopsy for Cancer, Jala
21(4) (2016) 599-608.
[44] S. Kreimer, A.M. Belov, I. Ghiran, S.K. Murthy, D.A. Frank, A.R. Ivanov, Mass-
Spectrometry-Based Molecular Characterization of Extracellular Vesicles: Lipidomics and
Proteomics, Journal of Proteome Research 14(6) (2015) 2367-2384.
[45] J. Lötvall, A.F. Hill, F. Hochberg, E.I. Buzás, D. Di Vizio, C. Gardiner, Y.S. Gho, I.V.
Kurochkin, S. Mathivanan, P. Quesenberry, S. Sahoo, H. Tahara, M.H. Wauben, K.W. Witwer,
C. Théry, Minimal experimental requirements for definition of extracellular vesicles and their
functions: a position statement from the International Society for Extracellular Vesicles, J.
Extracell. Vesicles 3 (2014) 10.3402/jev.v3.26913.
[46] Z. Andreu, M. Yáñez-Mó, Tetraspanins in Extracellular Vesicle Formation and Function,
Frontiers in Immunology 5 (2014) 442.
[47] J.M. Escola, M.J. Kleijmeer, W. Stoorvogel, J.M. Griffith, O. Yoshie, H.J. Geuze, Selective
enrichment of tetraspan proteins on the internal vesicles of multivesicular endosomes and
on exosomes secreted by human B-lymphocytes, J. Biol. Chem. 273(32) (1998) 20121-20127.
[48] M.J. Kleijmeer, W. Stoorvogel, J.M. Griffith, O. Yoshie, H.J. Geuze, Selective enrichment
of tetraspan proteins on the internal vesicles of multivesicular endosomes and on exosomes
secreted by human B-lymphocytes, The Journal of Biological Chemistry 273(32) (1998)
20121-20127.
[49] N.P. Hessvik, A. Llorente, Current knowledge on exosome biogenesis and release,
Cellular and Molecular Life Sciences 75(2) (2018) 193-208.

[50] S. Mathivanan, C.J. Fahner, G.E. Reid, R.J. Simpson, ExoCarta 2012: database of
exosomal proteins, RNA and lipids, Nucleic Acids Research 40(D1) (2012) D1241-D1244.
[51] H. Kalra, R.J. Simpson, H. Ji, E. Aikawa, P. Altevogt, P. Askenase, V.C. Bond, F.E. Borras, X.
Breakefield, V. Budnik, E. Buzas, G. Camussi, A. Clayton, E. Cocucci, J.M. Falcon-Perez, S.
Gabrielsson, Y.S. Gho, D. Gupta, H.C. Harsha, A. Hendrix, A.F. Hill, J.M. Inal, G. Jenster, E.M.
Kramer-Albers, S.K. Lim, A. Llorente, J. Lotvall, A. Marcilla, L. Mincheva-Nilsson, I. Nazarenko,
R. Nieuwland, E.N.M. Hoen, A. Pandey, T. Patel, M.G. Piper, S. Pluchino, T.S.K. Prasad, L.
Rajendran, G. Raposo, M. Record, G.E. Reid, F. Sanchez-Madrid, R.M. Schiffelers, P. Siljander,
A. Stensballe, W. Stoorvogel, D. Taylor, C. Thery, H. Valadi, B.W.M. van Balkom, J. Vazquez,
M. Vidal, M.H.M. Wauben, M. Yanez-Mo, M. Zoeller, S. Mathivanan, Vesiclepedia: A
Compendium for Extracellular Vesicles with Continuous Community Annotation, PLoS. Biol.
10(12) (2012) 5.
[52] D.K. Kim, B. Kang, O.Y. Kim, D.S. Choi, J. Lee, S.R. Kim, G. Go, Y.J. Yoon, J.H. Kim, S.C.
Jang, K.S. Park, E.J. Choi, K.P. Kim, D.M. Desiderio, Y.K. Kim, J. Lotvall, D. Hwang, Y.S. Gho,
EVpedia: an integrated database of high-throughput data for systemic analyses of
extracellular vesicles, J Extracell Vesicles 2 (2013).
[53] ThermoFisher, Total Exosome Isolation Reagent (from cell culture media).
https://www.thermofisher.com/order/catalog/product/4478359?SID=srch-srp-4478359).
[54] A. Shevchenko, H. Tomas, J. Havli, J.V. Olsen, M. Mann, In-gel digestion for mass
spectrometric characterization of proteins and proteomes, Nature Protocols 1 (2007) 2856.
[55] H.S. Berg, K.E. Seterdal, T. Smetop, R. Rozenvalds, O.K. Brandtzaeg, T. Vehus, E.
Lundanes, S.R. Wilson, Self-packed core shell nano liquid chromatography columns and
silica-based monolithic trap columns for targeted proteomics, J. Chromatogr. A 1498 (2017)
111-119.
[56] J. Van Deun, P. Mestdagh, R. Sormunen, V. Cocquyt, K. Vermaelen, J. Vandesompele, M.
Bracke, O. De Wever, A. Hendrix, The impact of disparate isolation methods for extracellular
vesicles on downstream RNA profiling, J. Extracell. Vesicles 3(1) (2014) 24858.
[57] B.-T. Pan, K. Teng, C. Wu, M. Adam, R.M. Johnstone, Electron microscopic evidence for
externalization of the transferrin receptor in vesicular form in sheep reticulocytes, The
Journal of Cell Biology 101(3) (1985) 942-948.
[58] B.-T. Pan, R.M. Johnstone, Fate of the transferrin receptor during maturation of sheep
reticulocytes in vitro: selective externalization of the receptor, Cell 33(3) (1983) 967-978.
[59] S.A. Melo, H. Sugimoto, J.T. O'Connell, N. Kato, A. Villanueva, A. Vidal, L. Qiu, E. Vitkin,
L.T. Perelman, C.A. Melo, Cancer exosomes perform cell-independent microRNA biogenesis
and promote tumorigenesis, Cancer Cell 26(5) (2014) 707-721.
[60] R.J. Lobb, M. Becker, S.W. Wen, C.S.F. Wong, A.P. Wiegmans, A. Leimgruber, A. Moller,
Optimized exosome isolation protocol for cell culture supernatant and human plasma, J.
Extracell. Vesicles 4 (2015) 11.
[61] P. Jenjaroenpun, Y. Kremenska, V.M. Nair, M. Kremenskoy, B. Joseph, I.V. Kurochkin,
Characterization of RNA in exosomes secreted by human breast cancer cell lines using next-
generation sequencing, PeerJ 1 (2013) e201.
[62] G. Palazzolo, N.N. Albanese, G. Di Cara, D. Gygax, M.L. Vittorelli, I. Pucci-Minafra,
Proteomic analysis of exosome-like vesicles derived from breast cancer cells, Anticancer
Research 32(3) (2012) 847-860.
[63] ThermoFisherScientific, Boldly go above and beyond.
https://tools.thermofisher.com/content/sfs/brochures/Exosomes-brochure.pdf, 2016
(accessed COL12585 0916.).

[64] F. Liu, O. Vermesh, V. Mani, T.J. Ge, S.J. Madsen, A. Sabour, E.-C. Hsu, G. Gowrishankar,
M. Kanada, J.V. Jokerst, R.G. Sierra, E. Chang, K. Lau, K. Sridhar, A. Bermudez, S.J. Pitteri, T.
Stoyanova, R. Sinclair, V.S. Nair, S.S. Gambhir, U. Demirci, The Exosome Total Isolation Chip,
ACS Nano 11(11) (2017) 10712-10723.

[65] V. Ciravolo, V. Huber, G.C. Ghedini, E. Venturelli, F. Bianchi, M. Campiglio, D. Morelli, A.
Villa, P.D. Mina, S. Menard, Potential role of HER2-overexpressing exosomes in countering
trastuzumab-based therapy, Journal of Cellular Physiology 227(2) (2012) 658-667.

[66] J.N. Higginbotham, M.D. Beckler, J.D. Gephart, J.L. Franklin, G. Bogatcheva, G.-J.
Kremers, D.W. Piston, G.D. Ayers, R.E. McConnell, M.J. Tyska, Amphiregulin exosomes
increase cancer cell invasion, Current Biology 21(9) (2011) 779-786.

[67] D.J. Clark, W.E. Fondrie, Z. Liao, P.I. Hanson, A. Fulton, L. Mao, A.J. Yang, Redefining the
breast cancer exosome proteome by tandem mass tag quantitative proteomics and
multivariate cluster analysis, Analytical Chemistry 87(20) (2015) 10462-10469.

[68] K. Sandvig, A. Llorente, Proteomic analysis of microvesicles released by the human
prostate cancer cell line PC-3, Molecular & Cellular Proteomics 11(7) (2012) M111. 012914.

[69] S. Mathivanan, J.W. Lim, B.J. Tauro, H. Ji, R.L. Moritz, R.J. Simpson, Proteomics analysis
of A33 immunoaffinity-purified exosomes released from the human colon tumor cell line
LIM1215 reveals a tissue-specific protein signature, Molecular & Cellular Proteomics 9(2)
(2010) 197-208.

[70] J.S. Redzic, T.H. Ung, M.W. Graner, Glioblastoma extracellular vesicles: reservoirs of
potential biomarkers, Pharmacogenomics and personalized medicine 7 (2014) 65-77.

[71] D. Thuringer, A. Hammann, N. Benikhlef, E. Fourmaux, A. Bouchot, G. Wettstein, E.
Solary, C. Garrido, Transactivation of the epidermal growth factor receptor by heat shock
protein 90 via Toll-like receptor 4 contributes to the migration of glioblastoma cells, The
Journal of biological chemistry 286(5) (2011) 3418-28.

[72] M.W. Graner, R.I. Cumming, D.D. Bigner, The Heat Shock Response and
Chaperones/Heat Shock Proteins in Brain Tumors: Surface Expression, Release, and Possible
Immune Consequences, The Journal of Neuroscience 27(42) (2007) 11214-11227.

[73] A. Wade, A.E. Robinson, J.R. Engler, C. Petritsch, C.D. James, J.J. Phillips, Proteoglycans
and their roles in brain cancer, The FEBS journal 280(10) (2013) 2399-2417.

[74] K.L. Mooney, W. Choy, S. Sidhu, P. Pelargos, T.T. Bui, B. Voth, N. Barnette, I. Yang, The
role of CD44 in glioblastoma multiforme, Journal of Clinical Neuroscience 34 (2016) 1-5.

[75] J. Kraan, P. van den Broek, C. Verhoef, D.J. Grunhagen, W. Taal, J.W. Gratama, S. Sleijfer,
Endothelial CD276 (B7-H3) expression is increased in human malignancies and distinguishes
between normal and tumour-derived circulating endothelial cells, British journal of cancer
111(1) (2014) 149-56.

[76] V.P. Tryndyak, O. Kovalchuk, I.P. Pogribny, Loss of DNA methylation and histone H4
lysine 20 trimethylation in human breast cancer cells is associated with aberrant expression
of DNA methyltransferase 1, Suv4-20h2 histone methyltransferase and methyl-binding
proteins, Cancer Biology & Therapy 5(1) (2006) 65-70.

[77] J. Beliakoff, L. Whitesell, Hsp90: an emerging target for breast cancer therapy, Anti-
Cancer Drugs 15(7) (2004) 651-662.

[78] J.T. Price, T. Tiganis, A. Agarwal, D. Djakiew, E.W. Thompson, Epidermal growth factor
promotes MDA-MB-231 breast cancer cell migration through a phosphatidylinositol 3⁠-
kinase and phospholipase C-dependent mechanism, Cancer Research 59(21) (1999) 5475-
5478.

[79] J. de Vrij, S.L.N. Maas, K.M.C. Kwappenberg, R. Schnoor, A. Kleijn, L. Dekker, T.M. Luider,
L.D. de Witte, M. Litjens, M.E. van Strien, E.M. Hol, J. Kroonen, P.A. Robe, M.L. Lamfers,
M.W. Schilham, M.L.D. Broekman, Glioblastoma-derived extracellular vesicles modify the
phenotype of monocytic cells, International Journal of Cancer 137(7) (2015) 1630-1642.
871