Obesogenic diet exposure alters uterine natural killer cell biology and impairs vasculature remodeling in mice

Jennet Baltayeva[1,2], Chaini Konwar[1,3], Barbara Castellana[1,2], Danielle L. Mara[1], Julian K. Christians[1,4], Alexander G. Beristain[1,2]

[1] British Columbia Children's Hospital Research Institute, Vancouver, Canada.
[2] Department of Obstetrics & Gynecology, The University of British Columbia, Vancouver, Canada.
[3] Department of Medical Genetics, The University of British Columbia, Vancouver, Canada.
[4] Department of Biological Sciences, Simon Fraser University, Burnaby, Canada.

Running Title: Effects of obesogenic diet on uterine natural killer (uNK) cell biology

To whom correspondence should be addressed: Alexander G. Beristain, British Columbia Children's Hospital Research Institute, Department of Obstetrics & Gynecology, The University of British Columbia, Vancouver, British Columbia, Canada. V5Z 4H4. Tel: (604) 875-3573; E-mail: aberista@mail.ubc.ca

Keywords: Pregnancy, natural killer cell, uterus, decidua, placenta, diet-induced obesity, spiral artery remodeling, high-fat/high-sucrose diet, gene expression, gene microarray

Funding: This work was supported by a BC Children's Hospital Healthy Starts Catalyst Grant (to AGB and JKC) and a Canadian Institutes of Health Research Open Operating Grant (201403MOP-325905-CIA-CAAA) (to AGB).

ABSTRACT

Pre-pregnancy obesity associates with adverse reproductive outcomes that impact maternal and fetal

health. While obesity-driven mechanisms linked to poor pregnancy outcomes remain unclear, it is

possible that obesity affects local uterine immune cells. Uterine immune cells, particularly uterine

natural killer cells (uNK), play central roles in orchestrating developmental events in pregnancy.

However, the effect of obesity on uNK biology is poorly understood. Using a high fat/high sugar diet

(HFD) obesogenic mouse model, we set out to examine the effects of maternal obesity on uNK

composition and establishment of the maternal-fetal interface. HFD exposure resulted in weight gain-

dependent increases in systemic inflammation and rates of fetal resorption. Within HFD mice, both

weight gain-dependent (diet-induced obese; DIO) and -independent (DIO-resistant; DIO-R) effects on

natural cytotoxicity receptor-1 frequency and uNK activity were observed. Importantly, HFD-

associated changes in uNK coincided with impairments in uterine artery remodeling in mid but not late

pregnancy. Comparison of uNK mRNA transcripts in control diet mice to gene signatures in DIO or

DIO-R identified differentially expressed genes that play roles in promoting activity/cytotoxicity and

impeding vascular biology. Together, this work provides new insight into how obesity, and particularly

how diet-induced weight-gain, may impact uNK-processes important in pregnancy.

INTRODUCTION

Obesity affects more than half a billion adults worldwide and has become a global epidemic[1].

Consistent with trends in the general population, rates of obesity among women of childbearing age are

increasing[2–4]. Pre-pregnancy obesity is recognized as a high-risk state because of its association with

multiple adverse obstetric and perinatal outcomes including miscarriage, preeclampsia, preterm birth,

and intrauterine fetal demise[5,6]. Providing the interface between the mother and developing baby, the

conceptus-derived placenta plays an essential role in the transfer of nutrients and gases between

maternal and fetal circulation. Importantly, placenta development and fetal health is impaired in obese

women[7,8]. In rodent models of obesity, obesogenic diets promote placental dysfunction, defined in part

by defects in utero-placental vascularization and altered trophoblast maturation[9–11]. However, the

cellular and molecular processes affected by obesity that underlie detrimental obesity-related

pregnancy outcomes are poorly understood.

Optimal placental function and overall pregnancy success is largely dependent upon tightly

regulated angiogenic and vascular remodeling programs that take place within the maternal-fetal

interface[12,13]. Early stages of uterine blood vessel growth, branching, pruning, and remodeling are

controlled by diverse populations of leukocytes that reside within the decidual microenvironment[14,15].

In early human and mouse pregnancy, uterine natural killer cells (uNK) constitute the most abundant

immune cell subtype within the uterine mucosa[16–18]. Genetic studies in mice show that uNK are major

contributors to uterine vascular growth and remodeling, where these uNK-related processes are

mediated in part by the production of pro-angiogenic factors (i.e. vascular endothelial growth factor;

VEGF) and tissue-remodeling cytokines (i.e. tumor necrosis factor-α; TNF$-\alpha$, type-II interferon)[19–21].

Obesity associates with adipose tissue immune cell infiltration, including macrophages, T and B

lymphocytes, mast cells, and neutrophils[22–24], and along with these changes, obesogenic environments

alter leukocyte activation by potentiating the secretion of pro-inflammatory cytokines that have effects

within both local and distal tissues[25]. In women, previous work shows that uNK frequencies decrease in

obesity[26]. Importantly, these cellular changes associate with increased rates of uNK degranulation,

changes in natural killer receptor expression, and decreased production of pro-angiogenic VEGF-A[27].

While these findings provide insight into how maternal obesity affects uNK processes important in

pregnancy, due to the heterogeneous nature of obesity within human populations, the need for a clearer

understanding of how excess adiposity impacts uNK biology and how these changes affect pregnancy

outcome is still needed.

In this study, we established a diet-based mouse model to investigate the impact of obesity on

pregnancy outcome, uNK function, and placental development in early-mid pregnancy. Mice subjected

to a chronic high-fat/high-sugar (HFD) diet showed increased levels of systemic inflammation and

increased rates of fetal resorption. Notably, uNK composition within HFD mice was different than in

mice fed a control diet, where alterations in state of uNK activity and receptor expression were

observed. Notably, these uNK-related differences associated with uterine artery remodeling

impairments within HFD mice. While diet exposure did not result in robust global gene expression

changes in uNK, specific genes important in vascular biology, inflammation/cytotoxic activation, and

metabolism were altered by HFD exposure. Together, our work provides novel insight into obesity-

induced changes in uNK function. These findings also establish insight into immunological

mechanisms impacted by maternal obesity that likely lead to alterations in utero-placental development

in early-mid pregnancy which in turn may contribute to the development of obesity-linked pregnancy

disorders.

RESULTS

Diet-induced weight gain drives systemic inflammation and potentiates fetal resorption

To investigate the effects of maternal obesity on uNK biology and placental development,

female C57BL/6J mice were fed either a control low-fat/no sucrose diet (LFD) or a HFD over a 13-

week period prior to mating; exposure to these diets was also maintained during pregnancy (Figure

1A). As expected, mice fed a HFD showed an increase in body weight (Figure 1B). However, there

was considerable heterogeneity in endpoint body weight within the HFD cohort, where weight gain

rates in ~37% of HFD mice (7/19) were indistinguishable from LFD controls (Figures 1C, 1D). This

observation led us to sub-categorize mice within the HFD group as either weight-gainers (diet-induced

obese mice; DIO) or as weight-gain resistant mice (DIO-resistant; DIO-R) (Figure 1C). Sub-

categorization of DIO and DIO-R mice was established following the example of Enriori *et al*[28], where

mice with endpoint weights > 3 standard deviations to the mean LFD weight were categorized as DIO.

Consistent with previous findings showing association of HFD-induced weight gain and increased

systemic inflammation[29,30], levels of pro-inflammatory TNFα in blood serum were higher in DIO mice

than in LFD controls at gestational day (Gd)10.5 (Figure 1E). Notably, a statistical difference was not

observed in TNFα serum levels between DIO-R and LFD mice, suggesting that the increase in TNFα is

due to increased adiposity and not exposure to HFD *per se* (Figure 1E).

Neither HFD exposure nor increased weight gain led to differences in placental or fetal

measurements, although a trend for DIO-R placentas to weigh less than control placentas was observed

(Supplemental Figure 1A, B). However, HFD exposure led to moderate, but significant increases in the

fetal resorption rate compared to control mice (18% vs. 5%), and this effect was primarily driven by the

DIO subgroup (Figure 1F). Together, our mouse model of maternal obesity shows that HFD-induced

weight gain in pregnancy correlates with increased systemic inflammation and impaired fetal viability.

1   High-fat diet exposure does not alter uNK frequencies in early pregnancy

Uterine NK in mice are defined as "tissue-resident" Dolichos biflorus agglutinin

(DBA)$^+$CD49a$^+$DX5$^{+/-}$ (tr-uNK) or "conventional" DBA$^-$CD49a$^-$DX5$^+$ (c-uNK), and these distinct

subtypes play specific roles in uterine neo-angiogenesis (i.e. tr-uNK) and artery remodeling (i.e. c-

uNK)[31]. In mid pregnancy, tr-uNK are the dominant uNK subtype comprising roughly 80% of all uNK,

while c-uNK make up a smaller proportion[32]. To this end, we first investigated whether HFD exposure

alters the number of tr-uNK in early pregnancy by histologically examining DBA$^+$ cells at Gd7.5 and

Gd10.5 of pregnancy; at Gd7.5, uNK begin to expand in number and this expansion peaks by Gd10.5,

at which point numbers subsequently decline[33]. While DBA$^+$ uNK in both HFD and LFD mucosa

underwent normal gestational age-related increases in numbers, uNK numbers were not affected by diet

(Figure 2A).

Following these histological assessments, proportions of total leukocytes (CD45$^+$) and uNK

(CD3$^-$CD122$^+$) were examined by flow cytometry in Gd10.5 implantation sites, as were specific tr-

uNK, c-uNK, and uNK precursor cell sub-populations (Figure 2B). HFD exposure showed a modest,

but non-significant effect on total CD45$^+$ immune cell proportions, where an increasing trend in CD45$^+$

cell frequency was observed only within the DIO subgroup (Figure 2C). Similarly, total uNK

frequencies were not impacted by HFD or diet-induced weight gain (Figure 2D). Consistent with this,

HFD did not exert any measurable differences in proportions of tr-uNK, although an increase in tr-uNK

frequency was observed within DIO compared to DIO-R mice; a similar though non-significant

relationship (P = 0.1) between tr-uNK proportions in DIO and LFD control mice was also observed

(Figures 2E). Conventional-uNK made up a smaller proportion (~10-20%) of uNK compared to the

dominant tr-uNK population, where frequencies were also not altered by diet or diet-induced weight

gain (Figure 2F). Lastly, the effect of HFD on uNK precursors (CD122$^+$/NK1.1$^-$/DX5$^-$)[34] was

measured, and consistent with the above findings, HFD exposure did not affect precursor frequencies

(Figure 2G).

Proportions of NCR1[+] uNK increase in response to high-fat diet

Natural cytotoxicity receptor 1 (NCR1) is essential for uNK maturation and function, and plays

critical roles in promoting uterine blood vessel remodeling in mice[35,36]. Thus, to further elucidate the

effects of maternal HFD on uNK biology, NCR1 uNK proportions and expression levels were

examined by flow cytometry. While HFD did not impact cell surface levels of NCR1 expression (MFI),

mice exposed to HFD did show increased frequencies of NCR1[+] uNK (Figures 3A, B, C). Further

analysis revealed that the increase in NCR1[+] uNK in HFD mice is primarily driven by the DIO-R sub-

group; DIO mice showed a modest but non-significant increase in NCR1[+] uNK proportions ($P = 0.09$)

(Figure 3B). These findings suggest that obesogenic diet exposure, and to a lesser extent diet-associated

weight gain, manifest in subtle changes in uNK composition defined by NCR1 expression. Given the

importance of NCR1 in vascular processes in pregnancy, diet-induced differences in NCR1 may

translate to alterations in uterine artery remodeling or cytotoxicity.

HFD exposure drives uNK activity

Since NCR1 associates with the promotion of uNK maturation and activation, we next set out to

examine if HFD exposure alters uNK activity[36]. To do this, two approaches were used:

Immunofluorescence microscopy facilitated measurement of uNK cell diameter/size as a readout of

maturity[36], and flow cytometry analysis measured cell surface expression of the early activation marker

CD69[37]. Because NCR1 is expressed predominantly by DBA[+] cells at Gd10.5[36], focus was given to

measuring DBA[+] cell diameters within histological sections of utero-placental tissues. While cell

diameters of DBA$^+$ uNK were not shown to be different between HFD and control LFD mice

(Supplemental Figure 1C), CD69 expression levels (MFI) were significantly higher in both total (CD3$^-$

CD122$^+$) and NCR1$^+$ uNK populations as a result of HFD exposure (Figures 4A, B, C). In line with our

findings that NCR1 proportions are higher within the DIO-R sub-group than in control LFD or DIO

mice, the increase in HFD-induced CD69 expression was primarily attributed to uNK within the DIO-

R, and not the DIO subgroup, the latter of which showed greater heterogeneity in CD69 expression

(Figures 4B, C). Overall these findings suggest that HFD exposure, independent of weight gain,

instructs an altered state of uNK activity.

HFD impairs uterine spiral artery remodeling in mid but not late pregnancy

HFD-induced differences in uNK NCR1$^+$ frequency and activity suggest that pregnancy

associated processes controlled by uNK, like placental development[31] and uterine artery remodeling[38],

may be affected by HFD. To assess this, we first examined if specific placental layers/zones at Gd14.5

are altered by HFD; Gd14.5 represents a developmental time-point when the mouse placenta is mature

and fully functional. Total placental area (mm^2), consisting of labyrinth zone (L), junctional zone (Jz),

and decidua (D)[39] were not different between control LFD and HFD mice (Figure 5A). Subgrouping of

HFD placentas into DIO and DIO-R categories showed that diet-induced weight gain also did not affect

overall placental area (Figure 5A). Moreover, HFD exposure did not result in changes in area within

individual placental layers (i.e. L, JZ, and D) (Figure 5B, C, D). However, discrepancies in labyrinth

area were identified between placentas of DIO and DIO-R mice, where the labyrinth area in DIO-R

mice was greater (Figure 5B). Together our findings indicate that HFD-related uNK changes in early-

mid gestation do not translate into major changes in placental morphology.

To examine whether HFD-induced uNK changes translate into alterations in uterine artery

remodeling, measurements of Gd10.5 uterine arterial wall and vessel lumen areas, as well as their

ratios, were next determined (Figure 6A). We also assessed decidual artery smooth muscle actin (SMA)

expression, localization, and vessel intactness via immunofluorescence microscopy at Gd7.5, just prior

to initiation of vascular remodeling, and at Gd10.5, when remodeling is at its peak and can be assessed

by smooth muscle integrity by staining for SMA[40,41]. At Gd10.5, HFD-exposed mice had smaller

vessels and thicker arterial walls (Figure 6A), and accordingly had significantly higher wall:lumen area

ratios (Figure 6B). Interestingly, both DIO and DIO-R mice showed this arterial remodeling phenotype

(Figure 6B). Examining SMA intactness at Gd7.5 by immunofluorescence microscopy, control LFD

mice showed a marginal but significant increase in artery SMA coverage over HFD mice (Figure 6C,

D). However, by Gd10.5, SMA staining within control mice was more sparse (i.e. smaller

SMA:Perimeter ratio) than in HFD mice; HFD arteries showed only a partial loss in SMA staining, and

this was reflected with higher SMA:Perimeter ratios (Figure 6D). By Gd14.5, differences in blood

vessel remodeling observed within HFD mice were no longer observed (Figure 6E). Specifically,

uterine arteries in both LFD and HFD mice showed dilated lumens and thin vessel walls indicative of

proper remodeling (Figure 6E). Together, these findings suggest that maternal obesity disrupts uNK-

mediated vascular remodeling in early-mid gestation, but that compensatory processes facilitating

proper artery remodeling by mid-late gestation may overcome these initial impairments.

HFD exposure leads to subtle changes in uNK gene expression

To gain insight into possible uNK-related mechanisms altered by HFD exposure, global gene

expression analyses were performed on mRNAs isolated from flow cytometry-sorted

CD45[+]/CD122[+]/CD3[-] uNK using gene microarrays (Figure 7A). Following standard probe filtering,

normalization, and batch correction (Supplemental Figure 2A), differential gene expression comparison

(FDR < 0.05), principal component (PCA), and hierarchical cluster analyses were performed between

control LFD, DIO, and DIO-R uNK subgroups. Surprisingly, differential gene expression analysis did

not identify any differentially expressed genes between the uNK subgroups (Supplemental Figure 2B),

likely owing to the underpowered and heterogeneous composition of HFD uNK. PCA clustering based

on global gene signatures further demonstrated that mouse uNKs exposed to HFD do not harbor unique

or drastic differences in gene transcription, where uNK sub-group clustering showed DIO and DIO-R

uNK to cluster randomly (Supplemental Figure 2C).

Despite the inability of a global gene signature approach to identify HFD-specific genes altered

in uNK, hierarchical clustering of LFD, DIO, and DIO-R uNK using a 578 gene signature derived from

raw *P*-value ranked genes (*P*<0.05, fold-change > 1.25) segregated uNK into diet-associated sub-

groups with 85-90% accuracy (Figure 7B). This finding suggests that HFD and associated weight gain

may subtly affect the expression of specific uNK genes that were previously not identifiable due to the

statistical stringency applied to the global high dimensional dataset. Of the selected 578 rank-ordered

genes, one-way ANOVA with post-hoc Tukey HSD identified 71 genes to be differentially expressed

within LFD, DIO, and DIO-R subgroups (Supplemental Table 1). Specifically, sets of genes linked

with uNK cytotoxicity and activation were expressed at higher levels within both DIO (*Cd200r4,*

Il18rap) and DIO-R (*Cd226*) subgroups (Figure 7C). In line with our findings that uterine vascular

remodeling was affected by high-fat diet, expression of pro-angiogenic/pro-remodeling *Vegfa*,

Adamts3, *Nr4a1* and *Pde5a* were shown to be altered in DIO and DIO-R subgroups compared to LFD

controls, with *Vegfa*-regulating *Anxa8* being inversely expressed within HFD samples compared to

controls (Figure 7D). Targeted qPCR analysis of *Vegfa* in a subset of the same uNK mRNA samples

used in the gene array supported the array's finding (Supplemental Figure 2D). Moreover, HFD also

resulted in alterations in levels of genes important in metabolism, where *H6pd* and *Hk2*, genes central

to glucose metabolism, and *Renbp*, regulator of amino-sugar metabolism, were up-regulated in DIO

and DIO-R subgroups (Figure 7E). Levels of *Scd1*, a gene important in lipid metabolism, were higher

in the LFD control group (Figure 7E). Taken together, while chronic HFD exposure does not lead to

robust global changes in uNK gene signatures, HFD does modestly affect the expression of select genes

important in regulating uNK-mediated cytotoxicity/activation, vascular biology, and metabolism. These

weight-gain dependent and independent differences provide insight into the biological underpinnings of

how maternal obesity affects uNK processes important in pregnancy.

DISCUSSION

In this study, we examine the effects of maternal HFD on uNK cell biology and uterine vascular

transformations in early and mid-pregnancy. We demonstrate that HFD does not alter total uNK

numbers or subsets of tissue-resident or conventional cells. However, weight gain-specific effects on tr-

uNK were observed, and additionally exposure to HFD results in a selective increase in NCR1$^+$ uNK

frequencies and overall uNK activity. These diet-related uNK changes associate with impairments in

uterine artery remodeling in early-mid pregnancy, but were no longer observed by mid-late pregnancy,

suggesting that a compensatory mechanism in uNK function may be taking place. While examination

of global gene expression signatures did not identify specific uNK genes or pathways altered by HFD,

gene signature-informed hierarchical clustering did segregate uNK according to diet exposure with

relative high accuracy, indicating that uNK genes and gene pathways are moderately affected in

obesogenic mice. Genes identified as being differentially expressed between control and HFD sub-

groups signified that uNK exposed to HFD harbor pro-cytotoxic and anti-angiogenic characteristics

that together provide new insight into how maternal obesity may alter uNK processes critical to early

pregnancy establishment and health.

To our knowledge, this is the first study to examine the effects of HFD on uNK cell proportions

and states of activation in pregnancy. Importantly, this work also examines the specific effects of HFD-

exposure and HFD-associated weight gain. While a previous study did report on the effects of maternal

Western diet exposure on NK properties within draining inguinal lymph nodes of the uterus[10], our

study directly examines uNK tissue-resident and conventional subsets within the decidual mucosa.

Because uNK are distinct from peripheral blood and lymphatic NKs[34,42,43], our findings generate new

insight into how maternal obesity potentially impacts immune cell function within the maternal-fetal

interface. While HFD exposure does not affect the frequency of total leukocytes or uNK subtypes

within the uterus, notably we show that HFD exposure increases both the proportion of NCR1[+] uNK

and uNK activation. Strikingly, differences in NCR1 frequency and uNK activation were primarily

observed in HFD mice that were resistant to diet-induced weight gain (i.e. DIO-R), suggesting that the

HFD-related features altered in uNK and uterine arteries may not necessarily be related to changes in

overall adiposity. Our finding that weight-gain related effects on elevated blood serum TNFα and

increased rates of fetal resorption may therefore be unrelated to changes in uNK biology. However, due

to inter- sub-group heterogeneity and overall modest sample size, further work is needed to investigate

the direct role of excessive adipose tissue (i.e. obesity) and diet-related factors like fatty acids and

metabolites that may also lead to uterine immune cell dysfunctions.

The impact of obesogenic-diets on fetal health and pregnancy outcome in rodents has not been

consistently reported on. Prior studies have reported that HFD exposure in pregnancy results in both

reduced[44,45] and increased[46,47] fetal weight, and reduced placental labyrinth[48] and junctional zone[44]

thickness, or no differences in placental weights[45,49,50]. In our study, placental and fetal weights and

placental/decidual areas were not affected by HFD. Notably, HFD exposure did result in increased rates

in fetal resorption, although this effect was modest. Important and somewhat counter to our findings

showing that DIO-R uNK have elevated activity and expression of NCR1, the increase in fetal

resorption is primarily attributed to DIO mice, and not weight gain-resistant DIO-R mice. The

underlying cause for this observation is unknown but does nonetheless highlight the importance of

needing to subcategorize mice subjected to experimental diet models of obesity as true weight gainers

or as weight gain-resistant mice. In our hands, DIO and DIO-R murine uNK showed modest, but

significant differences in specific parameters, and these differences were in part associated by weight

gain-specific differences in gene expression.

1    Our focus on NCR1 stems from previous work reporting that $Ncr1^{-/-}$ mice exhibit defects in

2  decidual angiogenesis and uterine vascular remodeling[36]. Notably, inhibition of NCR1 delays uNK cell

3  maturation and activation, reflected by smaller cell diameters and reductions in cytokine production[51].

4  Our finding that HFD led to inverse associations between frequencies of NCR1$^+$ uNK and degree of

5  vascular remodeling (i.e. elevated NCR1 frequencies and impaired arterial remodeling) is inconsistent

6  with what is known about the role of NCR1 in uNK biology. At least two possible explanations can

7  account for this discrepancy. Firstly, the increase in NCR1$^+$ uNK may reflect a level of compensation,

8  and this is supported by the fact that initial impairments in vascular structure are no longer observed by

9  Gd14.5.

10    A second possibility centers on the role of NCR1 as a key activating receptor important in

11  promoting cytotoxic functions of NK[52–54]. Perhaps, elevated NCR1 frequency indicates that uNK

12  exposed to HFD are either overly activated or are aberrantly cytotoxic. In line with this speculation, we

13  show that mRNA levels of the pro-cytotoxic receptor *Cd226* (encoding for DNAM-1) are elevated

14  within HFD uNK. DNAM-1 associates with NK maturation and memory and CD226$^+$ NK are more

15  cytotoxic and produce higher levels of inflammatory cytokines than do CD226$^-$ counterparts[55,56].

16  Intriguingly, *Cd226* levels are more abundant within DIO-R, while transcripts encoding for CD200R4,

17  a DAP12-associated activating receptor[57], are expressed at higher levels in DIO uNK, suggesting that

18  distinct mechanisms of activation are induced in relation to diet and/or adiposity/weight gain. Overall,

19  these possibilities are consistent with a recent finding in human uNK that correlates maternal obesity to

20  elevated rates of uNK degranulation[27]. However, it remains to be tested if obesity-linked hyper-

21  activation in uNK equates to increased killing and dysfunction of cells within the maternal-fetal

22  interface. Further, if indeed the case, cell-type specific uNK targets need also to be identified, as this

will help in our understanding of the biological consequences of heightened uNK activity in response

to maternal obesity and/or HFD.

Of interest, and consistent with previous findings in uNK from obese women[27], HFD exposure

in mice results in a decrease in *Vegfa* expression. While multiple cell types within the utero-placental

environment are responsible for VEGF-A production, including decidual stromal cells[58,59] and

macrophages[60], the transient nature of uNK in pregnancy and their links to vascular growth, branching,

and remodeling suggest that uNK deficiency in VEGF-A production could directly translate into

impairments in uterine vascular biology. VEGF-A is also a ligand for Flt-1 expressing trophoblasts,

where VEGF-A plays roles in controlling trophoblast survival and proliferation[61–63], however our

finding that HFD exposure does not impact the development of specific zones of the placenta indicates

that VEGF-A dysregulation may play a more significant and specific role in uterine artery

angiogenesis.

It is becoming widely appreciated that cellular metabolism plays a role in regulating NK cell

function and education[64]. A recent study demonstrated that glucose metabolism is required for receptor-

activated NK cell IFNγ production[65]. Accordingly, Michelet *et al*[66] showed that obesity-driven

metabolic paralysis results blunted NK cell cytotoxic responses. In line with these findings we

demonstrate an increase in expression of *H6pd* and *Hk2* and decreased levels of *Scd1* in DIO and DIO-

R uNKs, where *H6pd* and *Hk2* are important regulators of glycolysis and *Scd1* is involved in fatty acid

metabolism. Overall, it is likely that the obesogenic environment in HFD mice may induce a metabolic

switch in uNKs that impacts maturation and function that may ultimately alter maternal-fetal interface

remodeling and overall uNK education. Potential metabolic effects on uNK processes will be of high

interest to further examine as metabolic dysfunction and altered programming of uNK in human

pregnancies could in part contribute to impaired pregnancy outcomes.

uNKs are specialized immune cells that play an important role in coordinating reproductive

success. Deviations in normal uNK function may contribute to defective vascular modifications or

aberrant induction of cytotoxic potential at the maternal-fetal interface. Such uNK-related changes may

play roles in pregnancy complications. Our study provides insight into how exposure to an obesogenic

diet alters uNK activity and alters spiral artery remodeling, highlighting possible cellular mechanisms

underlying HFD exposure. Maternal obesity continues to pose serious consequences on reproductive

health, thus future studies aimed to further dissect how obesity-driven immune and cellular alterations

impact pregnancy health will be of substantial importance.

METHODS

Animals. Mouse experiments were carried out in accordance with the guidelines of the

Canadian Council on Animal Care and approved by SFU University Animal Care Committee (protocol

1094) and the UBC Animal Care Committee (protocol A17-0060). C57BL/6J mice were purchased

from the Jackson Laboratory (Bar Harbor, ME, USA). Animals were housed under standard conditions

in groups of max. 5 mice in individually ventilated cages with water and food available *ad libitum*,

under a 12:12h light/dark cycle.

High-fat/high sugar obesogenic model. At 9 weeks of age, females were placed on either a

high-fat, high-sucrose diet (HFHS; 45% kcal fat, 35% carbohydrate (including 17% kcal sucrose), 20%

kcal protein, 4.73 kcal/g, D12451, Research Diets, New Brunswick, NJ) or a nutrient-matched low-fat,

no-sucrose control diet (CON; 10% kcal fat, 70% kcal carbohydrate (corn starch and maltodextrin),

20% kcal protein, 3.85 kcal/g, D12450K, Research Diets). Females were kept on the experimental diets

for 13 weeks prior to being paired with male studs. The appearance of a plug the next morning was

taken as day 0.5 of pregnancy. Because not all females became pregnant when first paired, some had

been on the experimental diet for 13–19 weeks at the start of pregnancy. On days 7.5, 10.5 and 14.5 of

pregnancy females were euthanized and their blood collected via cardiac puncture. The utero-placental

tissues with multiple implantation sites were collected from each mouse.

ELISA measurements of serum TNFα. Serum was purified from whole blood of females

sacrificed at Gd10.5 of pregnancy using serum separator tubes following standard procedures. Serum

levels of TNFα were determined using commercially available Mouse TNFα ELISA kit (R&D

systems) according to manufacturer's protocol.

Decidual leukocyte preparations. For examination of uNK cell populations, murine

implantation sites were separated from uterine tissues and were mechanically and enzymatically

processed to obtain single-cell suspensions. In brief, tissues were washed extensively in cold

phosphate-buffered saline (PBS; pH 7.4), and following this, tissues were minced with a razor blade

and subjected to a 30-minute enzymatic digestion at 37 °C in 3 mL of 1:1 DMEM/F12 media (Gibco,

Grand Island, NY; 200 mM L-glutamine) with 1X-collagenase/hyaluronidase (10X stock; StemCell

Technologies, Vancouver, Canada), 80 μg/mL DNaseI (Sigma, St. Louis, MO),

penicillin/streptomycin, and Anti antimycotic solution (100X dilution, Gibco). Single cell suspensions

were passed through a 100μm strainer and decidual leukocyte enrichment was performed by Percoll

(GE Healthcare) density gradient (layered 40%/80%) centrifugation. To avoid red blood cell

contamination, cell mixture was briefly incubated with NH_4Cl. Decidual leukocytes were immediately

used for uNK isolation by fluorescence activated cell sorting (FACS) or cell-surface marker

characterization via flow cytometry analyses.

Flow cytometry. Flow cytometry analyses were carried out following a standard protocol. In

brief, $0.5x10^6$ isolated leukocytes were incubated with Fc Block (anti-CD16/CD32 antibody at 1:35

dilution in PBS; BD Pharmingen) for 10 min on ice to prevent non-specific binding. Cell suspensions

were further incubated with fluorescent-conjugated antibodies directed against cell-surface markers for

30 minutes at 4°C. The following antibodies were used: APC ef780 CD3e (1:50; clone 145-2C11;

eBioscience), ef450 CD45 (1:100; clone 30-F11; eBioscience), PerCP710 CD122 (1:100; clone TM-

b1; eBioscience), APC CD122 (1:35; clone TM-Beta 1; BD Pharmingen), APC CD49a (1:50; clone

HMα1; BioLegend), PeCy7 CD49b (1:100; clone DX5; eBioscience), APC CD69 (1:25; clone H1.2F3;

eBioscience), PeCy7 NKp46 (1:50; clone 29A1.4; eBioscience). Following staining, cells were re-

suspended in 7-AAD viability staining solution (at 1:25 dilution in PBS+ 2%FBS +1%

penicillin/streptomycin; eBioscience). Samples were acquired on an LSRII FACS (BD Biosciences)

and data was analysed by FlowJo 5.0 software (Tree Star, Inc., Ashland, OR, USA).

Immunohistochemistry, IF microscopy and morphometry. Implantation sites from Gd7.5,

Gd10.5 and Gd14.5were fixed in 4% paraformaldehyde for 24-48h followed by processing into

paraffin blocks. Six-µm sections were stained with hematoxylin and eosin (H&E) according to standard

methods. For IF staining, tissues underwent antigen retrieval by heating slides in a microwave for 5

minutes in 30-second intervals in a sodium citrate buffer (pH 6.0). Sections were then incubated in a

blocking solution (5% bovine serum albumin (BSA) in Tris-buffered saline with 0.05% Tween-20

(TBST)) for 1 hour at room temperature. Further, samples were incubated with the following

antibodies overnight at 4°C: rabbit polyclonal alpha-smooth muscle actin (1:200; Abcam) and

biotinylated DBA (1:100; Vector Laboratories). Slides were incubated with Alexa Fluor 568/488–

conjugated streptavidin and goat anti-rabbit secondary antibodies (Life Technologies) for 1 hour at

room temperature. Glass coverslips were mounted onto slides using ProLong Gold anti-fade reagent

containing DAPI (Life Technologies).

DBA$^+$ cell assessments. DBA$^+$ cell quantities in Gd7.5 and Gd10.5 implantation sites were

calculated based on DBA fluorescence intensity threshold. Slides were imaged using a 20x Plan-

Apochromat/0.80 NA or a 40x EC-Plan-Neofluar/0.9 PoI objective (Carl Zeiss). Data was processed

and analyzed using ZenPro software (Carl Zeiss). DBA$^+$ cell diameters in Gd10.5 implantation sites

were measured by randomly selecting 100 DBA$^+$ uNK cells from each implantation site. Slides were

imaged using a 40x EC-Plan-Neofluar/0.9 PoI objective (Carl Zeiss). Using ImageJ v1.50 software

(NIH), the diameters of these cells were measured across their longest and shortest axes; an average

of both values was used for statistical analysis. All images were obtained using Axiocam 506

monochrome digital camera (Carl Zeiss).

Spiral artery remodeling and SMA layer assessments. Spiral artery remodeling measurements

were performed on H&E stained slides. For Gd10.5, (HFD: n=9 implantation sites from 8 independent

pregnancies; LFD: n=11 implantation sites from 9 independent pregnancies) all of the spiral arteries

within the decidua were measured in triplicate (three sections 48μm apart from each other). For

Gd14.5, (HFD: n=18 implantation sites from 7 independent pregnancies; LFD: n=17 implantation sites

from 6 independent pregnancies) all of the spiral arteries within the decidua were measured in duplicate

(two sections 48μm apart from each other). Size of arterial walls and lumens was determined by

measuring cross-sectional areas with ImageJ v1.50 software (NIH). Wall:lumen ratios were calculated

by dividing the arterial wall area (μm^2) into the artery lumen area (μm^2). Slides were imaged using a

20x Plan-Apochromat/0.80 NA objective (Carl Zeiss). All images were obtained using Axiocam 105

color digital camera (Carl Zeiss). SMA layer intactness was measured in spiral arteries of Gd7.5

(HFD: n=15 arteries from 4 independent pregnancies; LFD: n=13 arteries from 4 independent

pregnancies) and Gd10.5 (HFD: n=19 arteries from 8 independent pregnancies; LFD: n=12 arteries

from 7 independent pregnancies). Slides were imaged using a 40x EC-Plan-Neofluar/0.9 PoI

objective (Carl Zeiss). Arterial length and SMA layer were measured using ImageJ v1.50 software

(NIH). SMA:Perimeter ratios were calculated by dividing SMA[+] vascular perimeter (μm) over total

arterial perimeter (μm). All images were obtained using Axiocam 506 monochrome digital camera

(Carl Zeiss).

Placental morphology assessments. H&E stained slides of Gd14.5 implantation sites were

used for placental morphology measurements. Total area (mm^2) and areas of each individual

placental layer (mm^2) were measured using ImageJ v1.50 software (NIH). To normalize for

differences in total placental area among the measurements, individual layer area was expressed as a

percentage of total area at each measurement point. Slides were imaged using 10x N-

Achroplan/0.25 Ph1 objective (Carl Zeiss). All images were obtained using Axiocam 105 color

digital camera (Carl Zeiss).

RNA extraction. Prior to RNA extraction, CD45+CD3-CD122+ uNK cells from Gd10.5

implantation sites were isolated by FACS using a FACSAria flow cytometry (BD Biosciences).

Total RNA was prepared from isolated uNKs using TRIzol LS reagent (Ambion) following by

RNeasy MinElute Cleanup (Qiagen) and DNase treatment (Ambion) according to manufacturers'

protocols. RNA purity was confirmed using a NanoDrop Spectrophotometer (Thermo Fisher

Scientific) and Agilent 2100 Bioanalyzer (Agilent). Only RNA samples having an RNA integrity

number (RIN) greater than 8.0 were used.

Microarray analysis. Total RNA samples extracted from mouse uNKs were sent to Génome

Québec Innovation Centre (McGill University, Montréal, Canada) for RNA quantification. Briefly,

mouse RNA samples were prepared for transcriptome profiling using the GeneChip ™ Pico Reagent

Kit (Affymetrix) as per manufacturer's protocol. Samples were run on the Clariom™ S Mouse Array to

measure gene expression at >20000 genes in the mouse genome. Raw data generated from the arrays

were read into R statistical software (version 3.5.1) with the Bioconductor *oligo*[67] package to convert

raw Affymetrix CEL files into an expression matrix of intensity values. Within oligo package samples

were background corrected, normalized, and log-transformed. Probe filtering identified 718 probes that

mapped to multiple gene symbols were excluded from the dataset. Additionally, 6923 Affymetrix

control probes and 749 probes that were not annotated to any given gene were also removed. A "soft"

intensity-based probe filtering[68], as recommended by *limma,* was performed to eliminate 2181 low

variance and low intensity probes. Together, these measures eliminated 10571 probes, leaving a total of

18558 probes remaining for further analysis. Using ComBat from *sva* package, known technical

variation associated with batch was removed. Following preprocessing, differential expression analysis

between groups was performed by applying a linear model to expression values using the *limma*

package's moderated t-statistics with empirical Bayesian variance estimation[69].

cDNA synthesis, and qPCR. RNA was reverse-transcribed using a first-strand cDNA

synthesis kit (Quanta Biosciences) and subjected to qPCR (ΔΔCT) analysis, using PerfeCTa SYBR

Green FastMix Low ROX (Quanta Biosciences) on an ABI ViiA 7 Real-Time PCR system (Thermo

Fisher Scientific). Forward and reverse primer sets used as previously described: *VEGFA* (F: 5'-

GTGCACTGGACCCTGGCTTTA-3', R: 5'- GGTCTCAATCGGACGGCAGTA-3')[70]; *HPRT* (F: 5'-

CTGGTGAAAAGGACCTCTCG-3', R: 5'-TGAAGTACTCATTATAGTCAAGGGCA-3')[71]. All raw

data were analyzed using QuantStudio V1.3 software (Thermo Fisher Scientific). The threshold cycle

(CT) values were used to calculate relative gene expression levels. Values were normalized to *HPRT*

transcripts.

Statistical analysis. Unless stated otherwise, statistical significance for all data was assessed by

unpaired two-tailed Student's t-tests. $P < 0.05$ was considered significant. Statistical analyses were

performed using Prism7 software (GraphPad Software, Inc.).

ACKNOWLEDGEMENTS

We are grateful to staff from Simon Fraser University Animal Care facility for their assistance.
We would like to extend our gratitude to BCCHRI Flow Core Facility Manager, Dr. Lisa Xu, for her
expertise and assistance in cell sorting of uNK cells. We are grateful to staff at Génome Québec
Innovation Centre at McGill University for their help with microarray hybridization.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

AUTHOR CONTRIBUTIONS

AGB and JKC designed the research. JB, CK, BC, DM, JKC and AGB performed the
experiments. JB, CK, WPR, AGB and JKC analysed the data and wrote the paper. All authors read and
approved the manuscript.

REFERENCES

1. World Health Organisation. WHO | Obesity and overweight. *WHO* (2017).

2. Statistics Canada. Canadian Community Health Survey: Obesity among children and adults. Available at: http://www.statcan.gc.ca/daily-quotidien/050706/dq050706a-eng.htm. (Accessed: 26th February 2018)

3. Ogden, C. L., Carroll, M. D., Kit, B. K. & Flegal, M. Prevalence of Childhood and Adult Obesity in the United States, 2011–2012. *Jama* **311,** 806–814 (2016).

4. Public Health Agency of Canada. Obesity in Canada: A joint report from the Public Health Agency of Canada and the Canadian Institute for Health Information. *Report No. Cat.: HP5-107/2011E-PDF.* (2011). Available at: https://www.canada.ca/en/public-health/services/health-promotion/healthy-living/obesity-canada.html. (Accessed: 26th February 2018)

5. Leddy, M. A., Power, M. L. & Schulkin, J. The impact of maternal obesity on maternal and fetal health. *Rev. Obstet. Gynecol.* **1,** 170–8 (2008).

6. Bautista-Castaño, I. *et al.* Maternal obesity in early pregnancy and risk of adverse outcomes. *PLoS One* (2013). doi:10.1371/journal.pone.0080410

7. Denison, F. C., Roberts, K. A., Barr, S. M. & Norman, J. E. Obesity, pregnancy, inflammation, and vascular function. *Reproduction* **140,** 373–385 (2010).

8. Schmatz, M., Madan, J., Marino, T. & Davis, J. Maternal obesity: The interplay between inflammation, mother and fetus. *Journal of Perinatology* **30,** 441–446 (2010).

9. Hayes, E. K. *et al.* Trophoblast invasion and blood vessel remodeling are altered in a rat model of lifelong maternal obesity. *Reprod. Sci.* **21,** 648–57 (2014).

10. Parker, V. J., Solano, M. E., Arck, P. C. & Douglas, A. J. Diet-induced obesity may affect the

uterine immune environment in early–mid pregnancy, reducing NK-cell activity and potentially compromising uterine vascularization. *Int. J. Obes.* **38,** 766–774 (2014).

11. Hayes, E. K. *et al.* Adverse fetal and neonatal outcomes associated with a life-long high fat diet: Role of altered development of the placental vasculature. *PLoS One* **7,** (2012).

12. Osol, G. & Moore, L. G. Maternal Uterine Vascular Remodeling During Pregnancy. *Microcirculation* **21,** 38–47 (2014).

13. Bardin, N., Murthi, P. & Alfaidy, N. Normal and Pathological Placental Angiogenesis. *BioMed Research International* **2015,** (2015).

14. Cartwright, J. E., Fraser, R., Leslie, K., Wallace, A. E. & James, J. L. Remodelling at the maternal-fetal interface: Relevance to human pregnancy disorders. *Reproduction* **140,** 803–813 (2010).

15. Faas, M. M. & de Vos, P. Uterine NK cells and macrophages in pregnancy. *Placenta* **56,** 44–52 (2017).

16. Moffett-King, A. Natural killer cells and pregnancy. *Nat. Rev. Immunol.* **2,** 656–663 (2002).

17. Felker, A. M. & Croy, B. A. Uterine natural killer cell partnerships in early mouse decidua basalis. *J. Leukoc. Biol.* **100,** 645–655 (2016).

18. Croy, B. A. *et al.* Imaging of vascular development in early mouse decidua and its association with leukocytes and trophoblasts. *Biol. Reprod.* **87,** 125 (2012).

19. Ashkar, A. A., Di Santo, J. P. & Croy, B. A. Interferon gamma contributes to initiation of uterine vascular modification, decidual integrity, and uterine natural killer cell maturation during normal murine pregnancy. *J. Exp. Med.* **192,** 259–70 (2000).

20. Barber, E. M. & Pollard, J. W. The uterine NK cell population requires IL-15 but these cells are not required for pregnancy nor the resolution of a Listeria monocytogenes infection. *J.*

Immunol. **171,** 37–46 (2003).

21.   Guimond, M.-J. *et al.* Absence of Natural Killer Cells during Murine Pregnancy Is Associated

with Reproductive Compromise in TgE26 Mice. *Biol. Reprod.* **56,** 169–179 (1997).

22.   Xu, H. Obesity and metabolic inflammation. *Drug Discovery Today: Disease Mechanisms* **10,**

(2013).

23.   Stolarczyk, E. Adipose tissue inflammation in obesity: a metabolic or immune response?

Current Opinion in Pharmacology **37,** 35–40 (2017).

24.   Patel, P. S., Buras, E. D. & Balasubramanyam, A. The role of the immune system in obesity

and insulin resistance. *Journal of Obesity* **2013,** (2013).

25.   Seijkens, T., Kusters, P., Chatzigeorgiou, A., Chavakis, T. & Lutgens, E. Immune cell crosstalk

in obesity: A key role for costimulation? *Diabetes* **63,** 3982–3991 (2014).

26.   Perdu, S. *et al.* Maternal obesity drives functional alterations in uterine NK cells. *JCI Insight*

1, (2016).

27.   Castellana, B. *et al.* Maternal obesity alters uterine NK activity through a functional

KIR2DL1/S1 imbalance. *Immunol. Cell Biol.* 805–819 (2018). doi:10.1111/imcb.12041

28.   Enriori, P. J. *et al.* Diet-Induced Obesity Causes Severe but Reversible Leptin Resistance in

Arcuate Melanocortin Neurons. *Cell Metab.* (2007). doi:10.1016/j.cmet.2007.02.004

29.   Heerwagen, M. J. R., Stewart, M. S., de la Houssaye, B. A., Janssen, R. C. & Friedman, J. E.

Transgenic Increase in N-3/N-6 Fatty Acid Ratio Reduces Maternal Obesity-Associated

Inflammation and Limits Adverse Developmental Programming in Mice. *PLoS One* (2013).

doi:10.1371/journal.pone.0067791

30.   Murabayashi, N. *et al.* Maternal high-fat diets cause insulin resistance through

inflammatory changes in fetal adipose tissue. *Eur. J. Obstet. Gynecol. Reprod. Biol.* (2013).

1   doi:10.1016/j.ejogrb.2013.02.003

2 31. Gaynor, L. M. & Colucci, F. Uterine natural killer cells: Functional distinctions and influence

3   on pregnancy in humans and mice. *Frontiers in Immunology* **8,** (2017).

4 32. Sojka, D. K. *et al.* Tissue-resident natural killer (NK) cells are cell lineages distinct from

5   thymic and conventional splenic NK cells. *Elife* (2014). doi:10.7554/eLife.01659

6 33. Croy, B. A., Yamada, A. T., DeMayo, F. J. & Adamson, S. L. *The Guide to investigation of mouse*

7   *pregnancy*. *The Guide to Investigation of Mouse Pregnancy* **1,** (2015).

8 34. Yadi, H. *et al.* Unique Receptor Repertoire in Mouse Uterine NK cells. *J. Immunol.* **181,**

9   6140–6147 (2008).

10 35. Felker AM, C. B. Natural cytotoxicity receptor 1 in mouse uNK cell maturation and function.

11   *Mucosal Immunol.* (2017). doi:10.1038/mi.2016.126

12 36. Felker, A. M., Chen, Z., Foster, W. G. & Croy, B. A. Receptors for non-MHC ligands contribute

13   to uterine natural killer cell activation during pregnancy in mice. *Placenta* **34,** 757–764

14   (2013).

15 37. Mosimann, B. *et al.* Natural killer cells and their activation status in normal pregnancy. *Int.*

16   *J. Reprod. Med.* **2013,** 906813 (2013).

17 38. Moffett, A. & Colucci, F. Uterine NK cells: Active regulators at the maternal-fetal interface.

18   *Journal of Clinical Investigation* **124,** (2014).

19 39. Coan, P. M., Ferguson-Smith, A. C. & Burton, G. J. Developmental Dynamics of the Definitive

20   Mouse Placenta Assessed by Stereology1. *Biol. Reprod.* **70,** 1806–1813 (2004).

21 40. Lima, P. D. A., Zhang, J., Dunk, C., Lye, S. J. & Anne Croy, B. Leukocyte driven-decidual

22   angiogenesis in early pregnancy. *Cellular and Molecular Immunology* **11,** 522–537 (2014).

23 41. Kieckbusch, J., Gaynor, L. M. & Colucci, F. Assessment of Maternal Vascular Remodeling

During Pregnancy in the Mouse Uterus. *J. Vis. Exp.* (2015). doi:10.3791/53534

42. Koopman, L. A. *et al.* Human Decidual Natural Killer Cells Are a Unique NK Cell Subset with

Immunomodulatory Potential. *J. Exp. Med.* **198,** 1201–1212 (2003).

43. Erick, T. K. & Brossay, L. Phenotype and functions of conventional and non-conventional

NK cells. *Current Opinion in Immunology* **38,** 67–74 (2016).

44. Mark, P. J. *et al.* A maternal high-fat diet in rat pregnancy reduces growth of the fetus and

the placental junctional zone, but not placental labyrinth zone growth. in *Journal of*

Developmental Origins of Health and Disease **2,** 63–70 (2011).

45. King, V., Hibbert, N., Seckl, J. R., Norman, J. E. & Drake, A. J. The effects of an obesogenic diet

during pregnancy on fetal growth and placental gene expression are gestation dependent.

Placenta **34,** 1087–1090 (2013).

46. Jones, H. N. *et al.* High-fat diet before and during pregnancy causes marked up-regulation of

placental nutrient transport and fetal overgrowth in C57/BL6 mice. *FASEB J.* **23,** 271–278

(2009).

47. Rosario, F. J., Kanai, Y., Powell, T. L. & Jansson, T. Increased placental nutrient transport in a

novel mouse model of maternal obesity with fetal overgrowth. *Obesity* **23,** 1663–1670

(2015).

48. Song, L. *et al.* Prenatal high-fat diet alters placental morphology, nutrient transporter

expression, and mtorc1 signaling in rat. *Obesity* **25,** 909–919 (2017).

49. Crew, R. C., Waddell, B. J. & Mark, P. J. Maternal obesity induced by a 'cafeteria' diet in the

rat does not increase inflammation in maternal, placental or fetal tissues in late gestation.

Placenta **39,** 33–40 (2016).

50. Kamimae-Lanning, A. N. *et al.* Maternal high-fat diet and obesity compromise fetal

hematopoiesis. *Mol. Metab.* **4,** 25–38 (2015).

51. Felker, A. M. & Croy, B. A. Natural cytotoxicity receptor 1 in mouse uNK cell maturation and

function. *Mucosal Immunol.* **10,** 1122–1132 (2017).

52. Glasner, A. *et al.* Recognition and Prevention of Tumor Metastasis by the NK Receptor

NKp46/NCR1. *J. Immunol.* (2012). doi:10.4049/jimmunol.1102461

53. Glasner, A. *et al.* Increased NK cell immunity in a transgenic mouse model of NKp46

overexpression. *Sci. Rep.* (2017). doi:10.1038/s41598-017-12998-w

54. Sheppard, S. *et al.* The Murine Natural Cytotoxic Receptor NKp46/NCR1 Controls TRAIL

Protein Expression in NK Cells and ILC1s. *Cell Rep.* **22,** 3385–3392 (2018).

55. Martinet, L. *et al.* DNAM-1 expression marks an alternative program of NK cell maturation.

Cell Rep. (2015). doi:10.1016/j.celrep.2015.03.006

56. Wagner, A. K. *et al.* Expression of CD226 is associated to but not required for NK cell

education. *Nat. Commun.* (2017). doi:10.1038/ncomms15627

57. Wright, G. J. *et al.* Characterization of the CD200 Receptor Family in Mice and Humans and

Their Interactions with CD200. *J. Immunol.* (2003). doi:10.4049/jimmunol.171.6.3034

58. Kim, M. *et al.* VEGF-A regulated by progesterone governs uterine angiogenesis and vascular

remodelling during pregnancy. *EMBO Mol. Med.* (2013). doi:10.1002/emmm.201302618

59. Sharma, S., Godbole, G. & Modi, D. Decidual Control of Trophoblast Invasion. *American*

Journal of Reproductive Immunology (2016). doi:10.1111/aji.12466

60. Lash, G. E. *et al.* Decidual macrophages: key regulators of vascular remodeling in human

pregnancy. *J. Leukoc. Biol.* (2016). doi:10.1189/jlb.1A0815-351R

61. Andraweera, P. H., Dekker, G. A. & Roberts, C. T. The vascular endothelial growth factor

family in adverse pregnancy outcomes. *Human Reproduction Update* (2012).

doi:10.1093/humupd/dms011

62.   Athanassiades, A., Hamilton, G. S. & Lala, P. K. Vascular endothelial growth factor stimulates

proliferation but not migration or invasiveness in human extravillous trophoblast. *Biol*

Reprod (1998). doi:10.1095/biolreprod59.3.643

63.   Zhou, Y. *et al.* Vascular endothelial growth factor ligands and receptors that regulate

human cytotrophoblast survival are dysregulated in severe preeclampsia and hemolysis,

elevated liver enzymes, and low platelets syndrome. *Am. J. Pathol.* (2002).

doi:10.1016/S0002-9440(10)62567-9

64.   Kobayashi, T. & Mattarollo, S. R. Natural killer cell metabolism. *Molecular Immunology*

(2017). doi:10.1016/j.molimm.2017.11.021

65.   Keppel, M. P., Saucier, N., Mah, A. Y., Vogel, T. P. & Cooper, M. A. Activation-Specific

Metabolic Requirements for NK Cell IFN-γ Production. *J. Immunol.* (2015).

doi:10.4049/jimmunol.1402099

66.   Michelet, X. *et al.* Metabolic reprogramming of natural killer cells in obesity limits

antitumor responses. *Nat. Immunol.* (2018). doi:10.1038/s41590-018-0251-7

67.   Carvalho, B. S. & Irizarry, R. A. A framework for oligonucleotide microarray preprocessing.

Bioinformatics (2010). doi:10.1093/bioinformatics/btq431

68.   Klaus, B. An end to end workflow for differential gene expression using Affymetrix

microarrays. *F1000Research* (2016). doi:10.12688/f1000research.8967.1

69.   Smyth, G. K. Linear Models and Empirical Bayes Methods for Assessing Differential

Expression in Microarray Experiments. *Stat. Appl. Genet. Mol. Biol.* (2004).

doi:10.2202/1544-6115.1027

70.   Futatsugi, K. *et al.* Obesity-induced kidney injury is attenuated by amelioration of aberrant

PHD2 activation in proximal tubules. *Sci. Rep.* **6,** (2016).

71.   Demarta-Gatsi, C. *et al.* Immunological memory to blood-stage malaria infection is

controlled by the histamine releasing factor (HRF) of the parasite. *Sci. Rep.* **7,** (2017).

FIGURE LEGENDS

Figure 1. Establishment of a mouse model of diet-induced maternal obesity. A) Schematic diagram

depicting the experimental workflow with low-fat, no sucrose (LFD) or high-fat, high sucrose (HFD)

diet regime, breeding strategy, pregnancy detection at gestational day (Gd) 0.5, and gestational

endpoints (Gd 7.5/10.5/14.5). Shown are uterine horns with multiple conception sites comprised of

utero-placental tissues that were allocated for downstream cellular analyses or histology. **B)** Body

weight gain prior to pregnancy of mice fed either HFD (black bars; n=20) or a matched LFD (grey

bars; n=20). Data represented as mean ± SEM. *p<0.05; **p<0.005. **C)** Body weight distribution of

LFD- and HFD-exposed mice following the 13-week diet regime. Red dashed line represents the mean

LFD weight + 3 SD that was used to subcategorize HFD mice into diet-induced obese (DIO) and diet-

induced obese resistant (DIO-R) groups. **D)** Average weekly weight gain rate (LFD: n=14, HFD:

n=19). Data represented as mean ± SD. *P*-values from Student's t-test. **E)** Maternal serum levels of

TNF-α at Gd10.5 in LFD (n=13), HFD (n=16), DIO (n=10), and DIO-R (n=6) mice. Data represented

as mean ± SD. *P*-values from Mann-Whitney test. **F)** Proportion of resorbed implantation sites per

mouse observed at Gd14.5 (LFD: n=13, HFD: n=13); Data represented as mean ± SD. *P*-values from

Student's t-test.

Figure 2. HFD exposure does not result in alterations of total uterine leukocytes or subsets of

uNK cells. A) Representative DBA immunofluorescence images of implantation sites at Gd7.5 (LFD:

n=9; HFD: n=13) and Gd10.5 (LFD: n=10; HFD: n=9) and scatter plots showing number of dolichos

biflorus agglutinin (DBA)+ uterine NK cells. Scale bar = 200μm. **B)** Flow cytometry gating strategy

identification and quantification of uterine immune cell populations, including total uNK (CD3e-

/CD45+/CD122+), tissue-resident (tr) (CD122+/CD49a+), conventional (c) (CD122+/DX5+), and uNK

precursors (CD122+/DX5-/NK1.1-). Representative scatter plots of **C)** total leukocyte (CD45+), **D)**

uterine NK, **E)** tr-uNK, **F)** c-uNK, and **G)** uNK precursors in LFD (n=14), HFD (n=18), DIO (n=11),

and DIO-R (n=7) mice. Data represented as mean ± SD; *P*-values from unpaired Student's t-test for

panels A, C-G.

1 **Figure 3. HFD and diet-induced weight gain alter frequencies of NCR1+ uNK cells. A)**

2 Representative flow cytometry plots of NCR1+ uNK proportions. Scatter plots show **B)** proportion of

3 NCR1+ uNK and **C)** NCR1 median fluorescent intensities (MFI) readings within uNK of LFD (n=14)

4 HFD (n=18), DIO (n=11), and DIO-R (n=7) mice.

Figure 4. High-fat diet and diet-induced weight-gain potentiates uNK activity. A) Representative

flow cytometry plots and gating strategy for analyzing proportions of CD69-expressing uNK and

NCR1+ uNK. **B)** Representative MFI histograms and scatter plots showing CD69 expression intensity

(MFI) within uNK. (LFD: n=14; HFD: n=18; DIO: n=11; DIO-R: n=7). **C)** Representative MFI

histograms and scatter plots showing expression levels of CD69 within NCR1+ uNK of LFD (n=14),

HFD (n=18), DIO (n=11), and DIO-R (n=7) mice. Data represented as mean ± SD; *P*-values from

unpaired Student's t-tests for panels B&C.

Figure 5. Exposure to HFD does not affect placental structure. A) Representative images of H&E-
stained placentas from HFD- and LFD-fed mice with outlined Labyrinth (L), Junctional zone (Jz) and
decidua (D). Scatter plots to the right show total placental area (mm^2) from LFD, HFD, DIO and DIO-
R groups. Scale bar = 1000μm. Scatter plots showing proportional measurements of **B)** labyrinth, **C)**
junctional zone and **D)** decidual areas relative to total placental area in LFD (n=28), HFD, (n=24), DIO
(n=16), and DIO-R (n=8) mice. Data represented as mean ± SD; *P*-values from unpaired Student's t-
tests for panels A-D.

Figure 6. HFD leads to impaired spiral artery remodeling in mid pregnancy. A) Images of H&E

stained Gd10.5 implantation sites. Hashed boxes and outlines show locations of spiral arteries being

measured and outline vessel wall and vessel lumen areas. **B)** Arterial wall area (μm^2) to lumen area

(μm^2) [wall:lumen] ratios from Gd10.5 LFD (n=11), HFD (n=9), DIO (n=4), and DIO-R (n=5) mice.

C) Representative immunofluorescence images of Gd7.5 and Gd10.5 implantation sites stained with

antibodies directed against smooth muscle actin (SMA; purple), and DBA (green). Nuclei are stained

with DAPI (white). SMA surrounding arteries is highlighted with yellow dashed-lines, and artery

lumens are indicated with blue-dashed lines. Scale bar = 40μm. **D)** Arterial SMA layer to arterial

lumen ratios from Gd7.5 LFD (n=13) and HFD (n=15) mice, and in Gd10.5 LFD (n=12) and HFD

(n=19) mice **E)** H&E images of GD14.5 decidual arteries in LFD (n=16), HFD (n=17), DIO (n=11),

and DIO-R (n=6) mice. Scale bar = 100μm. Data represented as mean ± SD; *P*-values from unpaired

Student's t-tests for panels B, D&E.

Figure 7. uNK gene expression profile of DIO and DIO-R vs LFD at Gd10.5. A) Schematic diagram depicting the workflow for gene expression analysis. uNK RNA was isolated from Gd10.5 implantation sites and hybridized onto Affymetrix microarray chip followed by analysis in R. **B)** Hierarchical clustering of LFD, DIO, and DIO-R uNK using a gene signature derived from nominal *P*-value ranking (*P*<0.05, fold-change > 1.25). Boxplots showing log-transformed, normalized expression levels for selected genes involved in **C)** immune function, **D)** vascular remodeling/angiogenesis and **E)** metabolism.

SUPPLEMENTAL FIGURE LEGENDS

Supplemental Figure 1. Scatter plots of **A)** placental and **B)** fetal weights at Gd14.5. (LFD: n=17;

HFD: n=16; DIO: n=10; DIO-R: n=6). **C)** DBA+ cell diameter measurements (µm) and

photomicrographs of DBA (DBA: red, DAPI/nuclei: blue) immunofluorescence staining showing the

approach for cell diameter measurements (yellow dashed lines). Scale bar = 10µm. (LFD: n=9 i.s.;

HFD: n=9 i.s.; DIO: n=4 i.s.; DIO-R: n=5 i.s.). Data represented as mean ± SD; *P*-values from unpaired

Student's t-test for panels A-C.

Supplemental Figure 2. uNK gene expression analysis. A) Principal component analysis (PCA) of

raw data, normalized data, filtered data and ComBat batch corrected data. **B)** Volcano plots following

differential gene expression analysis of HFD vs LFD, DIO vs LFD and DIO-R vs LFD groups. **C)** PCA

clustering based on global gene signatures in LFD, DIO and DIO-R groups. **D)** Scatter plot of relative

mRNA expression of *Vegfa* in Gd10.5 uNK (HFD: n=5; LFD: n=5). Data represented as mean ± SD;

P-values from unpaired Student's t-test.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A

HFD   LFD   Total area

B Labyrinth   C Junctional Zone   D Decidua

Figure 6

A

200μm

40μm

1000μm

LFD

200μm

40μm

1000μm

HFD

B

Gd10.5

Wall:lumen ratio

p = 0.01

LFD HFD

p = 0.04

p = 0.01

LFD DIO DIO-R

HFD

C

DAPI DBA SMA

Gd7.5

Gd10.5

LFD

HFD

D

SMA:Perimeter ratio

p = 0.02

p = 0.003

Gd 7.5 10.5 7.5 10.5

LFD HFD

E

LFD

HFD

Gd14.5

Wall:lumen ratio

LFD HFD

p = 0.08

LFD DIO DIO-R

HFD

Figure 7

A

Gd10.5   Fluorescence Activated Cell Sorting   RNA extraction   Microarray hybridization

B

- LFD
- DIO
- DIO-R

C   Cell-mediated immunity

D   Vascular biology

E   Metabolism