In vitro transcribed guide RNAs trigger an innate immune response via the RIG-I pathway

Short title: *In vitro* transcribed guide RNAs trigger an innate immune response

Beeke Wienert[1,2], Jiyung Shin[1,2], Elena Zelin[1,2], Kathleen Pestal[2] and Jacob E. Corn[1,2,#]

[1] Innovative Genomics Initiative, University of California, Berkeley, Berkeley, USA

[2] Department of Molecular and Cell Biology, University of California, Berkeley, Berkeley, USA

[#] Corresponding author: jcorn@berkeley.edu

1 Abstract

CRISPR-Cas9 genome editing is revolutionizing fundamental research and has great potential for

the treatment of many diseases. While editing of immortalized cell lines has become relatively

easy, editing of therapeutically relevant primary cells and tissues can remain challenging. One

recent advancement is the delivery of a Cas9 protein and an *in vitro* transcribed (IVT) guide RNA

(gRNA) as a precomplexed ribonucleoprotein (RNP). This approach allows editing of primary

cells such as T cells and hematopoietic stem cells, but the consequences beyond genome editing

of introducing foreign Cas9 RNPs into mammalian cells are not fully understood. Here we show

that the IVT gRNAs commonly used by many laboratories for RNP editing trigger a potent innate

immune response that can be several thousand times stronger than benchmark immune stimulating

ligands. IVT gRNAs are recognized in the cytosol through the RIG-I pathway but not the MDA5

pathway, thereby triggering a type I interferon response. Removal of the 5'-triphosphate from

gRNAs ameliorates inflammatory signaling and prevents the loss of viability associated with

genome editing in hematopoietic stem cells. The potential for Cas9 RNP editing to induce a potent

antiviral response indicates that care must be taken when designing therapeutic strategies to edit

primary cells.

Abbreviations:

Cas - CRISPR-associated

CIP - calf intestinal alkaline phosphatase

CRISPR - clustered, regularly interspaced, short palindromic repeat

dCas9 – nuclease-dead Cas9

HEK293 - Human embryonic kidney cells 293

HEK293T - Human embryonic kidney cells 293 SV40 large T antigen

HeLa – Henrietta Lacks cells

HSPCs – CD34$^+$ human hematopoietic stem and progenitor cells

IFNAR1 - Interferon Alpha And Beta Receptor Subunit 1

IFNβ/IFNB1 - Interferon beta

ISG15 - Interferon-stimulated gene 15

IVT – *in vitro* transcribed

KO - knockout

MAVS - mitochondrial activator of virus signaling

MDA5/IFIH1 - melanoma differentiation-associated gene 5/ Interferon Induced with Helicase C

Domain 1

PAMP - pathogen-associated molecular pattern

RIG-I/DDX58 - retinoic acid-inducible gene I/ DExD-H-box helicase 58

gRNA –guide RNA

SPRI - solid phase reversible immobilization

WT – wild type

Introduction

CRISPR (clustered, regularly interspaced, short palindromic repeat)-Cas (CRISPR-associated) genome editing has rapidly become a widely used tool in molecular biology laboratories. Its ease of use and high flexibility allows researchers to modify and edit genomes in cell lines (1), stem cells (2), animals and plants (3,4), and even human embryos (5). At least two components must be successfully delivered into cells during genome editing: the Cas protein, such as Cas9, and a guide RNA (gRNA) to direct the Cas protein to its target site. For *in vitro* cultured cells this can be done by transfecting plasmids encoding gRNA and Cas9 protein. However, transfection of plasmid DNA into sensitive cell types such as primary and stem cells is challenging and inefficient. The introduction of plasmids can also lead to undesired integration of DNA at the cut site (6), increased off-target activity through prolonged expression of the CRISPR-Cas9 components (7), and a delay in editing while the cell expresses gRNA and Cas protein (8).

The delivery of gRNA and Cas9 protein as a pre-complexed ribonucleoprotein (RNP) sidesteps issues related to plasmid expression and has proved to be a successful strategy to edit human primary cells, including T cells (9,10), hematopoietic stem cells (11−14), and neurons (15). This makes RNP editing a particularly attractive approach for therapeutic applications, but relatively little is known about the non-editing consequences of introducing a foreign gRNA and Cas9 protein. Human cells have evolved multiple defense mechanisms to guard against foreign components, and genome editing reagents have the potential to activate these systems. For example, recent data suggests that humans may have a pre-existing adaptive immune response to the Cas9 protein (16). But cellular responses to the gRNAs used to program Cas9 editing have so far not been well explored.

Cells respond to infection by RNA-viruses with an innate immune response that protects the host

cell from invading foreign genetic material (17). Foreign RNAs are recognized by pathogen-

associated molecular pattern (PAMP) binding receptors in the cytosol that include retinoic acid-

inducible gene I (RIG-I) and melanoma differentiation-associated gene 5 (MDA5) (18). This

triggers a cascade of events mediated by the mitochondrial activator of virus signaling (MAVS)

protein resulting in the transcriptional activation of type I interferons and interferon-stimulated

genes (ISGs) (19–21).

PAMPs usually contain exposed 5'-triphosphate ends (18), which may also be present in gRNAs

made via T7 *in vitro* transcription (IVT) (22,23). We asked whether IVT gRNAs cause an innate

immune response, and here show that RNP genome editing induces upregulation of interferon beta

(IFNβ) and interferon-stimulated gene 15 (ISG15) in a variety of human cell types. This activity

depends upon RIG-I and MAVS, but is independent of MDA5. The extent of the immune response

depends upon the protospacer sequence, but removal of the 5'-triphosphate from gRNAs avoids

stimulation of innate immune signaling. The potential for Cas9 RNP editing to induce an antiviral

response indicates that care must be taken when designing therapeutic strategies to edit primary

cells.

Results

To investigate if mammalian cells react to IVT gRNA/Cas9 with an innate immune response, we

first performed genome editing in human embryonic kidney 293 (HEK293) cells using Cas9 RNPs.

To separate innate immune response from genome editing, we performed these experiments with

a non-targeting gRNA that recognizes a sequence within BFP and has no known targets within the

human genome (24). Constant amounts of recombinant Cas9 protein were complexed with

different amounts of non-targeting IVT gRNA and RNPs were transfected into HEK293 cells using

CRISPRMAX lipofection reagent (25). We harvested cells 30h after transfection and measured

transcript levels of *IFNB1* and *ISG15* by qRT-PCR (**Figure 1A**). Introduction of gRNAs caused a

dramatic increase in both *IFNB1* and *ISG15* levels, and the presence of Cas9 protein did not have

an effect on the outcome. Cas9 on its own did not induce *IFNB1* or *ISG15* expression. To our

surprise, as little as 1 pmol of gRNA was sufficient to trigger a 30-50-fold increase in the

transcription of innate immune genes. We further found that the commonly-administered 100 pmol

of a gRNA can induce *IFNB1* by 4,000 fold, which is equal to or even stronger than induction by

canonical IFNβ inducers such as viral mRNA from Sendai Virus (26) or a Hepatitis C virus (HCV)

PAMP (20,27) (**Figure 1B**).

RNPs can be delivered into cells via different transfection methods and while lipofection is cost-

effective and easy to use, many researchers prefer electroporation for harder-to-transfect cells. We

wondered if the transfection method would affect the IFNβ response and compared gRNA

transfection via lipofection (Lipofectamine 2000 and RNAiMAX) to nucleofection (Lonza)

(**Figure 1C**). Lipofection led to a strong increase in *IFNB1* and *ISG15* transcript levels after as

little as 6h post transfection, and the response was sustained for up to 48 hours. Nucleofection also

caused an increase in innate immune signaling at early time points, but the response was much

milder than in lipofected samples and was greatly diminished by 48 hours.

Next, we asked if the innate immune response to gRNAs is a common phenomenon across different

cell types and compared IFNβ activation in seven commonly used human cell lines of various

lineages: HEK293T, HEK293, HeLa, Jurkat, HCT116, HepG2 and K562 (**Figure 2A**). While the

magnitude of induction varied between cell lines, all tested cell lines responded to IVT gRNA

transfection with activation of *IFNB1* expression. The sole exception was K562 cells, which have

a homozygous deletion of the *IFNA* and *IFNB1* genes (28). We also measured transcript levels of

two major cytosolic pathogen recognition receptors, RIG-I (*DDX58*) and MDA5 (*IFIH1*), and

noticed that all cell lines except K562 upregulated these transcripts in response to introduction of

gRNAs. We also confirmed these results on the protein level in HEK293 cells (**Figure 2B**).

The RIG-I and MDA5 receptors complement each other by recognizing different structures in

foreign cytosolic RNAs, but the exact nature of their ligands is not yet fully understood (29,30).

To investigate if IVT gRNAs are recognized via RIG-I or MDA5, we generated clonal knockout

(KO) cell lines for RIG-I, MDA5, and their downstream interaction partner MAVS in HEK293

cells using CRISPR-Cas9. As the expression of both RIG-I and MDA5 are themselves stimulated

by IFNβ, we confirmed successful KO after transfection with gRNAs by genomic PCR, Sanger

sequencing, and Western Blot (**Supplementary Figure 1A-C**). MAVS KO cells were confirmed

by Western Blot (**Supplementary Figure 1D**). Strikingly, activation of *IFNB1 e*xpression after

introduction of gRNAs was absent in RIG-I and MAVS KO cells, while MDA5 KO cells did not

show a significant decrease in *IFNB1* transcript levels (**Figure 2C**). This indicates that IVT gRNAs

are exclusively recognized through RIG-I to trigger a type I interferon response.

As the structural requirements of RIG-I ligands are still not completely understood, we wondered

if different 20 nucleotide protospacers in gRNAs vary in their potency to trigger an innate immune

response via RIG-I. We designed 10 additional non-targeting gRNAs that we *in vitro* transcribed

and transfected into HEK293 cells. Surprisingly, we found that the cells responded to different

protospacers with a wide range of differential *IFNB1* expression. Several gRNAs produced very

little innate immune response, and one gRNA (gRNA11) yielded no *IFNB1* activation at all

(**Figure 3A**). We speculated that the differential response may be correlated with the purity of the

RNA product after IVT or the stability of the secondary structure of the RNA (31). However, we

found that there was no obvious correlation between the immune response to certain gRNAs and

their purity, predicted protospacer secondary structure, full secondary structure including the

constant region, or predicted disruption of the constant region by mis-pairing with the protospacer

(**Supplementary Figure 2**). These results indicate that RIG-I recognition patterns of IVT gRNAs

are complex and difficult to predict *a priori* based on predicted properties of the variable

protospacer.

One well-established requirement of RIG-I ligands is the presence of a 5'-triphosphate group (32).

We asked if preparations that remove the 5' triphosphate might avoid or reduce the innate immune

response to IVT gRNAs. We first used a synthetic gRNA that lacks a 5'-triphosphate and verified

that this gRNA does not induce *IFNB1* expression when transfected into HEK293 cells (**Figure

3B**). Synthetic guide RNAs are becoming more commonplace, but are still an order of magnitude

more expensive than IVT of gRNAs. This limits their application for high-throughput interrogation

of gene function in primary cells. We therefore asked if treatment of IVT gRNA with calf intestinal

alkaline phosphatase (CIP) to remove the 5'-triphosphate would reduce *IFNB1* induction. We

found that CIP treatment significantly reduced the *IFNB1* response (**Figure 3B**). Using different

CIP treatment regimes, we further found that removal of the 5'-triphosphate must to be absolutely

complete and should be carried out rigorously to avoid IFN stimulation (**Supplementary Figure

A-B). We also compared purification of IVT gRNAs by solid phase reversible immobilization

(SPRI) beads to column purification and established that SPRI bead clean-up is not sufficient to

completely avoid an immune response even when more CIP is used (**Figure 3B and**

Supplementary Figure 3C). Taken together, these results indicate that 5'-triphosphate is a

necessary requirement for gRNA-induced *IFNB1* activation through RIG-I, but that additional

structural properties of the gRNAs also influence the magnitude of the immune response.

When a cell initiates an antiviral immune response, it also undergoes cellular stress that can affect

cell viability (33,34). Hence, we asked if there is a correlation between the IFNβ response and cell

viability after transfection with synthetic, IVT or CIP-treated IVT gRNA. Not surprisingly, the

viability of the very robust HEK293 cell line was not affected by the antiviral immune response

(**Supplementary Figure 3D**). We then turned to primary CD34$^+$ human hematopoietic stem and

progenitor cells (HSPCs), which are a much more sensitive cell type. We first nucleofected HSPCs

with RNPs targeting the *HBB* gene (11) and compared synthetic and IVT gRNA interferon

stimulation and cell viability post transfection. Double-strand breaks have been reported to cause

innate immune stimulation and can themselves cause decreases in cell fitness (35,36). Therefore,

we performed controls using nuclease-dead Cas9 (dCas9) to form RNPs. We saw a significant

decrease in cell viability of HSPCs in both IVT gRNA RNP samples which was associated with

an increase in IFN stimulated genes *ISG15* and *RIG-I* (**Figure 3C-D**). We did not see a substantial

difference in viability or ISG expression between Cas9 and dCas9 RNPs suggesting that nuclease

activity did not affect the immune response. Next, we asked if CIP treatment of gRNAs could

reverse the decrease in viability in HSPCs. We nucleofected HSPCs with dCas9 RNPs targeting a

non-coding intron of *JAK2* and compared synthetic, IVT and CIP-treated IVT gRNAs. Strikingly,

CIP treatment could completely restore the viability in HSPCs (**Figure 3E**).

Discussion

We have found that IVT gRNAs used with Cas9 RNPs for many genome editing experiments can

trigger a strong innate immune response in many mammalian cell types (**Figure 4**). Lipofection

results in a stronger and longer lasting response than nucleofection, possibly because lipofection

delivers gRNAs to the cytosol while nucleofection delivers mainly to the nucleus. Using isogenic

knockout clones, we found the gRNA-induced response is mediated via the anti-viral RIG-I

pathway and results in expression of genes that initiate an antiviral immune response. While

introduction of IFN-stimulating gRNAs does not affect viability in HEK293 cells, we found that

viability of primary HSPCs is negatively affected by the antiviral immune response.

These results have several implications. We suggest that the gene signature associated with type I

interferon stimulation should be considered when studying the transcriptome of recently edited

bulk populations of cells. Furthermore, all mammalian cells can both produce type I interferons

and also respond to them through the ubiquitously expressed receptor Interferon Alpha And Beta

Receptor Subunit 1 (IFNAR1) (37). Even cells that have not been successfully transfected with

RNPs could sense the INFβ produced by neighboring cells and activate downstream antiviral

defense mechanisms. This could be an important consideration during *in vivo* genome editing

applications as RNP delivery into one set of cells could provoke a wide-spread innate immune

response in the surrounding tissues.

We found that synthetic gRNAs completely circumvent the RIG-I mediated response, offering a

valuable path to avoid innate immune signaling during therapeutic editing. However, synthetic

gRNAs can become expensive when performing experiments that require testing or using many

gRNAs. We found that a cost-effective CIP treatment to remove the 5'-triphosphate before

transfection reduces the immune response and increases post-transfection viability in HSPCs.

Complete removal of the 5'-triphosphate is essential to restore viability in HSPCs to the same level

as observed with synthetic gRNAs. Thus, consideration of a potential innate immune stimulation

prior to choice of genome editing reagents, study design, and implementation of controls is critical

when performing genome editing using RNPs in mammalian cells.

Double-strand breaks have on their own been reported to induce an innate immune response (35).

However, we performed controls comparing WT Cas9 to a nuclease-dead Cas9 thereby showing

that the gRNA-mediated innate immune response and associated cell death in HSPCs is not caused

by double-strand breaks.

While we were preparing this manuscript for submission, the Kim group reported similar results

in HeLa cells and primary human CD4$^+$ T cells (38). They confirmed that the type I interferon

response is dependent on the presence of a 5'-triphopsphate group and that CIP treatment can

increase viability by avoiding the antiviral response. These results are very much in alignment with

our findings and extend the potential problem of innate immune signaling to additional cell types.

Our study adds extra depth by further outlining the mechanisms by which gRNAs are sensed. We

show that gRNA sensing depends upon RIG-I and MAVS, but MDA5 knockout cells are fully

capable of inducing IFNβ after IVT gRNA transfection. Hence, gRNA sensing is independent of

the MDA5 PAMP receptor. Furthermore, we show that in addition to a 5'-triphosphate, the

protospacer sequence is also critical to determine the intensity of the IFNβ response. Not only do

different gRNAs induce different innate immune responses, some gRNAs induce no response at

all. It has been proposed that 5'-basepaired RNA structures are required to activate antiviral

signaling via RIG-I, but we found no correlation between signaling and a variety of predicted RNA

properties, including secondary structure (31). Our results therefore suggest that the mechanism of

gRNA sensing by the RIG-I pathway is relatively complex, in that it requires 5' triphosphates but

that this moiety is not sufficient to induce the response. Future work to delineate the full set of

molecular features responsible for gRNA activation of innate immunity might yield accurate

predictors of innate immune signaling in general.

Materials and methods

In vitro transcription of gRNAs

gRNA was synthesized by assembly PCR and *in vitro* transcription as previously described (11).

Briefly, a T7 RNA polymerase substrate template was assembled by PCR from a variable 58-59

nt primer containing T7 promoter, variable gRNA guide sequence, and the first 15 nt of the non-

variable region of the gRNA (T7FwdVar primers, 10 nM, Supplementary Table 1; Supplementary

Table 2 for gRNA sequences), and an 83 nt primer containing the reverse complement of the

invariant region of the gRNA (T7RevLong, 10 nM), along with amplification primers

(T7FwdAmp, T7RevAmp, 200 nM each). The two long primers anneal in the first cycle of PCR

and are then amplified in subsequent cycles. Phusion high-fidelity DNA polymerase was used for

assembly (New England Biolabs). Assembled template was used without purification as a substrate

for *in vitro* transcription by T7 RNA polymerase using the HiScribe T7 High Yield RNA Synthesis

kit (New England Biolabs) following the manufacturer's instructions. Resulting transcription

reactions were treated with DNAse I (New England Biolabs), and RNA was purified by treatment

with a 5X volume of homemade solid phase reversible immobilization (SPRI) beads (comparable

to Beckman-Coulter AMPure beads) and elution in RNAse-free water.

<u>CIP treatment of IVT gRNAs</u>

When gRNAs were treated with Alkaline Calf intestinal phosphatase (CIP) (New England

Biolabs), 20U of CIP (2 μl) were added per 20 μl IVT reaction and samples were incubated at 37C

for 3h before proceeding to purification and DNAseI treatment. CIP-treated samples and

corresponding no CIP IVT gRNA controls were purified using a Qiagen RNeasy Mini Kit

(Qiagen). The detailed protocol and additional notes are available online

<u>(dx.doi.org/10.17504/protocols.io.nghdbt6).</u>

<i>In vitro</i> transcription of HCV PAMP and Sendai Virus DI RNA

HCV PAMP IVT template (20) was generated by annealing HCV fwd and rev (5 μM each) oligos

(Supplementary Table 1). 2 μl of the annealed product was used as DNA template in the

subsequent IVT reaction using HiScribe T7 High Yield RNA Synthesis kit (New England

Biolabs).

The plasmid containing the SeV DI RNA(26) was a gift from Prof. Peter Palese, Icahn School of

Medicine at Mount Sinai, New York. Plasmid was digested with HindII/EcoRI before IVT with

HiScribe T7 High Yield RNA Synthesis kit (New England Biolabs). The sequence of the IVT DI,

including the T7 promoter, hepatitis delta virus ribozyme, and the T7 terminator, is:

TAATACGACTCACTATA**ACCAGACAAGAGTTTAAGAGATATGTATCCTTTTAAAT**

TTTCTTGTCTTCTTGTAAGTTTTTCTTACTATTGTCATATGGATAAGTCCAAGAC

TTCCAGGTACCGCGGAGCTTCGATCGTTCTGCACGATAGGGACTAATTATTACG

AGCTGTCATATGGCTCGATATCACCCAGTGATCCATCATCAATCACGGTCGTGT

ATTCATTTTGCCTGGCCCCGAACATCTTGACTGCCCCTAAAATCTTCATCAAAA

TCTTTATTTCTTTGGTGAGGAATCTATACGTTATCTATGTATAATATCCTCAAA

CCTGTCTAATAAAGTTTTTGTGATAACCCTCAGGTTCCTGATTTCACGGGATGA

TAATGAAACTATTCCCAATTGAAGTCTTGCTTCAAACTTCTGGTCAGGGAATGA

CCCAGTTACCAATCTTGTGGACATAGATAAAGATAGTCTTGGACTTATCCATAT

GACAATAGTAAGAAAAACTTACAAGAAGACAAGAAAATTTAAAAGGATACATAT

CTCTTAAACTCTTGTCTGGTGGCCGGCATGGTCCCAGCCTCCTCGCTGGCGCCGGC

TGGGCAACATTCCGAGGGGACCGTCCCCTCGGTAATGGCGAATAGCATAACCCCTT

GGGGCCTCTAAACGGGTCTTGAGGGGTTTTTTG.

The sequence of the SeV DI is highlighted in boldface.

Both, HCV PAMP and SeV DI RNA were purified by treatment with a 5X volume of homemade

solid phase reversible immobilization (SPRI) beads (comparable to Beckman-Coulter AMPure

beads) and elution in RNAse-free water.

<u>Synthetic gRNAs</u>

Chemically synthesized gRNAs, which were purified using high-performance liquid

chromatography (HPLC), were purchased from Synthego.

<u>RNA quality control</u>

IVT gRNAs were analyzed using a Bioanalyzer. This was performed by the UC Berkeley

Functional Genomics Laboratory (FGL) core facility. gRNAs were denatured for 5 mins at 70C

before analysis on bioanalyzer.

<u>Cas9 protein preparation</u>

The Cas9 construct (pMJ915) contained an N-terminal hexahistidine-maltose binding protein

(His6-MBP) tag, followed by a peptide sequence containing a tobacco etch virus (TEV) protease

cleavage site. The protein was expressed in E. coli strain BL21 Rosetta 2 (DE3) (EMD

Biosciences), grown in TB medium at 16°C for 16h following induction with 0.5 mM IPTG. The

Cas9 protein was purified by a combination of affinity, ion exchange and size exclusion

chromatographic steps. Briefly, cells were lysed in 20 mM HEPES pH 7.5, 1M KCl, 10mM

imidazole, 1 mM TCEP, 10% glycerol (supplemented with protease inhibitor cocktail (Roche)) in

a homogenizer (Avestin). Clarified lysate was bound to Ni-NTA agarose (Qiagen). The resin was

washed extensively with lysis buffer and the bound protein was eluted in 20 mM HEPES pH 7.5,

100mM KCl, 300mM imidazole,1 mM TCEP 10% glycerol. The His6-MBP affinity tag was

removed by cleavage with TEV protease, while the protein was dialyzed overnight against 20 mM

HEPES pH 7.5, 300 mM KCl, 1 mM TCEP, 10% glycerol. The cleaved Cas9 protein was separated

from the fusion tag by purification on a 5 ml SP Sepharose HiTrap column (GE Life Sciences),

eluting with a linear gradient of 100 mM – 1 M KCl. The protein was further purified by size

exclusion chromatography on a Superdex 200 16/60 column in 20 mM HEPES pH 7.5, 150 mM

KCl and 1 mM TCEP. Eluted protein was concentrated to 40uM, flash-frozen in liquid nitrogen

and stored at -80°C.

<u>Culture and transfection of immortalized cell lines</u>

Cells were obtained from ATCC and verified mycoplasma-free (Mycoalert LT-07, Lonza).

HEK293, HEK293T, HCT116, HepG2 and HeLa cells were maintained in DMEM supplemented

with 10 % FBS and 100 μg/mL Penicillin-Streptomycin (all Gibco). K562 and Jurkat cells were

maintained in RPMI supplemented with 10% FBS and 100 μg/mL Penicillin-Streptomycin.

All transfections in cell lines were performed in 12-well cell culture dishes using $2x10^5$ cells per

transfection. For lipofection we used Lipofectamine™ CRISPRMAX™-Cas9, Lipofectamine®

RNAiMAX or Lipofectamine® 2000 Transfection Reagent (all Invitrogen) in reverse-

transfections according to the manufacturer's protocols. Unless stated otherwise, $2x10^5$ cells were

transfected with 50 pmol of RNA and harvested 24-30h post transfection for RNA extraction.

<u>Culture and transfection of primary HSPCs</u>

Human CD34+ HSPCs from mobilized peripheral blood (Allcells, Inc) were thawed and cultured

in StemSpan SFEM medium (StemCell Technologies) supplemented with StemSpan CC110

cocktail (StemCell Technologies) for 48 hours before nucleofection with dCas9 RNP (50pmol of

dCas9, 50 pmol of gRNA). $1.5x10^5$ HSPCs were pelleted (100 x g, 10 min) and resuspended in 20

μl Lonza P3 solution, mixed with 10ul dCas9 RNP, and nucleofected using ER100 protocol in

Lonza 4D nucleofector. Viability of the cells was measured 24 hours post nucleofection using

Trypan blue exclusion test. RNA was harvested 16 hours post nucleofection.

<u>RNA extraction, cDNA synthesis and qRT-PCR</u>

Cell cultures were washed with PBS prior to RNA extraction. Total RNA was extracted using

RNeasy Miniprep columns (Qiagen) according to the manufacturer's instructions including the

on-column DNAseI treatment (Qiagen). 1 µg of total RNA was used for subsequent cDNA

synthesis using iScript™ Reverse Transcription Supermix (Biorad). For qRT-PCR reactions, a

total of 20 ng of cDNA was used as a template and combined with primers (see Supplementary

Table 3) and SsoFast™ EvaGreen® Supermix (Biorad) and amplicons were generated using

standard PCR amplification protocols for 40 cycles on a StepOnePlus Real-Time PCR system

(Applied Biosystems). Ct values for each target gene were normalized against Ct values obtained

for *GAPDH* to account for differences in loading (ΔCt). To determine 'fold activation' of genes

ΔCt values for target genes were then normalized against ΔCt values for the same target gene for

mock-treated cells (ΔΔCt).

<u>Generation of knockout cell lines</u>

For CRISPR/Cas9 genome editing we used a plasmid encoding both the Cas9 protein and the

gRNA. pSpCas9(BB)-2A-GFP (px458) was a gift from Feng Zhang (Addgene plasmid #48138).

We designed gRNA sequences using the free CRISPR knockout design online tool from Synthego.

Two different gRNA sequences were designed for RIG-I and MDA5, respectively (see

Supplementary Table 3).

2×10^5 HEK293 cells were nucleofected with 2 µg of px458 plasmids containing both targeting

gRNAs in a 1:1 ratio using a Lonza 4D nucleofector (Lonza) with the manufacturer's

recommended settings. After 48h cells were harvested and subjected to fluorescence-activated cell

sorting (FACS). Cells expressing high levels of GFP were single-cell sorted into 96-well plates to

establish clonal populations.

For the screening process, genomic DNA from clonal populations was extracted using

QuickExtract solution (Lucigen). For KO of RIG-I and MDA5 we screened clones by genomic

PCR looking for a PCR product that is significantly smaller in size than that of WT HEK293 cells

(see Supplementary Table 4 for primers). PCR products were then Sanger sequenced by the UC

Berkeley DNA Sequencing facility using the forward primers of the PCR reaction as sequencing

primers.

<u>Western Blot</u>

Cells were harvested and washed with PBS. Cells were lysed in 1xRIPA buffer (EMD Millipore)

for 10 mins on ice. Samples were spun down at 14000xg for 15 mins and protein lysates were

transferred to a new tube. 50 μg of total protein was separated for size by SDS-PAGE and

transferred to a nitrocellulose membrane. Blots were blocked in 4% skim milk in 50 mm Tris-

HCl (pH 7.4), 150 mm NaCl, and 0.05% Tween 20 (TBST) and then probed for RIG-I,

MDA5, MAVS or GAPDH protein using antibodies against RIG-I (D14G6), MDA5 (D74E4),

MAVS (D5A9E) or GAPDH (14C10), respectively (all Cell Signaling Technologies). This was

followed by incubation with secondary antibody IRDye® 800CW Donkey anti-Rabbit IgG (Li-

Cor). Rainbow™ protein standards (GE Healthcare) were loaded in each gel for size estimation.

Blots were visualized using a Li-Cor Odyssey Clx (Li-Cor).

Acknowledgements

We would like to thank Prof. Peter Palese from the Icahn School of Medicine at Mount Sinai, New

York, for the Sendai virus DI IVT template DNA. The CRISPR/Cas9 genome-editing plasmid

px458 was a gift from Feng Zhang, Broad Institute, Cambridge, MA (Addgene plasmid # 48138).

References

References

1. Ran FA, Hsu PD, Wright J, Agarwala V, Scott DA, Zhang F. Genome engineering using

the CRISPR-Cas9 system. Nat Protoc. 2013 Nov;8(11):2281–308.

2. Zhang Z, Zhang Y, Gao F, Han S, Cheah KS, Tse H-F, et al. CRISPR/Cas9 Genome-

Editing System in Human Stem Cells: Current Status and Future Prospects. Mol Ther

Nucleic Acids. 2017 Dec 15;9:230–41.

3. Ceasar SA, Rajan V, Prykhozhij SV, Berman JN, Ignacimuthu S. Insert, remove or replace:

A highly advanced genome editing system using CRISPR/Cas9. Biochim Biophys Acta.

2016 Jun 24;1863(9):2333–44.

4. Bortesi L, Fischer R. The CRISPR/Cas9 system for plant genome editing and beyond.

Biotechnol Adv. 2015 Feb;33(1):41–52.

5. Ma H, Marti-Gutierrez N, Park S-W, Wu J, Lee Y, Suzuki K, et al. Correction of a

pathogenic gene mutation in human embryos. Nature. 2017 Aug 24;548(7668):413–9.

6. Gabriel R, Lombardo A, Arens A, Miller JC, Genovese P, Kaeppel C, et al. An unbiased

genome-wide analysis of zinc-finger nuclease specificity. Nat Biotechnol. 2011 Aug

7;29(9):816–23.

7. Chen Y, Liu X, Zhang Y, Wang H, Ying H, Liu M, et al. A Self-restricted CRISPR System

to Reduce Off-target Effects. Mol Ther. 2016;24(9):1508–10.

8. Kim S, Kim D, Cho SW, Kim J, Kim J-S. Highly efficient RNA-guided genome editing in

human cells via delivery of purified Cas9 ribonucleoproteins. Genome Res. 2014

Jun;24(6):1012–9.

9. Hultquist JF, Schumann K, Woo JM, Manganaro L, McGregor MJ, Doudna J, et al. A Cas9

Ribonucleoprotein Platform for Functional Genetic Studies of HIV-Host Interactions in

Primary Human T Cells. Cell Rep. 2016 Oct 25;17(5):1438–52.

10.   Schumann K, Lin S, Boyer E, Simeonov DR, Subramaniam M, Gate RE, et al. Generation

of knock-in primary human T cells using Cas9 ribonucleoproteins. Proc Natl Acad Sci

USA. 2015 Aug 18;112(33):10437–42.

11.   DeWitt MA, Magis W, Bray NL, Wang T, Berman JR, Urbinati F, et al. Selection-free

genome editing of the sickle mutation in human adult hematopoietic stem/progenitor cells.

Sci Transl Med. 2016 Oct 12;8(360):360ra134.

12.   Hendel A, Bak RO, Clark JT, Kennedy AB, Ryan DE, Roy S, et al. Chemically modified

guide RNAs enhance CRISPR-Cas genome editing in human primary cells. Nat

Biotechnol. 2015 Sep;33(9):985–9.

13.   Gundry MC, Brunetti L, Lin A, Mayle AE, Kitano A, Wagner D, et al. Highly efficient

genome editing of murine and human hematopoietic progenitor cells by crispr/cas9. Cell

Rep. 2016 Oct 25;17(5):1453–61.

14.   Dever DP, Bak RO, Reinisch A, Camarena J, Washington G, Nicolas CE, et al.

CRISPR/Cas9 β-globin gene targeting in human haematopoietic stem cells. Nature. 2016

Nov 17;539(7629):384–9.

15.   Staahl BT, Benekareddy M, Coulon-Bainier C, Banfal AA, Floor SN, Sabo JK, et al.

Efficient genome editing in the mouse brain by local delivery of engineered Cas9

ribonucleoprotein complexes. Nat Biotechnol. 2017 May;35(5):431–4.

16.   Charlesworth CT, Deshpande PS, Dever DP, Dejene B, Gomez-Ospina N, Mantri S, et al.

Identification of Pre-Existing Adaptive Immunity to Cas9 Proteins in Humans. BioRxiv.

2018 Jan 5;

17.   Akira S, Uematsu S, Takeuchi O. Pathogen recognition and innate immunity. Cell. 2006

Feb 24;124(4):783–801.

18.   Kell AM, Gale M. RIG-I in RNA virus recognition. Virology. 2015 May;479–480:110–21.

19.   Goubau D, Schlee M, Deddouche S, Pruijssers AJ, Zillinger T, Goldeck M, et al. Antiviral

immunity via RIG-I-mediated recognition of RNA bearing 5'-diphosphates. Nature. 2014

Oct 16;514(7522):372–5.

20.   Saito T, Owen DM, Jiang F, Marcotrigiano J, Gale M. Innate immunity induced by

composition-dependent RIG-I recognition of hepatitis C virus RNA. Nature. 2008 Jul

24;454(7203):523–7.

21.   Platanias LC. Mechanisms of type-I- and type-II-interferon-mediated signalling. Nat Rev

Immunol. 2005 May;5(5):375–86.

22.   Milligan JF, Groebe DR, Witherell GW, Uhlenbeck OC. Oligoribonucleotide synthesis

using T7 RNA polymerase and synthetic DNA templates. Nucleic Acids Res. 1987 Nov

11;15(21):8783–98.

23.   Milligan JF, Uhlenbeck OC. [5] Synthesis of small RNAs using T7 RNA polymerase.

RNA processing part A: general methods. Elsevier; 1989. p. 51–62.

24.   Richardson CD, Ray GJ, DeWitt MA, Curie GL, Corn JE. Enhancing homology-directed

genome editing by catalytically active and inactive CRISPR-Cas9 using asymmetric donor

DNA. Nat Biotechnol. 2016 Mar;34(3):339–44.

25.   Zuris JA, Thompson DB, Shu Y, Guilinger JP, Bessen JL, Hu JH, et al. Cationic lipid-

mediated delivery of proteins enables efficient protein-based genome editing in vitro and in

vivo. Nat Biotechnol. 2015 Jan;33(1):73–80.

26.   Martínez-Gil L, Goff PH, Hai R, García-Sastre A, Shaw ML, Palese P. A Sendai virus-

derived RNA agonist of RIG-I as a virus vaccine adjuvant. J Virol. 2013 Feb;87(3):1290–

300.

27.   Horner SM, Gale M. Regulation of hepatic innate immunity by hepatitis C virus. Nat Med.

2013 Jul;19(7):879–88.

28.   Diaz MO, Ziemin S, Le Beau MM, Pitha P, Smith SD, Chilcote RR, et al. Homozygous

deletion of the alpha- and beta 1-interferon genes in human leukemia and derived cell lines.

Proc Natl Acad Sci USA. 1988 Jul;85(14):5259–63.

29.   Runge S, Sparrer KMJ, Lässig C, Hembach K, Baum A, García-Sastre A, et al. In vivo

ligands of MDA5 and RIG-I in measles virus-infected cells. PLoS Pathog. 2014 Apr

17;10(4):e1004081.

30.   Yoneyama M, Kikuchi M, Matsumoto K, Imaizumi T, Miyagishi M, Taira K, et al. Shared

and unique functions of the DExD/H-box helicases RIG-I, MDA5, and LGP2 in antiviral

innate immunity. J Immunol. 2005 Sep 1;175(5):2851–8.

31.   Schmidt A, Schwerd T, Hamm W, Hellmuth JC, Cui S, Wenzel M, et al. 5'-triphosphate

RNA requires base-paired structures to activate antiviral signaling via RIG-I. Proc Natl

Acad Sci USA. 2009 Jul 21;106(29):12067–72.

32.   Hornung V, Ellegast J, Kim S, Brzózka K, Jung A, Kato H, et al. 5'-Triphosphate RNA is

the ligand for RIG-I. Science (80-). 2006 Nov 10;314(5801):994–7.

33.   Chawla-Sarkar M, Lindner DJ, Liu YF, Williams BR, Sen GC, Silverman RH, et al.

Apoptosis and interferons: role of interferon-stimulated genes as mediators of apoptosis.

Apoptosis. 2003 Jun;8(3):237–49.

34.   Onomoto K, Yoneyama M, Fung G, Kato H, Fujita T. Antiviral innate immunity and stress

granule responses. Trends Immunol. 2014 Sep;35(9):420–8.

35.   Yu Q, Katlinskaya YV, Carbone CJ, Zhao B, Katlinski KV, Zheng H, et al. DNA-damage-

induced type I interferon promotes senescence and inhibits stem cell function. Cell Rep.

2015 May 5;11(5):785–97.

36. Pietras EM, Lakshminarasimhan R, Techner J-M, Fong S, Flach J, Binnewies M, et al. Re-

entry into quiescence protects hematopoietic stem cells from the killing effect of chronic

exposure to type I interferons. J Exp Med. 2014 Feb 10;211(2):245–62.

37. de Weerd NA, Samarajiwa SA, Hertzog PJ. Type I interferon receptors: biochemistry and

biological functions. J Biol Chem. 2007 Jul 13;282(28):20053–7.

38. Kim S, Koo T, Jee H-G, Cho H-Y, Lee G, Lim D-G, et al. CRISPR RNAs trigger innate

immune responses in human cells. Genome Res. 2018 Feb 22;

39. Lorenz R, Bernhart SH, Höner Zu Siederdissen C, Tafer H, Flamm C, Stadler PF, et al.

ViennaRNA Package 2.0. Algorithms Mol Biol. 2011 Nov 24;6:26.

Figure 1

Figure 1: Transfection of IVT gRNAs into HEK293 cells triggers a type I interferon response.

(**A**) qRT-PCR analysis of *IFNB1* and *ISG15* transcript levels in HEK293 cells transfected with increasing amounts of gRNA with and without Cas9 protein. In the samples with Cas9, gRNAs were complexed with constant amounts (100 pmol) of Cas9 protein. Cells were harvested for RNA extraction 30h after transfection using CRISPRMAX transfection reagent. Ct values were normalized to Ct values of mock transfected HEK293 cells to determine fold activation. (**B**) qRT-PCR analysis of *IFNB1* transcript levels in HEK293 cells transfected with equimolar amounts (50 pmol) of IVT gRNA, Sendai Virus defective interfering (SeV DI) RNA or HCV PAMP,

 respectively. (**C**) qRT-PCR analysis of *IFNB1* and *ISG15* transcript levels in HEK293 cells over a

499   48h time course after transfection with 50 pmol via lipofection (Lipofectamine2000 or

500   RNAiMAX) or nucleofection, respectively. For all panels average values of three biological

501   replicates +/-SD are shown.

Figure 2

Figure 2: IVT gRNAs are recognized via the RIG-I pathway. (**A**) qRT-PCR analysis of increase

505   in *IFNB1* transcript levels (left) and transcript levels of the two main cytosolic RIG-I-like receptors

506   (*DDX58* and *IFIH1*) after introduction of IVT gRNA. Cell lines were ordered by responsiveness

507   to gRNA-mediated induction of *IFNB1* transcript levels. Cells were harvested for RNA extraction

508   30h after transfection. Ct values were normalized to Ct values of mock transfected cells for each

cell line to determine fold activation. *IFNB1* levels for K562 cells were too low to be determined

(n.d.). (**B**) Western Blot analysis for RIG-I and MDA5 expression of mock transfected and gRNA

transfected HEK293 cells after 48h. (**C**) qRT-PCR analysis of *IFNB1* transcript levels in HEK293

RIG-I (left), MDA5 (middle) and MAVS (right) knockout cells. Shown are three biological

replicates of three clonal populations of RIG-I, MDA5 or MAVS KO cells, respectively. *IFNB1*

levels for RIG-I KO clone #5 were too low to be determined (n.d.). For panels A and C, cells were

harvested for RNA extraction 30h after transfection using RNAiMAX transfection reagent.

Average values of three biological replicates +/-SD are shown. Statistical significances were

calculated by unpaired t-test (*p<0.0001, n.s.: not significant).

Figure 3

Figure 3. Protospacer and 5'-triphosphate determine the intensity of the gRNA-mediated

IFNβ response. (**A**) qRT-PCR analysis of *IFNB1* transcript levels in HEK293 cells transfected with equal amounts of gRNAs containing different 20 nucleotide protospacers. gRNAs were ordered by decreasing levels of *IFNB1* activation. gRNA1 refers to the gRNA that has been used in all previous experiments. (**B**) qRT-PCR analysis of *IFNB1* transcript levels in HEK293 cells transfected with synthetic, IVT and Calf intestine phosphatase treated (CIP) IVT gRNAs (gRNA1). After IVT and CIP treatment gRNAs were purified with SPRI beads or spin columns, respectively. Cells were harvested for RNA extraction 30h after transfection with RNAiMAX transfection reagent. Average values of three biological replicates +/-SD are shown. (**C**) Viability of human primary HSPCs 24h post nucleofection with no RNP and Cas9 or dCas9 RNPs. dCas9 or Cas9 were complexed with synthetic (syn) or IVT gRNA targeting the *HBB* gene. Viability was determined by Trypan blue exclusion test. (**D**) qRT-PCR analysis of *ISG15* and *DDX58* (RIG-I) transcript levels in human primary HSPCs 16h post nucleofection. dCas9 or Cas9 were complexed with synthetic or IVT gRNA targeting the *HBB* gene, respectively. Ct values were normalized against Ct of mock-nucleofected cells. Average values of two biological replicates +/-SD are shown. (**E**) Viability of human primary HSPCs 16h post transfection with RNPs. RNPs consisted of dCas9 complexed with synthetic, IVT or CIP-treated IVT gRNAs targeting a non-coding intron of *JAK2*. Viability was determined by Trypan blue exclusion test. Statistical significances were calculated by unpaired t-test (*$p<0.05$, **$p<0.01$, ***$p<0.0001$, n.s.: not significant).

Figure 4

Figure 4. Transfection of *in vitro* transcribed gRNAs induces a cytosolic immune response.

Proposed model of IVT gRNA recognition pathways in mammalian cells. IVT gRNAs carry a 5'-

triphosphate and when complexed with Cas9 protein and transfected into cells, cytosolic RNPs are

recognized by RIG-I triggering a cascade of activation events through the mitochondrial antiviral

signaling protein (MAVS). This results in phosphorylation of IRF3/7 and their shuttling into the

nucleus to activate expression of type I interferons (IFNα/β). This triggers the expression of

interferon-stimulated genes (ISGs). This innate immune response changes the transcriptome of the

cell and can cause cell stress and/or death which in turn might affect the editing outcomes.