Bacterial swarming reduces *Proteus mirabilis* and *Vibrio parahaemolyticus* cell stiffness and increases β-lactam susceptibility

George K. Auer[a], Piercen M. Oliver[b], Manohary Rajendram[b], Ti-Yu Lin[b], Qing Yao[c,d], Grant J. Jensen[c, d], and Douglas B. Weibel[a,b,e]*

[a] Department of Biomedical Engineering, University of Wisconsin-Madison, Madison, WI, 53706
[b] Department of Biochemistry, University of Wisconsin-Madison, Madison, WI, 53706
[c] Division of Biology and Biological Engineering, California Institute of Technology, Pasadena, CA 91125
[d] Howard Hughes Medical Institute, California Institute of Technology, Pasadena, CA 91125
[e] Department of Chemistry, University of Wisconsin-Madison, Madison, WI, 53706

* Author to whom correspondence should be addressed:
Douglas B. Weibel
Department of Biochemistry
6424 Biochemical Sciences Building
440 Henry Mall
Phone: +1 (608) 890-1342
Fax: +1 (608) 265-0764
E-mail: douglas.weibel@wisc.edu

Running title: Bacterial swarming reduces cell stiffness

Keywords: bacterial swarming, bacterial cell mechanics; peptidoglycan; osmotic pressure; antibiotics

Abstract

Swarmer cells of the gram-negative pathogenic bacteria *Proteus mirabilis* and *Vibrio*

parahaemolyticus become long (>10-100 μm) and multinucleate during their growth and

motility on polymer surfaces. We demonstrate increasing cell length is accompanied by

a large increase in flexibility. Using a microfluidic assay to measure single-cell

mechanics, we identified large differences in swarmer cell stiffness of (bending rigidity

of *P. mirabilis*, 9.6 x 10^{-22} N m²; *V. parahaemolyticus*, 9.7 x 10^{-23} N m²) compared to

vegetative cells (1.4 x 10^{-20} N m² and 3.2 x 10^{-22} N m², respectively). The reduction in

bending rigidity (~3-15 fold) was accompanied by a decrease in the average

polysaccharide strand length of the peptidoglycan layer of the cell wall from 28-30 to

19-22 disaccharides. Atomic force microscopy revealed a reduction in *P. mirabilis*

peptidoglycan thickness from 1.5 nm (vegetative) to 1.0 nm (swarmer) and electron

cryotomography indicated changes in swarmer cell wall morphology. *P. mirabilis* and *V.*

parahaemolyticus swarmer cells became increasingly sensitive to osmotic pressure and

susceptible to cell wall-modifying antibiotics (compared to vegetative cells)—they were

~30% more likely to die after 3 h of treatment with minimum inhibitory concentrations

of the β-lactams cephalexin and penicillin G. The adaptive cost of swarming is offset by

the increase in cell susceptibility to physical and chemical changes in their environment,

thereby suggesting the development of new chemotherapies for bacteria that leverage

swarming for colonization of hosts and survival.

Importance

Proteus mirabilis and *Vibrio parahaemolyticus* are bacteria that infect humans. To adapt to

environmental changes, these bacteria alter their cell morphology and move collectively

to access new sources of nutrients in a process referred to as 'swarming'. We found that

a change in the composition and thickness of the peptidoglycan layer of the cell wall

makes swarmer cells of *P. mirabilis* and *V. parahaemolyticus* more flexible (i.e., reduced

cell stiffness) and increases their sensitivity to osmotic pressure and cell-wall targeting

antibiotics (e.g., β-lactams). These results highlight the importance of assessing the

extracellular environment in determining antibiotic doses and the use of β-lactams

antibiotics for treating infections caused by swarmer cells of *P. mirabilis* and *V.*

parahaemolyticus.

1   **Introduction**

Bacteria have evolved a variety of mechanisms to adapt to their physical environment.

For example, in response to fluctuating environmental conditions, changes in

biochemistry and gene regulation can alter bacterial shape and increase cell fitness. Cell

filamentation is a commonly observed change in bacterial cell shape (1, 2) and has been

described as a mechanism that enables bacteria to evade predation by the innate

immune system during host infections (1).

In close proximity to surfaces, many bacteria alter their morphology and leverage

cell-cell physical contact to move collectively to access new sources of nutrients and

growth factors (3, 4). Referred to as 'swarming', this process is common among motile

bacteria and has been connected to bacterial pathogenesis and infections (3, 4). Swarmer

cells of *Salmonella enterica*, *Pseudomonas aeruginosa*, *Serratia marcescens*, and *Bacillus*

subtilis have reduced antibiotic susceptibility—compared to vegetative cells—to a

variety of drugs that alter protein translation, DNA transcription, and the bacterial cell

membrane and cell wall (5-8). The specific biochemical and biophysical mechanisms

underlying these observations are unknown.

Here, we describe physical changes in swarmer cells of the gram-negative

pathogenic bacteria *Proteus mirabilis* and *Vibrio parahaemolyticus* that have the opposite

effect: they increase the susceptibility of cells to cell wall-targeting clinical antibiotics.

We found that large changes in the length of *P. mirabilis* and *V. parahaemolyticus*

swarmer cells are accompanied by an increase in flexibility (i.e., a reduction in cell

stiffness) that enables long cells to pack together tightly and form cell-cell interactions;

maximizing cell-cell interactions promotes surface motility (9). Using biophysical,

biochemical, and structural techniques, we quantified changes in the structure and

composition of the *P. mirabilis* and *V. parahaemolyticus* cell wall in swarmer and

vegetative cells and characterized their susceptibility to osmotic changes and cell wall-

modifying antibiotics. Our results indicate that morphological changes that enable these

bacteria to adapt to new physical environments come at a significant fitness cost, as cells

become more susceptible to their chemical environment. In particular, changes in the

composition and thickness of *P. mirabilis* and *V. parahaemolyticus* swarmer cells make

them more sensitive to osmotic changes and to cell-wall modifying antibiotics, thereby

suggesting that these classes of drugs may be useful in treating infections of these

bacteria (e.g., in urinary tract infections).

Results

The bending rigidity of *P. mirabilis* and *V. parahaemolyticus* cells decreases during

swarming. During surface motility, *P. mirabilis* and *V. parahaemolyticus* cells grow into

swarmers that are characteristically long (10-100 μm) and present a high surface density

of flagella that enables them to translate through viscous environments (3, 10). We

found that these swarmer cells display an unusual phenotype that is rarely observed

among gram-negative bacteria: remarkable flexibility and a shape that is dynamically

altered by adjacent cell motion and collisions (Fig. 1). The ability of *P. mirabilis* swarmer

cells to maximize cell-cell contacts plays a role in their cooperative motility (10); our

observations indicate that flexibility enables these long cells to optimize packing into

multicellular structures that move cooperatively across surfaces.

Bacterial cell mechanics is generally attributed to the peptidoglycan (PG) layer of

the cell wall, which has a thickness of ~3-50 nm and surrounds the cytoplasmic

membrane (11). Very little is known about mechanical regulation in bacteria (12-17) and

we are unaware of studies connecting swarming to changes in cell mechanics. We

quantified changes in swarmer-cell stiffness using cell-bending assays in a reloadable,

poly(dimethylsiloxane) microfluidic system (Fig. 2 and S1) that is related to a method

developed previously (18). In bending assays, we applied a shear fluid force to multiple

filamentous cells, resulting in horizontal deflection of their cell tips (Fig. 2); fitting the

deflection data to a mechanical model provided us with a value of the (flexural)

bending rigidity of cells (Fig. S2). Introducing a reloadable mechanism enabled us to

perform rapid bending measurements of *P. mirabilis* and *V. parahaemolyticus* swarmer

cells after isolating them from swarm plates. Once removed from a surface, *P. mirabilis*

and *V. parahaemolyticus* swarmer cells dedifferentiate, grow, and divide to form

vegetative cells with a wild type length, requiring us to rapidly perform assays with

swarmer cells after their isolation from surfaces. As a point of comparison, we

filamented vegetative cells of *P. mirabilis* and *V. parahaemolyticus* using aztreonam—an

inhibitor of the division specific transpeptidase PBP3—to match the length of swarmer

cells (22.2 ±12.5 µm and 12.4 ± 8.2 µm respectively) and compared their bending rigidity

values to swarmer cells. As a control, we measured the bending rigidity of cells of

Escherichia coli strain MG1655 that we filamented using aztreonam, and determined the

value to be 4.2×10^{-20} N m^2 (Fig. 3); using a value for the thickness of the PG of 4-nm (19)

yields a Young's modulus of 26 MPa, which is close to values that have been reported

previously and supports the choice of using aztreonam to filament cells, as it apparently

has no effect on the bending rigidity of cells (12, 18). We found a substantial decrease in

the bending rigidity of swarmer cells of both *P. mirabilis* (~15-fold) and *V.*

parahaemolyticus (3-fold) compared to vegetative cells (Fig. 3), which is consistent with

our observations of the shape and behavior of these cells during swarming on surfaces.

V. parahaemolyticus vegetative cells were remarkably flexible: ~134-fold more so than *E.*

coli cells and ~3-fold more than *P. mirabilis* swarmer cells (Fig. 3).

To confirm that using aztreonam to inhibit PBP3 and produce filamentous cells

does not change the cross-linking density of PG at the division plane and alter cell

mechanics, we compared bending rigidity values of cells treated with aztreonam and

cells filamented by overexpressing SulA, a protein that prevents polymerization of the

division protein FtsZ and blocks cell division. Both mechanisms of filamenting *E. coli*

cells produced similar bending rigidity values: 4.0×10^{-20} N m^2 (SulA) and 4.2×10^{-20} N

m^2 (aztreonam) (Fig. S3A).

During swarming, *P. mirabilis* and *V. parahaemolyticus* cells dramatically increase

their density of flagella. Since flagella are attached to these cells through a protein

complex that extends through the PG (referred to as the basal body), it is possible that

these structures create local defects in the PG that alter its mechanical properties. To test

whether the density of flagella on swarming cells is responsible for the changes in cell

mechanics that we observe, we performed bending rigidity measurements (Fig. S3B) on

two K-12-derived strains of *E. coli* that have substantially different flagella densities

(Fig. S3C). Our results indicated no appreciable change in stiffness values (bending

rigidity of 4.0×10^{-20} N m^2 at high flagella density and 4.2×10^{-20} N m^2 at low flagella

density; Fig. S3B).

Overexpressing FlhDC—the heterohexameric activator that is important for

swarming—in vegetative cells growing in liquid produces a phenotype that replicates

many of the characteristics of swarmer cells, including increased cell length and flagella

density (20). A relationship between FlhDC production and cell mechanics in untested.

To test whether FlhDC is connected to changes in swarmer cell stiffness, we

overexpressed the protein from the plasmid-encoded genes *pflhDC* in filamentous cells

of *P. mirabilis* and measured their bending rigidity. We detected a ~1.5-fold difference in

bending rigidity between wildtype (1.4×10^{-20} N m^2) and *pflhDC*-containing *P. mirabilis*

vegetative cells (9.2×10^{-21} N m^2), and no difference between wildtype and *pflhDC*-

containing *P. mirabilis* swarmer cells (~9.6×10^{-22} N m^2) (Fig. 3). These results indicate

that FlhDC overexpression during swarming is a minor contributor to the mechanical

phenotype of swarming cells; the majority of the mechanical changes we observe arise

from another regulatory pathway(s).

***P. mirabilis* and *V. parahaemolyticus* swarmer cells are more sensitive to osmotic**

stress than vegetative cells. To complement cell stiffness measurements, we also

studied changes in the mechanics of swarmer cells and filamented vegetative cells by

measuring their changes in length and width in response to changes in osmotic

pressure, which cause cells to shrink (in a NaCl solution) and swell (in H$_2$O) (Fig. S4).

We filamented *P. mirabilis*, *V. parahaemolyticus*, and control *E. coli* cells using aztreonam

to create a range of cell lengths that matched the lengths of *P. mirabilis* and *V.*

parahaemolyticus swarmer cells. We expected that low values of bending rigidity and

compositional and structural changes in PG would cause swarmer cells to elongate in

response to changes in osmotic pressure (in comparison to filamented, vegetative cells).

We found that *P. mirabilis* and *V. parahaemolyticus* swarmer cell extension in response to

osmotic shock was significantly larger than filamented, vegetative cells (Fig. 4). To

circumvent the cellular production of osmolytes to protect cells from large changes in

osmotic pressure during shock (typically produced within 1 min) (21), we used a

microfluidic device to rapidly switch between H_2O and a NaCl solution and measured

changes in cell length before cells adapted (Fig. S4).

Osmotic shifts produced small changes in the length (Fig. 4A) and width (Fig.

4B) of vegetative filamented *P. mirabilis* cells compared to *E. coli* cells (Figs. 4A,B). In

contrast, *V. parahaemolyticus* vegetative cells substantially increased in cell length (Fig.

4C) compared to *E. coli* cells, with no observable change in cell width (Fig. 4D). In

response to osmotic upshifts (i.e., transitioning from H_2O to NaCl), *P. mirabilis* and *V.*

parahaemolyticus swarmer cells dramatically increased in cell length (Figs. 4A,C) (and to

a lesser extent, cell width for *P. mirabilis* but not for *V. parahaemolyticus*; Figs. 4B,D)

compared to vegetative cells. These results suggest that changes in *P. mirabilis* and *V.*

parahaemolyticus swarmer cell stiffness make them more responsive to osmotic changes.

Changes in PG composition of *P. mirabilis* and *V. parahaemolyticus* swarmer cells.

We next studied the structural and biochemical mechanisms underlying the change in

mechanical properties of swarmer cells. The connection between the structure of PG

and cell mechanical properties—dating back to early antibiotic studies that revealed the

penicillin binding proteins—suggested PG a starting point for these studies.

PG consists of the disaccharide building block β-(1,4)-N-acetylmuramic acid/N-

acetyl-glucosamine (MurNAc-GlcNAc) in which a pentapeptide is attached to each 3'-

OH group on MurNAc. Cross-linking between adjacent pentapeptides creates a mesh-

like polymeric layer, and altering its structure and composition affects cell-mechanical

properties (14, 15). To determine whether the PG composition of *P. mirabilis* and *V.*

parahaemolyticus cells changes during swarming, we isolated PG sacculi from vegetative

and swarmer cells and used ultra performance liquid chromatography-mass

spectrometry (UPLC-MS) to quantify its chemical composition (Fig. S5). As the PG

composition of *V. parahaemolyticus* has not yet been reported, we characterized its

muropeptide stem peptide using UPLC-MS/MS (Fig. S6 and Table S1). Similar to *E. coli*

(22) and *P. mirabilis* (23), *V. parahaemolyticus* has a PG structure that is conserved across

other gram-negative bacteria, in which the peptide stem consists of L-Ala-D-Glu-*meso*-

diaminopimelic acid (*meso*-DAP)-D-Ala-D-Ala (Fig. S6).

Compared to vegetative cells, *P. mirabilis* swarmer cells contained fewer

monomers (MurNAc-GlcNAc), more dimers, and more anhydrous-containing

saccharides, which are found at the terminating end of glycan polymers (Fig. 5A) (11).

We detected no differences in the relative abundance of trimers between swarmer and

vegetative cells of *P. mirabilis* (Fig. 5A). The increase in anhydrous-containing

saccharides that we observed in *P. mirabilis* swarmer cells was correlated with a

decrease in polysaccharide length (Fig. 5B). A similar increase in anhydrous-containing

saccharides and decreased length of polysaccharides occurred in *V. parahaemolyticus*

swarmer cells (Fig. 5A,B). We found no change in cross-linking density between

vegetative and swarmer cells of either *P. mirabilis or V. parahaemolyticus* (Fig. 5C).

Swarmer cells have a reduced PG thickness and display membrane defects. Changes

in the thickness of the PG layer and structure of the cell envelope may also explain the

observed decrease in swarmer cell stiffness. To identify changes in PG thickness of

swarmer cells, we isolated intact sacculi from *P. mirabilis* vegetative and swarmer cells

and measured the thickness of dried sacculi using tapping-mode atomic force

microscopy (AFM) (Fig. 6A). Differences in the nanoscopic appearance of the sacculi of

different cells were not observed by AFM (Fig. S7). The thickness of isolated, dry *P.*

mirabilis swarmer cell sacculi (1.0 ± 0.2 nm) was reduced ~1.5-fold compared to

vegetative cells (1.5 ± 0.2 nm) (Fig. 6A). *V. parahaemolyticus* swarmer cells (0.6 ± 0.1 nm)

exhibited a similar ~1.2-fold decrease in thickness compared to vegetative cells (0.8 ± 0.2

nm). Earlier AFM measurements of isolated sacculi indicated that dehydration reduced

the thickness of *Escherichia coli* PG by ~2x, which we used to estimate the dimensions of

hydrated PG from *P. mirabilis* (3.1 and 2.1 nm for vegetative and swarmer cells,

respectively) and *V. parahaemolyticus* (1.7 and 1.4 nm, respectively) (24). A comparison

of PG thickness and cell bending rigidity suggested that the relationship between these

data is approximately exponential (R^2=0.9874) (Fig. 6B).

A caveat with conversions between dried and hydrated values is that they are

most accurate for PG that best mimics the structure and composition (e.g., crosslinking

and polysaccharide composition) of the reference material: *E. coli* PG. Alterations in the

polysaccharide length of PG from *P. mirabilis* and *V. parahaemolyticus* may be more

elastic and stretch out during drying, thereby appearing to have a thickness that is

reduced compared to *E. coli* PG. Our control measurements on isolated, dry *E. coli*

sacculi yielded a thickness of 2.0 nm, which varies from the value of 3.0 nm published

by Yao et al. using the same technique (24). A difference of ~30% between these *E. coli*

measurements may arise for several reasons, including: variations in physical

conditions that impact AFM measurements, improvement in the resolution of AFMs,

and/or the precision of fitting force curves.

The relatively low variability in the values we measured for isolated sacculi from

P. mirabilis and *V. parahaemolyticus* cells (~16-25%) using AFM demonstrated a consistent

reduction in PG thickness for swarmer cells, suggesting that our measurements are

sufficient for comparing PG from vegetative and swarming cells, and demonstrating a

connection between PG and changes in cell mechanics. In contrast, the variability in

AFM measurements and analysis makes us less comfortable comparing absolute values

of PG thickness in our studies to those reported for bacteria in other papers.

To complement AFM measurements, we attempted to measure the thickness of

native PG using electron cryotomography (ECT) on intact vegetative and swarmer cells

(Figs. 6C,D, S8, and S9). Although we were unable to resolve the thickness of native PG

by direct ECT measurements, we noticed that the sub-tomogram-averaged ECT

volumes of the *P. mirabilis* cell wall (Fig. 6C) indicated the distance between the inner

and outer membranes of swarmer cells was smaller than in vegetative cells (Figs. 6C,D).

P. mirabilis vegetative cells had a characteristically smooth membrane (Fig. S8A,B) that

was similar to the presentation of the membrane found along the lateral, cylindrical

walls of swarmer cells (Fig. S8C,D). In contrast, the polar regions of both *P. mirabilis* and

V. parahaemolyticus swarmer cells had an undulating outer membrane suggestive of an

altered structure (Fig. S8E,F and S9D-F), and *V. parahaemolyticus* cells had significant

defects in their cell-envelope, including, membrane budding, vesicle formation, and

ruptured cell walls (Fig. S9D-F).

Increased susceptibility of *P. mirabilis* and *V. parahaemolyticus* swarmers to the cell

wall-targeting antibiotics. Although swarming colonies of bacteria display resistance to

many antibiotics (5), our experiments suggest *P. mirabilis* and *V. parahaemolyticus*

swarmer cells have changes in PG structure, composition, and properties that may

increase their susceptibility to cell wall-targeting antibiotics. To determine the

sensitivity of *P. mirabilis* and *V. parahaemolyticus* swarmers to the cell wall-targeting

antibiotics cephalexin (inhibits PBP3 and 4) (25) and penicillin G (inhibits PBP3, 4, and

6), we measured swarmer single-cell growth in the presence of the minimum inhibitory

concentration (MIC) of antibiotics in a microfluidic device. We were unable to measure

MICs using traditional protocols because the swarmer cells dedifferentiate at shorter

time scales than these protocols. Instead, we chose 3 h of incubation for our

measurement for three key reasons: 1) swarmer cells do not dedifferentiate over this

time period; 2) a past study demonstrated that treating cells with cephalexin for 3 h was

sufficient to kill ~100% of a population of *E. coli* cells (26); and 3) ~2 h of cephalexin

treatment was sufficient to observe cell lysis of *E. coli* cells measured by single cell

growth (27).

Using a microfluidic device enabled us to maintain the swarming phenotype of

cells during experiments and supply cells with a source of continuously replenished

fresh growth media to ensure exponential cell growth and a constant concentration of

antibiotics. We found that the survival of *P. mirabilis* (66%) and *V. parahaemolyticus* (64%)

vegetative cells treated with 1X MIC of cephalexin was slightly higher than *E. coli* cells

(55%) (Figs. 7A, S10). Treating *P. mirabilis* and *V. parahaemolyticus* swarmer cells with 1X

MIC of cephalexin reduced survival to 37% and 19%, respectively (Figs. 7A, S10),

indicating their increased susceptibility. Rates of cell survival in the presence of

penicillin G were similar to the use of cephelexin (Fig. 7A). We characterized dead cells

using microscopy to measure membrane blebbing, cell lysis, and changes in the

refractive index of cells.

Cell wall-targeting antibiotics are most effective against actively growing cells

(28, 29) and therefore increases in growth rate will reduce cell survival. To determine

whether a growth rate phenotype explains our observations, we normalized data for

cell length, and were unable to detect a change in the growth rate of swarmer cells

when they were treated with these two antibiotics (Fig. S11). We measured the mean

survival time of cells (the amount of time elapsed after treatment with drugs before cell

death) for *P. mirabilis* and *V. parahaemolyticus* swarmer cells treated with cephalexin and

penicillin G (Fig. 7B) and found that the survival time for these swarmer cells was lower

than that for vegetative cells (Fig. 7B), which is consistent with alterations in the PG.

Discussion

P. mirabilis is commonly associated with complicated urinary tract infections and

increased mortality in cases of bacteremia (30). Swarming is hypothesized to enable *P.*

mirabilis cell movement from the ureter to the kidney; in support of this hypothesis,

swarming deficient mutants have lower rates of host infection (31). We found that

during swarming, *P. mirabilis* and *V. parahaemolyticus* cells become more flexible due to

changes in the thickness and composition of the PG. Although we were unable to

precisely measure the thickness of isolated PG, the decrease in the distance between the

inner and outer membranes of swarmers may reflect a reduction in PG thickness.

Several recent studies suggest how changes in the cell wall influence cell stiffness. A

recent study revealed the connection between changes in the biochemical regulation of

D-Ala-D-Ala and *Pseudomonas aeruginosa* cell stiffness in cells in which this biochemical

machinery is altered (32). As our paper and the paper on P. aeruginosa demonstrate

(32), it is still difficult to quantitatively understand the scaling between the magnitude

of the biochemical changes in cells and the change in cell stiffness. The range of

techniques now available for quantitatively measuring single cell stiffness will enable

the development of this relationship.

We found that *P. mirabilis* and *V. parahaemolyticus* swarmer cells were more

susceptible to large changes in cell length than vegetative cells during osmotic shift

experiments. These changes are consistent with what would be expected from a

reduction in PG thickness in swarming cells. We also observed that the length of

polysaccharides was reduced in swarming cells, which would not obviously lead to

large changes in cell elongation during osmotic shifts if the organization of PG in these

cells follows what is known about the structure of PG in *E. coli* cells (33):

polysaccharides reported to be arranged circumferentially around cells with the peptide

cross-links oriented parallel to the long axis of cells. Instead, we would expect that the

decrease in polysaccharide lengths make swarmer cells more prone to changes in

diameter. A careful analysis of the data (Fig. 5) indicates that *P. mirabilis* swarmer cells

do increase in diameter during osmotic shifts (with no change occurring for vegetative

cells); no clear trend is observable with *V. parahaemolyticus* swarmer cells.

Another recent study demonstrated the role of the outer membrane in the cell

stiffness of gram-negative bacteria (34). Chemical and genetic changes in the outer

membrane increased the deformation of the cell envelope to stretching, bending, and

indentation forces and increased the amount of cell lysis arising during osmotic shock.

It is possible that the morphological changes in swarmer cell membranes that we

observed during ECT experiments indicate membrane perturbations that contribute

significantly to changes in cell stiffness. Swarming in *Proteus* and *Vibrio* is well known to

occur in parallel with the secretion of lipopolysaccharides (3, 4), the source of which is

presumably the outer membrane of these bacteria, which may alter cell stiffness.

This paper demonstrates for the first time the ability of bacteria to modulate their

cell stiffness (beyond L-forms). Presumably this phenotype (i.e., more flexible) conveys

an adaptive advantage for cells that is supported by observations from many labs that

the motility of communities of swarmer cells is enhanced when cell-cell contacts are

increased. The adaptive advantage of swarming in *P. mirabilis* and *V. parahaemolyticus*

however appears to be offset by a decrease in their fitness, as cells become more

sensitive to osmotic changes and cell wall-targeting antibiotics, thereby creating an

"Achilles heal" for targeting this phenotype in infectious diseases. This study highlights

the plasticity of bacteria (1), the continuing need for methods of evaluating

chemotherapies that measure efficacy against cells in a physiological state relevant to

specific infections, and the reconsideration of cell wall-targeting antibiotics for treating

UTIs in which *P. mirabilis* may be present.

Materials and Methods

Bacterial strains and cell culture. *P. mirabilis* strain HI4320, *V. parahaemolyticus* LM5674,

Escherichia coli MG1655 (CGSC #6300), and plasmids *pflhDC* (10) and *psulA*(18) were

used for experiments used in this paper. *P. mirabilis* was grown in PLB nutrient medium

consisting of 1% peptone (weight/volume), 0.5% yeast extract (w/v), and 1% NaCl (w/v).

V. parahaemolyticus was grown in nutrient medium (HI medium) consisting of 2.5%

heart infusion (w/v) and 2.5% NaCl (w/v). *E. coli* was grown in lysogeny broth (LB)

consisting of 1% tryptone (w/v), 0.5% yeast extract (w/v), and 1% NaCl (w/v). All strains

were grown at 30 °C with shaking at 200 rpm.

Immunostaining flagella on *E. coli* cells to determine flagella surface density. As a

control to determine the influence of the surface density of bacterial flagella on

measurements of bending rigidity, we imaged two *E. coli* MG1655 strains with different

flagella densities using a polyclonal flagellin antibody and an immunostaining

procedure for flagella visualization (35).

Calculating vegetative and swarmer cell division times. We prepared vegetative cells

by diluting an overnight culture 1:200 in fresh medium and grew the cells at 30 °C with

shaking at 200 rpm to an optical density (OD; λ=600 nm) of 0.6. We prepared swarmer

cells as described previously (35). We monitored the growth of *P. mirabilis*, *V.*

parahaemolyticus, and *E. coli* vegetative cells over 120 min at 30 °C in the microfluidic

flow chamber device described in the next section. We performed a similar experiment

with *P. mirabilis* and *V. parahaemolyticus* swarmers growing over 65 min at 30 °C to

determine the amount of time elapsed before cells divided. We collected images of cells

at 1-min intervals and determined the division time for a maximum of 10 generations

(for vegetative cells) and 4 division events (for swarmer cells).

Measuring the sensitivity of cells to osmotic shock in a microfluidic device. We

prepared filamentous vegetative cells of *P. mirabilis* and *V. parahaemolyticus* by diluting

an overnight culture of cells 1:200 into fresh medium. We then grew cultures at 30 °C

with shaking for 1 h, added aztreonam (MP Biomedicals) to a final concentration of 10

μg/mL and grew cells for an additional 70 min. Swarmer cells were prepared as

described previously (35). To maximize the number of cells attached to a surface in the

microfluidic device (flow chamber construction described below), we concentrated cells

to an OD_{600} of 8 in 100 μL of liquid nutrient medium.

We prepared the microfluidic device for experiments by flowing 10 μL of

undiluted Cell-Tak (Corning) into the device and incubated it for 10 min at 25 °C. Next,

we flowed 20 μL of a suspension of cells (OD_{600}=8) through the device, then repeated

with another 20 μL aliquot of cell suspension. To aid the adherence of the highly motile

swarmer cells to the Cell-Tak-coated surface, we centrifuged the device for 5 min at 300

x g in a centrifuge (Beckman Coulter) equipped with a swinging bucket rotor.

For osmotic shock experiments, we filled separate 1-mL syringes (BD) with a

solution of UltraPure distilled water (Invitrogen) or 1 M of a NaCl solution prepared in

ddH$_2$O and connected the syringes using a three-way valve. We mounted the

microfluidic device on a TE2000-E inverted microscope and imaged cells in the chamber

at 25 °C (decreasing the temperature reduced growth and cell division). Prior to osmotic

shock, we flowed 200 μL of fresh liquid media (PLB for *P. mirabilis*, LB for *E. coli*, and HI

for *V. parahaemolyticus*) through the device to remove cells that were not adhered to the

channel surface. While imaging, we flowed a 1 M NaCl solution through the channel

and observed cell plasmolysis, immediately after which we flowed ddH$_2$O through the

channel and observed cells elongating. We collected images of hundreds of single cells

after exposing them to three conditions: 1) isotonic culture (PLB, LB, HI); 2) hypertonic

shock (1 M NaCl); and 3) hypotonic shock (ddH$_2$O).

To determine changes in cell size and width, we extracted individual cell

contours from the images taken for each treatment using MATLAB 2014a (MathWorks)

and used MicrobeTracker (36) to determine cell lengths (*L*) and widths (*W*). From these

measurements we calculated ΔL ($L_{\text{hypotonic}} - L_{\text{hypertonic}}$) and ΔW ($W_{\text{hypotonic}} - W_{\text{hypertonic}}$) for

vegetative and swarmer cells of *P. mirabilis* and *V. parahaemolyticus*.

Determining the minimum inhibitory concentration (MIC) of vegetative cells of *P.*

mirabilis **and *V. parahaemolyticus.*** We used the micro-dilution protocol (37)

to determine the MIC of cephalexin in accordance with Clinical and Laboratory

Standards Institute. Briefly, we added 400 µg/mL of cephalexin (BP Biomedicals) or

penicillin G (BP Biomedicals) to the first well of a 96-well microplate (Nunc) and diluted

these antibiotics 2-fold across adjacent wells (wells #1-11); well 12 was a no-drug

control. We determined the MIC after 16 h of growth at 30 °C with shaking by

identifying the lowest concentration of cephalexin and penicillin G that inhibited cell

growth by visual inspection. The MIC was determined from three replicate plates. We

determined the MIC of cephalexin against *E. coli* MG1655 (1X MIC = 6.25 µg/mL, 32X

MIC = 200 µg/mL), *P. mirabilis* (100 µg/mL), and *V. parahaemolyticus* (50 µg/mL). The

MIC of penicillin G against *P. mirabilis* was 12.5 µg/mL.

Antibiotic treatment of vegetative and swarmer cells and measurement of their

growth in a microfluidic device. We prepared vegetative and swarmer cells of *P.*

mirabilis and *V. parahaemolyticus* as described above. Prior to use, we diluted cells 1:100

in fresh medium to a cell density that enabled us to image many individual cells

simultaneously in the microfluidic device described in the section above.

To prepare the microfluidic device for monitoring growth, we applied a 250 µm-

thick layer of PDMS prepolymer to the surface of (#1.5, 35 x 50 mm) cover glass

(FisherBrand) using a spincoater (Laurell Technologies) and polymerized the polymer

overnight at 60 °C. We subsequently removed a 6 mm x 4 mm rectangle of PDMS from

the center of the cover glass using a scalpel. We applied transparent tape to both sides

of this rectangular well and pipetted 100 μL of media containing a 2% (w/v) solution

UltraPure agarose (Invitrogen) into the well. We placed a (#1.5, 22 x 30 mm) cover glass

(FisherBrand) on top of the liquid agarose (to flatten the agarose surface), pressed the

cover glass against the transparent tape, and solidified the agarose at 25 °C. Next, we

removed the #1.5 cover glass, tape, and any residual agarose from the PDMS surface.

We pipetted 2 μL of a suspension of cells on the agarose pad surface, waited until the

excess liquid had evaporated or been absorbed by the agarose, carefully removed the

agarose layer, inverted it, and placed it back into the well such that the cells were

positioned between the agarose surface and the cover glass. We placed the PDMS flow

chamber used in the construction of the osmotic shock microfluidic device on the

PDMS-coated cover glass and aligned it such that the agarose pad was centered in the

flow channel (Fig. S12).

For antibiotic treatment of cells in the microfluidic cell growth device, we filled a

6-mL syringe (Norm-Ject) with cephalexin or penicillin G at a concentration

corresponding to 1X MIC dissolved in nutrient medium. We supplied a constant flow of

20 μL/min to the device using a syringe pump (Harvard Apparatus) and monitored the

growth of individual cells during antibiotic exposure using a TE2000-E inverted

microscope. The stage and objective heater were maintained at 30 °C. Images were

collected every 1 min for 3 h.

Due to the aberrant shape of cells treated with cephalexin and penicillin G (Fig.

S10), we were unable to use an automated script (e.g., MicrobeTracker) to determine cell

death. Instead, we visually determined the time of death for individual cells when they

exhibited three phenotypes that collectively indicate cell death: blebbing (membrane

swelling), lysis (bleb rupture), and disappearance of cells in phase contrast microscopy

(loss of cytoplasmic material).

Determining cell growth rates in the presence of antibiotics. We prepared vegetative

and swarmer cells of *P. mirabilis* and *V. parahaemolyticus* as described above. We

monitored individual cell growth at 30 °C in the presence of 1X MIC of cephalexin and

pencillin G in our microfluidic growth device (described above) by collecting an image

every 1 min for 15 min. To calculate the growth rate, we first extracted individual cell

contours at each time point using MicrobeTracker and determined cell length. To take

into account the differences in starting length between *P. mirabilis* and *V.*

parahaemolyticus cells, we normalized the change in length to the initial cell length

($\Delta L/L_0$) at each time point. To determine the growth rate for individual cells, we fit their

relative length over time to an exponential function using GraphPad Prism 6.0

(GraphPad Software).

Measuring cell envelope architecture and peptidoglycan thickness using electron

cryotomography. We prepared vegetative and swarmer cells of *P. mirabilis* and *V.*

parahaemolyticus and concentrated them to an OD_{600}=10. For ECT, we mixed vegetative

and swarmer cells with bovine serum albumin-treated 10-nm diameter gold particles

that served as fiducial markers, applied them to electron microscopy grids, and plunge-

froze them in a mixture of liquid ethane and propane, as described previously (38).

Grids were stored in liquid nitrogen until imaging.

We acquired images on a 300 KeV Polara transmission electron microscope (FEI)

with a GIF energy filter (Gatan) and a K2 Summit direct detector (Gatan). We collected

tilt series from -60° to +60° with 1° increments using UCSFtomo software with a defocus

of -10 μm and a total dosage of 190 $e^-/Å^2$ (39) at a magnification of 27500×. Tomograms

were calculated using IMOD software (40).

For sub-tomogram averaging, smooth and flat membrane regions were chosen

by eye; a volume of 40 × 70 × 12 voxels (62 nm × 109 nm × 19 nm) was centered using the

outer membrane and extracted. We aligned 38 extracted "membrane fragments" from

four *P. mirabilis* vegetative cells and 42 fragments from nine *P. mirabilis* swarmer cells

and averaged them in PEET (41). The densities from two averaged membranes were

scaled to match each other using IMOD (40), cross-averaged density profiles were

measured using ImageJ 1.50c (42), and figures were generated in OriginPro (OriginLab).

Determining the thickness of *P. mirabilis* and *V. parahaemolyticus* sacculi (isolated

peptidoglycan) by atomic force microscopy (AFM) in ambient conditions. We isolated

P. mirabilis and *V. parahaemolyticus* swarmer cells, concentrated them at 800 x *g* for 10

min, removed the supernatant, flash froze the cell pellet in liquid nitrogen, and stored it

at -80 °C. Swarmer cell pellets were thawed at 4 °C and pooled for isolation of sacculi

(intact peptidoglycan). To increase the efficiency of cell lysis prior to isolating sacculi,

we resuspended vegetative and swarmer cell pellets in 3 mL of cold 1X phosphate-

buffered saline (ThermoScientific), then lysed cells with a tip sonicator (Qsonica) for ~10

s at a power setting of 75%. We confirmed cell lysis using optical microscopy. We

isolated sonicated cells, resuspended sacculi in 20 (*V. parahaemolyticus* swarmers) or 200

μL of ddH$_2$O (all other cells), immediately flash froze the sacculi in liquid nitrogen, and

stored them at -80 °C.

To prepare sacculi for AFM, we transferred 10 μL of sacculi thawed at 4 °C to a

new microcentrifuge tube placed in a bath sonicator (Branson) that was cooled with ice

for 10 min to aid in the dispersal of sacculi without affecting sacculus architecture (43).

After sonication, we pipetted 10 μL of the sacculi onto freshly cleaved mica (Ted Pella),

dried the mica under nitrogen gas, washed it 3x with 1 mL of ddH$_2$O (filtered through a

0.2 μm-diameter pore filter (Corning)), dried the sacculi under nitrogen gas, and imaged

immediately after preparation.

We performed AFM using a Catalyst AFM (Bruker) operating in tapping mode in

ambient conditions (air) with an aluminum reflex-coated silicon AFM probe (Ted Pella;

k = 40 N/m). Before imaging, AFM probes were auto-tuned using Nanoscope 8.15

(Bruker). We collected all images at high resolution (512 x 512 pixels) with a scan speed

of 1 Hz and analyzed images using NanoScope Analysis 1.4 (Bruker). Prior to

determining sacculi thickness, we flattened (0^{th} order) all images to remove variations in

surface thickness. Thickness was determined perpendicular to the long axis; we avoided

surface debris, folds in the sacculi, and trapped material in sacculi (Fig. S7).

Determining the composition of peptidoglycan in *P. mirabilis* and *V.*

***parahaemolyticus* vegetative and swarmer cells using ultra high performance liquid**

chromatography/mass spectrometry (UPLC/MS). We prepared vegetative and swarmer

cells of *P. mirabilis* and *V. parahaemolyticus* as described above. To purify peptidoglycan

for UPLC/MS, we carried out a previously reported isolation technique for Gram-

negative bacteria (44) with the following modifications. Briefly, after trypsin (Sigma)

inactivation, we incubated the sacculi in 1 M HCl solution (Fluka) for 4 h at 37 °C to

remove any O-acetylation from the peptidoglycan, which is present in *P. mirabilis* (45).

Then, the sacculi were washed three times in ddH_2O, resuspended in 500 mM boric acid

(Sigma) [pH 9] to OD_{600}=3, and mixed with 1/10 the volume of mutanolysin (Sigma). The

sample was incubated 16 h at 37 °C with 200 rpm shaking. The next day, the samples

were centrifuged for 10 min at 9500 x g, pelleting the remaining insoluble material. The

supernatant was removed and put into a glass vial. To reduce the isolated muropeptide

fragments, we added 50 µL of 20 µg/mL sodium borohydride (Sigma) in 500 mM boric

acid [pH 9] and incubated the mixture for 30 min at 25 °C. We adjusted the pH of the

solution to 2-3 by adding 50% phosphoric acid (Fluka), then filtered the muropeptide

solution through a Duropore polyvinylidine fluoride filter (0.22-µm pores; Millex) into a

clean vial. Vials were immediately stored at -80 °C until use within 1 week of

muropeptide isolation.

For UPLC/MS, we injected 7.5 µL of purified muropeptides on a Cortecs 2.1 x 100

mm C18 column (Waters) packed with 1.6 µM-diameter particles and equipped with a

Cortecs C18 guard column (Waters). The column temperature was maintained at 52 °C

using an Acquity standard flow UPLC system equipped with an inline photodiode

array (Waters). For muropeptide separation by UPLC, we used solvent A (Optima

LCMS-grade water with 0.05% trifluoroacetic acid) and solvent B (30% (v/v) Optima

LCMS-grade methanol in Optima LCMS-grade water with 0.05% trifluoroacetic acid)

(Fisher Scientific). Muropeptides were eluted from the column with a gradient of

increasing solvent B (1 min hold at 1% B, ramp to 99% B over 59 min, hold at 99% B for

5 min, then decrease to 1% B over 1.5 min, then hold at 1% B for 4.5 min) at a flow rate

of 0.2 mL/min. We analyzed the eluent from the column using a Bruker MaXis Ultra-

High Resolution time-of-flight 4G mass spectrometer (Bruker Daltonic) with either an

MS method or a data-dependent, top 3 MS/MS method. For both methods, capillary

voltage was set to 4100 V, the nebulizer pressure was 2.0 bar, and the drying gas was set

to 6.0 L/m at 220 °C. Muropeptides were detected at $\lambda=205$ nm by MS.

We determined the peptidoglycan composition of *E. coli*, *P. mirabilis*, and *V.*

parahaemolyticus vegetative and swarmer cells by comparing MS/MS fragmentation

patterns using DataAnalysis version 4.2 (Bruker) (Fig. S6). Muropeptides were

identified according to mass values using DataAnalysis 4.2. We calculated muropeptide

masses using ChemDraw 14.0 (CambridgeSoft) (Table S1). We quantified the

corresponding UV ($\lambda=205$) absorbing peaks (Fig. S5) identified by MS (Bruker) from

which we calculated peptidoglycan cross-linking density and strand length (46) for each

species. Statistical significance was determined using GraphPad Prism 6.0

Fabrication of microfluidic devices for measuring cell bending and cell growth.

Masters for the cell bending device and the cell growth/osmotic shock device were

fabricated on separate 3" silicon wafers using SU8 photoresist that was exposed on a

Heidelberg μPG101 mask writer (Heidelberg Instruments, Heidelberg, Germany) and

developed. The bending device (Fig. S1) is a 2-layer device. The first SU8 layer consists

of ~1 μm tall channels for capturing cells. The wafer was then coated with the second

SU8 layer (~25 μm tall) that formed a central flow channel. The flow chamber (Fig. S12),

consists of a single channel with a length of 10 mm, a width of 5 mm, and a height of 50

μm. The volume of the flow chamber is ~10 μL. After developing the masters for both

devices, we used the masters to emboss layers of poly(dimethylsiloxane) (PDMS),

punched holes for inlets and outlets, cleaned the surfaces with tape. If the device was

attached to a glass coverslip it was treated with oxygen plasma and immediately sealed

against a plasma-treated # 1.5 cover glass (24 x 50 mm Fisherbrand) to form a

permanent seal.

Measuring the bending rigidity of *P. mirabilis*, *V. paramaemolyticus*, and *E. coli* cells.

We used streak velocimetry to determine the profile of fluid flow rates in the central

channel of the microfluidic device driven by gravity flow. We added fluorescently

labeled 0.22 μm microspheres (Polysciences) diluted ~1:10000 in ddH_2O containing

0.01% Brij-35 (Sigma) to the microfluidic channel and collected videos of the fluorescent

beads moving through the channel at focal planes 2 μm apart. We analyzed the movies

with custom-written code in Igor Pro 6.37 (WaveMetrics). Briefly, we applied a

Gaussian blur and threshold to each frame and used the thresholded image to establish

a region that was fit to a function based on a 2D Gaussian. We used the image exposure

time and length of the streaks—taking into account the size of the microspheres—to

calculate microsphere velocity. We mapped the velocity profile within the channel by

analyzing several hundred microspheres. We binned the velocity data into a 3D matrix

and fit it to a Poiseuille function, letting the velocity coefficient, height, and width float.

The velocity coefficient was used as an input to the cell bending fitting function.

The system for gravity flow pumping (Fig. S13) consisted of two 6-mL syringes

(Norm-Ject), one 60-mL syringe (BD), one 1-mL syringe (Norm-Ject), a ruler, and a VC-6

channel valve controller (Warner Instruments) connected to a VC-8 mini-valve system

(Warner Instruments) that drives a three-way solenoid valve (The Lee Company).

Syringe #1 (6-mL) was mounted to an immobile post, positioned 40 cm above the table

surface; this syringe was used to load cells. Syringe #2 (6-mL) was positioned 75-cm

above the table surface, was connected to a stage that could be raised and lowered

vertically, and was used to apply flow force in the device. Syringe #3 (60-mL) was

mounted to an immobile post, positioned 75-cm above the surface of the table, and was

used to apply an opposing flow force at the device outlet and to collect spent

media/cells. We attached two-way valves, a blunt-end needle, and tubing to each

syringe, including syringe #4 (1-mL). We joined the two 6-mL syringes using a Y-

junction connector that led to the VC-8-mini-valve system inlet. The outlet of the valve

system was connected to the microfluidic system. A ruler on an immobile post

indicated the '0 position' (no pressure).

For cell bending experiments, we prepared filamentous vegetative cells and

swarmer cells of *P. mirabilis* and *V. parahaemolyticus*; cells were normalized to an optical

density (OD; λ=600 nm) of 1. Prior to starting a measurement, syringes 1 and 2 were

flushed with ddH$_2$O and fresh medium. Syringe 2 was filled with 4 mL of medium,

syringe 3 was filled with 30 mL of ddH$_2$O, and syringe 4 was filled with 0.8 mL of

ddH$_2$O. We added cells to syringe 1 and flowed them through the tubing until they

reached the device outlet. To load the cells into the side channels of the device (from the

central channel), we applied a suction force using the 1-mL syringe. After the side

channels were loaded with cells, we adjusted the height of syringe 2 until no flow

occurred in the device. Syringe 2 was then raised 7 cm from the no-flow position (a

height of 0 cm). We collected images of cells in the channel when syringe 2 was at

positions 0 cm and 7 cm using a Zeiss Axiovert 100 inverted microscope (Zeiss)

equipped with an iXon3 CCD (Andor), a 63X Plan-APOCHROMAT oil objective (Zeiss),

and Micro-Manager 1.4.16. After collecting cell-bending deflections for all loaded cells,

we expelled cells from the side channels, flowed liquid through the device for 15 s, and

reloaded the side channels with new cells.

We analyzed images to determine cell deflection under flow using custom

image-analysis software written in Igor Pro 6.37. The cell-bending model

(Supplementary Information) is a differential equation that lacks an analytical solution

and thus requires calculation of a numerical solution. To determine the bending rigidity

of cells, we wrote custom fitting code in Igor Pro 6.37 that uses a variety of input

parameters, including channel dimensions, fluid velocity, bending rigidity of the cell,

cell radius, and cell length. The function numerically calculates the maximum deflection

based on our model. This function is integrated into a fitting algorithm to find a least-

squares solution to a dataset of maximum deflections versus cell lengths with bending

rigidity as the fitting parameter.

Our model for bending a cell under flow is based on the mechanics model of a

suspended rod or cantilever bending under its own weight (see DERIVATIONS in

Supplementary Information). Our experimental system was similar to that of Amir et al.

(47). Although laminar flow is perpendicular to the long axis of cells in this system, the

lateral deformations of cells that we measured were substantially larger (in our case, as

large as 10 μm; compared to < 1 μm in Amir et al. (47). For this reason, many of the

assumptions of the model presented by Amir et al. are not valid for our dataset. The

model we developed to extract the bending rigidity of the bacterium takes into account

the shape of the laminar flow profile, the angle of the cell against the flow profile, and

the arc length of the cell (longer cells tend to fold over and do not penetrate as deeply

into the flow profile). Our only major assumption is that the force of the bending

moment at a particular point along a cell is not applied as an integration to the end of

the cell, but is rather applied exactly one-half the distance (arc length) to the end of the

cell. We found that models that take into account the full integration of the bending

moment are computationally time-consuming and do not provide a substantial change

in the calculated cell shape (data not shown). Additional data and mathematical

derivations are described in Supplementary Information.

Acknowledgements. We thank Linda McCarter for *V. parahaemolyticus* LM5674,

Suckjoon Jun for the *psulA* plasmid, Cameron Scarlett and Molly Pellitteri-Hahn for

mass spectrometry support, and Julie Last for technical assistance with AFM

measurements. This research was supported by NIH grant 1DP2OD008735-01, National

Science Foundation grant DMR-1121288, a Mao Wisconsin Distinguished Graduate

Fellowship (to M.R.), and an NSF postdoctoral fellowship (#1202622 to P.M.O), and the

Howard Hughes Medical Institute.

References

1. Justice SS, Hunstad DA, Cegelski L, & Hultgren SJ (2008) Morphological plasticity as a bacterial survival strategy. *Nat Rev Microbiol* 6(2):162-168.

2. Rajendram M, *et al.* (2015) Anionic phospholipids stabilize RecA filament bundles in Escherichia coli. *Mol Cell* 60(3):374-384.

3. Copeland MF & Weibel DB (2009) Bacterial Swarming: A Model System for Studying Dynamic Self-assembly. *Soft Matter* 5(6):1174-1187.

4. Kearns DB (2010) A field guide to bacterial swarming motility. *Nat Rev Microbiol* 8(9):634-644.

5. Butler MT, Wang Q, & Harshey RM (2010) Cell density and mobility protect swarming bacteria against antibiotics. *Proc Natl Acad Sci U S A* 107(8):3776-3781.

6. Kim W, Killam T, Sood V, & Surette MG (2003) Swarm-cell differentiation in Salmonella enterica serovar typhimurium results in elevated resistance to multiple antibiotics. *J Bacteriol* 185(10):3111-3117.

7. Overhage J, Bains M, Brazas MD, & Hancock RE (2008) Swarming of Pseudomonas aeruginosa is a complex adaptation leading to increased production of virulence factors and antibiotic resistance. *J Bacteriol* 190(8):2671-2679.

8. Lai S, Tremblay J, & Deziel E (2009) Swarming motility: a multicellular behaviour conferring antimicrobial resistance. *Environ Microbiol* 11(1):126-136.

9. Hay NA, Tipper DJ, Gygi D, & Hughes C (1999) A novel membrane protein influencing cell shape and multicellular swarming of Proteus mirabilis. *J Bacteriol* 181(7):2008-2016.

10. Tuson HH, Copeland MF, Carey S, Sacotte R, & Weibel DB (2013) Flagellum density regulates Proteus mirabilis swarmer cell motility in viscous environments. *J Bacteriol* 195(2):368-377.

11. Vollmer W & Seligman SJ (2010) Architecture of peptidoglycan: more data and more models. *Trends Microbiol* 18(2):59-66.

12. Wang S, Arellano-Santoyo H, Combs PA, & Shaevitz JW (2010) Actin-like cytoskeleton filaments contribute to cell mechanics in bacteria. *Proc Natl Acad Sci U S A* 107(20):9182-9185.

13. Tuson HH, *et al.* (2012) Measuring the stiffness of bacterial cells from growth rates in hydrogels of tunable elasticity. *Mol Microbiol* 84(5):874-891.

14. Wheeler R, *et al.* (2015) Bacterial Cell Enlargement Requires Control of Cell Wall Stiffness Mediated by Peptidoglycan Hydrolases. *Mbio* 6(4):e00660.

15. Loskill P, *et al.* (2014) Reduction of the peptidoglycan crosslinking causes a decrease in stiffness of the Staphylococcus aureus cell envelope. *Biophys J* 107(5):1082-1089.

16. Auer GK, *et al.* (2016) Mechanical Genomics Identifies Diverse Modulators of Bacterial Cell Stiffness. *Cell Syst* 2(6):402-411.

17. Auer GK & Weibel DB (2017) Bacterial Cell Mechanics. *Biochemistry-Us* 56(29):3710-3724.

18. Amir A, Babaeipour F, McIntosh DB, Nelson DR, & Jun S (2014) Bending forces plastically deform growing bacterial cell walls. *Proc Natl Acad Sci U S A* 111(16):5778-5783.

19. Gan L, Chen S, & Jensen GJ (2008) Molecular organization of Gram-negative peptidoglycan. *Proc Natl Acad Sci U S A* 105(48):18953-18957.

20. Patrick JE & Kearns DB (2012) Swarming motility and the control of master regulators of flagellar biosynthesis. *Mol Microbiol* 83(1):14-23.

21. McLaggan D, Naprstek J, Buurman ET, & Epstein W (1994) Interdependence of K+ and glutamate accumulation during osmotic adaptation of Escherichia coli. *J Biol Chem* 269(3):1911-1917.

22. Desmarais SM, *et al.* (2015) High-throughput, highly sensitive analyses of bacterial morphogenesis using ultra performance liquid chromatography. *J Biol Chem* 290(52):31090-31100.

23. Strating H, Vandenende C, & Clarke AJ (2012) Changes in peptidoglycan structure and metabolism during differentiation of Proteus mirabilis into swarmer cells. *Can J Microbiol* 58(10):1183-1194.

24. Yao X, Jericho M, Pink D, & Beveridge T (1999) Thickness and elasticity of gram-negative murein sacculi measured by atomic force microscopy. *J Bacteriol* 181(22):6865-6875.

25. Eberhardt C, Kuerschner L, & Weiss DS (2003) Probing the catalytic activity of a cell division-specific transpeptidase in vivo with beta-lactams. *J Bacteriol* 185(13):3726-3734.

26. Chung HS, *et al.* (2009) Rapid beta-lactam-induced lysis requires successful assembly of the cell division machinery. *Proc Natl Acad Sci U S A* 106(51):21872-21877.

27. Yao Z, Kahne D, & Kishony R (2012) Distinct single-cell morphological dynamics under beta-lactam antibiotics. *Mol Cell* 48(5):705-712.

28. Tuomanen E, Cozens R, Tosch W, Zak O, & Tomasz A (1986) The Rate of Killing of Escherichia-Coli by Beta-Lactam Antibiotics Is Strictly Proportional to the Rate of Bacterial-Growth. *J Gen Microbiol* 132:1297-1304.

29. Spoering AL & Lewis K (2001) Biofilms and planktonic cells of Pseudomonas aeruginosa have similar resistance to killing by antimicrobials. *J Bacteriol* 183(23):6746-6751.

30. Chen CY, *et al.* (2012) Proteus mirabilis urinary tract infection and bacteremia: risk factors, clinical presentation, and outcomes. *J Microbiol Immunol Infect* 45(3):228-236.

31. Burall LS, *et al.* (2004) Proteus mirabilis genes that contribute to pathogenesis of urinary tract infection: Identification of 25 signature-tagged mutants attenuated at least 100-fold. *Infect Immun* 72(5):2922-2938.

32. Trivedi RR, *et al.* (2018) Mechanical genomic studies reveal the role of D-alanine metabolism in Pseudomonas aeruginosa cell stiffness. *mBio* 9:e01340-18.

33. Dobro MJ, *et al.* (2017) Uncharacterized bacterial structures revealed by electron cryotomography. *J Bacteriol* 199:e00100-17.

34. Rojas ER, *et al.* (2018) The outer membrane is an essential load-bearing element in Gram-negative bacteria. *Nature* 559: 617–621

35. Tuson HH, Copeland MF, Carey S, Sacotte R, & Weibel DB (2013) Flagellum Density Regulates Proteus mirabilis Swarmer Cell Motility in Viscous Environments. Journal of Bacteriology 195(2):368-377.

36. Sliusarenko O, Heinritz J, Emonet T, & Jacobs-Wagner C (2011) High-throughput, subpixel precision analysis of bacterial morphogenesis and intracellular spatio-temporal dynamics. Mol Microbiol 80(3):612-627.

37. CLSI (2000) Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically; Approved Standard (Clinical and Laboratory Standards Institute, Wayne, PA) Fifth Ed.

38. Tivol WF, Briegel A, & Jensen GJ (2008) An improved cryogen for plunge freezing. Microsc Microanal 14(5):375-379.

39. Zheng SQ, et al. (2007) UCSF tomography: an integrated software suite for real-time electron microscopic tomographic data collection, alignment, and reconstruction. J Struct Biol 157(1):138-147.

40. Kremer JR, Mastronarde DN, & McIntosh JR (1996) Computer visualization of three-dimensional image data using IMOD. J Struct Biol 116(1):71-76.

41. Nicastro D, et al. (2006) The molecular architecture of axonemes revealed by cryoelectron tomography. Science 313(5789):944-948.

42. Collins TJ (2007) ImageJ for microscopy. Biotechniques 43(1 Suppl):25-30.

43. Turner RD, Hurd AF, Cadby A, Hobbs JK, & Foster SJ (2013) Cell wall elongation mode in Gram-negative bacteria is determined by peptidoglycan architecture. Nat Commun 4.

44. Kuhner D, Stahl M, Demircioglu DD, & Bertsche U (2014) From cells to muropeptide structures in 24 h: peptidoglycan mapping by UPLC-MS. Sci Rep 4:7494.

45. Strating H, Vandenende C, & Clarke AJ (2012) Changes in peptidoglycan structure and metabolism during differentiation of Proteus mirabilis into swarmer cells. Can J Microbiol 58(10):1183-1194.

46. Glauner B (1988) Separation and Quantification of Muropeptides with High-Performance Liquid-Chromatography. Anal Biochem 172(2):451-464.

1   47.   Amir A, Babaeipour F, McIntosh DB, Nelson DR, & Jun S (2014) Bending forces
2   plastically deform growing bacterial cell walls. Proc Natl Acad Sci U S A
3   111(16):5778-5783.

4
5
6
7

FIGURES

Fig. 1. Images demonstrating the flexibility of *P. mirabilis* and *V. parahaemolyticus*

swarmer cells. (*A*) Time-series of *P. mirabilis* swarmer cells in a colony actively moving

across the surface of a 1.5% agarose gel. A representative cell, false colored green, has a

generally straight shape at t=0 s and is bent in half at t=0.98 s. Most of the cells in this

frame are bending during this imaging sequence. (*B*) A time-series of *V. parahaemolyticus*

swarmer cells in a colony actively moving across the surface of a 1.4% agarose gel. A

representative cell (false colored purple) has a generally straight shape at t=0 s and is

bent in half at t=4.02 s.

Fig. 1

Fig. 2. Using a reloadable microfluidic-based assay to determine bacterial cell stiffness.

(*A*) Schematic of the microfluidic channel used to apply a user-defined shear force to

bend filamentous or swarmer cells. Single-sided green arrows depict the flow of fluid

through the central channel; the parabolic flow profile of the fluid is shown. Double-

sided green arrows indicate the vacuum chamber used to load cells into side channels

and to empty the device. (*B*) Cartoon of a flexible bacterium (left) and a stiff bacterium

(right) under flow force (V_{flow}). x_{max} indicates the deflection of cells in the flow; $2r$ = cell

diameter; L = cell length in contact with the flow force. (*C*) Representative images of

filamentous cells of *P. mirabilis* in no flow (left) and flow (right) conditions (top) and *P.*

mirabilis swarmers (bottom). Purple dashed lines indicate the position of a cell tip under

no flow conditions and black dashed lines illustrate the position after flow is applied

using a gravity-fed mechanism. The arrow indicates the direction of fluid flow in the

channel.

A

B

C

Fig. 2

Fig. 3. *P. mirabilis* and *V. parahaemolyticus* swarmer cells have a lower bending rigidity

than vegetative cells. We measured the bending rigidity of *P. mirabilis* and *V.*

parahaemolyticus swarmer cells and filamentous vegetative cells in a microfluidic flow

assay and included measurements of vegetative *E. coli* cells. *P. mirabilis* swarmers

exhibited 15-fold lower bending rigidity than vegetative cells; *V. parahaemolyticus*

swarmers were 3-fold less rigid than vegetative cells. Overexpression of FlhDC (from

the plasmid-encoded *pflhDC*) had little effect on the stiffness of *P. mirabilis* vegetative

and swarmer cells. Error bars represent the 95% confidence interval of a fit to data. *n* >

100 cells, from at least 3 independent experiments.

2 **Fig. 3**

Fig. 4. Swarmer cells increase in cell extension during osmotic shock. We calculated ΔL

as (cell length in water – cell length in 1 M NaCl) and performed a similar calculation

for ΔW, substituting cell width. Cell length indicates length prior to osmotic shock. We

filamented all vegetative cells using aztreonam to grow them to lengths that were

comparable with *P. mirabilis* and *V. parahaemolyticus* swarmers. Lines indicate linear fits

to single-cell measurements (circles) of $n > 100$ cells from at least 3 independent

experiments. (*A*) *P. mirabilis* swarmer cells have an increase in extension (ΔL) under

osmotic shock compared to *E. coli* and *P. mirabilis* vegetative cells. (*B*) *V. parahaemolyticus*

vegetative and swarmer cells have an increase in extension (ΔL) under osmotic shock

compared to *E. coli*. (*C*) *P. mirabilis* swarmer cells have an increase in ΔW compared to *E.*

coli; *P. mirabilis* vegetative cells display a slight decrease in width and increased cell

length. (*D*) There was no observable change in ΔW of *V. parahaemolyticus* swarmer and

vegetative cells.

2 **Fig. 4**

Fig. 5. Alterations in the PG muropeptide composition of *P. mirabilis* and *V.*

parahaemolyticus swarmer cells. (*A*) UPLC-MS data reveal that the muropeptide

composition of *P. mirabilis* and *V. parahaemolyticus* vegetative and swarmer cells differs

slightly in the abundance of monomer, dimer, and anhydrous-terminated saccharides.

(*B*) We observed a relative increase in the amount of anhydrous-containing saccharides

in swarmers consistent with a decrease in polysaccharide strand length. (*C*) There was

no change in PG cross-linking of *P. mirabilis* and *V. parahaemolyticus* vegetative and

swarmer cells, although *V. parahaemolyticus* does display a lower level of cross-linking. *n*

= 3 biological replicates. Error bars represent the standard deviation of the mean.

For (*A-C*), significance was determined via two-way analysis of variance: $*P \leq 0.05$, $**P \leq$

0.01, $***P \leq 0.001$, $****P \leq 0.0001$, ns = not significant ($P > 0.05$).

2   **Fig. 5**

Fig. 6. AFM reveals the PG layer of *P. mirabilis* and *V. parahaemolyticus* swarmer cells is

thinner than in vegetative cells and ECT demonstrates a reduced membrane-to-

membrane distance. (*A*) Sacculi were isolated from cells, dried, and imaged by AFM.

The thickness of swarmer cell PG was reduced compared to vegetative cells. We

analyzed >65 vegetative cells of *E. coli*, *P. mirabilis*, and *V. parahaemolyticus*, >65 *P.*

mirabilis swarmer cells, and 7 *V. parahaemolyticus* swarmer cells. Error bars represent the

standard deviation of the mean. (*B*) Bending rigidity and cell wall thickness display an

approximately exponential relationship (R^2=0.9874). (*C*) Sub-tomogram-averaged ECT

volume images of the *P. mirabilis* vegetative (left) and swarmer (right) cell wall. Two

central slices of sub-tomogram average volume images with normalized image densities

are shown. Yellow dashed line indicates the orientation used for gray-value

measurements. ES, extracellular space; OM, outer membrane; PG, peptidoglycan; IM,

inner membrane; CP, cytoplasm. (*D*) The density profile of sub-tomogram-averaged

ECT volume images reveals a reduced membrane-to-membrane distance in swarmer

cells. The vertical axis is the normalized gray value with the darkest value equal to 0

and the lightest value equal to 1. The red dashed line denotes the average gray value of

the extracellular space and serves as a reference for the background; the blue shaded

area indicates the thickness of the putative PG layer.

Fig. 6

Fig. 7. Swarmer cells are more susceptible to antibiotics that target the cell wall than

vegetative cells. (*A*) Survival of cells treated with 1X MIC of cephalexin after 3 h of

incubation. We define percent survival as (cell count$_{no\ lysis}$/ cell count$_{total}$) x 100. *P.*

mirabilis and *V. parahaemolyticus* swarmer cells exhibit decreased levels (~30%) of

survival compared to vegetative cells; $n \geq 90$ cells from at least two independent

experiments. A similar decrease occurred when *P. mirabilis* was treated with penicillin

G; $n \geq 77$ cells from at least two independent experiments). Error bars represent the

standard deviation of the mean. (*B*) After exposure to cephalexin or penicillin G, the

survival time of *P. mirabilis* and *V. parahaemolyticus* swarmers was 2~3 fold lower than

for vegetative cells. Survival time was determined for ≥ 49 cells that lysed from at least 2

independent experiments. Significance in *A-B* was determined two-tailed t-test: *$P \leq$

0.05, **$P \leq 0.01$, ****$P \leq 0.0001$.

2 **Fig. 7**