Type I-F CRISPR-Cas distribution and array dynamics in *Legionella pneumophila.*

Shayna R. Deecker[1] and Alexander W. Ensminger[1,2]

[1] Department of Biochemistry, University of Toronto, Toronto, Ontario, Canada.

[2] Department of Molecular Genetics, University of Toronto, Toronto, Ontario, Canada.

Running title: *L. pneumophila* type I-F CRISPR-Cas

Abstract

In bacteria and archaea, several distinct types of CRISPR-Cas systems provide adaptive

immunity through broadly similar mechanisms: short nucleic acid sequences derived from

foreign DNA, known as spacers, engage in complementary base pairing with invasive genetic

elements setting the stage for nucleases to degrade the target DNA. A hallmark of type I

CRISPR-Cas systems is their ability to acquire spacers in response to both new and previously

encountered invaders (naïve and primed acquisition, respectively). Our phylogenetic analyses of

47 *L. pneumophila* type I-F CRISPR-Cas systems and their resident genomes suggest that many

of these systems have been horizontally acquired. These systems are frequently encoded on

plasmids and can co-occur with nearly identical chromosomal loci. We show that two such co-

occurring systems are highly protective and undergo efficient primed acquisition in the lab.

Furthermore, we observe that targeting by one system's array can prime spacer acquisition in the

other. Lastly, we provide experimental and genomic evidence for a model in which primed

acquisition can efficiently replenish a depleted type I CRISPR array following a mass spacer

deletion event.

Introduction

Microorganisms have evolved over millions of years to survive in harsh environments, backed in part by immune strategies that protect against foreign genetic elements, such as viral phages and foreign DNA elements (van Houte *et al.* 2016). Clustered regularly interspaced short palindromic repeats (CRISPR) coupled with associated *cas* genes form a potent adaptive immune response in numerous prokaryotic species (Barrangou *et al.* 2007; Brouns *et al.* 2008; Marraffini and Sontheimer 2008). These systems have been classified into six major types, which are further divided into various sub-types, based on their mechanism of action and Cas protein content (Makarova *et al.* 2011; 2015; Koonin *et al.* 2017).

A CRISPR-Cas response to invading DNA occurs in three distinct phases: adaptation, expression and interference (Barrangou *et al.* 2007; Brouns *et al.* 2008; Marraffini and Sontheimer 2008). In the adaptation phase, the CRISPR-Cas system acquires a DNA sequence (spacer) from the invader and integrates it into an array of spacers interspersed with repetitive sequences (Ishino *et al.* 1987; Mojica *et al.* 2000; Jansen, van Embden, Gaastra, and Schouls 2002b; Barrangou *et al.* 2007; Jackson *et al.* 2017). The spacers are generally derived from foreign elements whose infection was unsuccessful, such as defunct phage (Hynes *et al.* 2014), and form the basis of immunological memory for the bacterium. During the expression phase, the array is transcribed and processed to form CRISPR RNA (crRNA) molecules which form a surveillance complex with Cas proteins (Brouns *et al.* 2008; Haurwitz *et al.* 2010). Infection by an invading genetic element initiates the interference step, wherein the surveillance complex recognizes and binds the foreign DNA via base-pairing with the complementary crRNA and

cleaves it, effectively neutralizing the threat to the host (Brouns *et al.* 2008; Jore *et al.* 2011;

Wiedenheft *et al.* 2011; Westra *et al.* 2012).

Despite the sophistication of CRISPR-Cas systems, phages and other foreign DNA

elements can still escape CRISPR-Cas targeting (Samson *et al.* 2013; Stanley and Maxwell

2018). A common mechanism of escape is the accumulation of random mutations within the

foreign element which prevent complementary base pairing with crRNAs during interference

(Deveau *et al.* 2008; Semenova *et al.* 2011; Datsenko *et al.* 2012). CRISPR-Cas systems can

overcome this escape by acquiring new spacers; in fact, imperfect CRISPR targeting often leads

to a highly efficient "primed" acquisition response, providing an intrinsic mechanism to protect

against mutational escape (Swarts *et al.* 2012; Datsenko *et al.* 2012; Savitskaya *et al.* 2013;

Fineran *et al.* 2014; Richter *et al.* 2014). Primed acquisition has been studied in type I-B (Li,

Wang, Zhao, *et al.* 2014; Li, Wang, and Xiang 2014; Li *et al.* 2017), I-C (Rao *et al.* 2017), I-E

(Swarts *et al.* 2012; Datsenko *et al.* 2012; Savitskaya *et al.* 2013; Fineran *et al.* 2014; Xue *et al.*

2015; Jackson *et al.* 2019) and I-F (Richter *et al.* 2014; Vorontsova *et al.* 2015; Staals *et al.*

2016; Heussler *et al.* 2016; Jackson *et al.* 2019) CRISPR-Cas systems. Interference-driven

acquisition, or targeted acquisition, has also been observed in type I-C (Rao *et al.* 2017) and I-F

systems (Staals *et al.* 2016), wherein a primed acquisition response occurs against a target with a

perfect match to a spacer already within the array.

Legionella pneumophila is a gram-negative bacterium and the causative agent of

Legionnaires' disease (Brenner *et al.* 1979). Most isolates possess any of three different

CRISPR-Cas systems: type I-C, I-F and/or II-B (D'Auria *et al.* 2010; Rao *et al.* 2016). Our lab

has recently shown that all three types of CRISPR-Cas systems found in *L. pneumophila* isolates

are active (Rao *et al.* 2016) and we have characterized the targeted acquisition response for the

type I-C system (Rao *et al.* 2017). While much of our work to date has focused on the I-C

systems of *L. pneumophila*, the type I-F systems of this pathogen are highly protective,

remarkably diverse with respect to spacer content, and are frequently found on plasmids –

suggesting that they may be circulated via horizontal gene transfer (Rao *et al.* 2016). In this

study, we perform the first comprehensive phylogenetic analysis of the *L. pneumophila* type I-F

systems and test a model by which horizontal acquisition of a mobile type I-F CRISPR-Cas

system could replenish a collapsed chromosomal array.

Results

Phylogenetic analyses suggest widespread horizontal exchange of *L. pneumophila* type I-F

systems.

In order to explore the hypothesis that plasmid-based type I-F CRISPR-Cas systems in *L.*

pneumophila could be circulated via horizontal gene transfer, we bioinformatically examined the

diversity of type I-F CRISPR-Cas systems within this species. Leveraging both

CRISPRCasFinder (Couvin *et al.* 2018) and CRISPRDetect (Biswas *et al.* 2016), we surveyed

525 draft and 6 completed *L. pneumophila* genomes. In total, we identified 47 *L. pneumophila*

isolates that possessed type I-F systems (Table S1), including 5 that we had described previously

(Rao *et al.* 2016). We next performed two types of phylogenetic analysis: *cas1* phylogeny (Fig

1A), which placed each CRISPR-Cas system into one of six phylotypes; and core-genome

phylogeny (Fig 1B), which reflects the overall relatedness between each of the 47 isolates. A

comparison of the two trees indicates a clear phylogenetic incongruence suggesting that

horizontal acquisition has impacted the distribution of type I-F CRISPR-Cas systems within the

species.

Given the results of these phylogenetic analyses, we next examined the spacer

distribution across each of the arrays to determine the level of array diversification within each

of the six *L. pneumophila* type I-F *cas1* phylotypes. We aligned, clustered, and visualized each

of the 26 distinct *L. pneumophila* type I-F CRISPR arrays using CRISPRStudio (Dion *et al.*

2018) (Fig. 2). These analyses revealed patterns consistent with both spacer acquisition and

spacer loss, suggesting that both processes contribute to *L. pneumophila* type I-F CRISPR array

diversity.

The plasmid and chromosomal Lens CRISPR-Cas systems are active and adaptive

Given the evidence for spacer gain and loss that we observe in *L. pneumophila* type I-F

arrays, (Fig. 2), we next decided to examine array dynamics experimentally. We focused on *L.*

pneumophila str. Lens, which possesses two type I-F CRISPR-Cas systems: one on its

chromosome and one on an endogenous 60 Kb plasmid, pLPL (D'Auria *et al.* 2010; Rao *et al.*

2016). The two systems have a 97.6% Cas protein identity and the repeat units between the

spacers in the CRISPR array differ by only a single nucleotide (Rao *et al.* 2016) (Fig. 3). The

CRISPR arrays themselves are of different lengths (64 spacers for the chromosomal system and

53 for that contained on the pLPL plasmid). Despite a high degree of identity between the Cas

proteins, each array contains a completely unique set of spacers (D'Auria *et al.* 2010; Rao *et al.*

2016). The presence of two remarkably similar I-F systems in *L. pneumophila* str. Lens provided

us with an opportunity to examine targeted spacer acquisition in both of these largely

uncharacterized CRISPR-Cas systems and the interplay between them.

To assess CRISPR-Cas activity in both Lens type I-F systems, we performed an

established transformation efficiency assay (Marraffini and Sontheimer 2008) using two different

targeted protospacer sequences: one matching the most recently acquired spacer and one

matching a spacer from the middle of the array (chromosomal spacer 23 and pLPL spacer 50).

Consistent with active CRISPR-Cas protection, each of the protospacer-containing plasmids

exhibited significant reductions in transformation efficiencies relative to a scrambled protospacer

control (Fig. S1). These relative transformation efficiencies ranged from 1×10^{-2} to 1×10^{-4}, with

the most recently acquired spacers providing ~100-fold greater protection than spacers located in

the middle of each array.

To determine whether spacer acquisition occurs within the context of a perfectly matched

protospacer target, we pooled the transformed populations, passaged them on an automated

liquid handler for 20 generations without selection, extracted their genomic DNA, and screened

the leader end of the CRISPR array by PCR and agarose gel electrophoresis. While the

populations transformed with plasmids encoding either protospacer 23 (chromosome) or

protospacer 50 (pLPL plasmid) exhibited spacer acquisition in both Lens systems (Fig. 4), the

populations transformed with protospacer 1 plasmids exhibited spacer loss, with spacer

acquisition undetectable on a gel (Fig. 4). These data support our bioinformatic analyses, which

indicated that both spacer acquisition and spacer loss contribute to type I-F CRISPR array

diversity in *L. pneumophila* isolates (Fig. 2).

Our observation that the mid-array spacers of each type I-F system provided relatively

modest protection led us to ask whether these spacers could nevertheless drive primed

acquisition of new, more protective spacer sequences. To characterize the patterns of targeted

spacer acquisition in the chromosomal and pLPL CRISPR-Cas systems, we amplified the leader-

proximal region of each CRISPR array from wild-type populations that had been transformed

with the plasmids targeted by their relatively permissive mid-array spacers (chromosomal: spacer

23; pLPL: spacer 50). We Illumina sequenced these PCR products and used an established

bioinformatics pipeline (Rao *et al.* 2017) to identify newly acquired spacer sequences within

each read (Table 1). We then mapped the target of each new spacer to the priming plasmid

(Krzywinski *et al.* 2009) (Fig. 5).

Both the Lens type I-F CRISPR-Cas systems exhibited a biased distribution of acquired

spacers (Fig. 5, S2) consistent with what has been seen previously for the type I-F systems of

Pectobacterium astrosepticum (Richter *et al.* 2014; Staals *et al.* 2016) and *Pseudomonas*

aeruginosa (Vorontsova *et al.* 2015; Heussler *et al.* 2016). The majority of the newly targeted

protospacers clustered around the priming sequence on the targeted plasmid, with the non-

primed strand of DNA (the plus (+) strand) containing ~75% of these protospacers (Fig. 5).

Consistent with observations in other type I-F systems (Richter *et al.* 2014; Staals *et al.* 2016;

Heussler *et al.* 2016), switching the target sequence to the opposite strand led to an acquisition

pattern that mirrored the original distribution observed when the (-) strand contained the targeted

protospacer (Fig. S3). The spacer length distribution and PAM usage were also consistent with

previous observations of other type I-F systems (Mojica *et al.* 2009; Cady *et al.* 2012; Richter *et*

al. 2014; Vorontsova *et al.* 2015; Staals *et al.* 2016) (Fig. S3B, C, S4, S5). Taken together, these

data suggest that spacer acquisition is qualitatively similar between the chromosomal and pLPL

CRISPR-Cas systems.

Permissive targeting by one system can lead to primed acquisition in the other system

Since the chromosomal and pLPL CRISPR-Cas systems function in a very similar

manner during targeted acquisition and share a high degree of homology within their Cas genes

and repeat sequences, we speculated that priming of one system might lead to spacer acquisition

in the other. Specifically, we tested whether introducing a protospacer-containing plasmid

targeted by one CRISPR-Cas system would initiate a primed acquisition response in the second

system. Indeed, this led to efficient spacer acquisition on the second array (Fig. 6), with patterns

largely indistinguishable from what we previously observed on the cognate array (Fig. 5).

Primed repopulation of collapsed arrays

Based on our observations suggesting widespread horizontal inheritance of *L. pneumophila* type

I-F systems (Fig. 1), the diversity of spacer sequences (Fig. 2), and the ability of closely related

systems to prime each other (Fig. 6), we next asked whether coincident CRISPR-Cas might

provide a mechanism for replenishing collapsed chromosomal arrays. Spacer loss is one of

several outcomes when the targeting of a particular sequence becomes detrimental to bacterial

survival. In the lab, this occurs when we artificially "force" the coexistence of an efficiently

targeted plasmid and an active CRISPR-Cas system through selection (Fig. 4). Similar events are

also likely to occur randomly or when CRISPR-Cas systems acquire self-targeting spacers at a

low, but detectable rate (Fig. 7) (Yosef *et al.* 2012; Datsenko *et al.* 2012; Savitskaya *et al.* 2013;

Vorontsova *et al.* 2015; Staals *et al.* 2016; Rao *et al.* 2017). In the most extreme instances, an

entire array of spacers could be lost through recombination between the first and last repeats.

Normally, such a loss could only be reversed by the relatively inefficient mechanism of naïve

spacer acquisition (Yosef *et al.* 2012; Datsenko *et al.* 2012; Savitskaya *et al.* 2013). However, if

such a collapsed array could be restored through primed acquisition driven by a coincident

system, strains with multiple arrays would be inherently more resistant to the catastrophic loss of

CRISPR-Cas protection. Such a model makes a number of predictions (Fig. 7), which we sought

to bioinformatically and experimentally test.

First, we used allelic replacement to generate an *L. pneumophila* Lens strain in which the

entire chromosomal array was replaced by a single copy of its last repeat, mimicking what would

occur after complete spacer loss. Next, we transformed two independently derived array deletion

strains with the pLPL type I-F priming plasmid as above. Using PCR and Illumina sequencing,

we observed robust spacer acquisition in the formerly depleted chromosomal CRISPR array (Fig

8, Table S4), indicating that primed acquisition can replenish the completely collapsed array of a

coincidental type I-F system. As expected, the consensus repeat sequence of this replenished

array adopted the same alternate sequence as the last repeat of the array.

We next examined the CRISPR (repeat) sequences in each of the isolates used in our

earlier bioinformatic analyses for similar evidence of complete collapse followed by

replenishment. In many of the arrays, there is a consensus repeat that is found throughout the

majority of the array, with the last repeat in the array carrying a mutation (Table 2). This has

been observed before in other systems (Jansen, van Embden, Gaastra, and Schouls 2002a; 2002b;

Horvath *et al.* 2008; Lopez-Sanchez *et al.* 2012). However, for a number of the *L. pneumophila*

isolates, including Alcoy, JFIM01, LBAN01, and LBAV01, this is not the case. In these isolates,

their consensus repeat (found throughout the CRISPR array) is identical to their last repeat.

Intriguingly, the sequence that remains is identical to the last repeat found in other type I-F

isolates – what one would predict if they were the product of complete array collapse followed

by subsequent replenishment (Fig. 8).

Discussion

Horizontal gene transfer is a driving force in shaping bacterial biology and pathogenicity

(Ochman *et al.* 2000; Juhas 2013; Hall *et al.* 2017). With respect to *L. pneumophila*, comparative

genomic studies have highlighted the importance of horizontal gene transfer, mobile genetic

elements, and homologous recombination in shaping the bacterium's evolutionary trajectory

(Cazalet *et al.* 2004; 2008; D'Auria *et al.* 2010; Gomez-Valero *et al.* 2011; Sánchez-Busó *et al.*

2014; Gomez-Valero *et al.* 2014; McAdam *et al.* 2014; Burstein *et al.* 2016; David *et al.* 2017).

This aspect of *L. pneumophila* biology extends to the presence and maintenance of plasmid-

based CRISPR-Cas systems (D'Auria *et al.* 2010; Rao *et al.* 2016). Our bioinformatic analyses

suggest that CRISPR-Cas type I-F systems are horizontally distributed in this species (Fig. 1).

These CRISPR arrays have also undergone extensive spacer acquisition and some spacer loss

(Fig. 2). Horizontal acquisition of type I-F systems is well-established in other species. For

instance, in *Vibrio* species, 97% of identified type I-F systems are encoded on mobile genetic

elements (McDonald *et al.* 2019). Like *L. pneumophila*, it's hypothesized these systems have

been acquired by their host through horizontal gene transfer (McDonald *et al.* 2019). Our data,

however, suggest that such transfer is not merely a mechanism by which isolates acquire

CRISPR-Cas protection, but also a potential mechanism to maintain existing defensive

capabilities through the unique spacer dynamics provided by two inter-priming arrays. These

inter-priming arrays can be part of two different systems, as demonstrated by our data, but could

also occur in a system where two (or more) different arrays share a set of *cas* genes (Swarts *et al.*

2012; Datsenko *et al.* 2012; Staals *et al.* 2013; 2014; Majumdar *et al.* 2015; Elmore *et al.* 2015;

Silas *et al.* 2017). For example, in *E. coli* str. K12, it has been shown that two different CRISPR

arrays can be populated by the same set of *cas* genes (Swarts *et al.* 2012; Datsenko *et al.* 2012).

We propose that when a bacterium acquires a second, closely related CRISPR-Cas

system, it gains a mechanism by which depleted CRISPR arrays can be repopulated (Fig. 7). We

have modeled such an event in the lab, made predictions about what the signatures of such

events would be, and provide genomic data to suggest it may have occurred on several occasions

within our collection of sequenced isolates. One obvious line of future investigation would be to

observe whether such patterns are present in other species with mobilized type I CRISPR-Cas

and perhaps absent in instances where CRISPR-Cas is acquired primarily through vertical

inheritance.

Lastly, while we think of spacer loss as a predominantly negative event (loss of

protection), it is likely to play a more nuanced role in the maintenance of CRISPR-Cas activity.

Clearly, array length is the product of a dynamic process whose impact on adaptation,

expression, and interference remains largely unexplored. Many of the type I-F systems in *L.*

pneumophila have different array lengths, ranging from 8 spacers to 129 spacers, with an average

length of 61 spacers (Table 2). Toms and Barrangou have performed a global analysis of class I

CRISPR arrays and found that the average array length for type I-F systems was 33 spacers, with

statistically significant differences between the array lengths of different type I subtypes (Toms

and Barrangou 2017). Accordingly, if spacer acquisition is a driving force in array divergence, it

is likely coupled to spacer loss. Close examination of the mechanisms driving spacer loss in

these systems - and their subsequent impact on CRISPR-Cas functionality - will be crucial to

further testing the model of array diversification in *L. pneumophila*.

Methods and Materials

Bioinformatic analyses

Bioinformatic analyses of the Illumina sequence data were performed as described

previously (Rao *et al.* 2017). Briefly, the raw paired-end reads were merged using FLASH

(Magoc and Salzberg 2011), and any unpaired reads were subsequently quality trimmed using

Trimmomatic (Bolger *et al.* 2014). These processed reads were then combined and analyzed

using a Perl script (available upon request) that annotated existing spacers (S), newly acquired

spacers (X), repetitive sequences (R) and the downstream sequence (D). The newly acquired

spacers were aligned to the priming plasmid, the *L. pneumophila* str. Lens chromosome or the *L.*

pneumophila str. Lens plasmid using BLASTN (Altschul *et al.* 1990). The results from the

BLASTN alignment for the priming plasmid were then processed to obtain coverage per

nucleotide, and plotted on the reference sequence using Circos (Krzywinski *et al.* 2009).

For the bioinformatic analyses of the *L. pneumophila* type I-F system diversity, *L.*

pneumophila draft genomes and completed genomes were downloaded from the European

Nucleotide Archive and NCBI respectively. Type I-F CRISPR-Cas systems were identified using

CRISPRCasFinder (Couvin *et al.* 2018) and CRISPRDetect (Biswas *et al.* 2016). All genomes

with type I-F systems present were annotated using Prokka (Seemann 2014). For the core

genome phylogeny, pan-genome analysis was performed with Roary(Page *et al.* 2015). Both the

cas1 and core genome phylogenies were created with RAxML (Stamatakis 2014) and the

RAxML trees were condensed with MEGA7 (Kumar *et al.* 2016). CRISPR array alignments,

clustering and visualization were performed with CRISPRStudio (Dion *et al.* 2018).

Bacterial strains, plasmids and oligos used

The bacterial strains and plasmids used in this study are listed in supplementary table 2,

and the oligos used in this study are listed in supplementary table 3.

The priming plasmids were created by annealing oligos (see Table S3) to create the

protospacer insert with the canonical GG PAM (Mojica *et al.* 2009; Cady *et al.* 2012; Richter *et*

al. 2014; Vorontsova *et al.* 2015; Staals *et al.* 2016) and subsequently ligating the insert into an

ApaI/PstI-digested pMMB207 vector (Solomon *et al.* 2000). The scrambled control plasmid was

created in the same manner, except it contained a 32-nt scrambled sequence in place of a targeted

protospacer sequence.

Lens (chromsome) array deletion mutants were generated through allelic replacement.

Briefly, 1 kb of DNA upstream and 2 kb downstream of the CRISPR array were amplified by

PCR and stitched together to create an insert where the entire array, save for the last repeat, was

deleted. The insert was ligated into a pJB4648 plasmid. Overnight cultures of *L. pneumophila* str.

Lens were grown in ACES-buffered yeast extract (AYE) medium to an OD_{600} of ~4.0 using two-

day patches that were grown on charcoal-buffered ACES yeast extract (CYE) plates. Pellets from

4.0 ODU of culture underwent three washing steps: twice with 1 mL of ice-cold ultrapure water

and once with 1 mL of ice-cold 10% glycerol. The pellet was then re-suspended in 200 uL of ice-

cold 10% glycerol and 400 ng of plasmid was added to the sample. The solution was transferred

to an ice-cold electroporation cuvette with a 2 mm gap and electroporated with the following

settings: 2500 kV, 600 Ω and 25 mF. After electroporation, 800 uL of AYE medium was added

to each sample and the samples recovered for 3 hours at 37°C at 600 RPM in a shaking

incubator. The samples were plated on CYE plates supplemented with 15 μg mL^{-1} of gentamycin

and incubated at 37°C for 3 days. Surviving colonies were patched onto CYE + gentamycin

plates and grown at 37°C for 2 days. Patches were subsequently struck onto CYE plates

supplemented with sucrose and incubated at 37°C for 3 days. Surviving colonies were patched

onto CYE + sucrose plates, grown at 37°C for 2 days and screened by PCR to confirm the

deletion. Two independent clones were Illumina sequenced and used for subsequent

replenishment assays.

Transformation efficiency assay and population pool generation

The transformation efficiency assay was performed as we have described previously (Rao *et al.* 2016) with some modifications. Briefly, *L. pneumophila* str. Lens was electroporated as described above. The samples were plated in a dilution series on CYE plates supplemented with 5 μg mL^{-1} of chloramphenicol and incubated at 37°C for 3 days. The relative transformation efficiency for each targeted plasmid was calculated as a percentage of the transformation efficiency obtained from the scrambled control plasmid. Three biological replicates were performed for each transformation efficiency assay.

Population pools for spacer acquisition experiments were generated by mixing together ≥ 50 colonies per population from a newly transformed wild-type strain on CYE plates supplemented with 5 μg mL^{-1} of chloramphenicol using AYE medium supplemented with 5 μg mL^{-1} of chloramphenicol. Population pools were made in triplicate for each transformed plasmid.

Serial passaging on an automated liquid handler

The serial passaging of transformed *L. pneumophila* str. Lens populations was performed as described previously (Rao *et al.* 2016). Briefly, overnight cultures of the population pools in AYE medium supplemented with 5 μg mL^{-1} of chloramphenicol for plasmid maintenance were grown to an OD$_{600}$ of ~2.0. The culture was then back diluted to an OD$_{600}$ of ~0.0625 and grown in a flat-bottom 48-well plate (Greiner) in a shaking incubator at 37°C. A Freedom Evo 100 liquid handler (Tecan) connected to an Infinite M200 Pro plate reader (Tecan) measured the optical density of the plate every 20 minutes, until an OD$_{600}$ of ~2.0 was reached. The cultures were then automatically back diluted to an OD$_{600}$ of ~0.0625 in the adjacent well to continue growth, and the remaining culture was transferred to a 48-well plate that was kept at 4°C. In this

manner, each saved culture represented ~5 generations of growth. The passaging was done

without selection in AYE medium to allow for plasmid loss during passaging.

Genomic DNA extraction, PCR and agarose gel screening

Genomic DNA was extracted from the passaged cultures and the parental chromosome

Lens array deletion strains using a Machery-Nagel Nucleospin Tissue kit according to the

manufacturer's protocol. The extracted samples from passaged cultures were used as a template

in a 30-cycle PCR reaction with Econotaq Polymerase (Lucigen) to amplify the leader end of the

CRISPR array using primers listed in Table S3. The PCR products were then separated on a 3%

agarose gel to determine if spacer acquisition (or spacer loss) had occurred based on the presence

of an upper (or lower) band relative to the control sample.

Nextera library prep and Illumina sequencing

The extracted genomic DNA from passaged cultures was prepared for leader-end array

sequencing by performing a 20-cycle PCR using Kapa HiFi Polymerase (Kapa Biosystems) and

the primers listed in Table S3. The PCR products were purified using a Machery-Nagel

Nucleospin Gel and PCR Clean-up kit as per the manufacturer's instructions and normalized to 1

ng using the Invitrogen Quant-iT PicoGreen dsDNA assay. The DNA was then tagmented using

a Nextera XT tagmentation kit as per the manufacturer's instructions. The tagmented products

were sequenced with a paired-end (2 x 150 bp) sequencing run on an Illumina NextSeq platform

at the Centre for the Analysis of Genome Evolution and Function (CAGEF) at the University of

Toronto.

The genomic DNA from the parental chromosome Lens array deletion strains was

normalized to 1 ng using PicoGreen. The DNA was then tagmented using a Nextera XT

tagmentation kit as per the manufacturer's instructions. The tagmented products were sequenced

with paired-end (2 x 150 bp) sequencing in-house on an Illumina MiniSeq platform.

Data Accessibility

The raw Illumina reads have been deposited into the NCBI sequence read archive under

the BioProject PRJNA433194.

Acknowledgements

The authors thank Griffin Deecker (a volunteer high school student) for his assistance in

bioinformatically examining the diversity of I-F repeat sequences in *L. pneumophila*, Chitong

Rao for his contributions to experimental design and Kamran Rizzolo for discussions regarding

phylogenetic analyses. We also thank the Center for the Analysis of Genome Evolution and

Function (CAGEF) at the University of Toronto for performing Illumina sequencing. We thank

members of the Ensminger laboratory for their suggestions and careful reading of the

manuscript, in particular Beth Nicholson and Malene Urbanus. SRD is supported by a fellowship

from the Department of Biochemistry, University of Toronto. This work was supported by a

Project Grant from the Canadian Institutes of Health Research (PHT-148819), the Connaught

Fund (NR-2015-16), and an infrastructure grant from the Canada Foundation for Innovation and

the Ontario Research Fund (30364) to AWE.

Figures

Figure 1 | Phylogenetic analysis of type I-F CRISPR-Cas system diversity in *L.*

***pneumophila* reveals a horizontal distribution across isolates.** *L. pneumophila* draft and

completed genomes were analyzed using CRISPRDetect and CRISPRCasFinder to identify type

I-F systems present within the genomes. Isolates that possessed type I-F systems were subjected

to phylogenetic analyses of their *cas1* gene and core genome. The core genome alignment for the

examined isolates was determined by Roary. Isolate names are colour-coded based on the *cas1*

gene phylogeny to allow for comparison between the analyses. The isolates used in this analysis

and their accession numbers can be found in supplementary table 1. **A)** The *cas1* gene phylogeny

for *L. pneumophila* isolates with type I-F systems reveals six different cas1 groups. **B)** The core

genome phylogeny of the examined isolates is not congruent with the *cas1* phylogeny,

suggesting that the type I-F system was horizontally distributed rather than vertically inherited.

Note that Lens possesses two type I-F CRISPR-Cas systems, one on a plasmid (group Θ) and

one on its chromosome (group Φ). Also note that isolates with 100% nucleotide identity in their

core genome, as well as a shared *cas* gene group and redundant CRISPR array, have been

collapsed to one representative in the phylogeny. Isolates with unique CRISPR arrays are listed.

Collapsed isolates include:

+: FJEK01, FJEL01, FJEM01, FJEN01, FJEO01, FJEP01, FJEQ01, FJER01, FJES01, FJET01,

FJEU01, FJEV01, FJEX01, FJEY01, FJEZ01, FJFA01

++: FJBI01, FJBU01

+++: LAXR01

++++: FJEI01

+++++: FFI105, FFI337

Figure 2 | CRISPR array analysis suggests that spacer acquisition and spacer loss has

contributed to array diversification in *L. pneumophila* type I-F systems. *L. pneumophila*

isolates were subjected to a CRISPRStudio analysis to look at the spacer composition of their

CRISPR arrays. Grey boxes denote unique spacers, coloured boxes denote shared spacers and a

dashed line denotes spacer loss. The isolate colour coding scheme is based on the *cas1* grouping

from figure 1. Denoted isolates within the same *cas* gene group have 100% nucleotide identity in

their core genomes. Note that *cas1* group β appears to have spacer rearrangements present,

resulting in an imperfect alignment between the arrays.

Figure 3 | A comparison of the Lens chromosome and pLPL CRISPR-Cas systems. A) The

overall pairwise amino acid identity across the Cas proteins of the two systems is approximately

97%, with individual pairwise Cas protein identities ranging from 96% to 99%. **B)** A

CRISPRStudio alignment of the arrays for two systems shows completely unique spacer content

and a differing number of spacers. **C)** Analysis of the repeat sequences shows that the Lens

pLPL and chromosome systems have one SNP between their consensus repeats, in addition to

possessing a mutated last repeat in their arrays. Mutations are denoted in red.

Figure 4 | Spacer acquisition and spacer loss in both Lens CRISPR-Cas systems. Spacer

acquisition and loss were analyzed using a PCR-based screen where the leader-end of the

CRISPR array for both the control samples and the transformed samples was amplified with

system-specific primers to differentiate between the chromosomal Lens and the plasmid Lens

arrays and visualized on an agarose gel. Products from the transformed samples were compared

to the control, which contained untransformed genomic DNA. Bands representing spacer

acquisition and loss are indicated.

Figure 5 | Characterization of self-primed spacer acquisition in the two Lens CRISPR-Cas

systems. Bacterial transformants with targeted plasmids were passaged for 20 generations

without antibiotic selection to enrich for spacer acquisition; the leader end of the CRISPR array

was amplified and the amplicons were Illumina sequenced. Newly targeted protospacers were

obtained from the raw reads using an in-house bioinformatics pipeline and visualized with

Circos. The arrows in the simplified array schematic show the primer location for PCR

amplification prior to sequencing, while the star denotes the priming spacer. All data are the

average of three biological replicates. **A)** The distribution of newly targeted protospacers mapped

to the priming plasmid on the Circos plot reveals a strand bias in self-primed spacer acquisition

within the Lens (pLPL) CRISPR-Cas system. The height of the bars indicates the number of

spacers mapped to the position on the plasmid, up to 5% of total acquired spacers. **B)** The

distribution of newly targeted protospacers mapped to the priming plasmid shows a similar

pattern of self-primed spacer acquisition within the Lens (chromosome) CRISPR-Cas system to

that of its pLPL CRISPR-Cas counterpart. Labelling as in **(A)**.

Figure 6 | Characterization of cross-primed spacer acquisition in the two Lens CRISPR-

Cas systems. The experimental set-up is the same as described in figure 5. All data are the

average of three biological replicates. The distribution of newly targeted protospacers mapped to

the priming plasmid reveals cross-priming between the Lens (chromosome) and Lens (pLPL)

CRISPR-Cas systems. **(A)** shows Lens (chromosome) primed, Lens (pLPL) array examined

while **(B)** shows Lens (pLPL) primed, Lens (chromosome) array examined.

Figure 7 | Cross-priming between two similar CRISPR-Cas systems can repopulate a

collapsed CRISPR array. A bacterium with a CRISPR-Cas system could undergo a mass

spacer deletion event through homologous recombination between the first (black) and last

(white) repeat sequences. The remaining locus carries a single repeat (white). While such

"catastrophic collapse" events are likely to occur randomly at a certain rate, a driver of such a

collapse could be the acquisition of a self-targeting spacer (yellow), selecting for spacer loss.

Horizontal acquisition of a second CRISPR-Cas array (e.g. on a plasmid) is a first step towards

replenishing the primary array. If cross-priming can occur between this secondary array and the

collapsed array, the original CRISPR array is replenished, but bears an observable molecular scar

- conversion of all the repeats to the sequence of the last repeat (white).

Figure 8 | An experimentally depleted chromosomal CRISPR array can be replenished

through the activity of a plasmid-based array. A) Replenishment of a depleted array in the

lab. Allelic replacement was used to remove the entire array from the chromosomal Lens I-F

system, leaving behind a single, last repeat sequence (see materials and methods). This strain

was then transformed with a plasmid targeted by the pLPL (Lens plasmid-based) I-F array

previously shown to drive primed acquisition (pLPL protospacer 50; see Figure 4). Spacer

acquisition by the empty chromosomal array was analyzed using a PCR based screen where the

leader-end of the CRISPR array was amplified and visualized on an agarose gel. Products from

the transformed samples (samples 2 and 4) were compared to untransformed controls (samples 1

and 3). Samples 1 and 2 are from depleted array clone #1 and samples 3 and 4 are from depleted

array clone #2. **B)** Repeat signatures of depletion/replenishment. The repeat structure of the

experimentally replenished Lens CRISPR arrays resembles that of *L. pneumophila* str. Alcoy,

suggesting a similar array depletion/replenishment event may have occurred within the Alcoy

lineage. The frequency of acquiring one new spacer versus two new spacers following

replenishment in the Lens array depletion isolates was determined by Illumina sequencing.

References

Altschul, S. F., W. Gish, W. Miller, E. W. Myers, and D. J. Lipman, 1990 Basic Local
Alignment Search Tool. Journal of Molecular Biology 215: 403–410.

Barrangou, R., C. Fremaux, H. Deveau, M. Richards, P. Boyaval *et al.*, 2007 CRISPR provides
acquired resistance against viruses in prokaryotes. Science 315: 1709–1712.

Biswas, A., R. H. J. Staals, S. E. Morales, P. C. Fineran, and C. M. Brown, 2016 CRISPRDetect:
A flexible algorithm to define CRISPR arrays. BMC Genomics 1–14.

Bolger, A. M., M. Lohse, and B. Usadel, 2014 Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics 30: 2114–2120.

Brenner, D. J., A. G. Steigerwalt, and J. E. McDade, 1979 Classification of the Legionnaires'
Disease Bacterium: Legionella pneumophila, genus novum, species nova, of the Family
Legionellaceae, familia nova. Annals of Internal Medicine 90: 656–658.

Brouns, S. J. J., M. M. Jore, M. Lundgren, E. R. Westra, R. J. H. Slijkhuis *et al.*, 2008 Small
CRISPR RNAs guide antiviral defense in prokaryotes. Science 321: 960–964.

Burstein, D., F. Amaro, T. Zusman, Z. Lifshitz, O. Cohen *et al.*, 2016 Genomic analysis of 38
Legionella species identifies large and diverse effector repertoires. Nature Publishing Group
48: 167–175.

Cady, K. C., J. Bondy-Denomy, G. E. Heussler, and A. R. Davidson, 2012 The CRISPR/Cas
adaptive immune system of *Pseudomonas aeruginosa* mediates resistance to naturally
occurring and engineered phages. Journal of Bacteriology 194: 5728–5738.

Cazalet, C., S. Jarraud, Y. Ghavi-Helm, F. Kunst, P. Glaser *et al.*, 2008 Multigenome analysis
identifies a worldwide distributed epidemic Legionella pneumophila clone that emerged
within a highly diverse species. Genome Research 18: 431–441.

Cazalet, C., C. Rusniok, H. Brüggemann, N. Zidane, A. Magnier *et al.*, 2004 Evidence in the
Legionella pneumophila genome for exploitation of host cell functions and high genome
plasticity. Nat Genet 36: 1165–1173.

Couvin, D., A. Bernheim, C. Toffano-Nioche, M. Touchon, J. Michalik *et al.*, 2018
CRISPRCasFinder, an update of CRISRFinder, includes a portable version, enhanced
performance and integrates search for Cas proteins. Nucleic Acids Research 99: 7536–6.

D'Auria, G., N. Jimenez-Hernandez, F. Peris-Bondia, A. Moya, and A. Latorre, 2010 Legionella
pneumophila pangenome reveals strain-specific virulence factors. BMC Genomics 11: 181.

Datsenko, K. A., K. Pougach, A. Tikhonov, B. L. Wanner, K. V. Severinov *et al.*, 2012
Molecular memory of prior infections activates the CRISPR/Cas adaptive bacterial immunity
system. Nature Communications 3: 945–7.

David, S., L. Sánchez-Busó, S. R. Harris, P. Marttinen, C. Rusniok *et al.*, 2017 Dynamics and
impact of homologous recombination on the evolution of Legionella pneumophila (X.
Didelot, Ed.). PLoS Genet 13: e1006855–21.

Deveau, H., R. Barrangou, J. E. Garneau, J. Labonte, C. Fremaux *et al.*, 2008 Phage response to
CRISPR-encoded resistance in *Streptococcus thermophilus*. Journal of Bacteriology 190:
1390–1400.

Dion, M., S. Labrie, S. Shah, and S. Moineau, 2018 CRISPRStudio: A User-Friendly Software
for Rapid CRISPR Array Visualization. Viruses 10: 602–11.

Elmore, J., T. Deighan, J. Westpheling, R. M. Terns, and M. P. Terns, 2015 DNA targeting by
the type I-G and type I-A CRISPR–Cas systems of Pyrococcus furiosus. Nucleic Acids
Research gkv1140–11.

Fineran, P. C., M. J. H. Gerritzen, M. Suarez-Diez, T. Kunne, J. Boekhorst *et al.*, 2014
Degenerate target sites mediate rapid primed CRISPR adaptation. Proc Natl Acad Sci USA
111: E1629–E1638.

Gomez-Valero, L., C. Rusniok, S. Jarraud, B. Vacherie, Z. Rouy *et al.*, 2011 Extensive
recombination events and horizontal gene transfer shaped the Legionella pneumophila
genomes. BMC Genomics 12: 536.

Gomez-Valero, L., C. Rusniok, M. Rolando, M. Neou, D. Dervins-Ravault *et al.*, 2014
Comparative analyses of Legionella species identifies genetic features of strains causing
Legionnaires disease. Genome Biology 1–21.

Hall, J. P. J., M. A. Brockhurst, and E. Harrison, 2017 Sampling the mobile gene pool:
innovation via horizontal gene transfer in bacteria. Phil. Trans. R. Soc. B 372: 20160424–10.

Haurwitz, R. E., M. Jinek, B. Wiedenheft, K. Zhou, and J. A. Doudna, 2010 Sequence- and
structure-specific RNA processing by a CRISPR endonuclease. Science 329: 1355–1358.

Heussler, G. E., J. L. Miller, C. E. Price, A. J. Collins, and G. A. O'Toole, 2016 Requirements
for Pseudomonas aeruginosa Type I-F CRISPR-Cas Adaptation Determined Using a Biofilm
Enrichment Assay (T. J. Silhavy, Ed.). Journal of Bacteriology 198: 3080–3090.

Horvath, P., D. A. Romero, A. C. Coute-Monvoisin, M. Richards, H. Deveau *et al.*, 2008
Diversity, activity, and evolution of CRISPR loci in *Streptococcus thermophilus*. Journal of
Bacteriology 190: 1401–1412.

Hynes, A. P., M. Villion, and S. Moineau, 2014 Adaptation in bacterial CRISPR-Cas immunity
can be driven by defective phages. Nature Communications 5: 1–6.

Ishino, Y., H. Shinagawa, K. Makino, M. Amemura, and A. Nakata, 1987 Nucleotide sequence
of the *iap* gene, responsible for alkaline phosphatase isozyme conversion in *Escherichia coli*,
and identification of the gene product. Journal of Bacteriology 169: 5429–5433.

Jackson, S. A., N. Birkholz, L. M. Malone, and P. C. Fineran, 2019 Imprecise Spacer
Acquisition Generates CRISPR-Cas Immune Diversity through Primed Adaptation. Cell
Host and Microbe 1–16.

Jackson, S. A., R. E. McKenzie, R. D. Fagerlund, S. N. Kieper, P. C. Fineran *et al.*, 2017
CRISPR-Cas: Adapting to change. Science 356: eaal5056–11.

Jansen, R., J. D. van Embden, W. Gaastra, and L. M. Schouls, 2002a Identification of a novel
family of sequence repeats among prokaryotes. OMICS 6: 23–33.

Jansen, R., J. D. van Embden, W. Gaastra, and L. M. Schouls, 2002b Identification of genes that
are associated with DNA repeats in prokaryotes. Molecular Microbiology 43: 1565–1575.

Jore, M. M., M. Lundgren, E. van Duijn, J. B. Bultema, E. R. Westra *et al.*, 2011 Structural basis
for CRISPR RNA-guided DNA recognition by Cascade. Nature Publishing Group 18: 529–
536.

Juhas, M., 2013 Horizontal gene transfer in human pathogens. Critical Reviews in Microbiology
41: 101–108.

Koonin, E. V., K. S. Makarova, and F. Zhang, 2017 Diversity, classification and evolution of
CRISPR-Cas systems. Current Opinion in Microbiology 37: 67–78.

Krzywinski, M., J. Schein, I. Birol, J. Connors, R. Gascoyne *et al.*, 2009 Circos: an information
aesthetic for comparative genomics. Genome Research 19: 1639–1645.

Kumar, S., G. Stecher, and K. Tamura, 2016 MEGA7: Molecular Evolutionary Genetics
Analysis Version 7.0 for Bigger Datasets. Molecular Biology and Evolution 33: 1870–1874.

Li, M., L. Gong, D. Zhao, J. Zhou, and H. Xiang, 2017 The spacer size of I-B CRISPR is
modulated by the terminal sequence of the protospacer. Nucleic Acids Research 45: 4642–
4654.

Li, M., R. Wang, and H. Xiang, 2014 *Haloarcula hispanica* CRISPR authenticates PAM of a
target sequence to prime discriminative adaptation. Nucleic Acids Research 42: 7226–7235.

Li, M., R. Wang, D. Zhao, and H. Xiang, 2014 Adaptation of the *Haloarcula hispanica* CRISPR-
Cas system to a purified virus strictly requires a priming process. Nucleic Acids Research
42: 2483–2492.

Lopez-Sanchez, M.-J., E. Sauvage, V. Da Cunha, D. Clermont, E. Ratsima Hariniaina *et al.*,
2012 The highly dynamic CRISPR1 system of *Streptococcus agalactiae* controls the
diversity of its mobilome. Molecular Microbiology 85: 1057–1071.

Magoc, T., and S. L. Salzberg, 2011 FLASH: fast length adjustment of short reads to improve
genome assemblies. Bioinformatics 27: 2957–2963.

Majumdar, S., P. Zhao, N. T. Pfister, M. Compton, S. Olson *et al.*, 2015 Three CRISPR-Cas
immune effector complexes coexist in Pyrococcus furiosus. RNA 21: 1147–1158.

Makarova, K. S., D. H. Haft, R. Barrangou, S. J. J. Brouns, E. Charpentier *et al.*, 2011 Evolution
and classification of the CRISPR–Cas systems. Nature Reviews Microbiology 9: 467–477.

Makarova, K. S., Y. I. Wolf, O. S. Alkhnbashi, F. Costa, S. A. Shah *et al.*, 2015 An updated
evolutionary classification of CRISPR–Cas systems. Nature Reviews Microbiology 13: 722–
736.

Marraffini, L. A., and E. J. Sontheimer, 2008 CRISPR interference limits horizontal gene
transfer in Staphylococci by targeting DNA. Science 322: 1843–1845.

McAdam, P. R., C. W. Vander Broek, D. S. Lindsay, M. J. Ward, M. F. Hanson *et al.*, 2014
Gene flow in environmental Legionella pneumophila leads to genetic and pathogenic
heterogeneity within a Legionnaires   disease outbreak. Genome Biology 15: 1–10.

McDonald, N. D., A. Regmi, D. P. Morreale, J. D. Borowski, and E. F. Boyd, 2019 CRISPR-Cas
systems are present predominantly on mobile genetic elements in Vibrio species. 1–23.

Mojica, F. J. M., C. Díez-Villaseñor, J. García-Martínez, and C. Almendros, 2009 Short motif
sequences determine the targets of the prokaryotic CRISPR defence system. Microbiology
155: 733–740.

Mojica, F. J. M., C. Díez-Villaseñor, E. Soria, and G. Juez, 2000 Biological significance of a
family of regularly spaced repeats in the genomes of archaea, bacteria and mitochondria.
Molecular Microbiology 36: 244–246.

Ochman, H., J. G. Lawrence, and E. A. Groisman, 2000 Lateral gene transfer and the nature of
bacterial innovation. Nature 405: 299–304.

Page, A. J., C. A. Cummins, M. Hunt, V. K. Wong, S. Reuter *et al.*, 2015 Roary: rapid large-
scale prokaryote pan genome analysis. Bioinformatics 31: 3691–3693.

Rao, C., D. Chin, and A. W. Ensminger, 2017 Priming in a permissive type I-C CRISPR–Cas
system reveals distinct dynamics of spacer acquisition and loss. RNA 23: 1525–1538.

Rao, C., C. Guyard, C. Pelaz, J. Wasserscheid, J. Bondy-Denomy *et al.*, 2016 Active and
adaptive *Legionella* CRISPR-Cas reveals a recurrent challenge to the pathogen. Cellular
Microbiology 18: 1319–1338.

Richter, C., R. L. Dy, R. E. McKenzie, B. N. J. Watson, C. Taylor *et al.*, 2014 Priming in the
Type I-F CRISPR-Cas system triggers strand-independent spacer acquisition, bi-
directionally from the primed protospacer. Nucleic Acids Research 42: 8516–8526.

Samson, J. E., A. H. Magadan, M. Sabri, and S. Moineau, 2013 Revenge of the phages: defeating
bacterial defences. Nature Publishing Group 11: 675–687.

Savitskaya, E. E., E. Semenova, V. Dedkov, A. Metlitskaya, and K. V. Severinov, 2013 High-
throughput analysis of type I-E CRISPR/Cas spacer acquisition in *E. coli*. RNA Biology 10:
716–725.

Sánchez-Busó, L., I. Comas, G. Jorques, and F. González-Candelas, 2014 Recombination drives
genome evolution in outbreak-related Legionella pneumophila isolates. Nat Genet 46: 1205–
1211.

Seemann, T., 2014 Prokka: rapid prokaryotic genome annotation. Bioinformatics 30: 2068–2069.

Semenova, E., M. M. Jore, K. A. Datsenko, A. Semenova, E. R. Westra *et al.*, 2011 Interference
by clustered regularly interspaced short palindromic repeat (CRISPR) RNA is governed by a
seed sequence. Proc Natl Acad Sci USA 108: 10098–10103.

Silas, S., P. Lucas-Elio, S. A. Jackson, A. A.-C. Elife, 2017, 2017 Type III CRISPR-Cas systems
can provide redundancy to counteract viral escape from type I systems. eLife e27601.

Solomon, J. M., A. Rupper, J. A. Cardelli, and R. R. Isberg, 2000 Intracellular growth of
Legionella pneumophila in *Dictyostelium discoideum*, a system for genetic analysis of host-
pathogen interactions. Infect. Immun. 68: 2939–2947.

Staals, R. H. J., Y. Agari, S. Maki-Yonekura, Y. Zhu, D. W. Taylor *et al.*, 2013 Structure and
Activity of the RNA-Targeting Type III-B CRISPR-Cas Complex of Thermus thermophilus.
Molecular Cell 52: 135–145.

Staals, R. H. J., S. A. Jackson, A. Biswas, S. J. J. Brouns, C. M. Brown *et al.*, 2016 Interference-
driven spacer acquisition is dominant over naive and primed adaptation in a native CRISPR-
Cas system. Nature Communications 7: 1–13.

Staals, R. H. J., Y. Zhu, D. W. Taylor, J. E. Kornfeld, K. Sharma *et al.*, 2014 RNA Targeting by
the Type III-A CRISPR-Cas Csm Complex of Thermus thermophilus. Molecular Cell 56:
518–530.

Stamatakis, A., 2014 RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. Bioinformatics 30: 1312–1313.

Stanley, S. Y., and K. L. Maxwell, 2018 Phage-Encoded Anti-CRISPR Defenses. Annu. Rev.
Genet. 52: 445–464.

Swarts, D. C., C. Mosterd, M. W. J. van Passel, and S. J. J. Brouns, 2012 CRISPR interference
directs strand specific spacer acquisition (I. Mokrousov, Ed.). PLoS ONE 7: e35888–7.

Toms, A., and R. Barrangou, 2017 On the global CRISPR array behavior in class I systems.
Biology Direct 12: 1–12.

van Houte, S., A. Buckling, and E. R. Westra, 2016 Evolutionary ecology of prokaryotic immune
mechanisms. Microbiol. Mol. Biol. Rev. 80: 745–763.

Vorontsova, D., K. A. Datsenko, S. Medvedeva, J. Bondy-Denomy, E. E. Savitskaya *et al.*, 2015
Foreign DNA acquisition by the I-F CRISPR–Cas system requires all components of the
interference machinery. Nucleic Acids Research 43: 10848–10860.

Westra, E. R., P. B. G. van Erp, T. Künne, S. P. Wong, R. H. J. Staals *et al.*, 2012 CRISPR
immunity relies on the consecutive binding and degradation of negatively supercoiled
invader DNA by Cascade and Cas3. Molecular Cell 46: 595–605.

Wiedenheft, B., E. van Duijn, J. B. Bultema, S. P. Waghmare, K. Zhou *et al.*, 2011 RNA-guided
complex from a bacterial immune system enhances target recognition through seed sequence
interactions. Proc Natl Acad Sci USA 108: 10092–10097.

Xue, C., A. S. Seetharam, O. Musharova, K. V. Severinov, S. J. J. Brouns *et al.*, 2015 CRISPR
interference and priming varies with individual spacer sequences. Nucleic Acids Research
43: 10831–10847.

Yosef, I., M. G. Goren, and U. Qimron, 2012 Proteins and DNA elements essential for the
CRISPR adaptation process in *Escherichia coli*. Nucleic Acids Research 40: 5569–5576.

TABLE 1

The number of acquired spacers for the Lens (pLPL) and Lens (chromosome) CRISPR-Cas

systems and their respective targets during self-priming.

System	Number of acquired spacers[a]	% map to priming plasmid[a]	% map to Lens (chromosome)[a]	% map to Lens (pLPL)[a]	% unknown[a]
Lens (pLPL)	78,753	98.38	0.05	0.01	1.57
Lens (chromsosome)	86,789	98.66	0.01	0.00	1.33

[a] Data are the average of 3 biological replicates.

TABLE 2

The repeat sequences of *L. pneumophila* type I-F CRISPR-Cas systems.[a]

Strain	# of spacers in the array	Consensus Repeat (5'-3')[b]	Mutated last repeat (5'-3')[b]
Lens (chromosome)	64	GTTCACTGCCGCACAGG CAGCTTAGAAA	GTTCACTGCCGCACAGG CAGCTTAGAA**G**
FFI104	55	GTTCACTGCCGCACAGG CAGCTTAGAAA	GTTCACTGCCGCACAGG CAGCTTAGAA**G**
Alcoy	56	GTTCACTGCCGCACAGG CAGCTTAGAA**G**	--
Lens (pLPL)	53	GTTCACTGCCG**T**ACAGG CAGCTTAGAAA	GTTCACTGCCG**T**ACAGG CAGCTTAGAA**G**
Missisauga-2006 (plasmid)	74	GTTCACTGCCG**T**ACAGG CAGCTTAGAAA	GTTCACTGCCG**T**ACAGG CAGCTTAGAA**G**
C8_S (plasmid)	24	GTTCACTGCCG**T**ACAGG CAGCTTAGAAA	GTTCACTGCCG**T**ACAGG CAGCTTAGAA**G**
Calgary2017	8	GTTCACTGCCG**T**ACAGG CAGCTTAGAAA	GTTCACTGCCG**T**ACAGG CAGCTTAGAA**G**
FJAB01	59	GTTCACTGCCGCACAGG CAGCTTAGAAA	--[c]
FJAD01	48	GTTCACTGCCG**T**ACAGG CAGCTTAGAAA	GTTCACTGCCG**T**ACAGG CAGCTTAGAA**G**
FJAJ01	55	GTTCACTGCCG**T**ACAGG CAGCTTAGAAA	GTTCACTGCCG**T**ACAGG CAGCTTAGAA**G**
FJAM01	72	GTTCACTGCCG**T**ACAGG CAGCTTAGAAA	GTTCACTGCCG**T**ACAGG CAGCTTAGAA**G**
FJBW01	69	GTTCACTGCCG**T**ACAGG CAGCTTAGAAA	GTTCACTGCCG**T**ACAGG CAGCTTAGAA**G**
FJDB01	107	GTTCACTGCCGCACAGG CAGCTTAGAAA	GTTCACTGCCGCACAGG CAGCTTAGAA**G**
FJDP01	55	GTTCACTGCCG**T**ACAGG CAGCTTAGAAA	GTTCACTGCCG**T**ACAGG CAGCTTAGAA**G**
FJEH01	91	GTTCACTGCCGCACAGG CAGCTTAGAAA	GTTCACTGCCGCACAGG CAGCTTAGAA**G**
FJEJ01	56	GTTCACTGCCGCACAGG CAGCTTAGAAA	--[c]
FJMD01	47	GTTCACTGCCG**T**ACAGG CAGCTTAGAAA	GTTCACTGCCG**T**ACAGG CAGCTTAGAA**G**
JFID01	92	GTTCACTGCCGCACAGG CAGCTTAGAAA	GTTCACTGCCGCAT AGG CAG**TGGGTTG**
JFIM01	39	GTTCACTGCCGCACAGG CAGCTTAGAA**G**	--
LAYB01	94	GTTCACTGCCGCACAGG CAGCTTAGAAA	GTTCACTGCCGCACAGG CAGCTTAGAA**G**

LBAK01	53	GTTCACTGCCG**T**ACAGG CAGCTTAGAAA	GTTCACTGCCG**T**ACAGG CAGCTTAGAA**G**
LBAN01	54	GTTCACTGCCGCACAGG CAGCTTAGAA**G**	--
LBAV01	34	GTTCACTGCCGCACAGG CAGCTTAGAA**G**	--
UGOL01	129	GTTCACTGCCGCACAGG CAGCTTAGAAA	GTTCACTGCCGCACAGG CAGCTTAGAA**G**
UGOM01	53	GTTCACTGCCGCACAGG CAGCTTAGAAA	GTTCACTGCCGCACAGG CAGCTTAGAA**G**

[a] The isolates listed in this table reflect those present in the phylogeny from figure 1B

[b] Mutations in the last repeat relative to the consensus repeat, and in the two primary

consensus repeats relative to each other, are denoted in red

[c] While these strains do not possess a mutated last repeat, their 3[rd] last repeat is mutated

relative to the consensus repeat

Figure 1 | Phylogenetic analysis of type I-F CRISPR-Cas system diversity in *L. pneumophila* reveals a horizontal distribution across isolates. *L. pneumophila* draft and completed genomes were analyzed using CRISPRDetect and CRISPRCasFinder to identify type I-F systems present within the genomes. Isolates that possessed type I-F systems were subjected to phylogenetic analyses of their *cas1* gene and core genome. The core genome alignment for the examined isolates was determined by Roary. Isolate names are colour-coded based on the *cas1* gene phylogeny to allow for comparison between the analyses. The isolates used in this analysis and their accession numbers can be found in supplementary table 1. **A)** The *cas1* gene phylogeny for *L. pneumophila* isolates with type I-F systems reveals six different *cas1* groups. **B)** The core genome phylogeny of the examined isolates is not congruent with the *cas1* phylogeny, suggesting that the type I-F system was horizontally distributed rather than vertically inherited. Note that Lens possesses two type I-F CRISPR-Cas systems, one on a plasmid (group Θ) and one on its chromosome (group Φ). Also note that isolates with 100% nucleotide identity in their core genome, as well as a shared *cas* gene group and redundant CRISPR array, have been collapsed to one representative in the phylogeny. Isolates with unique CRISPR arrays are listed. Collapsed isolates include:
+: FJEK01, FJEL01, FJEM01, FJEN01, FJEO01, FJEP01, FJEQ01, FJER01, FJES01, FJET01, FJEU01, FJEV01, FJEX01, FJEY01, FJEZ01, FJFA01
++: FJBI01, FJBU01
+++: LAXR01
++++: FJEI01
+++++: FFI105, FFI337

Figure 2 | CRISPR array analysis suggests that spacer acquisition and spacer loss has contributed to array diversification in *L. pneumophila* **type I-F systems.** *L. pneumophila* isolates were subjected to a CRISPRStudio analysis to look at the spacer composition of their CRISPR arrays. Grey boxes denote unique spacers, coloured boxes denote shared spacers and a dashed line denotes spacer loss. The isolate colour coding scheme is based on the *cas1* grouping from figure 1. Denoted isolates within the same *cas* gene group have 100% nucleotide identity in their core genomes. Note that *cas1* group β appears to have spacer rearrangements present, resulting in an imperfect alignment between the arrays.

Figure 3 | A comparison of the Lens chromosome and pLPL CRISPR-Cas systems. A) The overall pairwise amino acid identity across the Cas proteins of the two systems is approximately 97%, with individual pairwise Cas protein identities ranging from 96% to 99%. **B)** A CRISPRStudio alignment of the arrays for two systems shows completely unique spacer content and a differing number of spacers. **C)** Analysis of the repeat sequences shows that the Lens pLPL and chromosome systems have one SNP between their consensus repeats, in addition to possessing a mutated last repeat in their arrays. Mutations are denoted in red.

Figure 4 | Spacer acquisition and spacer loss in both Lens CRISPR-Cas systems. Spacer acquisition and loss were analyzed using a PCR-based screen where the leader-end of the CRISPR array for both the control samples and the transformed samples was amplified with system-specific primers to differentiate between the chromosomal Lens and the plasmid Lens arrays and visualized on an agarose gel. Products from the transformed samples were compared to the control, which contained untransformed genomic DNA. Bands representing spacer acquisition and loss are indicated.

Figure 5 | Characterization of self-primed spacer acquisition in the two Lens CRISPR-Cas systems. Bacterial transformants with targeted plasmids were passaged for 20 generations without antibiotic selection to enrich for spacer acquisition; the leader end of the CRISPR array was amplified and the amplicons were Illumina sequenced. Newly targeted protospacers were obtained from the raw reads using an in-house bioinformatics pipeline and visualized with Circos. The arrows in the simplified array schematic show the primer location for PCR amplification prior to sequencing, while the star denotes the priming spacer. All data are the average of three biological replicates. **A)** The distribution of newly targeted protospacers mapped to the priming plasmid on the Circos plot reveals a strand bias in self-primed spacer acquisition within the Lens (pLPL) CRISPR-Cas system. The height of the bars indicates the number of spacers mapped to the position on the plasmid, up to 5% of total acquired spacers. **B)** The distribution of newly targeted protospacers mapped to the priming plasmid shows a similar pattern of self-primed spacer acquisition within the Lens (chromosome) CRISPR-Cas system to that of its pLPL CRISPR-Cas counterpart. Labelling as in **(A)**.

Figure 6 | Characterization of cross-primed spacer acquisition in the two Lens CRISPR-Cas systems. The experimental set-up is the same as described in figure 5. All data are the average of three biological replicates. The distribution of newly targeted protospacers mapped to the priming plasmid reveals cross-priming between the Lens (chromosome) and Lens (pLPL) CRISPR-Cas systems. **(A)** shows Lens (chromosome) primed, Lens (pLPL) array examined while **(B)** shows Lens (pLPL) primed, Lens (chromosome) array examined.

Catastrophic array collapse

HGT of secondary array

Cross-primed replenishment

Figure 7 | Cross-priming between two similar CRISPR-Cas systems can repopulate a collapsed CRISPR array. A bacterium with a CRISPR-Cas system could undergo a mass spacer deletion event through homologous recombination between the first (black) and last (white) repeat sequences. The remaining locus carries a single repeat (white). While such "catastrophic collapse" events are likely to occur randomly at a certain rate, a driver of such a collapse could be the acquisition of a self-targeting spacer (yellow), selecting for spacer loss. Horizontal acquisition of a second CRISPR-Cas array (e.g. on a plasmid) is a first step towards replenishing the primary array. If cross-priming can occur between this secondary array and the collapsed array, the original CRISPR array is replenished, but bears an observable molecular scar - conversion of all the repeats to the sequence of the last repeat (white).

Figure 8 | An experimentally depleted chromosomal CRISPR array can be replenished through the activity of a plasmid-based array. A) Replenishment of a depleted array in the lab. Allelic replacement was used to remove the entire array from the chromosomal Lens I-F system, leaving behind a single, last repeat sequence (see materials and methods). This strain was then transformed with a plasmid targeted by the pLPL (Lens plasmid-based) I-F array previously shown to drive primed acquisition (pLPL protospacer 50; see Figure 4). Spacer acquisition by the empty chromosomal array was analyzed using a PCR based screen where the leader-end of the CRISPR array was amplified and visualized on an agarose gel. Products from the transformed samples (samples 2 and 4) were compared to untransformed controls (samples 1 and 3). Samples 1 and 2 are from depleted array clone #1 and samples 3 and 4 are from depleted array clone #2. **B)** Repeat signatures of depletion/replenishment. The repeat structure of the experimentally replenished Lens CRISPR arrays resembles that of *L. pneumophila* str. Alcoy, suggesting a similar array depletion/replenishment event may have occurred within the Alcoy lineage. The frequency of acquiring one new spacer versus two new spacers following replenishment in the Lens array depletion isolates was determined by Illumina sequencing.