Measuring NDC80 binding reveals the molecular basis of tension-dependent kinetochore-microtubule attachments

AUTHORS: Tae Yeon Yoo[1,*], Jeong-Mo Choi[2], William Conway[3], Che-Hang Yu[4], Rohit V. Pappu[2], and Daniel J. Needleman[1,4]

AFFILIATIONS:

[1]Department of Molecular and Cellular Biology and Faculty of Arts and Sciences Center for Systems Biology, Harvard University, Cambridge, MA 02138, USA

[2]Department of Biomedical Engineering and Center for Biological Systems Engineering, Washington University in St. Louis, St. Louis, MO 63130, USA

[3]Department of Physics, Harvard University, Cambridge, MA 02138, USA

[4]John A. Paulson School of Engineering and Applied Sciences, Harvard University, Cambridge, MA 02138, USA.

*Correspondence: tyoo@fas.harvard.edu

Abstract: Proper kinetochore-microtubule attachments, mediated by the NDC80 complex, are required for error-free chromosome segregation. Erroneous attachments are corrected by the tension dependence of kinetochore-microtubule interactions. Here, we present a method, based on fluorescence lifetime imaging microscopy and Förster resonance energy transfer, to quantitatively measure the fraction of NDC80 complexes bound to microtubules at individual kinetochores in living human cells. We found that NDC80 binding is modulated in a chromosome autonomous fashion over prometaphase and metaphase, and is predominantly regulated by centromere tension. We show that this tension dependency requires phosphorylation of the N-terminal tail of Hec1, a component of the NDC80 complex, and the proper localization of Aurora B kinase, which modulates NDC80 binding. Our results lead to a mathematical

model of the molecular basis of tension-dependent NDC80 binding to kinetochore microtubules in vivo.

INTRODUCTION

Chromosome segregation errors lead to aneuploidy and micronuclei formation, which are closely

associated with cancer, infertility, and birth defects (Santaguida and Amon, 2015). Accurate chromosome

segregation is believed to result from a process that actively suppresses potential errors. The mechanism

of error correction remains unclear, but extensive evidence suggests that it is based on the regulation of

the attachment of microtubules to chromosome via the kinetochore, a protein complex assembled at

centromeres (Godek et al., 2015). Previous works suggested that error correction is largely due to the

detachment of kinetochore microtubules (kMTs) being regulated by the tension across centromeres,

which selectively destabilizes erroneous kMT attachments bearing low tension and stabilizes proper

attachments under high tension (Nicklas and Ward, 1994, Liu et al., 2009, Akiyoshi et al., 2010, Lampson

and Cheeseman, 2011, Godek et al., 2015). However, the molecular mechanism of the tension-dependent

regulation of kMT attachments is still poorly understood.

The highly conserved NDC80 complex is the major coupler of the kinetochore to microtubules

(Cheeseman et al., 2006, DeLuca et al., 2006). In human mitotic cells, ~240 NDC80 complexes are

recruited at the outer layer of each kinetochore (Suzuki et al., 2015) and interact with ~20 kMTs by

directly binding to them (Cheeseman and Desai, 2008, Maiato et al., 2004, Rieder, 1982). In vitro

experiments showed that the binding affinity of NDC80 for microtubules decreases upon the

phosphorylation of the N-terminal tail of Ndc80/Hec1 protein by Aurora B kinase (Cheeseman et al.,

2006, Zaytsev et al., 2014, Zaytsev et al., 2015), which may explain the contribution of Aurora B to error

correction (Tanaka et al., 2002). It is unclear how the biochemical activities of NDC80 and Aurora B

result in tension-dependent kMT detachment. The lack of techniques to measure the binding of the

NDC80 to kMTs in vivo has been a major obstacle to investigate this.

RESULTS

FLIM-FRET measures the fraction of donor-labeled NDC80 complexes engaged in FRET with

acceptor-labeled microtubules

Inspired by previous work (Posch et al., 2010), we sought to develop a Förster Resonance Energy

Transfer (FRET) based approach to directly measure the association between the NDC80 complex and

kinetochore microtubules (kMTs) in living cells. We engineered U2OS cells stably expressing Nuf2, a

subunit of the NDC80 complex, N-terminally labeled with a cyan fluorescent protein, mTurquoise2

(Figure 1A). In this same cell line, we also inserted a tetracysteine (TC) motif at the C-terminus of β-

tubulin (TUBB) using CRISPR-induced homologous recombination, which becomes fluorescent after

binding to the membrane-permeable dye FlAsH (Hoffmann et al., 2005) (Figure 1A and B). The small

size (six amino acids) of the TC motif minimizes the negative effects of labeling the C-terminus of

tubulin, allowing the engineered cells to successfully pass through mitosis (Andresen et al., 2004).

CRISPR-mediated endogenous tubulin tagging ensures low cell-to-cell variation and a high fraction of

labeled β-tubulin, which was estimated to be 26.1% ± 5.4% (SD) (Figure 1–figure supplement 1 and see

Supplemental experiments in Materials and Methods).

We used time-correlated single photon counting (TCSPC) fluorescence lifetime imaging

microscopy (FLIM) to quantitatively measure FRET between mTurquoise2 and TC-FlAsH in tissue

culture cells (Figure 1–figure supplement 2). TCSPC FLIM-FRET provides fluorescence decay curves of

the donor fluorophore at each pixel location. If a donor fluorophore has a single-exponential fluorescence

decay curve when not engaged in FRET, then when it is engaged in FRET the fluorescence decay curve

will also be single-exponential, but with a shorter lifetime. A pixel containing a mixture of such donor

fluorophores engaged in FRET and not engaged in FRET displays a fluorescence decay curve that is a

sum of two exponentials. Bayesian analysis of the fluorescence decay curves provides a bias-free

measurement of the relative fraction of the two exponentials, and hence the fraction of donor fluorophores

engaged in FRET (Yoo and Needleman, 2016, Kaye et al., 2017). In contrast to intensity-based FRET

1
Figure 1. FLIM-FRET measurement of NDC80-kMT binding in human tissue culture cells.
(A) Engineered U2OS cell expressing mTurquoise2-NDC80 and β-tubulin-TC-FlAsH. NDC80 (gray),
mTurquoise2 (blue) and TC-FlAsH (green).
(B) Two-photon microscopy images of the engineered U2OS cells not exposed to FlAsH (top) and
exposed to FlAsH (bottom). 3μm scale bar. mTurquoise2 (blue) and FlAsH (green).
(C) Example fluorescence decay curves of mTurquoise2-NDC80 in the engineered U2OS cells not
exposed to FlAsH (left, green circle) and exposed to FlAsH (right, orange triangle), plotted with the best-
fit single-exponential decay models (black and blue dotted lines). Corresponding weighted residuals (the
deviation of data from model, divided by the square root of the number of photons) are plotted below
after being smoothened to display systematic deviations.
(D) The fluorescence decay curve of mTurquoise2-NDC80 in the presence of FlAsH labeling (orange
triangle, same as (C)), plotted with the best-fit two-exponential model (blue solid line). The single-
exponential model fit to the fluorescence decay curve in the absence of FlAsH labeling (black dotted line)
plotted together for comparison. Corresponding smoothened weighted residual (described above) for the
two-exponential model is plotted below. Long- and short-lifetime exponentials correspond to the
mTurquoise2-NDC80 populations in non-FRET state and FRET state, respectively, and their relative
amplitudes give the fraction of each population. To facilitate the comparison, the fluorescence decay
curves in the absence and presence of FlAsH labeling were normalized such that they asymptotically
overlap.
Data points and source FLIM data are available in Figure1-Data (Yoo et al.).

methods, FLIM-FRET is capable of quantifying the fraction of donor fluorophores engaged in FRET

when donors and acceptors are differentially distributed in cells, and it is less prone to errors arising from

instrumental artefacts and photobleaching (Berezin and Achilefu, 2010).

We first characterized the fluorescence decay of mTurquoise2-NDC80 in the absence of FRET by

performing FLIM measurement on the engineered U2OS cells (mTurquoise2-NDC80/β-tubulin-TC) that

were not exposed to FlAsH (Figure 1B, top). We found that their fluorescence decays are well described

as a single exponential with a lifetime of 3.75 ns ± 0.09 ns (SD) (Figure 1C, left, and Figure 1−figure

supplement 3A). As discussed above, this single exponential decay profile is expected when the donor

fluorophores do not engage in FRET. We next measured the fluorescence decay of the mTurquoise2-

NDC80 in the presence of FlAsH labeling of microtubules. In this case, a single exponential provided a

poor fit to the data, exhibiting significant systematic deviations (Figure 1C, right). The fluorescence decay

in the presence of FlAsH labeling was well fit by a sum of two exponentials with lifetimes 3.71 ns ± 0.04

ns (SE) and 0.75 ns ± 0.12 ns (SE) (Figure 1D). The long lifetime of the two-exponential fit was

indistinguishable from the lifetime in the absence of FRET (p = 0.68, two-sided Z-test), and thus

corresponds to the non-FRET donor population. Therefore, the short-lifetime species is the FRET donor

population. The relative amplitude of the short- and long-lifetime exponentials are 0.13 ± 0.01 (SE) and

0.87 ± 0.01 (SE), respectively, thus 13% ± 1% (SE) of donor fluorophores are engaged in FRET.

FRET between mTurquoise2-NDC80 and FlAsH results from the NDC80-kMT binding

Having demonstrated our ability to measure FRET between mTurquoise2-NDC80 and FlAsH in tissue

culture cells, we explored if the FRET is due to the binding of NDC80 to kMTs. We first engineered an

alternative construct with mTurquoise2 conjugated to the distally located C-terminus of Nuf2, far

removed from kMTs. This alternative construct displayed only a single long-lifetime state in either the

presence or absence of TC-FlAsH, arguing that FRET does not result from non-specific interactions

(Figure 1−figure supplement 3B, C and E). Incubating cells with nocodazole to depolymerize

microtubules caused a reduction (p<10^{-10}, two-sided Z-test) of NDC80 FRET fraction from 13% ± 1%

(SE) to 3% ± 1% (SE) (Figure 1−figure supplement 3D). Thus, FRET strongly depends on the presence of

microtubules.

We next investigated if NDC80 that is close to kMTs, but not bound to them, can lead to

appreciable FRET. Answering this requires knowing the Förster radius between mTurquoise2 and TC-

FlAsH, which we determined to be 5.90 ± 0.10 nm (SE) through a combination of FLIM measurements

and Monte Carlo simulations (see Figure 1−figure supplement 4 and Supplemental experiments in

Materials and Methods). We next performed large-scale Monte Carlo simulations of mTurquoise2-

NDC80 at various distances between the calponin homology (CH) domain of Hec1/Ndc80 protein (the

NDC80 complex's microtubule binding domain adjacent to mTurquoise2 (Alushin et al., 2010)) and

FlAsH-labeled microtubules and simulated the fluorescence decay curves, which revealed that NDC80

more than 8 nm away from the kMT do not contribute to the short-lifetime FRET state (Figure 1−figure

supplement 5A and B). Thus, FRET only results when the CH domain of Hec1 is very close to the surface

of kMTs, consistent with the short-lifetime species being NDC80 complexes whose Hec1 CH domains

are bound to kMTs, an interpretation further supported by biological perturbation experiments described

below.

Even though FRET results only from NDC80 bound to kMTs, the measured FRET fraction is not

identical to the fraction of NDC80 bound to kMTs because not all tubulin heterodimers are labeled with

TC-FlAsH. Using large scale Monte Carlo simulations of mTurquoise2-NDC80 bound to FlAsH-labeled

microtubules, we generated fluorescence decay curves for various NDC80 binding fractions, and

estimated the resulting NDC80 FRET fractions from a fit to a two-exponential decay (see Figure 1−figure

supplement 5C and Supplemental experiments in Materials and Methods). We found that the NDC80

FRET fraction increases linearly with the NDC80 binding fraction with a slope of 0.42 ± 0.08, indicating

that 42% of attached mTurquoise2-NDC80 contribute to the short-lifetime FRET state (Figure 1−figure

supplement 5C). Thus, the measured FRET fraction of 13% in Figure 1D corresponds to 31% of NDC80 complexes being bound to kMTs.

NDC80-kMT binding is regulated in a chromosome-autonomous fashion throughout prometaphase

Using the FLIM-FRET measurements of NDC80-kMT binding, we first investigated how NDC80-kMT binding evolves over the course of mitosis. We found that the average NDC80-kMT binding gradually increases as mitosis progresses, with NDC80 FRET fraction rising from 7% in early prometaphase to 14% in late metaphase, and reaching about 18% in anaphase (corresponding to NDC80 binding fractions of 17% in prometaphase; 33% in late metaphase; and 43% in anaphase) (Figure 2A). This temporal change in NDC80-kMT binding may underlie the previously observed decrease in kMT turnover throughout mitosis (Kabeche and Compton, 2013, Zhai et al., 1995).

The change in the average NDC80-kMT binding over the course of error correction in prometaphase could be due to a cell cycle-dependent coordinated regulation of NDC80 across kinetochores (coordinated regulation), an independent modulation of NDC80 on different chromosomes (chromosome-autonomous regulation), or a combination of both. To investigate the contribution of chromosome-autonomous regulation, we sought to determine if different populations of kinetochores in prometaphase exhibit different NDC80-kMT binding. We compared the extent of the NDC80-kMT binding of kinetochores centered at the metaphase plate to those located off-centered (Figure 2B), and found that the centered kinetochores exhibit 2.0 ± 0.4 times higher NDC80 binding than the off-centered kinetochores. We next investigated how the NDC80-kMT binding of centered and off-centered kinetochores change with time in prometaphase. As mitosis progresses and the chromosomes align to the metaphase plate, the number of kinetochores in the center region increases while the number of kinetochores in the off-center region decreases (Figure 2C). NDC80-kMT binding continuously increases over time for the kinetochores located in the center region, but remains constant with the FRET fraction of ~7% for the kinetochores in the off-center region (Figure 2D). The observation of differences in

1
Figure 2. NDC80-kMT binding is regulated in a chromosome-autonomous fashion.
(A) Example cell images and time course of NDC80 FRET fraction from prometaphase to metaphase to
anaphase (n = 11 cells). Black squares are the mean, y-error bars are the SEM, and x-error bars are the SD
of the data points (green circles) in equally spaced time intervals. 5 µm scale bar.
(B) Kinetochores at each time point in prometaphase cells are divided into two groups, centered and off-
centered, based on their distances from the metaphase plate. Kinetochores less than 1 µm away from the
metaphase plate were classified as centered, and kinetochores more than 2.5 µm away were classified as
off-centered.
(C) Time course of the fraction of centered (green) and off-centered (orange) kinetochores in
prometaphase.
(D) Time course of NDC80 FRET fraction of centered (green circles) and off-centered (orange squares)
kinetochores in prometaphase (n = 11 cells, 2886 centered and 572 off-centered kinetochores). Data
points are the mean, y-error bars the SEM, and the x-error bars the SD in equally spaced time intervals.
Gray areas are the 95% confidence intervals for the linear fits.
Data points and source FLIM data are available in Figure2-Data (Yoo et al.).

NDC80 binding between different subpopulations of kinetochores strongly argues for the existence of chromosome-autonomous regulation, which might be modulated by tension, Aurora kinases A and B, pathways that control the conversion of lateral to end-on kMT attachments, or other factors (Godek et al., 2015, DeLuca et al., 2018). We speculate that the temporal increase in NDC80-kMT binding of centered kinetochores is due to the gradual decrease in the number of kinetochores with erroneous attachment that transiently lie on the metaphase plate (Magidson et al., 2011). The constant NDC80-kMT binding of off-centered kinetochores argues for a lack of temporal regulation of this subpopulation.

NDC80-kMT binding is positively correlated with centromere tension

After demonstrating that different population of kinetochores exhibits different NDC80-kMT binding throughout prometaphase, we next investigated the factors contributing to chromosome-autonomous regulation of the interaction between NDC80 and kMTs. Aligned chromosomes in U2OS cells oscillate around the metaphase plate, with microtubules attached to the leading and trailing kinetochores primarily depolymerizing and polymerizing, respectively (Tirnauer et al., 2002, Armond et al., 2015) (Figure 3A). The distance between sister kinetochores (referred to as K-K distance) fluctuates during the oscillation (Magidson et al., 2011), as the centromere deforms in response to the dynamic change in tension (Figure 3A). Therefore, chromosome oscillation provides a window to study how NDC80 binding is related to kMT dynamics and centromere tension in a physiologically relevant condition.

We first asked whether NDC80 binding is different on leading and trailing kinetochores. We acquired time-lapse movies of 17 metaphase cells, tracked their kinetochores, identified sister kinetochores by their relative motions (Figure 3B), and quantified the NDC80 binding fraction in groups of kinetochores with similar velocities using FLIM-FRET analysis. We found that the NDC80 FRET fraction is higher at trailing kinetochores (12.8% $\pm$ 0.5%, SEM) than leading kinetochores (11.4% $\pm$ 0.5%, SEM), regardless of their speeds (Figure 3C), suggesting that NDC80 preferentially binds to

1
Figure 3. NDC80-kMT binding is correlated with kMT dynamics and centromere tension.
(A) (left) kMTs predominantly depolymerize at leading kinetochores and polymerize at trailing
kinetochores. (right) K-K distance is a proxy for centromere tension. Measuring NDC80-kMT binding
along with the kinetochore movement and K-K distance therefore reveals how NDC80-kMT binding is
related to the kMT dynamics and centromere tension.
(B) Image of a metaphase cell with mTurquoise2-NDC80 (blue) and β-tubulin-TC-FlAsH (green), and
kinetochore tracking (yellow circles) and pairing (red lines) results. 3 μm scale bar.
(C) NDC80 FRET fraction vs. kinetochore speed for leading (green circle) and trailing (orange triangle)
kinetochores (n= 17 cells, 681 kinetochores/data point). Data points are the mean, y-error bars the SEM,
and the x-error bars the interquartile ranges within groups of kinetochores with similar velocities.
(D) NDC80 FRET fraction vs. K-K distance for untreated cells (green circle, n = 17 cells, 984
kinetochores/data point), cells treated with 10 μM taxol (orange triangle, n = 7 cells, 525
kinetochores/data point), and cells treated with 5 μM STLC (purple square, n = 16 cells, 493
kinetochores/data point). For STLC data, only poleward-facing kinetochores are plotted (see Figure 3-
supplement 2 for comparison between poleward and anti-poleward kinetochores). Data points are the
mean, y-error bars the SEM, and the x-error bars the interquartile ranges within groups of kinetochores
with similar K-K distances. Gray area is the 95% confidence interval for the linear fit to the combined
data.
(E) Histograms of K-K distances for the untreated (top, green), taxol-treated (middle, orange), and STLC-
treated (bottom, purple) cells. 3 μm scale bar in the cell images of mTurquoise2-NDC80 (blue) and beta-
tublin-TC-FlAsH (green). ***: p-value (Welch's t-test) less than 10^{-30}.
Data points and source FLIM data are available in Figure3-Data (Yoo et al.).

polymerizing kMTs in vivo. The preferential binding is statistically significant (p<0.02, two-sided Z-test),

yet small, presumably because leading and trailing kinetochores have a mixture of both polymerizing and

depolymerizing MTs (Armond et al., 2015). This differential binding of NDC80 provides an explanation

for the higher detachment rate of depolymerizing microtubules from kinetochores in vitro (Akiyoshi et al.,

2010), and may give insight into the nature of kMT attachments (Dumont et al., 2012).

The detachment rate of kMTs from kinetochores was shown to be reduced when tension was

increased using glass needles in classic micromanipulation experiments by Bruce Nicklas (Nicklas and

Koch, 1969). Since the NDC80 complex is the predominant coupler of the kinetochore to microtubules

(Cheeseman et al., 2006, DeLuca et al., 2006), we hypothesized that the tension-dependent detachment of

kMTs results from tension-dependent NDC80-kMT binding. To test this possibility, we next investigated

the correlation between NDC80 FRET fraction and centromere tension, inferred by K-K distance, during

chromosome oscillations. We used FLIM-FRET analysis to measure the NDC80 binding in groups of

sister kinetochores with similar K-K distances, and observed a highly statistically significant positive

correlation (p<0.005) between NDC80 FRET fraction and K-K distance (Figure 3D). We observed no

significant correlation between K-K distance and kinetochore velocity (p=0.75), arguing that NDC80

binding is independently regulated by these two factors (Figure 3–figure supplement 1). In the absence of

microtubules, the rest length of K-K distance in human cell is 0.73 μm ± 0.04 μm (Tauchman et al.,

2015), significantly shorter than the K-K distances during metaphase oscillations. Thus, in order to

investigate a wider range of K-K distance, we treated cells with taxol, a microtubule-stabilizing drug,

which greatly reduced K-K distances (0.90 μm ± 0.10 μm, taxol vs. 1.19 μm ± 0.19 μm, untreated, SD,

$p<10^{-30}$) (Figure 3E) as well as NDC80 FRET fraction (Figure 3D). As an alternative way to reduce the

tension, we inhibited Eg5 with 5 μM S-trityl-L-cysteine (STLC) (Skoufias et al., 2006). STLC-treated

cells form monopolar spindles with reduced K-K distances (0.87 μm ± 0.10 μm, SD, $p<10^{-30}$) (Figure 3E).

In these monopolar spindles, NDC80 FRET fractions from poleward-facing kinetochores was positively

correlated with the K-K distance (p<0.05), while anti-poleward-facing kinetochores displayed reduced

NDC80 FRET fraction with no significant correlation with K-K distance (p=0.46), presumably because

many of these kinetochores are monotelically attached from the poleward side (Figure 3-figure

supplement 2). The correlation between NDC80 FRET fraction and K-K distance was similar between

taxol- and STLC-treated cells (p=0.15, see Figure 3D), arguing that the relation between NDC80 FRET

fraction and K-K distance is insensitive to the precise perturbation used to reduce tension. Combining the

data of untreated, taxol-treated and STLC-treated (only poleward-facing kinetochores) cells, we found

that the NDC80 FRET fraction continually increases with K-K distance over the full range of K-K

distance (positive correlation, p<0.00005) (Figure 3D). The extent of variation of NDC80-kMT binding

with K-K distance is comparable to the extent of variation over the course of mitosis, from prometaphase

to anaphase onset, as well as the extent of difference between centered and off-centered kinetochores in

late prometaphase (compare Figure 3D with 2A and D).

Aurora B kinase regulates NDC80-kMT binding in a graded fashion in vivo

Aurora B kinase is one of the best characterized components of the error correction process, and

the N-terminal tail of the Hec1/Ndc80 protein in the NDC80 complex is a known substrate of Aurora B

kinase that contains nine phosphorylation sites (Tanaka et al., 2002, Biggins et al., 1999, Cheeseman et

al., 2006, DeLuca et al., 2006, Ciferri et al., 2008, Hauf et al., 2003). Previous biochemistry experiments

demonstrated that the phosphorylation state of Hec1 modulates its binding to microtubules in vitro

(Cheeseman et al., 2006, Zaytsev et al., 2014, Zaytsev et al., 2015). We used our FLIM-FRET technique

to investigate the relationship between Aurora B kinase activity and NDC80-kMT binding in cells. We

first added the ATP-competitive Aurora B inhibitor, ZM447439, to late prometaphase cells, and observed

a gradual increase in NDC80 FRET fraction over ~10 minutes, from 9% to nearly 18% (corresponding to

21% NDC80 binding fraction before Aurora B inhibition and 43% after the inhibition) (Figure 4A and

Figure 4–figure supplement 1A). Thus, Aurora B is a major modulator of NDC80-kMT binding in cells.

This modulation of NDC80-kMT binding could occur directly, through phosphorylation of the N-terminal

1
Figure 4. Aurora B kinase regulates NDC80-kMT binding in a graded fashion in vivo.
(A) (top) Cell images showing mTurquoise2-NDC80 (blue) and beta-tubulin-TC-FlAsH (green). (bottom)
Time course of NDC80 FRET fraction in response to Aurora B inhibition by 3 μM ZM447439 (n = 15
cells).
(B) Images of cells with mTurquoise2-NDC80 (blue) and beta-tubulin-TC-FlAsH (green) after depleting
endogenous Hec1 by siRNA and expressing siRNA-insensitive WT or three different phosphomimetic
mutants of Hec1: 9A-, 2D-, and 9D-Hec1 (see Methods).
(C) NDC80 binding fraction of cells whose endogenous Hec1 are replaced with WT or phosphomimetic
Hec1 (see Methods). Black dots are from individual cells and red error bars are mean ± SEM. n = 19, 22,
12, and 17 cells for WT-, 9A-, 2D-, and 9D-Hec1. *: p<0.1; **: p<0.01; ***: p<0.001; ****: p<0.0001.
(D) Time course of NDC80 FRET fraction of 2D-Hec1-expressing cells in response to Aurora B
inhibition by 3 μM ZM447439 (n = 12 cells).

(E) (top) Cell images color-coded with Aurora B sensor non-FRET fraction. (bottom) Time course of the non-FRET fraction of the cytoplasmic Aurora B FRET sensor in response to 3 μM ZM447439 (n = 10 cells).

In (A), (D) and (E), black squares and error bars are the weighted mean and SEM of the data points (green circles) in equally spaced time intervals of 1 minute. Red solid and dashed lines are the best-fit exponential decay models and their 95% confidence intervals, respectively. 5 μm scale bar for all images.

(F) NDC80 binding fraction (converted from NDC80 FRET fraction in (A)) plotted against the fraction of phosphorylated Aurora B phosphorylation sites in NDC80 (converted from Aurora B FRET sensor non-FRET fraction in (E)). Red solid and dashed lines are the best-fit NDC80-kMT binding model (derived in Mathematical modeling in Materials and Methods) and its 95% confidence interval.

Data points and source FLIM data are available in Figure4-Data (Yoo et al.).

tail of the Hec1/Ndc80 protein, or indirectly, through other Aurora B substrates which are at kinetochores

or which influence spindle assembly (Carmena et al., 2012, Krenn and Musacchio, 2015). To further

investigate the Aurora B modulation of NDC80 binding, we used non-phosphorylatable mutants of

Hec1/Ndc80 protein, in which all nine identified Aurora B target sites are mutated to either aspartic acid

(a phospho-mimicking mutation) or alanine (a phospho-null mutation) (DeLuca et al., 2011). We replaced

the endogenous Hec1 with wild-type Hec1 or the phosphomimetic mutant of Hec1 by RNAi knockdown

and rescue (Figure 4B, see Methods). The NDC80 binding fraction (converted from NDC80 FRET

fraction) decreased with the number of phospho-mimicking mutations, from 38%±1% with 9A-Hec1 (all

nine phosphorylation sites substituted with Ala) to 14%±1% with 2D-Hec1 (two sites, S44 and S55,

substituted with Asp while the others with Ala) and to 11%±1% with 9D-Hec1 (all nine sites substituted

with Asp) (mean±SEM, Figure 4C). The average NDC80 binding of WT-Hec1 was similar, but slightly

higher than 2D-Hec1, consistent with previous results arguing that on average there are zero to two sites

phosphorylated per Hec1/Ndc80 protein in prometaphase and metaphase (Zaytsev et al., 2014).

14   We next sought to determine if the increased binding of NDC80 to kMTs upon Aurora B

inhibition is caused by a change of phosphorylation of the N-terminal tail of Hec1/Ndc80 protein. To this

end, we added the Aurora B inhibitor, ZM447439, to cells with endogenous Hec1 replaced with 2D-Hec1,

whose nine identified Aurora B target sites in the N-terminal tail cannot be phosphorylated. After the

addition of ZM447439, the NDC80 FRET fraction increases by only 0.02 ± 0.01 (SE), which is

statistically significant ($p<0.001$), but substantially smaller than the increase observed in cells with

endogenous Hec1 (0.09 ± 0.01 (SE)) (compare Figure 4D to 4A). Thus, the modulation of NDC80

binding to kMTs by Aurora B predominantly occurs through the phosphorylation of the N-terminal tail of

Hec1/Ndc80.

23   We next quantified how Aurora B inhibition influences Aurora B activity in cells. We performed

FLIM measurement on a cytoplasmic Aurora B FRET biosensor (Fuller et al., 2008), which contains a

kinesin-13 family Aurora B substrate whose phosphorylation obstructs intramolecular FRET between

mTurquoise2 and YPet (Figure 4–figure supplement 1B). During ZM447439 treatment, we found a

continual reduction in the fraction of the Aurora B sensors in the non-FRET state, a proxy for Aurora B

phosphorylation, from 0.540 ± 0.007 (SEM) to 0.368 ± 0.012 (SEM) (Figure 4E and Figure 4–figure

supplement 1C). Nuf2-targeted Aurora B sensor responded to the ZM447439 treatment with similar

kinetics, arguing that the time scale of response to Aurora B inhibition is insensitive to the spatial location

of the substrate (Figure 4–figure supplement 1D). As the typical time scale of drug uptake is far slower

than typical phosphorylation/dephosphorylation kinetics (Thurber et al., 2014, Huang et al., 1997), it is

reasonable to assume that the phosphorylation level of Aurora B substrate is at steady state at each time

point, so plotting the measured NDC80 binding fraction (converted from FRET fraction) vs.

phosphorylated level (converted from Aurora B sensor non-FRET fraction) at each time point reveals

their relationship. This analysis showed a graded dependence of NDC80-kMT binding on

phosphorylation (Figure 4F), which is consistent with the impact of phosphomimetic Hec1 mutants on

NDC80 binding (compare Figure 4C and 4F) and results from previous in vitro assays (Zaytsev et al.,

2014, Zaytsev et al., 2015). The increased NDC80-kMT binding after Aurora B inhibition may underlie

the reduction in detachment of kMTs from kinetochores after Aurora B inhibition, observed in

photoactivation experiments (Cimini et al., 2006).

Haspin-dependent centromere-localized Aurora B is responsible for the tension dependency of

NDC80-kMT binding

Aurora B regulates kinetochore-microtubule interactions, but its contribution to the tension-dependent

stabilization of kinetochore-microtubule attachments is controversial (Campbell and Desai, 2013,

Salimian et al., 2011, Akiyoshi et al., 2010, Liu et al., 2009, Tanaka et al., 2002, Zaytsev et al., 2016,

Godek et al., 2015, Lampson and Cheeseman, 2011, Haase et al., 2017). We next sought to determine if

the correlation between NDC80-kMT binding and centromere tension that we observed (Figure 3D) is

caused by the phospho-regulation of NDC80 by Aurora B. After replacing the endogenous Hec1 protein

with 9A-Hec1, which cannot be phosphorylated on the nine mutated target sites, we no longer observed a

significant correlation between NDC80 FRET fraction and K-K distance (p=0.20) (Figure 5A). 9A-Hec1-

expressing cells displayed significantly larger K-K distance than unperturbed cells (1.36 µm ± 0.21 µm,

SD, $p < 10^{-30}$) (Figure 5B), consistent with previous studies (Tauchman et al., 2015, Zaytsev et al., 2014).

To investigate a wider range of K-K distance, we treated cells with taxol or STLC (Figure 5-supplement

figure 1A), as described above, and found no correlation over the full range of K-K distance (p=0.29,

Figure 5A and B). Since the strong correlation between NDC80 binding and K-K distance (Figure 3D) is

eliminated in the non-phosphorylatable Hec1 mutant 9A-Hec1 (Figure 5A), this argues that the

phosphorylation of the N-terminal tail of Hec1 is responsible for the correlation between NDC80-kMT

binding and K-K distance. As Aurora B is believed to be the primary kinase that phosphorylates the N-

terminal tail, this further suggests that the activity of Aurora B is responsible for the correlation between

NDC80-kMT binding and K-K distance.

Aurora B is localized to centromeres in prometaphase and metaphase (Carmena et al., 2012). We

next investigated if this localization is important for the correlation between NDC80-kMT binding and K-

K distance. We used the haspin kinase inhibitor, 5-iodotubercidin (5-ITu), which has previously been

shown to compromise the recruitment of Aurora B to inner-centromeres (Wang et al., 2012) (Figure 5C).

After 10 minutes of exposure of cells to 5-ITu, INCENP, a member of the chromosome passenger

complex (CPC), which also includes Aurora B (Carmena et al., 2012), was drastically reduced at

centromeres (Figure 5D). Treating cells with 5-ITu for over 15 minutes did not significantly alter the

average K-K distance (1.16 µm ± 0.18 µm, 5-ITu vs. 1.19 µm ± 0.19 µm, untreated, SD) or the overall

average fraction of NDC80 bound to kMTs (FRET fraction, 11.79% ± 0.02% 5-ITu vs. 11.87% ± 0.02%,

untreated, SEM), but eliminated the correlation between K-K distance and NDC80-kMT binding (p=0.41,

Figure 5E and F). In order to investigate a wider range of K-K distance, we treated cells with both taxol

and 5-ITu or with both STLC and 5-ITu (Figure 5-supplement figure 1B), and found no correlation over

the full range of K-K distance (p=0.96, Figure 5E and F). Our observation that the mislocalization of

1
2   **Figure 5. Haspin-dependent centromere-localized Aurora B is responsible for the tension**
3   **dependency of NDC80-kMT binding**

(A) NDC80 FRET fraction vs. K-K distance for 9A-Hec1-expressing cells with no drug treatment (green circle, n = 12 cells, 803 kinetochores/data point), with 10 μM taxol treatment (orange triangle, n = 9 cells, 1113 kinetochores/data point), or with 5 μM STLC treatment (purple square, n = 10 cells, 855 kinetochores/data point). For STLC data, only poleward-facing kinetochores are included (see Figure 5-supplement 1A for comparison between poleward-facing and anti-poleward-facing kinetochores). Data points are the mean, y-error bars the SEM, and the x-error bars the interquartile ranges within groups of kinetochores with similar K-K distances. Gray area is the 95% confidence interval for the linear fit to the combined data.

(B) Histograms of K-K distances for the untreated 9A-Hec1 cells (top, green), untreated cells with endogenous Hec1 (top, black line), 9A-Hec1 cells treated with taxol (middle, orange), and 9A-Hec1 cells treated with STLC (bottom, purple). 3 μm scale bar in the cell images of mTurquoise2-NDC80 (blue) and beta-tubulin-TC-FlAsH (green). ***: $p < 10^{-6}$ (Welch's t-test).

(C) Haspin kinase phosphorylates histone H3 at Thr3 (H3T3), which recruits the chromosome passenger complex (CPC, red) to centromeres. 5-Iodotubercidin (5-ITu) inhibits haspin kinase, thereby displacing Aurora B from centromeres.

(D) Spinning-disk confocal microscopy images of cells expressing mNeonGreen-Nuf2 (green) and INCENP-mCherry (red) before (top) and after (bottom) haspin inhibition by 10 μM 5-ITu treatment. 3 μm scale bar.

(E) NDC80 FRET fraction vs. K-K distance for cells treated with 10 μM 5-ITu (green circle, n = 15 cells, 1170 kinetochores/data point), for cells treated with both 10 μM 5-ITu and 10 μM taxol (orange triangle, n = 3 cells, 359 kinetochores/data point), and for cells treated with 10 μM 5-ITu and 5 μM STLC (purple square, n = 12 cells, 564 kinetochores/data point). For 5-ITu+STLC data, only poleward-facing kinetochores are included (see Figure 5-supplement 1B for comparison between poleward-facing and anti-poleward-facing kinetochores). Data points are the mean, y-error bars the SEM, and the x-error bars the interquartile ranges within groups of kinetochores with similar K-K distances. Gray area is the 95% confidence interval for the linear fit to the combined data.

(F) Histograms of K-K distances for the 5-ITu-treated (top, green), untreated (top, black line), 5-ITu+taxol-treated (middle, orange), and 5-ITu+STLC-treated cells (bottom, purple). 3 μm scale bar in the cell images of mTurquoise2-NDC80 (blue) and beta-tubulin-TC-FlAsH (green). ***: $p < 10^{-6}$ (Welch's t-test).

(G) NDC80 FRET fraction and (H) the non-FRET fraction of Nuf2-targeted Aurora B FRET sensor (proxy for Aurora B activity at NDC80) for different drug treatments. Each data point (gray circle) corresponds to an individual cell, and the error bar (red) shows the mean and SEM. P-values from two-sided Welch's t-test.

Data points and source FLIM data are available in Figure5-Data (Yoo et al.).

Aurora B from centromeres does not affect the K-K distance but compromises the correlation between

NDC80 binding and K-K distance argues that the correlation shown in Figure 3D is due to tension

causing an increase in NDC80-kMT binding, rather than NDC80-kMT binding causing an increase in

tension. Moreover, the tension dependency of NDC80-kMT binding in human tissue culture cells depends

on Aurora B recruitment to centromeres by haspin kinase, further arguing in favor of models in which

phosphorylation by Aurora B plays a central role in chromosome autonomous error correction.

We were surprised that the average NDC80-kMT binding does not significantly change after

mislocalizing Aurora B with 5-ITu (compare Figure 3D and 5E, and see Figure 5G). This suggests that

Aurora B can still act on NDC80 even after the concentration of Aurora B at centromeres is greatly

reduced. Consistent with this hypothesis, the Aurora B activity at kinetochores assessed by Nuf2-targeted

Aurora B FRET sensor was not changed by 5-ITu treatment (Figure 5H). Furthermore, when cells treated

with 5-ITu were exposed to the Aurora B inhibitor ZM447439, NDC80-kMT binding increased (to

NDC80 FRET fraction of 0.17 ± 0.01, SEM) and Aurora B activity at kinetochores decreased (to Aurora

B FRET sensor non-FRET fraction of 0.55 ± 0.01, SEM), indistinguishable from the levels in cells not

subject to 5-ITu exposed to ZM447439 (Figure 5G and 5H). Thus, tension dependency of NDC80-kMT

binding is conferred by Aurora B recruited to centromeres through a haspin-dependent pathway, while the

average level of NDC80-kMT binding is also set by Aurora B, but in a manner that is not dependent on

haspin.

The concentration of Aurora B at the location of NDC80 is dependent on centromere tension

The extent to which Aurora B phosphorylates NDC80 depends on the activity of Aurora B and the

concentration of Aurora B at NDC80. To further investigate how the haspin-dependent pool of Aurora B

confers tension dependency to NDC80-kMT binding, we next examined how Aurora B localization

depends on K-K distance. We used spinning-disk confocal microscopy to image mNeonGreen-Nuf2, to

locate NDC80, and INCENP-mCherry, to measure the distribution of Aurora B. We localized NDC80 to

sub-pixel accuracy and identified sister kinetochore pairs (see Figure 6A and Materials and Methods). For

each pair of kinetochores, we measured the intensity of INCENP-mCherry at the location of NDC80,

normalized on a cell-by-cell basis. Plotting the intensity of INCENP-mCherry at NDC80 as a function of

K-K distance revealed a highly significant anti-correlation ($p < 10^{-4}$, Figure 6B). To explore a

larger range of K-K distances, we treated cells with 10 μM taxol. Combining the data of untreated and

taxol-treated cells, we found that the intensity of INCENP-mCherry at NDC80 linearly decreases with K-

K distance over the full range of K-K distance ($p < 10^{-6}$). This observation suggests that the tension

dependency of NDC80 binding may result from the decrease of Aurora B at NDC80 with increasing K-K

distance.

We next investigated how the concentration of Aurora B at the location of NDC80 is influenced

by haspin inhibition. In the presence of 10 μM 5-ITu, the concentration of Aurora B at NDC80 was

greatly reduced and independent of K-K distance (Figure 6B). The lack of correlation between Aurora B

concentration at NDC80 and K-K distance may explain the lack of tension dependency between NDC80-

kMT binding and K-K distance upon haspin inhibition. We speculate that the finite concentration of

Aurora B at NDC80 after haspin inhibition is the pool of Aurora B that maintains the average level of

NDC80-kMT binding as described above.

A biophysical model of tension dependent NDC80-kMT binding

Taking together, our data suggest that the concentration of Aurora B at NDC80 determines the extent of

NDC80 phosphorylation, which in turn determines the level of NDC80-kMT binding. To further explore

this possibility, we sought to determine the relationship between Aurora B concentration at NDC80 and

NDC80-kMT binding. We plotted the NDC80-kMT binding (converted from the NDC80 FRET fraction

in Figure 3D and 5E) vs. the Aurora B concentration (converted from the normalized INCENP-mCherry

intensity in Figure 6D) for each K-K distance, both with and without haspin inhibition (Figure 7A, see

Materials and Methods). This revealed a highly nonlinear relationship: when the Aurora B concentration

1
Figure 6. The concentration of Aurora B at the location of NDC80 decreases with centromere
tension.
(A) Spinning-disk confocal microscopy image of mNeonGreen-Nuf2 (green) and INCENP-mCherry
(red). 3 µm scale bar. The location of NDC80 was determined to sub-pixel accuracy, using the
mNeonGreen-Nuf2 image. For each pair of sister kinetochores, the intensity of INCENP-mCherry at the
location of NDC80 was measured and normalized on a cell-by-cell basis.
(B) Normalized INCENP-mCherry intensity at the location of NDC80 were averaged within groups of
kinetochores with similar K-K distances, and plotted against the K-K distances for untreated (green
circles), taxol-treated (orange triangles), and 5-ITu-treated (purple squares) cells. Data points are the
mean, y-error bars the SEM, and the x-error bars the interquartile ranges. Black solid and dotted lines are
the linear fits to DMSO+taxol combined data and 5-ITu data, respectively.
906 kinetochore pairs in 9 cells, 599 pairs in 8 cells, and 680 pairs in 6 cells were analyzed for DMSO
control, taxol treatment, and 5-ITu treatment data, respectively.
Data points and source FLIM data are available in Figure6-Data (Yoo et al.).

1
Figure 7. A biophysical model of tension depedent NDC80-kMT binding
(A) Plot of NDC80 binding fraction, f_{bound}, (converted from NDC80 FRET fraction in Figure 3D and 5E)
vs. Aurora B concenration at NDC80, $[A]$ (converted from INCENP-mCherry intensity in Figure 6B, see
Materials and Methods). Data points (black cirles) are the mean and SEM. We constructed a
mathematical model that predicts NDC80 binding fraction from Aurora B concentration at NDC80
through three steps: intermolecular Aurora B auto-activation, NDC80 phosphorylation, and NDC80-kMT
binding. Red line shows the mathematical model fit to the data.
(B) NDC80 binding fraction vs. K-K distance before (left) and after (right) haspin inhibition by 5-ITu.
The data points (black circles) are adapted from Figure 3D and 5E. Red lines are the predictions from the
mathematical model.
Data points are available in Figure7-Data (Yoo et al.).

is lower than ~5 µM, the NDC80 binding fraction is independent of the Aurora B concentration, while for

higher concentrations, the NDC80 binding fraction decreases with the Aurora B concentration (Figure

7A). We constructed a mathematical model to determine if this nonlinear relationship can be explained by

the known biochemistry of Aurora B and NDC80 (Figure 7A) and observed change in the concentration

of Aurora B at NDC80 (Figure 6B). In this model, we assume that there are two independent pools of

Aurora B, haspin-dependent and haspin-independent, both of which engage in intermolecular

autoactivation by phosphorylation in *trans* (Zaytsev et al., 2016, Xu et al., 2010, Sessa et al., 2005,

Bishop and Schumacher, 2002), and are inactivated by phosphatases (Zaytsev et al., 2016, Sessa et al.,

2005, Kelly et al., 2007, Rosasco-Nitcher et al., 2008). The activated Aurora B phosphorylates NDC80,

which changes the binding affinity of NDC80 for kMTs (Cheeseman et al., 2006, Zaytsev et al., 2014,

Zaytsev et al., 2015). This model can be solved analytically, and is sufficient to account for the

relationship between NDC80 phosphorylation and NDC80-kMT binding (Figure 4F), and the relationship

between Aurora B concentration at NDC80 and NDC80-kMT binding (Figure 7A). In this model, the

nonlinear relationship between Aurora B concentration at NDC80 and NDC80-kMT binding ultimately

results from the activation dynamics of Aurora B: at low concentrations, dephosphorylation by

phosphatases overwhelm the in *trans* autoactivation, but above a threshold Aurora B concentration, A*,

these two processes balance, leading to steady state level of activated Aurora B that further increases with

increasing Aurora B concentration.

We next investigated if the same model can recapitulate the tension dependency of NDC80-kMT

binding. Inputting the measured linear relationship between Aurora B concentration at NDC80 and K-K

distance (Figure 6B) into the model reproduced the observed tension-dependent behavior of NDC80-kMT

binding (Figure 7B, left). Performing a similar procedure with the data from haspin inhibited cells (Figure

5E and 6B) revealed that the model successfully predicts both the level of NDC80-kMT binding upon

haspin inhibition and its independence on K-K distance (Figure 7B, right). Thus, this model provides a

self-consistent, quantitative explanation of how the tension dependency of NDC80-kMT binding results

from the biochemistry of Aurora B and NDC80, and the measured change in Aurora B concentration at

NDC80 with tension, without the need to invoke diffusible gradients or additional mechanochemistry.

DISCUSSION

In this study, we developed a method to quantitatively measure the binding of the NDC80 complex to

microtubules at individual kinetochores in human tissue culture cells. Our method uses TCSPC FLIM-

FRET which, in contrast to intensity-based FRET, allows quantitative measurements of the fraction of

molecules engaging in FRET, even with spatially varying concentrations of donors and acceptors. We

calibrated our measurements using control experiments and Monte Carlo simulations, allowing us to

convert the fraction of donor-labeled Nuf2 engaged in FRET to the fraction of NDC80 complexes bound

to kMTs. This technique can be extended to the quantitative assessment of other protein-protein

interactions in living cells.

Using this technique, we demonstrated that NDC80-kMT binding is regulated during

prometaphase in a chromosome-autonomous manner. We observed that NDC80-kMT binding is strongly

correlated to centromere tension, to an extent which is sufficient to account for the changes in NDC80-

kMT binding over the course of prometaphase and metaphase. We characterized how Aurora B modulates

NDC80-kMT binding in cells, which we found predominantly occurs through the phosphorylation of the

N-terminal tail of Hec1. We showed that the correlation between NDC80-kMT binding and centromere

tension is dependent on the phosphorylation of the N-terminal tail of Hec1. We determined the

concentration of Aurora B at the locations of NDC80, which decreases with increasing centromere

tension. Mislocalizing Aurora B by inhibiting haspin kinase eliminated the tension dependency of

NDC80-kMT binding, but did not change its average level. The observation that inhibiting haspin

removes the correlations between NDC80-kMT binding and tension, and between Aurora B localization

and tension, but does not affect the distribution of K-K distances, argues that these correlations are caused

by the influence of tension on NDC80-kMT binding and Aurora B localization. A simple mathematical

1. model of Aurora B autoactivation, and NDC80 phosphorylation and binding can quantitatively explain

2. these results. Taking together, this leads to a biophysical model of the tension dependency of NDC80-

3. kMT interactions, which arises from the nonlinearity of Aurora B autoactivation and the change in Aurora

4. B concentration at NDC80 with centromere tension.

5.   The FLIM-FRET technique developed in this study measures the fraction of NDC80 complexes

6. whose Hec1 CH domains are bound to kMTs. Hypothetically, changes in NDC80 binding fraction might

7. result from either alterations in the binding affinity of NDC80 or variations in the number of kMTs. Our

8. results argue that the increase in NDC80 binding with increasing tension is caused by changes in affinity,

9. because: (1) the addition of taxol causes a reduction in tension and a reduction in NDC80 binding (Figure

10. 3D and 3E), but an increase in the number of kMTs (McEwen et al., 1997); (2) taxol-treated and STLC-

11. treated cells exhibit the same correlation between tension and NDC80 binding (Figure 3D), suggesting

12. that the decrease in binding with decreasing tension occurs independently of the mechanism of

13. perturbation. Furthermore, previous estimates have found that each kMT can be contacted by

14. approximately 40 NDC80 complexes (Zaytsev et al., 2014), and since there are approximately 240

15. NDC80 complexes per kinetochore (Suzuki et al., 2015), roughly 6 kMTs are sufficient for every NDC80

16. complex to be within reach of potential binding sites. Thus, NDC80 binding is expected to be insensitive

17. to the number of kMTs if there are at least 6 kMTs. As on average there are ~20 kMTs per kinetochore in

18. human cells (Maiato et al., 2004, Rieder, 1982), small changes in the number of kMTs are unlikely to

19. modify the fraction of NDC80 complexes bound to kMTs. However, in circumstances where there are

20. very few kMTs, for example in early prometaphase, changes in the number of kMTs will lead to changes

21. in NDC80 binding. We hypothesize that the observed increase in NDC80 binding over the course of

22. prometaphase to metaphase results from the combination of changes in kMT number and the changes in

23. NDC80 binding affinity, due to the continual increase in tension at those times (Magidson et al., 2011).

24. The difference in NDC80 binding between poleward-facing and anti-poleward-facing kinetochores in

25. STLC-treated cells is likely caused by the difference in kMT numbers (Figure 3-figure supplement 2 and

Figure 5-figure supplement 1). This effect is particularly likely to explain the difference in NDC80

binding between sister kinetochores with 9A-Hec1, since these sister kinetochores are expected to exhibit

the same NDC80 binding affinity (Figure 5-figure supplement 1A).

Error correction is believed to result from the regulation of the detachment of kMTs from

kinetochores (Godek et al., 2015). As NDC80 is the primary coupler of kinetochores to microtubules

(Cheeseman et al., 2006, DeLuca et al., 2006), it is reasonable to hypothesize that the rate of kMT

detachment from kinetochores might largely be governed by NDC80-kMT binding. Consistent with this,

previous work showed that mutating NDC80 changes the number of kMTs in a manner that argues that

increasing NDC80-kMT binding increases the stability of kMTs (Guimaraes et al., 2008, Zaytsev et al.,

2014). Our work further supports the connection between NDC80-kMT binding and kMT stability by

comparing our results with previous measurements of the rate of kMT detachment from kinetochores:

NDC80-kMT binding increases during mitotic progression (Figure 2), while kMT stability increases

(Zhai et al., 1995, Kabeche and Compton, 2013); NDC80-kMT binding increases in response to Aurora B

inhibition (Figure 4), which causes an increase in kMT stability (Cimini et al., 2006); NDC80

preferentially binds to polymerizing kMTs over depolymerizing kMTs (Figure 3C), while reconstituted

kinetochores bind more strongly to polymerizing microtubules than depolymerizing microtubules

(Akiyoshi et al., 2010); NDC80-kMT binding increases with increasing tension (Figure 3D), and the

stability of kMTs increase with increasing tension (Nicklas and Koch, 1969, Akiyoshi et al., 2010). These

comparisons argue that the NDC80-kMT binding is a major determinant of the kMT detachment rate.

Hence, we propose that tension dependency of kMT detachment from kinetochores, which is believed to

underlie error correction, results from the tension dependency of NDC80-kMT binding. If correct, this

implies that error correction ultimately results from the nonlinear autoactivation of Aurora B and the

consequent phosphoregulation of NDC80-kMT binding. Further testing this proposal will require

additional quantitative measurements of kMT detachment, errors, and error correction, in combination

with measurement of NDC80-kMT binding using the FLIM-FRET method presented here.

1 MATERIALS AND METHODS

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
cell line (*Homo sapiens*)	U2OS	ATCC	HTB-96	
transfected construct (*Homo sapiens*)	pBABE-puro mTurquoise2-Nuf2	this paper		Nuf2 N-terminally labeled with mTurquoise2; in retroviral vector with puromycin selection marker
transfected construct (*Homo sapiens*)	pBABE-hygro mTurquoise2-Nuf2	this paper		Same as above, but with hygromycin selection marker
transfected construct (*Homo sapiens*)	pBABE-blast mTurquoise2-Nuf2	this paper		Same as above, but with blasticidin marker
transfected construct (*Homo sapiens*)	pBABE-blast Aurora B FRET sensor (mTurquoise2/YPet)	this paper		modified from Addgene #45215; Fuller et al. (2008)
transfected construct (*Homo sapiens*)	Nuf2-targeted Aurora B FRET sensor (mTurquoise2/Ypet)	this paper		modified from Addgene #45215; Fuller et al. (2008)
transfected construct (*Homo sapiens*)	mTurquoise2-TC	this paper		mTurquoise2 with tetracysteine motif at the C-terminus
transfected construct (Homo sapiens)	WT-Hec1-LSSmOrange	this paper		modified from WT-Hec1-GFP from Jennifer DeLuca
transfected construct (Homo sapiens)	9A-Hec1-LSSmOrange	this paper		modified from 9A-Hec1-GFP from Jennifer DeLuca
transfected construct (Homo sapiens)	2D(S44,55D)-Hec1-LSSmOrange	this paper		modified from 2D-Hec1-GFP from Jennifer DeLuca
transfected construct (Homo sapiens)	9D-Hec1-LSSmOrange	this paper		modified from 9D-Hec1-GFP from Jennifer DeLuca
transfected construct (*Homo sapiens*)	INCENP-mCherry	other		Gift from Michael Lampson
recombinant DNA reagent	pSpCas9(BB)-2A-GFP (pX458)	Ran et al. (2013)	Addgene: #48138	
sequence-based reagent	Donor single-stranded DNA for TC tag insertion at the C-terminus of TUBB	IDT		ssDNA: cgtctctgagtatcagcagtaccaggatgccaccgc agaagaggaggaggatttcggtgaggaggccgaa gaggaggcctGCTGTCCCGGCTGTTGct aaggcagagcccccatcacctcaggcttctcagttc ccttagccgtcttactcaactgcccctttcctctccctca ga; sgRNA target sequence: GAGGCCGAAGAGGAGGCCTA

sequence-based reagent	Hec1 siRNA	Qiagen	Cat#: SI02653567	
peptide, recombinant protein	TC-peptide	Genscript	Custom designed	Synthesized, Ac-AEEEACCPGCC-NH2
commercial assay or kit	Amaxa Cell Line Nucleofector Kit V	Lonza	Cat#:VCA-1003	
commercial assay or kit	Ingenio Electroporation Kit	Mirus	Cat#: MIR 50118	
commercial assay or kit	Lipofectamine RNAiMax	Thermo Fisher	Cat#:13778075	
chemical compound, drug	FlAsH-EDT2	Thermo Fisher	Cat#:T34561	
chemical compound, drug	1,2-Ethanedithiol (EDT)	Alfa Aesar	Cat#:540-63-6	
chemical compound, drug	ZM447439	Enzo Life Sciences	Cat#:BML-EI373	
chemical compound, drug	Paclitaxel (Taxol)	Enzo Life Sciences	Cat#:BML-T104	
chemical compound, drug	5-iodotubercidin (5-ITu)	Enzo Life Sciences	Cat#:BML-EI29	
chemical compound, drug	S-Trityl-L-cysteine	Sigma Aldrich	Cat#:164739-5G	
chemical compound, drug	Alexa Fluor 488	Thermo Fisher	Cat#:A20000	
chemical compound, drug	Sodium 2-mercaptoethanesulfonate	Sigma Aldrich	Cat#:M1511	
software, algorithm	Interactive kinetochore FLIM-FRET analysis GUI (MATLAB 2016)	This paper		http://doi.org/10.5281/zenodo.1198705
software, algorithm	Aurora B concentration at NDC80 analysis (Python 3)	This paper		http://doi.org/10.5281/zenodo.1198702
software, algorithm	CAMPARI (v2)	Pappu Lab		http://campari.sourceforge.net/V2/index.html
software, algorithm	Rosetta 3.8	RosettaCommons		RRID:SCR_015701
other	25-mm #1.5 poly-D-lysine coated round coverglass	neuVitro	Cat#:GG-25-1.5-pdl	
other	FluoroBrite DMEM	Thermo Fisher	Cat#:A1896701	
other	Microtubule structure	Zhang et al. (2015)	PDB 3JAS	
other	Human NDC80 bonsai decorated tubulin dimer	Alushin et al. (2010)	PDB 3IZ0	
other	mTurquoise structure	Stetten et al. (unpublished)	PDB 4B5Y	

Cell lines

3 U2OS cell lines (ATCC, HTB-96) were maintained in Dulbecco's modified Eagle's medium (DMEM,

4 Thermo Fisher) supplemented with 10% Fetal Bovine Serum (FBS, Thermo Fisher), and 50 IU ml^{-1}

5 penicillin and 50 μg ml^{-1} streptomycin (Thermo Fisher) at 37°C in a humidified atmosphere with 5% CO_2.

6 Cells were validated as mycoplasma free by PCR-based mycoplasma detection kit (Sigma Aldrich).

Live-cell imaging

All live-cell FLIM and spinning-disk confocal microscopy imaging were performed as follows. Cells

were grown on a 25-mm diameter, #1.5-thickness, round coverglass coated with poly-D-lysine (GG-25-

1.5-pdl, neuVitro) to 80~90% confluency. The cells were incubated in imaging media, which is

FluoroBrite™ DMEM (Thermo Fisher) supplemented with 4 mM L-glutamine (Thermo Fisher) and 10

mM HEPES, for 15~30 minutes before imaging. The coverglass was mounted on a custom-built

temperature controlled microscope chamber at 37°C, while covered with 1 ml of imaging media and 2 ml

of white mineral oil (VWR). An objective heater (Bioptech) was used to maintain the objective at 37°C.

We confirmed that the cells can normally divide longer than 6 hours in this condition. Only cells

displaying proper chromosome alignment, normal spindle morphology, and high signal-to-noise ratio

were selected for imaging and analysis.

NDC80-kMT FLIM-FRET measurement

mTurquoise2-NDC80/β-tubulin-TC-FlAsH stable U2OS cell line

A tetracysteine (TC) tag, CCPGCC, was genetically attached to the C-terminal end of tubulin beta class I

(TUBB), an isotype of β-tubulin that is predominantly expressed in U2OS (assessed by qPCR, data not

shown) and most other cancer cells (Leandro-Garcia et al., 2010). The attachment of the TC tag was

achieved by CRISPR-induced homologous recombination to ensure the consistent expression of labeled

β-tubulin. ssDNA (IDT) with TC tag (5'-TGCTGTCCCGGCTGTTGC-3') and ~80 bp-long homology

arms was used as a donor DNA. pSpCas9(BB)-2A-GFP (Addgene plasmid # 48138) (Ran et al., 2013)

was utilized as a backbone for the plasmid carrying a sgRNA (5'-GAGGCCGAAGAGGAGGCCUA-3')

and Cas9. The plasmid and the donor ssDNA were simultaneously delivered into U2OS cells by

electroporation (Nucleofector™ 2b and Amaxa Cell Line Nucleofector Kit V, Lonza). The insertion of the

TC tag was verified through a PCR-based genotyping with primers 5'-

GCATGGACGAGATGGAGTTCAC-3' and 5'-CCAGCCGTGTTTCCCTAAATAAG-3', qPCR, and a

fluorescence imaging after FlAsH-EDT$_2$ staining.

The U2OS cells expressing TC-tagged β-tubulin were further engineered to stably express Nuf2

N-terminally labeled with mTurquoise2 (Goedhart et al., 2012) by retroviral transfection, three times with

different antibiotic selections, 1 µg ml^{-1} puromycin, 2 µg ml^{-1} blasticidin, and 200 µg ml^{-1} hygromycin (all

from Thermo Fisher). The retroviral vectors and their information are available on Addgene (plasmid #:

80760, 80761, 80762). Monoclonal cell line was obtained by single cell sorting.

<u>FlAsH-EDT$_2$ staining</u>

The protocol for the association of FlAsH-EDT$_2$ with β-tubulin-TC in cell was adapted from the previous

study (Hoffmann et al., 2010) so as to maximize the labeling fraction while maintaining cell viability. The

engineered U2OS cells expressing β-tubulin-TC were grown to 80~90% confluency in a 30-mm cell

culture dish, and then were gently washed with Opti-MEM (Thermo Fisher) twice, and then stained in 2

ml Opti-MEM with 1 µM FlAsH-EDT$_2$ (Thermo Fisher) for 2 hours. To reduce the non-specific binding

of FlAsH, the stained cells were subsequently incubated in Opti-MEM containing 250 µM 1,2-

Ethanedithiol (EDT, Alfa Aesar) for 10 minutes, followed by a gentle wash with Opti-MEM. The cells

were incubated in DMEM with 10% FBS for 6~10 hours before imaging, because they were found to be

interphase-arrested for the first ~5 hours after the incubation with 250 µM EDT. Every buffers and media

above were pre-warmed at 37°C before use. All incubation steps were performed at 37°C in a humidified

atmosphere with 5% CO$_2$.

<u>FLIM measurement</u>

Schematic instrumental setup of FLIM is shown in Figure 1–figure supplement 2A, and more details can

be found in previous work (Yoo and Needleman, 2016). FLIM measurements were performed on a Nikon

Eclipse Ti microscope using two-photon excitation from a Ti:Sapphire pulsed laser (Mai-Tai, Spectral-

Physics) with an 80-MHz repetition rate and ~70-fs pulse width, a galvanometer scanner (DCS-120,

Becker & Hickl), TCSPC module (SPC-150, Becker & Hickl) and two hybrid detectors (HPM-100-40,

Becker & Hickl). Objective piezo stage (P-725, Physik Instrumente) and motorized stage (ProScan II,

Prior Scientific) were used to perform multi-dimensional acquisition, and a motor-driven shutter (Sutter

Instrument) was used to block the excitation laser between acquisitions. The wavelength of the excitation

laser was set to 865 nm. 470/24 and 525/30 bandpass emission filters (Chroma) were mounted on each

detector, and a dichroic beam splitter (FF506-Di03, Semrock) was used for the simultaneous detection of

mTurquoise2 and FlAsH fluorescence. The excitation laser was expanded to overfill the back-aperture of

a water-immersion objective (CFI Apo 40× WI, NA 1.25, Nikon). The power of the excitation laser was

adjusted to 1.1~1.5 mW at the objective. All the electronics were controlled by SPCM software (Becker

& Hickl) and μManager (Edelstein et al., 2014). Scanning area was set to either 13.75 μm × 13.75 μm or

27.5 μm × 27.5 μm, and the pixel size was set to 107 nm. Each image was acquired for 3~5 seconds of

integration time. Acquisition interval was set to 13 seconds for the FLIM-FRET data in Figure 3 and 5,

and 60~90 seconds for the FLIM-FRET data in Figure 2 and 4. Three or four z-sections, separated by 1

μm, were acquired for each time point. No photo-bleaching or photo-damage was observed in this

imaging condition. Mitotic phases were judged by the arrangement of kinetochores.

<u>Kinetochore tracking and pairing</u>

For the kinetochore FLIM-FRET measurements shown in Figure 2 to 5, custom-built MATLAB graphical

user interphase (GUI) (available at http://doi.org/10.5281/zenodo.1198705) was used to import Becker

and Hickl FLIM data, track kinetochores, identify kinetochore pairs, extract the FLIM curve from each

kinetochore, and estimate the FLIM parameters using a nonlinear least-squared fitting or Bayesian FLIM

analysis, as described below and in previous work (Yoo and Needleman, 2016). The GUI also allows the

users to scrutinize and manually correct the kinetochore trajectories and pairing. The kinetochore tracking

algorithm was adapted from a particle tracking algorithm (Pelletier et al., 2009), and the pair

identification was performed by selecting pairs of kinetochores with distances and velocity correlations in

predefined ranges. Drift correction was done by measuring correlation between two consecutive spindle

images. The velocity $v(t)$ of kinetochore (in Figure 3) was estimated from the position $x(t)$ using the

five-point method:

5
$$v(t) \approx \frac{-x(t + 2\Delta t) + 8x(t + \Delta t) - 8(t - \Delta t) + x(t - 2\Delta t)}{12\Delta t}$$

Leading and trailing kinetochores (in Figure 3) were determined based on the velocities and the relative

positions of paired sister kinetochores. The metaphase plate (in Figure 2) was determined by finding an

equidistant plane between the two spindle poles (that were manually located based on spindle images).

Poleward-facing and anti-poleward-facing kinetochores (in Figure 3 and 5) in STLC-treated cells were

determined based on the relative positions of paired sister kinetochores and the position of the spindle

pole, which was approximated by the average of the positions of kinetochores in each time point.

<u>Bayesian FLIM analysis</u>

Fluorescence decay curves from individual kinetochores at each time point contain only a few hundreds

of photons. In this low photon count regime, FLIM analysis with conventional least-squared nonlinear

regressions results in significantly biased estimate for the parameters (Kaye et al., 2017, Rowley et al.,

2016). Therefore, we used a Bayesian approach, which has been described and tested previously (Yoo and

Needleman, 2016, Kaye et al., 2017), and is briefly explained below.

Let θ be the set of parameters of the FLIM-FRET model, and $y = \{y_i\}$ be the observed FLIM

data, where y_i is the number of photons detected in the i-th time bin of the FLIM curve. Then the

posterior distribution of θ (assuming a uniform prior distribution) is

22
$$p(\theta|y) \propto \prod_{i=1}^{N} P(t_{ar} \in [(i-1)\Delta t, i\Delta t]|\theta)^{y_i}$$

1   where t_{ar} is the photon arrival time, and N is the number of time bins. Since the size of the time bin (Δt,

2   ~50 ps) is much smaller than the time scale of fluorescence decay (~ns), the probability that the arrival

3   time t_{ar} falls in the i-th time bin can be approximated by a Riemann sum:

4
$$P(t_{ar} \in [(i-1)\Delta t, i\Delta t]|\theta) \cong \sum_{k=(i-1)K+1}^{k=iK} h_\theta(k\widetilde{\Delta t})\widetilde{\Delta t}$$

5   where h_θ is the discretized FLIM model, $\widetilde{\Delta t}$ is the size of time bin with which instrument response

6   function (IRF) is measured, and the ratio $K = \frac{\Delta t}{\widetilde{\Delta t}}$ is the ADC ratio, which is set to 16 for our data.

7     $h_\theta(k\widetilde{\Delta t})$ can be written as the convolution between the IRF and an exponential decay model, g_θ:

8
$$h_\theta(k\widetilde{\Delta t}) = (IRF * (Ag_\theta + (1-A)))(k\widetilde{\Delta t})$$

9
$$\cong \sum_l mIRF[l - b_{shift}]\left(Ag_\theta\left((k-l)\widetilde{\Delta t}\right) + (1-A)\right)$$

where $mIRF$ is the IRF measured with the finest time bins of size $\widetilde{\Delta t}$, and b_{shift} is an integer parameter

that determines the approximate shift of measured IRF relative to the theoretical IRF. $(1-A)$ indicates

the relative contribution of noise that is uniformly distributed over time. The exponential decay model

$g_\theta(t_d)$ is set to $\exp\left(-\frac{t_d}{\tau}\right)$ for the single-exponential decay model or $(1 - f_{FRET})\,e^{-\frac{t_d}{\tau_D}} + f_{FRET}e^{-\frac{t_d}{\tau_{FRET}}}$

for the two-exponential decay model, where $0 \le f_{FRET} \le 1$ is the FRET fraction. The posterior

distribution was computed by Gibbs sampling if the number of free parameters is greater than 3, or by

grid sampling otherwise (for example, when both long and short lifetimes are fixed).

<u>NDC80 FRET fraction measurement procedures</u>

The instrument response function (IRF) was acquired by measuring second-harmonic generation from a

urea crystal. Negative control FLIM measurements on the engineered cells (mTurquoise2-NDC80/β-

tubulin-TC) not incubated with FlAsH were performed for every experiment and the fluorescence decay

curves extracted from kinetochores were analyzed with a single-exponential FLIM-FRET model to

determine the long non-FRET lifetime, which is usually 3.7 to 3.8 ns. The short FRET lifetime was

estimated by performing a two-exponential Bayesian FLIM-FRET analysis on the aggregated FLIM data

of kinetochores in each cell stained with FlAsH while fixing the non-FRET lifetime to the value pre-

determined from the negative control. Then we performed a two-exponential Bayesian FLIM-FRET

analysis, with both FRET and non-FRET lifetimes fixed to the predetermined values, on FLIM data from

each kinetochore. Kinetochores were grouped by time (Figure 2 and 4), positions (Figure 2B-D),

velocities (Figure 3C), and K-K distances (Figure 3D, 5A and 5E). The posterior distributions in a group

of kinetochores were multiplied and then marginalized to obtain the mean and SEM of the FRET fraction.

We previously confirmed that this way of combining posterior distribution gives an unbiased estimate of

the mean FRET fraction (Kaye et al., 2017). NDC80 binding fraction was calculated by dividing NDC80

FRET fraction by the conversion factor 0.42, which had been determined by the calibration shown in

Figure 1–figure supplement 5C.

Aurora B kinase activity measurement

An Aurora B FRET sensor was constructed by replacing CyPet in a previous construct (Addgene plasmid

45215) (Fuller et al., 2008) with mTurquoise2. The FRET sensor contains a kinesin-13 family Aurora B

substrate whose phosphorylation results in its binding to the forkhead-associated domain in the sensor,

which constrains the sensor to be in an open conformation and obstructs intramolecular FRET between

mTurquoise2 and YPet (Figure 4–figure supplement 1B). Hence, the non-FRET fraction of the Aurora B

FRET sensor is proportional to the Aurora B activity. The cytoplasmic Aurora B FRET sensor was stably

expressed in U2OS cells by retroviral transfection (plasmid available on Addgene, plasmid # 83286). The

Nuf2-targeted Aurora B FRET sensor was transiently transfected by electroporation (Nucleofector 2b,

Lonza; Ingenio Electroporation Kit, Mirus) a day before imaging. The non-FRET fraction of the Aurora B

FRET sensor was measured by FLIM-FRET in the same way as NDC80 FRET measurements described

above. The exponential decay models $y_{\text{binding}}(t) = A\left(1 - \exp\left(-\frac{I_{t \geq 0}t}{\tau}\right)\right) + c$ and $y_{\text{Aurora}}(t) =$

$A \exp\left(-\frac{I_{t \geq 0} t}{\tau}\right) + c$ were fitted to the time courses of NDC80 FRET fraction and FRET sensor non-FRET

fraction after ZM447439, respectively (Figure 4A, 4D, 4E and Figure 4-figure supplement 1D), where

$I_{t \geq 0}$ is equal to 0 if t is less than zero, and 1 otherwise. The estimated parameter values are given in the

table below:

Figure	Parameter	mean	95% CI
4A	A	0.088	(0.069,0.106)
	τ (min)	3.26	(1.31,5.21)
	c	0.089	(0.080,0.099)
4D	A	0.024	(0.011,0.038)
	τ (min)	0.50	(-0.70,1.71)
	c	0.059	(0.048,0.071)
4E	A	0.17	(0.16,0.18)
	τ (min)	1.95	(1.46,2.45)
	c	0.37	(0.36,0.38)
4-S1D	A	0.076	(0.061,0.090)
	τ (min)	1.12	(0.23,2.00)
	c	0.56	(0.55,0.57)

The fraction of Aurora B phosphorylation sites in NDC80, f_{phos} (x-axis of Figure 4F), was converted

from the non-FRET fraction of Aurora B FRET sensor, f_{sensor} (y-axis of Figure 4E), as follows. First, we

assumed that f_{sensor} increases linearly with f_{phos}. Our result (Figure 4C) and previous work (Zaytsev et

al., 2014) suggest that Ndc80 has about one phospho-residue out of nine phosphorylation sites in late

prometaphase, based on which we assumed that $f_{phos}^{WT} = 1/9$ before Aurora B inhibition, and $f_{phos}^{ZM} = 0$

after the full Aurora B inhibition. Since f_{sensor} were measured to be $f_{sensor}^{WT} = 0.540 \pm 0.007$ (SEM)

before Aurora B inhibition and $f_{sensor}^{ZM} = 0.368 \pm 0.012$ (SEM) after the full Aurora B inhibition (Figure

4E), we converted f_{sensor} to f_{phos} by:

13
$$f_{phos} = \frac{f_{sensor} - f_{sensor}^{ZM}}{f_{sensor}^{WT} - f_{sensor}^{ZM}} \left(f_{phos}^{WT} - f_{phos}^{ZM}\right) + f_{phos}^{ZM} = 0.646(f_{sensor} - 0.368)$$

14 The f_{bound} vs f_{phos} data in Figure 4F was fit using a NDC80 binding model:

15
$$f_{bound} = \left(1 + K_0 + K_0' f_{phos}\right)^{-1}$$

16 which is derived in Mathematical modeling section below.

Phosphomimetic Hec1 mutant experiments

We used three different non-phosphorylable mutant Hec1 (gift from Jennifer DeLuca) in which all nine

identified Aurora B target sites in the N-terminal tail are mutated to either Asp (phospho-mimicking

mutation) or Ala (phospho-null mutation) (DeLuca et al., 2011, Zaytsev et al., 2015, Zaytsev et al., 2014):

9A-Hec1 (all nine sites substituted with Ala), 2D-Hec1 (two sites, S44 and S55, substituted with Asp,

while the other seven sites with Ala), and 9D-Hec1 (all nine sites substituted with Asp). WT-Hec1 and the

mutant Hec1 are C-terminally labeled with LSSmOrange. LSSmOrange signal at kinetochores were

assessed to ensure the expression of the substituting Hec1 in cells.

Cells were grown to 50% confluence on a 10-cm petri dish in DMEM supplemented with 10%

FBS and Penicillin-Streptomycin (P/S) as described above. To knock down endogenous Hec1/Ndc80

protein, we used a FlexiTube siRNA duplex targeted to the 5' UTR of the Hec1 gene (5'-

TCCCTGGGTCGTGTCAGGAAA-3', QIAGEN Hs_KNTC2_7 SI02653567). We incubated 240 pmol

of the siRNA in 1.2 mL Opti-MEM (ThermoFisher 51985091) for 5 minutes with periodic flicking. We

simultaneously incubated 8 μL of Lipofectamine RNAiMax (ThermoFisher 13778030) in 1.2 mL Opti-

MEM for 5 minutes. We then combined the siRNA and Lipofectamine solutions and incubated at room

temperature for 30 minutes with periodic flicking. Prior to adding the siRNA-lipid complex, we washed

the cells once with PBS and then replaced the media with 8 mL Opti-MEM supplemented with 10% FBS.

We then added the entire 2.4 mL siRNA mixture to the cells dropwise and incubated the cells at 37°C for

30 hours. Following the incubation, we nucleofected 2 μg of plasmid encoding WT-, 9A-, 2D-, or 9D-

Hec1 along with an additional 30 pmol of Hec1 siRNA into 1 million cells using a Lonza Nucleofector

2b. We spread these cells evenly over three 35mm dishes containing 25mm poly-D-lysine coated

coverslips and 2 mL of Opti-MEM supplemented with 10% FBS and P/S. We incubated overnight at

37°C for 18 hours before staining with TC-FlAsH and FLIM-FRET imaging as described above.

Aurora B concentration at NDC80 measurement

<u>mNeonGreen-Nuf2/INCENP-mCherry U2OS cell</u>

mNeonGreen fluorescent protein (Shaner et al., 2013) was genetically attached to the N-terminal end of

Nuf2 by CRISPR-induced homologous recombination with an sgRNA (5'-

GAAAGACAAAGTTTCCATCTTGG-3') and mNeonGreen sequence (Allele Biotechnology) flanked by

2kb homology arms as a donor template. Monoclonal cell line was obtained by fluorescence-activated cell

sorting and screened by fluorescent microscopy imaging. The mNeonGreen-Nuf2 U2OS cell line was

transiently transfected with INCENP-mCherry (gift from Michael Lampson) by electroporation

(Nucleofector 2b and Amaxa Cell Line Nucleofector Kit V, Lonza) a day before imaging, using the

manufacturer's protocol.

<u>Spinning-disk confocal microscopy imaging</u>

Cells were imaged using a spinning-disk confocal microscope (Nikon Ti2000, Yokugawa CSU-X1) with

1.5x magnification lens and 1.2x tube lens, an EM-CCD camera (Hamamatsu), a 60x water-immersion

objective (Nikon), an objective piezo stage (P-725, Physik Instrumente), and motorized x-y stage

(ProScan II, Prior Scientific) controlled by μManager (Edelstein et al., 2014). A 488-nm laser and 514/30

filter were used to image mNeonGreen-Nuf2, and a 560-nm laser and 593/40 filter were used to image

INCENP-mCherry. 11-15 z-slices, separated by 2 μm, were taken for each time point. Three time points,

separated by a minute, were acquired before and after DMSO (for untreated data), 10 μM taxol, or 10 μM

5-ITu treatment.

<u>Aurora B concentration at NDC80 measurement</u>

Image analysis was performed by a custom Python code (available at

http://doi.org/10.5281/zenodo.1198702). Kinetochore identification was achieved by applying *trackpy*

package (github.com/soft-matter/trackpy) to mNeonGreen-Nuf2 fluorescence images. The sub-pixel

location of NDC80 was calculated by centroid estimation. Sister kinetochore pairs were determined based

on the relative positions of kinetochores and the INCENP-mCherry intensity between kinetochores. For

each identified kinetochore pair, INCENP-mCherry intensities at the NDC80 centroid locations, I_{NDC80},

and INCENP-mCherry intensity at the midpoint between two sister kinetochores, I_{mid}, were measured by

two-dimensional cubic interpolation with *scipy.interpolate.griddata* function. For each cell, we used

$\overline{I_{mid}}$, which is I_{mid} averaged over kinetochores in the images before chemical treatments, and

cytoplasmic background level, I_{bg}, to obtain normalized the INCENP-mCherry intensities at NDC80,

I_{NDC80}^{norm}, by:

$$I_{NDC80}^{norm} = \frac{I_{NDC80} - \overline{I_{mid}}}{\overline{I_{mid}} - I_{bg}}$$

Kinetochores with similar K-K distances were grouped in the same way as in Figure 3D and 5E, and then

the normalized INCENP-mCherry intensities at NDC80, I_{NDC80}^{norm}, were averaged within each group. The

normalized INCENP-mCherry intensity was converted to Aurora B concentration in Figure 7A by

assuming that $\overline{I_{mid}}$ corresponds to the peak Aurora B concentration, which was previously estimated to be

10 μM (Zaytsev et al., 2016).

Drug treatments

Cells were incubated with 5 μM Nocodazole (Sigma Aldrich) for >10 minutes for microtubule

depolymerization. Aurora B inhibition was performed by adding 3 μM of ZM447439 (Enzo Life

Sciences) during imaging. Taxol (Enzo Life Sciences) treatment was performed at 10 μM final

concentration for >10 minutes. S-Trityl-L-cysteine (STLC, Sigma Aldrich) treatment was performed at 5

μM final concentration for >60 minutes to induce monopolar spindles. For the haspin kinase inhibition,

cells were treated with 10 μM 5-iodotubercidin (5-ITu, Enzo Life Sciences) for >10 minutes. The double

treatment of 5-ITu and taxol or STLC was performed sequentially by treating cells with 10 μM taxol or 5

μM STLC and then adding 10 μM 5-ITu.

Mathematical modeling

Here we describe the mathematical model presented in Figure 7 in detail. The model predicts NDC80

binding fraction from Aurora B concentration at NDC80 in three steps: (1) Aurora B activation dynamics,

consisting of autoactivation in *trans* and deactivation, which determines the concentration of *active*

Aurora B from the concentration of Aurora B; (2) NDC80 phosphorylation, which is dependent on the

active Aurora B concentration; and (3) NDC80-kMT binding, which is governed by the phosphorylation

level of NDC80.

(1) Aurora B activation

In this section, we present a quantitative model for the relationship between the Aurora B concentration

(which we measured in Figure 6) and the *active* Aurora B concentration (which determines the steady-

state level of NDC80 phosphorylation). It has been previously argued that Aurora B activation is

predominately due to active Aurora B phosphorylating inactive Aurora B in *trans* (Zaytsev et al., 2016,

Xu et al., 2010, Sessa et al., 2005, Bishop and Schumacher, 2002), which we incorporate into our model.

We model Aurora B at the location of NDC80 as consisting of two separate pools: one that is dependent

on haspin, and the other that is not. We assume that those two Aurora B pools do not interact with each

other, and independently undergo auto-activation in *trans*. We further assume that the phosphatases

activity proceeds at a constant rate for each pool.

We denote the haspin-dependent and haspin-independent pools of Aurora B by A_{hd} and A_{hi},

respectively. Then the inter-molecular autoactivation by in *trans* phosphorylation and inactivation by

dephosphorylation for each of the two Aurora B pools are described by:

22
$$\begin{cases} A_x^{active} + A_x^{inactive} \xrightarrow{k_x^a} A_x^{active} + A_x^{active} \\ A_x^{active} \xrightarrow{k_x^d} A_x^{inactive} \end{cases} \qquad x = hd \text{ or } hi$$

1   where A_x^{active} and $A_x^{inactive}$ are the active and inactive Aurora B in pool x, respectively, and k_x^a and k_x^d

2   are the rates of Aurora B activation and deactivation for the pool x, respectively. Thus, an ordinary

3   differential equation (ODE) for active Aurora B concentration can be written as:

$$4 \quad \frac{\partial\left[A_x^{active}\right]}{\partial t} = k_x^a\left[A_x^{active}\right]\left[A_x^{inactive}\right] - k_x^d\left[A_x^{active}\right]$$

$$5 \quad = k_x^a\left[A_x^{active}\right]\left(\left[A_x\right] - \left[A_x^{active}\right]\right) - k_x^d\left[A_x^{active}\right]$$

$$6 \quad = k_x^a\left[A_x^{active}\right]\left(\left[A_x\right] - \frac{k_x^d}{k_x^a} - \left[A_x^{active}\right]\right)$$

7   where $\left[A_x\right] = \left[A_x^{active}\right] + \left[A_x^{inactive}\right]$ is the concentration of the pool x. The steady-state solution for this

8   ODE is:

$$9 \quad \left[A_x^{active}\right] = \begin{cases} 0, & [A] < K_x \\ \left[A_x\right] - K_x, & [A] \geq K_x \end{cases}$$

10   where $K_x \equiv k_x^d / k_x^a$ is the equilibrium constant for the Aurora B activation for pool x. We can infer that

11    $\left[A_{hi}^{active}\right] = [A_{hi}] - K_{hi}$ is positive, because Aurora B still acts on NDC80 after the removal of the

12   haspin-dependent pool (Figure 5). Therefore, the total concentration of active Aurora B at NDC80 can be

13   written as:

$$\left[A^{active}\right] = \left[A_{hd}^{active}\right] + \left[A_{hi}^{active}\right]$$

$$= \begin{cases} [A_{hi}] - K_{hi}, & [A] < K_{hd} + [A_{hi}] \\ [A] - K_{hd} - K_{hi}, & [A] \geq K_{hd} + [A_{hi}] \end{cases}$$

$$= \begin{cases} A^* - K, & [A] < A^* \\ [A] - K, & [A] \geq A^* \end{cases} \quad \text{(Eq. 1)}$$

14   where $[A] = [A_{hd}] + [A_{hi}]$ is the total concentration of Aurora B at NDC80, $K = K_{hd} + K_{hi}$, and $A^* =$

15    $K_{hd} + [A_{hi}]$ is a threshold Aurora B concentration, which is the minimum concentration of Aurora B

16   required for the activity of Aurora B to increase with its concentration.

18   (2) NDC80 phosphorylation

1   In this section, we present a mathematical model to relate the total concentration of active Aurora B at

2   NDC80, $[A^{active}]$, to the phosphorylation level of NDC80. Active Aurora B may phosphorylate multiple

3   Aurora B phosphorylation sites in each Ndc80 N-terminal tail (Guimaraes et al., 2008), which we

4   describe with the equations:

5
$$\begin{cases} A^{active} + \text{(dephosphorylated site)} \overset{k_p}{\rightarrow} A^{active} + \text{(phosphorylated site)} \\ \text{(phosphorylated site)} \overset{k_{dp}}{\longrightarrow} \text{(dephosphorylated site)} \end{cases}$$

6   The corresponding ODE for the number of phosphorylated sites is:

9
$$\frac{\partial N_p}{\partial t} = k_p[A^{active}]N_{dp} - k_{dp}N_p$$

10
$$= k_p[A^{active}](N - N_p) - k_{dp}N_p$$

7   where N_p, N_{dp}, and $N = N_p + N_{dp}$ is the number of phosphorylated sites, dephosphorylated sites, and the

8   total number of sites per kinetochore, respectively. The steady-state solution for the ODE gives:

$$f_{phos} = \frac{N_p}{N} = \left(1 + \frac{K_{phos}}{[A^{active}]}\right)^{-1} \tag{Eq. 2}$$

11   where f_{phos} is the fraction of phosphorylated sites, and $K_{phos} \equiv \frac{k_{dp}}{k_p}$ is the equilibrium constant for

12   NDC80 phosphorylation. Plugging Eq. 1 into Eq. 2 yields:

$$f_{phos} = \begin{cases} \left(1 + \frac{K_{phos}}{A^* - K}\right)^{-1}, & [A] < A^* \\ \left(1 + \frac{K_{phos}}{[A] - K}\right)^{-1}, & [A] \geq A^* \end{cases} \tag{Eq. 3}$$

14   (3) NDC80 binding

15   In this section, we present a model to relate the fraction of phosphorylated sites in NDC80 per

16   kinetochore, f_{phos}, to the fraction of NDC80 bound to kMTs (which we measure using FLIM-FRET).

17   Assuming that the number of available binding sites for NDC80 is constant, we may describe the NDC80

18   binding and unbinding by the following equations:

1
$$\begin{cases} (\text{NDC80 unbound}) \xrightarrow{k_{on}} (\text{NDC80 bound}) \\ (\text{NDC80 bound}) \xrightarrow{k_{off}} (\text{NDC80 unbound}) \end{cases}$$

2 The corresponding ODE for the number of NDC80 bound to kMTs is:

3
$$\frac{\partial n_{on}}{\partial t} = k_{on} n_{off} - k_{off} n_{on}$$

4
$$= k_{on}(n - n_{on}) - k_{off} n_{on}$$

5 where n_{on} and n_{off} are the number of NDC80 bound and unbound to kMTs, respectively, and $n =$

6 $n_{on} + n_{off}$ the total number of NDC80 per kinetochore. Solving for the steady state gives:

$$f_{bound} \equiv \frac{n_{on}}{n} = \left(1 + K_{binding}\right)^{-1} \qquad \text{(Eq. 4)}$$

where $K_{binding} = k_{off}/k_{on}$ is the equilibrium constant for the NDC80 binding, and f_{bound} is the NDC80

binding fraction.

The binding affinity of NDC80 decreases with the phosphorylation level of NDC80 (Figure 4) (Zaytsev et

al., 2015), arguing that $K_{binding}$ is a function of f_{phos}. Since f_{phos} is small in late prometaphase and

metaphase ($< 1/9$), we approximate the function by a first-order polynomial, i.e., $K_{binding}\left(f_{phos}\right) \approx K_0 +$

$K_0' f_{phos}$, and consequently Eq. 4 becomes:

$$f_{bound} = \left(1 + K_0 + K_0' f_{phos}\right)^{-1} \qquad \text{(Eq. 5)}$$

14 Combining Eq. 3 and 5, we have the relationship between the total Aurora B concentration $[A]$ and the

15 NDC80 binding fraction f_{bound} as:

$$f_{bound} = \begin{cases} \left(1 + K_0 + K_0'\left(1 + \dfrac{K_{phos}}{A^* - K}\right)^{-1}\right)^{-1}, & [A] < A^* \\[4mm] \left(1 + K_0 + K_0'\left(1 + \dfrac{K_{phos}}{[A] - K}\right)^{-1}\right)^{-1}, & [A] \geq A^* \end{cases} \qquad \text{(Eq. 6)}$$

We first determined the parameters K_0 and K_0' by fitting Eq. 5 to the NDC80 binding fraction vs.

phosphorylation level data in Figure 4F, which yielded $K_0 = 1.43 \pm 0.06$ (SE) and $K_0' = 18 \pm 2$ (SE). To

estimate the remaining three free parameters, K, K_{phos}, and A^*, we fit Eq. 6 to the NDC80 binding

fraction vs. Aurora B concentration at NDC80 data (Figure 7), and obtained $K = 3.5 \pm 0.4$ µM (SE),

$K_{phos} = 19 \pm 5$ µM (SE), and $A^* = 4.6 \pm 0.2$ µM (SE).

Supplemental experiments

<u>Measurement of the fraction of β-tubulin labeled with TC-FlAsH</u>

To measure the fraction of β-tubulin labeled with TC-FlAsH, we sought to determine the concentration of

labeled β-tubulin in the cell, and divide it by the total concentration of β-tubulin. We calculated the

concentration of labeled β-tubulin by combining 3D fluorescence microscopy to measure the total

fluorescence of β-tubulin-TC-FlAsH per cell, and fluorescence correlation spectroscopy (FCS) to measure

the fluorescence per molecule of TC-FlAsH.

3D fluorescence microscopy: We acquired z-stacks of β-tubulin-TC-FlAsH in mitotic cells using two-

photon fluorescence microscopy (Figure 1–figure supplement 1A), and then segmented the 3D images

using an active contour approach (Figure 1–figure supplement 1B). Assuming that the cytoplasmic

background results from FlAsH binding specifically to monomeric β-tubulin and nonspecifically to

cysteine-rich proteins freely diffusing in the cytoplasm, the average number of photons emitted from β-

tubulin-TC-FlAsH in microtubules is the difference between the average photon rate throughout the entire

cell (423 ± 33 ms^{-1}) and the average photon rate in the cytoplasm (determined from the mode of

fluorescence distribution within each segmented image, 327 ± 30 ms^{-1}), which is 96 ± 12 ms^{-1} (Figure 1–

figure supplement 1C). The instrumental setting of two-photon fluorescence microscopy was identical to

that of the FLIM system described above, where the imaging parameters are: laser wavelength, 865 nm;

excitation intensity, 3 mW; integration time, 3 seconds; z-stack separation, 0.5 µm; scanning area, 27.5

µm × 27.5 µm.

Fluorescence correlation spectroscopy: To convert the measured photon rate from fluorescence

microscopy to a measurement of the absolute concentration of β-tubulin-TC-FlAsH, we used two-photon

FCS to determine the volume of the point spread function (PSF) and the molecular brightness (i.e. the

number of photons emitted per molecule per second) of TC-FlAsH (Hess and Webb, 2002).

8   First, we performed an FCS measurement on 97-nM Alexa Fluor 488 (Thermo Fisher) in water.

9   FCS measurements were performed on the same instrumental setting as the 3D fluorescence microscopy

10   described above, with laser intensity 5 mW. 5 autocorrelation functions, each of which had been collected

11   for 300 seconds, were averaged, and then the following FCS model, $G_D(\tau)$, was fitted to the average

12   autocorrelation function:

13
$$G_D(\tau) = \frac{1}{V_{eff}\chi^2 C}\left(\frac{1}{1 + 8D\tau/w_{xy}^2}\right)\left(\frac{1}{1 + 8D\tau/w_z^2}\right)^{\frac{1}{2}} + G_{\infty}$$

14   where V_{eff} is the effective volume of PSF, C the concentration of fluorophores (which is 97 nM), χ^2 the

15   background noise correction factor (Hess and Webb, 2002), D the diffusion coefficient of Alexa Fluor 488,

16   which was previously estimated to be 435 µm^2/s (Petrasek and Schwille, 2008), and w_{xy} and w_z are the radial

17   and axial beam waists, respectively (Figure 1−figure supplement 1D). w_z can be written in terms of V_{eff} and

18   w_{xy}:

19
$$w_z = \left(\frac{2}{\pi}\right)^{3/2}\frac{V_{eff}}{w_{xy}^2}$$

Fitting the FCS model to the Alexa 488 FCS data estimated V_{eff} and w_{xy} to be 0.364 ± 0.004 µm^3 and 278

± 4 nm, respectively (Figure 1−figure supplement 1D).

We next performed an FCS measurement on a synthesized TC peptide labeled with FlAsH. 50

µM synthesized TC peptide (Ac-AEEEACCPGCC-NH$_2$, Genscript), 100 µM FlAsH-EDT$_2$, and 10 mM

2-mercaptoethanesulfonate (Sigma Aldrich) were incubated for an hour to associate TC peptide with

FlAsH, then diluted in the imaging buffer by 500 times, and prepared on a coverslip for FCS

measurement. The laser intensity was set to 3 mW. 6 autocorrelation functions, each of which had been

collected for 300 seconds, were averaged, and the following FCS model was fitted to the average

autocorrelation function to determine the number of fluorophores N in a focal volume V_{eff}:

6
$$G_D(\tau) = \frac{1}{N\chi^2}\left(\frac{1}{1 + \tau/\tau_D}\right)\left(\frac{1}{1 + (w_{xy}^2/w_z^2)(\tau/\tau_D)}\right)^{\frac{1}{2}} + G_\infty$$

7   while w_{xy} and w_z were fixed to the values determined from the FCS measurement on Alexa Fluor 488

8   (Figure 1–figure supplement 1E). The photon count collected during the FCS measurement was corrected

9   for background noise, and then divided by N to yield the molecular brightness of TC-FlAsH, 233.4 ± 9.3

10   s^{-1}. Using the estimated molecular brightness and the effective volume of the PSF, we calculated the

11   average concentration of the polymerized β-tubulin-TC-FlAsH to be

12
$$\frac{9.6 \times 10^4 \ s^{-1}}{(233.4 \ s^{-1})(0.364 \ \mu m^3)} = 1.13 \times 10^{21} m^{-3} = 1.88 \pm 0.13 \ \mu M$$

Calculating labeling ratio: A previous study (Dumontet et al., 1996) estimated the percentage of

polymerized β-tubulin in a mitotic human tissue culture cell to be $36\% \pm 7\%$. Combining this information

with our estimate of an average concentration of polymerized β-tubulin-TC-FlAsH of 1.88 ± 0.13 μM

leads to a total concentration of β-tubulin-TC-FlAsH of 1.88 μM $\times 100/36 \approx 5.22 \pm 1.08$ μM. Since the

total concentration of tubulin dimer in a tissue culture cell is ~20 μM (Hiller and Weber, 1978), we

estimated the fraction of labeled β-tubulin to be 5.22 μM/20 μM $\approx 26.1\% \pm 5.4\%$. This estimate makes

use of data obtained in different cell types, which may introduce inaccuracies. Systematic errors in the

total concentration of tubulin or the concentration of the β-tubulin-TC-FlAsH will produce proportional

errors in the estimates of the fraction of labeled β-tubulin, resulting in proportional systematic errors in

the conversion of NDC80 FRET fraction to NDC80 binding fraction.

<u>Förster radius estimation</u>

To measure the Förster radius R_0 of FRET between mTurquiose2 and TC-FlAsH, we created a construct

containing mTurquoise2 tethered to TC (mTurquoise2-TC), expressed it in U2OS cells, and acquired

fluorescence decays of mTurquoise2 using FLIM, which were well-described by a single-exponential

fluorescence decay with a lifetime of 3.75 ns ± 0.03 ns (SD) in the absence of FlAsH labeling (Figure 1−

figure supplement 4A). When FlAsH is added to these cells, FLIM measurements revealed the presence

of additional shorter-lifetime species, corresponding to mTurquoise2 molecules engaged in FRET with

TC-FlAsH (Figure 1−figure supplement 4A). Then we performed Monte Carlo protein simulations (which

are described below) to model the conformational ensemble of the flexible tether between mTurquoise2

and TC-FlAsH and obtain the distribution $p(r)$ of the distance r between mTurquoise2 and FlAsH

(Figure 1−figure supplement 4B). The fluorescence lifetime τ of donors engaged in FRET is related to the

donor-acceptor distance r by:

14
$$\tau(r; R_0) = \frac{\tau_D}{1 + \left(\frac{R_0}{r}\right)^6}$$

15 where τ_D is the fluorescence lifetime of the donor in non-FRET state (3.75 ns ± 0.03 ns), and R_0 is the

16 Förster radius. Therefore, the fluorescence decay $y(t)$ of mTurquoise2-TC-FlAsH can be modeled as:

17
$$y(t) = A_D \exp\left(-\frac{t}{\tau_D}\right) + A_{FRET} \int p(r) \exp\left(-\frac{t}{\tau(r; R_0)}\right) dr$$

where A_D is the population in the non-FRET state and A_{FRET} is that in the FRET state, both of which are

free parameters of the model along with R_0. This model (after convolved with the IRF) was fit to the

measured fluorescence decay curve of mTurquoise2-TC-FlAsH, allowing us to estimate the Förster radius

R_0 to be 5.90 nm ± 0.10 nm (Figure 1−figure supplement 4C).

<u>Characterization and calibration of NDC80-kMT FLIM-FRET measurements by Monte Carlo simulations</u>

Characterization of NDC80-kMT FRET vs NDC80-MT distance relationship: To characterize the FRET

between mTurquoise2-NDC80 and FlAsH-labeled microtubule when NDC80 is not bound to the

microtubule, we performed large-scale, atomistic Monte Carlo protein simulations to model the

conformational ensemble of the tether between mTurquoise2 and NDC80 and the disordered C-terminal

tails of twelve β-tubulins near the NDC80 complex (which is described below) (Figure 1−figure

supplement 5A). 4000 sets of positions of mTurquoise2 and TC were generated for each case where the

NDC80 bound to an inter- or intra-tubulin dimer interface was translated away from the microtubule by a

certain distance (0~15 nm, 0.5 nm increment) in a direction perpendicular to the microtubule surface. For

each randomly sampled set of distances between mTurquoise2 and TC-FlAsH, $\vec{r} = \{r_i\}$, the fluorescence

lifetime was calculated by:

11
$$\tau(\vec{r}) = \frac{\tau_D}{1 + \sum_{i=1}^{12} I_i \left(\frac{R_0}{r_i}\right)^6}$$

where $I_i \sim \text{Bernoulli}(f_{label})$ indicates whether or not the *i*-th TC motif is labeled with FlAsH-EDT$_2$; R_0 is

the Förster radius between mTurquoise2 and TC-FlAsH; and τ_D=3.75 ns is the non-FRET lifetime of

mTurquoise2 (Figure 1−figure supplement 5A). The estimated labeling fraction 26.1% and the Förster

radius 5.90 nm were used for f_{label} and R_0, respectively. For each NDC80-kMT distance, 2 million

fluorescence lifetimes were sampled, based on which we simulated 30 fluorescence decay curves of

mTurquoise2 by:

18
$$\text{Poisson}\left(A \int p(\vec{r}) \exp\left(-\frac{t}{\tau(\vec{r})}\right) d\vec{r}\right)$$

where the amplitude *A* of the fluorescence decay was set to 5000 (Figure 1−figure supplement 5A).

Single- and double-exponential decay models were fit to the simulated fluorescence decays (by maximum

likelihood method) (Figure 1−figure supplement 5B, bottom). The Bayesian information criterion (BIC)

was used as a criterion for model selection between the single- and double-exponential decay models. The

difference in BIC between single- and double-exponential models, $\Delta\text{BIC} = \text{BIC}_{1\text{expo}} - \text{BIC}_{2\text{expo}}$, was

plotted against the NDC80-kMT distance (Figure 1—figure supplement 5B, top right). ΔBIC is negative

when NDC80-kMT distance is larger than 8 nm, indicating that single-exponential model performs better

than double-exponential model in terms of the goodness of fit and the complexity of model (Figure 1—

figure supplement 5B).

NDC80-kMT FRET fraction calibration: To obtain the relationship between NDC80 FRET fraction and

NDC80 binding fraction, we performed large-scale Monte Carlo simulations to obtain 4000 sets of

distances between mTurquoise2 and TC-FlAsH, $\vec{r} = \{r_i\}$, for each case where mTurquoise2-NDC80 is

bound to the TC-tagged microtubule at the inter- or intra-dimer interface. Then we sampled 0.5 million

fluorescence lifetimes τ as described above, and simulated a fluorescence decay curve for the situation

where a fraction f_b of mTurquoise2-NDC80 are bound to microtubules and have lifetime $\tau(\vec{r})$, while the

other $1 - f_b$ are not bound and have lifetime τ_D (Figure 1—figure supplement 5C, green dots):

$$\text{Poisson}\left(A\left[(1 - f_b)\exp\left(-\frac{t}{\tau_D}\right) + f_b\int p(\vec{r})\exp\left(-\frac{t}{\tau(\vec{r})}\right)d\vec{r}\right]\right)$$

where the amplitude *A* of the fluorescence decay curve was set to 10,000 (Figure 1—figure supplement 5C,

left). The simulated fluorescence decay curve was then fit by two-exponential fluorescence decay model

(Figure 1—figure supplement 5C, red lines):

$$A'\left[(1 - f_{FRET})\exp\left(-\frac{t}{\tau_D}\right) + f_{FRET}\exp\left(-\frac{t}{\tau_{FRET}}\right)\right]$$

to acquire the FRET fraction, f_{FRET}. The data of NDC80 FRET fraction, f_{FRET}, vs NDC80 binding

fraction, f_b, was fit using a linear model $f_{FRET} = af_b$ (Figure 1—figure supplement 5C, right). To

determine the uncertainty in the slope *a* (gray area in Figure 1—figure supplement 5C, right), we repeated

the process above with the mean ± error values of R_0 and beta-tubulin labeling fraction f_{label}. As a result,

we obtained *a* = 0.42 ± 0.08, and used this calibration to convert NDC80 FRET fraction to NDC80

binding fraction.

1 <u>Monte Carlo protein simulations</u>

2 Atomistic simulations were performed by the CAMPARI (v2) package (Vitalis and Pappu, 2009b),

3 employing the ABSINTH implicit solvation model and forcefield paradigm (Vitalis and Pappu, 2009a) at

4 the intrinsic solvation (IS) limit (Das and Pappu, 2013) (unless stated otherwise), where the energy

5 function is simply a combination of Lennard-Jones energy and ABSINTH solvation energy.

7 *mTurquoise2-TC construct simulation*: For the Förster radius estimation, we ran 50 independent

8 simulations on the mTurquoise2-TC construct in spherical soft-wall boundary conditions with radius 100

9 Å. An input structure was used only for the folded mTurquoise2 domain (adapted from PDB 4B5Y), and

10 we employed CAMPARI to generate the tether and TC domains (GMDELYKYSDLFLNCCPGCCMEP)

11 from scratch. To prevent unphysical unfolding and/or conformational change of mTurquoise2, we

12 imposed constraints on internal degrees of freedom of residues in the folded region. Each simulation

13 consisted of 2×10^6 MC steps with sampling frequency of $(5,000\ \text{steps})^{-1}$, and the simulation temperature

14 was set to 400 K in order to scan a large structural ensemble. Note that the system is quickly relaxed in

15 the intrinsic solvation (IS) limit, and hence it does not require a long simulation time to reach

16 equilibration. The average coordinate of the alpha carbons of 4 residues before and after Trp66 was used

17 as the location of mTurquoise2 chromophore, and the average coordinate of alpha carbons of the four

18 cysteine residues in the TC motif was used as the location of TC-FlAsH.

20 *mTurquoise2-NDC80 and TC-labeled microtubule simulation*: We constructed a system consisting of the

21 NDC80 complex, 12 tubulin dimers as described in Figure 1−figure supplement 5A. The initial structures

22 of the system were constructed by combining the structures of microtubule (PDB 3JAS) (Zhang et al.,

23 2015) and one of bonsai-NDC80s attached to a tubulin dimer at the inter- and intra-dimer interfaces (PDB

24 3IZ0) (Alushin et al., 2010), and then incorporating the structures of disordered regions (tether, GMDEL

25 YKYSD LMET, and C-terminal tail+TC, SEYQQ YQDAT AEEEE DFGEE AEEEA CCPGC C)

generated by the loop modeling module of Rosetta 3.8 (Mandell et al., 2009, Leaver-Fay et al., 2011).

Clashes in the initial structure were removed by the Rosetta relaxation module (Nivón et al., 2013). For

folded regions (where we have structure information from PDB), we imposed constraints on internal

degrees of freedom as before. To prevent dissociation of microtubule into individual tubulins, we also

imposed a harmonic restraint potential on atoms at the interface of two different chains. If two atoms from

different chains are closer than 20 Å, the pair contributes an additional potential

$$E_{\mathrm{drest}}(i,j) = k\left(r_{ij} - r_{ij}^{0}\right)^{2},$$

where i and j are atomic indices, r_{ij} is the distance between two atoms i and j, r_{ij}^{0} is the initial distance

between i and j, and k is a force constant (set to 3.0 kcal/mol/Å^2). We employed spherical soft-wall

boundary conditions with radius 200 Å, and the simulation temperature of 400 K. We ran a relaxation

simulation for 2×10^6 MC steps, and a production simulation for another 2×10^6 MC steps with sampling

frequency of (500 steps)$^{-1}$. For the FRET efficiency vs NDC80-MT data (Figure 1–figure supplement

5B), we used the final structure of the relaxation simulation and translated the NDC80 complex along the

axis orthogonal to the microtubule surface by several distance values: from 0 to 15 nm by increment of

0.5 nm. For each system, a simulation of 2×10^6 MC steps was conducted to generate data with sampling

frequency of (500 steps)$^{-1}$.

Protein structure illustration

Protein structure illustrations were generated by The PyMOL Molecular Graphics System, Version 2.0

Schrödinger, LLC.

Quantification and Statistical Analysis

The statistical test used, sample size (number of cells and kinetochores), dispersion and precision

measures can be found in figure legends, Results, or below. All curve fittings, except FLIM data analysis

(which is separately explained above), were performed by Levenberg-Marquardt algorithm with residuals

weighted by the inverse of y-errors, and the corresponding 95% confidence intervals were calculated by

predint function in MATLAB. To assess the significance of correlation, we determined p-value from $1 -$

α, where α is the smallest confidence level that makes zero contained in the confidence interval of the

slope of the linear fit.

ACKNOWLEDGMENTS

We thank A. Murray and N. Kleckner for comments on the manuscript; J. DeLuca, M. Lampson, and I. Cheeseman for reagents; F. Rago for help with retroviral transfection; D. Kim for help with nucleofection; Needleman lab members and J. Oh for proof reading, comments and discussion.

COMPETING INTERESTS

The authors declare no competing financial interests.

REFERENCES

AKIYOSHI, B., SARANGAPANI, K. K., POWERS, A. F., NELSON, C. R., REICHOW, S. L., ARELLANO-SANTOYO, H., GONEN, T., RANISH, J. A., ASBURY, C. L. & BIGGINS, S. 2010. Tension directly stabilizes reconstituted kinetochore-microtubule attachments. *Nature,* 468**,** 576-9.

ALUSHIN, G. M., RAMEY, V. H., PASQUALATO, S., BALL, D. A., GRIGORIEFF, N., MUSACCHIO, A. & NOGALES, E. 2010. The Ndc80 kinetochore complex forms oligomeric arrays along microtubules. *Nature,* 467**,** 805-10.

ANDRESEN, M., SCHMITZ-SALUE, R. & JAKOBS, S. 2004. Short tetracysteine tags to beta-tubulin demonstrate the significance of small labels for live cell imaging. *Mol Biol Cell,* 15**,** 5616-22.

ARMOND, J. W., VLADIMIROU, E., ERENT, M., MCAINSH, A. D. & BURROUGHS, N. J. 2015. Probing microtubule polymerisation state at single kinetochores during metaphase chromosome motion. *J Cell Sci,* 128**,** 1991-2001.

BEREZIN, M. Y. & ACHILEFU, S. 2010. Fluorescence lifetime measurements and biological imaging. *Chem Rev,* 110**,** 2641-84.

BIGGINS, S., SEVERIN, F. F., BHALLA, N., SASSOON, I., HYMAN, A. A. & MURRAY, A. W. 1999. The conserved protein kinase Ipl1 regulates microtubule binding to kinetochores in budding yeast. *Genes Dev,* 13**,** 532-44.

BISHOP, J. D. & SCHUMACHER, J. M. 2002. Phosphorylation of the carboxyl terminus of inner centromere protein (INCENP) by the Aurora B Kinase stimulates Aurora B kinase activity. *J Biol Chem,* 277**,** 27577-80.

CAMPBELL, C. S. & DESAI, A. 2013. Tension sensing by Aurora B kinase is independent of survivin-based centromere localization. *Nature,* 497**,** 118-21.

CARMENA, M., WHEELOCK, M., FUNABIKI, H. & EARNSHAW, W. C. 2012. The chromosomal passenger complex (CPC): from easy rider to the godfather of mitosis. *Nat Rev Mol Cell Biol,* 13**,** 789-803.

CHEESEMAN, I. M., CHAPPIE, J. S., WILSON-KUBALEK, E. M. & DESAI, A. 2006. The conserved KMN network constitutes the core microtubule-binding site of the kinetochore. *Cell,* 127**,** 983-97.

CHEESEMAN, I. M. & DESAI, A. 2008. Molecular architecture of the kinetochore-microtubule interface. *Nat Rev Mol Cell Biol,* 9**,** 33-46.

CIFERRI, C., PASQUALATO, S., SCREPANTI, E., VARETTI, G., SANTAGUIDA, S., DOS REIS, G., MAIOLICA, A., POLKA, J., DE LUCA, J. G., DE WULF, P., SALEK, M., RAPPSILBER, J., MOORES, C. A., SALMON, E. D. & MUSACCHIO, A. 2008. Implications for kinetochore-microtubule attachment from the structure of an engineered Ndc80 complex. *Cell,* 133**,** 427-39.

CIMINI, D., WAN, X., HIREL, C. B. & SALMON, E. D. 2006. Aurora kinase promotes turnover of kinetochore microtubules to reduce chromosome segregation errors. *Curr Biol,* 16**,** 1711-8.

DAS, R. K. & PAPPU, R. V. 2013. Conformations of intrinsically disordered proteins are influenced by linear sequence distributions of oppositely charged residues. *Proceedings of the National Academy of Sciences,* 110**,** 13392-13397.

DELUCA, J. G., GALL, W. E., CIFERRI, C., CIMINI, D., MUSACCHIO, A. & SALMON, E. D. 2006. Kinetochore microtubule dynamics and attachment stability are regulated by Hec1. *Cell,* 127**,** 969-82.

DELUCA, K. F., LENS, S. M. & DELUCA, J. G. 2011. Temporal changes in Hec1 phosphorylation control kinetochore-microtubule attachment stability during mitosis. *J Cell Sci,* 124**,** 622-34.

DELUCA, K. F., MEPPELINK, A., BROAD, A. J., MICK, J. E., PEERSEN, O. B., PEKTAS, S., LENS, S. M. A. & DELUCA, J. G. 2018. Aurora A kinase phosphorylates Hec1 to regulate metaphase kinetochore-microtubule dynamics. *J Cell Biol,* 217**,** 163-177.

DUMONT, S., SALMON, E. D. & MITCHISON, T. J. 2012. Deformations within moving kinetochores reveal different sites of active and passive force generation. *Science,* 337**,** 355-8.

DUMONTET, C., DURAN, G. E., STEGER, K. A., MURPHY, G. L., SUSSMAN, H. H. & SIKIC, B. I. 1996. Differential expression of tubulin isotypes during the cell cycle. *Cell Motil Cytoskeleton,* 35**,** 49-58.

EDELSTEIN, A. D., TSUCHIDA, M. A., AMODAJ, N., PINKARD, H., VALE, R. D. & STUURMAN, N. 2014. Advanced methods of microscope control using muManager software. *J Biol Methods,* 1.

FULLER, B. G., LAMPSON, M. A., FOLEY, E. A., ROSASCO-NITCHER, S., LE, K. V., TOBELMANN, P., BRAUTIGAN, D. L., STUKENBERG, P. T. & KAPOOR, T. M. 2008. Midzone activation of aurora B in anaphase produces an intracellular phosphorylation gradient. *Nature,* 453**,** 1132-6.

GODEK, K. M., KABECHE, L. & COMPTON, D. A. 2015. Regulation of kinetochore-microtubule attachments through homeostatic control during mitosis. *Nat Rev Mol Cell Biol,* 16**,** 57-64.

GOEDHART, J., VON STETTEN, D., NOIRCLERC-SAVOYE, M., LELIMOUSIN, M., JOOSEN, L., HINK, M. A., VAN WEEREN, L., GADELLA, T. W., JR. & ROYANT, A. 2012. Structure-guided evolution of cyan fluorescent proteins towards a quantum yield of 93%. *Nat Commun,* 3**,** 751.

GUIMARAES, G. J., DONG, Y., MCEWEN, B. F. & DELUCA, J. G. 2008. Kinetochore-microtubule attachment relies on the disordered N-terminal tail domain of Hec1. *Curr Biol,* 18**,** 1778-84.

HAASE, J., BONNER, M. K., HALAS, H. & KELLY, A. E. 2017. Distinct Roles of the Chromosomal Passenger Complex in the Detection of and Response to Errors in Kinetochore-Microtubule Attachment. *Dev Cell,* 42**,** 640-654 e5.

HAUF, S., COLE, R. W., LATERRA, S., ZIMMER, C., SCHNAPP, G., WALTER, R., HECKEL, A., VAN MEEL, J., RIEDER, C. L. & PETERS, J. M. 2003. The small molecule Hesperadin reveals a role for Aurora B in correcting kinetochore-microtubule attachment and in maintaining the spindle assembly checkpoint. *J Cell Biol,* 161**,** 281-94.

HESS, S. T. & WEBB, W. W. 2002. Focal volume optics and experimental artifacts in confocal fluorescence correlation spectroscopy. *Biophys J,* 83**,** 2300-17.

HILLER, G. & WEBER, K. 1978. Radioimmunoassay for tubulin: a quantitative comparison of the tubulin content of different established tissue culture cells and tissues. *Cell,* 14**,** 795-804.

HOFFMANN, C., GAIETTA, G., BUNEMANN, M., ADAMS, S. R., OBERDORFF-MAASS, S., BEHR, B., VILARDAGA, J. P., TSIEN, R. Y., ELLISMAN, M. H. & LOHSE, M. J. 2005. A FlAsH-based FRET approach to determine G protein-coupled receptor activation in living cells. *Nat Methods,* 2**,** 171-6.

HOFFMANN, C., GAIETTA, G., ZURN, A., ADAMS, S. R., TERRILLON, S., ELLISMAN, M. H., TSIEN, R. Y. & LOHSE, M. J. 2010. Fluorescent labeling of tetracysteine-tagged proteins in intact cells. *Nat Protoc,* 5**,** 1666-77.

HUANG, H. B., HORIUCHI, A., GOLDBERG, J., GREENGARD, P. & NAIRN, A. C. 1997. Site-directed mutagenesis of amino acid residues of protein phosphatase 1 involved in catalysis and inhibitor binding. *Proc Natl Acad Sci U S A,* 94**,** 3530-5.

KABECHE, L. & COMPTON, D. A. 2013. Cyclin A regulates kinetochore microtubules to promote faithful chromosome segregation. *Nature,* 502**,** 110-3.

KAYE, B., FOSTER, P. J., YOO, T. Y. & NEEDLEMAN, D. J. 2017. Developing and Testing a Bayesian Analysis of Fluorescence Lifetime Measurements. *PLoS One,* 12**,** e0169337.

KELLY, A. E., SAMPATH, S. C., MANIAR, T. A., WOO, E. M., CHAIT, B. T. & FUNABIKI, H. 2007. Chromosomal enrichment and activation of the aurora B pathway are coupled to spatially regulate spindle assembly. *Dev Cell,* 12**,** 31-43.

KRENN, V. & MUSACCHIO, A. 2015. The Aurora B Kinase in Chromosome Bi-Orientation and Spindle Checkpoint Signaling. *Front Oncol,* 5**,** 225.

LAMPSON, M. A. & CHEESEMAN, I. M. 2011. Sensing centromere tension: Aurora B and the regulation of kinetochore function. *Trends Cell Biol,* 21**,** 133-40.

LEANDRO-GARCIA, L. J., LESKELA, S., LANDA, I., MONTERO-CONDE, C., LOPEZ-JIMENEZ, E., LETON, R., CASCON, A., ROBLEDO, M. & RODRIGUEZ-ANTONA, C. 2010. Tumoral and tissue-specific expression of the major human beta-tubulin isotypes. *Cytoskeleton (Hoboken),* 67**,** 214-23.

LEAVER-FAY, A., TYKA, M., LEWIS, S. M., LANGE, O. F., THOMPSON, J., JACAK, R., KAUFMAN, K. W., RENFREW, P. D., SMITH, C. A., SHEFFLER, W., DAVIS, I. W., COOPER, S., TREUILLE, A., MANDELL, D. J., RICHTER, F., BAN, Y.-E. A., FLEISHMAN, S. J., CORN, J. E., KIM, D. E., LYSKOV, S., BERRONDO, M., MENTZER, S., POPOVIĆ, Z., HAVRANEK, J. J., KARANICOLAS, J., DAS, R., MEILER, J., KORTEMME, T., GRAY, J. J., KUHLMAN, B., BAKER, D. & BRADLEY, P. 2011. Chapter nineteen - Rosetta3: An Object-Oriented Software Suite for the Simulation and Design of Macromolecules. *In:* JOHNSON, M. L. & BRAND, L. (eds.) *Methods in Enzymology.* Academic Press.

LIU, D., VADER, G., VROMANS, M. J., LAMPSON, M. A. & LENS, S. M. 2009. Sensing chromosome bi-orientation by spatial separation of aurora B kinase from kinetochore substrates. *Science,* 323**,** 1350-3.

MAGIDSON, V., O'CONNELL, C. B., LONCAREK, J., PAUL, R., MOGILNER, A. & KHODJAKOV, A. 2011. The spatial arrangement of chromosomes during prometaphase facilitates spindle assembly. *Cell,* 146**,** 555-67.

MAIATO, H., DELUCA, J., SALMON, E. D. & EARNSHAW, W. C. 2004. The dynamic kinetochore-microtubule interface. *J Cell Sci,* 117**,** 5461-77.

MANDELL, D. J., COUTSIAS, E. A. & KORTEMME, T. 2009. Sub-angstrom accuracy in protein loop reconstruction by robotics-inspired conformational sampling. *Nat Meth,* 6**,** 551-552.

MCEWEN, B. F., HEAGLE, A. B., CASSELS, G. O., BUTTLE, K. F. & RIEDER, C. L. 1997. Kinetochore fiber maturation in PtK1 cells and its implications for the mechanisms of chromosome congression and anaphase onset. *J Cell Biol,* 137**,** 1567-80.

NICKLAS, R. B. & KOCH, C. A. 1969. Chromosome micromanipulation. 3. Spindle fiber tension and the reorientation of mal-oriented chromosomes. *J Cell Biol,* 43**,** 40-50.

NICKLAS, R. B. & WARD, S. C. 1994. Elements of error correction in mitosis: microtubule capture, release, and tension. *J Cell Biol,* 126**,** 1241-53.

NIVÓN, L. G., MORETTI, R. & BAKER, D. 2013. A Pareto-Optimal Refinement Method for Protein Design Scaffolds. *PLOS ONE,* 8**,** e59004.

PELLETIER, V., GAL, N., FOURNIER, P. & KILFOIL, M. L. 2009. Microrheology of microtubule solutions and actin-microtubule composite networks. *Phys Rev Lett,* 102**,** 188303.

PETRASEK, Z. & SCHWILLE, P. 2008. Precise measurement of diffusion coefficients using scanning fluorescence correlation spectroscopy. *Biophys J,* 94**,** 1437-48.

POSCH, M., KHOUDOLI, G. A., SWIFT, S., KING, E. M., DELUCA, J. G. & SWEDLOW, J. R. 2010. Sds22 regulates aurora B activity and microtubule-kinetochore interactions at mitosis. *J Cell Biol,* 191**,** 61-74.

RAN, F. A., HSU, P. D., WRIGHT, J., AGARWALA, V., SCOTT, D. A. & ZHANG, F. 2013. Genome engineering using the CRISPR-Cas9 system. *Nat Protoc,* 8**,** 2281-308.

RIEDER, C. L. 1982. The formation, structure, and composition of the mammalian kinetochore and kinetochore fiber. *Int Rev Cytol,* 79**,** 1-58.

ROSASCO-NITCHER, S. E., LAN, W., KHORASANIZADEH, S. & STUKENBERG, P. T. 2008. Centromeric Aurora-B activation requires TD-60, microtubules, and substrate priming phosphorylation. *Science,* 319**,** 469-72.

ROWLEY, M. I., COOLEN, A. C., VOJNOVIC, B. & BARBER, P. R. 2016. Robust Bayesian Fluorescence Lifetime Estimation, Decay Model Selection and Instrument Response Determination for Low-Intensity FLIM Imaging. *PLoS One,* 11**,** e0158404.

SALIMIAN, K. J., BALLISTER, E. R., SMOAK, E. M., WOOD, S., PANCHENKO, T., LAMPSON, M. A. & BLACK, B. E. 2011. Feedback control in sensing chromosome biorientation by the Aurora B kinase. *Curr Biol,* 21**,** 1158-65.

SANTAGUIDA, S. & AMON, A. 2015. Short- and long-term effects of chromosome mis-segregation and aneuploidy. *Nat Rev Mol Cell Biol,* 16**,** 473-85.

SESSA, F., MAPELLI, M., CIFERRI, C., TARRICONE, C., ARECES, L. B., SCHNEIDER, T. R., STUKENBERG, P. T. & MUSACCHIO, A. 2005. Mechanism of Aurora B activation by INCENP and inhibition by hesperadin. *Mol Cell,* 18**,** 379-91.

SHANER, N. C., LAMBERT, G. G., CHAMMAS, A., NI, Y., CRANFILL, P. J., BAIRD, M. A., SELL, B. R., ALLEN, J. R., DAY, R. N., ISRAELSSON, M., DAVIDSON, M. W. & WANG, J. 2013. A bright monomeric green fluorescent protein derived from Branchiostoma lanceolatum. *Nat Methods,* 10**,** 407-9.

SKOUFIAS, D. A., DEBONIS, S., SAOUDI, Y., LEBEAU, L., CREVEL, I., CROSS, R., WADE, R. H., HACKNEY, D. & KOZIELSKI, F. 2006. S-trityl-L-cysteine is a reversible, tight binding inhibitor of the human kinesin Eg5 that specifically blocks mitotic progression. *J Biol Chem,* 281**,** 17559-69.

SUZUKI, A., BADGER, B. L. & SALMON, E. D. 2015. A quantitative description of Ndc80 complex linkage to human kinetochores. *Nat Commun,* 6**,** 8161.

TANAKA, T. U., RACHIDI, N., JANKE, C., PEREIRA, G., GALOVA, M., SCHIEBEL, E., STARK, M. J. & NASMYTH, K. 2002. Evidence that the Ipl1-Sli15 (Aurora kinase-INCENP) complex promotes chromosome bi-orientation by altering kinetochore-spindle pole connections. *Cell,* 108**,** 317-29.

TAUCHMAN, E. C., BOEHM, F. J. & DELUCA, J. G. 2015. Stable kinetochore-microtubule attachment is sufficient to silence the spindle assembly checkpoint in human cells. *Nat Commun,* 6**,** 10036.

THURBER, G. M., REINER, T., YANG, K. S., KOHLER, R. H. & WEISSLEDER, R. 2014. Effect of small-molecule modification on single-cell pharmacokinetics of PARP inhibitors. *Mol Cancer Ther,* 13**,** 986-95.

TIRNAUER, J. S., CANMAN, J. C., SALMON, E. D. & MITCHISON, T. J. 2002. EB1 targets to kinetochores with attached, polymerizing microtubules. *Mol Biol Cell,* 13**,** 4308-16.

VITALIS, A. & PAPPU, R. V. 2009a. ABSINTH: A new continuum solvation model for simulations of polypeptides in aqueous solutions. *Journal of Computational Chemistry,* 30**,** 673-699.

VITALIS, A. & PAPPU, R. V. 2009b. Chapter 3 Methods for Monte Carlo Simulations of Biomacromolecules. *In:* RALPH, A. W. (ed.) *Annual Reports in Computational Chemistry.* Elsevier.

WANG, F., ULYANOVA, N. P., DAUM, J. R., PATNAIK, D., KATENEVA, A. V., GORBSKY, G. J. & HIGGINS, J. M. 2012. Haspin inhibitors reveal centromeric functions of Aurora B in chromosome segregation. *J Cell Biol,* 199**,** 251-68.

XU, Z., VAGNARELLI, P., OGAWA, H., SAMEJIMA, K. & EARNSHAW, W. C. 2010. Gradient of increasing Aurora B kinase activity is required for cells to execute mitosis. *J Biol Chem,* 285**,** 40163-70.

YOO, T. Y., CHOI, J., YU, C., PAPPU, R. V. & NEEDLEMAN, D. J. Data from: Measuring NDC80 binding reveals the molecular basis of tension-dependent kinetochore-microtubule attachments. Dryad Data Repository.

YOO, T. Y. & NEEDLEMAN, D. J. 2016. Studying Kinetochores In Vivo Using FLIM-FRET. *Methods Mol Biol,* 1413**,** 169-86.

ZAYTSEV, A. V., MICK, J. E., MASLENNIKOV, E., NIKASHIN, B., DELUCA, J. G. & GRISHCHUK, E. L. 2015. Multisite phosphorylation of the NDC80 complex gradually tunes its microtubule-binding affinity. *Mol Biol Cell,* 26**,** 1829-44.

ZAYTSEV, A. V., SEGURA-PENA, D., GODZI, M., CALDERON, A., BALLISTER, E. R., STAMATOV, R., MAYO, A. M., PETERSON, L., BLACK, B. E., ATAULLAKHANOV, F. I., LAMPSON, M. A. & GRISHCHUK, E. L. 2016. Bistability of a coupled Aurora B kinase-phosphatase system in cell division. *Elife,* 5**,** e10644.

ZAYTSEV, A. V., SUNDIN, L. J., DELUCA, K. F., GRISHCHUK, E. L. & DELUCA, J. G. 2014. Accurate phosphoregulation of kinetochore-microtubule affinity requires unconstrained molecular interactions. *J Cell Biol,* 206**,** 45-59.

ZHAI, Y., KRONEBUSCH, P. J. & BORISY, G. G. 1995. Kinetochore microtubule dynamics and the metaphase-anaphase transition. *J Cell Biol,* 131**,** 721-34.

ZHANG, R., ALUSHIN, G. M., BROWN, A. & NOGALES, E. 2015. Mechanistic Origin of Microtubule Dynamic Instability and Its Modulation by EB Proteins. *Cell,* 162**,** 849-59.

SUPPLEMENTAL FIGURES

Figure 1–figure supplement 1. β-tubulin labeling fraction measurement
(A) Two-photon fluorescent microscopy images of 9 mitotic cells with β-tubulin-TC-FlAsH. 5 μm scale bar.
(B) Example 3D segmentation using active contour algorithm.
(C) (top left) The number of voxels in segmented 3D cell images, (top right) the total number of photons collected from entire cells, (bottom left) the total number of photons divided by the number of voxels and the measurement time, and (bottom right) the number of photons collected per second at a voxel in the cytoplasmic region. Gray lines are the average over the 9 different cells.
(D) and (E) FCS measurements on Alexa Fluor 488 and TC-FlAsH in solution, respectively. Black circles are averages over 5 or 6 autocorrelation functions, and red lines are a single-component FCS model fit to the averages. Corresponding weighted residuals (the difference between data and model, divided by the standard deviation of data) are plotted below to show any systematic deviations of the fitted models from the data.

Figure 1–figure supplement 2. Kinetochore FLIM-FRET measurement

(A) Illustration of fluorescence decay acquisition in a TCSPC (time-correlated single photon counting) FLIM system. A Ti:Sapphire pulsed laser is used for excitation and a photomultiplier tube (PMT) for detection. The photon arrival time, the difference in timing between the emitted photon and excitation laser pulse, is measured by TCSPC, which is accumulated to a fluorescence decay curve over many laser repetition periods.

(B) The fluorescence lifetime is the average time the fluorophore stays in the excited state. (left) When the donor fluorophore (D) is not engaged in FRET with an acceptor (A) fluorophore, the donor fluorophore has a single-exponential fluorescence decay with fluorescence lifetime $\tau_D = 1/(k_r+k_{nr})$, where k_r and k_{nr} are the radiative and non-radiative decay rates, respectively. (middle) FRET provides an additional relaxation pathway to the excited donor, reducing the fluorescence lifetime of the donor to $\tau_{FRET} = 1/(k_r+k_{nr}+k_{FRET})$, where k_{FRET} is the FRET rate. (right) The fluorescence decay of a mixture of donors engaged in FRET and not engaged in FRET is a sum of two exponentials with two different lifetimes, τ_D and τ_{FRET}, which corresponds to the non-FRET and FRET populations, respectively. The relative amplitude of the short-lifetime exponential decay provides the fraction of the FRET population, and the lifetime ratio provides the intrinsic FRET efficiency.

(C) TCSPC FLIM provides fluorescence decay curve at each pixel, and the total photon counts in each pixel provides a two-photon fluorescence intensity image. To quantify the FRET fraction at each kinetochore, kinetochores were identified based on the intensity image, then the fluorescence decay curves in the pixels within each kinetochore were summed. Then we performed a Bayesian analysis to

obtain the posterior distribution of the FRET fraction at each kinetochore. The posterior distributions of the kinetochores in a group of kinetochores were combined by multiplication to compute the mean and SEM of the FRET fraction for the group.

Figure 1–figure supplement 3. Negative control data for NDC80-kMT FLIM-FRET measurements
(A) to (D) Schematic descriptions, example cell images, and example mTurquoise2 fluorescence decay curves from three different FRET-negative control experiments and a nocodazole treatment experiment. mTurquoise2 fluorescence decay curves (blue circles) are plotted with best-fit single- (black dotted line) or two-exponential decay model (black solid line), and the associated weighted residuals are plotted below (blue curve). 3 μm scale bar.
(A) Negative control 1. Nuf2 N-terminally labeled with mTurquoise2, and no FlAsH labeling.
(B) Negative control 2. Nuf2 C-terminally labeled with mTurquoise2 (far from kMT), and no FlAsH labeling.
(C) Negative control 3. Nuf2 C-terminally labeled with mTurquoise2 (far from kMT), and β-tubulin C-terminally labeled with FlAsH.
(D) Nocodazole treatment experiment. Nuf2 N-terminally labeled with mTurquoise2, and β-tubulin C-terminally labeled with FlAsH. Cell was incubated with 5 μM nocodazole for >10 minutes to depolymerize microtubules.
(E) Boxplot of fluorescence lifetimes estimated from single-exponential models fit to the negative control fluorescence decays. n = 32, 11, and 6 cells for Neg Ctrl 1, 2, and 3, respectively.

Figure 1–figure supplement 4. Förster radius estimation by FLIM-FRET measurements and Monte Carlo protein simulations

(A) Fluorescence decay curves of cells expressing mTurquiose2-TC in the absence (green circle) and the presence (orange triangle) of FlAsH. A single-exponential model (black solid line) was fit to the fluorescence decay curve in the absence of FlAsH. For easier comparison, the fluorescence decay curves were normalized such that they asymptotically overlap.

(B) The conformational ensemble of the flexible tether between mTurquiose2 and TC were modeled by Monte Carlo protein simulations, and the distance, *r*, between mTurquiose2 (blue cartoon) and TC-FlAsH (green ball) was estimated. This distribution is denoted by *p(r)*.

(C) The measured fluorescence decay of mTurquiose2-TC-FlAsH (orange triangles, same as (A) but not normalized) plotted with the best-fit decay model (black dotted line, model described in the box and derived in Materials and Methods). Associated weighted residual (deviation of model from data, divided by the square root of the number of photons) plotted below. Fitting the decay model to the data estimated the Förster radius to be 5.90 nm ± 0.10 nm (SE).

Figure 1–figure supplement 5. Characterization and calibration of NDC80-kMT FLIM-FRET measurement

(A) The conformational ensemble of the flexible tether between mTurquoise2 and Nuf2 (red) and the disordered C-terminal tails of beta-tubulins around the NDC80 (green) were modeled by large-scale Monte Carlo protein simulations, which were then used to calculate the distances, r_i, between the mTurquoise2 and the TC motifs. FlAsH labeling was assigned to the TC motifs with 26.1% probability (which is the measured labeling fraction of beta-tubulin). Fluorescence lifetimes of the mTurquoise2 were calculated for randomly sampled sets of distances, $\vec{r} = (r_i)$, based on which fluorescence decay curves were simulated.

(B) (top) Fluorescence decay curves for various distances between NDC80 and MT were simulated and then were fit using single- and two-exponential decay models. Difference in Bayesian information criteria (BIC) between single- and double-exponential models is plotted against the NDC80-MT distance. Data points are mean and SD. (bottom) Example simulated fluorescence decay curves (green dots) for 0, 4, and 9 nm NDC80-MT distances are plotted with the best-fit single- (blue line) and two-exponential (red line) models. Corresponding smoothened weighted residuals plotted below.

(C) Fluorescence decay curves for various NDC80 binding fractions (f_b) were simulated and fit by using two-exponential decay model to estimate FRET fraction (f_{FRET}). (left) Three example simulated fluorescence decay curves (green dots) for 0%, 30%, and 60% binding fractions with the best-fit two-exponential models (red line), and the corresponding smoothened weighted residuals plotted below. (right) NDC80 FRET fractions (f_{FRET}) plotted against NDC80 binding fractions (f_b) (blue dots), and the linear fit (black line). Gray shaded area represents the uncertainty in the slope, which was determined from the uncertainties in the measured beta-tubulin labeling fraction and Förster radius (see Supplemental experiments in Materials and Methods).

Figure 3–figure supplement 1. K-K distance and kinetochore velocity are not correlated
(A) Each data point represents the fraction of leading kinetochores within a group of kinetochores with similar K-K distances. Gray region is the 95% confidence interval of the linear fit.
(B) Histogram of K-K distances of leading (green) and trailing (orange) kinetochores.

Figure 3–figure supplement 2. NDC80 FRET fraction of poleward- and anti-poleward-facing kinetochores in STLC-treated cells with monopolar spindles

NDC80 FRET fraction vs. K-K distance for poleward-facing kinetochores (purple square, same as Figure 3D) and anti-poleward-facing kinetochores (pink triangles) in cells treated with 5 µM STLC (n = 16 cells, 493 kinetochores/data point). Data points are the mean, y-error bars the SEM, and the x-error bars the interquartile ranges within groups of kinetochores with similar K-K distances. 3 µm scale bar. Gray area is the 95% confidence interval for the linear fit.

Figure 4–figure supplement 1. Supplemental data for Aurora B inhibition experiments
(A) Time course of NDC80 FRET fraction in response to 0.03% DMSO (n = 5 cells, negative control for Figure 4A).
(B) The design of Aurora B FRET biosensor. The FRET sensor contains a kinesin-13 family Aurora B substrate (gray) whose phosphorylation results in its binding to the forkhead-associated domain (FHA2, green) in the sensor, which constrains the sensor in the open non-FRET state. Therefore, measuring the non-FRET fraction of the FRET sensor allows the quantification of Aurora B activity.
(C) Time course of the non-FRET fraction of cytoplasmic Aurora B FRET sensor in response to 0.03% DMSO (n = 3 cells, negative control for Figure 4E).
(D) Time course of the non-FRET fraction of Nuf2-targeted Aurora B FRET sensor in response to 3 µM ZM447439 (n = 9 cells). Black squares and error bars are the weighted mean and SEM of the data points (green circles) in equally spaced time intervals of 1 minute. Red solid and dashed lines are the best-fit exponential model and its 95% confidence interval, respectively. 5 µm scale bars.

Figure 5–figure supplement 1. NDC80 FRET fraction of poleward and anti-poleward kinetochores in STLC-induced monopolar spindles of 9A-Hec1-expressing cells and haspin inhibited cells.
NDC80 FRET fraction vs. K-K distance for poleward-facing kinetochores (purple squares, same as Figure 5) and anti-poleward-facing kinetochores (pink triangles) in (A) 9A-Hec1-expressing cells treated with 5 μM STLC (n=10 cells, 855 kinetochores/data point) and (B) endogenous-Hec1-expressing cells treated both with 10 μM 5-ITu and 5 μM STLC (n = 12 cells, 564 kinetochores/data point). Data points are the mean, y-error bars the SEM, and the x-error bars the interquartile ranges within groups of kinetochores with similar K-K distances. 3 µm scale bar.