

Fusing Mobile Phone Sensing and Brain Imaging to Assess Depression in College Students: A Proof-of-Concept study

March 5th, 2018

Authors:

Jeremy F. Huckins¹, Alex W. daSilva¹, Rui Wang², Weichen Wang², Elin L. Hedlund¹, Eilis I. Murphy¹, Richard B. Lopez¹, Courtney Rogers¹, Paul E. Holtzheimer^{3,4}, William M. Kelley¹, Todd F. Heatherton¹, Andrew T. Campbell²

¹Department of Psychological and Brain Sciences, Dartmouth College, Hanover, NH

²Department of Computer Science, Dartmouth College, Hanover, NH

³National Center for PTSD, White River Junction, VT;

⁴Department of Psychiatry, Dartmouth-Hitchcock Medical Center, Lebanon, NH

Keywords:

Depression, mental health, smartphone, screen time, fMRI, resting-state

Abstract:

As smartphone usage has become increasingly prevalent in our society, so have rates of depression, particularly among young adults. Individual differences in smartphone usage patterns have been shown to reflect individual differences in underlying affective processes such as depression (Wang et al., *in press*), where depressive symptoms and duration of phone usage were previously linked. In the current study, we identified a positive relationship between smartphone screen time (e.g. phone unlock duration) and resting-state functional connectivity (RSFC) between the subgenual cingulate cortex (sgCC), a brain region implicated in depression and antidepressant treatment response, and regions of the ventromedial/orbitofrontal cortex, such that increased phone usage was related to stronger connectivity between these regions. We then used this cluster to constrain subsequent analyses looking at depressive symptoms in the same cohort and observed replication in a separate cohort. We believe the data and analyses presented here provide proof-of-concept analyses and a first step in combining functional brain activity and smartphone usage patterns to better understand issues related to mental health. Smartphones are a prevalent part of modern life and show promise in being a tool for mental health diagnostics and neuroscience research.

Introduction:

Smartphone usage has become nearly ubiquitous in daily life at a time when depression rates are concurrently rising, particularly among college students. Smartphones contain a variety of sensors that can allow researchers to passively measure various behaviors of the phone's user. Previous research has linked smartphone usage to self-reported depressive symptoms (Matar Boumosleh & Jaalouk, 2017; Twenge, Joiner, Rogers, & Martin, 2018; Wang et al., *in press*). In parallel, depressive symptoms have been linked to brain connectivity using resting-state functional connectivity (RSFC) MRI (Greicius et al., 2007). The current manuscript has multiple goals. First, is to provide a proof-of-concept for linking passive mobile smartphone sensing technologies to brain connectivity measures that have also been linked to self-reported depressive symptoms across two independent cohorts. Second, is to briefly describe a variety of methods which could be used to combine results across these various data types on a larger scale. Third, is to identify preliminary links between a key behavior inferred from sensing (e.g. smartphone screen time) and brain connectivity metrics.

Depression assessment

Depressive disorders affect over 300 million people worldwide and have been ranked among the largest contributors to global disability since the early 1990s and currently rate as the single largest contributor to global disability (Ustün, Ayuso-Mateos, Chatterji, Mathers, & Murray, 2004). Despite this, the diagnosis of depression has remained largely unchanged; further, a reliable means of identifying persons at risk of becoming depressed remains absent. Psychology, psychiatry and neuroscience have long relied upon self-reported surveys and in-person interviews to measure symptoms, diagnose mental health disorders and identify appropriate treatment strategies (Horwitz, Wakefield, & Lorenzo-Luaces, 2016). As a result of staggering costs inflicted at both individual and societal levels, clinicians and researchers set out to redefine the way mental disorders are conceptualized in hopes of creating innovative identification and prevention strategies. The aforementioned aims have been synthesized in a research framework known as RDoC (Research Domain Criteria). RDoC's objective is to incorporate information across all planes of analysis ranging from cellular level data to person level self-report survey data to provide of a holistic picture of mental disorders (NIMH). A core principle within the RDoC framework is the notion that neuroscience will inform future psychiatric classification schemes; in other words, aid in moving towards the establishment of a neural biomarker for depression. Thus, of great importance is understanding the complete range of human behavior (and neurological functioning) from typical to atypical (Insel et al., 2010). The Patient Health Questionnaire (PHQ, with four, eight and nine question versions) is a reliable, short survey which has been validated in clinical settings and can be used to assess self-reported symptoms of depression that cause significant impairment and subjective distress (Cameron, Crawford, Lawton, & Reid, 2008; Kroenke et al., 2001; Kroenke et al., 2009), an approach in keeping within the RDoC research framework, seeking to explain individual variance in symptoms across domains, constructs, and units of analysis. Future methods to accurately diagnose depression may hold promise with the inclusion of techniques that capitalize on the passive collection of behavioral data through mobile sensors (e.g. smartphones).

Passive sensing

Passive sensing using mobile smartphone technology allows for the assessment of daily activities by the smartphone user without continual effort on their part. This increases the

frequency with which data can be collected and may reduce self-report bias, which is often a problem in prompted surveys (Ben-Zeev, Scherer, Wang, Xie, & Campbell, 2015; Rosenman, Tennekoon, & Hill, 2011). Smartphone ownership has increased steadily over the last decade, with over 75% of the US population owning one (Smith, 2017). In parallel, depression rates have increased over the last decade (Twenge et al., 2018). Prevalence of both smartphone ownership and depression rates are often reported as being higher in college-age students (Eisenberg, Hunt, & Speer, 2013; Nielsen.com, 2016). Screen time, e.g. the amount of time that the screen is unlocked and being used is a relatively simple metric to calculate. Screen time and unlock duration will be used interchangeably henceforth.

Resting-State Functional Connectivity

Blood-oxygenation-level dependent (BOLD) functional magnetic resonance imaging (fMRI) is a non-invasive way to study activity in the human brain. Changes in BOLD signal are highly correlated with changes in neuronal activity in the local area, particularly local field potentials (Logothetis, Pauls, Augath, Trinath, & Oeltermann, 2001). Resting-state functional connectivity (RSFC) measures the relationship between the time-courses of different regions, often by using the correlation of the time-series. While connectivity across the whole brain, or “functional connectome” is fairly similar across individuals, there are small individual differences in connectivity between individuals which can be reliably observed across time. There are a variety of factors which may potentially influence RSFC, including genetics, experiences across the lifetime and current physiological and emotional state (Birn et al., 2013; Patriat et al., 2013; Poldrack et al., 2015; Richiardi et al., 2015; Shehzad et al., 2009; Sinclair et al., 2015; Zuo et al., 2014).

Depression and neuroimaging

RSFC has been used successfully to distinguish between healthy controls and depressed individuals, even going so far as to distinguish between subtypes of depressed individuals (Berman et al., 2013; Drysdale et al., 2016; Greicius et al., 2007; Kaiser, Andrews-Hanna, Wager, & Pizzagalli, 2015). Task-based studies of self-referential processing have revealed that the sgCC is preferentially involved in processing valenced self-referential information (Moran, Macrae, Heatherton, Wyland, & Kelley, 2006; Somerville, Heatherton, & Kelley, 2006). Additionally, this region has been associated with antidepressant treatment response, and an area proximal to this has been used as a site of deep-brain stimulation for treatment-resistant depression (Holtzheimer, 2012; Mayberg et al., 2005).

Combining RSFC and mobile smartphone passive-sensing technology

There are a wide-variety of approaches that can be taken when combining high-dimensional data from multiple modalities. We wanted to answer the following question: do smartphone sensing features previously identified as being related to a depression show correlations with resting-state functional connectivity from a region previously identified to have aberrant connectivity in depressed individuals? In the current proof-of-concept study we decided to take a targeted approach, selecting screen time with mobile smartphone (e.g. unlock duration), a feature previously shown to be linked to depressive symptoms (Twenge et al., 2018; Wang et al., *in press.*) and a brain area, the subgenual cingulate cortex (sgCC) which has previously been identified as having aberrant RSFC in depressed individuals, and more recently has been used as a target for deep brain stimulation for treatment resistant depression (Greicius et al., 2007;

Holtzheimer, 2012; Mayberg et al., 2005). Furthermore, if there are regions identified in the passive-sensing unlock duration analysis and RSFC analysis, do these regions also show similar connectivity patterns when looking at the same correlations with brief surveys of self-reported depressive symptoms (PHQ4/8)? We expect that they would. Alternatively, depression may be a summation of a variety of factors and may be better understood by interrogating passive-sensing mobile technology and neuroimaging than self-reported scales. Keeping within the RDoC matrix, we assess a variety of units of analysis including brain connectivity with fMRI (physiological), passive-sensing of phone usage (behaviors) and both computer-based and phone-based depression scales (self-report).

Methods:

Study design

In the current study two separate cohorts of first-year undergraduate students were enrolled and analyzed separately for test-retest comparison. Individuals were enrolled in three study components: neuroimaging, smartphone sensing/EMA and online surveys. Two modified versions of the PHQ-9 were used: PHQ-8 and PHQ-4. PHQ-8 is the same as PHQ-9 with the question about suicidal ideation removed. This question was removed because the surveys are not monitored in real-time. PHQ-4 is a four question survey which includes two questions from the PHQ-8 and two from the GAD-7 as to assess both depressive and anxiety related symptoms (Kurt Kroenke, Spitzer, Williams, & Löwe, 2009). They are used because of their brief form. They may miss some of the nuances that the other inventories pick up on but have been found to have high internal reliability (Cronbach's Alpha > 0.8) and are correlated with diagnoses of clinically relevant depression (Cameron et al., 2008; Khubchandani, Brey, Kotecki, Kleinfelder, & Anderson, 2016).

Individuals completed an online survey to assess study eligibility (safe for MRI per Dartmouth Brain Imaging Center guidelines, no contraindications that would lead to signal loss, and owned an Android or iOS smartphone). If an individual was eligible and interested in participating in the study, she or he completed a battery of online surveys, including the PHQ-8 through REDCap (Harris et al., 2009). Individuals were then scanned during the academic term and had the StudentLife application (Wang et al., 2014) installed on their phone at or near the time of scanning. In Cohort 1, StudentLife data was collected from the time of scanning until the end of the term. In Cohort 2, StudentLife data was collected from the time of scanning and data collection is currently ongoing but the data presented here only continued until the end of the first term.

StudentLife

A smartphone application, StudentLife is used in the current study to collect a variety of data about smartphone usage and mood from participants. The application is installed on a participant's phone (iOS or Android) and collects data from the GPS, microphone, accelerometer and lock/unlock status. Data from these sensors are processed create variables that assesses the day-to-day and week-by-week impact of workload on stress, sleep, activity, mood, sociability, mental well-being and academic performance of students (Wang et al., 2014). The workflow of the current study includes data collected through StudentLife, MRI scanning sessions and self-reported surveys (Figure 1).




Figure 1. Summary graphic of the study workflow in the current study, showing raw data collection from both smartphones (StudentLife, passive sensing) and MRI (resting-state functional connectivity, sgCC seed-based analysis). Calculated features were selected based on previous research. Survey data was collected both online (REDCap, PHQ-8) and smartphone (StudentLife, Ecological Momentary Assessments, PHQ-4) sources.

Ecological Momentary Assessments

Students were prompted once a week within the StudentLife application during the term to complete a few short surveys as Ecological Momentary Assessments (EMA, one of which was PHQ-4 (Shiffman, Stone, & Hufford, 2008)). In the current study we are focusing on PHQ-4, a modified, shorter version of the PHQ-8 which in four questions provides a glimpse of depressive and anxious symptoms (two questions related to each).

Subjects

Subjects were first-year undergraduate students recruited from the Dartmouth College community. Cohort 1 included 151 subjects (94 female, mean age = 19.59, std = 1.69, range = 18-28) which were all scanned during the first term of their first year at Dartmouth. Cohort 2 included 106 subjects (75 female, mean age = 18.25, std = 0.63, range = 18-22) which were all scanned during the first term of their first year at Dartmouth. One subject was removed from the study for having an incompatible phone and one MRI session was stopped due to not reporting a permanent top retainer.

See [Table 1](#) for a summary of the number of individuals included in each analysis, grouped by Cohort. Subjects were only included in each analysis if they met the minimum number of time-points for smartphone-based StudentLife data and each analysis and had resting-state functional connectivity that passed quality control (see RSFC analysis methods section below for further details). Subjects had normal or corrected-to-normal visual acuity. Each

subject provided informed consent in accordance with the guidelines set by the Committee for the Protection of Human Subjects at Dartmouth College and received either course credit or monetary compensation for participating in the study.

Table 1. Summary of the number of subjects in each analysis.

	Cohort 1	Cohort 2
Total Scanned	151	106
RSFC Data (Passed QC)	145	93
PHQ-8	65	89
PHQ-4 (1-Day)	84	89
Unlock Duration (20-Days)	77	77

RSFC data collection

Apparatus

Cohort 1 imaging was performed on a Philips Intera Achieva 3-Tesla scanner (Philips Medical Systems, Bothell, WA). Cohort 2 imaging was performed on a Siemens MAGNETOM Prisma 3-Tesla scanner (Siemens Medical Solutions, Malvern, PA). Data for both cohorts was collected using a 32-channel phased array head coil. During scanning, participants viewed a white fixation cross on a black background projected on a screen positioned at the head end of the scanner bore, which participants viewed through a mirror mounted on top of the head coil.

Cohort 1 imaging

Anatomic images were acquired using a high-resolution 3-D magnetization-prepared rapid gradient echo sequence (MP-RAGE; 160 sagittal slices; TE, 4.6 ms; TR, 9.9 ms; flip angle, 8°; voxel size, 1 x 1 x 1 mm). Resting-state functional images were collected using T2*-weighted fast field echo, echo planar functional imaging sensitive to BOLD contrast (TR= 2500 ms; TE= 35 ms; flip angle= 90°; 3 x 3 mm in-plane resolution; sense factor of 2). Functional scanning was performed in one or two runs; during each run, 240 brain volumes (36 slices, 3.5 mm slice thickness, 0.5 mm skip between slices) were acquired, allowing complete brain coverage. As such, each participant completed between 10 and 20 minutes of RSFC scanning.

Cohort 2 imaging

Anatomic images were acquired using a high-resolution 3-D magnetization-prepared rapid gradient echo sequence (MP-RAGE; 192 sagittal slices; TE, 2.32 ms; TR, 2300 ms; flip angle, 8°; voxel size, 1 x 1 x 1 mm) with a Grappa 2 acceleration factor. Resting-state functional images were collected using T2*-weighted fast field echo, echo planar functional imaging sensitive to BOLD contrast (TR= 1190 ms; TE= 32 ms; flip angle= 63°; 2.4 x 2.4 mm in-plane resolution; SMS factor of 4). Functional scanning was performed in one or two runs; during each run, 605 volumes (46 slices, 3 mm slice thickness, no skip between slices) were acquired, allowing complete brain coverage. As such, each participant completed 12 and 24 minutes of RSFC scanning. Data for cohort 2 was processed and organized into BIDS format with datalad (Gorgolewski et al., 2016; Halchenko et al., 2017).

RSFC analyses

All processing was performed using a standard previously published processing stream (Power et al., 2014) with two exceptions: frame-displacement (FD) threshold was set to 0.25mm (instead of 0.2mm) and 36 motion parameters (instead of 24) were used for motion regression. Functional images were preprocessed to reduce artifacts, including: (i) slice-timing correction, (ii) rigid body realignment to correct for head movement within and across runs, (iii) within-run intensity normalization such that the intensity of all voxels and volumes achieved a mode value of 1000 scale with 10 units equal to ~1% signal change, (iv) transformation to a standardized atlas space (3 mm isotropic voxels) based on (Talairach & Tournoux, 1988), (v) frame censoring, (vi) nuisance regression (excluding censored frames), (vii) interpolation, and (viii) bandpass filtering ($0.009 < f < 0.08\text{Hz}$) following Power et al. (2014). Final correlation calculations between time-courses were calculated based upon *uncensored* frames.

Neurosynth analysis and subgenual cingulate cortex seedmaps

To identify an unbiased subgenual cingulate cortex (sgCC) seed to create voxelwise functional seed maps, an automated meta-analysis was performed using Neurosynth for the term “subgenual” (Yarkoni, Poldrack, Nichols, Van Essen, & Wager, 2011). Subgenual cingulate cortex seed maps were created from a seed placed at 0, 25, -10 (MNI coordinates), which was the peak of the term “subgenual” as of February 17th, 2017 and are centered around BA 25. The mean time-course from this seed was correlated with the time-course from every voxel within the brain. These seed maps, i.e. maps of resting-state connectivity from the subgenual region, were produced for each individual that passed quality control (more than 5 minutes of uncensored frames, see above for more details).

Combining data

Since the version of the StudentLife application used in the current study generates 182 features automatically, and with RSFC it is possible to generate thousands of features, it is necessary to minimize the number of features compared given the relatively small size of the Cohorts ($N < 100$). To minimize the number of features inspected, unlock duration was the only feature inspected given its simplicity to calculate and previously-identified relationship with PHQ-8 (Wang et al., *in press*). While many features were identified, we specifically chose unlock duration (e.g. screen time) as a simple feature both to calculate and to conceptualize as it can be considered similar to total phone screen time.

For all surveys analyzed here, one time-point was sufficient for a subject to be included in the current analyses. If there were multiple responses to ecological momentary assessments (EMAs, e.g. surveys prompted by the application) over the course of the term those responses were averaged. Individuals were included in the unlock duration analysis if they had 20 days of quality data with more than 16 hours of quality unlock duration data for each day included.

Group analyses and statistics

SgCC seedmaps from Cohort 1 were correlated with unlock duration sampled from smartphone usage with the StudentLife application. For each analysis, the degrees of freedom was $N-2$, with N being listed in [Table 1](#). Results from the unlock duration and sgCC correlational analysis from Cohort 1 were volume corrected to account for multiple comparisons using

AFNI's 3dClustSim ACF function. Results from the sgCC/unlock duration analysis were used to restrict the regions investigated in further analyses.

Visualization

All results were transformed into MNI space (Montreal Neurological Institute) and mapped onto the Conte69 template for volume-based slices or inflated surfaces for visualization (Van Essen, Glasser, Dierker, Harwell, & Coalson, 2012). Group results were viewed in Connectome Workbench Version 1.1.1 (Marcus, Fotenos, Csernansky, Morris, & Buckner, 2010).

Results

Unlock duration correlated with sgCC connectivity

In Cohort 1 exploratory whole-brain analyses of the correlation between unlock duration and sgCC seedmaps identified a large cluster (1620 voxels, 43,740mm³) in the ventromedial prefrontal cortex with a positive linear relationship (Figure 2, S1). This cluster extended from the anterior caudate to medial frontal orbitofrontal cortex and dorsally to medial prefrontal cortex. Information about peaks within this cluster can be observed in Table 2. To determine if these results replicated in Cohort 2, the cluster identified in Cohort 1 was used as a mask and voxels which showed a significant positive relationship between unlock duration and sgCC connectivity in Cohort two were identified. This identified a small cluster at -9, 30, -15 (peak T = 2.67, voxel extent = 13) and another small cluster at 12, 63, 12 (peak T = 1.98, voxel extent = 5) (Figure S2).




Figure 2. Exploratory analysis correlation sgCC RSFC seedmaps correlated with mean unlock duration identified a cluster with a positive relationship to unlock duration in the ventromedial prefrontal cortex ($p < 0.05$, volume corrected using ACF to 0.01, $k > 489$) shown on inflated lateral (top left), medial (middle left) and ventral (right) cortical surfaces. The sgCC seed is represented as a black sphere.

Table 2. Exploratory analysis correlation sgCC RSFC seedmaps correlated with mean unlock duration (smartphone screen time) identified 1 cluster in the ventromedial prefrontal cortex ($p < 0.05$, volume corrected using AFNI's ACF to $p < 0.01$, $k > 489$, 1620 voxels total). Peaks identified from xjview 9.6, showing 3 maxima per cluster, at least 8mm apart.

Best Estimate of Region	X	Y	Z	T
Caudate	-15	21	-9	4.29
Caudate	12	21	-9	3.64
Anterior sgCC	6	33	-12	3.34

Self-reported depression symptoms correlated with sgCC connectivity

Previous research (Wang and colleagues, 2018) identified a relationship between depressive symptoms and unlock duration. To determine if depressive symptoms and unlock duration had overlap in the brain connectivity (seed based subgenual RSFC) regressions for both computer-based pre-screening (PHQ-8), phone based post-scanning (PHQ-4 as EMA) were performed. Results from both of these analyses were masked with the cluster identified in Cohort 1's sgCC/unlock duration analysis. PHQ-8 computer-based surveys correlated with sgCC connectivity maps identified clusters in both Cohorts and identified a cluster which overlapped between the two. Cohort 1 revealed three significant clusters, which were located in bilateral OFC with peaks located at -21, 42, -12 (peak T = 3.19, voxel extent = 130), 24, 51, -9 (peak T = 2.55, voxel extent = 21), and -21, 21, -21 (peak T = 2.75, voxel extent = 8) (Figure 3, Table 3). Cohort 2 revealed two significant cluster, which were located in left OFC and medial rectus at -15, 36, -9 (peak T = 3.35, voxel extent = 24) and 0, 54, -21 (peak T = 2.62, voxel extent 26) respectively. Overlap between significant clusters of the two Cohorts identified an 11-voxel cluster with located in left OFC, with a peak at -21, 39, -9 in Cohort 1 (peak T = 2.6) and -15, 33, -12 in Cohort 2 (peak T = 2.98). The peak of overlap from Cohort 1 is significant in the unlock duration analysis at $p < 0.01$ (T = 2.61) and significant at the Cohort 2 peak at $p < 0.005$ (T = 2.7) In addition to identifying a cluster with overlap between the two Cohorts for PHQ-8, visual inspection suggests proximal cortical regions beyond the area with overlap at a threshold of $p < 0.05$ and increased overlap, including right OFC at a more liberal threshold of $p < 0.1$.




Figure 3. PHQ-8 regression for sgCC connectivity seedmaps for A) Cohort 1, B) Cohort 2 (MNI Z of -3 to -27 in steps of 3 and C) overlap between Cohorts (MNI Z of -9 and -12), all masked by the results from Cohort 1 phone usage (unlock duration).

Table 3. Results for the correlation of sgCC RSFC seedmaps, correlated with PHQ-8, masked by phone screentime results (Top and Middle). Overlap between Cohort 1 and Cohort two for sgCC RSFC seedmaps correlated with PHQ-8 (Bottom).

Cohort 1						
Best Estimate of Region	X	Y	Z	T	Extent	
Left OFC	-21	42	-12	3.19	130	
	-21	51	-15	3.12	Subpeak	
	-6	48	-21	3.04	Subpeak	
Left OFC	-21	21	-21	2.75	8	
Right OFC	24	51	-9	2.55	21	
	18	42	-9	2.52	Subpeak	
Cohort 2						
Best Estimate of Region	X	Y	Z	T	Extent	
Left OFC	-15	36	-9	3.35	24	
Medial Rectus	0	54	-21	2.62	26	
Cohort Overlap						
Cohort	X	Y	Z	T	Extent	
1	-15	33	-12	2.6018	11	
2	-21	39	-9	2.98	11	

PHQ-4 EMAs correlated with sgCC connectivity maps identified peaks in Cohort 1 and 2, but there was no overlap in the clusters between the Cohorts (Table S1). In Cohort 1 there were 2 significant clusters identified, with peaks at -21, 72, -3 (peak T = 2.99, voxel extent = 7) and 9, 57, 3 (peak T = 2.31, voxel extent = 7). In Cohort 2 there were 4 significant clusters identified, with peaks at -15, 30 -12 (peak T = 3.71, voxel extent = 139), 9, 57, -24 (peak T = 2.38, voxel extent = 7), 18, 33, -24 (peak T = 2.11, voxel extent = 5), and 24, 39, -15 (peak T = 1.87, voxel extent = 13).

Discussion

In the current manuscript, we provide proof-of-concept linking passive smartphone metrics, active smartphone-based surveys of mental health and computer-based surveys of mental health with brain connectivity measures. Specifically, RSFC between the subgenual cingulate cortex, a region previously implicated in depression, and nearby ventral prefrontal regions, was strongly related to unlock duration, such that more connectivity was associated with more screen time, which has been implicated as being related to self-reported depressive symptoms. The link between RSFC and individual differences has long been established but extending that and combining it with an individual's behavior inferred from smartphone sensors

provides exciting new directions. While the results presented here are a relatively simple analysis of complex, highly dimensional data, we discuss some methods which could be used to combine these highly multivariate and complex datasets in exciting ways.

Phone-related screen time, which we estimate here by using unlock duration, has previously been shown to be related to self-reported depression levels (Twenge et al., 2018; Wang et al., *in press*). An exploratory analysis in Cohort 1 of the correlation between unlock duration and sgCC seedmaps identified a large cluster which extended from the anterior caudate to medial frontal orbitofrontal cortex and dorsally to medial prefrontal cortex, a result which was partially replicated in Cohort 2 even though the sampling rate for screen time was greatly reduced. Next, to determine if depressive symptoms showed a similar pattern of connectivity between sgCC and ventral prefrontal cortex the cluster from Cohort 1's unlock duration analysis was used as a mask with PHQ-8, a commonly used survey to assess depressive symptoms in the general population. Two small clusters of overlap were identified in the left OFC, one of them neighboring voxels that were identified to replicate in the unlock duration analysis between the Cohorts. While these clusters are not large and would not necessarily survive volume correction on their own, observing similar regions across Cohorts and analyses suggests that there is a link between depressive symptoms and related behaviors and sgCC-OFC connectivity, particularly left OFC that should be further investigated. The PHQ-4, which contains 2 depression questions and 2 anxiety questions, did not show the same robust relationship across both Cohorts, with no voxels overlapping, although Cohort 2 identified a cluster in the left OFC which overlapped with results observed with PHQ-8 in both Cohorts. Connectivity between the sgCC seed (BA 25), located at 0, 25, -10 and the left OFC region around -15, 33, -12 shows a consistent relationship between self-reported depressive symptoms and screen time, which has previously been associated with depression. Increased connectivity between sgCC, a region involved in processing of valenced information about the self (Moran et al., 2004) and OFC, which is involved in valuation and reward processing has been linked increased depressive symptoms and *screen time* across both Cohorts. It seems quite plausible that regions involved in valence processing related to the concept of self and a more general reward valuation processing region would have increased connectivity in individuals with higher depressive symptoms.

As a proof-of-concept, we have shown that resting-state functional connectivity of the brain, as measured with MRI, in two separate Cohorts of individuals, with two separate MRI's and two separate versions of the StudentLife applications show similarity in the results observed. The cluster identified with the unlock duration analysis covered an extent similar to that of the limbic network previously identified (Choi, Yeo, & Buckner, 2012; Yeo et al., 2011). Due to the constraints we imposed on the analysis, all of the subsequent results were within this area, but noticeably, many of the results were proximal to the left OFC, which is also a member of a set of nodes which are commonly activated during reward processing and can form their own system (Huckins et al., *submitted*) and is identified as a peak in reverse-inference meta-analyses of reward using Neurosynth (Yarkoni et al., 2011).

Limitations and Future Directions:

The current study is presented as a proof-of-concept and has several limitations and ways in which future research may provide a more comprehensive survey of the relationships between the diverse set of features provided from passive smartphone sensing, functional brain connectivity measures and self-reported measures of depression or other mental health metrics. The moderate sample size precludes the use of some machine learning methods. Furthermore,

the relatively small number of clinically depressed individual in the current sample weighs the results heavily on the RSFC and passive-sensing features from those individuals. Test-retest within the moderately sized samples allows for identification of factors with reliable cross-cohort replicability in RSFC both and passive-sensing features. Ideally similar sensing features could be collected across many sites, allow for identification and characterization of depressive subtypes that span across passive-sensing and RSFC as has been done by Drysdale and colleagues (2016) with RSFC and survey data. In the current study, particularly Cohort 2 in which data quality was actively monitored, we maintained a relatively large majority of individuals from those scanned (see Table 1). The sample sizes used here would have been considered relatively large several years ago. Increased sample sizes in the current study would help future analyses given the large number of features from both passive mobile smartphone sensing and RSFC. An outstanding question is if long-term changes in depressive symptoms can be better predicted by RSFC or smartphone sensing metrics at the initiation of the study or if changes in either of these over time parallel depressive symptoms. Ideally to assess this a large number of individuals would be tracked over multiple years. In the second Cohort our working group aims to track them over multiple years while eventually increasing the number of individuals enrolled. Furthermore, including multiple sites, as the ABCD study (Volkow et al., 2017) does, would increase applicability to a wider population. Multiple research sites are currently collecting MRI data, self-reported surveys and smartphone sensing metrics. As unresolved issue is what, exactly is the optimal approach to analyze the huge amounts of multivariate data produced by these methods.

Application changes Between cohorts

In the current study, unlock duration data collection changed between the cohorts. In Cohort 1, unlock duration was sampled for 1 minute out of every 4 minutes, while in Cohort 2 unlock duration was sampled for 1 minute every 10 minutes. This change was instituted to optimize battery life, a primary limitation to users keeping the StudentLife app on their phone. By decreasing the ratio of time sampled from 0.25 to 0.1, our ability to accurately estimate unlock duration decreases. As with all passive and active smartphone features, the ability to collect data must be weighed against the invasiveness to the user experience, either through app prompts or decreased battery life and phone speed.

Temporal factors related to school

The demands of the academic term provide a generally applicable path of stress which is shaped over the term. Avoiding, or potentially purposefully collecting MRI data during finals, which may be particularly stressful, or during popular social weekends may lead to changes in stress levels, sleep patterns and other variables which could alter connectivity patterns and self-reported behavioral data that would have otherwise been observed. In the study herein, we attempted to scan before finals and avoid well-known “party weekends”. Future studies may be able to capitalize on temporal differences in stress and depression levels by scanning at these peak times of stress or sleep deprivation and comparing that data to less stressful times, such as the beginning of the term.

Voxelwise Resting-State Functional Connectivity

A relatively simple first-pass method is to target specific region and feature pairs. If there are *a priori* hypotheses related to the topic of interest it may be possible to look at connectivity

from one region using seed maps or between a small number of regions and relate them to specific passive-sensing features. As shown here this is plausible but even correlating seed maps with 1 sensing variable leads to potential multiple comparisons issues based on the 50,000+ voxels in the brain using a 3mm³ voxel size. Recent statistical simulations have suggested an increased false-positive rate associated with older versions of 3dClustSim, a function of AFNI (Cox, Chen, Glen, Reynolds, & Taylor, 2017). Indeed, the authors of 3dClustSim now suggest using a different algorithm with the same program, the autocorrelation function (ACF) with a high p-value threshold per voxel to minimize the possibility of false-positives. In some datasets, at lower p-value thresholds ACF requires a much larger voxel-extent than the old version of 3dClustSim. The increased voxel-extent may make it difficult to identify smaller functional regions in a whole-brain regression using a lower per-voxel p-value threshold ($p < 0.05$). This shift in accepted methods decreases the rate of the false-positives but requires a larger expected functional region, a very strong effect size or a very large number of participants. Across all possible methods presented here there are a variety of factors which should be taken into consideration to decrease false positive rates. Having a large number of subjects to draw data will increase the portion of the population sampled.

If possible having two distinct Cohorts to analyze then looking for overlap in results between the Cohorts would decrease false positives due to random sampling, Cohort specific variance, and further increase the total size of individuals sampled. The above factors apply to most any study. With passive smartphone mobile sensing there are many features which can be measured or computed based on the intersection of multiple features. For example, “phone unlock duration” is a very simple metric, which measure the time that the smartphone was unlocked. This can be further broken down into location specific features, such as “phone unlock duration at dorm” or “phone unlock duration at study places” by looking at the intersection of location on a geo-tagged campus and “phone unlock duration. Given the large number of initial features that can be calculated, along with the nearly endless number of meta-features that can be generated, making sure that the feature is relatively simple to calculate and interpret should be at the forefront of anyone analyzing passive-mobile phone sensing features. Furthermore, such features should be validated to make sure they are actually measuring the effect or phenomena they are supposed to.

Typically, only features with sensing data from many days should be used to get a more stable estimate of that features’ value. While putting a sensing application of many students’ phones may seem like a plausible method for maximizing data collect, there are a variety of factors which can lead to reduced data collect, potentially rendering an individual’s sensing data unusable. Phone operating system (OS) updates can often change application permission or render the sensing application completely useless. To avoid this beta testing should be done as early as possible and new versions of the application that are compatible with the latest OS pushed to participants. Participant non-compliance or attrition is another important factor to consider. Individuals may delete the application, limit its permissions within the OS or otherwise limit the researcher's’ ability to accurately measure data. Clearly, it is the individual's choice to continue to participate in any study, particularly one where large amounts of data are being collected (anonymously) on their habits. It may be difficult for the researcher to determine if the individual has deleted the application or simply not uploaded their data in while. Finally, a rate of attrition is expected in all longitudinal studies and some individuals may simply decide that they do not wish to continue their participation in the study.

Whole-brain and network-based connectivity

A possible method to deal with the large number of comparisons related to voxelwise or whole-brain connectivity is to simply look at connectivity between a set of predefined regions or parcellation (Huckins et al., *submitted*) (Gordon et al., 2016; Poldrack et al., 2015; Power et al., 2011; Yeo et al., 2011). Connectivity between each pair of regions can be correlated with the sensing feature of interest. Unfortunately, many of the commonly used parcellations have many nodes, which increases the total number of comparisons in a nonlinear manner as the number of nodes increases. The number of comparisons can soon approach the number of comparisons evident when using voxelwise seed maps without methods such as voxel extent to appropriately correct for the associated multiple comparisons.

A simple but perhaps relatively unsophisticated method is to calculate mean connectivity within a functional system or network. The system or network would be determined off of data driven approach such community detection using a random walk technique like InfoMap (Rosvall & Bergstrom, 2008) or regions identified as being part of a coherent functional system using another method. In this approach, the mean of all Fisher r -to- z transformed correlation values between nodes of interest is calculated. For example, mean connectivity within the Cingulo-Opercular network would be calculated between all nodes or parcels belonging to that network. Between-network or system connectivity can also be calculated by taking the mean of all pairwise connections between the two networks of interest. This can greatly reduce the number of total connections observed, thus reducing the multiple comparisons problem mentioned under the whole-brain connectivity section. One drawback to this method is that it is not selective about which connections it is using in the calculation - specifically, that it may be and probably is including connections that are not physiologically or psychologically relevant.

A plausible way to reduce the number of connections by selecting ones that are likely to be “real”, such that information may actually travel through that connection on the neural level, even if not on a first-order or even second-order synapse. Multiple approaches have been taken to identify meaningful connections. Within or between networks there are likely to be positive and negative correlations, which then somewhat cancel out. One could take the absolute value of each connection before averaging across the network, but this would introduce bias in any connections with a distribution of correlation values that included positive and negative values. Values of correlation, or connectivity measures in the brain vary by orders of magnitude. Identifying a multiscale network backbone that accounts for important connections within and between communities, regardless of the connectivity strength would be a method to decrease the number of connections analyzed. One way of identifying the network backbone is to use the z -value from each connection as the weight, or amount of information that could travel between the two brain regions that the connectivity was estimated from. A group did just this (Serrano, Boguñá, & Vespignani, 2009), identifying connections which are statistically relevant across multiple scales of connectivity, work which has been extended non-parametrically (Foti, Hughes, & Rockmore, 2011). By identifying the network backbone for each individual, it may be plausible to identify a variety of subcategories or continuums of depression along which different symptom severities fall for each individual, along with passive smartphone monitoring will allow for greater insight into interactions of behavioral, self-report and physiological RDoC matrix criteria.

Data reduction techniques

A variety of techniques can be used to extract information from data that are both longitudinal and high-dimensional; that is, situations where the data are collected from participants at multiple time points and the number of covariates begins to approach, or even surpasses the number of subjects in the dataset (Cheng, Honda, Li, & Peng, 2014; Chu, Li, & Reimherr, 2016; L. Wang, Zhou, & Qu, 2012; Zipunnikov et al., 2014).

As has been mentioned repeatedly above, both with resting-state and passive smartphone sensing there are a large quantity of features and analyses that can be generated. In the current study we chose features that were reasonable based on previous data but are unlikely to be the optimal features that describe the relationship between depression, passive mobile sensing and brain connectivity. Multiple approaches could be taken with data from both sources. One approach which would greatly decrease the number of features that were necessary including trying to create a singular propensity metric, or biomarker of depression for both the resting-state fMRI data and a separate one for the sensing data then observing the relationship between the two. Alternatively, data reduction techniques such as independent component analysis could be applied to each group then the relationship between them could be measured. Many researchers have taken a “risk” or “propensity” score approach, where they generate models which contain predictive variables (gender, substance use, family history) pertinent to the outcome of interest and use the propensity score as a regressor when doing analyses at the group or individual difference level (Hansen et al., 2012; Stuart, 2010). This could be applied to smartphone data, but only once appropriate sensor features, and model have been calculated. By creating a unitary risk feature multiple comparison issues can be greatly mitigated. Data reduction techniques that account for variance that is common between two data modalities such as joint ICA, parallel ICA and CCA-Joint ICA, which has been implemented for combining high-dimension data across fMRI and genetic data (FusionICA, available from <http://mialab.mrn.org/software/fit/>).

Unresolved Questions about Directionality and Timing

In the current sample, resting-state fMRI data is from 1 time-point while mobile smartphone sensing data is dynamic and collects data over a longer period of time. An unresolved question is if changes in fMRI data across multiple sessions reflects or predicts changes in smartphone usage. Likely a more sensitive measure would be to do the reverse - using changes in smartphone usage, which is continuously monitored, to predict when there may be changes in brain connectivity as measured by fMRI. Changes in depressive symptoms have been successfully predicted with passive smartphone features (Wang et al., *in press*), and may be useful for signaling when an individual should be referred to clinical services or brought in for a subsequent fMRI session.

Moderating Factors of RSFC

RSFC has repeatedly been shown to be relatively stable across individuals and time, displaying similar network structure across thousands of individuals. While similar network structure and connectivity patterns are observed between sites, preprocessing methods, and Cohorts, differences between individuals are observed across individual differences in personality, affect and current mood have been related to alterations in RSFC. Furthermore, individual differences in the network structure on an individual level have been observed. Properly mapping individual differences in networks across the cortex would allow for better cross-subject alignment. The network assignment of particular regions may in itself be linked to

depressive symptoms, while lining up networks would allow for the proper comparison of networks across individuals. Additionally, the current state physiological state an individual is in, such as food satiety or caffeine status can influence their mood (Rogers & Lloyd, 1994) and has also been shown to influence an individual's brain connectivity (Poldrack et al., 2015). While there are a variety of factors that can influence RSFC, reliable individual differences across brain disorders have been observed in previous studies and here. As the predictive accuracy of RSFC or other neuroimaging methods increases the field may move closer to using MRI as a biomarker of depression, as has been done with physical pain (Atlas, Bolger, Lindquist, & Wager, 2010; Wager et al., 2013).

Conclusions:

In summary, we have identified proof-of-concept relationships between resting-state functional connectivity of the brain, web-based self-reported surveys of depressive symptoms (PHQ-8), a passive mobile smartphone sensing feature (unlock duration) and mobile smartphone based ecological momentary assessments of depressive symptoms (PHQ-4). An important mental health implication is that the amount of time spent using a phone is correlated with depressive symptoms. Further, these symptoms, both before time-of-scanning and after time-of-scanning, show a relationship with connectivity between areas implicated in depression, reward and processing of valenced self-relevant material. These results predominantly replicate across two separate cohorts, increasing the applicability and scope of the findings herein. Although the current results do not elucidate causality in the relationship between screen-time, depression and brain connectivity, future work should aim to do so, especially given recent changes to public policy, with professional groups such as the American Academy of Pediatrics providing suggesting screen-time limits and policy and investor groups calling on media device makers such as Apple and other phone makers. In the current work we replicate previous research, replicate results across multiple MRI's and cohorts all while combining data from a wide variety of sources. The analyses done here are by no-means comprehensive and we hope that the findings of this study and future research methods proposed herein are useful to a wide-range of researchers. Ultimately continuation and extensions of this research has the potential to provide important insights into mental health, as well as inform psychological treatments and other interventions.

Funding:

NIMH #'s 5R01MH059282-12 and 5R01DA022582-10.

Acknowledgements:

We would like to acknowledge Yaroslav O. Halchenko for assistance with MRI data conversion and BIDS organization and Terry Sackett for assistance with MRI data collection.

References

- Atlas, L. Y., Bolger, N., Lindquist, M. A., & Wager, T. D. (2010). Brain Mediators of Predictive Cue Effects on Perceived Pain. *Journal of Neuroscience*, *30*(39), 12964–12977. <https://doi.org/10.1523/JNEUROSCI.0057-10.2010>
- Ben-Zeev, D., Scherer, E. A., Wang, R., Xie, H., & Campbell, A. T. (2015). Next-generation psychiatric assessment: Using smartphone sensors to monitor behavior and mental health. *Psychiatric Rehabilitation Journal*, *38*(3), 218–226. <https://doi.org/10.1037/prj0000130>
- Berman, M. G., Yourganov, G., Askren, M. K., Ayduk, O., Casey, B. J., Gotlib, I. H., ... Jonides, J. (2013). Dimensionality of brain networks linked to life-long individual differences in self-control. *Nature Communications*, *4*, 1373. <https://doi.org/10.1038/ncomms2374>
- Birn, R. M., Molloy, E. K., Patriat, R., Parker, T., Meier, T. B., Kirk, G. R., ... Prabhakaran, V. (2013). The effect of scan length on the reliability of resting-state fMRI connectivity estimates. *NeuroImage*, *83*, 550–558. <https://doi.org/10.1016/j.neuroimage.2013.05.099>
- Cameron, I. M., Crawford, J. R., Lawton, K., & Reid, I. C. (2008). Psychometric comparison of PHQ-9 and HADS for measuring depression severity in primary care. *British Journal of General Practice*, *58*(546), 32–36. <https://doi.org/10.3399/bjgp08X263794>
- Cheng, M.-Y., Honda, T., Li, J., & Peng, H. (2014). Nonparametric independence screening and structure identification for ultra-high dimensional longitudinal data. *The Annals of Statistics*, *42*(5), 1819–1849. <https://doi.org/10.1214/14-AOS1236>
- Choi, E. Y., Yeo, B. T. T., & Buckner, R. L. (2012). The organization of the human striatum estimated by intrinsic functional connectivity. *Journal of Neurophysiology*, *108*(8), 2242–2263. <https://doi.org/10.1152/jn.00270.2012>
- Chu, W., Li, R., & Reimherr, M. (2016). Feature screening for time-varying coefficient models with ultrahigh-dimensional longitudinal data. *The Annals of Applied Statistics*, *10*(2), 596–617. <https://doi.org/10.1214/16-AOAS912>
- Cox, R. W. ., Chen, G., Glen, D. R. ., Reynolds, R. C. ., & Taylor, P. A. . (2017). fMRI clustering and false-positive rates. *Proceedings of the National Academy of Sciences*, *114*(17), E3370–E3371. <https://doi.org/10.1073/pnas.1614961114>
- Drysdale, A. T., Grosenick, L., Downar, J., Dunlop, K., Mansouri, F., Meng, Y., ... Liston, C. (2016). Resting-state connectivity biomarkers define neurophysiological subtypes of depression. *Nature Medicine*, (December). <https://doi.org/10.1038/nm.4246>
- Eisenberg, D., Hunt, J., & Speer, N. (2013). Mental health in American colleges and universities: variation across student subgroups and across campuses. *The Journal of Nervous and Mental Disease*, *201*(1), 60–67. <https://doi.org/10.1097/NMD.0b013e31827ab077>
- Foti, N. J., Hughes, J. M., & Rockmore, D. N. (2011). Nonparametric Sparsification of Complex Multiscale Networks. *PLoS ONE*, *6*(2), e16431. <https://doi.org/10.1371/journal.pone.0016431>
- Gordon, E. M. E. M., Laumann, T. O. T. O., Adeyemo, B., Huckins, J. F. J. F., Kelley, W. M. W. M., & Petersen, S. E. (2016). Generation and Evaluation of a Cortical Area Parcellation from Resting-State Correlations. *Cerebral Cortex*, *26*(1), 288–303. <https://doi.org/10.1093/cercor/bhu239>
- Gorgolewski, K. J., Auer, T., Calhoun, V. D., Craddock, R. C., Das, S., Duff, E. P., ... Milham, M. P. (2016). The brain imaging data structure, a format for organizing and describing outputs of neuroimaging experiments. *Scientific Data*, *3*, 160044. <https://doi.org/10.1038/sdata.2016.44>

- Greicius, M. D., Flores, B. H., Menon, V., Glover, G. H., Solvason, H. B., Kenna, H., ... Schatzberg, A. F. (2007). Resting-State Functional Connectivity in Major Depression: Abnormally Increased Contributions from Subgenual Cingulate Cortex and Thalamus. *Biological Psychiatry*, 62(5), 429–437. <https://doi.org/10.1016/j.biopsych.2006.09.020>
- Halchenko, Y., Hanke, M., Poldrack, B., Debanjum, Alteva, G., gors, jason, ... Christian, H. (2017). datalad/datalad 0.9.1. <https://doi.org/10.5281/ZENODO.1000098>
- Hansen, R. A., Dusetzina, S. B., Ellis, A. R., Stürmer, T., Farley, J. F., & Gaynes, B. N. (2012). Risk of adverse events in treatment-resistant depression: propensity-score-matched comparison of antidepressant augment and switch strategies. *General Hospital Psychiatry*, 34(2), 192–200. <https://doi.org/10.1016/j.genhosppsy.2011.10.001>
- Harris, P. A., Taylor, R., Thielke, R., Payne, J., Gonzalez, N., & Conde, J. G. (2009). Research electronic data capture (REDCap)—A metadata-driven methodology and workflow process for providing translational research informatics support. *Journal of Biomedical Informatics*, 42(2), 377–381. <https://doi.org/10.1016/j.jbi.2008.08.010>
- Holtzheimer, P. E. (2012). Subcallosal Cingulate Deep Brain Stimulation for Treatment-Resistant Unipolar and Bipolar Depression. *Archives of General Psychiatry*, 69(2), 150. <https://doi.org/10.1001/archgenpsychiatry.2011.1456>
- Horwitz, A. V., Wakefield, J. C., & Lorenzo-Luaces, L. (2016). *History of Depression*. (R. J. DeRubeis & D. R. Strunk, Eds.) (Vol. 1). Oxford University Press. <https://doi.org/10.1093/oxfordhb/9780199973965.013.2>
- Insel, T., Cuthbert, B., Garvey, M., Heinssen, R., Pine, D. S., Quinn, K., ... Wang, P. (2010). Research domain criteria (RDoC): toward a new classification framework for research on mental disorders. *The American Journal of Psychiatry*, 167(7), 748–751. <https://doi.org/10.1176/appi.ajp.2010.09091379>
- Kaiser, R. H., Andrews-Hanna, J. R., Wager, T. D., & Pizzagalli, D. A. (2015). Large-Scale Network Dysfunction in Major Depressive Disorder: A Meta-analysis of Resting-State Functional Connectivity. *JAMA Psychiatry*, 72(6), 603–611. <https://doi.org/10.1001/jamapsychiatry.2015.0071>
- Khubchandani, J., Brey, R., Kotecki, J., Kleinfelder, J., & Anderson, J. (2016). The Psychometric Properties of PHQ-4 Depression and Anxiety Screening Scale Among College Students. *Archives of Psychiatric Nursing*, 30(4), 457–462. <https://doi.org/10.1016/j.apnu.2016.01.014>
- Kroenke, K., Spitzer, R. L., & Williams, J. B. (2001). The PHQ-9: validity of a brief depression severity measure. *Journal of General Internal Medicine*, 16(9), 606–613. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11556941>
- Kroenke, K., Spitzer, R. L., Williams, J. B. W., & Löwe, B. (2009). An ultra-brief screening scale for anxiety and depression: the PHQ-4. *Psychosomatics*, 50(6), 613–621. <https://doi.org/10.1176/appi.psy.50.6.613>
- Kroenke, K., Strine, T. W., Spitzer, R. L., Williams, J. B. W., Berry, J. T., & Mokdad, A. H. (2009). The PHQ-8 as a measure of current depression in the general population. *Journal of Affective Disorders*, 114(1–3), 163–173. <https://doi.org/10.1016/j.jad.2008.06.026>
- Logothetis, N. K., Pauls, J., Augath, M., Trinath, T., & Oeltermann, A. (2001). Neurophysiological investigation of the basis of the fMRI signal. *Nature*, 412(6843), 150–157. <https://doi.org/10.1038/35084005>
- Marcus, D. S., Fotenos, A. F., Csernansky, J. G., Morris, J. C., & Buckner, R. L. (2010). Open access series of imaging studies: longitudinal MRI data in nondemented and demented older

- adults. *Journal of Cognitive Neuroscience*, 22(12), 2677–2684.
<https://doi.org/10.1162/jocn.2009.21407>
- Matar Boumosleh, J., & Jaalouk, D. (2017). Depression, anxiety, and smartphone addiction in university students- A cross sectional study. *PLOS ONE*, 12(8), e0182239.
<https://doi.org/10.1371/journal.pone.0182239>
- Mayberg, H. S., Lozano, A. M., Voon, V., McNeely, H. E., Seminowicz, D., Hamani, C., ... Kennedy, S. H. (2005). Deep Brain Stimulation for Treatment-Resistant Depression. *Neuron*, 45(5), 651–660. <https://doi.org/10.1016/j.neuron.2005.02.014>
- Moran, J. M., Macrae, C. N., Heatherton, T. F., Wyland, C. L., & Kelley, W. M. (2006). Neuroanatomical evidence for distinct cognitive and affective components of self. *Journal of Cognitive Neuroscience*, 18(9), 1586–1594. <https://doi.org/10.1162/jocn.2006.18.9.1586>
- Nielsen.com. (2016). MILLENNIALS ARE TOP SMARTPHONE USERS. Retrieved from <http://www.nielsen.com/us/en/insights/news/2016/millennials-are-top-smartphone-users.html>
- Patriat, R., Molloy, E. K., Meier, T. B., Kirk, G. R., Nair, V. A., Meyerand, M. E., ... Birn, R. M. (2013). The effect of resting condition on resting-state fMRI reliability and consistency: a comparison between resting with eyes open, closed, and fixated. *NeuroImage*, 78, 463–473. <https://doi.org/10.1016/j.neuroimage.2013.04.013>
- Poldrack, R. A., Laumann, T. O., Koyejo, O., Gregory, B., Hover, A., Chen, M.-Y., ... Mumford, J. A. (2015). Long-term neural and physiological phenotyping of a single human. *Nature Communications*, 6, 8885. <https://doi.org/10.1038/ncomms9885>
- Power, J. D., Cohen, A. L., Nelson, S. M., Wig, G. S., Barnes, K. A., Church, J. A., ... Petersen, S. E. (2011). Functional network organization of the human brain. *Neuron*, 72(4), 665–678. <https://doi.org/10.1016/j.neuron.2011.09.006>
- Power, J. D., Mitra, A., Laumann, T. O., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E. (2014). Methods to detect, characterize, and remove motion artifact in resting state fMRI. *NeuroImage*, 84, 320–341. <https://doi.org/10.1016/j.neuroimage.2013.08.048>
- Richiardi, J., Altmann, A., Milazzo, A.-C., Chang, C., Chakravarty, M. M., Banaschewski, T., ... Tahmasebi, A. (2015). Correlated gene expression supports synchronous activity in brain networks. *Science*, 348(6240), 1241–1244. <https://doi.org/10.1126/science.1255905>
- Rogers, P. J., & Lloyd, H. M. (1994). Nutrition and mental performance. *Proceedings of the Nutrition Society*, 53(2), 443–456. <https://doi.org/10.1079/PNS19940049>
- Rosenman, R., Tennekoon, V., & Hill, L. G. (2011). Measuring bias in self-reported data. *International Journal of Behavioural and Healthcare Research*, 2(4), 320. <https://doi.org/10.1504/IJBHR.2011.043414>
- Rosvall, M., & Bergstrom, C. T. (2008). Maps of random walks on complex networks reveal community structure. *Proceedings of the National Academy of Sciences of the United States of America*, 105(4), 1118–1123. <https://doi.org/10.1073/pnas.0706851105>
- Serrano, M. A., Boguñá, M., & Vespignani, A. (2009). Extracting the multiscale backbone of complex weighted networks. *Proceedings of the National Academy of Sciences of the United States of America*, 106(16), 6483–6488. <https://doi.org/10.1073/pnas.0808904106>
- Shehzad, Z., Kelly, A. M. C., Reiss, P. T., Gee, D. G., Gotimer, K., Uddin, L. Q., ... Milham, M. P. (2009). The resting brain: unconstrained yet reliable. *Cerebral Cortex (New York, N.Y. : 1991)*, 19(10), 2209–2229. <https://doi.org/10.1093/cercor/bhn256>
- Shiffman, S., Stone, A. A., & Hufford, M. R. (2008). Ecological momentary assessment. *Annual Review of Clinical Psychology*, 4, 1–32. Retrieved from

- <http://www.ncbi.nlm.nih.gov/pubmed/18509902>
- Sinclair, B., Hansell, N. K., Blokland, G. A. M., Martin, N. G., Thompson, P. M., Breakspear, M., ... McMahon, K. L. (2015). Heritability of the Network Architecture of Intrinsic Brain Functional Connectivity. *NeuroImage*. <https://doi.org/10.1016/j.neuroimage.2015.07.048>
- Smith, A. (2017). *Record shares of Americans now own smartphones, have home broadband*. Retrieved from <http://www.pewresearch.org/fact-tank/2017/01/12/evolution-of-technology/>
- Somerville, L. H., Heatherton, T. F., & Kelley, W. M. (2006). Anterior cingulate cortex responds differentially to expectancy violation and social rejection. *Nature Neuroscience*, 9(8), 1007–1008. <https://doi.org/10.1038/nn1728>
- Stuart, E. A. (2010). Matching Methods for Causal Inference: A Review and a Look Forward. *Statistical Science*, 25(1), 1–21. <https://doi.org/10.1214/09-STS313>
- Talairach, J., & Tournoux, P. (1988). *Co-Planar Stereotaxis Atlas of the Human Brain*. Direct (Vol. 270). New York: Thieme Medical Publishers, Inc.
- Twenge, J. M., Joiner, T. E., Rogers, M. L., & Martin, G. N. (2018). Increases in Depressive Symptoms, Suicide-Related Outcomes, and Suicide Rates Among U.S. Adolescents After 2010 and Links to Increased New Media Screen Time. *Clinical Psychological Science*, 6(1), 3–17. <https://doi.org/10.1177/2167702617723376>
- Ustün, T. B., Ayuso-Mateos, J. L., Chatterji, S., Mathers, C., & Murray, C. J. L. (2004). Global burden of depressive disorders in the year 2000. *The British Journal of Psychiatry: The Journal of Mental Science*, 184, 386–392. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/15123501>
- Van Essen, D. C., Glasser, M. F., Dierker, D. L., Harwell, J., & Coalson, T. (2012). Parcellations and hemispheric asymmetries of human cerebral cortex analyzed on surface-based atlases. *Cerebral Cortex (New York, N.Y. : 1991)*, 22(10), 2241–2262. <https://doi.org/10.1093/cercor/bhr291>
- Volkow, N. D., Koob, G. F., Croyle, R. T., Bianchi, D. W., Gordon, J. A., Koroshetz, W. J., ... Weiss, S. R. B. (2017). The conception of the ABCD study: From substance use to a broad NIH collaboration. *Developmental Cognitive Neuroscience*. <https://doi.org/10.1016/j.dcn.2017.10.002>
- Wager, T. D., Atlas, L. Y., Lindquist, M. A., Roy, M., Woo, C.-W., & Kross, E. (2013). An fMRI-Based Neurologic Signature of Physical Pain. *New England Journal of Medicine*, 368(15), 1388–1397. <https://doi.org/10.1056/NEJMoa1204471>
- Wang, L., Zhou, J., & Qu, A. (2012). Penalized Generalized Estimating Equations for High-Dimensional Longitudinal Data Analysis. *Biometrics*, 68(2), 353–360. <https://doi.org/10.1111/j.1541-0420.2011.01678.x>
- Wang, R., Chen, F., Chen, Z., Li, T., Harari, G., Tignor, S., ... Campbell, A. T. (2014). StudentLife. In *Proceedings of the 2014 ACM International Joint Conference on Pervasive and Ubiquitous Computing - UbiComp '14 Adjunct* (pp. 3–14). New York, New York, USA: ACM Press. <https://doi.org/10.1145/2632048.2632054>
- Wang, R., Wang, W., DaSilva, A., Huckins, J. F., Kelley, W. M., Heatherton, T. F., & Campbell, A. T. (n.d.). Tracking Depression Dynamics in College Students using Mobile Phone and Wearable Sensing. *Proc. ACM Interact. Mob. Wearable Ubiquitous Technology*.
- Yarkoni, T., Poldrack, R. A., Nichols, T. E., Van Essen, D. C., & Wager, T. D. (2011). Large-scale automated synthesis of human functional neuroimaging data. *Nature Methods*, 8(8), 665–670. <https://doi.org/10.1038/nmeth.1635>
- Yeo, B. T. T., Krienen, F. M., Sepulcre, J., Sabuncu, M. R., Lashkari, D., Hollinshead, M., ...

- Buckner, R. L. (2011). The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *Journal of Neurophysiology*, *106*(3), 1125–1165.
<https://doi.org/10.1152/jn.00338.2011>
- Zipunnikov, V., Greven, S., Shou, H., Caffo, B. S., Reich, D. S., & Crainiceanu, C. M. (2014). Longitudinal high-dimensional principal components analysis with application to diffusion tensor imaging of multiple sclerosis. *The Annals of Applied Statistics*, *8*(4), 2175–2202.
<https://doi.org/10.1214/14-AOAS748>
- Zuo, X.-N., Anderson, J. S. J. S., Bellec, P., Birn, R. M. R. M., Biswal, B. B. B. B., Blautzik, J., ... Milham, M. P. M. P. (2014). An open science resource for establishing reliability and reproducibility in functional connectomics. *Scientific Data*, *1*, 140049.
<https://doi.org/10.1038/sdata.2014.49>

Supplemental Material




Figure S1. Exploratory analysis correlating sgCC RSFC seedmaps correlated with mean unlock duration identified a cluster with a positive relationship to unlock duration in the ventromedial prefrontal cortex ($p < 0.05$, volume corrected using ACF to 0.01, $k > 489$) shown on axial slices.






Figure S2. Replication of the relationship between unlock duration and sgCC connectivity between Cohorts 1 and 2. The large cluster identified in Cohort 1 to have a positive relationship with unlock duration was used to restrict the area interrogated in Cohort 2, showing positive clusters in sgCC/vmPFC (sagittal view, X = -10) and MPFC (axial view, Z = 12). Replication clusters were observed at $p < 0.05$, with 5 or more contiguous voxels.



$p < 0.05$

$p < 0.001$

Figure S3. PHQ-4 regression for sgCC connectivity seedmaps for A) Cohort 1, B) Cohort 2 masked by the results from unlock duration in Cohort 1. No common regions were found between the two analyses.

Table S1. Exploratory analysis identifying peaks with a positive relationship from sgCC RSFC seedmaps correlated with phone-based surveys of depressive symptoms (EMA form of PHQ-4). Only clusters showing overlap with the Cohort 1 unlock duration analysis, with a positive relationship between PHQ-4 and connectivity with greater than 5 contiguous voxels are reported here.

Cohort 1					
Best Estimate of Region	X	Y	Z	T	Extent
Superior Frontal Gyrus	-21	72	-3	2.99	7
Medial PFC	9	57	3	2.31	7
Cohort 2					
Best Estimate of Region	X	Y	Z	T	Extent
Ventromedial PFC	-15	30	-12	3.71	139
	-9	51	-18	2.87	Subpeak
	3	48	-21	2.49	Subpeak
Anterior Medial OFC	9	57	-24	2.38	7
Right OFC	18	33	-24	2.11	5
Right OFC	24	39	-15	1.87	13
	15	51	-15	1.72	Subpeak
No Cohort Overlap Observed					