Perspective

Cost of resistance: an unreasonably expensive concept

Thomas Lenormand[1,2], Noémie Harmand[1], Romain Gallet[3]

[1] UMR 5175 CEFE, CNRS - Université Montpellier - Université P. Valéry - EPHE, Montpellier Cedex 5, France

[2] Radcliffe Institute for advanced study, Harvard University, Byerly Hall, 8 Garden Street, Cambridge, MA 02138

[3] INRA - UMR BGPI, Cirad TA A-54/K Campus International de Baillarguet 34398 Montpellier Cedex 5 France

Abstract

The cost of resistance, or the fitness effect of resistance mutation in absence of the drug, is a very widepsread concept in evolutionary genetics and beyond. It has represented an important addition to the simplistic view that resistance mutations should solely be considered as beneficial mutations. Yet, this concept also entails a series of serious difficulties in its definition, interpretation and current usage. In many cases, it may be simpler, clearer, and more insightful to study, measure and analyze the fitness effects of mutations across environments and to better distinguish those effects from 'pleiotropic effects' of those mutations.

Introduction

The study of resistance to antibiotics, insecticides, acaricides, fungicides, herbicides, chemotherapy drugs etc. is, for obvious reasons, a very active field of research. We are in the middle of a "crisis," which has important consequences for public health and agriculture (e.g. World Health Organization 2014; Ventola 2015). It is yet unclear whether we will be able to deal with "superbugs", "superweeds" and other "supermicrobes" in the near future. Studies have focused intensely on the genetic, cellular, and biochemical mechanisms responsible for resistance, but also on the fitness effect of those mutations.

Resistance mutations as beneficial mutations

Resistance evolution is a particular case of the more general situation of adaptation to new environmental conditions. Processes of adaptation have been intensely studied from Darwin's time, since refined by powerful population genetic concepts that have been put forward in the modern synthesis (Orr 2005). In brief, resistance mutation are beneficial mutations. They spread because they confer an obvious fitness advantage in presence of the drug (the antibiotic, insecticide, acaricide, fungicide etc.). By definition, a resistance mutation allows for survival while the susceptible wild-type simply dies when exposed to the drug. Hence, classically, the fitness benefit of a resistance mutation (relative to a susceptible one) depends on the fraction of the population exposed to the drug (since it determines the fraction of surviving susceptible genotypes). This is a black and white outcome, and it seems that there is little to understand beyond this obvious reasoning. Yet, in natural / real conditions (i.e. unlike in ecotoxicological tests performed in the laboratory), it is not very clear which dose of the drug is relevant. Concentrations of drugs vary at different temporal and geographical scales, within bodies (Levison and Levison 2009), within microhabitats, within regions etc (Thiele-Bruhn 2003; Depledge 2011). Notwithstanding, it is often assumed that a resistance mutation is associated with a selection coefficient measuring the rate at which it is expected to change in frequency in populations, like any other beneficial mutation. This selective advantage is not easy to estimate in the field, but is often thought to represent an inherent property of the mutation itself. However, this advantage must depend in some ways on the exposure to the drug, which is an environmental variable, and not a property of the mutation.

The context dependence of fitness effects

This point leads to a very simple and obvious idea: the selective effects of mutations depend on ecological conditions. This conclusion is somewhat trivial (Bell 2008), but it contradicts the naïve view that tends to essentialize the properties of mutations or genotypes (i.e. that characterize mutationnal properties as intrinsic). There is no such thing as a single selective effect of a mutation or a genotype. A mutation is not inherently beneficial, deleterious, or neutral. Rather, it depends on the ecological conditions, the genetic background, and on other alleles considered in comparison. Even the 'dominance' of mutations in diploids can be highly context and trait dependent (Bourguet *et al.* 1996; Manna *et al.* 2011). It is important to reiterate this point for several reasons. The first is that it tends to be simply ignored, even if this context dependence is never directly challenged. For instance, in molecular evolution, it

is customary to assign a mutation, without further specifications, in three categories: deleterious, neutral or beneficial (Eyre-Walker and Keightley 2007). Secondly, it makes little sense to study the selective effects of mutations while neglecting the diversity of real ecological conditions and the diversity of genetic backgrounds. Of course, it is always possible to think in terms of average, and this is the usual view when dealing with the diversity of genetic backgrounds in sexual populations (where recombination ensures that mutations 'experience' effectively the range of possible backgrounds)(Lande 1983; Chevin and Hospital 2008). Yet, for ecological variation, taking such an average may be very misleading. For example, a resistance mutation can spread somewhere (in a treated environment), but not elsewhere (in a non-treated environment), generating a situation of polymorphism (Jain and Bradshaw 1966; Suckling and Khoo 1993; Carrière *et al.* 1994; Guillemaud *et al.* 1998; Lenormand *et al.* 1998, 1999; Lenormand and Raymond 2000; Neve and Powles 2005; Labbé *et al.* 2009). Differences of effects across environments matters, not only their average, provided that the spatial scale of dose variation is larger than the scale of dispersal (Lenormand 2002).

The cost of resistance

This is where the concept of 'cost of resistance' became important. This cost is defined as the selection coefficient of resistance mutations in absence of treatments (or similarly in absence of predator, parasite or pathogens when considering resistance in the context of biotic interactions). The idea of a "lower adaptive value" of resistant genotypes in the absence of treatment can be traced back quite far (e.g. in Dobzhansky 1951). The term "reproductive disadvantage" or no specific term is used in this context in the 60s and 70s, in empirical or theoretical papers (Abedi and Brown 1960; Gillespie 1975; Hickey and McNelly 1975; Antonovics 1977; Georghiou and Taylor 1977; Curtis *et al.* 1978). The term 'cost of resistance' is relatively recent in comparison and started to be widespread only in the 80s, especially in the context of plant resistance to herbivores (Windle and Franz 1979; Leonard and Czochor 1980; Simms and Rausher 1987) or bacteria resistance to phage (Lenski 1988). It is in particular used in the influential paper of Anderson and May (1982) on coevolution and resistance to pathogens. Curiously, when dealing with resistance evolution in an abiotic context (i.e. to pesticides or antibiotics), the term 'cost of resistance' still refers in the 80s to the economic, not the evolutionary, cost of resistance (i.e. the extra monetary cost due to increasing pesticide dosage consecutive to resistance evolution), even in papers by May,

referring to evolutionary cost as 'back selection' (May and Dobson 1986). The first mentions of an evolutionary cost of resistance in an abiotic context seem to be in (Jacobs *et al.* 1988) about herbicide resistance, although the term "cost of tolerance" was used in studies of heavy metal tolerance in plants a bit earlier (Bradshaw 1984; Wilson 1988). The cost of resistance was then intensely investigated in the 90s (Bergelson and Purrington 1996; Andersson and Levin 1999 for reviews), with several methodological improvements (e.g. separating costs from effect of linked variation). By the end of the 90s, most pesticide resistance management models included the cost of resistance, i.e. the fact that resistance mutations could be selected against in non-treated environments (Lenormand and Raymond 1998). As this short historical overview shows, the concept of cost of resistance is relatively recent and was not used before the 80s. In particular, all the paper on local adaptation, clines and all the field of ecological genetics developed without the need to refer to this concept. For instance, in classical models such as Levene model (Levene 1953) or cline models (Haldane 1948; Nagylaki 1975), it was sufficient to talk about the selective effects of alleles in different environments. Why was the concept of cost of resistance introduced? At first sight, the concept seems asymmetrical and not very useful. For instance, nobody talks of the "cost of susceptibility" in treated environments, although it would be equally legitimate.

A likely reason is that the concept was helpful to bring attention to the fact that a mutation could be both beneficial or deleterious, depending on circumstances, something well known in ecological genetics, but somewhat ignored in resistance studies. It helped introduce some ecology in the understanding of the fitness effect of resistance mutations. This can have important consequences as the cost of resistance is a powerful force that can keep resistance in check (see e.g. Curtis *et al.* 1978 for an early model). Considering cost is also very important to predict the dynamics of resistance mutations in heterogeneous (treated and non-treated) environments.

A second reason is that the concept of cost was tightly associated to the notion of trade-off among traits, an idea borrowed from life history theory. When resistance is viewed as a trait, or a "defense" function, it is natural to consider that it may trade-off with other organismal traits and functions (e.g. in terms of resource). In other words, resistance mutations should influence many traits, i.e. have pleiotropic effects. Besides resistance, the variation of all these traits is likely to be deleterious, and therefore represent a 'cost', since these traits were previously optimized by natural selection. This connection with life history theory is entirely explicit in the first papers mentioning the concept (Simms and Rausher 1987; Lenski 1988;

Wilson 1988; Smith *et al.* 1991; Bergelson and Purrington 1996) and led to idea that the cost of resistance was caused by the pleiotropic effects of resistance mutations. Although the idea of trade-off among traits was initially present in this interpretation, it is often forgotten today: resistance mutations are simply viewed as 'pleiotropic'. Naturally, only considering pleiotropy, it seems very natural to think that the cost can evolve to be reduced, or even eliminated (which is also directly suggested by the word 'cost' itself). For instance, one could consider that compensatory evolution should attenuate these unwanted pleiotropic effects and restore optimal values for all traits, eliminating the cost. At first sight, the best proof for this reasoning is that cost-free resistance mutations are sometimes found along with the existence of modifier loci mitigating or even eliminating costs (McKenzie *et al.* 1982; Davies *et al.* 1996; Lenski 1998; Andersson and Levin 1999; Andersson and Hughes 2010; Melnyk *et al.* 2015).

Today, the term 'cost of resistance' is widely used, but the concept suffers from several ambiguity that can be understood in the light of this short history. First, the concept seems unnecessary to study adaptation in different environments (it would be sufficient to simply consider fitness effects in each environment). It also introduces an asymmetry, which is quite arbitrary, and somewhat misleading (susceptibility too is costly). Second, it conflates effects across traits (pleiotropy) and effects across environments, which can also be misleading. Third, the word 'cost', still reflects an essentialization of mutation/genotypes. The deeply engrained view "one mutation – one fitness effect" was not really challenged by the introduction of the 'cost' idea. It was merely replaced by the idea that one resistance mutation corresponded to two important characteristics: its benefit and its cost.

For these reasons, we think that the concept of cost of resistance presents important shortcomings, to the point, that it is now becoming a problem and hindrance to conceptually clarify the process of adaptation. We now try to explain these issues in more details.

Costs of resistance are not pleiotropic effects

The first problem with the concept of cost is its interpretation in terms of pleiotropic effects of mutations. To be very clear, a simple situation is sketched below where adaptation is represented by the optimization of many traits simultaneously, like in Fisher's model of adaptation or multivariate models of stabilizing selection in quantitative genetics (Lande 1980; Hartl and Taubes 1998; Orr 1998; Martin and Lenormand 2006a), but with two

environments (Martin and Lenormand 2006b, 2015). Figure 1 uses only two traits, which is sufficient to make the argument.

Figure 1. Graph of treated and non-treated environments, with distinct phenotypic requirements (phenotypic optima A and O, respectively) in a two-traits space. Assuming a wild-type positioned in O, the resistance mutation R brings the phenotype closer to A. Relative to the wild type, R is therefore a beneficial mutation. Its cost is usually defined as its fitness in the non-treated environment relative to the wild type, which corresponds to the distance between R and O on the figure. The direction of the two optima (OA axis) defines a phenotypic trait of 'resistance'. Variation of trait(s) orthogonal to this axis may be defined as pleiotropic effects.

Let us label "O" the optimal phenotype in the non-treated environment and "A" the phenotypic optimum in the treated environment. For simplicity, we can assume that fitness declines with the distance from the peak in any given environment. Again, this simplified model could be more complex but the core argument does not require the use of more complex situation or assumptions. We can also assume that the wild type is very close to O, as one would expect from the effect of past selection in absence of drug, and represents a resistance mutation by a vector pointing from O to R, where R is a phenotype closer to A than to O. The mutant is beneficial (relative to the wild type) in the treated environment because the distance AR is smaller than the distance AO. The difference between these two distances scales with the selection coefficient of the resistance mutation in the treated environment. The OA axis, by definition, represents the phenotypic direction of the 'resistance' phenotype. Point P_1 on this axis such that $AR = AP_1$ allows representation of the phenotypic point that would confer the same benefit in the treated environment compared to our resistance mutation R, but that would only alter the phenotype in the exact direction of the optimum A. In other words, the distance OP_1 scales with the selective advantage of the resistance mutation in the treated environment, relative to the wild type. The cost of resistance is shown by the distance

OR, as it is defined as the fitness effect of the resistance mutation in the non-treated environment (again, relative to the wild type). What about the pleiotropic effects? Here, the OA axis represents the phenotypic axis of resistance and therefore, the orthogonal direction represents all other traits (here, there is only one other trait, because we consider only a two dimensional trait space but in with n phenotypic traits, there would be $n - 1$ such traits). Hence, the 'other' pleiotropic effects all project on these axes. Furthermore, note the point P_2, the projection of point R on the axe OA. The vector RP_2 represents the pleiotropic effects of the resistance mutation. Should these effects be totally compensated, the phenotype would be in P_2 and it would indeed enjoy a greater fitness in both the treated and non-treated environments (since $AP_2 < AR=AP_1$ and $OP_2 < OR$, respectively).

This simple geometric argument indicates several things. First, pleiotropic effects and the 'cost of resistance' are two different things – biologically and geometrically – contrary to what is usually considered. Pleiotropic effects will be eventually compensated through the well-known process of "amelioration" when the population reaches the phenotypic optimum after remaining exposed to the treated environment and involves new resistance mutations, compensatory mutations or a mixture of mutations with the two properties (Cohan *et al.* 1994; Schrag *et al.* 1997; Lenski 1998; Levin *et al.* 2000; Schoustra *et al.* 2006; MacLean and Vogwill 2015). Cost evolution will be quite different, and may occur if the population is exposed, at least part time, to the non-treated environment (by evolution of plasticity or inducible response, e.g. Nguyen *et al.* 1989; Foucault *et al.* 2009). Second, the full compensation of these pleiotropic effects does not reduce the cost of resistance to zero. Indeed, biologically speaking, it is likely that acquiring 'resistance' requires changing at least one trait, and thus, this trait becomes suboptimal in the original environment for the resistant mutant. This is irreducible and corresponds to the idea that different phenotypic requirement necessarily involve the occurrence of a fitness trade-off. In any case, there is no reason to believe that the deleterious pleiotropic effects of a resistance mutations at a given drug dose are equal to the cost at dose zero. Finally, it points out that the fitness effects of mutations are not a fixed property of that mutation. It also depends largely on the environments (here the positions of optima). We discuss now this idea in more details.

Resistance mutations do not have a cost

From an experimentalist perspective, defining a non-treated and treated environment is straightforward. You first select a given environment, then you can either add or not add the

drug. With this definition, it is possible to make a very clear and clean experiment demonstrating the effect of the drug, with a control. Yet, the problem is that there is virtually an infinite set of possible environment to start with. Which pair of treated/non-treated environment is relevant? This is difficult to know. It is difficult to represent the complexity of natural conditions in controlled experiment in the laboratory. Even trying to determine which environment corresponds to the environment to which an organism has been adapting to is challenging. For instance, using 'absolute' demographic performance to answer this question may not be reliable. For instance, habitat quality varies and can even obscure the relationship between 'absolute' measures of fitness and environment variables (Gallet *et al.* 2014). For instance, *E. coli* in the gut of humans evolved for a long time at 37°C, yet it grows faster at slightly higher temperatures. Hence, the variation in growth rate (often taken as an absolute measure of fitness) may not be used so easily to infer the environment where adaptation took place. There are many other examples like this.

In fact, it is quite straightforward to see that the 'cost' of a mutation will be different in varying non-treated environments. There is not a cost, but as many costs as there are different (non-treated) environments (Gassmann *et al.* 2009; Vila-Aiub *et al.* 2009; Angst and Hall 2013), which may also be revealed by different compensatory evolution in different environments (Björkman *et al.* 2000). Worse, this cost of resistance may not even actually be a "cost". In other words, the resistance mutation may be favourable in both the treated and non-treated environment, relative to the wild type (see examples in Kassen and Bataillon 2006; Melnyk *et al.* 2015). This can occur for many reasons, but globally will occur often when the wild-type is not well adapted neither to the treated nor to the non-treated environment (Martin and Lenormand 2015). Such a situation is illustrated on Figure 2. Let keep the position of the wild type and resistance mutant in O and R, respectively. Let's also keep the position of the optimal phenotype in the treated environment (in A). Now, lets consider that the optimal phenotype in the non-treated environment is not O, as in Figure 1, but is B. Reporting the point P_3 such that $BP_3 = BR$, we see that the distance to the non-treated optimum is greater for the wild type than for the mutant (*i.e.* BR < BO). The difference between BR and BO actually corresponds to OP_3. When the resistance mutant is favourable in both the treated and non-treated environments (like in this example), the term 'cost' becomes confusing since it implies talking about a 'negative cost'. It would also, in this case, be unclear to interpret costs as deleterious pleiotropic effects (since there is no deleterious effect in the first place). If costs are not pleiotropic effects and not even costly (meaning

deleterious), the terminology and its usual interpretation start obscuring things instead of clarifying them.

Figure 2. Graph of treated and non-treated environments, with distinct phenotypic requirements (phenotypic optima A and B, respectively) in a two-traits space. Relative to a wild-type positioned in O, the resistance mutation R brings the phenotype closer to A. As in Figure 1, R is therefore a beneficial mutation relative to the wild type O. However, its cost is now "negative" in the non-treated environment, as R is also closer to B compared to the wild type O.

Such situation happens when RB ≤ OB i.e. when the wild-type and the resistance mutant are at least equally distant from the non-treated optimal phenotype. Because such situations are quite common (either because we cannot properly reproduce natural conditions experimentally, or because wild type genotypes are not well adapted to their environment), it is perhaps not very surprising that sometimes "cost free" resistance mutation or even "negative cost" are found. All these situations can occur but are unrelated to the pleiotropic effects of the resistance mutation (as defined in the previous section). Importantly, finding an absence of cost or even 'negative costs' do not indicate necessarily that there is no phenotypic trade-off between the treated and non-treated environment (and indeed, on the figure 2, A and B are distinct points, indicating that there is a phenotypic trade-off, despite that no cost is detected). It can just indicate that the wild-type reference is not adapted well to the non-treated environment. At this point, one might argue that all this confusion arises because the mutation R was not a "resistance" mutation to begin with. The mutation R is beneficial in both the treated and non- treated environment, and so, it may be better interpreted as a beneficial mutation to the non-treated environment than as a 'resistance' mutation. For instance, if we are talking of a bacteria mutant showing this property in a laboratory test, we

may want to say that the mutation R corresponds to adaptation to the "laboratory condition", not really to the drug *per se*. But, then, how is a resistance mutation truly defined?

What is a resistance mutation?

If resistance mutations cannot be defined by the fact that they are beneficial in the treated environment, relative to wild type (as we did up to now), then, how do we classify them? In fact, it might be possible to define them more specifically by saying that they are beneficial in the treated environment relatively to wild type, provided that the wild type is perfectly well adapted to the corresponding non-treated environment. In principle, this definition makes sense, as it avoids conflating adaptation to conditions that are shared by both treated and non-treated environments, with adaptation to the drug itself. Yet, with this definition, the mutation R illustrated on Figure 2 would still be a beneficial mutation. R is closer to A (the optimum with the drug) than would a wild type well adapted to the non-treated environment (distance RA < AB). In such a case, the phenotypic direction corresponding to resistance would be the AB axis (and the pleiotropic effects best defined on axes orthogonal to AB). This would be in general clearer and more insightful. With such a definition, it is possible to distinguish mutation R and R' on Figure 3 for instance. Both would be beneficial in both treated and non-treated environment, relative to a wild-type in O, but only R would be beneficial in the treated environment relative to a wild-type in B.

Figure 3. Sketch of treated and non-treated environments, with distinct phenotypic requirements (phenotypic optima A and B, respectively) in a two-traits space. Relative to a wild-type positioned in O, both the resistance mutation R and R' brings the phenotype closer to A. As in Figure 1, they are therefore both beneficial mutations relative to the wild type O. However, if resistance is defined by comparing to a wild-type well adapted to the non-treated

environment (i.e. to a wild type located on B), then only mutation R would be beneficial, and therefore could qualify as being a resistance mutation. With this comparison, costs would also be better defined (i.e. would be always positive).

The problem is that measures of fitness will be made against a wild-type and it may not be straightforward to determine whether this wild type is well adapted to the non-treated environment (but not impossible, as e.g. in cases of experimental evolution in the laboratory where the wild type can be chosen as the type that evolved in the non treated environment for a long time). Without this knowledge, it may be very difficult to classify resistance mutations versus mere beneficial mutations (see e.g. Marcusson *et al.* 2009 for an example of this problem).

There is yet another difficulty lurking in the vast range of possible natural situations. Just as there are many non-treated environments, there may be many treated environments as well. In particular, any drug may be added in different quantities or concentrations to any given environment. Here resides a problem, which is rarely addressed in studies on resistance: do different drug concentrations correspond to different intensity of selection (Milesi *et al.* 2016)? Or do they also correspond to different optimum phenotypes (Harmand *et al.* 2017)?

Figure 4. Sketch of a non-treated environment (phenotypic optimum in B) and two treated environments, with different doses of drug (optima A₁ and A₂). Two mutations are illustrated, with fitness effects compared to a wild type well adapted to the non-treated environment (i.e.

located at B). Using the definition from Figure 3, R_1 is a resistance mutation with respect to environment A_1, but not with respect to environment A_2, and reciprocally for mutation R_2. Nevertheless, both mutations have the same cost.

If different drug doses correspond to different optima (say A_1 and A_2 on Figure 4), it is fairly easy to imagine two mutations R_1 and R_2 that would qualify as resistance mutations, in each of the two environments, but not in the other. On Figure 4, R_1 is a resistance mutation for drug dose 1 (optimum A_1), but not for drug dose 2 (optimum A_2), and reciprocally for R_2. This difference does not arise because these mutations have different "costs". In fact, the two mutations illustrated on Figure 4 have the same cost (the same fitness in non-treated environment B). This situation occurs because phenotypic requirement is different with different doses, such that a phenotypic change can be favourable in one environment (here one dose), but not another. 'Resistance mutations' can differ not only in their benefit and costs at a given drug dose, but also in their fitness effects at other doses. When optima for doses are different, there are many underlying trade-offs, which are not necessarily caused by differing costs in the non-treated environment. Trade-offs among doses are not captured by studying costs. A common view is that mutations conferring strong resistance (i.e. resistance to a high dose) may carry strong costs (Melnyk *et al.* 2015), explaining perhaps why they may not be beneficial at a lower dose. This may well be true, but not necessarily. A mutation favourable at high dose may be deleterious at low dose, irrespectively of its cost. This is the case illustrated on Figure 4. Studying "cost" and "benefit" may give the illusion that all trade-offs are understood. In fact, this is not the case: all trade-offs among doses will be missed. These problems of course mainly arise when different doses correspond to different optima, a situation which requires to be demonstrated empirically. However, the possibility cannot be ignored *a priori* without investigating it. Furthermore, admitting that natural processes have some degree of continuity, it is likely that differing doses end up corresponding to different phenotypic optima. Indeed, it is difficult to conceive that adding a vanishingly small quantity of drug suddenly shifts away phenotypic requirements, and that further increases in doses only change the selection intensity around that shifted phenotypic peak. In any case, representing evolution of resistance as convergence to a phenotypic optimum has received some empirical support (Bataillon *et al.* 2011; Sousa *et al.* 2012; Harmand *et al.* 2017) and may capture well the dynamics of adaptation.

Summary and conclusion

Taking into account the 'cost of resistance' has been a major progress because it is essential to distinguish the fitness effects of resistance mutations in treated versus non-treated environments. However, this cost is also very expensive conceptually as it is associated with too many simplifications, to the point that it may even be misleading. Its interpretation in terms of pleiotropic effects is very unclear. First, pleiotropic effects may be better defined as effects projected on phenotypic axes orthogonal to resistance phenotype, than in terms of fitness effect in the non-treated environment (see Figure 1). Second, there are as many costs as there are non-treated environments. The idea that there is a cost associated with a resistance mutation is an extreme and naïve simplification. Third, costs of resistance are ill defined when several precautions are not taken. For instance, failing to measure costs relative to a well-adapted wild type to the non-treated environment can lead to absurd notions, such as 'negative costs' that serve no conceptual clarification (Figure 2). In practice, these precautions can be very difficult to meet, as the degree of adaptation of the wild type to the non-treated environment will be in general difficult to assess. Fourth, the concept of cost leads to an oversimplified and often erroneous view of trade-offs across environments. Finding costs equal to zero (*i*) cannot be used to say that there is no trade-offs between treated and non-treated environments; (*ii*) cannot be interpreted by saying that pleiotropic effects were compensated; (*iii*) completely misses the possibility that fitness trade-off may occur among different doses. Finding that resistance mutations are not favourable at all doses cannot be attributed to their cost. For instance, high resistance mutations may not be beneficial at low doses, not because they have a high cost, but simply because they do not match phenotypic requirements that are optimal at low dose. Overall, it may be safer in most cases to simply discuss and measure the fitness effects of mutation in different environments and to carefully consider the role of the reference genotype/phenotype when interpreting relative fitnesses.

All these points hold for many other situations of adaptation besides resistance, but this is perhaps where the vocabulary and the conceptual issues are the most acute and widespread. Differences among selective conditions and the occurrence of pleiotropy are both important ideas in evolutionary ecology. However, they cannot be solely summarized by assigning resistance mutations a 'benefit' and 'a cost' and essentialize their properties. There is a problem with a reduction of evolutionary thinking to a cost-benefit thinking, with "fitness" as a universal currency, valid regardless of ecological conditions. Although fitness is a general concept and a universal currency for adaptation, different conditions entail different fitnesses

and possibly different phenotypic requirements (different adaptations). The vocabulary that we use should not oversimplify these ideas.

Acknowledgments

We thank G. Martin for insightful discussions.

References

Abedi Z. H., Brown A. W. A., 1960 Development and reversion of DDT resistance in *Aedes aegypti*. Can. J. Genet. Cytol. 2: 252–261.

Anderson R. M., May R. M., 1982 Coevolution of hosts and parasites. Parasitology 85: 411–426.

Andersson D. I., Levin B. R., 1999 The biological cost of antibiotic resistance. Curr. Opin. Microbiol. 2: 489–93.

Andersson D. I., Hughes D., 2010 Antibiotic resistance and its cost: is it possible to reverse resistance? Nat. Rev. Microbiol. 8: 260–271.

Angst D. C., Hall A. R., 2013 The cost of antibiotic resistance depends on evolutionary history in *Escherichia coli*. BMC Evol. Biol. 13.

Antonovics J., 1977 Metal tolerance in plants: perfecting an evolutionary paradigm. In: Institute for Environmental Studies U of T (Ed.), *International Conference on Heavy Metals in the Environment, Toronto, Ontario, Canada*, Toronto, pp. 169–186.

Bataillon T., Zhang T., Kassen R., 2011 Cost of adaptation and fitness effects of beneficial mutations in *Pseudomonas fluorescens*. Genetics 189: 939–49.

Bell G., 2008 *Selection*. Oxford Univ. Press, Oxford.

Bergelson J., Purrington C. B., 1996 Surveying Patterns in the Cost of Resistance in Plants. Am. Nat. 148: 536–558.

Björkman J., Nagaev I., Berg O. G., Hughes D., Andersson D. I., 2000 Effects of environment on compensatory mutations to ameliorate costs of antibiotic resistance. Science (80-.). 287: 1479–1482.

Bourguet D., Prout M., Raymond M., 1996 Dominance of insecticide resistance presents a plastic response. Genetics 143: 407–416.

Bradshaw A. D., 1984 Adaptation of plants to soils containing toxic metals - a test for conceit. In: Evered D, Collins GM (Eds.), *Origins and development of adaptation*, Pitman Books, London, p. 273.

Carrière Y., Deland J., Roff D., Vincent C., 1994 Life-history costs associated with the evolution of insecticide resistance. Proc. R. Soc. London B Biol. Sci. 258.

Chevin L. M., Hospital F., 2008 Selective sweep at a quantitative trait locus in the presence of background genetic variation. Genetics 180: 1645–1660.

Cohan F. M., King E. C., Zawadzki P., 1994 Amelioration of the deleterious pleiotropic effects of an adaptive mutation in *Bacillus subtilis*. Evolution 48: 81–95.

Curtis C. F., Cook L. M., Wood R. J., 1978 Selection for and against insecticide resistance and possible methods of inhibiting the evolution of resistance in mosquitoes. Ecol. Entomol. 3: 228–273.

Davies A. G., Game A. Y., Chen Z., Williams T. J., Goodall S., *et al.*, 1996 Scalloped wings is the Lucilia cuprina Notch homologue and a candidate for the Modifier of fitness and asymmetry of diazinon resistance. Genetics 143: 1321–1337.

Depledge M., 2011 Reduce drug waste in the environment. Nature 478: 36.

Dobzhansky T., 1951 *Genetics and the origin of species*. Columbia university press, New York.

Eyre-Walker A., Keightley P. D., 2007 The distribution of fitness effects of new mutations. Nat. Rev. Genet. 8: 610–618.

Foucault M. L., Courvalin P., Grillot-Courvalin C., 2009 Fitness cost of VanA-type vancomycin resistance in methicillin-resistant *Staphylococcus aureus*. Antimicrob. Agents Chemother. 53: 2354–2359.

Gallet R., Latour Y., Hughes B. S. B. S., Lenormand T., 2014 The dynamics of niche evolution upon abrupt environmental change. Evolution 68: 1257–1269.

Gassmann A. J., Carrière Y., Tabashnik B. E., 2009 Fitness Costs of Insect Resistance to *Bacillus thuringiensis*. Annu. Rev. Entomol. 54: 147–163.

Georghiou G. P., Taylor C. E., 1977 Genetic and biological influences in the evolution of insecticide resistance. J. Econ. Entomol. 70: 319–323.

Gillespie J. H., 1975 Natural selection for resistance to epidemics. Ecology 56: 493–495.

Guillemaud T., Lenormand T., Bourguet D., Chevillon C., Pasteur N., *et al.*, 1998 Evolution of resistance in *Culex pipiens*: allele replacement and changing environment. Evolution 52: 443–453.

Haldane J. B. S., 1948 The theory of a cline. J. Genet. 48: 277–284.

Harmand N., Gallet R., Jabbour-Zahab R., Martin G., Lenormand T., 2017 Fisher's geometrical model and the mutational patterns of antibiotic resistance across dose gradients. Evolution 71: 23–37.

Hartl D. L., Taubes C. H., 1998 Towards a theory of evolutionary adaptation. Genetica 103: 525–533.

Hickey D. A., McNelly T., 1975 Competition Between Metal Tolerant and Normal Plant Populations ; A Field Experiment on Normal Soil. Evolution 29: 458–464.

Jacobs B. F., Duesing J. H., Antonovics J., Patterson D. T., 1988 Growth performance of triazine-resistant and -susceptible biotypes of *Solanum nigrum* over a range of temperatures. Can. J. Bot. 66: 847–850.

Jain S. K., Bradshaw A. D., 1966 Evolutionnary divergence among adjacent plant populations. I. The evidence and its theoretical analysis. Heredity 21: 407–441.

Kassen R., Bataillon T., 2006 Distribution of fitness effects among beneficial mutations before selection in experimental populations of bacteria. Nat. Genet. 38: 484–488.

Labbé P., Sidos N., Raymond M., Lenormand T., 2009 Resistance Gene Replacement in the mosquito *Culex pipiens*: fitness estimation from long-term cline series. Genetics 182: 303–12.

Lande R., 1980 The Genetic Covariance between Characters Maintained by Pleiotropic Mutations. Genetics 94: 203–215.

Lande R., 1983 The response to selection on major and minor mutations affecting a metrical trait. Heredity 50: 47–65.

Lenormand T., Guillemaud T., Bourguet D., Raymond M., 1998 Evaluating gene flow using selected markers: a case study. Genetics 149: 1383–1392.

Lenormand T., Raymond M., 1998 Resistance management: the stable zone strategy. Proc. R. Soc. London B 265: 1985–1990.

Lenormand T., Bourguet D., Guillemaud T., Raymond M., 1999 Tracking the evolution of

insecticide resistance in the mosquito *Culex pipiens*. Nature 400: 861–864.

Lenormand T., Raymond M., 2000 Analysis of clines with variable selection and variable migration. Am. Nat. 155: 70–82.

Lenormand T., 2002 Gene flow and the limits to natural selection. Trends Ecol. Evol. 17: 183–189.

Lenski R. E., 1988 Experimental studies of pleiotropy and epistasis in *Escherichia coli*. I. Variation in competitive fitness among mutant resistant to virus T4. Evolution 42: 425–432.

Lenski R. E., 1998 Bacterial evolution and the cost of antibiotic resistance. Int. Microbiol. 1: 265–70.

Leonard K. J., Czochor R. J., 1980 Theory of genetic interactions among populations of plants and their pathogens. Annu. Rev. Phytopathol. 18: 237–258.

Levene H., 1953 Genetic equilibrium when more than one ecological niche is available. Am. Nat. 87: 331–333.

Levin B. R., Perrot V., Walker N., 2000 Compensatory mutations, antibiotic resistance and the population genetics of adaptive evolution in bacteria. Genetics 154: 985–997.

Levison M. E., Levison J. H., 2009 Pharmacokinetics and pharmacodynamics of antibacterial agents. Infect Dis Clin North Am. 23: 791–797.

MacLean R. C., Vogwill T., 2015 Limits to compensatory adaptation and the persistence of antibiotic resistance in pathogenic bacteria. Evol. Med. Public Heal. 2015: 4–12.

Manna F., Martin G., Lenormand T., 2011 Fitness landscapes: An alternative theory for the dominance of mutation. Genetics 189.

Marcusson L. L., Frimodt-Møller N., Hughes D., 2009 Interplay in the selection of fluoroquinolone resistance and bacterial fitness. PLoS Pathog. 5.

Martin G., Lenormand T., 2006a A general multivariate extension of Fisher's geometrical model and the distribution of mutation fitness effects across species. Evolution 60: 893–907.

Martin G., Lenormand T., 2006b The fitness effect of mutations across environments: a survey in the light of fitness landscape models. Evolution 60: 2413–2427.

Martin G., Lenormand T., 2015 The fitness effect of mutations across environments: Fisher's

geometrical model with multiple optima. Evolution 69: 1433–1447.

May R. M., Dobson A. P., 1986 Population Dynamics and the Rate of Evolution of Pesticide Resistance. In: *Pesticide Resistance: Strategies and Tactics for Management*, The National Academies Press, Washington, DC, pp. 170–193.

McKenzie J. A., Whitten M. J., Adena M. A., 1982 The effect of genetic background on the fitness of diazinon resistance genotypes of the australian sheep blowfly, *Lucilia cuprina*. Heredity 49: 1–9.

Melnyk A. H., Wong A., Kassen R., 2015 The fitness costs of antibiotic resistance mutations. Evol. Appl. 8: 273–283.

Milesi P., Lenormand T., Lagneau C., Weill M., Labbé P., 2016 Relating fitness to long-term environmental variations in natura. Mol. Ecol. 25: 5483–5499.

Nagylaki T., 1975 Conditions for the existence of clines. Genetics 80: 595–615.

Neve P., Powles S., 2005 High survival frequencies at low herbicide use rates in populations of *Lolium rigidum* result in rapid evolution of herbicide resistance. Heredity 95: 485–492.

Nguyen T. N., Phan Q. G., Duong L. P., Bertrand K. P., Lenski R. E., 1989 Effects of carriage and expression of the Tn10 tetracycline-resistance operon on the fitness of *Escherichia coli* K12. Mol. Biol. Evol. 6: 213–225.

Orr H. A., 1998 The population genetics of adaptation: The distribution of factors fixed during adaptive evolution. Evolution 52: 935–949.

Orr H. A., 2005 The genetic theory of adaptation: A brief history. Nat. Rev. Genet. 6: 119–127.

Schoustra S. E., Debets a J. M., Slakhorst M., Hoekstra R. F., 2006 Reducing the cost of resistance; experimental evolution in the filamentous fungus *Aspergillus nidulans*. J. Evol. Biol. 19: 1115–27.

Schrag S. J., Perrot V., Levin B. R., 1997 Adaptation to the fitness costs of antibiotic resistance in *Escherichia coli*. Proc. Biol. Sci. 264: 1287–91.

Simms E. L., Rausher M. D., 1987 Costs and Benefits of Plant Resistance to Herbivory. Am. Nat. 130: 570–581.

Smith P., Townsend M. G., Smith R. H., 1991 A Cost of Resistance in the Brown Rat?

Reduced Growth Rate in Warfarin-Resistant Lines. Funct. Ecol. 5: 441–447.

Sousa A., Magalhães S., Gordo I., Sousa A., Magalha S., 2012 Cost of antibiotic resistance and the geometry of adaptation. Mol. Biol. Evol. 29: 1417–28.

Suckling D. M., Khoo J. G. I., 1993 Cline in frequency of azinphosmethyl resistance in light brown apple moth (Lepidoptera: Tortricidae). J. Econ. Entomol. 86: 1308–1316.

Thiele-Bruhn S., 2003 Pharmaceutical antibiotic compounds in soils - a review. J. plant Nutr. soil Sci. 166: 145–167.

Ventola C. L., 2015 The antibiotic resistance crisis: part 1: causes and threats. P T 40: 277–83.

Vila-Aiub M. M., Neve P., Powles S. B., 2009 Fitness costs associated with evolved herbicide resistance alleles in plants. New Phytol. 184: 751–767.

Wilson J. B., 1988 The cost of heavy-metal tolerance: an example. Evolution 42: 408–413.

Windle P. N., Franz E. H., 1979 The Effects of Insect Parasitism on Plant Competition : Greenbugs and Barley. Ecology 60: 521–529.

World Health Organization, 2014 *Antimicrobial Resistance Global Report on Surveillance*.