Perspective

Cost of resistance: an unreasonably expensive concept

Thomas Lenormand[1,2], Noémie Harmand[1], Romain Gallet[3]

[1] UMR 5175 CEFE, CNRS - Université Montpellier - Université P. Valéry - EPHE, Montpellier Cedex 5, France

[2] Radcliffe Institute for advanced study, Harvard University, Byerly Hall, 8 Garden Street, Cambridge, MA

02138

[3] INRA - UMR BGPI, Cirad TA A-54/K Campus International de Baillarguet 34398 Montpellier Cedex 5 France

Abstract

The cost of resistance, or the fitness effect of resistance mutation in absence of the drug,
is a very widepsread concept in evolutionary genetics and beyond. It has represented an
important addition to the simplistic view that resistance mutations should solely be
considered as beneficial mutations. Yet, this concept also entails a series of serious
difficulties in its definition, interpretation and current usage. In many cases, it may be
simpler, clearer, and more insightful to study, measure and analyze the fitness effects of
mutations across environments and to better distinguish those effects from 'pleiotropic
effects' of those mutations.

Introduction

The study of resistance to antibiotics, insecticides, acaricides, fungicides, herbicides,
chemotherapy drugs etc. is, for obvious reasons, a very active field of research. We are in the
middle of a "crisis," which has important consequences for public health and agriculture (e.g.
World Health Organization 2014; Ventola 2015). It is yet unclear whether we will be able to
deal with "superbugs", "superweeds" and other "supermicrobes" in the near future. Studies
have focused intensely on the genetic, cellular, and biochemical mechanisms responsible for
resistance, but also on the fitness effect of those mutations.

Resistance mutations as beneficial mutations

Resistance evolution is a particular case of the more general situation of adaptation to new environmental conditions. Processes of adaptation have been intensely studied from Darwin's time, since refined by powerful population genetic concepts that have been put forward in the modern synthesis (Orr 2005). In brief, resistance mutation are beneficial mutations in treated environments. They spread because they confer an obvious fitness advantage in presence of the drug (the antibiotic, insecticide, acaricide, fungicide etc.). By definition, a resistance mutation allows for survival while the susceptible wild-type simply dies when exposed to the drug. Hence, classically, the fitness benefit of a resistance mutation (relative to a susceptible one) depends on the fraction of the population exposed to the drug (since it determines the fraction of surviving susceptible genotypes). This is a black and white outcome, and it seems that there is little to understand beyond this obvious reasoning. Yet, in natural / real conditions (*i.e.* unlike in ecotoxicological tests performed in the laboratory), it is not very clear which dose of the drug is relevant. Concentrations of drugs vary at different temporal and geographical scales, within bodies (Levison and Levison 2009), within microhabitats, within regions *etc.* (Thiele-Bruhn 2003; Depledge 2011). Notwithstanding, it is often assumed that a resistance mutation is associated with a selection coefficient measuring the rate at which it is expected to change in frequency in populations, like any other beneficial mutation. This selective advantage is not easy to estimate in the field, but is often thought to represent an inherent property of the mutation itself. However, this advantage must depend in some ways on the exposure to the drug, which is an environmental variable, and not a property of the mutation.

The context dependence of fitness effects

This point leads to a very simple and obvious idea: the selective effects of mutations depend on ecological conditions. This conclusion is somewhat trivial (Bell 2008), but it contradicts the naïve view that tends to essentialize the properties of mutations or genotypes (*i.e.* that characterize mutationnal properties as intrinsic). There is no such thing as a single selective effect of a mutation or a genotype. A mutation is not inherently beneficial, deleterious, or neutral. Rather, it depends on the ecological conditions, the genetic background, and on other alleles considered in comparison. Even the 'dominance' of mutations in diploids can be highly context and trait dependent (Bourguet *et al.* 1996; Manna *et al.* 2011). It is important to reiterate this point for several reasons. The first is that it tends to be simply ignored, even if

this context dependence is never directly challenged. For instance, in molecular evolution, it
is customary to assign a mutation, without further specifications, in three categories:
deleterious, neutral or beneficial (Eyre-Walker and Keightley 2007). Secondly, it makes little
sense to study the selective effects of mutations while neglecting the diversity of real
ecological conditions and the diversity of genetic backgrounds. Of course, it is always
possible to think in terms of average, and this is the usual view when dealing with the
diversity of genetic backgrounds in sexual populations (where recombination ensures that
mutations "experience" effectively the range of possible backgrounds, Lande 1983; Chevin
and Hospital 2008). Yet, for ecological variation, taking such an average may be very
misleading. For example, a resistance mutation can spread somewhere (in a treated
environment), but not elsewhere (in a non-treated environment), generating a situation of
polymorphism (Jain and Bradshaw 1966; Suckling and Khoo 1993; Carrière *et al.* 1994;
Guillemaud *et al.* 1998; Lenormand *et al.* 1998, 1999; Lenormand and Raymond 2000; Neve
and Powles 2005; Labbé *et al.* 2009). Differences of effects across environments matters, not
only their average, provided that the spatial scale of dose variation is larger than the scale of
dispersal (Lenormand 2002) or that temporal fluctuations occur at scales exceeding generation
time (e.g. Cvijović *et al.* 2015).

The cost of resistance

In presence of such heterogeneity with both treated and non-treated environments, the concept
of 'cost of resistance' becomes important. This cost is defined as the selection coefficient of
resistance mutations in absence of treatments (or similarly in absence of predator, parasite or
pathogens when considering resistance in the context of biotic interactions). The idea of a
"lower adaptive value" of resistant genotypes in the absence of treatment can be traced back
quite far (e.g. in Dobzhansky 1951). The term "reproductive disadvantage" or no specific
term is used in this context in the 60s and 70s, in empirical or theoretical papers (Abedi and
Brown 1960; Gillespie 1975; Hickey and McNelly 1975; Antonovics 1977; Georghiou and
Taylor 1977; Curtis *et al.* 1978). The term 'cost of resistance' is relatively recent in
comparison and started to be widespread only in the 80s, especially in the context of plant
resistance to herbivores (Windle and Franz 1979; Leonard and Czochor 1980; Simms and
Rausher 1987) or bacteria resistance to phage (Lenski 1988). It is in particular used in the
influential paper of Anderson and May (1982) on coevolution and resistance to pathogens.
Curiously, when dealing with resistance evolution in an abiotic context (*i.e.* to pesticides or

antibiotics), the term 'cost of resistance' still refers in the 80s to the economic, not the

evolutionary, cost of resistance (*i.e.* the extra monetary cost due to increasing pesticide dosage

consecutive to resistance evolution), even in papers by May, referring to evolutionary cost as

'back selection' (May and Dobson 1986). The first mention of an evolutionary cost of

resistance in an abiotic context seem to be in Jacobs *et al.* (1988) about herbicide resistance,

although the term "cost of tolerance" was used in studies of heavy metal tolerance in plants a

bit earlier (Bradshaw 1984; Wilson 1988). The cost of resistance was then intensely

investigated in the 90s (Bergelson and Purrington 1996; Andersson and Levin 1999 for

reviews), with several methodological improvements (e.g. separating costs from effect of

linked variation). By the end of the 90s, most pesticide resistance management models

included the cost of resistance, *i.e.* the fact that resistance mutations could be selected against

in non-treated environments (Lenormand and Raymond 1998). As this short historical

overview shows, the concept of cost of resistance is relatively recent and was not used before

the 80s. In particular, all the paper on local adaptation, clines and all the field of ecological

genetics developed before the 80s without the need to refer to this concept. For instance, in

classical models such as Levene model (Levene 1953) or cline models (Haldane 1948;

Nagylaki 1975; Endler 1977), it was sufficient to talk about the selective effects of alleles in

different environments. Why was the concept of cost of resistance introduced? At first sight,

the concept seems asymmetrical and not very useful. For instance, nobody talks of the "cost

of susceptibility" in treated environments, although it would be equally legitimate.

A likely reason is that the concept was helpful to bring attention to the fact that a mutation

could be both beneficial or deleterious depending on circumstances, something well known in

ecological genetics but somewhat ignored in resistance studies. It helped introduce some

ecology in the understanding of the fitness effect of resistance mutations. This can have

important consequences as the cost of resistance is a powerful force that can keep resistance

in check (see e.g. Curtis *et al.* 1978 for an early model). Considering cost is also very

important to predict the dynamics of resistance mutations in heterogeneous (treated and non-

treated) environments.

A second reason is that the concept of cost was tightly associated to the notion of trade-off

among traits, an idea borrowed from life history theory. When resistance is viewed as a trait,

or a "defense" function, it is natural to consider that it may trade-off with other organismal

traits and functions (e.g. in terms of resource). In other words, resistance mutations should

influence many traits, *i.e.* have pleiotropic effects. Besides resistance, the variation of all these

traits is likely to be deleterious, and therefore represent a 'cost', since these traits were

previously optimized by natural selection. This connection with life history theory is entirely

explicit in the first papers mentioning the concept (Simms and Rausher 1987; Lenski 1988;

Wilson 1988; Smith *et al.* 1991; Bergelson and Purrington 1996) and led to idea that the cost

of resistance was caused by the pleiotropic effects of resistance mutations. Although the idea

of trade-off among traits was initially present in this interpretation, it is often forgotten today:

resistance mutations are simply viewed as 'pleiotropic'. Naturally, only considering

pleiotropy, it seems very natural to think that the cost can evolve to be reduced, or even

eliminated (which is also directly suggested by the word 'cost' itself). For instance, one could

consider that compensatory evolution should attenuate these unwanted pleiotropic effects and

restore optimal values for all traits, eliminating the cost. At first sight, the best proof for this

reasoning is that cost-free resistance mutations are sometimes found along with the existence

of modifier loci mitigating or even eliminating costs (McKenzie *et al.* 1982; Davies *et al.*

1996; Lenski 1998; Andersson and Levin 1999; Andersson and Hughes 2010; Melnyk *et al.*

2015).

Today, the term 'cost of resistance' is widely used, but the concept suffers from several

ambiguity that can be understood in the light of this short history. First, the concept seems

unnecessary to study adaptation in different environments (it would be sufficient to simply

consider fitness effects in each environment). It also introduces an asymmetry, which is quite

arbitrary, and somewhat misleading (susceptibility too is costly). Second, it conflates effects

across traits (pleiotropy) and effects across environments, which can also be misleading.

Third, the word 'cost', still reflects an essentialization of mutation/genotypes. The deeply

engrained view "one mutation – one fitness effect" was not really challenged by the

introduction of the 'cost' idea. It was merely replaced by the idea that one resistance mutation

corresponded to two important characteristics: its benefit and its cost.

For these reasons, we think that the concept of cost of resistance presents important

shortcomings, to the point, that it is now becoming a problem and hindrance to conceptually

clarify the process of adaptation. We now try to explain these issues in more details.

Costs of resistance are not equivalent to pleiotropic effects

The first problem with the concept of cost is its interpretation in terms of pleiotropic effects of

mutations. To be very clear, a simple situation is sketched below where adaptation is

represented by the optimization of many traits simultaneously, like in Fisher's model of

adaptation or multivariate models of stabilizing selection in quantitative genetics (Lande
1980; Hartl and Taubes 1998; Orr 1998; Martin and Lenormand 2006a), but with two
environments (Martin and Lenormand 2006b, 2015). Figure 1 uses only two traits, which is
sufficient to make the argument and discuss the issue of pleiotropy. It is straightforward to
relate this type of fitness landscape model to the more traditional one-dimensional 'dose-
response' models (see Box 1). Representing evolution of resistance as convergence to a
phenotypic optimum has received some empirical support (Bataillon *et al.* 2011; Sousa *et al.*
2012; Harmand *et al.* 2017, 2018) and may capture well the dynamics of adaptation.

Figure 1. Graph of treated and non-treated environments, with distinct phenotypic requirements
(phenotypic optima A and O, respectively) in a two-trait space. Assuming a wild-type
positioned in O, the resistance mutation R brings the phenotype closer to A. Relative to the
wild type, R is therefore a beneficial mutation in the treated environment. Its cost is usually
defined as its fitness in the non-treated environment relative to the wild type, which depends on
the distance between R and O on the figure. Note that the cost (and all fitness measures on this
figure, and similar figure below) depends on Euclidian distances in phenotypic space, and a
mapping function converting this distance to fitness (*i.e.* the cost is not distance OR, but a
monotonic function of this distance). The mapping is left implicit on the figure, but can be
thought as a third orthogonal axis representing fitness for each trait 1 – trait 2 combination,
which defines a "fitness landscape". The coloured inset figure represents such a fitness
mapping in 3D. Fitness values, when projected on the phenotypic space correspond to
isofitness curves (like altitude on a geographic map is indicated by contour lines). For instance
all phenotypes on the light grey circles have the same fitness than mutation R in the treated
environment (optimum A). The direction of the two optima (OA axis) defines a phenotypic
trait of 'resistance'. Variation of trait(s) orthogonal to this axis may be defined as pleiotropic

effects. Point P_1 on OA axis is such that $AR = AP_1$. It represents the phenotypic point that

would confer the same fitness in the treated environment compared to the mutation R, but that

would only alter the phenotype in the exact direction of the optimum A. Point P_2 is the

orthogonal projection of R on OA axis. It represents the phenotypic point that would be

reached or if all the pleiotropic effects of the mutation R were compensated (e.g. by subsequent

compensatory mutations).

Let us label "O" the optimal phenotype in the non-treated environment and "A" the

phenotypic optimum in the treated environment. For simplicity, we can assume that fitness

monotonously declines with the (Euclidian) distance from the peak in any given environment.

Again, this simplified model could be more specific (with a particular mapping of distance to

fitness) or complex but the core argument does not require the use of more complex situation

or assumptions. We can also assume that the wild type is very close to O, as one would expect

from the effect of past selection in absence of drug, and represents a resistance mutation by a

vector pointing from O to R, where R is a phenotype closer to A than to O. The mutant is

beneficial (relative to the wild type) in the treated environment because the distance AR is

smaller than the distance AO. The difference between these two distances scales with the

selection coefficient of the resistance mutation in the treated environment. The OA axis, by

definition, represents the phenotypic direction of the 'resistance' phenotype. Point P_1 on this

axis such that $AR = AP_1$ allows representation of the phenotypic point that would confer the

same benefit in the treated environment compared to our resistance mutation R, but that

would only alter the phenotype in the exact direction of the optimum A. In other words, the

distance OP_1 scales with the selective advantage of the resistance mutation in the treated

environment, relative to the wild type. The cost of resistance depends on the distance OR, as it

is defined as the fitness effect of the resistance mutation in the non-treated environment

(again, relative to the wild type). What about the pleiotropic effects? Here, the OA axis

represents the phenotypic axis of resistance and therefore, the orthogonal direction represents

all other traits (here, there is only one other trait, because we consider only a two dimensional

trait space but in with n phenotypic traits, there would be $n - 1$ such traits). Hence, the 'other'

pleiotropic effects all project on these axes orthogonal to OA. Furthermore, note the point P_2,

the projection of point R on the OA axis. The vector RP_2 represents the pleiotropic effects of

the resistance mutation. Should these effects be totally compensated, the phenotype would be

in P_2 and it would indeed enjoy a greater fitness in both the treated and non-treated

environments (since $AP_2 < AR = AP_1$ and $OP_2 < OR$, respectively).

This simple geometric argument indicates several things. First, pleiotropic effects and the
'cost of resistance' are two different things – biologically and geometrically – contrary to
what is usually considered. Pleiotropic effects will be eventually compensated through the
well-known process of "amelioration" when the population reaches the phenotypic optimum
after remaining exposed to the treated environment and involves new resistance mutations,
compensatory mutations or a mixture of mutations with the two properties (Cohan *et al.* 1994;
Schrag *et al.* 1997; Lenski 1998; Levin *et al.* 2000; Schoustra *et al.* 2006; MacLean and
Vogwill 2015). Cost evolution will be quite different, and may occur if the population is
exposed, at least part time, to the non-treated environment (by evolution of plasticity or
inducible response, e.g. Nguyen *et al.* 1989; Foucault *et al.* 2009). Second, the full
compensation of these pleiotropic effects does not reduce the cost of resistance to zero.
Indeed, biologically speaking, it is likely that acquiring 'resistance' requires changing at least
one trait, and thus, this trait becomes suboptimal in the original environment for the resistant
mutant. This is irreducible and corresponds to the idea that different phenotypic requirements
necessarily involve the occurrence of a fitness trade-off. In any case, there is no reason to
believe that the deleterious pleiotropic effects of a resistance mutation at a given drug dose are
equal to the cost of resistance at dose zero. Finally, it points out that the fitness effects of
mutations are not a fixed property of that mutation. It also depends largely on the
environments (here the positions of optima). We discuss now this idea in more details.

Box 1. Correspondence between a fitness landscape model and a drug-response model. On the
left panel, a fitness landscape model is illustrated as in Figure 1 to 4, in a two-traits phenotypic
space. Four environments are represented with increasing concentration of a drug (with optima

0, 1, 2, 3). Environment with optimum 0 (in green) represents the environment without drug, while environment 1, 2, 3 represent increasing concentrations of the drug. In this model, fitness depends on the Euclidian distance to the optimum, and a mapping function (see inset of Fig. 1). It is possible to set a threshold value for absolute fitness below which a phenotype cannot persist / grow. This threshold is indicated by circles (the colour of the circle corresponds to the different environments). In many cases, resistant mutants can have a positive growth rate in absence of drug (while the reverse is not true: susceptible phenotype do not grow in presence of the drug). Hence the threshold contours will be often nested (but it is possible to imagine cases where this is not the case). In this representation, it is easy to see that an absolute fitness criterion (= being within the threshold contour) is not synonymous with adaptation (= being close to the optimum). The dot S represents the position of a susceptible phenotype, while R_1 and R_2 represent two resistance mutants. The right panel illustrates the 'dose response' curves relating, dose to absolute fitness, for each phenotype (S, R_1, R_2). For instance, R_1 is within the threshold contour of dose 1, but not of dose 2 and 3. Hence, its dose-response is zero for doses above dose 1. Note also that R_1 is further apart from optimum 0, compared to S. Hence, its absolute fitness is lower than that of S at dose 0. This correspondence shows that it is entirely possible (and straightforward) to relate fitness modes to more traditional one-dimensional dose-response models. Furthermore, since LD_{50}, IC_{50}, MIC or other ecotoxicological measures can be defined using dose response, they can also be defined in the fitness landscape model. Note however that these measures have been rightly criticized as being only partial fitness summaries (Regoes et al. 2004; Sampah et al. 2011; Wen et al. 2016). They are also often obtained in absence of competition, or often concern only a particular life stage. Note also that absolute measures of fitness are often more appropriate (than relative fitness) when dealing with the demography of the treated species (Day *et al.* 2015). However, relative fitness is in general more relevant to study environment specialization, where the "cost of resistance" matters.

Resistance mutations do not have *"a cost"*

From an experimentalist perspective, defining a non-treated and treated environment is straightforward. You first select a given environment, then you can either add the drug or not. With this definition, it is possible to make a very clear and clean experiment demonstrating the effect of the drug, with a control. Yet, the problem is that there is virtually an infinite set of possible environments to start with. Which pair of treated/non-treated environments is relevant? This is difficult to know. It is difficult to represent the complexity of natural conditions in controlled experiment in the laboratory. Even trying to determine which environment corresponds to the environment to which an organism has been adapting to is challenging. For instance, using 'absolute' demographic performance to answer this question

may not be reliable. For instance, habitat quality varies and can even obscure the relationship

between 'absolute' measures of fitness and environment variables (Gallet *et al.* 2014). For

instance, *E. coli* in the gut of humans evolved for a long time at 37°C, yet it grows faster at

slightly higher temperatures in laboratory conditions (Gonthier *et al.* 2001). Hence, the

variation in growth rate (often taken as an absolute measure of fitness) may not be used so

easily to infer the environment where adaptation took place.

In fact, it is quite straightforward to see that the cost of a mutation will be different in varying

non-treated environments. There is not "a" cost, but as many costs as there are different (non-

treated) environments (Gassmann *et al.* 2009; Vila-Aiub *et al.* 2009; Angst and Hall 2013;

Gifford *et al.* 2016), which may also be revealed by different compensatory evolution in

different environments (Björkman *et al.* 2000). Worse, this cost of resistance may not even

actually be positive, challenging the usage of the word "cost" itself (in the usual economic

sense, a cost is necessarily positive, or it would not be a "cost" in the first place). In other

words, the resistance mutation may be favourable in both the treated and non-treated

environment, relative to the wild type (see examples in Kassen and Bataillon 2006; Melnyk *et*

al. 2015). This can occur for many reasons, but globally will occur often when the wild-type

is not well adapted neither to the treated nor to the non-treated environment (Martin and

Lenormand 2015). Such a situation is illustrated on Figure 2. Let keep the position of the wild

type and resistance mutant in O and R, respectively. Let's also keep the position of the

optimal phenotype in the treated environment (in A). Now, lets consider that the optimal

phenotype in the non-treated environment is not O, as in Figure 1, but is B. Reporting the

point P_3 such that $BP_3 = BR$, we see that the distance to the non-treated optimum is greater for

the wild type than for the mutant (*i.e.* BR < BO). The difference between BR and BO actually

corresponds to OP_3. When the resistance mutant is favourable in both the treated and non-

treated environments (like in this example), the term 'cost' becomes confusing since it implies

talking about a 'negative cost'. It would also, in this case, be unclear to interpret costs as

deleterious pleiotropic effects (since there is no deleterious effect in the first place). If costs

are not pleiotropic effects and not even costly (meaning deleterious), the terminology and its

usual interpretation start obscuring things instead of clarifying them.

Figure 2. Graph of treated and non-treated environments, with distinct phenotypic requirements (phenotypic optima A and B, respectively) in a two-trait space. Relative to a wild-type positioned in O, the resistance mutation R brings the phenotype closer to A. As in Figure 1, R is therefore a beneficial mutation relative to the wild type O in the treated environment (with optimum A). However, its cost is now "negative" in the non-treated environment, as R is also closer to B compared to the wild type O. The point P3 is such that $BP_3 = BR$. The distance to the non-treated optimum B is greater for the wild type O than for the mutant R (*i.e.* BR < BO). The distance difference between BR and BO corresponds to OP_3. As in Figure 1, all fitness measures depend on the phenotypic distances illustrated and a mapping that could be represented as a third orthogonal axis representing fitness. This fitness axis is not shown. All phenotypes on the light grey circles have the same fitness than mutation R in the treated environment (optimum A, large circle) and non-treated environment (optimum B, small circle).

Such situation happens when RB ≤ OB *i.e.* when the wild-type and the resistance mutant are at least equally distant from the non-treated optimal phenotype. Because such situations are quite common (either because we cannot properly reproduce natural conditions experimentally, or because wild type genotypes are not well adapted to their environment), it is perhaps not very surprising that sometimes "cost free" resistance mutation or even "negative cost" are found. All these situations can occur but are unrelated to the pleiotropic effects of the resistance mutation (as defined in the previous section). Importantly, finding an absence of cost or even 'negative costs' do not indicate necessarily that there is no phenotypic trade-off between the treated and non-treated environment (and indeed, on the Figure 2, A and B are distinct points, indicating that there is a phenotypic trade-off, despite that no cost is detected). It can just indicate that the wild-type reference is not adapted well to the non-treated environment. At this point, one might argue that all this confusion arises because the mutation R was not a "resistance" mutation to begin with. The mutation R is beneficial in both the treated and non- treated environment, and so, it may be better interpreted as a

334 beneficial mutation to the non-treated environment than as a 'resistance' mutation. For
335 instance, if we are talking of a bacteria mutant showing this property in a laboratory test, we
336 may want to say that the mutation R corresponds to adaptation to the "laboratory condition",
337 not really to the drug *per se*. But, then, how is a resistance mutation truly defined?

What is a resistance mutation?

If resistance mutations cannot be defined by the fact that they are beneficial in the treated
environment, relative to wild type (as we did up to now), then, how do we classify them? In
fact, it might be possible to define them more specifically by saying that they are beneficial in
the treated environment relatively to wild type, provided that the wild type is perfectly well
adapted to the corresponding non-treated environment. In principle, this definition makes
sense, as it avoids conflating adaptation to conditions that are shared by both treated and non-
treated environments, with adaptation to the drug itself. Yet, with this definition, the mutation
R illustrated on Figure 2 would still be a beneficial mutation in the treated environment. R is
closer to A (the optimum with the drug) than would a wild type well adapted to the non-
treated environment (distance RA < AB). In such a case, the phenotypic direction
corresponding to resistance would be the AB axis (and the pleiotropic effects best defined on
axes orthogonal to AB). This would be in general clearer and more insightful. With such a
definition, it is possible to distinguish mutation R and R' on Figure 3 for instance. Both would
be beneficial in both treated and non-treated environment, relative to a wild-type in O, but
only R would be beneficial in the treated environment relative to a wild-type in B.

Figure 3. Sketch of treated and non-treated environments, with distinct phenotypic requirements (phenotypic optima A and B, respectively) in a two-trait space. Relative to a wild-type positioned in O, both the resistance mutation R and R' brings the phenotype closer to A. As in Figure 1, they are therefore both beneficial mutations, to the treated environment, relative to the wild type O. However, if resistance is defined by comparing to a wild-type well adapted to the non-treated environment (*i.e.* to a wild type located on B), then only mutation R would be beneficial, and therefore could qualify as being a resistance mutation. With this comparison, costs would also be better defined (*i.e.* would always be positive). As with Figure 1 and 2, fitness depends on phenotypic distance and a mapping, which is not illustrated but would correspond to a third orthogonal axis.

The problem is that measures of fitness will be made against a wild-type and it may not be straightforward to determine whether this wild type is well adapted to the non-treated environment (but not impossible, as e.g. in cases of experimental evolution in the laboratory where the wild type can be chosen as the type that evolved in the non treated environment for a long time). Without this knowledge, it may be very difficult to classify mutations that are specifically beneficial in the treated environment (*i.e.* "resistance" mutations as defined here), versus mutations that are beneficial in both treated and non-treated environments (see e.g. Marcusson *et al.* 2009 for an example of this problem).

There is yet another difficulty lurking in the vast range of possible natural situations. Just as there are many non-treated environments, there may be many treated environments as well. In particular, any drug may be added in different quantities or concentrations to any given environment. Here resides a problem, which is rarely addressed in studies on resistance: different drug concentrations can correspond to different intensity of selection (Milesi *et al.* 2016), but they can also correspond to different optimum phenotypes (Harmand *et al.* 2017). This possibility needs to be demonstrated empirically (as in Harmand *et al.* 2018), and cannot be ignored *a priori*. Admitting that natural processes have some degree of continuity, it is likely that differing doses end up corresponding to different phenotypic optima. Indeed, it is difficult to conceive that adding a vanishingly small quantity of drug suddenly shifts away phenotypic requirements, and that further increases in doses only change the selection intensity around that shifted phenotypic peak.

Figure 4. Sketch of a non-treated environment (phenotypic optimum in B) and two treated
environments, with different doses of drug (optima A_1 and A_2). Two mutations are illustrated,
with fitness effects compared to a wild type well adapted to the non-treated environment (*i.e.*
located at B). Using the definition from Figure 3, R_1 is a resistance mutation with respect to
environment A_1, but not with respect to environment A_2, and reciprocally for mutation R_2.
Nevertheless, both mutations have the same cost. As with Figure 1 to 3, fitness depends on
phenotypic distance and a mapping, which is not illustrated but would correspond to a third
orthogonal axis.

If different drug doses correspond to different optima (say A_1 and A_2 on Figure 4), it is fairly
easy to imagine two mutations R_1 and R_2 that would qualify as resistance mutations, in each
of the two environments, but not in the other. On Figure 4, R_1 is a resistance mutation for drug
dose 1 (optimum A_1), but not for drug dose 2 (optimum A_2), and reciprocally for R_2. This
difference does not arise because these mutations have different "costs". In fact, the two
mutations illustrated on Figure 4 have the same cost (the same fitness in non-treated
environment B). This situation occurs because phenotypic requirement is different with
different doses, such that a phenotypic change can be favourable in one environment (here
one dose), but not another. 'Resistance mutations' can differ not only in their benefit and costs
at a given drug dose, but also in their fitness effects at other doses. When optima for doses are
different, there are many underlying trade-offs, which are not necessarily caused by differing
costs in the non-treated environment (Harmand *et al.* 2018). Trade-offs among doses are not
captured by studying costs. A common view is that mutations conferring strong resistance
(*i.e.* resistance to a high dose) may carry strong costs (Melnyk *et al.* 2015), explaining perhaps

410 why they may not be beneficial at a lower dose. This may well be true, but not necessarily

411 (Harmand *et al.* 2017). A mutation favourable at high dose may be deleterious at low dose,

412 irrespectively of its cost. This is the case illustrated on Figure 4. Studying "cost" and "benefit"

413 at one particular dose may give the illusion that all trade-offs are understood. In fact, this is

414 not the case: all trade-offs among doses will be missed.

Summary, conclusion and practical implications

Taking into account the 'cost of resistance' has been a major progress because it is essential to

distinguish the fitness effects of resistance mutations in treated versus non-treated

environments. However, this cost is also very expensive conceptually as it is associated with

too many simplifications, to the point that it may even be misleading. In practical terms,

should the term 'cost of resistance' be avoided? The term is extremely widespread and in

many cases, it has at least the merit to attract the attention to the fact that fitness effects are

different in different environments. In most cases and depending on context, the more robust

concepts of fitness trade-off across environments or pleiotropy could be used, although the

concision of the expression "cost of resistance" will be difficult to match. We hope that this

perspective will help correcting some of the sloppy usage of the term and dismiss the implicit

expectations based on this terminology.

First, the interpretation of "cost" in terms of pleiotropic effects is very unclear. Pleiotropic

effects may be better defined as effects projected on phenotypic axes orthogonal to resistance

phenotype, than in terms of fitness effect in the non-treated environment (see Figure 1). In

practical terms, this indicates that the fitness effects across environments should be better

distinguished from pleiotropic effects. Pleiotropy is not necessarily the cause of fitness trade-

off. A single trait can present different optimal values in different environments and primary

resistance traits can exhibit a trade-off without having a pleiotropic effect. This clarifies our

expectation about the process of adaptation: compensatory evolution is not expected to

systematically reduce the "cost of resistance" to zero.

Second, there are as many costs as there are non-treated environments. The idea that there is

one cost associated with a resistance mutation is an extreme and naïve simplification. In

practice, this indicates that studying the fitness effects of resistance mutations "in the wild"

(*i.e.* beyond the simplified laboratory conditions) is very important.

Third, costs of resistance are ill defined when several precautions are not taken. For instance,

failing to measure costs relative to a well-adapted wild type to the non-treated environment

can lead to absurd notions, such as 'negative costs' that serve no conceptual clarification

(Figure 2). In practice, these precautions can be very difficult to meet, as the degree of

adaptation of the wild type to the non-treated environment will be in general difficult to

assess, but steps can be taken in that direction (e.g. by carefully choosing the reference

genotype and the reference environment). In any case, this issue is important to keep in mind

when interpreting results.

Fourth, the concept of cost leads to an oversimplified and often erroneous view of trade-offs

across environments. Finding costs equal to zero (*i*) cannot be used to say that there is no

trade-offs between treated and non-treated environments; (*ii*) cannot be interpreted by saying

that pleiotropic effects were compensated; (*iii*) completely misses the possibility that fitness

trade-off may occur among different doses. In practice, finding a mutation with no "cost" (or

a "negative cost"), should not be surprising. It does not prove the existence of "Darwinian

demons"; it should not be used to imply that there are no trade-off across environments and

therefore that resistance management strategies will necessary fail. As we have shown, this

finding can simply result from a particular choice for the reference genotype and

environment. Considering fitness effects across the range of possible doses is also important

beyond the simplified conditions of most lab-based ecotoxicological tests and

ecotoxicological fitness proxies such as LD50, MIC, dose responses *etc*. Importantly, finding

that resistance mutations are not favourable at all doses cannot be attributed to their "cost".

For instance, high resistance mutations may not be beneficial at low doses, not because they

have a high cost, but simply because they do not match phenotypic requirements that are

optimal at low dose. Here again, the concept of cost leads in practice to biased expectations.

Finally, observing that a resistance mutation does not decrease in frequency after an arrest of

treatment, is not proving that there is "no cost" or no trade-off across environments. Other

causes of frequency changes must be first investigated (*i.e.* effect of gene flow, effect of

residual or hidden treatments, drift, indirect selection) as well as possible ascertainment biases

(low power to detect slow frequency change).

Overall, it may be safer in most cases to simply discuss and measure the fitness effects of

mutation in different environments and to carefully consider the role of the reference

genotype/phenotype when interpreting relative fitnesses. All these points hold for many other

situations of adaptation besides resistance, but this is perhaps where the vocabulary and the

conceptual issues are the most acute and widespread. Differences among selective conditions

and the occurrence of pleiotropy are both important ideas in evolutionary ecology. However,

they cannot be solely summarized by assigning resistance mutations a 'benefit' and 'a cost' and essentialize their properties. There is a problem with a reduction of evolutionary thinking to a cost-benefit thinking, with "fitness" as a universal currency, valid regardless of ecological conditions. Although fitness is a general concept and a universal currency for adaptation, different conditions entail different fitnesses and possibly different phenotypic requirements (different adaptations). The vocabulary that we use should not oversimplify these ideas.

Acknowledgments

We thank Helen Alexander, Danna Gifford, Inês Fragata and Claudia Bank for comments on the manuscript. We thank G. Martin for insightful discussions.

References

Abedi Z. H., Brown A. W. A., 1960 Development and reversion of DDT resistance in *Aedes aegypti*. Can. J. Genet. Cytol. 2: 252–261.

Anderson R. M., May R. M., 1982 Coevolution of hosts and parasites. Parasitology 85: 411–426.

Andersson D. I., Levin B. R., 1999 The biological cost of antibiotic resistance. Curr. Opin. Microbiol. 2: 489–93.

Andersson D. I., Hughes D., 2010 Antibiotic resistance and its cost: is it possible to reverse resistance? Nat. Rev. Microbiol. 8: 260–271.

Angst D. C., Hall A. R., 2013 The cost of antibiotic resistance depends on evolutionary history in *Escherichia coli*. BMC Evol. Biol. 13.

Antonovics J., 1977 Metal tolerance in plants: perfecting an evolutionary paradigm. In: Institute for Environmental Studies U of T (Ed.), *International Conference on Heavy Metals in the Environment, Toronto, Ontario, Canada*, Toronto, pp. 169–186.

Bataillon T., Zhang T., Kassen R., 2011 Cost of adaptation and fitness effects of beneficial mutations in *Pseudomonas fluorescens*. Genetics 189: 939–49.

Bell G., 2008 *Selection*. Oxford Univ. Press, Oxford.

Bergelson J., Purrington C. B., 1996 Surveying Patterns in the Cost of Resistance in Plants. Am. Nat. 148: 536–558.

Björkman J., Nagaev I., Berg O. G., Hughes D., Andersson D. I., 2000 Effects of environment

on compensatory mutations to ameliorate costs of antibiotic resistance. Science 287: 1479–1482.

Bourguet D., Prout M., Raymond M., 1996 Dominance of insecticide resistance presents a plastic response. Genetics 143: 407–416.

Bradshaw A. D., 1984 Adaptation of plants to soils containing toxic metals - a test for conceit. In: Evered D, Collins GM (Eds.), *Origins and development of adaptation*, Pitman Books, London, p. 273.

Carrière Y., Deland J., Roff D., Vincent C., 1994 Life-history costs associated with the evolution of insecticide resistance. Proc. R. Soc. London B Biol. Sci. 258.

Chevin L. M., Hospital F., 2008 Selective sweep at a quantitative trait locus in the presence of background genetic variation. Genetics 180: 1645–1660.

Cohan F. M., King E. C., Zawadzki P., 1994 Amelioration of the deleterious pleiotropic effects of an adaptive mutation in *Bacillus subtilis*. Evolution 48: 81–95.

Curtis C. F., Cook L. M., Wood R. J., 1978 Selection for and against insecticide resistance and possible methods of inhibiting the evolution of resistance in mosquitoes. Ecol. Entomol. 3: 228–273.

Cvijović I., Good B. H., Jerison E. R., Desai M. M., 2015 Fate of a mutation in a fluctuating environment. Proc. Natl. Acad. Sci. U. S. A. 112: E5021-8.

Davies A. G., Game A. Y., Chen Z., Williams T. J., Goodall S., *et al.*, 1996 Scalloped wings is the Lucilia cuprina Notch homologue and a candidate for the Modifier of fitness and asymmetry of diazinon resistance. Genetics 143: 1321–1337.

Day T., Huijben S., Read A. F., 2015 Is selection relevant in the evolutionary emergence of drug resistance? Trends Microbiol. 23: 126–33.

Depledge M., 2011 Reduce drug waste in the environment. Nature 478: 36.

Dobzhansky T., 1951 *Genetics and the origin of species*. Columbia university press, New York.

Endler J. A., 1977 *Geographic variation, speciation and clines*. Princeton University Press, Princeton.

Eyre-Walker A., Keightley P. D., 2007 The distribution of fitness effects of new mutations. Nat. Rev. Genet. 8: 610–618.

Foucault M. L., Courvalin P., Grillot-Courvalin C., 2009 Fitness cost of VanA-type
vancomycin resistance in methicillin-resistant *Staphylococcus aureus*. Antimicrob.
Agents Chemother. 53: 2354–2359.

Gallet R., Latour Y., Hughes B. S. B. S., Lenormand T., 2014 The dynamics of niche
evolution upon abrupt environmental change. Evolution 68: 1257–1269.

Gassmann A. J., Carrière Y., Tabashnik B. E., 2009 Fitness Costs of Insect Resistance to
Bacillus thuringiensis. Annu. Rev. Entomol. 54: 147–163.

Georghiou G. P., Taylor C. E., 1977 Genetic and biological influences in the evolution of
insecticide resistance. J. Econ. Entomol. 70: 319–323.

Gifford D., Moss E., MacLean R., 2016 Environmental variation alters the fitness effects of
rifampicin resistance mutations in *Pseudomonas aeruginosa*. Evolution 70: 725–730.

Gillespie J. H., 1975 Natural selection for resistance to epidemics. Ecology 56: 493–495.

Gonthier A., Guerin-Faublee V., Tilly B., Delignette-Muller M.-L., 2001 Optimal growth
temperature of O157 and non-O157 Escherichia coli strains. Lett. Appl. Microbiol. 33:
352–356.

Guillemaud T., Lenormand T., Bourguet D., Chevillon C., Pasteur N., *et al.*, 1998 Evolution
of resistance in *Culex pipiens*: allele replacement and changing environment. Evolution
52: 443–453.

Haldane J. B. S., 1948 The theory of a cline. J. Genet. 48: 277–284.

Harmand N., Gallet R., Jabbour-Zahab R., Martin G., Lenormand T., 2017 Fisher's
geometrical model and the mutational patterns of antibiotic resistance across dose
gradients. Evolution 71: 23–37.

Harmand N., Gallet R., Martin G., Lenormand T., 2018 Evolution of bacteria specialization
along an antibiotic dose gradient. Evol. Lett. in press.

Hartl D. L., Taubes C. H., 1998 Towards a theory of evolutionary adaptation. Genetica 103:
525–533.

Hickey D. A., McNelly T., 1975 Competition Between Metal Tolerant and Normal Plant
Populations ; A Field Experiment on Normal Soil. Evolution 29: 458–464.

Jacobs B. F., Duesing J. H., Antonovics J., Patterson D. T., 1988 Growth performance of
triazine-resistant and -susceptible biotypes of *Solanum nigrum* over a range of

temperatures. Can. J. Bot. 66: 847–850.

Jain S. K., Bradshaw A. D., 1966 Evolutionnary divergence among adjacent plant
populations. I. The evidence and its theoretical analysis. Heredity 21: 407–441.

Kassen R., Bataillon T., 2006 Distribution of fitness effects among beneficial mutations
before selection in experimental populations of bacteria. Nat. Genet. 38: 484–488.

Labbé P., Sidos N., Raymond M., Lenormand T., 2009 Resistance Gene Replacement in the
mosquito *Culex pipiens*: fitness estimation from long-term cline series. Genetics 182:
303–12.

Lande R., 1980 The Genetic Covariance between Characters Maintained by Pleiotropic
Mutations. Genetics 94: 203–215.

Lande R., 1983 The response to selection on major and minor mutations affecting a metrical
trait. Heredity 50: 47–65.

Lenormand T., Guillemaud T., Bourguet D., Raymond M., 1998 Evaluating gene flow using
selected markers: a case study. Genetics 149: 1383–1392.

Lenormand T., Raymond M., 1998 Resistance management: the stable zone strategy. Proc. R.
Soc. London B 265: 1985–1990.

Lenormand T., Bourguet D., Guillemaud T., Raymond M., 1999 Tracking the evolution of
insecticide resistance in the mosquito *Culex pipiens*. Nature 400: 861–864.

Lenormand T., Raymond M., 2000 Analysis of clines with variable selection and variable
migration. Am. Nat. 155: 70–82.

Lenormand T., 2002 Gene flow and the limits to natural selection. Trends Ecol. Evol. 17:
183–189.

Lenski R. E., 1988 Experimental studies of pleiotropy and epistasis in *Escherichia coli*. I.
Variation in competitive fitness among mutant resistant to virus T4. Evolution 42: 425–
432.

Lenski R. E., 1998 Bacterial evolution and the cost of antibiotic resistance. Int. Microbiol. 1:
265–70.

Leonard K. J., Czochor R. J., 1980 Theory of genetic interactions among populations of plants
and their pathogens. Annu. Rev. Phytopathol. 18: 237–258.

Levene H., 1953 Genetic equilibrium when more than one ecological niche is available. Am.

Nat. 87: 331–333.

Levin B. R., Perrot V., Walker N., 2000 Compensatory mutations, antibiotic resistance and
the population genetics of adaptive evolution in bacteria. Genetics 154: 985–997.

Levison M. E., Levison J. H., 2009 Pharmacokinetics and pharmacodynamics of antibacterial
agents. Infect Dis Clin North Am. 23: 791–797.

MacLean R. C., Vogwill T., 2015 Limits to compensatory adaptation and the persistence of
antibiotic resistance in pathogenic bacteria. Evol. Med. Public Heal. 2015: 4–12.

Manna F., Martin G., Lenormand T., 2011 Fitness landscapes: An alternative theory for the
dominance of mutation. Genetics 189.

Marcusson L. L., Frimodt-Møller N., Hughes D., 2009 Interplay in the selection of
fluoroquinolone resistance and bacterial fitness. PLoS Pathog. 5.

Martin G., Lenormand T., 2006a A general multivariate extension of Fisher's geometrical
model and the distribution of mutation fitness effects across species. Evolution 60: 893–
907.

Martin G., Lenormand T., 2006b The fitness effect of mutations across environments: a
survey in the light of fitness landscape models. Evolution 60: 2413–2427.

Martin G., Lenormand T., 2015 The fitness effect of mutations across environments: Fisher's
geometrical model with multiple optima. Evolution 69: 1433–1447.

May R. M., Dobson A. P., 1986 Population Dynamics and the Rate of Evolution of Pesticide
Resistance. In: *Pesticide Resistance: Strategies and Tactics for Management*, The
National Academies Press, Washington, DC, pp. 170–193.

McKenzie J. A., Whitten M. J., Adena M. A., 1982 The effect of genetic background on the
fitness of diazinon resistance genotypes of the australian sheep blowfly, *Lucilia cuprina*.
Heredity 49: 1–9.

Melnyk A. H., Wong A., Kassen R., 2015 The fitness costs of antibiotic resistance mutations.
Evol. Appl. 8: 273–283.

Milesi P., Lenormand T., Lagneau C., Weill M., Labbé P., 2016 Relating fitness to long-term
environmental variations *in natura*. Mol. Ecol. 25: 5483–5499.

Nagylaki T., 1975 Conditions for the existence of clines. Genetics 80: 595–615.

Neve P., Powles S., 2005 High survival frequencies at low herbicide use rates in populations

of *Lolium rigidum* result in rapid evolution of herbicide resistance. Heredity 95: 485–
492.

Nguyen T. N., Phan Q. G., Duong L. P., Bertrand K. P., Lenski R. E., 1989 Effects of carriage
and expression of the Tn10 tetracycline-resistance operon on the fitness of *Escherichia*
coli K12. Mol. Biol. Evol. 6: 213–225.

Orr H. A., 1998 The population genetics of adaptation: The distribution of factors fixed
during adaptive evolution. Evolution 52: 935–949.

Orr H. A., 2005 The genetic theory of adaptation: A brief history. Nat. Rev. Genet. 6: 119–
127.

Regoes R. R., Wiuff C., Zappala R. M., Garner K. N., Baquero F., *et al.*, 2004
Pharmacodynamic functions: a multiparameter approach to the design of antibiotic
treatment regimens. Antimicrob. Agents Chemother. 48: 3670–6.

Sampah M. E. S., Shen L., Jilek B. L., Siliciano R. F., 2011 Dose-response curve slope is a
missing dimension in the analysis of HIV-1 drug resistance. Proc. Natl. Acad. Sci. U. S.
A. 108: 7613–8.

Schoustra S. E., Debets a J. M., Slakhorst M., Hoekstra R. F., 2006 Reducing the cost of
resistance; experimental evolution in the filamentous fungus *Aspergillus nidulans*. J.
Evol. Biol. 19: 1115–27.

Schrag S. J., Perrot V., Levin B. R., 1997 Adaptation to the fitness costs of antibiotic
resistance in *Escherichia coli*. Proc. Biol. Sci. 264: 1287–91.

Simms E. L., Rausher M. D., 1987 Costs and Benefits of Plant Resistance to Herbivory. Am.
Nat. 130: 570–581.

Smith P., Townsend M. G., Smith R. H., 1991 A Cost of Resistance in the Brown Rat?
Reduced Growth Rate in Warfarin-Resistant Lines. Funct. Ecol. 5: 441–447.

Sousa A., Magalhães S., Gordo I., 2012 Cost of antibiotic resistance and the geometry of
adaptation. Mol. Biol. Evol. 29: 1417–28.

Suckling D. M., Khoo J. G. I., 1993 Cline in frequency of azinphosmethyl resistance in light
brown apple moth (Lepidoptera: Tortricidae). J. Econ. Entomol. 86: 1308–1316.

Thiele-Bruhn S., 2003 Pharmaceutical antibiotic compounds in soils - a review. J. plant Nutr.
soil Sci. 166: 145–167.

Ventola C. L., 2015 The antibiotic resistance crisis: part 1: causes and threats. P T 40: 277–
83.

Vila-Aiub M. M., Neve P., Powles S. B., 2009 Fitness costs associated with evolved herbicide
resistance alleles in plants. New Phytol. 184: 751–767.

Wen X., Gehring R., Stallbaumer A., Riviere J. E., Volkova V. V., 2016 Limitations of MIC
as sole metric of pharmacodynamic response across the range of antimicrobial
susceptibilities within a single bacterial species. Sci. Rep. 6: 37907.

Wilson J. B., 1988 The cost of heavy-metal tolerance: an example. Evolution 42: 408–413.

Windle P. N., Franz E. H., 1979 The Effects of Insect Parasitism on Plant Competition :
Greenbugs and Barley. Ecology 60: 521–529.

World Health Organization, 2014 *Antimicrobial Resistance Global Report on Surveillance*.