1

Defects in Piwi or germ granules trigger a common program of sterility that is independent of transposon activation

Short title: Germ granules may orchestrate Piwi sterility

Authors: Katherine Kretovich Billmyre[1,2]†, Bree Heestand[1,2,3]†, Maya Spichal[1,2]†, Stephen Frenk[1,2]† and Shawn Ahmed[1,2,3]*

Affiliations:

[1]Department of Genetics, University of North Carolina, Chapel Hill, NC, USA

[2]Department of Biology, University of North Carolina, Chapel Hill, NC, USA

[3]Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill, NC, USA

†These authors contributed equally to this manuscript

*Corresponding author: shawn@med.unc.edu

Abstract:

Defective Piwi/piRNA genome silencing leads to heterochromatin dysfunction and immediate sterility in many species. Sterile Piwi mutants experience transposon expression and transposon-induced genomic instability, although the cause of Piwi mutant sterility remains uncertain. *C. elegans* germ cells deficient for Piwi pathway genome silencing factors transmit a form of heritable stress that induces sterility after growth for several generations, allowing comparisons of animals that are close to sterility but fertile with sterile siblings. Sterile Piwi pathway mutants displayed inconsistent increases in DNA damage signaling but consistently altered perinuclear liquid droplets termed germ granules. Germ granule dysfunction did not elicit significant levels of transposon expression but was sufficient to induce a range of idiosyncratic phenotypes associated with sterile Piwi pathway genome silencing mutants, including germline atrophy, reproductive arrest and univalents in oocytes. Expression of genes that perturb germ granule structure was not compromised in sterile Piwi pathway mutants, suggesting that a post-transcriptional mechanism regulates the stability of Piwi mutant germ granules. We propose that sterility in response to transgenerational deficiency for Piwi and in response to acute dysfunction of germ granules occur as a consequence of a common reproductive arrest mechanism. The germ granule abnormalities of sterile Piwi pathway mutants therefore suggest germ granule dysfunction as a cause of Piwi mutant sterility.

Keywords: Germ granules, reproductive arrest, small RNAs, heterochromatin, germ cell immortality, Heritable Stress, Piwi

Introduction:

Germ cells give rise to the mortal somatic tissues while maintaining themselves in a

pristine condition that allows them to be propagated for an indefinite number of

generations. This capacity for self-renewal is termed "germ cell immortality", and

functions to protect the germline from forms of damage that can result in sterility [1].

Defects in *Caenorhabditis elegans* (*C. elegans)* genes that promote germline

immortality cause a Mortal Germline *(mrt)* phenotype, where robust fertility occurs for

several generations but deteriorates in later generations, finally culminating in complete

sterility. The transgenerational sterility of *mrt* mutants implies that their germ cells may

age as they proliferate over the generations such that they accumulate of a form of

damage that ultimately evokes sterility. Identifying the transgenerational damage that

accumulates in *mrt* mutants and the cause of sterility are central challenges for

understanding the basis of germ cell immortality.

The initial germline immortality pathway defined in *C. elegans* was caused by

deficiency for telomerase, a reverse transcriptase that maintains telomere stability by

adding telomere repeats sequences to chromosome termini de novo. *C. elegans*

mutants that are defective for telomerase-mediated telomere repeat addition display

several phenotypes that correlate with telomerase deficiency. The first is progressive

telomere erosion as detected by Southern blotting, which represents the

transgenerational damage that accumulates in these mutants. The second is that end-

to-end chromosome fusions accumulate in these strains as they become sterile [2,3].

The sterility of *C. elegans* telomerase mutants is therefore believed to occur as a result

of genome catastrophe induced by telomere dysfunction, although the precise cause of

telomerase mutant sterility remains uncertain [4–7], possibly the creation of circular

chromosomes that are toxic to meiosis [4]. Similar to the transgenerational proliferation

defects of *C. elegans* telomerase mutants, primary human somatic cells that are

cultured in vitro have a limited proliferative capacity that is associated with telomerase

dysfunction, progressive telomere erosion and ultimately telomere dysfunction, which

triggers entry into an irreversible state of cell cycle arrest termed senescence [5,6].

Confidence that senescence of primary human fibroblasts that proliferate in vitro is

caused by telomere dysfunction was obtained by acute knockdown of the telomere

capping protein TRF2, which led to immediate telomere deprotection of telomeres

whose lengths were normal, accompanied by telomere fusions and senescence [7].

A distinct fertility pathway, P-M Hybrid Dysgenesis, was discovered in the 1970's

as a temperature-sensitive *Drosophila* fertility defect that was linked to defects in

transposon silencing [8,9]. In these experiments, Drosophila M strain females were

mated with males of the more recently isolated P strain to yield F1 hybrid progeny with

high levels of P element transposition and temperature-sensitive sterility. The reduced

fertility of P-M F1 hybrids has been attributed to lack of piRNA-mediated genome

silencing for specific transposable elements [10], which are normally repressed by Piwi.

Piwi is a conserved Argonaute protein that is enriched in germ cells and interacts with

thousands of small RNAs termed piRNAs to repress transposons and foreign nucleic

acids that represent parasitic threats to the integrity of the genome [11–13]. Deficiency

for Piwi proteins leads to immediate sterility in *Drosophila*, zebrafish and mouse, and

this correlates with transposon expression and increased levels of DNA damage in

sterile F2 animals, which implies that transposon-induced genome instability could

cause sterility of Piwi mutants [9,13,14]. However, analysis of effects of transposon

copy number on P-M hybrid dysgenesis suggests that transposon-induced genome

instability may only explain a minor fraction of P-M hybrid sterility [15]. Therefore, the

cause of sterility in Piwi mutants is uncertain and remains an unsolved problem in

experimental biology.

Deficiency for the *C. elegans* Piwi ortholog *prg-1* compromises germ cell

immortality [18], rather than inducing the immediate sterility phenotype that is observed

in Piwi mutants in some other species. The reason for the transgenerational delay in the

onset of sterility in *prg-1*/Piwi mutants and in related Piwi pathway mutants of *C.*

elegans is not clear, but it is possible that genomic silencing marks that are induced by

the Piwi/piRNA pathway are more robustly maintained in germ cells of *C. elegans* if the

Piwi silencing system is inactivated. The transgenerational damage that accumulates in

C. elegans Piwi mutants may therefore correspond to progressive deterioration of Piwi-

mediated heterochromatin.

Fertility defects that occur in response to heterochromatin dysfunction in *C.*

elegans have been associated with increased levels of RNA-DNA hybrids and of

transposon-induced DNA damage [16,17]. *C. elegans prg-1* mutants display

progressively increased levels of germline apoptosis when grown for successive

generations, and apoptosis promotes germ cell atrophy for sterile *rsd-2, rsd-6* and *prg-1*

mutants [18–20]. However, very low levels of transposition occur in late-generation *prg-*

1 mutants, and in *C. elegans mutator* mutants, which do not display transgenerational

sterility, high levels of transposition do occur suggesting transposition is not sufficient to

induce transgenerational sterility [18]. Furthermore, fertility can be restored to minor

fraction of sterile late-generation *prg-1* mutants [18,20]. Together, these results suggest

that the sterility of *prg-1*/Piwi mutants is unlikely to be a consequence of debilitating

levels of transposon-induced DNA damage. However, it remains possible that

significant levels of DNA damage are unleashed when *C. elegans prg-1* mutants

become sterile.

The late-generation sterility phenotype of *C. elegans prg-1*/Piwi mutants was

recently shown to elicit a striking but pleiotropic germ cell degeneration phenotype at

the generation of sterility, as L4 larvae mature into 1 day old adults, such that most

animals display 'atrophied' germlines that retain only a small population of mitotic germ

cells, whereas some germlines are 'empty' and lack germ cells altogether, some

germlines are 'short' such that they have fewer germ cells than normal but possess both

mitotic and meiotic germ cells, and some germlines are sterile but 'normal' in size such

that they are not smaller than germlines of fertile late-generation *prg-1* mutant siblings

that are not yet sterile [20]. Consistently, temperature-sensitive Piwi pathway genome

silencing mutants *rsd-2* and *rsd-6* give rise to late-generation sterile adults that exhibit

germline arms that are normal in size but also give rise to small and atrophied germline

arms that are promoted by apoptosis [19]. Moreover, atrophied germ lines of sterile *prg-*

1 mutants can regrow on day 2 of adulthood, and a small fraction of constitutively sterile

5 day old *prg-1* mutant adults can become fertile again if they are shifted to a distinct

food source [20]. The ability of sterile late-generation of *C. elegans prg-1*/Piwi mutant

adults to become fertile suggests that Piwi mutant sterility is a reversible form of

reproductive arrest termed Adult Reproductive Diapause, which can be induced in

response to environmental stresses such as starvation [22–24]. We therefore proposed

that late-generation sterility of *prg-1*/Piwi mutants may occur in response to

accumulation of a heritable stress that is transmitted by germ cells [20].

Some Argonaute proteins that regulate epigenomic pathways are located in

perinuclear germ granules, which are conserved germline structures akin to liquid

droplets that interact with nuclear pore complexes and are necessary for fertility [25]. *C.*

elegans germ granules are termed P granules based on antibodies that label the P

blastomere that generates the germ cell lineage [26]. Core P granule proteins have

been shown to play redundant functions making it difficult to eliminate P granules [27].

Here we study the late-generation sterility phenotype of Piwi pathway genome silencing

mutants and find that it is associated with abnormalities of germ granules, and we

demonstrate that acute germ granule dysfunction is sufficient to phenocopy the

reproductive arrest phenotypes of sterile Piwi mutants.

Results:

Genome instability in Piwi pathway genome silencing mutants

Transgenerational sterility in *C. elegans* telomerase mutants that lack the ability to

maintain sequences that cap chromosome ends is caused by high levels of telomere

fusions and associated genome damage [2]. Many *mrt* mutants only become sterile

when grown at the non-permissive temperature 25°C [28], some of which are deficient

for Piwi pathway-mediated genome silencing [1,16,19,21,29–31]. However, the

deficiency for the Piwi Argonaute protein PRG-1 or the nuclear RNA interference (RNAi)

proteins NRDE-1 or NRDE-4 results in progressive sterility at any temperature

[16,18,30–33]. We studied the sterility of Piwi pathway mutants that were temperature-

sensitive for germ cell immortality at 25°C and also non-conditional Piwi pathway

mutants at 20°C. We compared fertile late-generation Piwi pathway mutants with late

generation siblings that were sterile, in an effort to learn about the cause of their sterility.

Although DNA damage has been suggested not to contribute significantly to the

sterility of *prg-1* mutants based on several lines of evidence, the levels of DNA damage

signaling in sterile animals have not been previously studied. We therefore examined

prg-1 mutants grown at 20°C across several generations for the presence of the DNA

damage response, as detected by an antibody to phosphorylated S/TQ, a protein

modification that is created by two sensors of DNA damage, ATM and ATR, which are

phosphatadylisonisol-3-kinase like protein kinases [34,35]. We observed a significant

increase in the fraction of germ cells with upregulated DNA damage response for late-

generation fertile and sterile *prg-1* mutants (~3- to 6-fold) when compared to either wild-

type controls or to early-generation *prg-1* mutants (Fig. 1a,b). However, the fraction of

germ cells with an activated DNA damage response was not consistently elevated in

sterile *prg-1*/Piwi mutants in comparison to fertile late-generation siblings (Fig. 1b). We

also tested temperature-sensitive Piwi pathway mutants and found that the wild-type

DNA damage response was upregulated by growth at 25°C but that this level was not

further elevated in germlines of sterile late-generation *rsd-6, nrde-2* or *hrde-1* mutant

adults (Fig. 1b,c).

Piwi pathway genome silencing mutants exhibit P granule defects at sterility

Deficiency for PRG-1 causes delocalization of the P granule protein PGL-1 [36], and

combined loss of the epigenomic regulators SPR-5 and LET-418, or loss of H3K9

methylation modifier SET-2 and the nuclear RNAi pathway, results in immediate sterility

that is associated with disruption of P granules [37,38]. Whilst characterizing germ line

defects of sterile *rsd-6* mutants by immunofluorescence, we found that a P granule

antibody exhibited altered staining patterns in sterile but not fertile late-generation *rsd-6*

mutant siblings (Fig 2d-d'). We also found that sterile late-generation *hrde-1* and *nrde-2*

temperature-sensitive mutant germ lines displayed P granule defects (Fig 2c-c', S1),

which were absent or comparatively minor in fertile late-generation mutant adults that

were close to the generation of sterility (Fig. 2c-c', S1, Table 1). When compared to

wild-type controls, P granule defects in the sterile mutants ranged from a loss of P

granules in a subset of cells to varying amounts of P granule disorganization (Fig. 2,

S1)*. Sterile *hrde-1(tm1200)* mutant animals displayed the most severe P granule

phenotype, as the majority of sterile *hrde-1* mutant germ cells exhibited little to no

staining (Fig. S1). We also examined the temperature sensitive *mrt* mutant *rbr-*

2(tm1231), which encodes a histone 3 lysine 4 demethylase that promotes genomic

silencing and likely functions downstream of small RNAs to promote germ cell

immortality [39]. Similarly to *hrde-1* and *nrde-2* mutants, late-generation fertile *rbr-2*

mutant animals did not exhibit obvious P granule defects, while loss of P granules was

observed in many *rbr-2* mutant germ cells at sterility (Fig. 2e-e').

Germ line atrophy occurs in some but not all *prg-1* mutants as L4 larvae develop

into sterile adults [20]. Large-scale apoptosis contributes to this germ cell degeneration

phenotype [20], and P granules have been shown to vanish in cells that undergo

apoptosis [40,41]. However, we observed widespread P granule disorganization in

sterile germlines of *rsd-6* mutants that were wildtype in size and had not undergone

significant amounts of apoptosis (Fig. 2d).

nrde-1 and *nrde-4* mutant animals displayed a distinct sterility phenotype,

whereby only a subpopulation of sterile germline arms displayed P granule defects

(60% and 22% respectively (Fig. S1, Table 1). This suggests that loss of *nrde-1* and

nrde-4, which act downstream of multiple small RNA genome silencing pathways, may

dampen the pronounced P granule defects that are observed in germline arms of most

sterile Piwi pathway mutants.

Although most Piwi pathway mutants do not display severe fertility defects until

the generation of sterility, late-generation *prg-1*/Piwi mutants display very small brood

sizes for many generations prior to sterility [20]. By immunofluorescence, we observed a

wide range of P granule defects in both sterile and fertile late-generation *prg-1* mutant

germ cells (Fig. 2b, S1), such that many germlines exhibited strong P granule

disorganization that either resulted in an irregular distribution of or a total absence of P

granules (Fig. 2b-b', S1).

C. elegans P granules are adjacent to Mutator bodies, which house Mutator

proteins that recruit RNA-dependent RNA polymerase to create secondary effector

siRNA populations in response to primary siRNAs [42]. *mutator* mutants are

dysfunctional for secondary siRNA biogenesis and remain fertile indefinitely at low

temperatures [18], but display an immediate highly penetrant fertility defect at 25^{o}C that

could be caused by dysfunction of Piwi pathway-mediated heterochromatin [43]. We

examined germlines of sterile *mut-14* mutant adults at the restrictive temperature of

25°C and observed pronounced P granule defects in 87% of animals (Fig. 2f-f', Table

1).

We conclude that germ granule morphology was consistently altered in most

sterile Piwi pathway mutants (Fig. 2, Table 1), even though there was qualitative

variability in the nature of the germ granule defects in distinct Piwi pathway mutants.

These results contrast with levels of DNA damage signaling that were either unchanged

or inconsistently increased in sterile Piwi pathway mutants in comparison to fertile late-

generation siblings (Fig. 1).

Germ granule dysfunction is sufficient to elicit phenotypes of sterile Piwi pathway

genome silencing mutants

We initially addressed the significance of P granule dysfunction in sterile late-generation

small RNA genome silencing mutants using an RNA interference clone that

simultaneously targets four P granule subunits *pgl-1*, *pgl-3*, *glh-1*, *glh-4* [27,44]. This

resulted in many second generation (G2) sterile adults at 25°C (80%) (Fig. 3a). The

germlines of G2 P granule RNAi L4 larvae were normal in size in comparison to control

worms. However, a pronounced germ cell atrophy phenotype occurred as many P

granule-depleted animals matured into sterile adults, resulting in a significant change in

their germline profile (Fig. 3b, p=3.93E-24). Furthermore, oocyte univalents were

previously observed in sterile but not fertile late-generation *rsd-6* mutants [19], and

45.5% of oocytes of sterile P granule depleted adults contained 7-12 univalents (Fig. 3c,

e-h).

Deficiency for *pgl-1* phenocopies of reproductive arrest of sterile *prg-1* mutants

The germline phenotypes observed in P granule-depleted worms mimicked those of sterile *prg-1*/Piwi mutant adults (Fig. 3a-c), which can regrow their atrophied germlines and become fertile, suggesting that germ granule dysfunction might be sufficient to cause a form of reproductive arrest termed Adult Reproductive Diapause [20]. To test this hypothesis, we used a mutant deficient for the P granule component *pgl-1*, which displays a pronounced sterility phenotype if shifted to the restrictive temperature of 25°C [45]. While most first generation (G1) *pgl-1* animals at 25°C displayed normal germlines at the L4 larval stage, there was a significant shift in the germline profiles with 52% of day 1 adults showing germline atrophy (Fig. 3b, p=1.04E-35), similar to P granule RNAi knockdown and sterile *prg-1* mutants [20]. Furthermore, when oocytes of sterile *pgl-1* mutant adults were scored, we found that 56.7% of oocytes had univalents, in agreement with the oocyte univalents observed in response to P granule depletion by RNAi (Fig. 3c).

Some G1 progeny of *pgl-1* mutants whose parents were shifted from 20°C to 25°C were fertile and gave rise to G2 25°C animals that were uniformly sterile. 25°C G2 *pgl-1* mutant L4 larvae had germline profiles that displayed more severe germ cell proliferation defects than those of G1 L4 larvae, and day 1 adults showed a striking 54% increase in empty germlines (Fig. 3d, p=2.44E-05). Therefore, germlines of G1 progeny of *pgl-1* mutant mothers that were shifted to the restrictive temperature of 25°C for a single generation mimicked the germ cell degeneration phenotypes of sterile Piwi pathway mutants as L4 larvae matured into adults, whereas 25°C G2 generation *pgl-1* mutant larvae displayed more pronounced germ cell proliferation defects.

We asked if the fertility of sterile 25°C G1 *pgl-1* mutants could be restored by

shifting them to the permissive temperature of 20°C. Sterile G1 *pgl-1* mutants that

lacked embryos were shifted to 20°C on day 1 of adulthood and their germlines were

scored after 48 hours, which revealed a significant change in their germline profiles

(p=0.007), with a shift from atrophied germlines towards large germlines when

compared to sterile G1 *pgl-1* sibling controls that remained sterile at 25°C (Fig. 3i). In

agreement with the regrowth of some germline arms, we observed that some sterile G1

pgl-1 mutant adults that were shifted to 20°C for several days had laid oocytes and

dead embryos, and that a few even gave rise to progeny (Fig. 3j). In contrast, sterile

pgl-1 mutant control adults that were kept at 25°C never gave rise to oocytes, dead

embryos or living progeny (Fig. 3j). The frequency of fertility observed in this

circumstance is very similar to that observed for sterile 5 day old *prg-1* mutant adults

that exit Adult Reproductive Diapause in response to an alternative food source [20].

Therefore, P granule dysfunction may induce a transient and potentially adaptive form

of reproductive arrest termed Adult Reproductive Diapause.

Transcriptional consequences of germ granule dysfunction and small RNA

genome silencing mutant sterility

The sterility of Piwi/piRNA mutants correlates with expression of transposons and

associated genome instability [9,13,14,46]. In an effort to deduce how germ granule

dysfunction might occur in small RNA genome silencing mutants, we sequenced RNA

from the nuclear RNAi defective mutants *nrde-1*, *nrde-2* and *nrde-4*. These mutants are

all defective in nuclear silencing in response to exogenous dsRNA triggers [30,47,48],

and at 25°C they all have Mortal Germlines [31]. However, at 20°C *nrde-1* and *nrde-4*

mutants become progressively sterile but *nrde-2* mutants remain fertile indefinitely [31].

We therefore focused on *nrde-1* and *nrde-4* mutants because they have relatively large

brood sizes at sterility at 20°C, such that large cohorts of L4 larvae that are poised to

become sterile can be collected. In contrast, *prg-1* mutants develop very low brood

sizes in late generations [20], and it is difficult to obtain large numbers of synchronous

animals that are poised to become sterile. We therefore prepared RNA from early-

generation and sterile-generation *nrde-1* and *nrde-4* mutants at 20°C and compared this

with RNA from wildtype and *nrde-2* mutant controls.

We found that sterile generation *nrde-1* and *nrde-4* mutant L4 larvae showed

strong upregulation of similar transposon classes in comparison to early generation

nrde-1 or *nrde-4* mutant L4 larvae or to wildtype controls (Fig. 4a). We identified 18

transposon classes that were upregulated at least two-fold in both *nrde-1* and *nrde-4*,

13 of which were CER retrotransposons, which is consistent with previous

transgenerational analysis of *hrde-1* mutants [49]. However, we also found that *nrde-2*

mutant controls that do not become sterile at 20°C displayed strong upregulation of

similar transposon loci, even in early generations. Of the 18 transposons upregulated in

both *nrde-1* and *nrde-4*, 17 were also upregulated in *nrde-2* compared with wildtype. As

a comparison, we analyzed published RNA-seq data from *prg-1(n4357)* mutant animals

[16]. Previous papers have reported that distinct classes of transposons are upregulated

in *prg-1* and *nrde-2* mutants [16,18,33,49], and we confirmed this result by showing that

MIRAGE1 was the only transposon upregulated in both *nrde-1*, *nrde-4* and *prg-1*

mutants (Fig. 4a).

We compared gene expression data from sterile generation *nrde-1* or *nrde-4*

mutant L4 larvae with late-generation *prg-1* mutants [16] and compared this with RNA

from wildtype or *nrde-2* mutant control L4 larvae in order to identify common genes

whose expression is specifically induced in L4 larvae that are poised to become sterile

and that might explain the P granule abnormalities at sterility. Late-generation *prg-1*

mutant and sterile generation *nrde-1* and *nrde-4* mutant gene expression was very

different from wildtype or *nrde-2* mutant controls RNA. Spermatogenic genes were over-

represented among genes significantly upregulated in sterile generation *nrde-1* and

nrde-4 mutant larvae ($p = 1e-15$, Chi Square test for both mutants), which is consistent

with what has been previously reported for *spr-5* mutants that become progressively

sterile [50]. However, there was no general tendency of an up- or down-regulation of

any gene category for *prg-1*, *nrde-1* and *nrde-4* mutants. Only 11 genes were

significantly upregulated in all three mutants, and there were no commonly

downregulated genes (Fig. 4c). We identified a set of 20 genes that were significantly

upregulated at least fourfold in *prg-1* and either *nrde-1* or *nrde-4*. which included many

pud (protein upregulated in dauer) genes that could be an indicator for genes

upregulated due to a general stress response of the organism (Table S2). This may be

consistent with our recent observation that DAF-16, which promotes dauer formation

and stress resistance [51], is responsible for several phenotypes of sterile generation

prg-1 mutant adults, including germ cell atrophy [20].

RNA-seq data sets have been previously obtained for germlines of P granule

defective L4 larvae and adults, but gene expression was not significantly different from

wildtype for L4 larvae, whereas somatic genes were overexpressed in P granule

defective adults because these animals displayed a germline to soma transformation

[44]. We compared sterile generation *nrde-1* and *nrde-4* RNA from L4 larvae with RNA

from germlines of P granule defective L4 larvae [44] and found no genes that were up or

downregulated in both *nrde-1/nrde-4* and P granule defective larvae. Moreover, only two

out of the 18 transposons upregulated in *nrde-1/nrde-4* were also upregulated in P

granule defective L4 larvae (Fig. 4a). We conclude that although varying degrees of P

granule disorganization occur at the generation of sterility for Piwi pathway genome

silencing mutants, and that although acute dysfunction of P granules based on

simultaneous knockdown of four P granule components leads to germ line degeneration

phenotypes that mimic those of sterile generation Piwi pathway mutants, that acute P

granule dysfunction does not cause immediate overt effects on expression of

heterochromatic segments of the genome or on genes in a manner that mimics Piwi

pathway mutant transcriptomes that are poised to become sterile.

Knockdown of a number of genes has been shown to disrupt *C. elegans*

germline P granule structure, in a manner that could be similar to what we observe in

sterile Piwi pathway mutant adult germlines [52]. We compared the expression of genes

known to perturb P granule structure with genes whose expression is significantly

altered in late-generation *prg-1* mutants or in sterile generation *nrde-1* and *nrde-4*

mutant L4 larvae. We examined gene expression in late- or sterile-generation *prg-1*,

nrde-1 or *nrde-4* mutants but could find no consistent changes in in genes whose

knockdown leads to reduced or eliminated P granule structures in wildtype animals [52].

Our analyses indicate that *prg-1* and *nrde-1/nrde-4* mutants have fundamentally

different gene expression profiles at or near sterility. We conclude that although the

sterility phenotype of *prg-1* and *nrde-1/nrde-4* is physiologically similar in terms of germ

cell degeneration at the L4 stage of development, it is very different on a molecular

level, which could be consistent with our observations that pronounced P granule

defects are observed in fertile and sterile late-generation *prg-1* mutant germ lines as

opposed to pronounced P granule defects in sterile but not fertile late-generation adults

of deficient for other Piwi pathway genes. We also note the relatively mild effects on P

granules in sterile *nrde-1* and *nrde-4* mutants, where only a subset of nuclei are

affected. Therefore, although deficiency for *prg-1* and *nrde-1/nrde-4* disrupts small

RNA-mediated genome silencing and compromises germ cell immortality, these genes

have distinct functions with regards to small RNA regulation of the genome such that we

observed only a small number of overlapping transcriptional effects.

Discussion:

Previous work in the field of Piwi pathway mutant sterility has demonstrated that the

immediate sterility that is observed in several species correlates with transposon

expression and genomic instability [9,13,14]. However, analysis of our transgenerational

model of Piwi pathway mutant sterility in *C. elegans* indicates that DNA damage, as

reflected by the activities of the ATM and ATR DNA damage response kinases, is

unlikely to be the central cause of sterility, as levels of DNA damage were not

consistently increased in sterile generation animals in comparison to fertile late-

generation siblings. These data complement previous observations that suggest that

DNA damage may not cause the sterility of *prg-1*/Piwi mutants [18], and provide an

important contrast with germ granule morphology of sterile Piwi pathway mutants.

We addressed the mechanism that evokes Piwi sterility by discovering

pronounced germ granule defects in most sterile but not fertile late-generation Piwi

pathway mutant animals. Disrupting germ granules directly by RNAi-knockdown of four

germ granule components or by shifting the *pgl-1* germ granule mutant to restrictive

temperature led to germline degeneration phenotypes that phenocopied those of sterile

generation Piwi pathway mutants [20]. Furthermore, our data suggest that univalent

chromosomes that are specifically observed in oocytes of some sterile Piwi pathway

genome silencing mutants are a secondary consequence of P granule dysfunction that

indirectly disrupts a meiotic process [53,54]. We also found that some sterile *pgl-1*

mutants could recover at permissive temperature, resulting in growth of the germline

and fertility (Fig. 3i-j), as previously observed for sterile late-generation *prg-1* mutants

whose germlines were in a quiescent state of Adult Reproductive Diapause [20]. This

implies that sterility in response to transgenerational dysfunction of the *C. elegans* Piwi

pathway and in response to acute dysfunction of germ granules lead to a common

reproductive arrest program. Defects in telomerase-mediated telomere repeat addition

results in progressive telomere erosion in the context of human somatic cell

proliferation, which leads to the accumulation of critically short dysfunctional telomeres

and senescence [5,6]. Based on this analogy, and on the ability of acute telomere

uncapping to similarly induce senescence [7], we suggest that the common reproductive

arrest program caused by acute germ granule dysfunction and by transgenerational

deficiency for Piwi suggest that the germ granule morphology defects of sterile Piwi

mutants could be responsible for their sterility. However, it is formally possible that the

germ granule defects that occur in sterile but not fertile late-generation Piwi pathway

mutants are a consequence, rather than a cause, of the reproductive arrest program

that is responsible for Piwi mutant sterility. Even if this were the case, the changes in

germ granule morphology of sterile late-generation Piwi pathway mutants are likely to

be closely related to the cause of Piwi mutant sterility.

The endogenous functions of cytoplasmic liquid droplet compartments such as

germ granules, P bodies and stress granules, which contain common protein

components, are not well understood [55]. Although germ granules have been well

studied in the context of germ cell biology and fertility, we report the first evidence to our

knowledge that connects germ granules to a complex, pleiotropic reproductive arrest

mechanism that we recently demonstrated occurs for sterile *C. elegans prg-1*/Piwi

mutants [20]. Given that reproductive arrest has been linked to stress responses in

biology, for example in response to starvation or seasonal change [23,24], we suggest

that it is not coincidental that germ granules share components with other cytoplasmic

liquid compartments like P bodies and stress granules, which are well known to respond

to stress. We speculate that the release of RNA or protein components from germ

granules could promote reproductive arrest in this context.

Lack of commonly upregulated transposons in late-generation Piwi pathway

mutants and in germ granule defective animals suggests that the reproductive arrest

that they experience is not simply a response to transposon desilencing. Given that *prg-*

*1/*Piwi mutant germ cells may transmit a form of heritable stress that occurs in response

to heterochromatin dysfunction [20], and given that we observed germ granule

disorganization in Piwi pathway mutants that generally coincided precisely with the

generation of sterility, our results suggest two possible models regarding the cause of

Piwi mutant sterility. First, the Piwi/piRNA pathway could play a direct role in the

maintenance of germ granules, which are liquid droplet composites of protein and RNA

whose biogenesis and stability may depend on protein-RNA interactions [55]. Small

RNAs created in response to the Piwi/piRNA pathway could interact with mRNAs in

germ granules to facilitate their maintenance in a manner that becomes

transgenerationally mis-regulated in Piwi pathway mutants. In this first scenario (Fig.

5b), disruption of germ granules themselves would represent the cause of reproductive

arrest, as we found for *pgl-1* mutants (Fig. 3). Some Argonaute proteins that modulate

gene expression in response to small RNAs, such as CSR-1, PRG-1 or WAGO-1, are

localized to P granules [32,33,56–58]. In addition, previous analysis of sterile *prg-1*

mutants, created by crossing *prg-1; mutator* double mutants together, resulted in adult

germlines with a range of sizes analogous to those observed in sterile late-generation

prg-1 mutants [20,59]. This led to the hypothesis that misrouting of small RNAs

associated with pro- and anti-silencing small RNA pathways could be the cause of

sterility in Piwi pathway genome silencing mutants [36,59], which may be consistent

with our observations of germ granule abnormalities in sterile Piwi pathway mutants.

However, the misrouting of small RNAs hypothesis was not previously suggested or

shown to be related to changes to germ granule structure or function [36,59].

A second scenario that could explain the sterility of *C. elegans* Piwi pathway

mutants is that transgenerational dysfunction of the Piwi/piRNA pathway could lead to

heterochromatin defects that result in the misregulation of an endogenous biochemical

pathway in late-generation *C. elegans* Piwi pathway mutants, and that this then perturbs

germ granule structure or function in a manner that elicits reproductive arrest (Fig. 5b).

When RNA from sterile generation *nrde-1* and *nrde-4* mutant L4 larvae and late-

generation *prg-1* mutants was examined, we failed to observe consistent depletion of

any gene whose knockdown is known to disrupt P granule formation [52], suggesting a

post-transcriptional reproductive arrest mechanism that could be consistent with either

scenario depicted in Fig. 5b. Nevertheless, we favor scenario 2 because deficiency for

the *C. elegans* H3K4 demethlases *spr-5* or *rbr-2*, which function within nuclei

downstream of small RNAs to promote genomic silencing, elicit a transgenerational

sterility phenotype that is similar to that of Piwi pathway silencing mutants [39].

However, we also note that heterochromatin itself is transcribed in a manner that

facilitates the maintenance of small RNA populations [60,61].

Possibly consistent with our observations, sterile mouse mutants with defects in

piRNA biogenesis have been reported to display germ granule defects [62,63]. We

suggest that germ granule dysfunction could explain the sterility observed in P-M hybrid

dysgenesis and the immediate sterility phenotype of Piwi mutants in several species

[9,13,14]. Environmental stresses such as starvation can also evoke reproductive arrest

[22,23], and our data suggest that germ granule dysfunction could represent a general

mechanism for orchestrating stress-induced reproductive arrest. We note that there has

been a rise in human infertility in recent decades that has been attributed to factors

such as psychological stress, caffeine consumption and pollutants [64,65].

The sterility of *prg-1*/Piwi mutants is a pleiotropic, non-uniform phenotype that

involves dramatic levels of germ cell atrophy and can be followed by germ line regrowth

and reproduction for a small fraction of sterile animals [20]. We suggest that this

phenotype, while non-uniform and inefficient in terms of its reversibility, is an effective

form of reproductive arrest that promotes survival of wild nematode populations that are

composed of numerous individuals. Therefore, even if the reproductive arrest

mechanism of sterile Piwi mutants is inefficient, it is likely to be under significant

evolutionary pressure.

Our results establish a novel framework for considering inheritance in the context

of epigenetic stress and how this can evoke transgenerational sterility. We suggest that

the sterility we describe in response to epigenome dysfunction represents a

developmental arrest mechanism to protect the genome or epigenome in times of

stress. We note that transposons and viruses are likely to be in a continuous arms race

between parasite and host, where components of germ granules that promote Piwi

pathway genome silencing and protect the host are likely to be targeted [52]. It may

therefore be reasonable to hypothesize the evolution of a reproductive arrest

mechanism in the context of an attack on the Piwi silencing system that is predicated on

disrupting the integrity of germ granules themselves.

Materials and Methods

497
Strains

All strains were cultured at 20°C or 25°C on Nematode Growth Medium (NGM) plates

seeded with *E. coli* OP50. Strains used include Bristol N2 wild type, *hrde-1(tm1200) III,*

rsd-6(yp11) I, nrde-1(yp4) III, nrde-2(gg95) II, nrde-4 (gg131) IV, prg-1 (n4357) I, prg-1

(tm872) I, mut-14 (pk730) V, and *rbr-2(tm1231) IV.*

RNAi Assay

Feeding RNAi plates harboring host bacteria HT115(DE3) engineered to express "quad"

dsRNA (targeting P granules) were obtained from Susan Strome [27,44]. L1 larvae were

placed onto freshly prepared feeding RNAi plates with dsRNA induced by 1 mM IPTG

(isopropyl-β-D(-)-thiogalactopyranoside) and were transferred after 1 generation at 25°C

and collected at G2 adults as described in Knutson et al 2017 [44] for DAPI staining,

oocyte and germline analysis.

<u>DAPI staining and Scoring</u>

DAPI staining was performed as previously described [28]. Briefly, L4 larvae were

selected from sibling plates and sterile adults were singled as late L4s, observed 24

hours later for confirmed sterility, and then stained 48 hours after collection. Univalents

were scored by counting DAPI bodies in the -1 to -4 oocytes. Germline profiles were

scored using the method outlined in Heestand et al 2018 [20].

<u>Statistical Analysis</u>

Statistical analysis was performed as previously described [20]. Briefly, statistical

analysis was performed using the R statistical environment [66]. For germline

phenotypes, contingency tables were constructed and pairwise Chi Square tests with

Bonferroni correction was used to determine significant differences in in germline

phenotype distributions. The significance of the DNA damage response analysis was

determined by a Kruskal-Wallis test, followed by a Mann-Whitney test between

individual samples. P-values were adjusted with Bonferroni correction when multiple

comparisons were performed.

RNA extraction and sequencing

Animals were grown at 20°C on 60 mm NGM plates seeded with OP50 bacteria. RNA

was extracted using Trizol (Ambion) followed by isopropanol precipitation. Library

preparation and sequencing was performed at the UNC School of Medicine High-

Throughput Sequencing Facility (HTSF). Libraries were prepared from ribosome-

depleted RNA and sequenced on an Illumina Hiseq 2500.

RNA-seq Analysis

The following publicly available RNA-seq datasets were download from the Gene

Expression Omnibus (https://www.ncbi.nlm.nih.gov/geo/): GSE92690 (P granule RNAi

experiment) and GSE87524 (*prg-1* experiment). Adapter trimming was performed as

required using the bbduk.sh script from the BBmap suite [67] and custom scripts. Reads

were then mapped to the *C. elegans* genome (WS251) using hisat2 [68] with default

settings and read counts were assigned to protein-coding genes using the

featureCounts utility from the Subread package [69] . For multimapping reads, each

mapping locus was assigned a count of 1/n where n=number of hits. Differentially

expressed genes were identified using DESeq2, and were defined as changing at least

2-fold with FDR-corrected p-value < 0.01. For analysis of transposon RNAs, reads were

mapped to the *C. elegans* transposon consensus sequences downloaded from

Repbase (http://www.girinst.org/repbase/) with bowtie [70] using the options -M 1 -v 2.

Transposons with fewer than 10 counts in each sample were excluded from further

analysis. Counts were normalized to the total number of mapped reads for each library

for the prg-1 dataset, or to the total number of non-ribosomal mapped reads for all other

datasets. A pseudocount of 1 was added to each value to avoid division by zero errors.

Analysis of sequencing data and plot creation was performed using the R statistical

computing environment [66].

Accession numbers: RNA-seq data reported in this study have been submitted to the

GEO database and will be available at the time of publication.

Immunofluorescence

Adult hermaphrodites raised at 20°C or 25°C were dissected in M9 buffer and flash

frozen on dry ice before fixation for 1 min in methanol at -20°C. After washing in PBS

supplemented with 0.1% Tween-20 (PBST), primary antibody diluted in in PBST was

used to immunostain overnight at 4 °C in a humid chamber. Primaries used were 1:50

OIC1D4 (Developmental Studies Hybridoma Bank). Secondary antibody staining was

performed by using a Cy3 donkey anti-mouse or Cy-5 donkey anti-rabbit overnight at

4°C. All images were obtained using a LSM 710 laser scanning confocal and were

taken using same settings as control samples. Images processed using ImageJ.

DNA Damage Assay

Worms that were close to sterility were isolated and defined as sterile if they did not

have any offspring as day 3 adults at 20˚C or day 2 adults at 25˚C. Fertile siblings of

sterile worms were defined as 'close to sterility'. The presence of the DNA damage

response was determined by using a phospho-specific antibody targeting the

phosphorylated consensus target site of ATM and ATR kinases (pS/TQ) (Cell Signaling

Technology). This antibody has only been shown to stain a DNA damage response in

the germline and was used as previously described [35].

Acknowledgments: We thank members of the Ahmed lab for critical reading of the

manuscript and Jacinth Mitchell for wildtype control, *nrde-1* and *nrde-2* RNA-seq data.

Some strains were provided by the CGC, which is funded by NIH Office of Research

Infrastructure Programs (P40 OD010440). This study was supported by NIH grants F32

GM120809 (K.B) and RO1 GM083048 (S.A). M.S. was supported by a DFG postdoc

fellowship.

Competing interests: Authors declare no competing interests.

Author contributions: K.B., B.H., M.S. and S.F. performed experiments. S.F. analyzed

the data. K.B., B.H., M.S., S.F. and S.A. wrote manuscript.

References:

1.   Smelick C, Ahmed S. Achieving immortality in the C. elegans germline. Ageing Res Rev. 2005;4: 67–82.

doi:10.1016/j.arr.2004.09.002

2.   Meier B, Clejan I, Liu Y, Lowden M, Gartner A, Hodgkin J, et al. trt-1 is the Caenorhabditis elegans

catalytic subunit of telomerase. PLoS Genet. Public Library of Science; 2006;2: e18.

doi:10.1371/journal.pgen.0020018

3.   Lowden MR, Meier B, Lee TWS, Hall J, Ahmed S. End joining at Caenorhabditis elegans telomeres.

Genetics. 2008;180: 741–754. doi:10.1534/genetics.108.089920

4.   Nakamura TM, Cooper JP, Cech TR. Two modes of survival of fission yeast without telomerase. Science.

1998;282: 493–6. Available: http://www.ncbi.nlm.nih.gov/pubmed/9774280

5.   Shay JW, Wright WE. Senescence and immortalization: role of telomeres and telomerase. Carcinogenesis.

2005;26: 867–874. doi:10.1093/carcin/bgh296

6.   Sharpless NE, DePinho RA. How stem cells age and why this makes us grow old. Nat Rev Mol Cell Biol.

2007;8: 703–713. doi:10.1038/nrm2241

7.   Karlseder J, Broccoli D, Dai Y, Hardy S, de Lange T. p53- and ATM-dependent apoptosis induced by

telomeres lacking TRF2. Science. 1999;283: 1321–5. Available:

http://www.ncbi.nlm.nih.gov/pubmed/10037601

8.   Kidwell MG, Kidwell JF, Sved JA. Hybrid Dysgenesis in DROSOPHILA MELANOGASTER: A

Syndrome of Aberrant Traits Including Mutation, Sterility and Male Recombination. Genetics. 1977;86:

813–33. Available: http://www.ncbi.nlm.nih.gov/pubmed/17248751

9.   Kelleher ES. Reexamining the P-Element Invasion of Drosophila melanogaster Through the Lens of piRNA

Silencing. Genetics. 2016;203: 1513–1531. doi:10.1534/genetics.115.184119

10.   Brennecke J, Malone CD, Aravin AA, Sachidanandam R, Stark A, Hannon GJ. An epigenetic role for

maternally inherited piRNAs in transposon silencing. Science. 2008;322: 1387–92.

doi:10.1126/science.1165171

11.   Aravin AA, Hannon GJ, Brennecke J. The Piwi-piRNA pathway provides an adaptive defense in the

transposon arms race. Science. 2007;318: 761–4. doi:10.1126/science.1146484

12.   Luteijn MJ, Ketting RF. PIWI-interacting RNAs: from generation to transgenerational epigenetics. Nat Rev

Genet. Nature Publishing Group; 2013;14: 523–534. doi:10.1038/nrg3495

13.   Siomi MC, Sato K, Pezic D, Aravin AA. PIWI-interacting small RNAs: the vanguard of genome defence.

Nat Rev Mol Cell Biol. Nature Publishing Group; 2011;12: 246–258. doi:10.1038/nrm3089

14.   Juliano C, Wang J, Lin H. Uniting Germline and Stem Cells: The Function of Piwi Proteins and the piRNA

Pathway in Diverse Organisms. Annu Rev Genet. 2011;45: 447–469. doi:10.1146/annurev-genet-110410-

132541

15.   Srivastav SP, Kelleher ES. Paternal Induction of Hybrid Dysgenesis in *Drosophila melanogaster* Is Weakly

Correlated with Both *P* -Element and *hobo* Element Dosage. G3: Genes|Genomes|Genetics.

2017;7: 1487–1497. doi:10.1534/g3.117.040634

16.   McMurchy AN, Stempor P, Gaarenstroom T, Wysolmerski B, Dong Y, Aussianikava D, et al. A team of

heterochromatin factors collaborates with small RNA pathways to combat repetitive elements and germline

stress. Elife. 2017;6. doi:10.7554/eLife.21666

17.   Zeller P, Padeken J, van Schendel R, Kalck V, Tijsterman M, Gasser SM. Histone H3K9 methylation is

dispensable for Caenorhabditis elegans development but suppresses RNA:DNA hybrid-associated repeat

instability. Nat Genet. 2016;48: 1385–1395. doi:10.1038/ng.3672

18.   Simon M, Sarkies P, Ikegami K, Doebley A-L, Goldstein LD, Mitchell J, et al. Reduced insulin/IGF-1

signaling restores germ cell immortality to Caenorhabditis elegans Piwi mutants. Cell Rep. 2014;7: 762–73.

doi:10.1016/j.celrep.2014.03.056

19.   Sakaguchi A, Sarkies P, Simon M, Doebley A-L, Goldstein LD, Hedges A, et al. Caenorhabditis elegans

RSD-2 and RSD-6 promote germ cell immortality by maintaining small interfering RNA populations. Proc

Natl Acad Sci U S A. National Academy of Sciences; 2014;111: E4323-31. doi:10.1073/pnas.1406131111

20.   Heestand B, Simon M, Frenk S, Titov D, Ahmed S. Transgenerational Sterility of Piwi Mutants Represents

a Dynamic Form of Adult Reproductive Diapause. Cell Rep. Elsevier; 2018;23: 156–171.

doi:10.1016/j.celrep.2018.03.015

21.   Spracklin G, Fields B, Wan G, Vijayendran D, Wallig A, Shukla A, et al. Identification and Characterization

of Caenorhabditis elegans RNAi Inheritance Machinery. Genetics. 2017; Available:

http://www.genetics.org/content/early/2017/05/22/genetics.116.198812

22.   Angelo G, Van Gilst MR. Starvation Protects Germline Stem Cells and Extends Reproductive Longevity in

C. elegans. Science (80-). 2009;326: 954–958. doi:10.1126/science.1178343

23.   Padilla PA, Ladage ML. Suspended animation, diapause and quiescence. Cell Cycle. 2012;11: 1672–1679.

doi:10.4161/cc.19444

24.   Tatar M, Yin C. Slow aging during insect reproductive diapause: why butterflies, grasshoppers and flies are

650   like worms. Exp Gerontol. 2001;36: 723–38. Available: http://www.ncbi.nlm.nih.gov/pubmed/11295511

651   25.   Updike DL, Hachey SJ, Kreher J, Strome S. P granules extend the nuclear pore complex environment in the

652   *C. elegans* germ line. J Cell Biol. 2011;192: 939–948. doi:10.1083/jcb.201010104

653   26.   Strome S, Wood WB. Immunofluorescence visualization of germ-line-specific cytoplasmic granules in

654   embryos, larvae, and adults of Caenorhabditis elegans. Proc Natl Acad Sci U S A. 1982;79: 1558–62.

655   Available: http://www.ncbi.nlm.nih.gov/pubmed/7041123

656   27.   Updike DL, Knutson AK, Egelhofer TA, Campbell AC, Strome S. Germ-Granule Components Prevent

657   Somatic Development in the C. elegans Germline. Curr Biol. 2014;24: 970–975.

658   doi:10.1016/j.cub.2014.03.015

659   28.   Ahmed S, Hodgkin J. MRT-2 checkpoint protein is required for germline immortality and telomere

660   replication in C. elegans. Nature. 2000;403: 159–64. doi:10.1038/35003120

661   29.   Ashe A, Sapetschnig A, Weick E-M, Mitchell J, Bagijn MP, Cording AC, et al. piRNAs can trigger a

662   multigenerational epigenetic memory in the germline of C. elegans. Cell. Elsevier; 2012;150: 88–99.

663   doi:10.1016/j.cell.2012.06.018

664   30.   Buckley BA, Burkhart KB, Gu SG, Spracklin G, Kershner A, Fritz H, et al. A nuclear Argonaute promotes

665   multigenerational epigenetic inheritance and germline immortality. Nature. 2012;489: 447–451.

666   doi:10.1038/nature11352

667   31.   Burkhart KB, Guang S, Buckley BA, Wong L, Bochner AF, Kennedy S. A pre-mRNA-associating factor

668   links endogenous siRNAs to chromatin regulation. PLoS Genet. 2011;7: e1002249.

669   doi:10.1371/journal.pgen.1002249

670   32.   Batista PJ, Ruby JG, Claycomb JM, Chiang R, Fahlgren N, Kasschau KD, et al. PRG-1 and 21U-RNAs

671   interact to form the piRNA complex required for fertility in C. elegans. Mol Cell. 2008;31: 67–78.

672   doi:10.1016/j.molcel.2008.06.002

33.   Das PP, Bagijn MP, Goldstein LD, Woolford JR, Lehrbach NJ, Sapetschnig A, et al. Piwi and piRNAs Act

Upstream of an Endogenous siRNA Pathway to Suppress Tc3 Transposon Mobility in the Caenorhabditis

elegans Germline. Mol Cell. 2008;31: 79–90. doi:10.1016/j.molcel.2008.06.003

34.   Kim ST, Lim DS, Canman CE, Kastan MB. Substrate specificities and identification of putative substrates

of ATM kinase family members. J Biol Chem. 1999;274: 37538–43. Available:

http://www.ncbi.nlm.nih.gov/pubmed/10608806

35.   Vermezovic J, Stergiou L, Hengartner MO, d'Adda di Fagagna F. Differential regulation of DNA damage

response activation between somatic and germline cells in Caenorhabditis elegans. Cell Death Differ.

2012;19: 1847–1855. doi:10.1038/cdd.2012.69

36.   Phillips CM, Brown KC, Montgomery BE, Ruvkun G, Montgomery TA. piRNAs and piRNA-Dependent

siRNAs Protect Conserved and Essential C. elegans Genes from Misrouting into the RNAi Pathway. Dev

Cell. 2015;34: 457–465. doi:10.1016/j.devcel.2015.07.009

37.   Käser-Pébernard S, Müller F, Wicky C. LET-418/Mi2 and SPR-5/LSD1 cooperatively prevent somatic

reprogramming of C. elegans germline stem cells. Stem cell reports. Elsevier; 2014;2: 547–59.

doi:10.1016/j.stemcr.2014.02.007

38.   Robert VJ, Mercier MG, Bedet C, Janczarski S, Merlet J, Garvis S, et al. The SET-2/SET1 Histone H3K4

Methyltransferase Maintains Pluripotency in the Caenorhabditis elegans Germline [Internet]. Cell Reports.

2014. doi:10.1016/j.celrep.2014.09.018

39.   Alvares SM, Mayberry GA, Joyner EY, Lakowski B, Ahmed S. H3K4 demethylase activities repress

proliferative and postmitotic aging. Aging Cell. 2014;13: 245–253. doi:10.1111/acel.12166

40.   Sung M, Kawasaki I, Shim Y-H. Depletion of *cdc-25.3* , a *Caenorhabditis elegans* orthologue of *cdc25* ,

increases physiological germline apoptosis. FEBS Lett. 2017;591: 2131–2146. doi:10.1002/1873-

3468.12717

41.   Min H, Shim Y-H, Kawasaki I. Loss of PGL-1 and PGL-3, members of a family of constitutive germ-

granule components, promotes germline apoptosis in C. elegans. J Cell Sci. 2016;129: 341–353.

doi:10.1242/jcs.174201

42.   Phillips CM, Montgomery TA, Breen PC, Ruvkun G. MUT-16 promotes formation of perinuclear mutator

foci required for RNA silencing in the C. elegans germline. Genes Dev. Cold Spring Harbor Laboratory

Press; 2012;26: 1433–44. doi:10.1101/gad.193904.112

43.   Ketting RF, Haverkamp TH, van Luenen HG, Plasterk RH. Mut-7 of C. elegans, required for transposon

silencing and RNA interference, is a homolog of Werner syndrome helicase and RNaseD. Cell. 1999;99:

133–41. Available: http://www.ncbi.nlm.nih.gov/pubmed/10535732

44.   Knutson AKK, Egelhofer T, Rechtsteiner A, Strome S. Germ Granules Prevent Accumulation of Somatic

Transcripts in the Adult Caenorhabditis elegans Germline. Genetics. 2017;206. Available:

http://www.genetics.org/content/early/2017/03/02/genetics.116.198549

45.   Kawasaki I, Shim YH, Kirchner J, Kaminker J, Wood WB, Strome S. PGL-1, a predicted RNA-binding

component of germ granules, is essential for fertility in C. elegans. Cell. 1998;94: 635–45. Available:

http://www.ncbi.nlm.nih.gov/pubmed/9741628

46.   Brennecke J, Malone CD, Aravin AA, Sachidanandam R, Stark A, Hannon GJ. An Epigenetic Role for

Maternally Inherited piRNAs in Transposon Silencing. Science (80-). 2008;322: 1387–1392.

doi:10.1126/science.1165171

47.   Guang S, Bochner AF, Burkhart KB, Burton N, Pavelec DM, Kennedy S. Small regulatory RNAs inhibit

RNA polymerase II during the elongation phase of transcription. Nature. 2010;465: 1097–1101.

doi:10.1038/nature09095

48.   Guang S, Bochner AF, Pavelec DM, Burkhart KB, Harding S, Lachowiec J, et al. An Argonaute Transports

siRNAs from the Cytoplasm to the Nucleus. Science (80-). 2008;321: 537–541.

doi:10.1126/science.1157647

49.   Ni JZ, Kalinava N, Chen E, Huang A, Trinh T, Gu SG. A transgenerational role of the germline nuclear

RNAi pathway in repressing heat stress-induced transcriptional activation in C. elegans. Epigenetics

Chromatin. BioMed Central; 2016;9: 3. doi:10.1186/s13072-016-0052-x

50.   Katz DJ, Edwards TM, Reinke V, Kelly WG. A C. elegans LSD1 demethylase contributes to germline

immortality by reprogramming epigenetic memory. Cell. NIH Public Access; 2009;137: 308–20.

doi:10.1016/j.cell.2009.02.015

51.   Murphy CT, Hu PJ. Insulin/insulin-like growth factor signaling in C. elegans. WormBook. 2013; 1–43.

727   doi:10.1895/wormbook.1.164.1

728   52.   Updike D, Strome S. P Granule Assembly and Function in Caenorhabditis elegans Germ Cells. J Androl.

729   2010;31: 53–60. doi:10.2164/jandrol.109.008292

730   53.   Loidl J. Conservation and Variability of Meiosis Across the Eukaryotes. Annu Rev Genet. Annual Reviews

731   ; 2016;50: 293–316. doi:10.1146/annurev-genet-120215-035100

732   54.   Hillers K.J., Jantsch V., Martinez-Perez E., Yanowitz J.L. Meiosis. 2017. doi:10.1895/wormbook.1.178.1

733   55.   Protter DSW, Parker R. Principles and Properties of Stress Granules. Trends Cell Biol. 2016;26: 668–679.

734   doi:10.1016/j.tcb.2016.05.004

735   56.   Wang G, Reinke V. A C. elegans Piwi, PRG-1, Regulates 21U-RNAs during Spermatogenesis. Curr Biol.

736   2008;18: 861–867. doi:10.1016/j.cub.2008.05.009

737   57.   Claycomb JM, Batista PJ, Pang KM, Gu W, Vasale JJ, van Wolfswinkel JC, et al. The Argonaute CSR-1

738   and its 22G-RNA cofactors are required for holocentric chromosome segregation. Cell. 2009;139: 123–34.

739   doi:10.1016/j.cell.2009.09.014

740   58.   Gu W, Shirayama M, Conte D, Vasale J, Batista PJ, Claycomb JM, et al. Distinct Argonaute-Mediated 22G-

741   RNA Pathways Direct Genome Surveillance in the C. elegans Germline. Mol Cell. 2009;36: 231–244.

742   doi:10.1016/j.molcel.2009.09.020

743   59.   de Albuquerque BFM, Placentino M, Ketting RF. Maternal piRNAs Are Essential for Germline

744   Development following De Novo Establishment of Endo-siRNAs in Caenorhabditis elegans. Dev Cell.

745   2015;34: 448–56. doi:10.1016/j.devcel.2015.07.010

746   60.   Martienssen R, Moazed D. RNAi and Heterochromatin Assembly. Cold Spring Harb Perspect Biol. 2015;7:

747   a019323. doi:10.1101/cshperspect.a019323

748   61.   Slotkin RK, Martienssen R. Transposable elements and the epigenetic regulation of the genome. Nat Rev

749   Genet. 2007;8: 272–285. doi:10.1038/nrg2072

62.   Zheng K, Wang PJ. Blockade of Pachytene piRNA Biogenesis Reveals a Novel Requirement for

Maintaining Post-Meiotic Germline Genome Integrity. Schimenti JC, editor. PLoS Genet. 2012;8:

e1003038. doi:10.1371/journal.pgen.1003038

63.   Chuma S, Hosokawa M, Kitamura K, Kasai S, Fujioka M, Hiyoshi M, et al. Tdrd1/Mtr-1, a tudor-related

gene, is essential for male germ-cell differentiation and nuage/germinal granule formation in mice. Proc Natl

Acad Sci. 2006;103: 15894–15899. doi:10.1073/pnas.0601878103

64.   Younglai E V., Holloway AC, Foster WG. Environmental and occupational factors affecting fertility and

IVF success. Hum Reprod Update. 2005;11: 43–57. doi:10.1093/humupd/dmh055

65.   Homan GF, Davies M, Norman R. The impact of lifestyle factors on reproductive performance in the

general population and those undergoing infertility treatment: a review. Hum Reprod Update. 2007;13: 209–

223. doi:10.1093/humupd/dml056

66.   R Core Team. R: A language and environment for statistical   computing [Internet]. R Foundation for

Statistical Computing. Vienna, Austria; 2013. Available: http://www.r-project.org/

67.   Bushnell B. BBMap short-read aligner [Internet]. [cited 22 Apr 2018]. Available:

https://sourceforge.net/projects/bbmap/

68.   Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. TopHat2: accurate alignment of

transcriptomes in the presence of insertions, deletions and gene fusions. Genome Biol. 2013;14: R36.

doi:10.1186/gb-2013-14-4-r36

69.   Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning sequence

reads to genomic features. Bioinformatics. 2014;30: 923–930. doi:10.1093/bioinformatics/btt656

70.   Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient alignment of short DNA

sequences to the human genome. Genome Biol. BioMed Central; 2009;10: R25. doi:10.1186/gb-2009-10-3-

r25

Figure 1: DNA Damage Response is variable in Piwi pathway genome silencing

mutants prior to and at sterility. (A) Staining of pS/TQ in wildtype, *prg-1(n4357) and*

prg-1(tm876) mutants. pS/TQ(grey and yellow) and DAPI(blue) **(B)** Quantification of

pS/TQ staining in 20°C wildtype and *prg-1* mutants at different levels of fertility. **(C)**

Quantification of pS/TQ staining in 25°C wildtype and small RNA genomic silencing

mutants at sterility. P-values were obtained by comparing sample groups in a Mann-

Whitney U test (p-value: $P > 0.05$ = n. s., $P < 0.05$ = *, $P = <0.01$ = **, $P =< 0.001$ = ***

compared to wildtype)

Figure 2: At sterility, partial loss of P granule staining occurred in *prg-1* and

temperature-sensitive mutants. Germlines of sterile Day 2-3 adult animals were

stained using the OIC1D4 antibody against P granules (green) and DAPI (blue). **(A-A')**

Control animals contain uniform puncta of P granule staining surrounding each nuclei.

(B-F) Late generation fertile *prg-1, rsd-6*, *nrde-2*, *rbr-2*, and *mut-14* animals displayed

relatively normal P granule staining in the germlines. **(B'-F')** *prg-1*, *nrde-2*, *rsd-6, rbr-2,*

and *mut-14* sterile animals all exhibited some degree of germ cell P granule loss.

Figure 3: Disruption of P granules by RNAi or loss of *pgl-1* resulted in similar

phenotypes. (A) G2 wild-type (N2) worms on Empty Vector (EV) or P granule RNAi at

25°C were scored for sterility, embryonic lethality, or fertility. n=16-30 worms per

condition, 2 independent experiments. **(B)** G1 L4 *pgl-1* mutants and G2 L4 P granule

RNAi treated animals stained by DAPI contained mostly large germlines while adults

exhibited a range of germline phenotypes from empty to large. **(C)** G1 *pgl-1* mutants

and G2 P granule RNAi treated animals contained oocytes with increased numbers of

univalents at sterility as measured by DAPI bodies n=151 wild-type, 81 P granule RNAi,

37 *pgl-1*. **(D)** Stained DAPI germlines of sterile G2 L4 and Day 1 *pgl-1* mutants. **(E-H)**

Representative DAPI images of germline and oocyte defects in empty vector treated

animals **(E,F)** and P granule RNAi treated animals **(G,H)** scale bar=10μm. **(I)** *pgl-1* G1

worms at 25°C were either DAPI stained as Day 3 adults (blue) or shifted to 20°C at

Day 1 and DAPI stained at Day 3 (red). **(I)** No sterile day 1 *pgl-1* G1 at 25°C laid

oocytes or progeny (n=35 plates, 350 worms). Sterile day 1 *pgl-1* shifted to 20°C (n=50

plates, 500 worms) gave rise to plates with oocytes and rarely progeny. Plates with

oocytes contained both oocytes and dead embryos. Plates with progeny contained

embryos that hatched to L1. For B,D,I *** $p<0.0001$, * $p=0.007$ Fisher's Chi Square test

with Bonferroni correction, n=total germlines scored, error bars are S.E.M.

Figure 4: Transcriptome profiling of germ granule deficient larvae of Piwi pathway

genome silencing mutants *nrde-1* and *nrde-4*. (A) Log2-fold changes in transposon

transcript abundance in mutant vs wild-type (VW), late vs early generation (LVE) or P

granule RNAi-treated vs control (PGD VC). Transposons upregulated at least two-fold in

nrde-1, *nrde-4* or *prg-1(n4357)* are shown. **(B)** Enrichment of germline-expressed genes

in genes upregulated in *nrde-1* and *nrde-4* mutants. **(C)** Number of genes up and

downregulated in *nrde-1* and *nrde-4* mutants.

Figure 5: Model for P granule dysfunction in Piwi pathway genome silencing

mutants. (**A**) Model of P granules and associated small RNA-mediated genome

silencing proteins. Green circles contain P granule components. Mutants in this paper

that are temperature-sensitive for germ cell immortality have blue outlines. Non-

conditional mutants have red outlines. **(B)** Model for transgenerational sterility in

response to accumulation of Heritable Stress in Piwi pathway genome silencing

mutants, which ultimately compromises P granule integrity (top), and model for

induction of an analogous form of reproductive arrest by acute dysfunction of P granule

components (bottom).

Table 1

Genotype	<20% loss	>20%loss	p-value
wt	32	0	
prg-1 (tm872)	1	7	<.0001
prg-1 (n4357)	3	1	=.11
hrde-1 (tm1200)	0	31	<.0001
rsd-6 (yp11)	0	32	<.0001
nrde-2(gg95)	0	25	<.0001
rbr-2 (tm1231)	0	32	<.0001
nrde-4 (gg131)	17	5	=.008
mut-14 (pk730)	1	6	<.0001
nrde-1 (yp4)	4	6	<.0001

Table 1: Quantification of P granule defects in the germline.

Sterile Piwi pathway genome silencing mutants were stained with P granule antibody.
Germlines arms were semi-qualitatively scored for P granules. If at least 20% of the arm
was missing P granules if was scored as loss and if 80% or more of the arm contained
P granules on DAPI-stained nuclei it was considered normal. Statistics were performed
using Fishers exact test.
842

Supplemental Information

Fig. S1. Related to Figure 2. P granule defects in *prg-1, nrde- 4, hrde-1 and nrde-1*.
Germlines of sterile Day 2-3 adult animals were stained using the OIC1D4 antibody against P

granules (green) and DAPI (blue). (A-A') Control animals contain uniform puncta of P granule staining surrounding each nuclei. (B-E) Late generation fertile *prg-1, nrde-4, hrde-1*, and *nrde-1* animals displayed relatively normal P granule staining in the germlines. (B'-E') sterile *prg-1, nrde-4, hrde-1*, and *nrde-1* germ cell P granules.

	Total Worms Scored	% Fertile	% Sterile
pgl-1 G1 at 25°C	91	22%	78%
pgl-1 G2 at 25°C	71	0%	100%

Table S1: Related to Figure 3. Fertility of *pgl-1* mutants at 25°C. 20°C P0 *pgl-1* mutants were shifted to 25°C and fertility was measured for the G1 generation. Fertile G1 worms were cloned and the fertility of the G2 generation was measured.

gene	*prg-1*	*nrde-1*	*nrde-4*
C06B3.7	4.331128036	1.732445354	2.669978253
dhs-26	3.88587979	3.752818032	1.793937273
bath-45	3.604851731	5.5002102	2.319444032
fbxb-97	3.555301139	3.002992933	3.278539846
irg-2	3.519633623	2.605415087	2.189754742
pud-4	3.226292885	2.956152303	4.407697185
pud-3	3.097392415	3.143142026	4.328678969
R09E10.2	2.745900789	5.100997358	4.251177196
Y53F4B.5	2.636725232	5.821227047	4.162462556
K08D12.6	2.631068263	2.955437046	0.277803871
R03H10.6	2.469258519	4.82098578	3.295830487
Y58A7A.5	2.397260966	1.795895	2.143861408
pud-1.1	2.299685196	2.556376909	3.17781014
T22B7.7	2.276777478	3.01143186	2.207919501
fbxa-224	2.248998131	4.226992478	1.979961678
swt-6	2.144598564	3.153313277	0.277029378
pud-2.1	2.119890425	2.781615959	2.880747229
pud-2.2	2.094123631	2.779337879	2.796566876
pud-1.2	2.080250623	2.559790088	3.000944771
catp-3	2.014970724	2.436698711	2.674464594

Table S2: Related to Figure 4. Genes upregulated in late-generation *prg-1* mutants and in sterile *nrde-1* and *nrde-4* mutant L4 larvae.