1

A reproductive arrest program triggered by defects in Piwi and germ granules

Short title: Germ granule dysfunction phenocopies Piwi sterility

Authors: Maya Spichal[1,2]†, Bree Heestand[1,2,3]†, Katherine Kretovich Billmyre[1,2]†, Stephen Frenk[1,2]† and Shawn Ahmed[1,2,3*]

Affiliations:

[1]Department of Genetics, University of North Carolina, Chapel Hill, NC, USA

[2]Department of Biology, University of North Carolina, Chapel Hill, NC, USA

[3]Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill, NC, USA

†These authors contributed equally to this manuscript

*Corresponding author: shawn@med.unc.edu

Abstract:

In several species, Piwi/piRNA genome silencing defects lead to immediate sterility accompanied by heterochromatin dysfunction and transposon-induced genomic instability, which may cause Piwi mutant sterility. In *C. elegans,* Piwi pathway mutants transmit a heritable stress through germ cells that induces sterility after growth for several generations. We found that sterile Piwi pathway mutant germ cells displayed inconsistent increases in DNA damage but consistently altered perinuclear germ granules that are known to promote fertility. Germ granule dysfunction did not elicit transposon expression but was sufficient to induce multiple phenotypes found in sterile Piwi silencing mutants, including germline atrophy and regrowth. Furthermore, loss of the germ granule component PGL-1 accelerated sterility in response to deficiency for *prg-1*/Piwi. Restoration of germ granule function to sterile *pgl-1* mutants restored their fertility. Together, our results suggest that germ granule defects may promote an adult reproductive arrest phenotype that is responsible for Piwi/piRNA mutant sterility.

Keywords: Germ granules, reproductive arrest, small RNAs, heterochromatin, germ cell immortality, heritable stress, Piwi

Introduction:

Germ cells give rise to the mortal somatic tissues while maintaining themselves in a pristine condition that allows them to be propagated for an indefinite number of generations. This capacity for self-renewal is termed "germ cell immortality", and

functions to protect the germline from forms of damage that can result in sterility

(Smelick and Ahmed, 2005a). Defects in *Caenorhabditis elegans* (*C. elegans)* genes

that promote germline immortality cause a Mortal Germline (Mrt) phenotype, where

robust fertility occurs for several generations but then deteriorates and finally culminates

in complete sterility. The transgenerational sterility of *mrt* mutants implies that their germ

cells may age as they proliferate over the generations such that they accumulate a form

of damage that ultimately evokes sterility. Identifying the transgenerational damage that

accumulates in *mrt* mutant germ cells and the mechanism(s) by which this damage

induces sterility are central questions relevant to understanding the basis of germ cell

immortality.

The initial germline immortality pathway defined in *C. elegans* was promoted by

telomerase, a reverse transcriptase that maintains telomere length and stability by

adding *de novo* telomere repeats to chromosome termini. *C. elegans* mutants that are

defective for telomerase-mediated telomere repeat addition display progressive

telomere erosion. In later generations, some telomeres become critically short and fuse

together, leading to accumulation of end-to-end chromosome fusions (Lowden et al.,

2008; Meier et al., 2006). The precise cause of telomerase mutant sterility remains

uncertain (Karlseder et al., 1999; Nakamura et al., 1998; Sharpless and DePinho, 2007;

Shay and Wright, 2005), but might be the creation of circular chromosomes that are

toxic in meiosis (Nakamura et al., 1998).

Most human somatic cells can proliferate *in vitro* for a limited number of divisions

before they enter a state of irreversible cell cycle arrest termed senescence (Sharpless

and DePinho, 2007; Shay and Wright, 2005). Senescent cells are large and flattened in

comparison to proliferating cells, they express senescence-associated beta-

galactosidase and the tumor suppressor protein p16, and they secrete growth factors

such as IL6 and IL8 cytokines that act in a cell-non-autonomous manner to promote

aging (Tchkonia et al., 2013).The limited proliferative capacity of primary human

somatic cells is called 'replicative aging'. Lack of telomerase is thought to cause

replicative aging in humans, because 1) most primary human somatic cells lack

detectable telomerase activity *in vitro* (Counter et al., 1992; Shay and Wright, 2000), 2)

primary human somatic cells experience progressive telomere erosion as they divide

(Harley et al., 1990), 3) senescent cells display DNA damage response foci at

telomeres that activate p53 to induce senescence (Fagagna et al., 2003; Herbig et al.,

2004), and 4) because senescence does not occur if telomerase activity is restored by

expression of the telomerase reverse transcriptase TERT, which is typically silenced in

human somatic cells (Bodnar et al., 1998). Support for the model that telomere attrition

induces senescence in humans was provided by acute knockdown of the telomere

capping protein TRF2, which results in deprotection of telomeres that are normal in

length, accompanied by DNA damage response foci at telomeres and senescence

(Karlseder et al., 1999). Together, the above data support a model that telomere

uncapping results in DNA damage response that activates p53 to cause senescence in

human somatic cells.

Other forms of stress can accumulate to cause replicative aging of mammalian

somatic cells. For example, expression of the p16 protein induces senescence and

increases by orders of magnitude in mouse and human tissues with age, even though

mouse somatic cells express telomerase (Krishnamurthy et al., 2004, 2006; Sharpless

and DePinho, 2007). Forms of stress that accumulate during replicative aging of

mammalian cells may be revealed by analysis of *C. elegans mortal germline* mutants,

which display transgenerational sterility phenotypes that could reflect 'replicative aging'

of germ cells. In this regard, deficiency for Piwi/piRNAs and related small RNA-

mediated genome silencing proteins like RSD-6 and HRDE-1 results in

transgenerational sterility in the absence of end-to-end chromosome fusions, indicating

that telomere uncapping does not cause sterility of Piwi/piRNA genome silencing

mutants (Billmyre et al., 2019; Buckley et al., 2012; Sakaguchi et al., 2014; Simon et al.,

2014a).

P-M Hybrid Dysgenesis was discovered in the 1970's as a temperature-sensitive

Drosophila sterility defect that was linked to high levels of transposition (Kelleher, 2016;

Kidwell et al., 1977), which occurs in response to lack of piRNA-mediated genome

silencing of specific transposable elements (Brennecke et al., 2008a). Piwi is a

conserved Argonaute protein that interacts with thousands of piRNAs in germ cells to

repress expression of transposons and foreign nucleic acids that represent parasitic

threats to the integrity of the genome (Aravin et al., 2007; Luteijn and Ketting, 2013;

Siomi et al., 2011). Deficiency for Piwi proteins leads to immediate sterility in

Drosophila, zebrafish and mouse, which correlates with increased transposon

expression and DNA damage in sterile animals (Juliano et al., 2011; Kelleher, 2016;

Siomi et al., 2011). Although this implies that transposon-induced genome instability

could cause sterility of Piwi mutants, analysis of *Drosophila* strains with variable levels

of transposons suggested that transposon-induced genome instability may only explain

a minor fraction of P-M hybrid sterility (Srivastav and Kelleher, 2017). Therefore, the

cause of sterility in Piwi mutants remains an unsolved problem in experimental biology.

Deficiency for the *C. elegans* Piwi ortholog *prg-1* compromises germ cell

immortality (Simon et al., 2014b), rather than inducing the immediate sterility phenotype

that is observed in Piwi mutants in some other species. The reason for the

transgenerational delay in the onset of sterility in *prg-1*/Piwi mutants and in related Piwi

pathway mutants of *C. elegans* is not clear, but it is possible that genomic silencing

marks that are induced by the Piwi/piRNA pathway are more robustly maintained in

germ cells of *C. elegans* if the Piwi silencing system is inactivated. The

transgenerational damage that accumulates in *C. elegans* Piwi mutants may therefore

correspond to progressive deterioration of Piwi-mediated heterochromatin.

Heterochromatin dysfunction in *C. elegans* has been associated with increased

levels of transposon and tandem repeat expression, with RNA-DNA hybrid formation,

and with transposon-induced DNA damage (McMurchy et al., 2017; Padeken et al.,

2019; Zeller et al., 2016). However, very low levels of transposition occur in late-

generation *prg-1* mutants, and *C. elegans mutator* mutants, which display high levels of

transposition, do not become sterile if grown at low temperatures, suggesting that

transposition is unlikley to cause transgenerational sterility of *prg-1*/Piwi mutants (Simon

et al., 2014b).

Sterile late-generation *C. elegans prg-1*/Piwi mutants display an unusual

pleiotropic germ cell degeneration phenotype as L4 larvae mature into 1 day old adults,

such that most animals display 'atrophied' germlines that contain a small population of

mitotic germ cells, whereas some germlines are 'empty' and lack germ cells altogether,

some germlines are 'short' such that they have fewer germ cells than normal but

possess both mitotic and meiotic germ cells, and some germlines are sterile but 'normal'

in size such that they are not smaller than germlines of fertile late-generation *prg-1*

mutant siblings (Heestand et al., 2018). Similarly, growth of *C. elegans* Piwi pathway

mutants *rsd-6*, *nrde-2* and *hrde-1* at the restrictive temperature of 25°C for several

generations results in sterile late-generation L4 larvae with germlines that are

predominantly normal in size, but these larvae mature into sterile adults that possess

germline arms with variable sizes, including small, atrophied and empty (Billmyre et al.,

2019; Sakaguchi et al., 2014). The histone 3 lysine 4 demethylase RBR-2 also

promotes germ cell immortality at 25°C and may promote genome silencing

downstream of small RNAs (Alvares et al., 2014).

Atrophied germ lines of sterile *prg-1* mutants can regrow on day 2 of adulthood,

and fertility can be restored to a minor fraction of sterile late-generation *prg-1* mutants

by altering their food source (Heestand et al., 2018; Simon et al., 2014a). This implies

that Piwi mutant sterility is a reversible form of reproductive arrest, also known as Adult

Reproductive Diapause or Reproductive Quiescence, which can be induced in response

to severe environmental stresses such as starvation (Angelo and Van Gilst, 2009;

Padilla and Ladage, 2012; Tatar and Yin, 2001). We therefore proposed that late-

generation sterility of *prg-1*/Piwi mutants may occur in response to accumulation of a

heritable stress that is transmitted by germ cells (Heestand et al., 2018).

Some Argonaute proteins that regulate silencing of the epigenome, like PRG-

1/Piwi, are located in germ granules, which are perinuclear structures that interact with

nuclear pore complexes and have been previously demonstrated to be necessary for

fertility (Fig. 6A) (Updike et al., 2011). *C. elegans* germ granules are termed P granules

based on antibodies that label the P blastomere that generates the germ cell lineage

(Strome and Wood, 1982). Here we study the late-generation sterility phenotype of Piwi

pathway genome silencing mutants and find that it is associated with abnormalities of

germ granules. We show that germ granule dysfunction can induce reproductive arrest

and that germ granule dysfunction interacts with Piwi deficiency to promote sterility.

Results:

Genome instability in Piwi pathway genome silencing mutants

Transgenerational sterility in *C. elegans* telomerase mutants that lack the ability to

maintain sequences that cap chromosome ends is caused by high levels of telomere

fusions and associated genome damage (Meier et al., 2006). Many *mrt* mutants only

become sterile when grown at the non-permissive temperature 25°C (Ahmed and

Hodgkin, 2000), some of which are deficient for Piwi pathway-mediated genome

silencing (Ashe et al., 2012; Buckley et al., 2012; Burkhart et al., 2011; McMurchy et al.,

2017; Sakaguchi et al., 2014; Smelick and Ahmed, 2005b; Spracklin et al., 2017).

However, the deficiency for the Piwi Argonaute protein PRG-1 or the nuclear RNA

interference (RNAi) proteins NRDE-1 or NRDE-4 results in progressive sterility at any

temperature (Batista et al., 2008; Buckley et al., 2012; Burkhart et al., 2011; Das et al.,

2008; McMurchy et al., 2017; Simon et al., 2014b). We asked if DNA damage is directly

associated with the sterility of Piwi pathway mutants by comparing fertile late-generation

Piwi pathway mutants with late generation siblings that were sterile, using both

temperature-sensitive Piwi pathway mutants that become sterile at 25°C and non-

conditional mutants that become sterile when grown at 20°C.

Although DNA damage has been suggested not to contribute significantly to the

sterility of *prg-1* mutants (Simon et al., 2014b), DNA damage signaling in sterile *prg-1*

mutant animals has not been previously studied. We therefore examined early and late-

generation *prg-1* mutants grown at 20°C across several generations for activation of the

DNA damage response, as detected by an antibody to phosphorylated S/TQ, a protein

modification that is created by two sensors of DNA damage, ATM and ATR, which are

phosphatadylisonisol-3-kinase like protein kinases (Kim et al., 1999; Vermezovic et al.,

2012). We observed a significant increase in the fraction of germ cells with upregulated

DNA damage response for late-generation fertile and sterile *prg-1* mutants (~3- to 6-

fold) when compared to either wild type controls or to early-generation *prg-1* mutants

(Fig. 1A,B). However, the fraction of germ cells with an activated DNA damage

response was not consistently elevated in sterile *prg-1*/Piwi mutants in comparison to

fertile late-generation siblings (Fig. 1B). We also tested temperature-sensitive Piwi

pathway mutants and found that the wild-type DNA damage response was upregulated

by growth at 25°C but that this level was not further elevated in germlines of sterile late-

generation *rsd-6, nrde-2* or *hrde-1* mutant adults (Fig. 1B,C).

Piwi pathway genome silencing mutants exhibit germ granule defects at sterility

If secondary siRNAs are depleted from *prg-1* mutants and then restored by

crossing distinct *prg-1; mutator* double mutants together, this causes immediate sterility

for *prg-1* mutant F1 cross progeny that possess germ cells with germ granule defects

206
Figure 1: DNA Damage Response is variable in Piwi pathway genome silencing

mutants prior to and at sterility. (A) Staining of pS/TQ in wild type, *prg-1(n4357) and*

prg-1(tm876) mutants. pS/TQ(grey and yellow) and DAPI(blue) **(B)** Quantification of

pS/TQ staining in 20°C wild type and *prg-1* mutants at different levels of fertility. **(C)**

Quantification of pS/TQ staining in 25°C wild type and small RNA genomic silencing

mutants at sterility. P-values were obtained by comparing sample groups in a Mann-

Whitney U test (p-value: P > 0.05 = n. s., P < 0.05 = *, P = <0.01 = **, P =< 0.001 = ***

compared to wild type).

and a range of adult germline sizes that are similar to those of sterile late-generation

Piwi pathway mutants (de Albuquerque et al., 2015; Phillips et al., 2015). Furthermore,

combined loss of the epigenomic regulators SPR-5 and LET-418, or loss of H3K9

methylation modifier SET-2 and the nuclear RNAi pathway, results in immediate sterility

that is associated with disruption of germ granules (P granules in *C. elegans*) (Käser-

Pébernard et al., 2014; Robert et al., 2014). Although most Piwi pathway mutants do not

display severe fertility defects until the generation of sterility, late-generation *prg-1*/Piwi

mutants display very small brood sizes for many generations prior to sterility (Heestand

et al., 2018). By immunofluorescence with a P granule antibody, we observed that P

granules were frequently strongly reduced or absent in nuclei of sterile late-generation

prg-1 mutant germ cells (hereafter referred to as marked P granule dysfunction) (Fig.

2B, S1, Table 1), which was less common in fertile *prg-1(n4357)* mutant siblings while

prg-1(tm872) mutants showed more consistent P granule defects right before and at

sterility (Fig. 2B-B', Table 1). Consistently, growth of the temperature-sensitive Mrt

mutants *rsd-6*, *nrde-2*, *hrde-1* and *rbr-2* at the restrictive temperature of 25°C resulted in

sterile late-generation animals with marked P granule dysfunction, whereas their fertile

late-generation siblings did not exhibit pronounced P granule defects (Fig. 2C, D, E, C',

D', E', S1, Table 1).

Germ line atrophy occurs for many but not all sterile Piwi pathway mutants as L4

larvae develop into adults (Billmyre et al., 2019; Heestand et al., 2018). Large-scale

Figure 2: At sterility, partial loss of P granule staining occurred in *prg-1* and temperature-sensitive mutants. Germlines of sterile Day 2-3 adult animals were stained using the OIC1D4 antibody against P granules (green) and DAPI (blue). **(A)** Control animals contain uniform puncta of P granule staining surrounding each nuclei.(B) *mut-14* mutants display an immediate sterility when shifted to 25˚C for one generation and display P granule abnormalities **(C-E)** Late generation fertile *prg-1,nrde-2* and *rbr-2* animals displayed relatively normal P granule staining in the germlines. **(C'-E')** *prg-1*, *nrde-2 and rbr-2* sterile animals all exhibited some degree of germ cell P granule loss.

apoptosis contributes to but is not essential for this germ cell degeneration phenotype (Heestand et al., 2018), and P granules have been shown to vanish in cells that undergo apoptosis (Min et al., 2016; Sung et al., 2017). However, we observed marked P granule dysfunction in germlines of sterile late-generation *nrde-2* and some *rsd-6* mutant adults that were wild type in size and had not undergone significant amounts of apoptosis (Fig. 2D, D'; Fig. S1F, F'). *nrde-1* and *nrde-4* mutant animals displayed a sterility phenotype where many sterile germline arms displayed P granule defects (90% and 87% respectively) (Fig. S1, Table 1). P granule defects in *nrde* mutants included P granule dissociation from the nuclear envelope into the cytoplasm and P granule aggregations as well as P granule loss. Therefore, dysfunction of *nrde-1* and *nrde-4*, which act downstream of multiple small RNA genome silencing pathways (Fig. 6A), might influence P granules differently than we observed in the germline arms of most sterile Piwi pathway mutants.

261   *C. elegans* P granules are adjacent to Mutator bodies, which house Mutator

262 proteins that recruit RNA-dependent RNA polymerase to create secondary effector

263 siRNA populations in response to primary siRNAs (Phillips et al., 2012). *mutator*

264 mutants are dysfunctional for secondary siRNA biogenesis and remain fertile indefinitely

265 at low temperatures (Simon et al., 2014b), but display an immediate highly penetrant

266 fertility defect at 25°C that could be caused by dysfunction of Piwi pathway-mediated

267 heterochromatin (Ketting et al., 1999). We examined germlines of sterile *mut-14(pk738)*

268 mutant adults at the restrictive temperature of 25°C and observed pronounced P

269 granule defects in 66.67% of animals with no offspring and 100% P granule defects in

270 animals that laid eggs that could not develop into adults, here referred to as 'fertile'(Fig.

271 2B', Table 1).

273 Germ granule dysfunction phenocopies sterile Piwi pathway genome silencing

274 mutants

275 We addressed the significance of P granule dysfunction in sterile late-generation small

276 RNA genome silencing mutants using an RNA interference clone that simultaneously

277 targets four P granule subunits *pgl-1*, *pgl-3*, *glh-1*, *glh-4* (Fig. 6A) (Knutson et al., 2017;

278 Updike et al., 2014). This resulted in many second generation (G2) sterile adults at

279 25°C (80%) (Fig. 3A). The germlines of G2 P granule RNAi L4 larvae were normal in

280 size in comparison to control worms. However, a pronounced germ cell atrophy

281 phenotype occurred as many P granule-depleted animals matured into sterile adults,

282 resulting in a significant change in their germline profile (Fig. 3B, p=3.93E-24).

Figure 3: Disruption of P granules by RNAi or by deficiency for *pgl-1* results in

similar phenotypes. (A) G2 wildtype (N2) worms on Empty Vector (EV) or P granule

RNAi at 25°C were scored for sterility, embryonic lethality, or fertility. n=16-30 worms

per condition, 2 independent experiments. **(B)** Germline phenotypes of G1 L4 *pgl-1*

mutants and G2 L4 P granule RNAi treated animals stained by DAPI. **(C)** G1 *pgl-1*

mutants and G2 P granule RNAi treated animals contained oocytes with increased

numbers of univalents at sterility as measured by DAPI bodies n=151 wild type, 81 P

granule RNAi, 37 *pgl-1*. **(D)** Stained DAPI germlines of sterile G2 L4 and Day 1 *pgl-1*

mutants. **(E-H)** Representative DAPI images of germline and oocyte defects in empty

vector treated animals **(E,F)** and P granule RNAi treated animals **(G,H)** scale

bar=10µm. **(I)** *pgl-1* G1 worms at 25°C were either DAPI stained as Day 3 adults (blue)

or shifted to 20°C at Day 1 and DAPI stained at Day 3 (red). **(I)** No sterile day 1 *pgl-1*

G1 at 25°C laid oocytes or progeny (n=35 plates, 350 worms). Sterile day 1 *pgl-1*

shifted to 20°C (n=50 plates, 500 worms) gave rise to plates with oocytes and rarely

progeny. Plates with oocytes contained both oocytes and dead embryos. Plates with

progeny contained embryos that hatched to L1. For B,D,I *** p<0.0001, * p=0.007

Fisher's Chi Square test with Bonferroni correction, n=total germline arms scored, error

bars are S.E.M.

Furthermore, oocyte univalents were previously observed in sterile but not fertile late-

generation *rsd-6* mutants (Sakaguchi et al., 2014), and 45.5% of oocytes of sterile P

granule depleted adults contained 7-12 univalents (Fig. 3C, E-H), consistent with a

previous report (Billmyre et al., 2019).

The germline phenotypes observed in P granule-depleted worms mimicked those

of sterile *prg-1*/Piwi mutant adults (Fig. 3A-C), which can regrow their atrophied

germlines and become fertile (Heestand et al., 2018). We tested the hypothesis that

germ granule dysfunction might be sufficient to cause reproductive arrest using a

mutant deficient for the P granule component *pgl-1*, which remains fertile at low

temperatures but displays a pronounced sterility phenotype if shifted to the restrictive

temperature of 25°C (Kawasaki et al., 1998). Most first generation (G1) *pgl-1* animals at

25°C displayed normal germlines at the L4 larval stage, but there was a significant shift

in the germline profiles with 52% of day 1 adults showing germline atrophy (Fig. 3B,

p=1.04E-35), similar to P granule RNAi knockdown and sterile *prg-1* mutants (Heestand

et al., 2018). Furthermore, when oocytes of sterile *pgl-1* mutant adults were scored, we

found that 56.7% of oocytes had univalents, in agreement with the oocyte univalents

observed in response to P granule depletion by RNAi (Fig. 3C).

Some G1 progeny of *pgl-1* mutants whose parents were shifted from 20°C to

25°C were fertile and gave rise to G2 25°C animals that were uniformly sterile. 25°C G2

pgl-1 mutant L4 larvae had germline profiles that displayed more severe germ cell

proliferation defects than those of G1 L4 larvae, and day 1 adults showed a striking 54%

increase in empty germlines (Fig. 3D, p=2.44E-05). Therefore, germlines of G1 progeny

of *pgl-1* mutant mothers that were shifted to the restrictive temperature of 25°C for a

single generation mimicked the germ cell degeneration phenotypes of sterile Piwi

pathway mutants as L4 larvae matured into adults, whereas 25°C G2 generation *pgl-1*

mutant larvae displayed more pronounced germ cell proliferation defects.

We asked if the fertility of sterile 25°C G1 *pgl-1* mutants could be restored by

shifting them to the permissive temperature of 20°C. Sterile G1 *pgl-1* mutants that

lacked embryos were shifted to 20°C on day 1 of adulthood and their germlines were

scored after 48 hours, which revealed a significant change in their germline profiles

(p=0.007), with a shift from atrophied germlines towards large germlines when

compared to sterile G1 *pgl-1* sibling controls that remained sterile at 25°C (Fig. 3I). In

agreement with the regrowth of some germline arms, we observed that some sterile G1

pgl-1 mutant adults that were shifted to 20°C for several days had laid oocytes and

dead embryos, and that a few even gave rise to progeny (Fig. 3J). In contrast, sterile

pgl-1 mutant control adults that were kept at 25°C never gave rise to oocytes, dead

embryos or living progeny (Fig. 3J).

Transcriptional consequences of germ granule dysfunction and small RNA

genome silencing mutant sterility

The sterility of Piwi/piRNA mutants correlates with expression of transposons and

associated genome instability (Brennecke et al., 2008b; Juliano et al., 2011; Kelleher,

2016; Siomi et al., 2011). We therefore studied RNA from the nuclear RNAi defective

mutants *nrde-1*, *nrde-2* and *nrde-4* (Buckley et al., 2012; Guang et al., 2008, 2010) that

have Mortal Germlines at 25°C (Burkhart et al., 2011). However, at 20°C *nrde-1* and

nrde-4 mutants become progressively sterile but *nrde-2* mutants remain fully fertile

indefinitely (Burkhart et al., 2011). *nrde-1* and *nrde-4* mutants have relatively large

brood sizes at sterility at 20°C, such that large cohorts of L4 larvae that are poised to

become sterile can be collected. In contrast, *prg-1* mutants develop very low brood

sizes in late generations (Heestand et al., 2018), which makes it difficult to obtain many

synchronous animals that are poised to become sterile. We therefore prepared RNA

from early-generation and sterile-generation *nrde-1* and *nrde-4* mutants at 20°C and

compared this with RNA from wild type and *nrde-2* mutant controls.

We found that sterile generation *nrde-1* and *nrde-4* mutant L4 larvae showed

strong upregulation of similar transposon classes in comparison to early generation

nrde-1 or *nrde-4* mutant L4 larvae or to wild type controls (Fig. 5A). We identified 18

transposon classes that were upregulated at least two-fold in both *nrde-1* and *nrde-4*,

13 of which were CER retrotransposons, which is consistent with previous

transgenerational analysis of *hrde-1* mutants (Ni et al., 2016). However, we also found

that *nrde-2* mutant controls that do not become sterile at 20°C displayed strong

upregulation of similar transposon loci, even in early generations. Of the 18 transposons

upregulated in both *nrde-1* and *nrde-4*, 17 were also upregulated in *nrde-2* compared

with wild type. Previous work reported that distinct classes of transposons are

upregulated in *prg-1* and *nrde-2* mutants (Das et al., 2008; McMurchy et al., 2017; Ni et

al., 2016; Simon et al., 2014b). Consistently, we analyzed published RNA-seq data from

prg-1(n4357) mutant animals (McMurchy et al., 2017) found that MIRAGE1 was the only

transposon upregulated in both *nrde-1*, *nrde-4* and *prg-1* mutants (McMurchy et al.,

2017) (Fig. 4A). Therefore, although deficiency for *prg-1* and *nrde-1/nrde-4* disrupts

small RNA-mediated genome silencing and compromises germ cell immortality, these

genes have distinct functions with regards to small RNA-mediated regulation of the

genome.

We compared gene expression data from sterile generation *nrde-1* or *nrde-4*

mutant L4 larvae with late-generation *prg-1* mutants (McMurchy et al., 2017) and with

wild type or *nrde-2* mutant control L4 larvae in order to identify common genes whose

expression is specifically induced in *nrde* mutant L4 larvae that are poised to become

sterile and that might explain the P granule abnormalities at sterility. Late-generation

Figure 4: Transcriptome profiling of germ granule deficient larvae of Piwi pathway

genome silencing mutants *nrde-1* and *nrde-4*. (A) Log2-fold changes in transposon

transcript abundance in mutant vs wild type (VW), late vs early generation (LVE) or P

granule RNAi-treated vs control (PGD VC). Transposons upregulated at least two-fold in

nrde-1, *nrde-4* or *prg-1(n4357)* are shown. **(B)** Enrichment of germline-expressed genes

in genes upregulated in *nrde-1* and *nrde-4* mutants. **(C)** Number of genes up and

downregulated in *nrde-1* and *nrde-4* mutants.

prg-1 mutant and sterile generation *nrde-1* and *nrde-4* mutant gene expression was

very different from wild type or *nrde-2* mutant controls RNA. Spermatogenic genes were

over-represented among genes significantly upregulated in sterile generation *nrde-1*

and *nrde-4* mutant larvae (p = 1e-15, Chi Square test for both mutants), which is

consistent with what has been previously reported for *spr-5* mutants that become

progressively sterile (Katz et al., 2009). However, there was no general tendency of an

up- or down-regulation of any gene category for *prg-1*, *nrde-1* and *nrde-4* mutants. Only

11 genes were significantly upregulated in all three mutants, and there were no

commonly downregulated genes, which suggests that *prg-1* and *nrde-1/nrde-4* mutants

have fundamentally different gene expression profiles at or near sterility (Fig. 4C). We

identified a set of 20 genes that were significantly upregulated at least fourfold in *prg-1*

and either *nrde-1* or *nrde-4*, which included *pud* (protein upregulated in dauer) genes

that might be consistent with a role for DAF-16/Foxo in *prg-1* mutant sterility (Table S2)

(Buckley et al., 2012; Simon et al., 2014a). However, we could not find consistent

changes in any gene whose knockdown perturbs P granule structure (Updike and

Strome, 2010), which suggests that altered germ granule structure of sterile Piwi

pathway mutants is a post-transcriptional response.

RNA-seq data sets were previously obtained for germlines of P granule defective

L4 larvae that were poised to become sterile, but gene expression was not significantly

different from wild type (Knutson et al., 2017). We compared RNA from germlines of P

granule defective L4 larvae (Knutson et al., 2017) with RNA from sterile generation

nrde-1 and *nrde-4* L4 larvae and found no genes that were up or downregulated in both

nrde-1/nrde-4 and P granule defective larvae. Moreover, only two of 18 transposons

upregulated in *nrde-1/nrde-4* were also upregulated in P granule defective L4 larvae

(Fig. 4A). We conclude although acute dysfunction of P granules based on

simultaneous knockdown of four P granule components leads to germ line degeneration

phenotypes that mimic those of sterile generation Piwi pathway mutants, it does not

cause immediate overt effects on expression of heterochromatic segments of the

genome or on genes in a manner that mimics the transcriptomes of Piwi pathway

mutants that are poised to become sterile. Interestingly, the differences between *nrde-*

1/nrde-4 RNA data and *prg-1* also correlate with differences that we observed in P

granule abnormalities between these mutants (Figure 2, S1). This indicates that the

sterility observed in nrde and PIWI mutants causes P granule defects resulting from

different molecular pathways.

423   Deficiency for *pgl-1* enhances *prg-1*/Piwi mutant sterility

424   Given that deficiency for *pgl-1* or RNAi knockdown of P granule components at 25°C

425   mimics Piwi mutant sterility, we performed an RNAi screen for P granule components

whose knockdown induces sterility in early-generation *prg-1* mutants or in *pgl-1* mutants
grown at the permissive temperature of 20^{o}C. We knocked down several P-granule
components for 10 generations in wild type control animals and in strains deficient for
either *prg-1* or the P-granule component *pgl-1* (Fig 5A). Several RNAi clones led to
immediate sterility for all genotypes tested, and we also identified P granule
components whose knockdown induced sterility the *pgl-1* mutant. Specifically, we found
that RNAi of *prg-1*/Piwi induced sterility in *pgl-1* mutants in generation F4 (Fig 5B). We
studied the germline phenotypes of sterile *pgl-1* mutants treated with *prg-1* RNAi and
observed that they had a mixture of empty, atrophied and short germlines (Fig. 5B, D).

To confirm the interaction of *pgl-1* mutation with RNAi knockdown of *prg-1*, we
used outcrossed mutations in *prg-1* and *pgl-1* to create *prg-1; pgl-1* double mutants and
compared them to the single mutant outcrossed controls. We found that *prg-1* (20/20)
and *pgl-1* (20/20) single mutant controls were fertile for 10 generations when grown at
20^{o}C. In contrast, *prg-1; pgl-1* double mutant F2 adults were fertile but gave rise to F3
animals that were almost all sterile (41/45) (Fig 5C). We stained sterile *prg-1; pgl-1*
double mutant F3 with DAPI and found that on Day 1 62.9% of germlines were either
atrophied or empty (Fig 5B).

Discussion:

Previous work has demonstrated that Piwi mutant sterility correlates with transposon
expression, transposition and genomic instability in several species (Juliano et al., 2011;
Kelleher, 2016; Siomi et al., 2011). We studied DNA damage signaling in *C. elegans*
Piwi pathway mutants and found that this was not consistently increased in sterile

Figure 5: Deficiency for *pgl-1* results in rapid sterility of *prg-1* mutants. (A)

Germline immortality assay with wild type, *prg-1(pk2298), prg-1(n4357)* and *pgl-*

1(bn101) on different P-granule RNAi clones. 2 replicates of 3 L1 larvae were

transferred every week or after 2 generations and plates were scored as sterile when no

more than 3 worms were present after 1 week. Light red indicates sterility for 1 of 2

replicate plates while dark red indicates sterility for both replicates. When one plate

became sterile worms from the second plate were used for 2 replicates. (B) Germline

sizes of wild type and *pgl-1(bn101)* on *prg-1* RNAi after 4 generations and of *prg-1; pgl-*

1 double mutant F3 animals. (C) *prg-1; pgl-1* double mutants display marked sterility in

the F3 generation.(D) Germline comparison of wild type and *pgl-1(bn101)* on *prg-1*

RNAi after 4 generations.

animals in comparison to fertile late-generation siblings. These data complement

previous observations that suggest that DNA damage induced by transposition is

unlikely to cause the transgenerational sterility of *prg-1*/Piwi mutants (Simon et al.,

2014b). A recent study revealed that defects in H3K9 methylation caused by *met-2* and

set-25 histone methyltransferases cause R-loops and mutations at tandem repeats in

the *C. elegans* genome, reduced fertility, and synthetic sterility with the ATR homolog

atl-1 (Padeken et al., 2019) However, the sterility of the Piwi mutants we studied was

not associated with high levels of ATR signaling, suggesting that Piwi sterility may be

distinct from the reduced fertility caused by loss of H3K9me2 and me3 (Padeken et al.,

2019).

We discovered pronounced germ granule defects in sterile but to a lesser extent

in fertile late-generation Piwi pathway mutants and tested their significance by

disrupting germ granule proteins (Fig. 3). We found that shifting a *pgl-1* germ granule

mutant to restrictive temperature, led to germline degeneration phenotypes that

phenocopied those of sterile generation Piwi pathway mutants (Heestand et al., 2018).

We also found that restoring germ granule function sterile *pgl-1* mutant adults, by

shifting them to the permissive temperature of 20°C, resulted in growth of the germline

and fertility (Fig. 3I-J).

Although sterility in response to germ granule dysfunction has previously been

considered to be a pathological condition that results in a germline-to-soma

transformation (Knutson et al., 2017), our results indicate that germ granule dysfunction

can induce reproductive arrest in adults (Fig. 3, 6C). As environmental stresses such as

starvation can also evoke reproductive arrest (Angelo and Van Gilst, 2009; Padilla and

Ladage, 2012), we speculate that germ granule dysfunction might represent a general

mechanism for orchestrating stress-induced reproductive arrest.

Sterile late-generation *prg-1*/Piwi pathway mutants display a broad range of

germline phenotypes that are observed in response to germ granule dysfunction,

including germ cell atrophy, empty germlines, univalent chromosomes in oocytes,

germline regrowth and reproductive arrest (Heestand et al., 2018). Together with the

germ granule defects observed in sterile Piwi pathway mutants, our results suggest that

Piwi mutant sterility is a form of reproductive arrest that may be caused by germ granule

dysfunction (Fig. 6B). If the germ granule defects in sterile Piwi pathway mutants were a

consequence rather than a cause of Piwi mutant sterility, then we would have expected

that disrupting germ granule function would lead to a distinct form of sterility that did not

bear hallmarks of Piwi mutant sterility like germ cell atrophy and reproductive arrest

(Fig. 6C). Another reason that germ granule defects of sterile Piwi pathway mutants are

unlikely to be a consequence of Piwi mutant sterility is that we observed some sterile

Piwi pathway mutants with large germlines that did not display obvious P granule

defects (Fig. 2, S1). We also found that deficiency for the P granule component *pgl-1*

dramatically accelerated the sterility in response to deficiency for *prg-1*/Piwi, indicating

that these proteins function redundantly to promote germ cell immortality. Knockdown of

the germ granule components *gld-1* and *pgl-3* also induced F4 generation sterility for

pgl-1 mutants, which suggests a common function PRG-1, GLD-1 and PGL-3 in

promoting transgenerational fertility in the context of germ granules that is consistent

with their known physical interactions (Akay et al., 2013; Chen et al., 2016)

Although P granules and associated Mutator bodies contain proteins like PRG-

1/Piwi and a host of other factors that promote small RNA biogenesis (Billi et al., 2012),

we did not observe strong expression of heterochromatic transposons that are normally

silenced by small RNAs in L4 larvae that are poised to become sterile because of germ

granule dysfunction. This observation is consistent with the premise that small RNA-

induced genomic silencing marks can be maintained in the absence of small RNA

biogenesis.

Previous analysis showed that crosses that restored secondary siRNAs to *prg-1*

mutants created sterile F1 *prg-1/*Piwi mutant progeny that possessed adult germlines

with sizes analogous to those of sterile late-generation *prg-1* mutants and in disruption

of P granules (de Albuquerque et al., 2015; Heestand et al., 2018). This led to the

hypothesis that misrouting of small RNAs associated with pro- and anti-silencing small

RNA pathways could be the cause of sterility in Piwi pathway genome silencing mutants

(de Albuquerque et al., 2015; Phillips et al., 2015), which agrees with our observations

of germ granule abnormalities in sterile but not fertile late-generation Piwi pathway

mutants. Consistently, sterile mouse mutants with defects in piRNA biogenesis have

been reported to display germ granule defects (Chuma et al., 2006; Zheng and Wang,

2012). We suggest that germ granule dysfunction could explain the sterility observed in

P-M hybrid dysgenesis and the immediate sterility phenotype of Piwi mutants in several

species (Juliano et al., 2011; Kelleher, 2016; Siomi et al., 2011).

When RNA from sterile generation *nrde-1* and *nrde-4* mutant L4 larvae and late-

generation *prg-1*/Piwi mutants was examined, we failed to observe consistent depletion

of any gene whose knockdown is known to disrupt P granule formation (Updike and

Strome, 2010), suggesting a post-transcriptional reproductive arrest mechanism (Fig.

4B). We propose that transgenerational dysfunction of the Piwi/piRNA pathway in *C.*

elegans results in heterochromatin defects and transcription of 'stressful RNA' from one

or more genomic loci in later generations, which triggers a reproductive arrest program

that perturbs germ granule structure (Fig. 6B). A role for defective heterochromatin as a

cause of Piwi mutant sterility is supported by the *C. elegans* mutants *spr-5* and *rbr-2*,

which encode H3K4 demethylases that function within nuclei downstream of small

RNAs to promote genomic silencing (Fig. 2). *spr-5* and *rbr-2* mutants display

transgenerational sterility phenotypes that mirror those of Piwi pathway silencing

mutants (Alvares et al., 2014; Billmyre et al., 2019).

With regards to the model of telomerase deficiency as a cause of replicative

aging in mammalian cells, we suggest that germ granule dysfunction in Piwi mutants

may be analogous to telomere uncapping in the context of senescence (Sharpless and

DePinho, 2007; Shay and Wright, 2000). Acute germ granule dysfunction is sufficient to

mimic Piwi mutant sterility (Fig. 3, 6C), similar to knockdown of the TRF2 telomere

capping protein being sufficient to induce acute telomere uncapping and senescence

(Karlseder et al., 1999). In the case of replicative aging of mammalian cells, progressive

telomere erosion is the source of damage that ultimately induces telomere uncapping.

In the case of transgenerational Piwi pathway mutant sterility, it remains unclear what is

the precise source of 'heritable stress' that accumulates over generations to induce

sterility, although this is likely to be related to transgenerational desilencing of

heterochromatin/transposons (Fig. 5A).

Our results establish a novel framework for considering inheritance in the context

of epigenetic stress and how this can evoke sterility. We suggest that Piwi mutant

sterility is a response to epigenome dysfunction that arrests germ cell development in

an effort to protect the genome or epigenome in times of stress. Transposons and

viruses are likely to be in a continuous arms race between parasite and host, where

components of germ granules that promote Piwi pathway genome silencing and protect

the host are likely to be targeted (Updike and Strome, 2010). It may therefore be

reasonable to hypothesize the evolution of a reproductive arrest mechanism in the

context of an attack on the Piwi silencing system that is predicated on disrupting the

integrity of germ granules themselves.

Materials and Methods

Strains

All strains were cultured at 20°C or 25°C on Nematode Growth Medium (NGM) plates

seeded with *E. coli* OP50. Strains used include Bristol N2 wild type, *hrde-1(tm1200) III,*

nrde-1(yp4) III, nrde-2(gg95) II, nrde-4 (gg131) IV, prg-1 (n4357) I, prg-1 (tm872) I, prg-

Figure 6: Model for P granule dysfunction in Piwi pathway genome silencing

mutants. (**A**) Model of P granules and associated small RNA-mediated genome

silencing proteins. Blue circles contain P granule components used in this paper.

Mutants used this paper are in lavender. Argonaute proteins are hexagons. Non-

conditional mutants have red outlines. **(B)** Model for transgenerational sterility in

response to accumulation of Heritable Stress in Piwi pathway genome silencing

mutants, which ultimately compromises P granule integrity. **(C)** Model for induction of

reproductive arrest by acute dysfunction of P granule components.

1(pk2298) I, pgl-1(bn101) IV, mut-14 (pk738) V, rbr-2(tm1231) IV, HIS-72::mNeonGreen

(LP230)*.*

<u>RNAi Assay</u>

Feeding RNAi plates harboring host bacteria HT115(DE3) engineered to express *"*quad*"*

dsRNA (targeting *pgl-1*, *pgl-3*, *glh-1*, *glh-4*) were obtained from Susan Strome (Knutson

et al., 2017; Updike et al., 2014). L1 larvae were placed onto freshly prepared feeding

RNAi plates with dsRNA induced by 1 mM IPTG (isopropyl-β-D(-)-

thiogalactopyranoside) and were transferred after 1 generation at 25°C and collected at

G2 adults as described in Knutson et al 2017 (Knutson et al., 2017) for DAPI staining,

oocyte and germline analysis.

For RNAi screen, L1 worms containing the *his-72*:: *mNeonGreen* transgene were

arrested in M9 media and then pipetted at a density of 20 worms per plate and

maintained at 20°C. At the L4 stage and during adulthood worms were scored for

germline size using Leica M205 fluorescence microscope.

For the P-granule germline immortality assay 3 L1 worms were transferred every week

on two replicate plates per condition at 20˚C. Plates were counted as sterile when no

more than 3 worms were present after one week. Partial sterility was counted when one

out of two plates became sterile while both replicate plates were sterile for full sterility.

DAPI staining and Scoring

DAPI staining was performed as previously described (Ahmed and Hodgkin, 2000).

Briefly, L4 larvae were selected from sibling plates and sterile adults were singled as

late L4s, observed 24 hours later for confirmed sterility, and then stained 48 hours after

collection. Univalents were scored by counting DAPI bodies in the -1 to -4 oocytes.

Germline profiles were scored using the method outlined in Heestand et al 2018

(Heestand et al., 2018).

Statistical Analysis

Statistical analysis was performed as previously described (Heestand et al., 2018).

Briefly, statistical analysis was performed using the R statistical environment (R Core

Team, 2013). For germline phenotypes, contingency tables were constructed and

pairwise Chi Square tests with Bonferroni correction was used to determine significant

differences in in germline phenotype distributions. The significance of the DNA damage

response analysis was determined by a Kruskal-Wallis test, followed by a Mann-

Whitney test between individual samples. P-values were adjusted with Bonferroni

correction when multiple comparisons were performed.

RNA extraction and sequencing

Animals were grown at 20°C on 60 mm NGM plates seeded with OP50 bacteria. RNA

was extracted using Trizol (Ambion) followed by isopropanol precipitation. Library

preparation and sequencing was performed at the UNC School of Medicine High-

Throughput Sequencing Facility (HTSF). Libraries were prepared from ribosome-

depleted RNA and sequenced on an Illumina Hiseq 2500.

RNA-seq Analysis

The following publicly available RNA-seq datasets were download from the Gene

Expression Omnibus (https://www.ncbi.nlm.nih.gov/geo/): GSE92690 (P granule RNAi

experiment) and GSE87524 (*prg-1* experiment). Adapter trimming was performed as

required using the bbduk.sh script from the BBmap suite (Bushnell) and custom scripts.

Reads were then mapped to the *C. elegans* genome (WS251) using hisat2 (Kim et al.,

2013) with default settings and read counts were assigned to protein-coding genes

using the featureCounts utility from the Subread package (Liao et al., 2014) . For

multimapping reads, each mapping locus was assigned a count of 1/n where n=number

of hits. Differentially expressed genes were identified using DESeq2, and were defined

as changing at least 2-fold with FDR-corrected p-value < 0.01. For analysis of

transposon RNAs, reads were mapped to the *C. elegans* transposon consensus

sequences downloaded from Repbase (http://www.girinst.org/repbase/) with bowtie

(Langmead et al., 2009) using the options -M 1 -v 2. Transposons with fewer than 10

counts in each sample were excluded from further analysis. Counts were normalized to

the total number of mapped reads for each library for the prg-1 dataset, or to the total

number of non-ribosomal mapped reads for all other datasets. A pseudocount of 1 was

added to each value to avoid division by zero errors. Analysis of sequencing data and

plot creation was performed using the R statistical computing environment (R Core

Team, 2013).

Accession numbers: RNA-seq data reported in this study have been submitted to the

GEO database and will be available at the time of publication.

Immunofluorescence

Adult hermaphrodites raised at 20°C or 25°C were dissected in M9 buffer and flash

frozen on dry ice before fixation for 1 min in methanol at -20°C. After washing in PBS

supplemented with 0.1% Tween-20 (PBST), primary antibody diluted in in PBST was

used to immunostain overnight at 4 °C in a humid chamber. Primaries used were 1:50

OIC1D4 (Developmental Studies Hybridoma Bank). Secondary antibody staining was

performed by using a Cy3 donkey anti-mouse or Cy-5 donkey anti-rabbit overnight at

4°C. All images were obtained using a LSM 710 laser scanning confocal and were

taken using same settings as control samples. Images processed using ImageJ.

Marked P granule dysfunction was scored by eye as a loss of greater or less than 20%

loss of P granules.

<u>DNA Damage Assay</u>

Worms that were close to sterility were isolated and defined as sterile if they did not

have any offspring as day 3 adults at 20°C or day 2 adults at 25°C. Fertile siblings of

sterile worms were defined as 'close to sterility'. The presence of the DNA damage

response was determined by using a phospho-specific antibody targeting the

phosphorylated consensus target site of ATM and ATR kinases (pS/TQ) (Cell Signaling

Technology). This antibody has only been shown to stain a DNA damage response in

the germline and was used as previously described (Vermezovic et al., 2012).

Acknowledgments: We thank members of the Ahmed lab for critical reading of the

manuscript and Jacinth Mitchell for wild type control, *nrde-1* and *nrde-2* RNA-seq data.

Some strains were provided by the CGC, which is funded by NIH Office of Research

Infrastructure Programs (P40 OD010440). This study was supported by NIH grants F32

GM120809 (K.B) and RO1 GM083048 (S.A). M.S. was supported by a DFG postdoc

fellowship.

Competing interests: Authors declare no competing interests.

Author contributions: K.B., B.H., M.S. and S.F. performed experiments. S.F. analyzed

the data. K.B., B.H., M.S., S.F. and S.A. wrote manuscript.

References:

Ahmed, S., and Hodgkin, J. (2000). MRT-2 checkpoint protein is required for germline

immortality and telomere replication in C. elegans. Nature *403*, 159–164.

Akay, A., Craig, A., Lehrbach, N., Larance, M., Pourkarimi, E., Wright, J.E., Lamond, A.,

Miska, E., and Gartner, A. (2013). RNA-binding protein GLD-1/quaking genetically

interacts with the mir-35 and the let-7 miRNA pathways in Caenorhabditis elegans.

Open Biol. *3*, 130151–130151.

de Albuquerque, B.F.M., Placentino, M., and Ketting, R.F. (2015). Maternal piRNAs Are

Essential for Germline Development following De Novo Establishment of Endo-siRNAs

in Caenorhabditis elegans. Dev. Cell *34*, 448–456.

Alvares, S.M., Mayberry, G.A., Joyner, E.Y., Lakowski, B., and Ahmed, S. (2014). H3K4

demethylase activities repress proliferative and postmitotic aging. Aging Cell *13*, 245–

253.

Angelo, G., and Van Gilst, M.R. (2009). Starvation Protects Germline Stem Cells and

Extends Reproductive Longevity in C. elegans. Science (80-.). *326*, 954–958.

Aravin, A.A., Hannon, G.J., and Brennecke, J. (2007). The Piwi-piRNA pathway

provides an adaptive defense in the transposon arms race. Science *318*, 761–764.

Ashe, A., Sapetschnig, A., Weick, E.M., Mitchell, J., Bagijn, M.P., Cording, A.C.,

Doebley, A.L., Goldstein, L.D., Lehrbach, N.J., Le Pen, J., et al. (2012). PiRNAs can

trigger a multigenerational epigenetic memory in the germline of C. elegans. Cell *150*,

88–99.

Batista, P.J., Ruby, J.G., Claycomb, J.M., Chiang, R., Fahlgren, N., Kasschau, K.D.,

Chaves, D.A., Gu, W., Vasale, J.J., Duan, S., et al. (2008). PRG-1 and 21U-RNAs

Interact to Form the piRNA Complex Required for Fertility in C. elegans. Mol. Cell *31*,

67–78.

Billi, A.C., Freeberg, M.A., and Kim, J.K. (2012). piRNAs and siRNAs collaborate in

Caenorhabditis elegans genome defense. Genome Biol. *13*, 164.

Billmyre, K.K., Doebley, A.-L., Spichal, M., Heestand, B., Belicard, T., Sato-Carlton, A.,

Flibotte, S., Simon, M., Gnazzo, M., Skop, A., et al. (2019). The meiotic phosphatase

GSP-2/PP1 promotes germline immortality and small RNA-mediated genome silencing.

PLOS Genet. *15*, e1008004.

Bodnar, A.G., Ouellette, M., Frolkis, M., Holt, S.E., Chiu, C.P., Morin, G.B., Harley, C.B.,

Shay, J.W., Lichtsteiner, S., and Wright, W.E. (1998). Extension of life-span by

introduction of telomerase into normal human cells. Science *279*, 349–352.

Brennecke, J., Malone, C.D., Aravin, A.A., Sachidanandam, R., Stark, A., and Hannon,

G.J. (2008a). An epigenetic role for maternally inherited piRNAs in transposon silencing.

Science *322*, 1387–1392.

Brennecke, J., Malone, C.D., Aravin, A.A., Sachidanandam, R., Stark, A., and Hannon,

G.J. (2008b). An Epigenetic Role for Maternally Inherited piRNAs in Transposon

Silencing. Science (80-.). *322*, 1387–1392.

Buckley, B.A., Burkhart, K.B., Gu, S.G., Spracklin, G., Kershner, A., Fritz, H., Kimble, J.,

Fire, A., and Kennedy, S. (2012). A nuclear Argonaute promotes multigenerational

epigenetic inheritance and germline immortality. Nature *489*, 447–451.

Burkhart, K.B., Guang, S., Buckley, B.A., Wong, L., Bochner, A.F., and Kennedy, S.

(2011). A pre-mRNA-associating factor links endogenous siRNAs to chromatin

regulation. PLoS Genet. *7*, e1002249.

Bushnell, B. BBMap short-read aligner.

Chen, J.-X., Cipriani, P.G., Mecenas, D., Polanowska, J., Piano, F., Gunsalus, K.C., and Selbach, M. (2016). *In Vivo* Interaction Proteomics in *Caenorhabditis elegans* Embryos Provides New Insights into P Granule Dynamics. Mol. Cell. Proteomics *15*, 1642–1657.

Chuma, S., Hosokawa, M., Kitamura, K., Kasai, S., Fujioka, M., Hiyoshi, M., Takamune, K., Noce, T., and Nakatsuji, N. (2006). Tdrd1/Mtr-1, a tudor-related gene, is essential for male germ-cell differentiation and nuage/germinal granule formation in mice. Proc. Natl. Acad. Sci. *103*, 15894–15899.

Counter, C.M., Avilion, A.A., LeFeuvre, C.E., Stewart, N.G., Greider, C.W., Harley, C.B., and Bacchetti, S. (1992). Telomere shortening associated with chromosome instability is arrested in immortal cells which express telomerase activity. EMBO J. *11*, 1921–1929.

Das, P.P., Bagijn, M.P., Goldstein, L.D., Woolford, J.R., Lehrbach, N.J., Sapetschnig, A., Buhecha, H.R., Gilchrist, M.J., Howe, K.L., Stark, R., et al. (2008). Piwi and piRNAs act upstream of an endogenous siRNA pathway to suppress Tc3 transposon mobility in the Caenorhabditis elegans germline. Mol. Cell *31*, 79–90.

Fagagna, F. d'Adda di, Reaper, P.M., Clay-Farrace, L., Fiegler, H., Carr, P., von Zglinicki, T., Saretzki, G., Carter, N.P., and Jackson, S.P. (2003). A DNA damage checkpoint response in telomere-initiated senescence. Nature *426*, 194–198.

Guang, S., Bochner, A.F., Pavelec, D.M., Burkhart, K.B., Harding, S., Lachowiec, J., and Kennedy, S. (2008). An Argonaute Transports siRNAs from the Cytoplasm to the

Nucleus. Science (80-.). *321*, 537–541.

Guang, S., Bochner, A.F., Burkhart, K.B., Burton, N., Pavelec, D.M., and Kennedy, S.

(2010). Small regulatory RNAs inhibit RNA polymerase II during the elongation phase of

transcription. Nature *465*, 1097–1101.

Harley, C.B., Futcher, A.B., and Greider, C.W. (1990). Telomeres shorten during ageing

of human fibroblasts. Nature *345*, 458–460.

Heestand, B., Simon, M., Frenk, S., Titov, D., and Ahmed, S. (2018). Transgenerational

Sterility of Piwi Mutants Represents a Dynamic Form of Adult Reproductive Diapause.

Cell Rep. *23*, 156–171.

Herbig, U., Jobling, W.A., Chen, B.P.C., Chen, D.J., and Sedivy, J.M. (2004). Telomere

shortening triggers senescence of human cells through a pathway involving ATM, p53,

and p21(CIP1), but not p16(INK4a). Mol. Cell *14*, 501–513.

Juliano, C., Wang, J., and Lin, H. (2011). Uniting Germline and Stem Cells: The

Function of Piwi Proteins and the piRNA Pathway in Diverse Organisms. Annu. Rev.

Genet. *45*, 447–469.

Karlseder, J., Broccoli, D., Dai, Y., Hardy, S., and de Lange, T. (1999). p53- and ATM-

dependent apoptosis induced by telomeres lacking TRF2. Science *283*, 1321–1325.

Käser-Pébernard, S., Müller, F., and Wicky, C. (2014). LET-418/Mi2 and SPR-5/LSD1

cooperatively prevent somatic reprogramming of C. elegans germline stem cells. Stem

Cell Reports *2*, 547–559.

Katz, D.J., Edwards, T.M., Reinke, V., and Kelly, W.G. (2009). A C. elegans LSD1

demethylase contributes to germline immortality by reprogramming epigenetic memory.

Cell *137*, 308–320.

Kawasaki, I., Shim, Y.-H., Kirchner, J., Kaminker, J., Wood, W.B., and Strome, S.

(1998). PGL-1, a Predicted RNA-Binding Component of Germ Granules, Is Essential for

Fertility in C. elegans. Cell *94*, 635–645.

Kelleher, E.S. (2016). Reexamining the P-Element Invasion of Drosophila melanogaster

Through the Lens of piRNA Silencing. Genetics *203*, 1513–1531.

Ketting, R.F., Haverkamp, T.H., van Luenen, H.G., and Plasterk, R.H. (1999). Mut-7 of

C. elegans, required for transposon silencing and RNA interference, is a homolog of

Werner syndrome helicase and RNaseD. Cell *99*, 133–141.

Kidwell, M.G., Kidwell, J.F., and Sved, J.A. (1977). Hybrid Dysgenesis in DROSOPHILA

MELANOGASTER: A Syndrome of Aberrant Traits Including Mutation, Sterility and

Male Recombination. Genetics *86*, 813–833.

Kim, D., Pertea, G., Trapnell, C., Pimentel, H., Kelley, R., and Salzberg, S.L. (2013).

TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions

and gene fusions. Genome Biol. *14*, R36.

Kim, S.T., Lim, D.S., Canman, C.E., and Kastan, M.B. (1999). Substrate specificities

and identification of putative substrates of ATM kinase family members. J. Biol. Chem.

274, 37538–37543.

Knutson, A.K., Egelhofer, T., Rechtsteiner, A., and Strome, S. (2017). Germ Granules

Prevent Accumulation of Somatic Transcripts in the Adult Caenorhabditis elegans

Germline. Genetics.

Krishnamurthy, J., Torrice, C., Ramsey, M.R., Kovalev, G.I., Al-Regaiey, K., Su, L., and

Sharpless, N.E. (2004). Ink4a/Arf expression is a biomarker of aging. J. Clin. Invest.

114, 1299–1307.

Krishnamurthy, J., Ramsey, M.R., Ligon, K.L., Torrice, C., Koh, A., Bonner-Weir, S., and

Sharpless, N.E. (2006). p16INK4a induces an age-dependent decline in islet

regenerative potential. Nature *443*, 453–457.

Langmead, B., Trapnell, C., Pop, M., and Salzberg, S.L. (2009). Ultrafast and memory-

efficient alignment of short DNA sequences to the human genome. Genome Biol. *10*,

R25.

Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose

program for assigning sequence reads to genomic features. Bioinformatics *30*, 923–

930.

Lowden, M.R., Meier, B., Lee, T.W.S., Hall, J., and Ahmed, S. (2008). End joining at

Caenorhabditis elegans telomeres. Genetics *180*, 741–754.

Luteijn, M.J., and Ketting, R.F. (2013). PIWI-interacting RNAs: from generation to

transgenerational epigenetics. Nat. Rev. Genet. *14*, 523–534.

McMurchy, A.N., Stempor, P., Gaarenstroom, T., Wysolmerski, B., Dong, Y.,

Aussianikava, D., Appert, A., Huang, N., Kolasinska-Zwierz, P., Sapetschnig, A., et al.

(2017). A team of heterochromatin factors collaborates with small RNA pathways to

combat repetitive elements and germline stress. Elife *6*.

Meier, B., Clejan, I., Liu, Y., Lowden, M., Gartner, A., Hodgkin, J., and Ahmed, S.

(2006). trt-1 is the Caenorhabditis elegans catalytic subunit of telomerase. PLoS Genet.

2, 187–197.

Min, H., Shim, Y.-H., and Kawasaki, I. (2016). Loss of PGL-1 and PGL-3, members of a

family of constitutive germ-granule components, promotes germline apoptosis in C.

elegans. J. Cell Sci. *129*, 341–353.

Nakamura, T.M., Cooper, J.P., and Cech, T.R. (1998). Two modes of survival of fission

yeast without telomerase. Science *282*, 493–496.

Ni, J.Z., Kalinava, N., Chen, E., Huang, A., Trinh, T., and Gu, S.G. (2016). A

transgenerational role of the germline nuclear RNAi pathway in repressing heat stress-

induced transcriptional activation in C. elegans. Epigenetics and Chromatin *9*, 1–15.

Padeken, J., Zeller, P., Towbin, B., Katic, I., Kalck, V., Methot, S.P., and Gasser, S.M.

(2019). Synergistic lethality between BRCA1 and H3K9me2 loss reflects satellite

derepression. Genes Dev. 1–16.

Padilla, P.A., and Ladage, M.L. (2012). Suspended animation, diapause and

quiescence. Cell Cycle *11*, 1672–1679.

Phillips, C.M., Montgomery, T.A., Breen, P.C., and Ruvkun, G. (2012). MUT-16

promotes formation of perinuclear mutator foci required for RNA silencing in the C.

elegans germline. Genes Dev. *26*, 1433–1444.

Phillips, C.M., Brown, K.C., Montgomery, B.E., Ruvkun, G., and Montgomery, T.A.

(2015). piRNAs and piRNA-Dependent siRNAs Protect Conserved and Essential

C. elegans Genes from Misrouting into the RNAi Pathway. Dev. Cell *34*, 457–465.

R Core Team (2013). R: A language and environment for statistical   computing. R

Found. Stat. Comput.

Robert, V.J., Mercier, M.G., Bedet, C., Janczarski, S., Merlet, J., Garvis, S., Ciosk, R.,

and Palladino, F. (2014). The SET-2/SET1 Histone H3K4 Methyltransferase Maintains

Pluripotency in the Caenorhabditis elegans Germline.

Sakaguchi, A., Sarkies, P., Simon, M., Doebley, A.-L., Goldstein, L.D., Hedges, A.,

Ikegami, K., Alvares, S.M., Yang, L., LaRocque, J.R., et al. (2014). Caenorhabditis

elegans RSD-2 and RSD-6 promote germ cell immortality by maintaining small

interfering RNA populations. Proc. Natl. Acad. Sci. U. S. A. *111*, E4323-31.

Sharpless, N.E., and DePinho, R.A. (2007). How stem cells age and why this makes us

grow old. Nat. Rev. Mol. Cell Biol. *8*, 703–713.

Shay, J.W., and Wright, W.E. (2000). Hayflick, his limit, and cellular ageing. Nat. Rev.

Mol. Cell Biol. *1*, 72–76.

Shay, J.W., and Wright, W.E. (2005). Senescence and immortalization: role of

telomeres and telomerase. Carcinogenesis *26*, 867–874.

Simon, M., Sarkies, P., Ikegami, K., Doebley, A.L., Goldstein, L.D., Mitchell, J.,

Sakaguchi, A., Miska, E.A., and Ahmed, S. (2014a). Reduced Insulin/IGF-1 Signaling

Restores Germ Cell Immortality to Caenorhabditis elegans Piwi Mutants. Cell Rep. *7*,

762–773.

Simon, M., Sarkies, P., Ikegami, K., Doebley, A.-L., Goldstein, L.D., Mitchell, J.,

Sakaguchi, A., Miska, E.A., and Ahmed, S. (2014b). Reduced Insulin/IGF-1 Signaling

Restores Germ Cell Immortality to Caenorhabditis elegans Piwi Mutants. Cell Rep. *7*,

762–773.

Siomi, M.C., Sato, K., Pezic, D., and Aravin, A.A. (2011). PIWI-interacting small RNAs:

the vanguard of genome defence. Nat. Rev. Mol. Cell Biol. *12*, 246–258.

Smelick, C., and Ahmed, S. (2005a). Achieving immortality in the C. elegans germline.

Ageing Res. Rev. *4*, 67–82.

Smelick, C., and Ahmed, S. (2005b). Achieving immortality in the C. elegans germline.

Ageing Res. Rev. *4*, 67–82.

Spracklin, G., Fields, B., Wan, G., Vijayendran, D., Wallig, A., Shukla, A., and Kennedy,

S. (2017). Identification and Characterization of Caenorhabditis elegans RNAi

Inheritance Machinery. Genetics.

Srivastav, S.P., and Kelleher, E.S. (2017). Paternal Induction of Hybrid Dysgenesis in

Drosophila melanogaster Is Weakly Correlated with Both *P* -Element and *hobo* Element

Dosage. G3: Genes|Genomes|Genetics *7*, 1487–1497.

Strome, S., and Wood, W.B. (1982). Immunofluorescence visualization of germ-line-

specific cytoplasmic granules in embryos, larvae, and adults of Caenorhabditis elegans.

Proc. Natl. Acad. Sci. U. S. A. *79*, 1558–1562.

Sung, M., Kawasaki, I., and Shim, Y.-H. (2017). Depletion of *cdc-25.3* , a

Caenorhabditis elegans orthologue of *cdc25* , increases physiological germline

apoptosis. FEBS Lett. *591*, 2131–2146.

Tatar, M., and Yin, C. (2001). Slow aging during insect reproductive diapause: why

butterflies, grasshoppers and flies are like worms. Exp. Gerontol. *36*, 723–738.

Tchkonia, T., Zhu, Y., van Deursen, J., Campisi, J., and Kirkland, J.L. (2013). Cellular

senescence and the senescent secretory phenotype: therapeutic opportunities. J. Clin.

Invest. *123*, 966–972.

Updike, D., and Strome, S. (2010). P Granule Assembly and Function in Caenorhabditis

elegans Germ Cells. J. Androl. *31*, 53–60.

Updike, D.L., Hachey, S.J., Kreher, J., and Strome, S. (2011). P granules extend the

nuclear pore complex environment in the *C. elegans* germ line. J. Cell Biol. *192*, 939–

948.

Updike, D.L., Knutson, A.K., Egelhofer, T.A., Campbell, A.C., and Strome, S. (2014).

Germ-Granule Components Prevent Somatic Development in the C. elegans Germline.

Curr. Biol. *24*, 970–975.

Vermezovic, J., Stergiou, L., Hengartner, M.O., and d'Adda di Fagagna, F. (2012).

Differential regulation of DNA damage response activation between somatic and

germline cells in Caenorhabditis elegans. Cell Death Differ. *19*, 1847–1855.

Zeller, P., Padeken, J., van Schendel, R., Kalck, V., Tijsterman, M., and Gasser, S.M.

(2016). Histone H3K9 methylation is dispensable for Caenorhabditis elegans

development but suppresses RNA:DNA hybrid-associated repeat instability. Nat. Genet.

48, 1385–1395.

Zheng, K., and Wang, P.J. (2012). Blockade of Pachytene piRNA Biogenesis Reveals a

Novel Requirement for Maintaining Post-Meiotic Germline Genome Integrity. PLoS

Genet. *8*, e1003038.

Table 1

Percent of germlines showing P granule abnormalities

	fertile	sterile
20˚C		
wt (n=12)	0.00%	NA
prg-1(n4357) (n=16)	36.36%	100.00%
prg-1(tm872) (n=21)	87.50%	92.31%
nrde-4(gg131) (n=13)	50.00%	90.91%
nrde-1(yp4) (n=13)	20.00%	87.50%
25˚C		
wt (n=35)	0.00%	NA
mut-14(pk738) (n=15)	NA	85%
rsd-6(yp11) (n=32)	0.00%	40.00%
nrde-2(gg95) (n=25)	0.00%	100.00%
rbr-2(tm1231) (n=32)	0.00%	100.00%
hrde-1(tm1200) (n=31)	0.00%	100.00%

Table 1: Quantification of P granule defects in the germline.

Sterile Piwi pathway genome silencing mutants were stained with P granule antibody.
Germlines arms were semi-qualitatively scored for P granules. If 20% or more of the the
germline arm exhibited P granules abnormalities (complete or partial loss of P granules)
germlines were scored as abnormal.

Supplemental Information

OICD/DAPI

Late generation fertile 20˚C   Sterile 20˚C

Late generation fertile 25˚C   Sterile 25˚C

Fig. S1. Related to Figure 2. P granule defects in *prg-1, nrde- 4, hrde-1 and nrde-1*.
Germlines of sterile Day 2-3 adult animals were stained using the OIC1D4 antibody
against P granules (green) and DAPI (blue). (A) Control animals contain uniform puncta
of P granule staining surrounding each nuclei. (B-F) Late generation fertile *prg-
1(tm876), nrde-4(gg131), nrde-1(yp4)*, *hrde-1(tm1200)* and *rsd-6(yp11)* and animals
displayed P granule staining similar to wild type except for *prg-1(tm876).* (B'-F') Sterile
prg-1(tm876), nrde-4(gg131), nrde-1(yp4), *hrde-1(tm1200)* and *rsd-6(yp11)* with P
granule abnormalities.

	Total Worms Scored	% Fertile	% Sterile
pgl-1 G1 at 25°C	91	22%	78%
pgl-1 G2 at 25°C	71	0%	100%

Table S1: Related to Figure 3. Fertility of *pgl-1* mutants at 25°C. 20°C P0 *pgl-1*
mutants were shifted to 25°C and fertility was measured for the G1 generation. Fertile
G1 worms were cloned and the fertility of the G2 generation was measured.

gene	*prg-1*	*nrde-1*	*nrde-4*
C06B3.7	4.331128036	1.732445354	2.669978253
dhs-26	3.88587979	3.752818032	1.793937273
bath-45	3.604851731	5.5002102	2.319444032
fbxb-97	3.555301139	3.002992933	3.278539846
irg-2	3.519633623	2.605415087	2.189754742
pud-4	3.226292885	2.956152303	4.407697185
pud-3	3.097392415	3.143142026	4.328678969
R09E10.2	2.745900789	5.100997358	4.251177196
Y53F4B.5	2.636725232	5.821227047	4.162462556
K08D12.6	2.631068263	2.955437046	0.277803871
R03H10.6	2.469258519	4.82098578	3.295830487
Y58A7A.5	2.397260966	1.795895	2.143861408
pud-1.1	2.299685196	2.556376909	3.17781014
T22B7.7	2.276777478	3.01143186	2.207919501
fbxa-224	2.248998131	4.226992478	1.979961678
swt-6	2.144598564	3.153313277	0.277029378
pud-2.1	2.119890425	2.781615959	2.880747229
pud-2.2	2.094123631	2.779337879	2.796566876
pud-1.2	2.080250623	2.559790088	3.000944771
catp-3	2.014970724	2.436698711	2.674464594

**Table S2: Related to Figure 4. Genes upregulated in late-generation *prg-1* mutants
and in sterile *nrde-1* and *nrde-4* mutant L4 larvae.**