1

6 # Transgenerational sterility of small RNA genome silencing mutants in

7 # response to germ granule dysfunction

10
11 Katherine Kretovich Billmyre[1,2]*, Bree Heestand[1,2,3]*, Maya Spichal[1,2]* and Shawn
12 Ahmed[1,2,3]
13
14

15 [1]Department of Genetics, University of North Carolina, Chapel Hill, NC, USA
16 [2]Department of Biology, University of North Carolina, Chapel Hill, NC, USA
17 [3]Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill, NC, USA
18 To whom correspondence should be addressed. E-mail: shawn@med.unc.edu
19
20 * These authors contributed equally to this manuscript

Abstract

Deficiency for small RNA nuclear silencing proteins that promote heterochromatin maintenance

leads to defects in germ cell immortality where fertile mutant strains become sterile over

successive generations. The transgenerational sterility of small RNA nuclear silencing is

associated with mis-wiring of small RNA pathways and with progressive deterioration of

heterochromatin, which leads to increased levels of transposon expression, transposition and

RNA-DNA hybrids. However, the cause of transgenerational sterility in small RNA nuclear

silencing mutants is not understood. We found that sterile small RNA genome silencing mutants

display altered perinuclear germ granules. Moreover, germ granule dysfunction was sufficient

to induce phenotypes associated with sterile small RNA genome silencing mutants, including

germline atrophy, reproductive diapause and univalents in oocytes. We conclude that germ

granules, which are intimately associated with small RNA nuclear silencing proteins and RNAs,

orchestrate the transgenerational sterility of small RNA genome silencing mutants.

Introduction

Germ cells give rise to the mortal somatic tissue while maintaining themselves in a pristine

condition that allows them to be propagated for indefinite number of generations. This capacity

for self-renewal is termed "germ cell immortality", and functions to protect the germline from

stressors which can result in sterility[1]. *C. elegans* mutations in genes that promote small RNA-

mediated genome silencing (Fig. 4a) can cause a Mortal Germline *(mrt)* phenotype, where

fertility occurs for several generations but leads to a deterioration of fertility in late-generations

and finally to complete sterility. Many *mrt* mutants exhibit a conditional transgenerational

sterility phenotype, such that they only become sterile when grown at the non-permissive

temperature 25˚C[1]. Examples of conditional *mrt* mutants include a nuclear RNAi defective

mutant, *nrde-2,* a number of heritable RNAi mutants, including *hrde-1*, and two RNAi defective

mutants, *rsd-2* and *rsd-6*[2–7]. Other *mrt* mutants display progressive sterility at any temperature,

including *prg-1*/piRNA mutants and nuclear RNAi mutants *nrde-1* and *nrde-4*[4–6,8–10].

Piwi is a conserved germ line Argonaute protein that interacts with thousands of small RNAs

termed piRNAs to repress transposons and foreign nucleic acids that represent parasitic threats

to the integrity of the genome. Deficiency for the *C. elegans* Piwi ortholog *prg-1* was recently

shown to elicit a striking germ cell atrophy phenotype at the generation of sterility, as L4 larvae

mature into 1 day old adults[11]. Moreover, atrophied germ lines of sterile *prg-1* mutants can

regrow on day 2 of adulthood, and a small fraction of sterile adults can become fertile again if

animals are shifted to a distinct food source[11]. This indicates that the late-generation sterility of

prg-1/Piwi mutants is a form of reproductive arrest termed Adult Reproductive Diapause[12].

Temperature-sensitive small RNA genome silencing mutants *rsd-2* and *rsd-6* also exhibit small

and empty germlines at the generation of sterility, indicating that the transgenerational sterility

of small RNA genome silencing mutants elicits consistent developmental consequences[2].

Although these *mrt* mutants likely suffer from transgenerational deterioration of

heterochromatin, the underlying cause of the late-generation sterility of these mutants is not

understood.

The fertility defects in response to heterochromatin dysfunction have been associated with

increased levels DNA damage[4,13]. Moreover, the fertility of late-generation *prg-1* mutants could

be improved by RNAi knockdown of the Mirage transposon, suggesting a link between

transposition and fertility defects of *prg-1* mutants[4]. Furthermore, RNA-DNA hybrids that result

from collisions between replication forks and RNA polymerase were increased in response to

heterochromatin defects[4,13]. *prg-1* mutants display progressively increased levels of germline

apoptosis when grown for successive generations, and at the generation of sterility apoptosis

promotes germ cell atrophy for *rsd-2*, *rsd-6* and *prg-1* mutants[2,10,11]. At the same time, the

frequency of spontaneous mutation in either *prg-1* or *rsd-6 mrt* mutants is minimal[2,7,10].

Moreover, temperature shift experiments have indicated that the trigger of the

transgenerational fertility defects in such mutants is reversible and therefore not the

consequence of (transposon-induced) changes to the genome[7]. Finally, high levels of

transposition in *C. elegans mutator* mutants that experience significant levels of transposition

do not display progressive sterility phenotype if propagated at low temperatures[10].

Another possible cause of small RNA mutant germline atrophy is disruption in proper function

of the pro- and anti-silencing of the *C. elegans* genome [14,15]. Major proteins involved in these

pathways such as PRG-1 and CSR-1, respectively, are located in perinuclear germ granules,

which are conserved germline structures that consist of liquid droplets that interact with

nuclear pore complexes (NPCs) and are necessary for fertility [16]. In *C. elegans*, germ granules

are termed P-granules based on antibodies that label the P blastomere that generates the germ

cell lineage[17].

Here we study the late-generation sterility phenotype of small RNA-mediated genome silencing

mutants and link this to dysfunction of P granules, which play a central role in modulating the

process of small RNA genome silencing itself.

Results

Genome instability in small RNA mutants

Transgenerational sterility in *C. elegans* telomerase mutants that lack the ability to maintain

sequences that cap chromosome ends is caused by high levels of telomere fusions and

associated damage to chromosome termini[18]. We examined *prg-1* mutants grown at 20°C at

progressive generations for the presence of the DNA damage response and observed that late-

generation *prg-1* mutants and sterile *prg-1* mutants showed a significant 3 to 6 fold

upregulation of DNA damage response when compared to wild-type controls or to early

generation *prg-1* mutants (Fig. 1a). However, the number of cells with DNA damage was not

consistently increased in sterile *prg-1* mutant samples, when compared to fertile, late-

generation controls (Fig. 1a). We also tested levels of DNA damage response in the

temperature-sensitive small RNA mutants *rsd-6, nrde-2* and *hrde-1*. We found that the wild-

type control DNA damage response was upregulated in many cells at 25°C but that this level did

not change in sterile *rsd-6, nrde-2* or *hrde-1* mutant germlines sterile *prg-1* (Fig. 1b). Therefore,

although the DNA damage response is transgenerationally upregulated in *prg-1* mutants, it is

unlikely to be the cause of sterility in small RNA genome silencing mutants.

Small RNA mutants exhibit P-granule defects at sterility

Whilst we were characterizing the germ line defects present sterile *rsd-6* small RNA genomic

silencing mutants in immunofluorescence assays, we happened to use an antibody against P-

granules as a positive control. Surprisingly, we found there was an altered P-granule staining

pattern when compared to fertile r*sd-6* mutant siblings. This finding was interesting as PRG-1

localizes to P-granules and deficiency for PRG-1 causes delocalization of the structural P-granule

protein PGL-1[19]. Additionally, combined loss of the epigenomic regulators SPR-5 and LET-418

results in immediate sterility that is associated with germ cell to soma transformation that

disrupts P granule structures, and the H3K9 methylation modifier SET-2 can cause a loss of

perinuclear P-granules in a context where the nuclear RNAi pathway acts in parallel to promote

germ cell identity[20,21]. We therefore asked if sterility in small RNA genomic mutants was

consistently linked with P-granule dysfunction by testing sterile small RNA mutants *hrde-1*, *rsd-*

6, and *nrde-2* and found that sterile animals display P-granule dysfunction, but that P-granule

defects were absent or comparatively minor in fertile late-generation animals that were close

to the generation of sterility (Fig. 2c-d, S1, Table 1). When compared to wild-type controls, P-

granule dysfunction in the sterile mutants ranged from a loss of P-granules in a subset of cells

to varying amounts of P-granule disorganization (Fig. 2c-d)*.* Sterile *hrde-1* mutant animals

displayed the most severe P-granule phenotype, as the majority of sterile *hrde-1* mutant germ

cells exhibited little to no staining (Fig. S1).

Large-scale apoptosis occurs in some but not all *prg-1* mutants as L4 larvae to adult

transition at generation of sterility[11], and P-granules have been shown to vanish in cells that

undergo apoptosis[22,23]. However, we saw widespread P granule dysfunction in sterile germlines

of *rsd-6* mutants that were wildtype in size and had not undergone significant levels of

apoptosis (Fig. 2d). Moreover, sterile *nrde-4* mutant germlines did not show obvious defects to

the structure of P-granules, indicating that the sterility of small RNA genome silencing mutants

is not always associated with overt effects on P granule structure (Fig. S1).

In contrast to many small RNA genome silencing mutants where fertility defects are not

severe until the generation of sterility, late-generation *prg-1* mutants display very small brood

sizes prior to sterility[4,11]. By immunofluorescence, late-generation *prg-1* mutants had a wide

range of P-granule defects that could be identified in fertile and sterile animals (Fig. 2b, S1).

Many germlines exhibited strong P-granule dysfunction that either resulted in a total absence

of P-granules or irregular distribution (Fig. 2b, S1). Often these strong P-granule defects

coincided with sterile animals, though some sterile *prg-1* germlines displayed mild defects.

We next examined a temperature sensitive mutant, *rbr-2(tm1231)*, which encodes a histone 3

lysine 4 demethylase known to promote genomic silencing and is thought to function

downstream of small RNAs to promote genomic silencing[24]. Sterile *rbr-2* animals displayed loss

of P-granules in many germ cells, but late-generation fertile *rbr-2* animals did not exhibit

obvious P-granule defects (Fig. 2e).

C. elegans P-granules are adjacent to Mutator bodies[28], where small RNA-mediated silencing

pathway proteins function to amplify primary siRNA populations into effector secondary

siRNAs. *mutator* mutants are dysfunctional for RDRP-mediated biogenesis of effector small RNA

and although they remain fertile indefinitely at low temperatures, they display an immediate

highly penetrant sterility phenotype at 25°C that could be caused by dysfunction of small RNA-

mediated heterochromatin.

P-granule dysfunction is sufficient to elicit phenotypes of sterile small RNA silencing mutants

Given that P-granule loss has been shown to cause a germline to soma transformation in

conjunction with short germlines in adults [25,26], we assessed the consequences of an RNA

interference clone, that simultaneously targets four P-granule subunits *pgl-1*, *pgl-3*, *glh-1*, *glh-4*

[25,26]. Knockdown of P-granules by RNAi resulted in F2 sterile animals at 25°C (80%) (Fig. 3a).

When *prg-1* mutants became sterile their germline profile shifts specifically at adulthood from

large L4 larval stage germlines to adult only atrophy[11]. Interestingly, sterile P-granule RNAi F2

adults at 25°C displayed a range of germline phenotypes similar to those observed in our small

RNA silencing sterile adults[11] (Fig. 3b).

172   To confirm that the apparent germline degeneration phenotypes of P-granule depleted

adults were not due to failure of germ cells to proliferate in early development, we collected L4

larval stage F2 animals and assessed germline development. The germlines of F2 P-granule RNAi

L4 larvae were normal in size in comparison to control worms, but a significant change in their

germline profile occurred as these animals matured into adults with pronounced germ cell

atrophy occurring at adulthood (Fig. 3b, p=3.93e-24). Furthermore, for sterile P-granule

depleted worms that contained oocytes, 45.5% of oocytes contained 7-12 univalents (Fig. 3c, e-

h), revealing a phenocopy of the oocyte univalents that was previously observed for oocytes of

late-generation sterile *rsd-6, hrde-1* and *nrde-2* mutants[2].

prg-1 mutants phenocopy and _pgl-1_ mutants

If P-granule defects cause the late-generation sterility of small RNA genome silencing mutants,

we hypothesized that the germline atrophy phenotypes observed in P-granule knockdown

worms might have the potential to mimic the delayed fertility phenomenon for sterile _prg-1_

mutant adults, which can regrow atrophied germlines to become fertile[11]. We therefore used

the temperature-sensitivity sterility of _pgl-1_ mutants that become sterile in the F1 generation

at 25˚C but can be fertile at low temperatures. While most F1 _pgl-1_ animals at 25˚C displayed

normal germlines at the L4 larval stage, there was a significant shift in the germline profiles

with 52% of day 1 adults showing germline atrophy (Fig. 3b, p=1.04E-35), similar to P-granule

RNAi knockdown and sterile _prg-1_ mutants[11]. Furthermore, when these sterile _pgl-1_ worms

were scored for univalents in worms that contained oocytes, we found that 56.7% of oocytes

had univalents, in agreement with the oocyte univalent observed for P-granule RNAi oocytes

(Fig. 3c). 20% of F1 _pgl-1_ mutants gave rise to maternally depleted P-granule F2 animals and

these were uniformly sterile. F2 L4 larval stage germline profiles displayed more severe germ

cell proliferation defects than the L4 F1 generation, and day 1 adults showed a striking 54%

increase in empty germlines (Fig. 3d). This suggest that F1 generation _pgl-1_ mutants mimic the

germ cell degeneration phenotypes of sterile small RNA genome silencing mutants at the L4 to

adult transition, whereas maternal depletion of _pgl-1_ elicits more pronounced effects on germ

cell proliferation at earlier stages of development and at the L4/adult transition.

When sterile F1 _pgl-1_ day 1 adults were shifted to 20˚C, some animals were capable of

regrowing their germlines. We scored the germline profiles of sterile F1 _pgl-1_ adults shifted to

20°C after 48 hours and saw a significant change compared to controls (Fig. 3i, p=0.007). The

germline profile shifted from a high incidence of atrophy germlines towards an increase in large

germlines when compared to sterile F1 *pgl-1* sibling controls that remained at 25°C (Fig. 3i). In

agreement with this data, several days after 1 day old sterile F1 *pgl-1* mutant adults at 25°C

were shifted to 20°C, we observed that some laid oocytes and dead embryos and rarely

progeny, whereas sterile day 1 *pgl-1* at 25°C never gave rise to oocytes, dead embryos or

progeny (Fig. 3j). The frequency of fertility in this circumstance is very similar to that of 5 day

old sterile *prg-1* mutant adults that exit Adult Reproductive Diapause in response to an altered

food source[11]. These results indicate that severe P-granule dysfunction and associated sterility

can mimic a state of reproductive quiescence that does not abolish fertility.

Discussion

Previous work has established that a subset of small RNA genome silencing mutants display a

transgenerational sterility phenotype in conjunction with desilencing of various repetitive

segments of the genome[2,6,10]. However, the molecular mechanism that evokes

transgenerational sterility has remained an open question. We show that *prg-1* and several

other small RNA mutants show P-granule defects at sterility[27]. Given that these mutants

experience desilencing of heterochromatic transposons, their sterility might occur in response

to an increased DNA damage response induced by transposition or by R-loop formation.

However, we observed that sterile late-generation small RNA mutants at 25°C displayed similar

levels of DNA damage response as wild-type controls. Consistently, while levels of DNA damage

increased in late-generation *prg-1* mutants, these levels did not consistently increase at

sterility.

We examined the integrity of P-granules in sterile small RNA genome silencing mutants and

found that P-granule defects occur in sterile animals of most late-generation *mortal germline*

mutants examined. Our data further show that a direct knockdown of P-granules by RNAi was

sufficient to mimic the germline atrophy and meiotic recombination defects that are observed

for small RNA mutants. The similarities between the sterility phenotypes of *pgl-1* mutants, P-

granule RNAi treated animals and small RNA silencing mutants imply that P-granule defects are

likely to cause sterility in small RNA mutants. Furthermore, our data indicate that univalent

chromosomes that are observed in oocytes of some sterile small RNA genome silencing animals

are a likely secondary consequence of P-granule dysfunction that indirectly disrupts a process

that promotes meiotic recombination[29,30]. Interestingly, when *pgl-1* mutants were shifted to

the permissive temperature of 20°C, there is a decrease in empty and atrophied germlines, a

corresponding increase in the number of large germlines and a few give rise to progeny. These

results closely resemble the regrowth of the germline and the fertility observed for sterile late-

generations *prg-1* mutants that are in Adult Reproductive Diapause[11]. Taken together our

results suggest that P-granule dysfunction is likely to be the mechanism that promotes the

reproductive arrest phenotype termed Adult Reproductive Diapause.

Small RNA silencing proteins such as CSR-1, PRG-1 or HRDE-1 are localized to P-granules, and it

has been hypothesized that small RNA dysfunction in P-granules could be the cause of sterility

in small RNA genome silencing mutants[14,19]. Our evidence supports a loss of P-granule structure

and function, not DNA damage response, as the trigger of sterility in late-generation *mrt*

mutants (Fig. 4b). Our results establish a novel framework to consider inheritance in the

context of epigenetic stress and how this can evoke transgenerational sterility. Future efforts to

understand how epigenomic defects lead to P-granule mis-regulation may provide insight into

the precise cause of sterility in small RNA silencing mutants. We suggest that the sterility in

response to epigenome dysfunction is consistent with a developmental arrest mechanism to

protect the genome or epigenome in times of stress. We note that transposons and viruses that

are central targets of the small RNA genome silencing system are likely in a continuous arms

race between parasite and host, where components of P-granules are likely to be targeted[27]. It

is therefore reasonable to hypothesize and arrest mechanism in the context of an attack on the

P-granule silencing system, which is predicated on the integrity of the P-granules themselves.

Methods

Strains:

All strains were cultured at 20°C or 25°C on Nematode Growth Medium (NGM) plates seeded

with *E. coli* OP50. Strains used include Bristol N2 wild type, *hrde-1(tm1200) III, rsd-6(yp11) I,*

nrde-2(gg95) II, and *rbr-2(tm1231) IV.*

RNAi Assay:

Feeding RNAi plates harboring host bacteria HT115(DE3) engineered to express "quad" dsRNA

(targeting P-granules) were obtained from Susan Strome [25,26]. L1 larvae were placed onto

freshly prepared feeding RNAi plates with dsRNA induced by 1 mM IPTG (isopropyl-β-D(-)-

thiogalactopyranoside) and were transferred after 1 generation at 25°C and collected at F2

adults as described in Knutson et al., 2017 for DAPI staining, oocyte and germline analysis.

DAPI staining and Scoring:

DAPI staining was performed as previously described[18]. Briefly, L4 larvae were selected from

sibling plates and sterile adults were singled as late L4s, observed 24 hours later for confirmed

sterility, and then stained 48 hours after collection. Univalents were scored by counting DAPI

bodies in the -1 to -4 oocytes. Germline profiles were scored using the method outlined in

Heestand et al 2018.

Statistical Analysis

Statistical analysis was performed as previously described[11]. Briefly, statistical analysis was

performed using the R statistical environment[31]. For germline phenotypes, contingency tables

were constructed and pairwise Chi Square tests with Bonferroni correction was used to

determine significant differences in in germline phenotype distributions.

Immunofluorescence:

Adult hermaphrodites raised at 20°C or 25°C were dissected in M9 buffer and flash frozen on

dry ice before fixation for 1 min in methanol at -20°C. After washing in PBS supplemented with

0.1% Tween-20 (PBST), primary antibody diluted in in PBST was used to immunostain overnight

at 4 °C in a humid chamber. Primaries used were 1:50 OIC1D4 (Developmental Studies

Hybridoma Bank). Secondary antibody staining was performed by using an Cy3 donkey anti-

mouse or Cy-5 donkey anti-rabbit overnight at 4°C. All images were obtained using a LSM 710

laser scanning confocal and were taken using same settings as control samples. Images

processed using ImageJ.

DNA Damage Assay:

Worms that were close to sterility were isolated and defined as sterile if they did not have any

offspring as day 2 adults at 25˚C. The presence of the DNA damage response by using a

phospho-specific antibody targeting the phosphorylated consensus target site of ATM and ATR

kinases (pS/TQ). This antibody has only been shown to stain a DNA damage response in the

germline[32].

Scoring of P-granules:

The number of P-granules was determined in 60 representative DAPI-stained nuclei of the

middle section of a z-stack taken at 40x for 3 different germlines per condition. P-granule

counts were attributed to different number counts per nucleus (0, 1-4, 5-8, >8).

Author Contributions

KB, MS and BH performed experiments. KB, MS, BH and SA wrote manuscript.

References

1. Smelick, C. & Ahmed, S. Achieving immortality in the C. elegans germline. *Ageing Res.*

Rev. **4,** 67–82 (2005).

2. Sakaguchi, A. *et al.* Caenorhabditis elegans RSD-2 and RSD-6 promote germ cell

immortality by maintaining small interfering RNA populations. *Proc. Natl. Acad. Sci. U. S.*

A. **111,** E4323-31 (2014).

3. Ashe, A. *et al.* piRNAs can trigger a multigenerational epigenetic memory in the germline

of C. elegans. *Cell* **150,** 88–99 (2012).

4. McMurchy, A. N. *et al.* A team of heterochromatin factors collaborates with small RNA

pathways to combat repetitive elements and germline stress. *Elife* **6,** (2017).

5. Buckley, B. A. *et al.* A nuclear Argonaute promotes multigenerational epigenetic

inheritance and germline immortality. *Nature* **489,** 447–451 (2012).

6. Burkhart, K. B. *et al.* A pre-mRNA-associating factor links endogenous siRNAs to

chromatin regulation. *PLoS Genet.* **7,** e1002249 (2011).

7. Spracklin, G. *et al.* Identification and Characterization of Caenorhabditis elegans RNAi

Inheritance Machinery. *Genetics* (2017).

8. Batista, P. J. *et al.* PRG-1 and 21U-RNAs interact to form the piRNA complex required for

fertility in C. elegans. *Mol. Cell* **31,** 67–78 (2008).

9. Das, P. P. *et al.* Piwi and piRNAs Act Upstream of an Endogenous siRNA Pathway to

Suppress Tc3 Transposon Mobility in the Caenorhabditis elegans Germline. *Mol. Cell* **31,**

79–90 (2008).

10.   Simon, M. *et al.* Reduced insulin/IGF-1 signaling restores germ cell immortality to

Caenorhabditis elegans Piwi mutants. *Cell Rep.* **7,** 762–73 (2014).

11.   Heestand, B., Simon, M., Frenk, S., Titov, D. & Ahmed, S. Transgenerational sterility of

Piwi mutants represents a dynamic form of Adult Reproductive Diapause. *Cell Rep.* **in**

press, (2018).

12.   Angelo, G. & Van Gilst, M. R. Starvation Protects Germline Stem Cells and Extends

Reproductive Longevity in C. elegans. *Science (80-.).* **326,** 954–958 (2009).

13.   Zeller, P. *et al.* Histone H3K9 methylation is dispensable for Caenorhabditis elegans

development but suppresses RNA:DNA hybrid-associated repeat instability. *Nat. Genet.*

48, 1385–1395 (2016).

14.   de Albuquerque, B. F. M., Placentino, M. & Ketting, R. F. Maternal piRNAs Are Essential

for Germline Development following De Novo Establishment of Endo-siRNAs in

Caenorhabditis elegans. *Dev. Cell* **34,** 448–56 (2015).

15.   Phillips, C. M., Brown, K. C., Montgomery, B. E., Ruvkun, G. & Montgomery, T. A. piRNAs

and piRNA-Dependent siRNAs Protect Conserved and Essential C. elegans Genes from

Misrouting into the RNAi Pathway. *Dev. Cell* **34,** 457–465 (2015).

16.   Updike, D. L., Hachey, S. J., Kreher, J. & Strome, S. P granules extend the nuclear pore

complex environment in the *C. elegans* germ line. *J. Cell Biol.* **192,** 939–948 (2011).

17.   Strome, S. & Wood, W. B. Immunofluorescence visualization of germ-line-specific

cytoplasmic granules in embryos, larvae, and adults of Caenorhabditis elegans. *Proc.*

Natl. Acad. Sci. U. S. A. **79,** 1558–62 (1982).

18.   Ahmed, S. & Hodgkin, J. MRT-2 checkpoint protein is required for germline immortality

and telomere replication in C. elegans. *Nature* **403,** 159–64 (2000).

19. Phillips, C. M., Brown, K. C., Montgomery, B. E., Ruvkun, G. & Montgomery, T. A. piRNAs

1. Phillips, C. M., Brown, K. C., Montgomery, B. E., Ruvkun, G. & Montgomery, T. A.

piRNAs and piRNA-Dependent siRNAs Protect Conserved and Essential C. elegans Genes

from Misrouting into the RNAi Pathway. Dev. Cell 34, 457–465 (2015).and piRNA-Dep.

Dev. Cell **34,** 457–465 (2015).

20. Käser-Pébernard, S., Müller, F. & Wicky, C. LET-418/Mi2 and SPR-5/LSD1 cooperatively

prevent somatic reprogramming of C. elegans germline stem cells. *Stem cell reports* **2,**

547–59 (2014).

21. Robert, V. J. *et al. The SET-2/SET1 Histone H3K4 Methyltransferase Maintains*

Pluripotency in the Caenorhabditis elegans Germline. *Cell Reports* **9,** (2014).

22. Sung, M., Kawasaki, I. & Shim, Y.-H. Depletion of *cdc-25.3* , a *Caenorhabditis elegans*

orthologue of *cdc25* , increases physiological germline apoptosis. *FEBS Lett.* **591,** 2131–

2146 (2017).

23. Min, H., Shim, Y.-H. & Kawasaki, I. Loss of PGL-1 and PGL-3, members of a family of

constitutive germ-granule components, promotes germline apoptosis in C. elegans. *J. Cell*

Sci. **129,** 341–353 (2016).

24. Alvares, S. M., Mayberry, G. A., Joyner, E. Y., Lakowski, B. & Ahmed, S. H3K4 demethylase

activities repress proliferative and postmitotic aging. *Aging Cell* **13,** 245–253 (2014).

25. Knutson, A. K. K., Egelhofer, T., Rechtsteiner, A. & Strome, S. Germ Granules Prevent

Accumulation of Somatic Transcripts in the Adult Caenorhabditis elegans Germline.

Genetics **206,** (2017).

26.   Updike, D. L., Knutson, A. K., Egelhofer, T. A., Campbell, A. C. & Strome, S. Germ-Granule

Components Prevent Somatic Development in the C. elegans Germline. *Curr. Biol.* **24,**

970–975 (2014).

27.   Updike, D. & Strome, S. P Granule Assembly and Function in Caenorhabditis elegans

Germ Cells. *J. Androl.* **31,** 53–60 (2010).

28.   Phillips, C. M., Montgomery, T. A., Breen, P. C. & Ruvkun, G. MUT-16 promotes formation

of perinuclear mutator foci required for RNA silencing in the C. elegans germline. *Genes*

Dev. **26,** 1433–44 (2012).

29.   Loidl, J. Conservation and Variability of Meiosis Across the Eukaryotes. *Annu. Rev. Genet.*

50, 293–316 (2016).

30.   Hillers K.J., Jantsch V., Martinez-Perez E. & Yanowitz J.L. in (2017).

doi:10.1895/wormbook.1.178.1

31.   R Core Team. R: A language and environment for statistical   computing. *R Foundation for*

Statistical Computing (2013).

32.   Vermezovic, J., Stergiou, L., Hengartner, M. O. & d'Adda di Fagagna, F. Differential

regulation of DNA damage response activation between somatic and germline cells in

Caenorhabditis elegans. *Cell Death Differ.* **19,** 1847–1855 (2012).

Figures and Figure Legends

Figure 1

Figure 1: DNA Damage Response is variable in small RNA genome silencing mutants prior to

and at sterility. (A) Staining of pS/TQ in wildtype, *prg-1(n43) and prg-1(tm)* mutants.

pS/TQ(red) and DAPI(blue) **(B)** Quantification of pS/TQ staining in 20°C wildtype and *prg-1*

mutants at different levels of fertility. **(C)** Quantification of pS/TQ staining in 25°C wildtype and

small RNA genomic silencing mutants at sterility.

Figure 2

Figure 2: At sterility, partial loss of P-granule staining occurred in *prg-1* and temperature-

sensitive mutants. Germlines of sterile Day 2-3 adult animals were stained using the OIC1D4

antibody against P-granules (green) and DAPI (blue). **(A-A')** Control animals contain uniform

puncta of P-granule staining surrounding each nuclei. **(B-E)** Late generation fertile *prg-1, rsd-6,*

nrde-2, and *rbr-2* animals displayed relatively normal P-granule staining in the germlines. **(B'-E')**

prg-1, *nrde-2*, *rsd-6* and *rbr-2* sterile animals all exhibited loss of germ cell P-granules. (N=30)

Figure 3

Figure 3: Disruption of P-granules by RNAi or loss of *pgl-1* resulted in a similar germline

atrophy phenotype at 25°C. (A**)** F2 wild-type (N2) worms on Empty Vector (EV) or P-granule

RNAi at 25°C were scored for sterility, embryonic lethality, or fertility. 80% of worms on P-

granule RNAi were sterile. (B) F1 L4 *pgl-1* mutants and F2 L4 P-granule RNAi treated animals

stained by DAPI contained mostly large germlines while adults exhibited a range of germline

phenotypes from empty to large. *** p<0.0001 Fisher's Chi Square test with Bonferroni

correction n=total germlines scored **(C)** F1 *pgl-1* mutants and F2 P-granule RNAi treated animals

contained oocytes with increased numbers of univalents at sterility as measured by DAPI bodies

n=151 wild-type, 81 P-granule RNAi, 37 *pgl-1*. (D) Stained DAPI germlines of sterile F2 L4 *pgl-1*

contained a range of germline phenotypes from large to empty. Sterile F2 day 1 *pgl-1* mutants

stained by DAPI contained predominantly empty and atrophy germlines. n=total germlines

scored. **(E-H)** Representative DAPI images of germline and oocyte defects in empty vector

treated animals **(E,F)** and P-granule RNAi treated animals **(G,H)** scale bar=10um. **(I)** *pgl-1* F1

worms at 25°C were either DAPI stained as Day 3 adults (blue) or shifted to 20°C at Day 1 and

DAPI stained at Day 3 (red) ** p=0.007 Fisher's Chi Square test with Bonferroni correction.

n=total germlines scored **(I)** No sterile day 1 *pgl-1* F1 at 25°C laid oocytes or progeny (n=35

plates, 350 total worms). Sterile day 1 *pgl-1* shifted to (20°C n=50 plates, 500 worms) gave rise

to plates with oocytes and rarely progeny. Plates with oocytes contained both oocytes and

dead embryos. Plates with progeny contained embryos that hatched to L1.

Figure 4: Model for P-granule dysfunction in small RNA genome silencing mutants

(A**)** Model for small RNA genome silencing mutants B) Model for accumulating heritable stress

in small RNA genome silencing mutants and P-granule function.

Supplemental Figure 1

Supplemental Figure 1: P-granule defects in *prg-1(tm), nrde- 4* and *hrde-1*. Germlines of sterile

Day 2-3 adult animals were stained using the OIC1D4 antibody against P-granules (green) and

DAPI (blue). **(A-A')** Control animals contain uniform puncta of P-granule staining surrounding

each nuclei. **(B-D)** Late generation fertile *prg-1(tm), nrde-4*, and *hrde-1* animals displayed

relatively normal P-granule staining in the germlines. **(B'-D')** sterile *prg-1(tm), nrde-4*, and *hrde-*

1 germ cell P-granules.

Genotype	Normal	Loss (>20%)	p-value
N2	32	0	
hrde-1 (tm1200)	0	31	<.0001
rsd-6 (yp11)	0	32	<.0001
nrde-2(gg95)	0	25	<.0001
rbr-2 (tm1231)	0	32	<.0001

448 **Table 1: Quantification of P-granule defects in the germline**