1

The IMMUNE-ASSOCIATED NUCLEOTIDE-BINDING 9 protein is a regulator of

basal immunity in *Arabidopsis thaliana*

Yuanzheng Wang[1,2], Tabata Rosas-Diaz[1], Carlos Caceres-Moreno[1,2], Rosa Lozano-

Duran[1] and Alberto P. Macho[1,*].

[1]Shanghai Center for Plant Stress Biology, CAS Center for Excellence in Molecular

Plant Sciences; Shanghai Institutes of Biological Sciences, Chinese Academy of

Sciences, Shanghai 201602, China.

[2]University of Chinese Academy of Sciences, Beijing, China.

Running title: IAN9 is a regulator of plant basal immunity

Highlight: IAN9 and its interactor IAP1 act as negative regulators of basal immunity

against bacterial pathogens in *Arabidopsis thaliana*.

* Corresponding author: Alberto P. Macho,

 alberto.macho@sibs.ac.cn; (+86) 02157078280

Submission date: March 6th 2018

Word count: 5956

Figures: 6 (2 in colour in print)

Supporting figures: 19

Supporting tables: 3

Abstract

A robust regulation of plant immune responses requires multitude of positive and negative regulators that act in concert. The immune-associated nucleotide-binding (IAN) gene family members are associated with immunity in different organisms, although no characterization of their function has been carried out to date in plants. In this work, we analyzed the expression patterns of *IAN* genes and found that *IAN9* is repressed upon pathogen infection or treatment with immune elicitors. *IAN9* encodes a plasma membrane-localized protein that genetically behaves as a negative regulator of immunity. A novel *ian9* mutant generated by CRISPR/Cas9 shows increased resistance to *Pseudomonas syringae*, while transgenic plants overexpressing *IAN9* show a slight increase in susceptibility. *In vivo* immunoprecipitation of IAN9-GFP followed by mass spectrometry analysis revealed that IAN9 associates with a previously uncharacterized C3HC4-type RING finger domain-containing protein that we named IAP1, for <u>I</u>AN9-<u>a</u>ssociated <u>p</u>rotein 1, which also acts as a negative regulator of basal immunity. Interestingly, neither *ian9* or *iap1* mutant plants show any obvious developmental phenotype, suggesting that they display enhanced inducible immunity rather than constitutive immune responses. Since both *IAN9* and *IAP1* have orthologs in important crop species, they could be suitable targets to generate plants more resistant to diseases caused by bacterial pathogens without yield penalty.

Keywords: Arabidopsis, CRISPR/Cas9, IAN, plant disease, plant immunity, negative regulation, protein complex, *Pseudomonas syringae*.

Introduction

The plant immune system comprises an intricate network of receptors and regulators aimed at keeping the cellular homeostasis in the absence of pathogen threat and responding rapidly to biotic stimuli in order to prevent infection. Plants have evolved to perceive pathogen-derived molecules that constitute signals of a potential invasion, also called invasion patterns (Cook et al, 2015). Conserved microbial molecules constitute good targets for recognition by plants; some of these molecules have been shown to be perceived by plant cells as pathogen-associated molecular patterns (PAMPs; Boller and Felix, 2009). PAMPs are perceived at the cell surface by transmembrane pattern-recognition receptors (PRRs; Zipfel, 2014). PRRs act in coordination with several regulators and additional proteins that mediate signal transduction (Couto and Zipfel, 2016), including mitogen-activated protein kinases (MAPKs), calcium-dependent protein kinases (CDPKs), receptor-like cytoplasmic kinases (RLCKs), and respiratory burst oxidase homologs (RBOHs) (Macho and Zipfel, 2014; Bigeard et al, 2015). Downstream responses include the production of the phytohormone salicylic acid and antimicrobial compounds, the deposition of callose at the cell wall, and extensive transcriptional reprogramming (Boller and Felix, 2009). The activation of immunity is aimed at preventing the proliferation of the perceived pathogen, and prepares plant cells to mount an efficient defence response against subsequent biotic threats. However, defence is costly, in terms of energy investment and the concomitant disruption to the normal developmental program (Huot et al, 2014; Stael et al, 2015), and as such needs to be tightly regulated. For this reason, immune responses are inducible, and negative regulators ensure a firm control of their activated state. On the other hand, it has been demonstrated that activation of defence and inhibition of growth can be uncoupled, so that active defence and growth can occur simultaneously, indicating that developmental alterations are the consequence of an active process, and not necessarily of limiting resources (De Wit et al, 2013; Campos et al, 2016; Scheres and van der Putten, 2017).

Pathogens have developed strategies to manipulate plant cells in order to proliferate inside plant tissues. These include the suppression of plant immunity, the alteration of the physical environment, and the acquisition of nutrients to support pathogenic lifestyle (Win et al, 2012). In Gram-negative bacterial pathogens, the most important virulence factor is the type-III secretion system (T3SS), which injects effector proteins directly inside plant cells (type-III-secreted effectors, T3Es). The manipulation of

plant cellular functions by T3Es is essential for bacteria to proliferate and is required
for the development of disease (Macho and Zipfel, 2015; Macho, 2016). However,
some plants have evolved intracellular receptors that can perceive T3E activities as
an indication of pathogen invasion, hence becoming resistant to bacterial infection.
These receptors contain nucleotide-binding and leucine-rich repeat domains (NLRs;
Khan et al, 2016). NLR activation contributes to the development of defence
responses similar to those established after PRR activation, but are often more
intense, and sometimes lead to local cell death, which prevents further pathogen
proliferation and spread (Chiang and Coaker, 2015). Both PRRs and NLRs constitute
the basis of plant innate immunity, and are the major determinants of basal immunity
in plants.

Guanosine triphosphate (GTP)-binding proteins are regulators of various biological
processes in eukaryotic cells, such as such as signal transduction, cell proliferation,
cytoskeletal organization, and intracellular membrane trafficking, and are classified
into numerous families (Takai et al, 2001; Vernoud et al, 2003). IMMUNE-
ASSOCIATED NUCLEOTIDE-BINDING/GTPases OF IMMUNITY-ASSOCIATED
PROTEINS (IAN/GIMAP) proteins comprise a sub-family of GTPase-like proteins
that has been found in anthozoans, vertebrates, and angiosperm plants (Nitta and
Takahama, 2007; Weiss et al, 2013). In vertebrates, proteins from the IAN/GIMAP
family regulate the development and homeostasis of T cells and are associated to
autoimmunity (Nitta and Takahama, 2007). The transcriptional regulation of genes
encoding IAN/GIMAP proteins has been linked to immunity in different organisms: in
mice, *IAN/GIMAP* genes are mostly expressed in immune tissues (Nitta et al, 2006),
and they have also been reported as induced in corals after treatment with the
bacterial immune elicitor MDP (Weiss et al, 2013). In Arabidopsis, IAN/GIMAP
proteins are defined by the presence of an avrRpt2-induced gene 1 (AIG1) domain,
and contain conserved GTP-binding domains, including a P-loop motif known to bind
GTP/GDP in Ras GTPases (Bourne et al, 1991), and coiled-coil motifs (Liu et al,
2008). Originally, *AtAIG1* (also known as *IAN8*) was defined as a gene
overexpressed in response to the avirulent bacterial strain *Pseudomonas syringae*
pv. *maculicola* (*Pma*) expressing the effector AvrRpt2 (Reuber and Ausubel, 1996).
Additionally, computational analysis of transcriptomic data has revealed that the
transcription of other *IAN* family members responds to different biotic stimuli:
nematode infection induces the expression of *IAN3* and *IAN11*, while the
transcription of *IAN11*, *IAN12* and *IAN13* is reduced by infection with *Myzus persicae*
(Liu et al, 2008).

Despite the accumulating evidences that associate *IAN* genes with immunity in

plants, no characterization of their function in this process has been carried out to

date. Here, we analyzed the expression patterns of *IAN* genes and found that the

expression of *IAN9* is repressed upon pathogen infection or treatment with immune

elicitors. Further characterization indicated that *IAN9* encodes a plasma membrane-

localized protein and that it genetically behaves as a negative regulator of immunity.

In vivo immunoprecipitation of IAN9-GFP followed by mass spectrometry analysis

revealed that IAN9 associates with a C3HC4-type RING finger domain-containing

protein that we named IAP1, for <u>I</u>AN9-<u>a</u>ssociated <u>p</u>rotein 1. Interestingly, our results

show that IAP1, like IAN9, negatively regulates immunity, raising the idea that these

two proteins may work together in the control of immune responses.

Materials and Methods

Plant materials and growth conditions

Arabidopsis seeds were sterilized with bleach solution (20% bleach and 0.1% Triton

X-100) for 2-3 minutes, then washed with sterile water 4-5 times and sown on solid ½

MS medium (2.21 g Murashige & Skoog Basal Medium with Vitamins, 15 g Sucrose,

7 g Agar. 1 L). After stratification at 4°C for 3 days, the plates were transferred to a

growth chamber (22°C, 16 h light/8 h dark) for germination and growth. For

experiments involving mature Arabidopsis plants, sterile seeds were firstly stratified

at 4°C for 3 days, then transferred to soil. Plants for bacterial infection assays were

cultivated in a short day chamber (22°C, 8 h light/16 h dark photoperiod, 65%

humidity). Plants for *Agrobacterium* transformation were grown in a long day growth

room (22°C, 16 h light/8 h dark photoperiod).

Bacterial infections

Pseudomonas syringae pv tomato DC3000 (*Pst* DC3000) and DC3000 (AvrRpt2)

were streaked on selective ½ salt LB plates (10 g Tryptone, 5 g Yeast Extract, 5 g

NaCl, 15 g Agar per 1 L) and cultivated at 22°C for 2 days. For *Pst* DC3000, bacteria

were collected from the plates into sterile water, the OD was adjusted to a value of

0.1 (5×10^7 cfu/ml) or 0.2 (10^8 cfu/ml), and silwet-L77 was added to a final

concentration of 0.03% before performing spray inoculation onto 3-4 week-old soil-

grown Arabidopsis under short day conditions. The plants were then covered with

cling wrap for 3 hours. The whole plants were harvested in 1.5 ml microcentrifuge

tubes and weighed. For *Pst* DC3000 (AvrRpt2), the OD was adjusted to 0.1 (5×10^7

cfu/ml) or 0.001 (10^5 cfu/ml). The bacterial suspension was pressure-infiltrated into 5-

6 week-old short day-grown Arabidopsis leaves with a needleless syringe (3 leaves

per plant), and leaf discs were collected into 1.5 ml tubes using a leaf punch at 3

days post-inoculation. In both cases, plant tissues were ground and homogenized in

sterile water before plating serial dilutions on selective ½ salt LB agar plates. The

plates were placed at 28°C for 1.5 days and the bacterial growth was calculated as

colony-forming units.

Treatments with immune elicitors

Sterile seeds were sown on ½ solid MS medium to germinate and grown for 3-4 days

in long day conditions. Seedlings were then transferred into ½ liquid MS medium in

12-well culture plates (Thermo Fisher Scientific, Waltham, MA, US), and grown for

another 6-7 days. Every well contained 3 seedlings, and every experiment included 3

independent samples from 3 independent wells. The flg22 peptide or salicylic acid
were added into the liquid medium to a final concentration of 100 nM and 0.5 mM,
respectively. All the plates were incubated on a shaker for 5-10 minutes following
addition of the elicitor. Samples were harvested into 1.5 ml tubes at different time
points after treatment, as indicated in the figures.
184

Root growth assay
Sterile seeds were sown on ½ solid MS medium and stratified at 4°C for 3 days.
Then the plates were placed into a growth chamber for 1 day. The germinated
seedlings were transferred to the new ½ solid MS medium (square plates). Square
plates were placed vertically into a growth chamber and root length was measured 5
days later.
191

RNA isolation, RT-PCR, and RT-qPCR
For RNA extraction, plant tissues were collected in 1.5 ml microfuge tubes with one
metal bead and the tubes were immediately placed into liquid nitrogen. Samples
were ground thoroughly with a TissueLyser (QIAGEN, Duesseldorf, Nordrhein-
Westfalen, Germany) for 1 minute, and placed back in liquid nitrogen. Total RNA was
extracted with the E.Z.N.A. Plant RNA kit (Omega Bio-Tek, Norcross, GA, US) with
in-column DNA digestion and an additional sample treatment with DNAase (Omega
Bio-Tek, Norcross, GA, US). First-strand cDNA was synthesized using the iScript
cDNA synthesis kit (Bio-Rad, Hercules, CA, US) with a volume of 20 µl. For RT-PCR,
the PCR reaction was performed in 50 µl using the Q5 Hot Start High-Fidelity DNA
polymerase (New England Biolabs, Ipswich, MA, US) with 35 or 38 cycles. For
quantitative RT-PCR (RT-qPCR) , PCR reactions were performed in 20 µl using the
iTaq Universal SYBR Green Supermix kit (Bio-Rad, Hercules, CA, US) in the
StepOnePlus Real-Time PCR System (Applied Biosystems, Foster City, CA, US).
Data were analyzed with Excel and GraphPad Prism 6.
207

Confocal microscopy imaging
Cotyledons from 3-4 day-old long day-grown Arabidopsis seedlings were imaged
using a Confocal Laser Scanning Microscope (CLSW) Platform: Leica TCS SP8
(Leica, Mannheim, Germany). The GFP was excited with an argon laser at 488 nm,
and its emission was detected at 500-550 nm. For plasmolysis assays, cotyledons
were placed into a 1 M NaCl solution on glass slides, and GFP was observed and
recorded after 5-10 minutes. For FM4-64 staining, cotyledons from 4 day-old
seedlings were cut and soaked into 5 ng/µl FM4-64 solution as described previously

(Beck et al., 2012) for 5 minutes; samples were then transferred to water on glass
slides and covered with coverslips. For dual channel simultaneous observation, the
fluorescence signal of FM4-64 was excited with an argon laser at 561 nm and its
emission was observed at 580-650 nm; the GFP was excited at 488 nm and
observed at 500-550 nm. For co-localization assays, 3 week-old *N. benthamiana*
leaves were co-infiltrated with *Agrobacterium* GV3101 carrying plasmids to express
RFP-IAN9 and *GFP-IAP1*. Leaves were co-infiltrated with the GV3101 strain carrying
plasmids to express *RFP-IAN9* and *GFP* as control. Three-mm leaf discs were
punched from the whole leaf 24 hours post infiltration and transferred to water on
glass slides. For dual channel image acquisition, GFP and RFP were excited at 488
nm and 561nm respectively; emission was collected at 500-550 nm for GFP and
580-650nm for RFP.
228

MAPK activation and western blot assays
For protein extraction, Arabidopsis seedlings or leaf discs from *N. benthamiana* were
ground with a Tissue Lyser (QIAGEN, Hilden, Nordrhein-Westfalen, Germany).
Samples were then re-suspended in lysis buffer [2x loading buffer: 100 mM Tris-HCl
(pH 6.8), 10% Glycerol, 2% SDS, 0.03% bromophenol blue], boiled at 95°C for 10
minutes, and centrifuged at 14,000 g for 5 minutes before loading in SDS-PAGE
acrylamide gels. Western blots were performed using anti-GFP (Sigma G6795), anti-
Luciferase (Sigma L0159), anti-Mouse IgG-Peroxidase (Sigma A2554), and anti-
Rabbit IgG-Peroxidase (Sigma A0545).
MAPK activation assays were performed as previously described (Macho et al,
2012), with minor modifications. Briefly, 7 day-old Arabidopsis seedlings grown on
solid ½ MS were transferred to water and then treated with 100 nM flg22 for 10
minutes after vacuuming 5 minutes (15 minutes in total). Anti-pMAPK [Phospho-
p44/42 MAPK (Erk1/2) (Thr202/Tyr204) XP Rabbit mAb; Cell Signaling 4370] was
dissolved in 3% gelatin (Sigma G7041) and used to hybridize the membranes. All
membraneswere stained with Ponceau stain (Sangon Biotech, Shanghai, China) to
verify equal loading.
246

ROS burst
ROS burst assays were conducted as previously described (Sang and Macho, 2017).
Briefly, 4 mm leaf discs from 5-6 week-old Arabidopsis plants grown in short day
conditions were transferred to 96-well microplates (PerkinElmer, Waltham, MA, US)
with 100 µl Mili-Q water per well and incubated overnight. Water was then removed
and ROS burst was elicited by adding 100 µl of a solution containing 100 ng flg22,

100 µM luminol, and 20 µg/mL horseradish peroxidase. The luminescence was

recorded over 40 minutes using a Thermo Scientific VARIOSKAN FLASH (Thermo

Fisher Scientific, Waltham, MA, US).

Co-IP and large-scale IP for LC-MS/MS

Leaves from 3-4 week-old *N. benthamiana* plants were co-infiltrated with

Agrobacterium GV3101 carrying plasmids to express *GFP-IAP1* and *Cluc-IAN9*;

leaves co-infiltrated with GV3101 carrying plasmids to express *GFP-AIP1* and *Cluc*

or *GFP* and *Cluc* were used as controls. Total proteins were extracted 24 hours later,

and immunoprecipitation was performed with GFP-trap beads (Chromotek, Am

Klopferspitz, Planegg-Martinsried, Germany) as described previously (Sang et al,

2016). Proteins were stripped from the beads by boiling in 50 µl SDS loading buffer

for 10 mins. Immunoprecipitated proteins were separated on SDS-PAGE acrylamide

gels and western blots were performed as described above. Large-scale

immunoprecipitation assays for LC-MS/MS were performed as described before

(Kadota et al. 2016; Sang et al. 2016), using 5 g of 10 day-old Arabidopsis seedlings

before or after treatment with 100 nM flg22 or 0.5 mM SA.

Results

Expression analysis of the *IAN* gene family upon bacterial infection reveals differential expression patterns for *IAN7, IAN8,* and *IAN9*

The IAN protein family in Arabidopsis is composed of 13 members (Liu et al, 2008; Figure 1A). Phylogenetic analysis of IAN amino acid sequences shows a clear separation in different subgroups (Figure 1A, Figure S1). Previous reports have described changes in expression of *IAN* gene family members upon different biotic stimuli (Reuber and Ausubel, 1996; Liu et al, 2008). In order to characterize the transcriptional response of *IAN* genes to bacterial infection, we inoculated Arabidopsis seedlings with the virulent pathogen *Pseudomonas syringae* pv. *tomato* (*Pst*) DC3000. Three *IAN* genes showed differential expression patterns upon bacterial infection: *IAN7* and *IAN8* showed a significant up-regulation, while *IAN9* showed a significant down-regulation (Figure 1B). Regarding other *IAN* members, we did not detect mRNA for *IAN1/2/4/6/10/11/12/13,* suggesting that these genes are not expressed in 10 day-old Arabidopsis seedlings in our experimental conditions, while transcript abundance for *IAN3/5* was not influenced by pathogen infection (data not shown). The pathogen-induced up-regulation of *IAN8* is reminiscent of the originally reported up-regulation of this gene by bacteria expressing the effector AvrRpt2, which induces activation of the plant NLR RPS2 (Reuber and Ausubel, 1996). Accordingly, we found that the differential expression of *IAN7/8* and *IAN9* also takes place upon infection of Arabidopsis rosette leaves with *Pst* expressing *AvrRpt2* (Figure 1C). The particular expression pattern of *IAN9* among the *IAN* gene family suggests an exclusive function for IAN9 rather than functional redundancy with other IAN family members. This idea is supported by the fact that *IAN9* constitutes a specific phylogenetic group within the *IAN* gene family (Figure 1A). For these reasons, we decided to focus our attention on IAN9.

***IAN9* expression is reduced upon chemical activation of plant immunity**

IAN9 is broadly expressed in cotyledons, hypocotyls, and roots of Arabidopsis seedlings (Figure S2). To dissect the bacteria-induced repression of *IAN9* transcription, we sought to determine whether perception of purified immune elicitors affects *IAN9* expression. For this purpose, we first treated Arabidopsis seedlings with the flagellin-derived peptide flg22, which is widely used as an elicitor of immune responses. Our results show that *IAN9* expression is significantly reduced one hour after flg22 treatment (Figure 2A). The perception of different invasion patterns, including flg22, leads to the production of the phytohormone salicylic acid (SA),

which is a key player in the activation of plant immunity against biotrophic pathogens
(Vlot et al., 2009). Treatment with SA for three hours led to a reduction of *IAN9*
expression, although the abundance of *IAN9* transcripts returned to basal levels after
a 6-hour SA treatment (Figure 2B), revealing a transient down-regulation of *IAN9*
transcript abundance upon SA treatment. Finally, to determine whether the reduction
on *IAN9* transcription upon flg22 treatment depends on the downstream SA
accumulation (Tsuda et al., 2008), we performed flg22 treatment in the SA-depleted
sid2/NahG plants, deficient in pathogen-induced SA biosynthesis (Wildermuth et al.
2001) and constitutively expressing the bacterial salicylate hydroxylase NahG, which
degrades SA (Delaney et al., 1994). Interestingly, the flg22-triggered down-regulation
of *IAN9* transcript abundance was also observed in *sid2/NahG* plants, suggesting
that SA is not required for the flg22-induced transcriptional repression of *IAN9*.
320
IAN9 localizes to the plasma membrane through its C-terminal part
In order to investigate the subcellular localization of IAN9, we generated stable
transgenic Arabidopsis lines expressing an N-terminal GFP-tagged IAN9 protein (see
below for a detailed characterization of these lines), and used confocal microscopy to
determine the localization of GFP-IAN9. Contrary to free GFP, which shows a
nuclear/cytoplasmic localization, GFP-IAN9 specifically localized at the cell periphery
(Figure 3A). This localization was similar to that observed for well-characterized
plasma membrane (PM)-localized proteins, such as the brassinosteroid (BR)
receptor BRI1 (Wang et al., 2001; Figure 3A). To determine whether GFP-IAN9 is
associated to membranes, we used the lipophilic fluorescent dye FM4-64, which is
rapidly incorporated into membranes upon contact with plant cells (Fischer-Parton et
al., 2000; Bolte et al., 2004). Our results show that GFP-IAN9 fluorescence co-
localizes with FM4-64-labeled compartments (Figure 3B and 3C), similar to BRI1-
GFP fluorescence, and different from free GFP (Figure 3B and 3C). To further
confirm that IAN9 localizes at the PM, we performed plasmolysis assays by treating
plant tissues with 1 M NaCl. Upon plasmolysis, both GFP-IAN9 and BRI1-GFP were
detected in Hechtian strands, which are associated to PM retractions from the cell
wall (Figure 3D). Altogether, our microscopy analysis indicates that IAN9 localizes at
the PM in plant cells.
The C-terminal domain is not conserved among IAN proteins (Figure S1).
Interestingly, the C-terminal sequence of IAN9 shows an over-representation of
positively charged amino acids (KKLRENLERAEKETKELQKKLGKCINL; 33.3% of
R/K), not present in other IAN proteins (Figure S1 and S3A), which could mediate an
interaction with the negatively charged phospholipids of the PM. To test this

hypothesis, we generated Arabidopsis stable transgenic lines expressing a truncated

GFP-IAN9 version lacking the 27 C-terminal amino acids (IAN9ΔC-27). The IAN9ΔC-

27 version lost the specific PM localization seen for wild type GFP-IAN9, and was

mostly detected in the cytoplasm (Figure S3B). Additionally, we found that, when

GFP is fused to the C-terminal end of IAN9 (IAN9-GFP), this protein loses its specific

PM localization, and is mostly found in the cytoplasm (Figure S3B). Compared to

IAN9, the IAN8 C-terminal part has a lower representation of positively charged

aminoacids (18.5%; Figure S3A), and we found that a N-terminal GFP-tagged IAN8

(GFP-IAN8) localizes to the cytoplasm. Altogether, these results suggest that the

exclusive C-terminal domain of IAN9 is required for its localization at the PM.

Generation of *IAN9* knock-out and overexpression lines

Public repositories for Arabidopsis T-DNA insertion lines contain four independent

lines with T-DNA insertions in the *IAN9* genomic locus: *SAIL_167_B02*,

SALK_534_B01, *SALK_144369,* and *GK-146B08*. For all these lines, the insertion

site is located in the predicted promoter region of *IAN9*. Among them, *GK-146B08* is

the line harboring the nearest T-DNA insertion to the *IAN9* start codon (Figure S4A),

and therefore we chose this line for further analyses. Sequencing analysis results

showed that the insertion site is located 76 bp upstream of the 5'-UTR of the *IAN9*

gene, and 319 bp upstream of the *IAN9* start codon (Figure S4A and S4B). RT-PCR

and RT-qPCR analyses showed that this mutant has approximately a 3-fold

reduction in the amount of *IAN9* transcripts (Figure S4C and S4D), indicating that this

line is a knockdown *ian9* mutant. To further characterize this line, we determined the

IAN9 transcript levels upon bacterial inoculation. Surprisingly, we found that *IAN9*

transcript levels increased significantly upon inoculation with *Pst* or *Pst* (AvrRpt2),

reaching higher levels than those observed in wild type plants (Figure S4E) and an

opposite regulation (Figure 1). These findings indicate that the T-DNA insertion in the

IAN9 promoter generates an aberrant expression pattern of *IAN9* in this line,

rendering it unusable for the functional characterization of this gene.

In order to perform genetic analysis of the contribution of IAN9 to plant immunity, we

generated *ian9* mutant lines using CRISPR/Cas9-assisted genome editing (Feng et

al, 2013; Mao et al, 2013). We selected the best predicted target site in the *IAN9*

gene sequence for recognition by the Cas9/sgRNA complex (Figure S5), and

performed the targeted mutagenesis as explained in the methods section.

Sequencing of the resulting line showed an addition of 1 base pair in the second

exon of *IAN9*, creating a premature stop codon 366 base pairs downstream of the

start codon (Figure S6A), which generates a truncated protein with a disrupted GTP-

binding domain. Upon selection of seedlings containing the *ian9* mutation in
homozygosis, we selected a line for which where the Cas9 gene was segregated out
(Figure S6B); this Cas9-free *ian9* line was used for further experiments. Additionally,
as mentioned before, we generated Arabidopsis transgenic lines overexpressing
IAN9 in a Col-0 wild type background, using a 35S promoter to express a *GFP-IAN9*
fusion. We selected two independent homozygous lines that accumulated detectable
amounts of GFP-IAN9 fusion protein (Figure S6C), and higher transcript levels of
IAN9 compared to those in Col-0 wild type (OE-*GFP-IAN9#3-10* and OE-*GFP-*
IAN9#7-1) (Figure S6D). As controls, we selected two independent homozygous
lines expressing free GFP, which did not show changes in *IAN9* transcription and
accumulated detectable levels of free GFP (Figure S6C and S6D). Interestingly,
although the 35S promoter led to high *IAN9* expression compared to Col-0 wild type,
bacterial infection still caused a reduction in *IAN9* transcript levels (Figure S7),
suggesting a post-transcriptional regulation of the abundance of *IAN9* transcripts.
Overexpression of *IAN9* did not affect *IAN8* expression in basal conditions or upon
bacterial infection (Figure S7).
398

Plant growth and early immune responses are not affected by altered
expression of *IAN9*
Both *IAN9* overexpressing or *ian9* knockout seedlings and adult plants were visually
indistinguishable from the wild type (Figure S8 and S9). Given the predicted
association of the IAN family to immune responses and the alteration in *IAN9*
expression upon elicitation with flg22, we sought to determine whether IAN9 is
involved in early PTI responses, namely the flg22-triggered burst of reactive oxygen
species (ROS), and the activation of a cascade of mitogen-activated protein kinases
(MAPKs) (Boller & Felix, 2009, Macho & Zipfel, 2014). Our results show that
mutation or overexpression of *IAN9* did not have a detectable impact in the dynamics
or total accumulation of ROS upon flg22 treatment (Figure S8B, S8C and Figure
S10). Similarly, neither mutant nor overexpression (OE) lines showed differences in
flg22-triggered MAPK activation compared to wild type plants or free GFP-expressing
controls (Figure S8D and S8E). Taken together, these results suggest that
alterations of *IAN9* expression do not affect early PTI responses.
414

IAN9 negatively regulates plant immunity against *Pst* DC3000
To test whether IAN9 contributes to plant immunity against bacterial pathogens, we
performed surface inoculation of the *ian9* mutant line with *Pst* DC3000 and
determined bacterial replication in plant tissues. Our results show that knockout

mutation of *IAN9* increased plant resistance against *Pst* DC3000, causing a 13-fold

reduction on bacterial titers (Figure 4A). In order to test whether the increase in

resistance also occurs in a context of ETI, we syringe-infiltrated Arabidopsis rosette

leaves with *Pst* (AvrRpt2). However, no differences were detected in terms of

replication of this strain (Figure 4B). To confirm that the observed increased

resistance to *Pst* DC3000 is really due to the absence of *ian9*, we generated

complementation lines expressing *GFP-IAN9*, driven by a *35S* promoter, in the *ian9*

knockout mutant background, where the GFP-IAN9 protein accumulated and

localized to the PM (Figure S11A and S11B). Expression of GFP-IAN9 in the *ian9*

background was able to rescue the level of growth of *Pst* DC3000 to that observed in

wild type plants (Figure S11C), confirming the association of IAN9 with the observed

increased resistance. Interestingly, transgenic lines overexpressing GFP-IAN9 in a

wild type background showed a tendency to support higher bacterial loads compared

to wild type or GFP-expressing lines, suggesting that IAN9 overexpression

suppresses plant immunity against *Pst* DC3000, although such tendency was not

always reproducible or statistically significant across eight independent biological

repeats (Figure S12).

Identification of IAN9-interacting proteins

To characterize further the mode of action of IAN9, we searched for proteins

physically associated with IAN9 in plant cells using Arabidopsis seedlings expressing

GFP-IAN9 (line OE-*GFP-IAN9#3-10*) and seedlings expressing free GFP as control.

Upon GFP immunoprecipitation (IP) using agarose beads coupled to an anti-GFP

nanobody (GFP-Trap beads), we detected associated proteins using liquid

chromatography coupled to tandem mass-spectrometry (LC-MS/MS). In order to

detect potential dynamic interactions, we additionally treated seedlings for one hour

with flg22 or SA before immunoprecipitation. Two independent biological replicates

showed a large number of proteins physically associated with IAN9, which we filtered

using the following criteria: 1. Presence in both biological replicates; 2. Detection of

two or more exclusive unique peptides; 3. Absence in the GFP control. After applying

these filters, a total of 14 proteins were identified as IAN9 candidate interactors

(Table S1).

Among these candidate interactors, we drew our attention to an uncharacterized

protein encoded by the AT1G18660 gene (Table S1), which we named IAP1 for

IAN9-ASSOCIATED PROTEIN1. Interestingly, IAP1 associated with IAN9 in both

replicates of IP + LC-MS/MS, but the number of detected IAP1 peptides decreased in

samples treated with SA, while it did not change significantly in samples treated with
flg22 (Table S1). Domain analysis predicts the presence of three tetratricopeptide-
like helical (TPR) domains, a C3HC4-type RING finger domain, and an ATP-
dependent protease La (LON) domain in this protein (Figure S13A). Upon
Agrobacterium-mediated transient expression in *Nicotiana benthamiana*, a GFP-IAP1
fusion protein localized at the cell periphery, nucleus (weakly), and around the
nucleus; the latter most likely corresponds to the usual endoplasmic reticulum
localization of PM-localized proteins overexpressed in this system (Figure S13B and
S13C). Co-expression with RFP-IAN9 showed that both proteins co-localize upon
transient expression (Figure 5A). To confirm the interaction between IAN9 and IAP1,
we performed a targeted co-IP using GFP-IAP1 and a N-terminal fusion of IAN9 to
the C-terminal domain of luciferase (Cluc-IAN9). Co-IP assays show that GFP-IAP1
interacts with Cluc-IAN9, but not with Cluc alone (Figure 5B); Cluc-IAN9 does not
interact with free GFP (Figure S14). To determine whether the IAN9-IAP1 interaction
is direct, we performed a luciferase complementation assay, transiently co-
expressing Cluc-IAN9 and IAP1 fused to the N-terminal domain of luciferase (IAP1-
Nluc). A positive control co-expressing AtSGT1b-Nluc and Cluc-AtRAR1 showed a
strong luciferase signal (Figure S15A), as described before (Chen et al., 2008). On
the contrary, tissues co-expressing Cluc-IAN9 and Nluc-IAP1 did not show any
detectable luciferase signal (Figure S15A), although both proteins accumulated
(Figure S15B). As an alternative technique to detect direct interactions, we employed
FRET-FLIM by co-expressing GFP-IAP1 and RFP-IAN9. However, no difference in
GFP fluorescence lifetime was detected when compared to control samples (Figure
S15C). Although an influence of the tags cannot be ruled out, these results suggest
that the interaction observed for IAP1 and IAN9 is most likely indirect.

IAP1 negatively regulates plant immunity against *Pst* DC3000
Public repositories for Arabidopsis T-DNA insertion lines contain two independent
lines with T-DNA insertions in the *IAP1* genomic locus: *SALK_119114* and
SALK_093498 (Figure S16A). We isolated homozygous lines containing these
insertions (Figure S16B) and confirmed the absence of *IAP1* transcripts (Figure
S16C), naming these lines *iap1-1* and *iap1-2*, respectively (Figure S16A). Both lines
displayed wild-type-like growth and development when grown on soil under short-day
conditions (Figure 6A), although mutant seedlings showed a slightly reduced root
length compared to Col-0 wild type when grown in vertical MS plates (Figure S17).
To determine whether mutations in *IAP1* have an impact on plant resistance against
bacterial pathogens, we performed surface inoculation of the *iap1* mutant lines with

Pst DC3000. Our results show that both *iap1-1* and *iap1-2* mutant lines are more

resistant than Col-0 wild type against *Pst* DC3000, showing a 19-fold reduction on

bacterial loads (Figure 6B). However, none of these lines showed differences after

inoculation with *Pst* expressing AvrRpt2 (Figure 6C). Interestingly, these results

resemble those obtained during the characterization of the *ian9* knockout mutant line

(Figure 4), suggesting that both proteins are involved in the negative regulation of

basal immunity against bacterial pathogens, and may act in the same pathway

through physical association.

Discussion

In this work, we characterized the transcription patterns of *IAN* family members in Arabidopsis upon infection with the bacterial pathogen *Pto DC3000*. *IAN7* and *IAN8*, which seem to form a separate phylogenetic subgroup, are up-regulated upon bacterial infection. On the contrary, *IAN9* seems to form a specific subgroup and shows a unique down-regulation upon bacterial infection. Infection with a virulent bacterial pathogen, such as *Pto DC3000*, triggers changes in plant cells caused by bacterial virulence activities (e.g. alteration of plant processes by T3Es) and basal immune responses. Therefore, formally, transcriptional changes upon infection could be associated with bacterial virulence or plant immunity. In this case, the down-regulation of *IAN9* expression upon bacterial infection seems associated with the activation of immunity, since we detected similar expression patterns upon treatment with elicitors of immunity, namely flg22 and SA. Interestingly, we detected a reduction in the amount of *IAN9* transcripts after bacterial infection even in transgenic lines where *IAN9* overexpression is driven by a constitutive *35S* promoter, suggesting a post-transcriptional regulation of *IAN9* transcript abundance. Although the perception of flg22 leads to the production of SA (Tsuda et al., 2008), our data show that the flg22-triggered reduction of *IAN9* transcription takes place in *sid2/NahG* plants, indicating that this transcriptional change is independent of the observed SA-triggered reduction of *IAN9* transcription. SA-deficient mutants are partially affected in flg22-triggered induction of defence-related genes, probably caused by lower levels of the flg22 receptor FLS2 (Yi et al., 2014). Our results indicate that these lower levels of FLS2 in *sid2/NahG* plants are nevertheless sufficient to trigger the *IAN9* transcriptional repression, suggesting that *IAN9* transcription may be highly sensitive to external biotic stimuli.

Besides showing different expression patterns, the subcellular localization of IAN9 (mostly at the PM) is also different from that of IAN8 (mostly in the cytoplasm). This differential localization further suggests distinct functions for these two IAN family members. The C-terminal domain of IAN9, required for PM localization, is not present in other IAN proteins. This could indicate that the IAN9 mechanism for PM attachment is exclusive within the IAN family, although other IAN proteins could localize at the PM through different means.

Mutation of *IAN9* causes increased resistance to *Pto* DC3000, and this phenotype is rescued in complementation lines. It is worth considering the possibility that IAN9 is

guarded by NLRs as it has been shown for some regulators of immunity (Khan et al,
2016), therefore explaining the increased resistance in the *ian9* knockout mutant.
However, the absence of developmental phenotypes or constitutive immune
responses makes this possibility unlikely. Additionally, overexpression lines show a
tendency for increased susceptibility, although this phenotype was not always
significant or reproducible. The lack of robustness in this phenotype could suggest
that *IAN9* transcript abundance is not rate limiting; however, it could also be due to
the potential post-transcriptional regulation of *IAN9* transcript abundance that we
have seen in the *IAN9* overexpression lines, which show a reduction in the amount of
IAN9 transcripts after bacterial infection. The transient nature of *IAN9* transcriptional
repression upon SA treatment, returning to basal levels after 6 hours, may indicate
that the negative regulation exerted by IAN9 could be required to contribute to the
repression of immune responses after immunity has been established.

Using IP followed by LC-MS/MS, we found IAP1 as a protein associated with IAN9.
IAP1 contains three tetratricopeptide-like helical (TPR) domains, a C3HC4-type
RING finger domain, and an ATP-dependent protease La (LON) domain. Upon
transient expression, IAP1 co-localizes with IAN9 in *N. benthamiana* cells. Genetic
analysis indicates that both IAN9 and IAP1 negatively regulate immunity against *Pto*
DC3000. Given their physical association, it is possible that both proteins belong to a
protein complex involved in the negative regulation of basal immunity against
bacterial pathogens. Although we confirmed the association between IAN9 and IAP1
using targeted IP, we failed to detect a direct physical interaction using luciferase
complementation or FRET-FLIM assays. Specific limitations of interaction techniques
may be hindering the detection of a direct interaction; however, these data may also
indicate that the interaction between IAN9 and IAP1 is indirect, perhaps mediated by
other scaffolding components in the same protein complex. The Arabidopsis
Interactome Database (http://interactome.dfci.harvard.edu/A_thaliana/index.php) has
reported several interactors for IAP1 in Y2H assays (Table S2). Interestingly, several
of these interacting proteins are transcription factors, and others have predicted
nuclear localization or undergo nucleo-cytoplasmic re-localization associated to their
activity (Table S2). Considering that IAP1 interacts with several proteins with nuclear
activities, it is tempting to speculate that the IAN9/IAP1 complex could associate with
transcriptional regulators in the absence of biotic stress, acting as negative
regulators of immune responses, and the complex may dissociate upon pathogen
infection to allow for the activation of executor immune responses that restrict
pathogen proliferation. This hypothetical model is in agreement with the observation

that IAN9 does not seem to regulate early elicitor-induced responses (namely flg22-
triggered ROS burst and MAPK activation). Given that *ian9* or *iap1* mutants do not
show auto-immunity phenotypes, the final activation of immune responses may
require additional activation by other immune regulators, of which the activity could
be facilitated by the absence of IAN9 or IAP1.

Altogether, our results reveal IAN9 and IAP1 as suitable targets for biotechnological
approaches to generate crops with increased disease resistance to bacterial
pathogens, since both IAN9 and IAP1 have orthologs in agriculturally important crop
species (Figure S18 and S19). Current regulations hinder the use of transgenic
plants to generate disease-resistant crops. Alternative strategies consider the
mutation of negative regulators of immunity, although this often leads to fitness costs
or auto-immunity phenotypes, which obstruct their applicability in agriculture.
Interestingly, IAN9 and IAP1 behave as negative regulators of basal immunity, since
ian9 and *iap1* knockout plants show increased resistance to bacterial infection, but
they do not show obvious differences compared to wild-type plants in terms of size or
development. Our *IAN9* mutagenesis approach shows that it is feasible to design
CRISPR/Cas9-mediated strategies to obtain stable non-transgenic mutant plants
with increased resistance to pathogens and no obvious developmental defects,
paving the way to a potential application in breeding for disease resistance.

Acknowledgements
We thank members of the Macho and Lozano-Durán laboratories for helpful
discussions; Jianming Li, Jian-Min Zhou and Jian-Kang Zhu for sharing biological
materials; and Xinyu Jian for technical and administrative assistance during this
work. We thank the PSC Cell Biology core facility for assistance with confocal
microscopy, and the PSC Proteomics core facility for LC-MS/MS analysis. This work
was supported by the Shanghai Center for Plant Stress Biology (Chinese Academy
of Sciences). Research in the Macho laboratory is also supported by the National
Natural Science Foundation of China (NSFC; grant 31571973) and the Chinese 1000
Talents Program. Research in the Lozano-Durán laboratory is also supported by the
National Natural Science Foundation of China (NSFC; grant 31671994) and the 100
Talents Program from the Chinese Academy of Sciences. CCM is sponsored by a
CAS-TWAS President's Fellowship for International PhD Students. TR-D is the
recipient of a President's International Fellowship Initiative (PIFI) postdoctoral
fellowship (No. 2016PB042) from the Chinese Academy of Sciences. The authors
have no conflict of interest to declare.
613

Supplementary data

Figure S1. Alignment of IAN proteins.

Figure S2. Expression of *IAN9* in *Arabidopsis* seedlings.

Figure S3. The C-terminal region of IAN9 is important for its PM localization.

Figure S4. Characterization of the GK-146B08 line with an insertion in the

promoter of *IAN9*.

Figure S5. Exact mutation site in the *ian9* mutant generated by CRISPR/Cas9.

Figure S6. Characterization of the *ian9* mutant and *IAN9* over-expression lines.

Figure S7. Transcription of *IAN8* is not affected by over-expression of *IAN9*.

Figure S8. Plant growth and early immune responses are not affected by

altered expression of *IAN9*.Figure S9. Seedlings of the *ian9* mutant and *IAN9*

over-expression do not show developmental defects.

Figure S10. Flg22-triggered ROS dynamics are not affected by alterations in

***IAN9* expression.**

Figure S11. Analysis of the *ian9/35S::GFP-IAN9* complementation line.

Figure S12. Transgenic lines overexpressing GFP-IAN9 in a wild type

background show a tendency to support higher bacterial loads compared to

wild type or GFP-expressing lines.

Figure S13. Characterization of IAP1.

Figure S14. Cluc-IAN9 associates with GFP-IAP1 but not with free GFP in *N.*

***benthamiana* leaves.**

Figure S15. IAN9 and IAP1 do not interact directly in *N. benthamiana* leaves.

Figure S16. Characterization of the *iap1* mutant alleles.

Figure S17. Seedlings of the *iap1* mutant lines show slightly reduced root

elongation.

Figure S18. *IAN9* orthologs in different plant species.

Figure S19. *IAP1* orthologs in different plant species.

Table S1. Proteins associated to GFP-IAN9 identified after affinity purification

followed by LC-MS/MS analysis.

Table S2. IAP1 interactors according to the Arabidopsis Interactome Database.

Table S3. List of primers used in this study

Methods S1

Figure Legends

Figure 1. Transcription patterns of the *IAN* gene family in Arabidopsis.

(A) Phylogenetic tree of IAN proteins. (B and C) Relative expression of *IAN7, IAN8*

and *IAN9* in 9-10 days-old Arabidopsis seedlings, 6 hours after inoculation with *Pst*

DC3000 (B) or *Pst* DC3000 (AvrRpt2) (C). Real-time quantitative PCR results were

normalized with *UBIQUITIN 10* (*UBQ10*, AT4G05320). Values are means ± SEM

(n=3). Statistical differences were calculated using one-way ANOVA (p<0.01). Each

experiment was performed three times with similar results.

Figure 2. Treatment with flg22 or salicylic acid causes a reduction on the

transcription of *IAN9*.

Relative expression of *IAN9* in 9-10 days-old Arabidopsis seedlings. Seedlings in (A)

and (C) were treated with 100 nM flg22 for 1 hour. Seedlings in (B) were treated with

0.5 mM salicylic acid for 3 or 6 hours. Real-time quantitative PCR results were

normalized with *UBIQUITIN 10* (*UBQ10*, AT4G05320). Values are means ± SEM

(n=3). Statistical differences were calculated using one-way ANOVA (p<0.01). Each

experiment were performed three times with similar results.

Figure 3. GFP-IAN9 localizes to the plasma membrane in *Arabidopsis*

(A) Confocal images of GFP, GFP-IAN9, and BRI1-GFP in *Arabidopsis* cotyledon

epidermal cells. Bar=10 μm. (B) Confocal images of GFP, GFP-IAN9, and BRI1-GFP

in *Arabidopsis* cotyledon epidermal cells. *Arabidopsis* seedlings were treated FM4-64

for 5 minutes before confocal imaging, and the FM4-64 signal is shown in red. Bar=5

μm. (C) Fluorescence intensity through the thin lines shown in (B). (D) Confocal

images of GFP, GFP-IAN9, and BRI1-GFP in *Arabidopsis* cotyledon epidermal cells

after plasmolysis (5-minute treatment with 1M NaCl). Red arrows indicate the

presence of Hectian strands. Bar=10 μm.

Figure 4. IAN9 negatively regulates plant immunity against *Pst* DC3000.

(A) Growth of surface (spray)-inoculated *Pto* DC3000 (OD_{600}=0.1) in wild-type (WT)

Col-0 and *ian9* mutant plants, 3 days post-inoculation (dpi). Experiments repeated

more than three times with similar results. (B) Growth of *Pto* DC3000 (AvrRpt2)

(OD_{600}=0.001) infiltrated with a needleless syringe into wild-type (WT) Col-0 and *ian9*

mutant plants, 3 days post-inoculation (dpi). Experiments performed twice with

similar results. (A and B) Data were represented as means ± SE (n=8). Statistical

differences were calculated using a Student's t-test. "ns" indicates no significant
difference, and asterisk indicated significant difference (p<0.05).
688

Figure 5. IAP1 interacts with IAN9 in *N. benthamiana* leaves.
(A) Confocal microscopy images showing the co-localization of GFP-IAP1 and RFP-
IAN9 in *N. benthamiana* leaves. Bar=10 μm. (B) Cluc or Cluc-IAN9 was co-
expressed with GFP-IAP1 in *N. benthamiana* before immunoprecipitation using GFP-
trap beads. Immunoblots were analysed using anti-luc or anti-GFP antibody.
Molecular weight (kDa) marker bands are indicated for reference. The experiments
were repeated three with similar results.
696

Figure 6. IAP1 negatively regulates plant immunity against *Pst* DC3000.
(A) Photography of four week-old Col-0 wild type (WT), *iap1-1*, and *iap1-2* plants,
grown at a 8 h light/16 h dark photoperiod. Scare bar is 0.5 cm. (B) Growth of surface
(spray)-inoculated *Pto* DC3000 (OD_{600}=0.1) in wild-type (WT) Col-0, *iap1-1,* and *iap1-
2* mutant plants, 3 days post-inoculation (dpi). Experiments repeated more than three
times with similar results. (C) Growth of *Pto* DC3000 (AvrRpt2) (OD_{600}=0.001)
infiltrated with a needleless syringe into wild-type (WT) Col-0, *iap1-1,* and *iap1-2*
mutant plants, 3 days post-inoculation (dpi). Experiments performed twice with
similar results. (B and C) Data were represented as means ± SE (n=8). Statistical
differences were calculated using a Student's t-test. "ns" indicates no significant
difference, and asterisk indicated significant difference (p<0.05).

Bibliography

Beck M, Zhou J, Faulkner C, MacLean D, Robatzek S. 2012. Spatio-temporal cellular

dynamics of the Arabidopsis flagellin receptor reveal activation status-dependent

endosomal sorting. *Plant Cell* 24: 4205-4219.

Bigeard J, Colcombet J, Hirt H. 2015. Signaling mechanisms in pattern-triggered

immunity (PTI). *Mol Plant* 8: 521-539.

Boller T, Felix G. 2009. A Renaissance of Elicitors: Perception of Microbe-Associated

Molecular Patterns and Danger Signals by Pattern-Recognition Receptors. Annu Rev

Plant Biol 60: 379-406.

Bolte S, Talbot C, Boutte Y, Catrice O, Read ND, Satiat-Jeunemaitre B. 2004. FM-

dyes as experimental probes for dissecting vesicle trafficking in living plant cells. *J*

Microsc 214: 159-173.

Bourne HR, Sanders DA, McCormick F. 1991. The GTPase superfamily: conserved

structure and molecular mechanism. *Nature* 349: 117-127.

Campos ML, Yoshida Y, Major IT, de Oliveira Ferreira D, Weraduwage SM,

Froehlich JE, Johnson BF, Kramer DM, Jander G, Sharkey TD and Howe GA.

(2016). Rewiring of jasmonate and phytochrome B signalling uncouples plant growth-

defense tradeoffs. *Nature Comms*, 7, 12570.

Chen H, Zou Y, Shang Y, Lin H, Wang Y, Cai R, Tang X, Zhou JM. 2008. Firefly

luciferase complementation imaging assay for protein-protein interactions in plants.

Plant Physiol 146: 368-376.

Chiang YH, Coaker G. 2015.Effector Triggered Immunity: NLR Immune Perception

and Downstream Defense Responses. *The Arabidopsis Book*. American Society of

Plant Biologists, 13: e0183.

Clough SJ, Bent AF. 1998. Floral dip: a simplified method for Agrobacterium-

mediated transformation of Arabidopsis thaliana. *Plant J* 16: 735-743.

Cook DE, Mesarich CH, Thomma BPHJ. 2015. Understanding Plant Immunity as a
Surveillance System to Detect Invasion. *Annual Review of Phytopathology*, Vol 53:
541-563.
754

755 Couto D, Zipfel C. 2016. Regulation of pattern recognition receptor signalling in
756 plants. *Nat Rev Immunol* 16: 537-552.
757

758 de Wit M, Spoel SH, Sanchez-Perez GF, Gommers CMM, Pieterse CMJ, Voesenek
759 LACJ, Pierik R. 2013. Perception of low red:far-red ratio compromises both salicylic
760 acid- and jasmonic acid-dependent pathogen defences in Arabidopsis. *Plant J* 75:
761 90-103.
762

763 Delaney TP, Uknes S, Vernooij B, Friedrich L, Weymann K, Negrotto D, Gaffney T,
764 Gut-Rella M, Kessmann H, Ward E, et al. 1994. A central role of salicylic Acid in
765 plant disease resistance. *Science* 266: 1247-1250.
766

767 Feng Z, Zhang B, Ding W, Liu X, Yang DL, Wei P, Cao F, Zhu S, Zhang F, Mao Y, et
768 al. 2013. Efficient genome editing in plants using a CRISPR/Cas system. *Cell Res*
769 23: 1229-1232.
770

771 Fischer-Parton S, Parton RM, Hickey PC, Dijksterhuis J, Atkinson HA, Read ND.
772 2000. Confocal microscopy of FM4-64 as a tool for analysing endocytosis and
773 vesicle trafficking in living fungal hyphae. *J Microsc* 198: 246-259.
774

775 Huot B, Yao J, Montgomery BL, He SY. 2014. Growth-defense tradeoffs in plants: a
776 balancing act to optimize fitness. *Mol Plant* 7: 1267-1287.
777

778 Kadota Y, Macho AP, Zipfel C. 2016. Immunoprecipitation of Plasma Membrane
779 Receptor-Like Kinases for Identification of Phosphorylation Sites and Associated
780 Proteins. *Methods Mol Biol* 1363: 133-144.
781

782 Khan M, Subramaniam R, Desveaux D. 2016. Of guards, decoys, baits and traps:
783 pathogen perception in plants by type III effector sensors. *Curr Opin Microbiol* 29: 49-
784 55.
785

Liu C, Wang T, Zhang W, Li X. 2008. Computational identification and analysis of
immune-associated nucleotide gene family in Arabidopsis thaliana. *J Plant Physiol*
165: 777-787.

Macho AP. 2016. Subversion of plant cellular functions by bacterial type-III effectors:
beyond suppression of immunity. *New Phytol* 210: 51-57.

Macho AP, Boutrot F, Rathjen JP, Zipfel C. 2012. Aspartate oxidase plays an
important role in Arabidopsis stomatal immunity. *Plant Physiol* 159: 1845-1856.

Macho AP, Zipfel C. 2014. Plant PRRs and the activation of innate immune signaling.
Mol Cell 54: 263-272.

Macho AP, Zipfel C. 2015. Targeting of plant pattern recognition receptor-triggered
immunity by bacterial type-III secretion system effectors. *Curr Opin Microbiol* 23: 14-
22.

Mao Y, Zhang H, Xu N, Zhang B, Gou F, Zhu JK. 2013. Application of the CRISPR-
Cas system for efficient genome engineering in plants. *Mol Plant* 6: 2008-2011.

Nakagawa T, Kurose T, Hino T, Tanaka K, Kawamukai M, Niwa Y, Toyooka K,
Matsuoka K, Jinbo T, Kimura T. 2007. Development of series of gateway binary
vectors, pGWBs, for realizing efficient construction of fusion genes for plant
transformation. *J Biosci Bioeng* 104: 34-41.

Nitta T, Nasreen M, Seike T, Goji A, Ohigashi I, Miyazaki T, Ohta T, Kanno M,
Takahama Y. 2006. IAN family critically regulates survival and development of T
lymphocytes. *PLoS Biol* 4: e103.

Nitta T, Takahama Y. 2007. The lymphocyte guard-IANs: regulation of lymphocyte
survival by IAN/GIMAP family proteins. *Trends Immunol* 28: 58-65.

Reuber TL, Ausubel FM. 1996. Isolation of Arabidopsis genes that differentiate
between resistance responses mediated by the RPS2 and RPM1 disease resistance
genes. *Plant Cell* 8: 241-249.

Sang Y, Macho AP. 2017. Analysis of PAMP-Triggered ROS Burst in Plant Immunity.

Methods Mol Biol 1578: 143-153.

Sang Y, Wang Y, Ni H, Cazale AC, She YM, Peeters N, Macho AP. 2016. The

Ralstonia solanacearum type III effector RipAY targets plant redox regulators to

suppress immune responses. *Mol Plant Pathol.*

Scheres B, van der Putten WH. 2017. The plant perceptron connects environment to

development. *Nature* 543: 337-345.

Stael S, Kmiecik P, Willems P, Van Der Kelen K, Coll NS, Teige M, Van Breusegem

F. 2015. Plant innate immunity--sunny side up? *Trends Plant Sci* 20: 3-11.

Takai Y, Sasaki T, Matozaki T. 2001. Small GTP-binding proteins. *Physiol Rev* 81:

153-208.

Tsuda K, Glazebrook J, Katagiri F. 2008. The interplay between MAMP and SA

signaling. *Plant Signal Behav* 3: 359-361.

Vernoud V, Horton AC, Yang Z, Nielsen E. 2003. Analysis of the small GTPase gene

superfamily of Arabidopsis. *Plant Physiol* 131: 1191-1208.

Vlot AC, Dempsey DA, Klessig DF. 2009. Salicylic Acid, a multifaceted hormone to

combat disease. *Annu Rev Phytopathol* 47: 177-206.

Wang ZY, Seto H, Fujioka S, Yoshida S, Chory J. 2001. BRI1 is a critical component

of a plasma-membrane receptor for plant steroids. *Nature* 410: 380-383.

Weiss Y, Foret S, Hayward DC, Ainsworth T, King R, Ball EE, Miller DJ. 2013. The

acute transcriptional response of the coral Acropora millepora to immune challenge:

expression of GiMAP/IAN genes links the innate immune responses of corals with

those of mammals and plants. *BMC Genomics* 14: 400.

Wildermuth MC, Dewdney J, Wu G, Ausubel FM. 2001. Isochorismate synthase is

required to synthesize salicylic acid for plant defence. *Nature* 414: 562-565.

Win J, Chaparro-Garcia A, Belhaj K, Saunders DG, Yoshida K, Dong S, Schornack
S, Zipfel C, Robatzek S, Hogenhout SA, et al. 2012. Effector biology of plant-
associated organisms: concepts and perspectives. *Cold Spring Harb Symp Quant*
Biol 77: 235-247.
862
863 Yi SY, Shirasu K, Moon JS, Lee SG, Kwon SY. 2014. The activated SA and JA
864 signaling pathways have an influence on flg22-triggered oxidative burst and callose
865 deposition. *PLoS One* 9: e88951.
866
867 Zipfel C. 2014. Plant pattern-recognition receptors. *Trends Immunol* 35: 345-351.
868
869

A

WT   *iap1-1*   *iap1-2*

B

Log₁₀(CFU/mg FW)

*

WT   *iap1-1*   *iap1-2*

C

Log₁₀(CFU/cm²)

ns

WT   *iap1-1*   *iap1-2*