Limit distribution of the quartet balance index for Aldous's $\beta \geq 0$–model

Krzysztof Bartoszek

November 21, 2018

Abstract

This paper builds up on T. Martínez–Coronado, A. Mir, F. Rosselló and G. Valiente's work "A balance index for phylogenetic trees based on quartets", introducing a new balance index for trees. We show here that this balance index, in the case of Aldous's $\beta \geq 0$–model, convergences weakly to a distribution that can be characterized as the fixed point of a contraction operator on a class of distributions.

Keywords : Balance index; contraction method; phylogenetic tree; tree shape; weak convergence

1 Introduction

Phylogenetic trees are key to evolutionary biology. However, they are not easy to summarize or compare as it might not be obvious how to tackle their topologies, understood as the internal branching structure of the trees. Therefore, many summary indices have been proposed in order to "project" a tree into $\mathbb{R}$. Such indices have as their aim to quantify some property of the tree and one of the most studied properties is the symmetry of the tree. Tree symmetry is commonly captured by a balance index. Multiple balance indices have been proposed, Sackin's (Sackin, 1972), Colless' (Colless, 1982) or the total cophenetic index (Mir et al., 2013). A compact introduction to phylogenetics, containing in particular a list of tree asymmetry measures (p. 562–564), can be found in Felsenstein (2004)'s book. This work accompanies

a newly proposed balance index—the quartet index (QI, Martínez-Coronado et al., 2018b).

One of the reasons for introducing summary indices for trees is to use them for significance testing—does a tree come from a given probabilistic model. Obtaining the distribution (for a given n–number of contemporary species, i.e. leaves of the tree, or in the limit $n \to \infty$) of indices is usually difficult and often is done only for the "simplest" Yule (pure–birth Yule, 1924) tree case and sometimes uniform model (see e.g. Aldous, 1991; Steel and McKenzie, 2001).

Using the contraction method, central limit theorems were found for various balance indices, like the total cophenetic index (Yule model case Bartoszek, 2018) and jointly for Sackin's and Colless' (in the Yule and uniform model cases Blum et al., 2006). Furthermore, Blum and François (2006) showed that Sackin's index has the same weak limit as the number of comparisons of the quicksort algorithm (Hoare, 1962), both after normalization of course.

Chang and Fuchs (2010) consider the number of occurrences of patterns in a tree, where a pattern is understood as "any subset of the set of phylogenetic trees of fixed size jk. For a tree with n leaves such a pattern will satisfy the recursion

$$X_{n,k} \overset{\mathcal{D}}{=} X_{L_n,k} + X^*_{n-L_n,k}$$

where $X_{n,k}$, $X^*_{n,k}$ and L_n are independent, $X_{n,k} \overset{\mathcal{D}}{=} X^*_{n,k}$ and L_n is the size of the left subtree branching from the root. For the Yule and uniform models they derived central limit theorems (normal limit distribution) with Berry–Esseen bounds and Poisson approximations in the total variation distance.

Even though the pure–birth model seems to be very widespread in the phylogenetics community, more complex models need to be studied, especially in the context of tree balance. From Roch and Snir (2013)'s Lemma 4 it can be deduced that Yule trees have to be rather balanced— as the maximum quartet weight (maximum of number of randomly placed marks along branches over induced subtrees on four leaves) is asymptotically proportional to the expectation of the tree's height.

In this work here, using the contraction method, we show convergence in law of the (scaled and centred) quartet index and derive a representation (as a fixed point of a particular contraction operator) of the weak–limit. Remarkably, this is possible not only for the Yule tree case but for Aldous's

more general β–model (in the $\beta \geq 0$ regime).

The paper is organized as follows. In Section 2 we introduce Aldous's β–model and the quartet index. In Section 3 we prove our main result— Thm. 3.1 via the contraction method. When studying the limit behaviour of recursive–type indices for pure–birth binary trees one has that for each internal node the leaves inside its clade are uniformly split into to sub–clades as the node splits. However, in Aldous's β–model this is not the case, the split is according to a BetaBinomial distribution, and a much finer analysis is required to show weak–convergence, with n, of the recursive–type index to the fixed point of the appropriate contraction. Theorem 3.1 is not specific for the quartet index but covers a more general class of models, where each internal node split divides its leaf descendants according to a BetaBinomial distribution (with $\beta \geq 0$). In Section 4 we apply Thm. 3.1 to the quartet index and characterize its weak limit. Then, in Section 5 we illustrate the results with simulations. Finally, in the Appendix we provide R code used to simulate from this weak limit.

2 Preliminaries

2.1 Aldous's β–model for phylogenetic trees

Birth–death models are popular choices for modelling the evolution of phylogenetic trees. However, Aldous (1996, 2001) proposed a different class of models—the so–called β–model for binary phylogenetic trees.

The main idea behind this model is to consider a (suitable) family $\{q_n\}_{n=2}^{\infty}$ of symmetric, $q_n(i) = q_n(n - i)$, probability distributions on the natural numbers. In particular $q_n : \{1, \ldots, n - 1\} \to [0, 1]$. The tree grows in a natural way. The root node of a n–leaf tree defines a partition of the n nodes into two sets of sizes i and $n - i$ ($i \in \{1, \ldots, n-1\}$). We randomly choose the number of leaves of the left subtree, $L_n = i$, according to the distribution q_n and this induces the number of leaves, $n - L_n$, in the right subtree. We then repeat recursively in the left and right subtrees, i.e. splitting according to the distributions q_{L_n} and q_{n-L_n} respectively. Notice that due to q_n's symmetry the terms left and right do not have any particular meaning attached.

Aldous (1996) proposed to consider a one–parameter, $-2 \leq \beta \leq \infty$, family of probability distributions,

$$q_n(i) = \frac{1}{a_n(\beta)} \frac{\Gamma(\beta+i+1)\Gamma(\beta+n-i+1)}{\Gamma(i+1)\Gamma(n-i+1)}, \quad 1 \le i \le n-1, \quad (1)$$

where $a_n(\beta)$ is the normalizing constant and $\Gamma(\cdot)$ the Gamma function. We may actually recognize this as the BetaBinomial$(n, \beta+1, \beta+1)$ distribution and represent

$$q_n(i) = B(\beta+1, \beta+1)^{-1} \int_0^1 \left(\binom{n}{i} \tau^i (1-\tau)^{n-i} \right) \tau^\beta (1-\tau)^\beta \mathrm{d}\tau, \quad (2)$$

where $B(a, b)$ is the Beta function with parameters a and b. Writing informally, from the form of the probability distribution function, Eq. (2), we can see that if we would condition under the integral on τ, then we obtain a binomially distributed random variable. This is a key observation that is the intuition for the analysis presented here.

Particular values of β correspond to some well known models. The uniform tree model is represented by $\beta = -3/2$, and the pure birth, Yule, model by $\beta = 0$.

Of particular importance to our work is the limiting behaviour of the scaled size of the left (and hence right) subtree, $n^{-1}L_n$. Aldous (1996) characterizes the asymptotics in his Lemma 3,

Lemma 2.1 *(Aldous, 1996, Lemma 3 for $\beta > -1$)*

1. $\beta = \infty$, $n^{-1}L_n \xrightarrow{\mathcal{D}} \frac{1}{2}$;

2. $-1 < \beta < \infty$, $n^{-1}L_n \xrightarrow{\mathcal{D}} \tau_\beta$, where τ_β has the Beta distribution

$$f(x) = \frac{\Gamma(2\beta+2)}{\Gamma^2(\beta+1)} x^\beta (1-x)^\beta, \quad 0 < x < 1. \quad (3)$$

2.2 Quartet index

Martínez-Coronado et al. (2018b) introduced a new type of balance index for discrete (i.e. without branch lengths) phylogenetic trees—the quartet index. This index is based on considering the number of so–called quartets of each

type made up by the leaves of the tree. A (rooted) quartet is the induced subtree from choosing some four leaves. We should make a point here about the used nomenclature. Usually in the phylogenetic literature a quartet is an unrooted tree on four leaves (e.g. Semple and Steel, 2003). However, here we consider rooted trees and following Martínez-Coronado et al. (2018b) by (rooted) quartet we mean a rooted tree on four leaves. We will from now on write quartet for this, dropping the "rooted" clarification.

For a given tree T, let $\mathcal{P}_4(T)$ be the set of quartets of the tree. Then, the quartet index of T is defined as

$$QI(T) = \sum_{\mathcal{P}_4(T)} QI(Q), \tag{4}$$

where $QI(Q)$ assigns a predefined value to a specific quartet. When the tree is a binary one (as here) there are only two possible topologies on four leaves (see Fig. 1). Following Martínez-Coronado et al. (2018b) (their Table 1) we assign the value 0 to K_4 quartets and 1 to B_4 quartets. Therefore, the QI for a binary tree (QIB) will be

$$QIB(T) = \text{number of } B_4 \text{ quartets of } T. \tag{5}$$

Figure 1: The two possible rooted quartets for a binary tree. Left: K_4 the four leaf rooted caterpillar tree (also known as a comb or pectinate tree), right: B_4 the fully balanced tree on four leaves (also known as a fork, see e.g. Chor and Snir, 2007, for some nomenclature).

Importantly for us Martínez-Coronado et al. (2018b) show in their Lemma 4 that for $n > 4$, the quartet index has a recursive representation as

$$QIB(T_n) = QIB(T_{L_n}) + QIB(T_{n-L_n}) + \binom{L_n}{2}\binom{n-L_n}{2}, \tag{6}$$

where T_n is the tree on n leaves.

Martínez-Coronado et al. (2018b) considered various models of tree growth, Aldous's β–model, Ford's α–model (Ford, 2005, but see also Martínez-Coronado et al. (2018a)) and Chen–Ford–Winkel's α–γ–model (Chen et al., 2009). In this work we will focus on the Aldous's $\beta \geq 0$–model of tree growth and characterize the limit distribution, as the number of leaves, n, grows to infinity, of the QI. We will take advantage of the recursive representation of Eq. (6) that allows for the usage of the powerful contraction method.

We require knowledge of the mean and variance of the QI and Martínez-Coronado et al. (2018b) show for Aldous's β–model that these are (their Corollaries 4 and 7)

$$
\begin{array}{rcl}
\mathrm{E}\left[QIB(T_n)\right] & = & \frac{3\beta+6}{7\beta+18}\binom{n}{4} \\[2mm]
\mathrm{Var}\left[QIB(T_n)\right] & = & \frac{(\beta+2)\left(2\beta^2+9\beta+12\right)}{2(7\beta+18)^2(127\beta^3+1383\beta^2+4958\beta+5880)}n^8 + O(n^7).
\end{array}
\tag{7}
$$

3 Contraction method approach

Consider the space D of distribution functions with finite second moment and first moment equalling 0. On D we define the Wasserstein metric

$$
d(F,G) = \inf \|X - Y\|_2
$$

where $\|\cdot\|_2$ denotes the L_2 norm and the infimum is over all $X \sim F, Y \sim G$. Notice that convergence in d induces convergence in distribution.

For $r \in \mathbb{N}_+$ define the transformation $S : D \to D$ as

$$
S(F) = \mathcal{L}\left(\tau^r Y' + (1-\tau)^r Y'' + C(\tau)\right),
\tag{8}
$$

where Y', Y'', τ are independent, $Y', Y'' \sim F$, $\tau \in [0,1]$ whose distribution is not a Dirac δ at 0 nor 1, satisfying, for all n,

$$
2\sum_{i=1}^{n} p_{n,i}\left(\frac{i}{n}\right)^{2r} < 1,
\tag{9}
$$

where $p_{n,i} = P((i-1)/n < \tau \leq i/n)$ and the function $C(\cdot)$ is of the form

$$
C(\tau) = \sum_{r_1+r_2 \leq r} C_{r_1,r_2} \tau^{r_1}(1-\tau)^{r_2}
\tag{10}
$$

for some constants C_{r_1, r_2} and furthermore satisfies $E[C(\tau)] = 0$. By Rösler (1992)'s Thms. 3 and 4 S is well defined, has a unique fixed point and for any $F \in D$ the sequence $S^n(F)$ converges exponentially fast in the d metric to S's fixed point. Using the exact arguments to show Rösler (1991)'s Thm. 2.1 one can show that the map S is a contraction. Only the Lipschitz constant of convergence will differ being $\sqrt{C_\tau}$, where $C_\tau = \max\{E[\tau^{2r}], E[(1 - \tau)^{2r}]\}$ in our case. Notice that as $\tau \in [0, 1]$ and is non–degenerate at the edges, then $C_\tau < 1$ and we have a contraction.

We now state the main result of our work. We show weak convergence, with a characterization of the limit for a class of recursively defined models.

Theorem 3.1 *(cf. Rösler, 1991, Thm. 3.1) For $n \geq 2$, $\beta > 0$ let $L_n \in \{1, \ldots, n - 1\}$ be such that $(L_n - 1)$ is $\mathrm{BetaBinomial}(n - 2, \beta + 1, \beta + 1)$ distributed and $\tau \sim \mathrm{Beta}(\beta + 1, \beta + 1) =: F_\tau$ distributed. Starting from the Dirac δ at 0, i.e. $Y_1 = 0$ and convention $\mathrm{BetaBinomial}(0, \beta + 1, \beta + 1) = \delta_0$, for $r \in \mathbb{N}_+$ such that the condition of Eq. (9) is met with the previous choice of F_τ, define recursively the sequence of random variables,*

$$Y_n = \left(\frac{L_n}{n}\right)^r Y_{L_n} + \left(1 - \frac{L_n}{n}\right)^r Y_{n - L_n} + C_n(L_n),$$

where the function $C_n(\cdot)$ is of the form

$$C_n(i) = n^{-r} \left(\sum_{r_1 + r_2 + r_3 \leq r} C_{r_1, r_2, r_3} i^{r_1}(n - i)^{r_2} n^{r_3} + h_n(i)\right), \qquad (11)$$

where $E[C_n(L_n)] = 0$ and $\sup_i n^{-r} h_n(i) \to 0$. If $E[Y_n^2]$ is uniformly bounded then, the random variable Y_n converges in the Wasserstein d–metric to the random variable Y_∞ whose distribution satisfies the unique fixed point of S (Eq. 8).

Notice that as $Y_1 = 0$ and by the definition of the recursion we will have $E[Y_n] = 0$ for all n.

The Yule tree case will be the limit of $\beta = 0$ and this case the proof of the result will be more straightforward (as commented on in the proof of Thm. 3.1).

Notice that $L_n/n \xrightarrow{D} \tau$. It would be tempting to suspect that Thm. 3.1 should be the conclusion of a general result related to the contraction method (as presented in Eq. (8.12), p. 351 Drmota, 2009). However, to the best of

my knowledge, general results assume L_2 convergence of L_n/n (e.g. Thm. 8.6, p. 354 Drmota, 2009), while in our phylogenetic balance index case we will have only convergence in distribution. In such a case it seems that convergence has to be proved case by case (e.g. examples in Rachev and Rüschendorf, 1995). Here we show the convergence of Thm. 3.1 along the lines of Rösler (1991).

We first derive a lemma that controls the non–homogeneous part of the recursion, i.e. $C_n(\cdot)$ as defined in Eq. (11).

Lemma 3.1 *(cf. Rösler, 1991, Prop. 3.2) Let $C_n : \{1, \ldots, n-1\} \to \mathbb{R}$ be as in Eq. (11). Then*

$$\sup_{x \in [0,1)} \left| C_n(\lfloor (n-1)x \rfloor + 1) - C(x) \right| \leq \sup_i n^{-r} h_n(i) + O(n^{-1}). \qquad (12)$$

PROOF For $1 \leq \lfloor (n-1)x \rfloor + 1 \leq n-1$ and writing $i = \lfloor (n-1)x \rfloor + 1$ we have due to the representation of Eqs. (10) and (11)

$$\left| C_n(\lfloor (n-1)x \rfloor + 1) - C(x) \right| \leq \max\{C_{r_1,r_2}\} \left(\left| \left(\tfrac{i}{n}\right)^r - x^r \right| + \left| \left(1 - \tfrac{i}{n}\right)^r - (1-x)^r \right| + \right.$$
$$\left. \sum_{r_1+r_2 \leq r} \left| \left(\tfrac{i}{n}\right)^{r_1} \left(1 - \tfrac{i}{n}\right)^{r_2} - x^{r_1}(1-x)^{r_2} \right| \right) + \sup_i n^{-r} h_n(i).$$

Bounding the individual components, as by construction x cannot differ from i/n by more than $1/n$

$$\left| \left(\tfrac{i}{n}\right)^r - x^r \right| \leq \sup_{|y-z|<1/n} \left| y^r - z^r \right| \leq \sup_{0 \leq y \leq 1} \left| y^r - (y + n^{-1})^r \right| = O(n^{-1})$$

$$\left| \left(1 - \tfrac{i}{n}\right)^r - (1-x)^r \right| \leq \sup_{|y-z|<1/n} \left| (1-y)^r - (1-z)^r \right| \leq \sup_{0 \leq y \leq 1} \left| y^r - (y + n^{-1})^r \right| = O(n^{-1})$$

$$\left| \left(\tfrac{i}{n}\right)^{r_1} \left(1 - \tfrac{i}{n}\right)^{r_2} - x^{r_1}(1-x)^{r_2} \right| \leq \sup_{\epsilon \in [-n^{-1}, n^{-1}]} \left| y^{r_1}(1-y)^{r_2} - (y+\epsilon)^{r_1}(1-y-\epsilon)^{r_2} \right|$$
$$= \left| y^{r_1}(1-y)^{r_2} - \left(\sum_{k_1=0}^{r_1} \binom{r_1}{k_1} y^{r_1-k_1} \epsilon^{k_1} \right) \left(\sum_{k_2=0}^{r_2} \binom{r_2}{k_2} y^{r_2-k_2} \epsilon^{k_2} \right) \right|$$
$$= \left| \left(\sum_{k_1=1}^{r_1} \binom{r_1}{k_1} y^{r_1-k_1} \epsilon^{k_1} \right) + \left(\sum_{k_2=1}^{r_2} \binom{r_2}{k_2} y^{r_2-k_2} \epsilon^{k_2} \right) + \left(\sum_{k_1=1}^{r_1} \binom{r_1}{k_1} y^{r_1-k_1} \epsilon^{k_1} \right) \left(\sum_{k_2=1}^{r_2} \binom{r_2}{k_2} y^{r_2-k_2} \epsilon^{k_2} \right) \right|$$
$$= O(n^{-1})$$

□

Lemma 3.2 *(cf. Rösler, 1991, Prop. 3.3) Let a_n, b_n, $p_{n,i}$, $n \in \mathbb{N}$ be three sequences such that $0 \leq b_n \to 0$ with n, $0 \leq p_{n,i} \leq 1$,*

$$0 \leq a_{n+1} \leq 2 \sum_{i=1}^{n} p_{n,i} \left(\frac{i}{n} \right)^R \left(\sup_{i \in \{1,\ldots,n\}} a_i \right) + b_n. \tag{13}$$

and

$$0 < 2 \sum_{i=1}^{n} p_{n,i} \left(\frac{i}{n} \right)^R = C < 1.$$

Then $\lim_{n\to\infty} a_n = 0$.

PROOF The proof is exactly the same as Rösler (1991)'s proof of his Proposition 3.3. In the last step we will have with $a := \limsup a_n < \infty$ the sandwiching for all $\epsilon > 0$

$$0 \leq a \leq C(a + \epsilon).$$

□

Having Lemmata 3.1 and 3.2 we turn to showing Thm. 3.1.

PROOF[Proof of Thm. 3.1] Denote the law of Y_n as $\mathcal{L}(Y_n) = G_n$. We take Y_∞ and Y'_∞ independent and distributed as G_∞, the fixed point of S. Then, for $i = 1, \ldots, n - 1$ we choose independent versions of Y_i and Y'_i. We need to show $d^2(G_n, G_\infty) \to 0$. As the metric is the infimum over all pairs of random variables that have marginal distributions G_n and G_∞ the obvious choice is to take Y_n, Y_∞ such that L_n/n will be close to τ for large n. Rösler (1991) was considering the Yule model ($\beta = 0$) and there $\tau \sim \text{Unif}[0,1]$ and L_n is uniform on $\{1, \ldots, n - 1\}$. Hence, $\lfloor (n-1)\tau \rfloor + 1$ will be uniform on $\{1, \ldots, n - 1\}$, remember $P(\tau = 1) = 0$, and $L_n/n \overset{D}{=} (\lfloor (n-1)\tau \rfloor + 1)/n$. However, when $\beta > 0$ the situation complicates. For a given n, $(L_n - 1)$ is BetaBinomial$(n - 2, \beta + 1, \beta + 1)$ distributed (cf. Eq. 1 and Aldous (1996) Eqs. 1 and 3). Hence, if $\tau \sim \text{Beta}(\beta + 1, \beta + 1)$ and $(L_n - 1) \sim$ BetaBinomial$(n - 2, \beta + 1, \beta + 1)$ we do not have $L_n/n \overset{D}{=} (\lfloor (n-1)\tau \rfloor + 1)/n$ exactly. We may bound the Wasserstein metric by any coupling that retains

the marginal distributions of the two random variables. Therefore, from now on we will be considering a version, where conditional on τ, the random variable $(L_n - 1)$ is Binomial$(n-2, \tau)$ distributed. Let r_n be any sequence such that $r_n/n \to 0$ and $n/r_n^2 \to 0$, e.g. $r_n = n \ln^{-1} n$. Then, by Chebyshev's inequality

$$P\left(|L_n - \mathrm{E}[L_n|\tau]| \geq r_n \Big| \tau\right) \leq \frac{n\tau(1-\tau)}{r_n^2} \leq \frac{n}{4r_n^2} \to 0.$$

We now want to show $d^2(G_n, G_\infty) \to 0$ and we will exploit the above coupling in the bound

$$d^2(G_n, G_\infty) \leq \mathrm{E}\left[\left(\left(\left(\tfrac{L_n}{n}\right)^r Y_{L_n} - \tau^r Y_\infty\right) + \left(\left(\tfrac{n-L_n}{n}\right)^r Y_{n-L_n} - (1-\tau)^r Y'_\infty\right) + (C_n(L_n) - C(\tau))\right)^2\right]$$

$$= \mathrm{E}\left[\left(\left(\tfrac{L_n}{n}\right)^r Y_{L_n} - \tau^r Y_\infty\right)^2\right] + \mathrm{E}\left[\left(\left(\tfrac{n-L_n}{n}\right)^r Y_{n-L_n} - (1-\tau)^r Y'_\infty\right)^2\right] + \mathrm{E}\left[(C_n(L_n) - C(\tau))^2\right],$$

where $Y_\infty, Y'_\infty \sim G_\infty$ are independent. Remember that $\mathrm{E}[Y_i] = \mathrm{E}[Y_\infty] = 0$ so that the expectation of the cross products disappears.

Our main step is to have a bound where the L_n/n term is replaced by some transformation of τ. Let $\tilde{r}_n$ be a (appropriate) random integer in $\{\pm 1, \ldots, \pm \lceil r_n \rceil\}$ and we may write (with the chosen coupling of L_n and τ),

$$\mathrm{E}\left[\left(\left(\tfrac{L_n}{n}\right)^r Y_{L_n} - \tau^r Y_\infty\right)^2\right] = \mathrm{E}\left[\mathrm{E}\left[\left(\left(\tfrac{L_n}{n}\right)^r Y_{L_n} - \tau^r Y_\infty\right)^2 \Big| \tau\right]\right]$$

$$= \mathrm{E}\left[\mathrm{E}\left[\left(\left(\tfrac{\lfloor (n-1)\tau \rfloor + 1 + \tilde{r}_n}{n}\right)^r Y_{L_n} - \tau^r Y_\infty\right)^2 \Big| |L_n - \mathrm{E}[L_n]| \leq r_n, \tau\right] P(|L_n - \mathrm{E}[L_n]| \leq r_n \Big| \tau)\right]$$

$$+ \mathrm{E}\left[\mathrm{E}\left[\left(\left(\tfrac{L_n}{n}\right)^r Y_{L_n} - \tau^r Y_\infty\right)^2 \Big| |L_n - \mathrm{E}[L_n]| \geq r_n, \tau\right] P(|L_n - \mathrm{E}[L_n]| \geq r_n \Big| \tau)\right]$$

$$\leq \mathrm{E}\left[\left(\left(\tfrac{\lfloor (n-1)\tau \rfloor + 1 + \tilde{r}_n}{n}\right)^r Y_{L_n} - \tau^r Y_\infty\right)^2\right] + \tfrac{n}{4r_n^2} \mathrm{E}\left[\left(\left(\tfrac{L_n}{n}\right)^r Y_{L_n} - \tau^r Y_\infty\right)^2\right]$$

$$= \mathrm{E}\left[\left(\left(\left(\tfrac{\lfloor (n-1)\tau \rfloor + 1}{n}\right)^r + r\tfrac{\tilde{r}_n}{n}\left(\tfrac{\lfloor (n-1)\tau \rfloor + 1 + \xi_{\tilde{r}_n}}{n}\right)^{r-1}\right) Y_{L_n} - \tau^r Y_\infty\right)^2\right]$$

$$+ \tfrac{n}{4r_n^2} \mathrm{E}\left[\left(\left(\tfrac{L_n}{n}\right)^r Y_{L_n} - \tau^r Y_\infty\right)^2\right]$$

$$
= \mathrm{E}\left[\left(\left(\tfrac{\lfloor(n-1)\tau\rfloor+1}{n}\right)^r Y_{L_n} - \tau^r Y_\infty\right)^2\right] + r^2 n^{-2}\, \mathrm{E}\left[\tilde{r}_n^2\left(\tfrac{\lfloor(n-1)\tau\rfloor+1+\xi_{\tilde{r}_n}}{n}\right)^{2(r-1)} Y_{L_n}^2\right]
$$
$$
+ 2rn^{-1}\, \mathrm{E}\left[\tilde{r}_n\left(\tfrac{\lfloor(n-1)\tau\rfloor+1+\xi_{\tilde{r}_n}}{n}\right)^{r-1} Y_{L_n} \cdot \left(\left(\tfrac{\lfloor(n-1)\tau\rfloor+1}{n}\right)^r Y_{L_n} - \tau^r Y_\infty\right)\right]
$$
$$
+ \tfrac{n}{4r_n^2}\, \mathrm{E}\left[\left(\left(\tfrac{L_n}{n}\right)^r Y_{L_n} - \tau^r Y_\infty\right)^2\right],
$$

where $\xi_{\tilde{r}_n} \in (0, \tilde{r}_n)$ is (a random variable) such that the mean value theorem holds (for the function $(\cdot)^r$). As Y_n, Y_∞ have uniformly bounded second moments, $0 \le \xi_{\tilde{r}_n} \le \tilde{r}_n \le r_n \le n$, by assumption $r_n/n \to 0$ and $n/r_n^2 \to 0$, we have that

$$
\left(r\tfrac{r_n}{n}\right)^2 \mathrm{E}\left[\left(\tfrac{\lfloor(n-1)\tau\rfloor+1+\xi_{\tilde{r}_n}}{n}\right)^{2(r-1)} Y_{L_n}^2\right] + \tfrac{n}{4r_n^2}\, \mathrm{E}\left[\left(\left(\tfrac{L_n}{n}\right)^r Y_{L_n} - \tau^r Y_\infty\right)^2\right]
$$
$$
+ 2r\tfrac{r_n}{n}\, \mathrm{E}\left[\left(\tfrac{\lfloor(n-1)\tau\rfloor+1+\xi_{\tilde{r}_n}}{n}\right)^{r-1} Y_{L_n} \cdot \left(\left(\tfrac{\lfloor(n-1)\tau\rfloor+1}{n}\right)^r Y_{L_n} - \tau^r Y_\infty\right)\right] \to 0
$$

and hence for some sequence $u_n \to 0$ we have,

$$
\mathrm{E}\left[\left(\left(\tfrac{L_n}{n}\right)^r Y_{L_n} - \tau^r Y_\infty\right)^2\right] \le \mathrm{E}\left[\left(\left(\tfrac{\lfloor(n-1)\tau\rfloor+1}{n}\right)^r Y_{L_n} - \tau^r Y_\infty\right)^2\right] + u_n.
$$

Remembering the assumption $\sup_i n^{-r} h_n(i) \to 0$, the other component can be treated in the same way as $\mathrm{E}\left[\left(\left(\tfrac{L_n}{n}\right)^r Y_{L_n} - \tau^r Y_\infty\right)^2\right]$ with conditioning on τ and then controlling by r_n and Chebyshev's inequality L_n's deviation from its expected value. We therefore have for some sequence $v_n \to 0$

$$
d^2(G_n, G_\infty) \le \mathrm{E}\left[\left(\left(\tfrac{\lfloor(n-1)\tau\rfloor+1}{n}\right)^r Y_{L_n} - \tau^r Y_\infty\right)^2\right] + \mathrm{E}\left[\left(C_n(\lfloor(n-1)\tau\rfloor+1) - C(\tau)\right)^2\right]
$$
$$
+ \mathrm{E}\left[\left(\left(\tfrac{n-\lfloor(n-1)\tau\rfloor-1}{n}\right)^r Y_{n-L_n} - (1-\tau)^r Y_\infty'\right)^2\right] + v_n.
$$

Consider the first term of the right–hand side of the inequality and denote $d_{n-1}^2 := \sup_{i \in \{1,\dots,n-1\}} d^2(G_i, G_\infty)$

$$\mathrm{E}\left[\left(\left(\tfrac{\lfloor(n-1)\tau\rfloor+1}{n}\right)^{r}Y_{L_n}-\tau^{r}Y_{\infty}\right)^{2}\right]=\mathrm{E}\left[\sum_{i=1}^{n-1}\mathbb{1}_{(i-1)/(n-1)<\tau\le i/(n-1)}\left(\left(\tfrac{i}{n}\right)^{r}Y_{L_n}-\tau^{r}Y_{\infty}\right)^{2}\right]$$

$$\le\sum_{i=1}^{n-1}p_{n-1,i}\left(\tfrac{i}{n}\right)^{2r}\mathrm{E}\left[(Y_{L_n}-Y_{\infty})^{2}\right]=\sum_{i=1}^{n-1}p_{n-1,i}\left(\tfrac{i}{n}\right)^{2r}d_{n-1}^{2},$$

where $p_{n,i}=P((i-1)/(n-1)<\tau\le i/(n-1))$. Invoking Lemmata 3.1, 3.2 and using the assumption of Eq. (9) with $R=2r$ we have

$$d^{2}(G_n,G_{\infty})\le 2\sum_{i=1}^{n-1}p_{n-1,i}\left(\tfrac{i}{n}\right)^{2r}d_{n-1}^{2}+\left(n^{-r}\sup_{i}h_n(i)\right)^{2}+v_n+O(n^{-2})\to 0.$$

$\square$

4 Limit distribution of the quartet index for Aldous's $\beta\ge 0$–model trees

We show here that the QIB of Aldous's $\beta\ge 0$–model trees satisfies the conditions of Thm. 3.1 with $r=4$ and hence the QIB has a well characterized limit distribution. We define a centred and scaled version of the QIB for Aldous's $\beta\ge 0$–model tree on $n\ge 4$ leaves

$$Y_n^Q=n^{-4}\left(QIB(T_n)-\frac{3\beta+6}{7\beta+18}\binom{n}{4}\right). \tag{14}$$

We now specialize Thm. 3.1 to the QIB case and assume $Y_1=Y_2=Y_3=0$ for completeness

Theorem 4.1 *The sequence of random variables Y_n^Q random variable for trees generated by Aldous's β–model with $\beta\ge 0$ converges with $n\to\infty$ in the Wasserstein d–metric (and hence in distribution) to a random variable $Y_Q\sim\mathcal{Q}\equiv G_{\infty}$ satisfying the following equality in distribution*

$$Y_Q\overset{\mathcal{D}}{=}\tau^4 Y_Q'+(1-\tau)^4 Y_Q''+\frac{3\beta+6}{24(7\beta+18)}\left(\tau^4+(1-\tau)^4\right)-\frac{3\beta+6}{24(7\beta+18)}+\frac{1}{4}\tau^2(1-\tau)^2, \tag{15}$$

where $\tau\sim F_{\tau}$ is distributed as the Beta distribution of Eq. (3), $Y_Q,Y_Q',Y_Q''\sim \mathcal{Q}$ and Y_Q',Y_Q'',τ are all independent.

PROOF Denote by $P_3(x, y)$ a polynomial of degree at most three in terms of the variables x, y. From the recursive representation of Eq. (6) for $n > 4$

$$Y_n^Q = n^{-4}\left(QIB(T_{L_n}) - \tfrac{3\beta+6}{(7\beta+18)}\binom{L_n}{4} + QIB(T_{n-L_n}) - \tfrac{3\beta+6}{(7\beta+18)}\binom{n-L_n}{4} + \binom{L_n}{2}\binom{n-L_n}{2}\right.$$
$$\left. + \tfrac{3\beta+6}{(7\beta+18)}\binom{L_n}{4} + \tfrac{3\beta+6}{(7\beta+18)}\binom{n-L_n}{4} - \tfrac{3\beta+6}{(7\beta+18)}\binom{n}{4}\right)$$
$$= \left(\tfrac{L_n}{n}\right)^4 Y_{L_n}^Q + \left(1 - \tfrac{L_n}{n}\right)^4 Y_{n-L_n}^Q + \tfrac{1}{4}\left(\tfrac{L_n}{n}\right)^2\left(1 - \tfrac{L_n}{n}\right)^2 + \tfrac{3\beta+6}{24(7\beta+18)}\left(\tfrac{L_n}{n}\right)^4$$
$$+ \tfrac{3\beta+6}{24(7\beta+18)}\left(1 - \tfrac{L_n}{n}\right)^4 - \tfrac{3\beta+6}{24(7\beta+18)} + n^{-4}P_3(n, L_n).$$

We therefore have $r = 4$ and

$$C_n(i) = \tfrac{1}{4}\left(\tfrac{i}{n}\right)^2\left(1 - \tfrac{i}{n}\right)^2 + \tfrac{3\beta+6}{24(7\beta+18)}\left(\left(\tfrac{i}{n}\right)^4 + \left(1 - \tfrac{i}{n}\right)^4\right) - \tfrac{3\beta+6}{24(7\beta+18)}$$
$$+ n^{-4}P_3(n, i).$$

By the scaling and centring we know that $EY_n^Q = 0$ and $E\left(Y_n^Q\right)^2$ is uniformly bounded by Eq. (7). For the Beta law of τ we need to examine for all i

$$p_{n,i} := P\left(\frac{i-1}{n} \leq \tau < \frac{i}{n}\right) = \frac{\Gamma(2\beta+2)}{\Gamma^2(\beta+1)}\int\limits_{(i-1)/n}^{i/n} x^\beta(1-x)^\beta \mathrm{d}x.$$

We consider two cases

1. $\beta > 0$, we have to check if the condition of Eq. (9) is satisfied. Let

$$B_x(\beta+1, \beta+1) = \int\limits_0^x u^\beta(1-u)^\beta \mathrm{d}u$$

be the incomplete Beta function. Then,

$$\begin{aligned}
p_{n,i} &= \tfrac{\Gamma(2\beta+2)}{\Gamma^2(\beta+1)}\left(B_{i/n}(\beta+1, \beta+1) - B_{(i-1)/n}(\beta+1, \beta+1)\right) \\
&= n^{-1}\tfrac{\Gamma(2\beta+2)}{\Gamma^2(\beta+1)}B'_\xi(\beta+1, \beta+1)
\end{aligned}$$

by the mean value theorem for some $\xi \in ((i-1)/n, i/n)$. Obviously

$$B'_\xi(\beta+1, \beta+1) = \xi^\beta (1-\xi)^\beta \le \left(\frac{i}{n}\right)^\beta \left(1 - \frac{i-1}{n}\right)^\beta$$

and now

$$\sum_{i=1}^n p_{n,i} \left(\frac{i}{n}\right)^R \le \frac{\Gamma(2\beta+2)}{\Gamma^2(\beta+1)} n^{-1} \sum_{i=1}^n \left(\frac{i}{n}\right)^R \left(\frac{i}{n}\right)^\beta \left(1 - \frac{i-1}{n}\right)^\beta \to \frac{\Gamma(2\beta+2)}{\Gamma^2(\beta+1)} \int_0^1 u^{\beta+R}(1-u)^\beta \mathrm{d}u$$

$$= \frac{\Gamma(2\beta+2)}{\Gamma^2(\beta+1)} \frac{\Gamma(\beta+R+1)\Gamma(\beta+1)}{\Gamma(2\beta+R+2)} = \frac{\Gamma(2\beta+2)}{\Gamma(\beta+1)} \frac{\Gamma(\beta+R+1)}{\Gamma(2\beta+R+2)}.$$

$$(16)$$

Take $1 < R_1 < R_2$ and consider the ratio

$$A = \frac{\Gamma(2\beta+2)}{\Gamma(\beta+1)} \frac{\Gamma(\beta+R_2+1)}{\Gamma(2\beta+R_2+2)} \left(\frac{\Gamma(2\beta+2)}{\Gamma(\beta+1)} \frac{\Gamma(\beta+R_1+1)}{\Gamma(2\beta+R_1+2)}\right)^{-1} = \frac{\Gamma(\beta+R_2+1)}{\Gamma(2\beta+R_2+2)} \frac{\Gamma(\beta)}{\Gamma(\beta)} \frac{\Gamma(2\beta+R_1+1)}{\Gamma(\beta+R_1+1)}$$

$$= \frac{B(\beta+1+R_2,\beta)}{B(\beta+1+R_1,\beta)}.$$

The ratio $A < 1$ as the Beta function is decreasing in its arguments—hence the derived upper bound in Eq. (16) is decreasing in R. For $R = 1$ the bound equals

$$\frac{\Gamma(2\beta+2)}{\Gamma(\beta+1)} \frac{\Gamma(\beta+2)}{\Gamma(2\beta+3)} = \frac{\Gamma(2\beta+2)}{\Gamma(\beta+1)} \frac{(\beta+1)\Gamma(\beta+1)}{(2\beta+2)\Gamma(2\beta+2)} = \frac{(\beta+1)}{2(\beta+1)} = \frac{1}{2}$$

and hence for all $R > 1$ and all $\beta > 0$

$$\sum_{i=1}^n p_{n,i} \left(\frac{i}{n}\right)^R < \frac{1}{2}.$$

As in our case we have $r \ge 1$, then for $R = 2r \ge 2$ the assumptions of Lemma 3.2 are satisfied and the statement of the theorem follows through.

2. $\beta = 0$, then directly $p_{n,i} = n^{-1}$, Eq. (9) and assumptions of Lemma 3.2 are immediately satisfied and the statement of the theorem follows through. This is the Yule model case, in which the proof of the counterpart of Thm. 3.1 is much more straightforward, as mentioned before.

□

Remark 4.1 *When $\beta < 0$ the process L_n/n seems to have a more involved asymptotic behaviour (cf. Lemma 3 of Aldous, 1996, in $\beta \leq -1$ case). Furthermore, the bounds applied here do not hold for $\beta < 0$. Therefore, this family of tree models (including the important uniform model, $\beta = -3/2$) deserves a separate study with respect to its quartet index.*

5   Comparing with simulations

To verify the results we compared the simulated values from the limiting theoretical distribution of Y_Q with scaled and centred values of Yule tree QI values. The 500–leaf Yule trees were simulated using the `rtreeshape()` function of the apTreeshape R package and Tomás Martínez–Coronado's in–house Python code. Then, for each tree the QI value was calculated by Gabriel Valiente's and Tomás Martínez–Coronado's in–house programs. The raw values $QIB(\text{Yule}_{500})$ were scaled and centred as

$$Y_n^Q = 500^{-4} \left(QIB(\text{Yule}_{500}) - \frac{1}{3}\binom{500}{4} \right).$$

The Y_Q values were simulated using the proposed by Bartoszek (2018) heuristic Algorithm 3 (R code in Appendix). The results of the simulation are presented in Fig. 2.

Acknowledgments

I would like to thank the whole Computational Biology and Bioinformatics Research Group of the Balearic Islands University for hosting me on multiple occasions, introducing me to problems related with the quartet index and for valuable comments on this manuscript. The simulated values of the quartet index for the Yule tree were provided by Gabriel Valiente and Tomás Martínez–Coronado. I was supported by the Knut and Alice Wallenberg Foundation and am by the Swedish Research Council (Vetenskapsrådet) grant no. 2017–04951. My visits to the Balearic Islands University were partially supported by the the G S Magnuson Foundation of the

Figure 2: Left: histogram of scaled and centred simulated values of the QIB for the Yule tree, Y_n^Q, right: histogram of Y_Q for the Yule model, $\beta = 0$. The mean, variance, skewness and excess kurtosis of the simulated values are $-3.177 \cdot 10^{-6}$, $6.321 \cdot 10^{-6}$, -0.308, -0.852 (left, simulated values) and $1.682 \cdot 10^{-5}$, $6.38 \cdot 10^{-6}$, -0.317, -0.834 (right, theoretical values by Bartoszek (2018)'s heuristic Algorithm 2 with recursion depth 15). For $\beta = 0$ the leading constant of the variance in Eq. (7) is $5/(24 \cdot 33075) \approx 6.299 \cdot 10^{-6}$.

References

D. Aldous. *The Continuum Random Tree II: An Overview*, pages 23–70. London Mathematical Society Lecture Note Series. Cambridge University Press, 1991.

D. Aldous. Probability distributions on cladograms. In *Random Discrete Structures*, pages 1–18. Springer–Verlag, 1996.

D. Aldous. Stochastic models and descriptive statistics for phylogenetic trees. *Statist. Sci.*, 16:23–34, 2001.

K. Bartoszek. Exact and approximate limit behaviour of the Yule tree's cophenetic index. *Math. Biosci.*, 303:26–45, 2018.

M. G. B. Blum and O. François. On statistical tests of phylogeny imbalance: The Sackin and other indices revisited. *Math. Biosci.*, 195:141–153, 2006.

M. G. B. Blum, O. François, and S. Janson. The mean, variance and limiting distribution of two statistics sensitive to phylogenetic tree balance. *Ann. Appl. Probab.*, 16(4):2195–2214, 2006.

H. Chang and M. Fuchs. Limit theorems for patterns in phylogenetic trees. *J. Math. Biol.*, 60:481–512, 2010.

B. Chen, D. Ford, and M. Winkel. A new family of Markov branching trees: the alpha–gamma model. *Electron. J. Probab.*, 14:400–430, 2009.

B. Chor and S. Snir. Analytic solutions of maximum likelihood on forks of four taxa. *Math. Biosci.*, 208:347–358, 2007.

D. H. Colless. Review of "Phylogenetics: the theory and practise of phylogenetic systematics". *Syst. Zool.*, 31:100–104, 1982.

M. Drmota. *Random Trees: an Interplay between Combinatorics and Probability*. Springer–Verlag, 2009.

J. Felsenstein. *Inferring Phylogenies*. Sinauer Associates Inc., Sundarland, U.S.A., 2004.

D. J. Ford. Probabilities on cladograms: introduction to the alpha model. *ArXiv e-prints*, 2005.

C. A. R. Hoare. Quicksort. *Comput. J.*, 5:10–15, 1962.

T. Martínez-Coronado, A. Mir, and F. Rosselló. The probabilities of trees and cladograms under Ford's α–model. *ArXiv e-prints*, 2018a.

T. Martínez-Coronado, A. Mir, F. Rosselló, and G. Valiente. A balance index for phylogenetic trees based on quartets. *ArXiv e-prints*, 2018b.

A. Mir, F. Rosselló, and L. Rotger. A new balance index for phylogenetic trees. *Math. Biosci.*, 241(1):125–136, 2013.

S. T. Rachev and L. Rüschendorf. Probability metrics and recursive algorithms. *Adv. Appl. Probab.*, 27:770–779, 1995.

S. Roch and S. Snir. Recovering the treelik trend of evolution despite extensive lateral gene transfer: a probabilisitic analysis. *J. Comp. Biol.*, 20(2): 93–112, 2013.

U. Rösler. A limit theorem for "Quicksort". *Theor. Inf. Applic.*, 25(1): 85–100, 1991.

U. Rösler. A fixed point theorem for distributions. *Stoch. Proc. Appl.*, 42: 195–214, 1992.

M. J. Sackin. "Good" and "bad" phenograms. *Syst. Zool.*, 21:225–226, 1972.

C. Semple and M. Steel. *Phylogenetics.* Oxford University Press, 2003.

M. Steel and A. McKenzie. Properties of phylogenetic trees generated by Yule–type speciation models. *Math. Biosci.*, 170:91–112, 2001.

G. U. Yule. A mathematical theory of evolution: based on the conclusions of Dr. J. C. Willis. *Philos. T. Roy. Soc. B*, 213:21–87, 1924.

Appendix: R code for simulating from the limit distribution of the quartet index

```
fCtau_QIB<-function(x,beta=0){
    PB4beta<-(3*beta+6)/(7*beta+18);
    PB4beta*(x^4)/24+PB4beta*((1-x)^4)/24-PB4beta/24+0.25*x*x*(1-x)^2
}

fdistribution_limitQIB<-function(num.iter=10,popsize=10000,Y0=0){
    replicate(popsize,fdraw_limitQIB(num.iter,Y0))
}

fdraw_limitQIB<-function(num.iter=15,Y0=0){
    res<-0
    if (num.iter==0){
        Y1<-Y0
        Y2<-Y0
    }
    else{
        Y1<-fdraw_limitQIB(num.iter-1,Y0)
        Y2<-fdraw_limitQIB(num.iter-1,Y0)
    }
    tau<-runif(1)
    res<-((tau)^4)*Y1+((1-tau)^4)*Y2+fCtau_QIB(tau)
    res
}

popsize<-10000 ## size of sample for histogram
num.iter<-15 ## depth of the recursion
Y0 <- 0 ## initial value

vlimitQIB<-fdistribution_limitQIB(num.iter=num.iter,popsize=popsize)
```