Synaptonemal complex architecture facilitates the chromosome-specific regulation of

recombination in Drosophila

Cori K. Cahoon*, G. Matthew Heenan*, Zulin Yu*, Jay R. Unruh*, Sean McKinney*, and R. Scott

Hawley*,‡

* Stowers Institute for Medical Research, Kansas City, MO, United States of America

‡ Department of Molecular and Integrative Physiology, University of Kansas Medical Center,

Kansas City, KS, United States of America

Corresponding Author:

R. Scott Hawley

Stowers Institute for Medical Research

Kansas City, MO 64110

Phone (816) 926-4427

Fax (816) 926-2060

rsh@stowers.org

17
18

Key words: synaptonemal complex, meiosis, homologous recombination, crossover, crossover

interference

Abstract

In Drosophila, meiotic recombination is initiated by the formation of programmed DNA double-

strand breaks (DSBs), which occur within the context of the synaptonemal complex (SC). To

better understand the role of the SC in mediating recombination we created an in-frame deletion

mutant in $c(3)G$ (deleting amino acids L340 to N550, denoted as $c(3)G^{cc\Delta 1}$), which encodes the

major transverse filament protein of the SC. Although $c(3)G^{cc\Delta 1}$ oocytes assemble ribbon-like

SC and exhibit normal DSB formation, the euchromatic SC precociously disassembles into

fragments that persist until mid to late pachytene in both $c(3)G^{cc\Delta 1}$ heterozygotes and

homozygotes. Centromeric SC, however, is unaffected in both genotypes. Thus, $c(3)G^{cc\Delta 1}$ is a

separation-of-function mutant that establishes different functional and structural requirements

between euchromatic and centromeric SC. Our data also demonstrate that the chromosome

arms differ in their sensitivity to $c(3)G^{cc\Delta 1}$-induced perturbations in the SC. The X chromosome is

distinctly sensitive to these perturbations, such that euchromatic pairing and crossing over are

altered in $c(3)G^{cc\Delta 1}$ heterozygotes and severely reduced in $c(3)G^{cc\Delta 1}$ homozygotes. On the

autosomes, crossovers are shifted to centromere-proximal regions and crossover interference is

defective in both $c(3)G^{cc\Delta 1}$ homozygotes and heterozygotes. However, only $c(3)G^{cc\Delta 1}$

homozygotes display a progressive loss of euchromatic pairing in distal autosomal regions,

suggesting that discontinuity in the euchromatic SC—rather than failed pairing—might cause the

altered crossover distribution. These phenotypes reveal that different chromatin states or

regions have differing requirements to maintain both the SC and homologous pairing.

Furthermore, $c(3)G^{cc\Delta 1}$ is the first mutant in Drosophila to demonstrate that the SC appears to

facilitate the regulation of recombination frequency and distribution differently on each

chromosome.

Author Summary

Chromosome segregation errors during meiosis are the leading cause of miscarriage and birth defects in humans. To prevent these errors from occurring, meiotic cells have evolved multiple mechanisms to ensure that each gamete receives exactly half the number of chromosomes. During meiosis I, this is accomplished by forming a crossover between homologous chromosomes, which is facilitated by a large protein complex called the synaptonemal complex (SC). The SC is assembled between homologous chromosomes during early prophase I, and it is unclear how the SC regulates the position and number of crossovers each homolog receives. To better understand the role of the SC in mediating recombination, we created an in-frame deletion mutant in *Drosophila melanogaster* in the gene encoding the C(3)G protein, the major transverse filament protein of the SC. Although mutant oocytes assemble ribbon-like SC, the SC along the chromosome arms precociously disassembles in early meiosis. Surprisingly, the SC around the centromeres is unaffected in these mutants, suggesting that the requirements for SC formation may differ depending on where the SC is located along the chromosomes. Our data also demonstrate that the chromosome arms differ in their sensitivity to the mutant-induced perturbations of the SC in both crossing over and homolog pairing.

INTRODUCTION

At the beginning of meiotic prophase, homologous chromosomes must identify each other, pair,

and initiate recombination. Meiotic recombination is critical for the proper segregation of

homologous chromosomes during meiosis I. The failure to properly exchange genetic

information will frequently result in missegregation of the chromosomes leading to eggs and

sperm with the incorrect number of chromosomes. Indeed, errors in meiotic chromosome

segregation are the leading cause of miscarriage and aneuploidy in humans, which can result in

chromosomal disorders such as Down syndrome and Turner syndrome (reviewed in [1]).

To initiate recombination, a series of programmed DNA double-strand breaks (DSBs)

are formed by the topoisomerase-like enzyme Spo11 [2]. Most organisms make a large excess

of DSBs—in some cases 30 times more than necessary—since only a small fraction of DSBs

will be repaired into crossovers [reviewed in 3]. Some organisms appear to use DSBs to identify

homology, and it is thought that creating more DSBs than necessary helps to ensure that

homologs are able to properly pair and that each chromosome receives at least one crossover

[reviewed in 4].

Neither DSBs nor crossovers are randomly distributed along chromosome arms, but

rather seem to be placed in a fairly ordered manner. Fine-scale genomic analysis of the

nonrandom distribution of DSBs in mice and yeast has indicated that the location of DSBs

involves a complicated series of factors working together on both a local DNA sequence level

and on a global chromatin architecture level [5, 6]. Moreover, the nonrandom distribution of the

resulting crossovers varies depending on the organism. Some organisms, like Drosophila, place

most of their crossovers in the middle of the chromosome arms, while other organisms, like

grasshoppers, planarians and nematodes, position crossovers at the distal regions of

chromosome arms [reviewed in 7] [8]. This suggests that there may be multiple mechanisms

that can determine the position of DSBs along the chromosome arms and that only certain

DSBs can become crossovers.

In Drosophila, approximately one crossover forms per chromosome arm [9, 10]. The

tight regulation of crossover number is controlled by multiple processes, including crossover

interference, the centromere effect, and region-specific constraints [reviewed in 11]. Once a

crossover has formed, crossover interference functions to suppress the formation of additional

crossovers nearby. Similarly, the centromere effect and regional constraints function to prevent

the formation of crossovers in or next to specific chromatin states, such as the centromere and

heterochromatin. All of these processes function over megabases of DNA to constrain the

crossover number in each meiosis to an average of six crossovers across the three crossover-

competent chromosomes in Drosophila. Exactly how interference, the centromere effect, and

region-specific constraints function is not well understood.

Double-strand break formation and crossing over in Drosophila occur within the context

of full-length synaptonemal complex (SC) (Fig 1A) [15, 16]. The SC, a large protein complex

that assembles between paired homologous chromosomes, is required for the formation of

crossovers. It is therefore possible that the SC may play a role in regulating the crossover fate

of DSBs. Additionally, the SC connects two homologs along their entire lengths, making it an

ideal candidate structure for monitoring DSB formation and crossover maturation occurring at

different places along the chromosomes. Indeed, several studies have strongly suggested the

SC plays a role in mediating crossover interference [17, 18].

The synaptonemal complex is arranged in three parts: lateral elements (LEs), a central

region (CR), and a central element (CE) (Fig 1B). The LEs assemble on each homolog

alongside the axis components, which consist of cohesin and cohesin-like proteins that are

thought to establish the meiotic chromatin architecture [reviewed in 19, 20]. The LEs interact

with the CR proteins, which occupy the space between the two homologs. Transverse filament

proteins span the distance of the CR, thereby connecting the LEs of each homolog. In the

middle of the CR is the CE, which consists of proteins that are thought to help stabilize the

transverse filaments.

Genetic analyses have identified multiple components of the Drosophila SC, and recent

advances in superresolution microscopy have allowed SC components to be precisely

positioned within the overall SC structure (Fig 1B). The Drosophila SC forms two layers that are

mirror images of one another [21]. C(3)G, the major transverse filament protein of the

Drosophila SC [22], forms a homodimer, with the C-terminal ends of each subunit positioned in

the LE and the N-terminal ends in the CE [23]. The N terminus of one homodimer is then

thought to interact with the N terminus of another homodimer to span the distance between the

two homologs, thus connecting the two homologous LEs [23]. It is currently unclear exactly how

C(3)G interacts with the LE. An additional CR protein, Corolla, was recently identified [24].

Corolla, resides within the CR as two parallel tracks and has been shown by yeast two-hybrid

analysis to interact with the CE protein Corona (CONA) [21, 24, 25], which also forms two

parallel tracks within the CE [21, 26]. C(3)G, Corolla, and CONA depend on each other in order

to assemble the SC, and all three proteins are required for the formation of crossovers [22, 24-

26].

The first step toward dissecting the functional anatomy of the C(3)G protein involved

creating in-frame deletion mutations, which showed that the N- and C-terminal globular domains

of C(3)G are required to properly assemble the SC and for robust crossover formation [27].

Here, we describe the effects of a mutant, $c(3)G^{cc\Delta 1}$, with an in-frame deletion of a coiled-coil

domain. To properly characterize this mutant, we created a CRISPR/Cas9 allele, eliminating

any of the common concerns regarding levels of expression that are inherent when using a

randomly inserted transgene. (A transgene in-frame deletion mutant, similar to the one in this

manuscript, has been previously created, but was never extensively analyzed [22]).

$c(3)G^{cc\Delta 1}$ removes 211 amino acids from the largest coil-coiled domain of C(3)G which,

as a homozygote, shortens the width of the SC by ~40 nm and thus reduces the distance

between the homologs. Both heterozygotes and homozygotes of $c(3)G^{cc\Delta 1}$ are able to assemble

the SC, display normal DSB formation, and affect neither the persistence of SC at the

centromere nor centromere clustering. However, euchromatic SC disassembles prematurely in

both genotypes, resulting in fragmented or discontinuous SC along the length of the

chromosomes. This premature SC disassembly affects both homolog pairing and recombination,

but to different degrees on each chromosome. $c(3)G^{cc\Delta 1}$ is the first Drosophila mutant to

illustrate a chromosome-specific regulation of recombination, where each chromosome has

different requirements for the SC in maintaining homolog pairing and facilitating recombination.

RESULTS

Novel loss-of-function allele of *c(3)G* affects the width of the SC

A large deletion in the yeast transverse filament protein Zip1 decreases the distance between

the LEs, suggesting that Zip1 is directly involved in determining the width of the SC [28]. Since

Zip1 and C(3)G are functional homologs of each other, we postulated that a similar deletion in

C(3)G may alter the Drosophila SC in a similar manner to the Zip1 deletion in yeast. To that

end, we created a CRISPR/Cas9-mediated in-frame deletion in the Drosophila *c(3)G* gene (see

Methods) that removes 211 amino acids from the second coiled-coil domain in C(3)G (Fig 1B).

We call this mutant $c(3)G^{cc\Delta 1}$.

Based on the 1.485-angstrom size of an amino acid in a coiled-coil, the mathematically

calculated decrease in SC width expected by deleting 211 amino acids from a coiled-coil

configuration is ~63 nm. However, when using single-molecule superresolution techniques such

as stochastic optical reconstruction microscopy (STORM), which require primary and secondary

antibodies, the size of an antibody complex (20–30 nm) will introduce uncertainty into the

precise positioning of the structure being imaged. To account for this, we added the size of the

primary and secondary antibody complex, which recognizes the C-terminus C(3)G, to the

calculated decrease in SC width resulting in an expected decrease of ~43–53 nm in $c(3)G^{cc\Delta 1}$

homozygotes.

Using STORM on intact germaria with an antibody that recognizes the C-terminal end of

C(3)G, we observed two tracks of the C(3)G C-termini, suggesting that the SC was able to

assemble both C(3)G homodimers (Fig 2). Additionally, when we measured the distance

between these two tracks, referred to as the SC width, it was decreased by ~40 nm in $c(3)G^{cc\Delta 1}$

homozygotes (Fig 2; wild type, 119 nm ± SE 1.0; $c(3)G^{cc\Delta 1}$, 78.6 nm ± SE 0.3; P<0.0001), which

closely matched the expected decrease in SC width. This suggests that, like Zip1, C(3)G is

responsible for determining the distance between the LEs in Drosophila.

STORM analysis of $c(3)G^{cc\Delta 1}$ heterozygotes, in which flies express one $c(3)G^{cc\Delta 1}$ version

and one wild type (WT) version of C(3)G, showed an intermediate change in SC width (Fig 2;

107 nm ± SE 0.4; P<0.0001 when compared to WT). This suggests that $c(3)G^{cc\Delta 1}$ heterozygotes

incorporate a combination of $c(3)G^{cc\Delta 1}$ and WT versions of C(3)G into the SC. One possible way

to accomplish this heterogeneity would be to alternate the C(3)G tetramers such that every

other tetramer is the $c(3)G^{cc\Delta 1}$ mutant. In this case, we would expect the profiles on either C-

terminal side of C(3)G to be thicker. However, this is not likely to be the case because the

thicknesses of the C-terminal C(3)G signal are nearly the same for the $c(3)G^{cc\Delta 1}$ heterozygotes

and homozygotes (WT: 62 nm; $c(3)G^{cc\Delta 1}$ heterozygote: 49 nm; $c(3)G^{cc\Delta 1}$ homozygotes: 43 nm).

It is also possible that $c(3)G^{cc\Delta 1}$ and WT C(3)G monomers are able to dimerize and create mixed

C(3)G homodimers, or that the N termini of $c(3)G^{cc\Delta 1}$ homodimers can interact with the N termini

of WT homodimers in the CE of the SC to create a mixed tetramer. Both possibilities would

result in the intermediate SC width seen in $c(3)G^{cc\Delta 1}$ heterozygotes. Given the heterogeneous

nature of STORM data, it is impossible to distinguish between these possibilities. Further

studies with alternative methodologies will be needed to gain a more detailed understanding of

this ultrastructure.

Both $c(3)G^{cc\Delta 1}$ heterozygotes and homozygotes initiate SC assembly at the centromeres

during the pre-meiotic mitotic divisions (region 1) and at the onset of meiosis in zygotene/early

pachytene (region 2A) assembled ribbon-like SC with normal kinetics (Fig 3). Moreover, further

analysis of the SC in region 2A by measuring the intensity of the antibody staining of Corolla

between the mutants and WT showed that although the SC appeared to be ribbon-like in the

mutants the intensity was decreased compared to WT (Fig S1). This suggested either that both

$c(3)G^{cc\Delta 1}$ heterozygotes and homozygotes may not be assembling as much SC as in WT or that

both mutants have altered the tripartite structure of the SC. However, based on our ability to

resolve two C(3)G C-terminal tracks in the STORM data and on Corolla being able to assemble

into ribbon-like SC in the deconvolution microscopy data (Fig 3, Figure S1), we do not believe

that the decrease in SC intensity indicates a lack of tripartite SC. All of the known components

of the SC in Drosophila are co-dependent on each other to be able to assemble, thus if the

structure of the SC was failing in these mutants then we would expect both C(3)G and Corolla to

be unable to assemble into ribbon-like SC [24, 25].

$c(3)G^{cc\Delta 1}$ affects the persistence of SC along the chromosome arms but not at the

centromere

Previous studies have shown that *c(3)G* null mutants initiate DSBs, but at reduced levels

compared to WT [16, 29]. This suggests that some DSBs will only form in the presence of SC.

In *c(3)G*^{ccΔ1} heterozygotes and homozygotes, DSB formation appears to be normal. To identify

DSBs, we used an antibody that recognizes the phosphorylated form of the histone variant

H2AV (referred to as γH2AV), and this phosphorylation mark is one of the first events that

occurs following the formation of DSBs [30]. At zygotene/early pachytene (region 2A) all 16-cells

within the cyst will induce DSBs, and for this assay we scored only DSBs in the meiotic cells,

were identified by the presence of the SC [16]. Nurse cells also induce DSBs, but they do not

assemble the SC and were excluded from this analysis [16].

In an otherwise WT background, both *c(3)G*^{ccΔ1} heterozygotes and homozygotes

displayed near WT levels of γH2AV foci in zygotene/early pachytene (region 2A) (Fig S2; WT=

average 6 ± SE 0.6; *c(3)G*^{ccΔ1} homozygotes= average 6 ± SE 0.6, P=0.77; *c(3)G*^{ccΔ1}

heterozygotes= average 7 ± SE 0.8, P=0.69). By early pachytene (region 2B), the number of

γH2AV foci was reduced to ~1 focus in WT (average 1 ± SE 0.3) and *c(3)G*^{ccΔ1} heterozygotes

(average 1 ± SE 0.2, P=0.06). *c(3)G*^{ccΔ1} homozygotes displayed a wider range of γH2AV foci

with an average of 3 DSBs (± SE 0.5) in early pachytene (region 2B). This was statistically

different from WT (P=0.036), but it is unclear if this indicates a delay in DSB repair or the

initiation of more DSBs. By mid pachytene (region 3) all γH2AV foci were absent in both

c(3)G^{ccΔ1} homozygotes (P=0.54) and heterozygotes (P=0.17). Thus, while *c(3)G*^{ccΔ1}

homozygotes may have some anomalies in DSB repair or late-DSB initiation in early pachytene,

by mid pachytene the repair of these breaks follows similar to WT timing.

In WT, euchromatic SC initiates disassembly in mid/late pachytene (stages 5–7) and the

SC is fully disassembled from the chromosome arms by stages 8–9. Interestingly, both *c(3)G*^{ccΔ1}

homozygotes and heterozygotes initiated premature disassembly of euchromatic SC as cells

progressed into early and mid pachytene (Fig 3). *c(3)G*^{ccΔ1} homozygotes began to initiate

euchromatic SC disassembly in early pachytene (region 2B), whereas $c(3)G^{cc\Delta 1}$ heterozygotes

started disassembling euchromatic SC in mid pachytene (region 3). Furthermore, this pattern

was recapitulated when the antibody intensity of Corolla was quantified in each mutant at each

stage of meiosis (Fig S1). Thus, the presence of one WT copy of C(3)G in a $c(3)G^{cc\Delta 1}$

heterozygote delays the onset of premature SC disassembly but cannot fully rescue the defect,

demonstrating the semi-dominant nature of the $c(3)G^{cc\Delta 1}$ mutant. In both homozygotes and

heterozygotes, the fragmented SC persisted until stages 5–7 (mid/late pachytene) and was fully

disassembled by stages 8–9, similar to when euchromatic SC is normally disassembled from

the chromosome arms (Fig 3; Fig S1). Surprisingly, $c(3)G^{cc\Delta 1}$ mutants were able to maintain SC

at the centromeres, suggesting that the SC at the centromere is unaffected by this deletion.

In Drosophila, centromeres begin to homologously pair and initiate the assembly of

centromeric SC during the four premeiotic mitotic divisions (region 1) (Fig 1A, Fig 3)[13]. At the

transition from the premeiotic divisions to meiotic prophase (region 2A) the paired centromeres

cluster together into 1–3 masses, and this process is dependent on SC assembly [12, 14]. Since

$c(3)G^{cc\Delta 1}$ homozygotes and heterozygotes were able to maintain the SC at the centromeres with

WT kinetics, we assayed centromere clustering to determine if this SC was functional. To do

this, we counted the number of centromere foci in nuclei containing SC using the centromeric

nucleosome Centromere identifier (CID) as a marker for the centromeres. In all of the stages

analyzed, centromere clustering was unaffected by the presence of $c(3)G^{cc\Delta 1}$ (Fig S3, Table

S1). Also, $c(3)G^{cc\Delta 1}$ did not display a dominant effect on centromere clustering when assayed as

a heterozygote (Fig S3, Table S1). For $c(3)G^{cc\Delta 1}$ homozygotes, we do observe an increase in

the number of nuclei with 3 centromeric foci in early pachytene. However, in the $c(3)G^{68}$ null

mutant, most of the nuclei have 4 or more CID foci at this stage, which does not occur in the

$c(3)G^{cc\Delta 1}$ mutants [14]. Thus, these data demonstrate that centromeric SC is not dependent on

the deleted amino acids in $c(3)G^{cc\Delta 1}$ to function in clustering centromeres.

255 ***c(3)G^{ccΔ1}* causes defects in both recombination and euchromatic pairing on the *X***

256 **chromosome**

257 Previous studies have shown that in most (but not all) organisms the SC is required for the

258 formation of crossovers [reviewed in 19, 31]. In Drosophila, Gowen showed that flies

259 heterozygous for the *c(3)G* null mutation are indistinguishable from WT with respect to

260 recombination on all three crossover-competent chromosomes, demonstrating that one copy of

261 C(3)G is sufficient for normal recombination [32]. Moreover, flies homozygous for the *c(3)G* null

262 mutation are completely unable to repair DSBs into either crossovers or noncrossover gene

263 conversions on all three crossover competent chromosomes [16, 22, 32-34]. Therefore, the

264 question of how these breaks are repaired remains open [34].

265 Unlike *c(3)G* null mutants, *c(3)G^{ccΔ1}* does initially assemble ribbon-like SC, which

266 prematurely disassembles into fragments as the cells progress through early to mid pachytene.

267 To determine if the SC assembled in *c(3)G^{ccΔ1}*-bearing oocytes is able to promote crossing over,

268 we assayed for the presence of crossovers in both *c(3)G^{ccΔ1}* homozygotes and heterozygotes.

269 Interestingly, *c(3)G^{ccΔ1}* homozygotes and heterozygotes displayed different phenotypes with

270 regard to recombination on the *X* chromosome (Fig 4A; Table 1). Both genotypes were able to

271 from at least some crossovers, suggesting that the region deleted in *c(3)G^{ccΔ1}* is not required to

272 promote crossover formation per se. However, the crossovers made in *c(3)G^{ccΔ1}* heterozygotes

273 displayed an altered distribution along the length of the *X* chromosome, while *c(3)G^{ccΔ1}*

274 homozygotes severely reduced the frequency of crossing over on the *X*.

275 In WT, the majority of crossovers are known to occur in the middle one-third of the

276 chromosome arms and are inhibited near the centromeres and telomeres. Indeed, in the WT

277 control for our *X* chromosome recombination assay, most of the crossovers occurred between

278 the markers *vermillion* (*v*) and *forked* (*f*), which reside in the medial one-third of the *X*

chromosome (Fig 4A; Table 1 and S2). However, in $c(3)G^{cc\Delta1}$ heterozygotes, the crossover

distribution was shifted toward the centromere-proximal region of the chromosome, such that

most crossovers occurred between the *forked* and *yellow*$^+$ (y^+) markers that encompass the

centromere. This crossover distribution shift resulted in a corresponding decrease in crossovers

in the distal chromosome regions, between the markers *scute* (*sc*) and *crossveinless* (*cv*). Also,

the overall map length of the *X* chromosome was slightly decreased in $c(3)G^{cc\Delta1}$ heterozygotes

(51.2 cM in $c(3)G^{cc\Delta1}$ heterozygotes vs. 63 cM in WT; Table 1). Thus, $c(3)G^{cc\Delta1}$ heterozygotes

appear to alter the distribution of crossovers, without greatly changing the total number of

crossovers on the *X*.

In contrast, $c(3)G^{cc\Delta1}$ homozygotes severely reduced crossing over on the *X*

chromosome (Fig 4A; Table 1). The total map length of the *X* chromosome decreased by more

than 80%, to 11.8 cM from 63.0 cM in WT, with the majority of the reduction in the medial and

distal intervals. However, even with this reduction the distribution of crossovers seems to

display the same centromere-proximal shift as in $c(3)G^{cc\Delta1}$ heterozygotes.

A more comprehensive approach for understanding these reductions in exchange lies in

the application of Weinstein's method to calculate the frequency of bivalents that underwent

zero (E_0), one (E_1), or two (E_2) crossover events [36]. This method accounts for the fact that a

single recombination event between two homologs will produce two single crossover and two

noncrossover chromatids (only one of which is recovered) and that a double crossover event

(depending on whether it is a two-strand, a three-strand, or a four-strand double) will produce

some combination of noncrossover, single crossover, or double crossover chromatids (again,

only one of which is recovered). In WT, the frequency of noncrossover bivalents (or E_0

bivalents) was 18.9%, while in $c(3)G^{cc\Delta1}$ homozygotes it was 79.2%—a four-fold increase in the

frequency of an oocyte receiving a noncrossover bivalent (Table 1). The remaining ~20% of *X*

chromosomal bivalents in $c(3)G^{cc\Delta1}$ homozygotes did undergo a single crossover event, creating

single crossover (or E_1) bivalents and the vast majority of these single crossovers occurred in

centromere-proximal intervals (Fig 4A).

The presence of SC between homologous chromosomes is thought to keep them paired

throughout pachytene, which promotes crossing over between two homologous chromosomes.

Since the SC is prematurely disassembled in $c(3)G^{cc\Delta 1}$, it is possible that homologous

chromosome pairing may be lost before crossovers can be established, which may explain the

altered crossover distribution. We analyzed homolog pairing using euchromatic FISH probes at

distal (3C3–3C7) and proximal (15C1–15D6) regions on the X chromosome (Fig 4B; Table S2

and S3). A locus was considered paired if the distance between the two FISH probe foci was

less than 0.75 µm and unpaired if foci exhibited a distance 0.75 µm or greater.

In $c(3)G^{cc\Delta 1}$ heterozygotes, both distal and proximal probes were paired in nearly 100%

of the oocytes in zygotene/early pachytene (region 2A) (Fig 4B, Table S3). However, the distal

probe was progressively less paired as the cells progressed to early pachytene (region 2B) and

mid pachytene (region 3). The proximal probe remained nearly 100% paired throughout these

regions. Thus, the centromere-proximal shift in recombination was correlated with a loss of

distal euchromatic pairing and a maintenance of proximal euchromatic pairing on the X

chromosome in $c(3)G^{cc\Delta 1}$ heterozygotes.

Unlike $c(3)G^{cc\Delta 1}$ heterozygotes, $c(3)G^{cc\Delta 1}$ homozygotes displayed a progressive loss of

euchromatic pairing at both proximal and distal loci on the X chromosome, suggesting that the X

chromosomes may be unpaired by mid pachytene (Fig 4B, Table S3). Based on our FISH

assay, the proximal locus was able to maintain pairing more effectively than the locus detected

by the distal probe, which may explain why the few crossovers observed in $c(3)G^{cc\Delta 1}$

homozygotes occur in the centromere proximal region of the X chromosome. Indeed, we

propose that the loss of homolog pairing caused by a progressive loss of X chromosome SC is

the reason for the significant decrease in *X* recombination observed in *c(3)G^{ccΔ1}* homozygotes. If

we are correct in this asserion, then the observation that homolog pairing is not drastically

reduced in *c(3)G^{ccΔ1}* homozygotes until early pachytene (region 2B) suggests that crossover

designation on at least the *X* chromosome may be occurring either multiple times throughout

pachytene or later in pachytene than was previously thought.

Autosomal crossover distribution is shifted to centromere-proximal regions

Meiotic mutants that decrease recombination on the *X* chromosome also display similar

decreases in recombination on the autosomes [22, 27, 29, 37, 38]. It was therefore surprising

that both *c(3)G^{ccΔ1}* homozygotes and heterozygotes displayed robust recombination on the

autosomes with the map lengths of each autosome increasing by more than 10 map units,

suggesting that each chromosome received more crossovers than normally would occur in WT

(Fig 4C,E; Tables 2 and 3). Indeed, to our knowledge, *c(3)G^{ccΔ1}* is the first mutant in Drosophila

to exhibit such dramatic differences in terms of its effects on recombination on the *X*

chromosome compared to the autosomes. Moreover, the biological significance of finding that

the *X* chromosome may be more sensitive to *c(3)G^{ccΔ1}*-induced changes in the SC than the

autosomes may be that the various chromosome arms differ in their requirements for proper SC

structure and function.

Although overall recombination on the autosomes was robust, the crossover distribution

was shifted to the centromere-proximal regions in both *c(3)G^{ccΔ1}* homozygotes and

heterozygotes. On the 3^{rd} chromosome, we observed a slight decrease in recombination in the

distal interval [*roughoid–hairy (ru–h)*] of both homozygotes and heterozygotes (Fig 4E; Table 3),

while on the 2^{nd} chromosome, only *c(3)G^{ccΔ1}* homozygotes displayed a decrease in the distal

interval *net-decapentaplegic (net–dpp)* (Fig 4C; Table 2).

However, the most striking effects of the $c(3)G^{cc\Delta1}$ mutants on autosomal crossover

distribution can be seen by considering the intervals that span their centromeres (Fig 4C,E;

Tables 2, 3 and S2). On the 2^{nd} chromosome, exchange in the pericentromeric *purple* to

cinnabar (*pr–cn*) interval increased more than three fold in both heterozygotes and

homozygotes when compared to WT. Moreover, on the 3^{rd} chromosome, both $c(3)G^{cc\Delta1}$

heterozygotes and homozygotes also showed a greater than 300% increase in recombination

frequency for the centromere spanning interval *scarlet* to *curled* (*st–cu*). Indeed, for the 3^{rd}

chromosome more than one third of the crossover events occurred in the pericentromeric *st–cu*

interval (as compared to 12% in wild type). Thus, both autosomes displayed a substantial

increase in pericentromeric recombination frequency compared to WT, suggesting that the

$c(3)G^{cc\Delta1}$ mutation may alleviate constraints that mediate recombination in such intervals, such

as centromere effect, which normally functions to restrict recombination in these intervals.

Since the *X* chromosome displayed both a centromere-proximal shift in recombination

and a premature loss in homolog pairing, we wondered if the discontinuous SC in $c(3)G^{cc\Delta1}$ was

affecting homolog pairing on the autosomes in a similar manner. In $c(3)G^{cc\Delta1}$ homozygotes,

these decreases in recombination were paralleled by progressive decreases in euchromatic

pairing for the distal probe on each autosome (Fig 4C–F; Tables 2, 3 and S3). Both the proximal

and medial probes remained relatively well paired on the autosomes throughout zygotene to

mid pachytene (region 2A to region 3) in $c(3)G^{cc\Delta1}$ homozygotes.

$c(3)G^{cc\Delta1}$ heterozygotes were able to maintain chromosome pairing along the entire

length of both the 2^{nd} and 3^{rd} chromosomes throughout zygotene to mid pachytene (region 2A to

region 3) (Fig 4D,F; Table S3). Because both $c(3)G^{cc\Delta1}$ homozygotes and heterozygotes

displayed the same altered crossover distribution, but only $c(3)G^{cc\Delta1}$ homozygotes lost pairing in

the distal chromosomal regions, the loss of homolog pairing in the distal regions of the

autosomes is likely not the cause of the centromere-proximal shift in crossing over. Rather, our

observations in $c(3)G^{cc\Delta 1}$ heterozygotes suggest that discontinuity of the SC may cause the shift

in crossover distribution. It is possible that because centromeric SC appears to be unaffected by

the $c(3)G^{cc\Delta 1}$ mutation, the SC near the centromeric heterochromatin is more stable, thus

making it more suitable for crossovers.

$c(3)G^{cc\Delta 1}$ homozygotes and heterozygotes exhibit greatly reduced crossover interference

It is known that intact SC is required for crossover interference, which acts to regulate the

number of crossovers formed on each chromosome arm by inhibiting crossovers from forming

within several megabases of each other [reviewed in 4, 31]. In other organisms, mutants that

display fragmented SC and/or chromosome axes have defects in crossover interference that

result in an increase in the number of crossovers [reviewed in 4]. On the *X* chromosome neither

$c(3)G^{cc\Delta 1}$ homozygotes nor heterozygotes displayed an increase in crossing over, in fact

$c(3)G^{cc\Delta 1}$ homozygotes caused a severe reduction in crossing over (Fig 4A). Thus, it is not

surprising that crossover interference is unaffected on the *X* chromosome in either $c(3)G^{cc\Delta 1}$

homozygotes or heterozygotes; but rather the observed loss of recombination is the result of a

loss of homolog pairing and/or synapsis (Table 1).

However, unlike the *X* chromosome, the autosomes displayed an increase in the

frequency of crossing over in $c(3)G^{cc\Delta 1}$ homozygotes and heterozygotes suggesting that

crossover interference may be affected (Fig 4C,E). Indeed, crossover interference is reduced in

$c(3)G^{cc\Delta 1}$ mutants on chromosome *2*. $c(3)G^{cc\Delta 1}$ heterozygotes have a complete or near complete

loss of crossover interference across the entire chromosome (Table 2). This decrease in

interference allows for an increase in the occurrence of double and triple crossovers (Table 2)

and results in an increase in map length [WT = 52.2 cM; $c(3)G^{cc\Delta 1}$ homozygotes = 67.6 cM;

c(3)G$^{cc\Delta 1}$ heterozygotes = 72.8 cM]. It is possible that the decreased interference on the 2nd

chromosome results from the fragmented SC observed in the *c(3)G$^{cc\Delta 1}$* mutants.

On chromosome *3, c(3)G$^{cc\Delta 1}$* homozygotes displayed levels of crossover interference

similar to WT in the distal regions of the chromosomes (Table 3). However, in the centromere-

proximal regions [*thread/scarlet/curled* (*th/st/cu*)], the effect of interference seemed to get

significantly weaker (WT = –4.5; *c(3)G$^{cc\Delta 1}$* homozygotes = –12.6). While we did not expect to

observe positive crossover interference across the centromere due to the centromere effect and

other potential factors functioning to suppress crossing over in this region, the large increase in

negative crossover interference was unexpected. Furthermore, this increase in negative

crossover interference may explain the observed increase in map length for *c(3)G$^{cc\Delta 1}$*

homozygotes on chromosome *3* (WT = 50.9 cM; *c(3)G$^{cc\Delta 1}$* homozygotes = 59.6 cM). Curiously,

c(3)G$^{cc\Delta 1}$ heterozygotes displayed near-WT levels of crossover interference across all the

intervals assayed on chromosome *3* (Table 3). Thus, the observed increase in map length in

c(3)G$^{cc\Delta 1}$ heterozygotes cannot be explained by a loss in crossover interference (WT = 50.9 cM;

c(3)G$^{cc\Delta 1}$ heterozygotes = 63.8 cM).

Chromosome segregation is unaffected by the *c(3)G$^{cc\Delta 1}$* mutation

It is well known that a decrease in recombination on the *X* chromosome should result in an

increase in frequency of *X* chromosome nondisjunction, such that the frequency of *X*

chromosome nondisjunction is equal to the frequency of E_0 cubed [39]. This reflects the

requirement for a pair of nonexchange autosomes to mediate missegregation of nonexchange *X*

chromosomes [40, 41]. Curiously, we did not observe this expected increase in the frequency of

X nondisjunction in *c(3)G$^{cc\Delta 1}$* homozygotes (Table S4).

Moreover, mutants that decrease recombination on the *X* chromosome and increase the

frequency of *X* chromosome nondisjunction typically also decrease recombination on the

423 autosomes [37, 38]. However, $c(3)G^{cc\Delta 1}$ homozygotes showed robust recombination on the

424 autosomes (Fig 4). Previous studies have shown that the distributive system, which normally

425 functions to segregate the obligately achiasmate 4^{th} chromosomes, can accurately segregate

426 two achiasmate chromosomes, such as the *X* and 4^{th} chromosomes [41]. Therefore, in $c(3)G^{cc\Delta 1}$

427 homozygotes, it is likely that the lack of nondisjunction is due to the distributive system's ability

428 to accurately segregate the two achiasmate *X* and 4^{th} chromosomes. The mechanism of how

429 the distributive system accomplishes accurate segregation without a crossover is unclear, thus

430 future studies are needed to elucidate this mechanism.

DISCUSSION

433 Although our understanding of SC structure has been greatly facilitated by advancements in

434 superresolution microscopy, which have revealed a more precise three-dimensional model of

435 SC ultrastructure, there is still much to learn about SC function. The multiple roles the SC plays

436 during meiotic prophase illustrate the functional importance of this protein complex in ensuring

437 the successful transmission of genetic information from one generation to the next. However,

438 because the SC is integral for the proper execution of so many meiotic processes, it has been

439 difficult to elucidate which SC proteins or protein domains are required for which functions.

440 Here, we show that a deletion in the large coiled-coil domain of C(3)G reveals two different

441 requirements for the SC: first, the deleted coiled-coil region in $c(3)G^{cc\Delta 1}$ is required to maintain

442 both SC structure along the chromosome arms and homolog pairing during meiotic prophase;

443 and second, the SC's ability to facilitate recombination depends upon its full-length assembly. In

444 addition, each chromosome appears to respond differently, on both local and global levels, to

445 structural changes in the SC.

446 **Centromeric SC versus euchromatic SC**

The difference between the kinetics of SC assembly and disassembly along the chromosome

arms versus at the centromeres suggests that the SC at the centromeres might be structurally

different than the SC along the chromosome arms. Indeed, it is known that the SC at the

centromeres requires certain proteins that euchromatic SC does not. For example, the cohesion

complex proteins ORD, SUNN, and SOLO are required for assembly of centromeric SC but not

the assembly of euchromatic SC [42-45], while euchromatic—but not centromeric—SC requires

the LE protein C(2)M to completely synapse homologs [12]. We now show in $c(3)G^{cc\Delta1}$ mutants

that a large region of the major coiled-coil domain in C(3)G is also dispensable for both

centromeric SC assembly and centromere clustering.

This difference between the SC at the euchromatin and at the centromere is not unique

to Drosophila. In yeast, the transverse filament protein Zip1 also persists long after the SC along

the chromosome arms has disassembled, and it is necessary for accurate achiasmate

chromosome segregation [46, 47]. Similarly, mice spermatocytes retain CR components of the

SC at the centromeres long after the SC has disassembled from the chromosome arms [48]. In

both mice and Drosophila, it is unclear why the SC persists at the centromeres, but it is possible

that centromeric SC in these organisms may function in achiasmate chromosome segregation,

as it does in yeast. Although errors in Drosophila female meiosis and mice spermatocytes are

rare, they do occur. Thus, persistence of centromeric SC may act as a backup segregation

mechanism to ensure proper chromosome segregation when a rare error does occur.

Chromosome-specific regulation of recombination

All previously characterized mutants in Drosophila that affect recombination on the *X*

chromosome show a similar, if not identical, effect on the autosomes. Prior to $c(3)G^{cc\Delta1}$, all

known SC mutants abolished or severely reduced recombination on all three crossover

chromosomes [22, 24, 25, 27]. $c(3)G^{cc\Delta1}$ is unique when compared to other SC mutants

because it differentially affects recombination on each chromosome, which suggests that each

chromosome has different rules for positioning crossovers along each chromosome arm [9].

While subtler than the differences observed between the *X* chromosome and the

autosomes, the differences between the two autosomes in $c(3)G^{cc\Delta 1}$ was indeed surprising.

Similar differences have been observed before. Miller et al. found that chromosome *2* displayed

a stronger effect of crossover interference than chromosome *3* [9]. They noticed that the 2^{nd}

chromosome rarely had double crossovers (out of 100 flies, only 14 double crossovers were

detected) and the double crossovers were positioned on average 11.5 Mb apart [9]. Meanwhile,

the 3^{rd} chromosome exhibited nearly twice as many double crossovers as the 2^{nd} chromosome

(out of 100 flies, 27 double crossovers were detected), and the double crossovers were

positioned on average 11.0 Mb apart [9, 49].

In $c(3)G^{cc\Delta 1}$ homozygotes and heterozygotes, both of the autosomes display defects in

crossover interference that result in an increase in the number of crossovers, but the 2^{nd}

chromosome seems to display a stronger effect. Perhaps this result is best interpreted in the

context of Miller et al.'s characterization of double crossovers on the two autosomes [9].

Because the 2^{nd} chromosome relies heavily on crossover interference to regulate crossing over,

a loss of crossover interference, like that observed in $c(3)G^{cc\Delta 1}$ homozygotes and

heterozygotes, would cause a large increase in the number of crossovers. Perhaps, then, the 3^{rd}

chromosome places more emphasis on an alternative mechanism, such as the centromere

effect, to regulate crossing over instead of crossover interference. This may explain why the

autosomes display subtle differences in recombination when crossover interference is

diminished to a similar degree in $c(3)G^{cc\Delta 1}$ homozygotes and heterozygotes.

Additionally, $c(3)G^{cc\Delta 1}$ demonstrates the importance of assaying multiple chromosomes

when mutants are defective in recombination. Although numerous previous characterizations of

recombination-defective mutants in Drosophila have only measured recombination on one

chromosome [16, 22, 24], our data from the $c(3)G^{cc\Delta 1}$ homozygotes illustrate how misleading it

can be to assay only one chromosome. In this case, the X chromosome is more sensitive to the

$c(3)G^{cc\Delta 1}$ mutation than are the autosomes. Differences between the sex chromosomes and the

autosomes are not specific to Drosophila. For example, studies in *Caenorhabditis elegans* found

that the X chromosome initiated meiotic DNA replication later than the autosomes [50, 51].

While the X chromosome and the autosomes seemed to homologously pair during the same

small window of time in *C. elegans*, complete synapsis of the X chromosome was delayed

and/or slower than the autosomes [50]. Currently, it is unknown in Drosophila if X chromosome

synapsis is delayed, but our data support a view that initiating a synapsis delay of the X

chromosome may result in a later initiation of recombination on the X chromosome.

Multiple factors work in concert to regulate recombination, and the SC appears to play a

vital role in this regulation. The SC is required for crossovers in all organisms that assemble the

structure, and $c(3)G^{cc\Delta 1}$ has demonstrated an additional role of the Drosophila SC in regulating

the positions of crossovers. Future studies are needed to determine the mechanism of how

C(3)G and the SC regulate recombination. Additionally, $c(3)G^{cc\Delta 1}$ is a large deletion mutant, thus

future experiments dissecting this deletion may reveal insights into this mechanism.

MATERIALS AND METHODS

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the Lead Contact, R. Scott Hawley (RSH@stowers.org).

Experimental model and subject details

The Drosophila stocks used in the foregoing assays were kept at a humidity-controlled 25°C

and on standard food. All mention of "wild type" (WT) refers to the genotype: *y w; +/+; +/+; sv^{spa-}

^{pol}, unless stated otherwise. The key resource table contains a list of all the fly stocks used in

this manuscript.

CRISPR/Cas9 generation of the c(3)G^{ccΔ1} deletion flies

To aid in screening for *c(3)G^{ccΔ1}* mutant flies, we incorporated a piggy BAC transposon carrying

a 3xP3-DsRed that expresses in the fly eye into the intron directly downstream of the *c(3)G^{ccΔ1}*

deletion [intron 5 of *c(3)G*] in the homologous repair template plasmid. The piggy BAC

transposon plasmid was constructed to have flanking AarI and SapI restriction sites (Addgene

51434). We used PCR to obtain two fragments of *c(3)G* from the Drosophila genome that

flanked the position where the piggy BAC would be inserted and added in either an AarI or SapI

restriction site. The ~2600-bp fragment upstream of the piggy BAC insertion site contained AarI

sites and was obtained using these primers: Forward,

tataCACCTGCattaCCGAcgctagtggctcctagagttcag; Reverse,

gcagCACCTGCgcggTTAAtgaaaaagaatttataagtcttaccattaggttatc. The ~1000-bp fragment

downstream of the piggy BAC insertion site contained SapI sites and was obtained using these

primers: Forward, gccgGCTCTTCNTAAccttttttctacaaaatgatttatt; Reverse,

gtatGCTCTTCNCGGtcatcaaaacatagtttagtatcg.

To insert these fragments into the piggy BAC plasmid, the plasmid and the downstream

SapI-containing PCR was digested with SapI (also called LguI from ThermoFisher ER1931),

phosphatase treated (Antarctic phosphatase, NEB M0289S), and ligated together using T4

ligase (NEB M0202S). The upstream AarI-containing PCR fragment was TOPO cloned using

the Zero Blunt TOPO kit (ThermoFisher 451245). Then, TOPO AarI plasmid was cut using the

restriction enzyme HindIII (NEB R0104S), which removes a 708-bp fragment from *c(3)G*. The

cut TOPO AarI plasmid was then phosphatase treated and ligated back together to create the

c(3)G^{ccΔ1} deletion. Then, this plasmid was digested with AarI (ThermoFisher ER1581) to

generate a ~1900-bp fragment containing the *c(3)G^{ccΔ1}* deletion, which was cloned into the

piggy BAC plasmid containing the downstream SapI *c(3)G* fragment.

A CRISPR target sequence was selected from the flyCRISPR Optimal Target Finder

(http://tools.flycrispr.molbio.wisc.edu/targetFinder/). Only a single site upstream of the *c(3)G^{ccΔ1}*

deletion was selected (AAAGCTTTGTTGGCCTGTATTGG) and constructed into the pU6-BbsI-

chiRNA guide RNA (gRNA) plasmid (Addgene 45946). Sense

(CTTCGAAAGCTTTGTTGGCCTCTAT) and antisense (AAACATAGAGGCCAACAAAGCTTTC)

oligonucleotides were ordered from IDT and cloned into the gRNA plasmid as described by the

flyCRISPR subcloning pU6-gRNA protocol (http://flycrispr.molbio.wisc.edu/protocols/gRNA).

Three single nucleotide polymorphisms (SNPs) were made in the CRISPR target sequence (the

mutated bases are shown in bold: ccaataga**a**gc**g**aat**a**aaagcttt) in the *c(3)G^{ccΔ1}* homologous repair

template plasmid to prevent Cas9 from cutting this plasmid. These SNPs were made using the

Quik Change II XL Site-Directed Mutagenesis Kit (Agilent Technology, 200521) and the reaction

was performed as described in the kit protocol. The gRNA and *c(3)G^{ccΔ1}* homologous repair

template plasmid were sent to Genetivision (Houston, Texas) for injection into *y m[VASA-Cas9-*

3xGFP]ZH-2A-3xRFP w^{1118}/FM7c flies (BLM 51323). Genetivision injected the gRNA plasmid at

250 ng/µl and the *c(3)G^{ccΔ1}* homologous repair template at 500 ng/µl.

c(3)G^{ccΔ1} was isolated by crossing the G0 injected flies to *y w; Pr/TM3; sv^{spa-pol}*, then the

F1 progeny were screened for expression of dsRed in the fly eyes. Unfortunately, the genomic

locus marking the *VASA-Cas9* transgene is marked with RFP, so only F1 males were screened

for dsRed expression. PCR and Sanger sequencing were used to confirm the *c(3)G^{ccΔ1}* deletion

and a transposase (BLM 32073) was crossed in to excise the transposon from the gene. Those

flies that successfully excised the transposon were PCR and Sanger sequenced to confirm the

absence of the transposon and presence of the $c(3)G^{cc\Delta1}$ deletion. Then, the $c(3)G^{cc\Delta1}$

chromosome was allowed to freely recombine with a multiply marked third chromosome to

remove any potential off-target mutations caused by removal of the piggy Bac and/or Cas9. Two

$c(3)G^{cc\Delta1}$ fly stocks were established and used for all the assays in this manuscript: (1) carrying

all the markers from *3L* [*roughoid (ru), hairy (h), thread (th), scarlet (st)]* and two markers from

3R [curled (cu), claret (ca)] in addition to the $c(3)G^{cc\Delta1}$ deletion*,* and (2) only carrying *ca* with the

$c(3)G^{cc\Delta1}$ deletion.

Method details

Immunohistochemistry for DeltaVision and STORM microscopy

Ovary fixation and immunofluorescence for DeltaVision imaging were performed as detailed in

Lake et al. 2015 [29]. Briefly, ovaries were dissected in PBS with 10% Tween (PBST) for 15 min

and fixed in a 2% formaldehyde solution with heptane [165 µl PBS, 10 µl nonidet-P40 (Sigma-

Aldrich, 11332473001), 600 µl heptane and 25 µl of a 16% formaldehyde (Electron Microscopy

Sciences, 15710)] for 20 min. Then, ovaries were washed 3 times for 10 min each with PBST.

For experiments when only the early stages of oogenesis (stages 1–6) were analyzed, the

ovaries were clipped to remove the later stages of oogenesis. However, for the experiments

analyzing stages 7–9, the ovaries were rapidly pipetted to separate out the stages. Next,

ovaries were blocked in PBST+0.1% bovine serum albumin (BSA) for at least 1 hr, the block

was removed, and fresh PBST was added with the primary antibodies and left rocking overnight

at 4ºC. Primary antibodies used in this manuscript include mouse anti-C(3)G C-terminus (1A8–

1G2; used at 1:500; [23]), affinity-purified rabbit anti-Corolla (used a 1:2000; [24]), rat anti-CID

(used at 1:3000; from Claudio Sunkel), and mouse anti-γH2AV (used at 1:500; Iowa Hybridoma

Bank, UNC93-5.2.1; [30]). The next day, the primary antibodies were removed and the ovaries

were washed 3 times for 15 min each with PBST. Then, secondary antibodies were added in

PBST and incubated for 2 hr, rocking at room temperature. All secondary antibodies were used

at 1:500, and the secondary antibodies used were Alexa Fluor 488 goat anti-mouse

(ThermoFisher, A11001), Alexa Fluor 555 goat anti-mouse (ThermoFisher, A21422), Alexa

Fluor 647 goat anti-mouse (ThermoFisher, A21235), Alexa Fluor 488 goat anti-rabbit

(ThermoFisher, A11008), Alexa Fluor 555 goat anti-rabbit (ThermoFisher, A21428), Alexa Fluor

647 goat anti-rat (ThermoFisher, A21434), and Alexa Fluor 555 goat anti-rat (ThermoFisher,

A21247). After 1 hr and 45 min, 5 µl of 100X 4'6-diamididino-2-phenylindole (DAPI) was added

and left to incubate with the secondaries for 15 min. Ovaries were then washed with PBST 3

times for 15 min and mounted in Prolong Gold (Life Technologies, P36930) using an NA 1.5

glass coverslip and frosted glass slides.

For STORM of C(3)G, ovary fixation and immunofluorescence was performed as

described above with minor changes. The primary antibody was a mouse monoclonal anti-

C(3)G C-terminal (1A8–1G2, used at 1:500, [23]) and the secondary antibody was Alexa Fluor

647 mouse (used at 1:500; ThermoFisher, A21235). Since samples had to be imaged the same

day they were mounted, secondary antibody incubation was left rocking on the nutator overnight

at 4ºC. Then, the sample was washed 2 times in PBST for 15 min each. No DAPI was added to

these samples. Following the washes, the sample was optically cleared using 2,2-thiodiethanol

(TDE; VWR, 700008-210). The sample was rocked on the nutator at room temperature for 10

min each in 10% TDE in PBS, followed by 20% TDE in PBS, and finally in 50% TDE in PBS. For

mounting, a mixture of TDE and VECTASHIELD (VWR, 101098-042) was used since it had

been shown that low concentrations of VECTASHIELD allowed for blinking of the Alexa Fluor

647 dye [52]. We found that for WT samples, a mixture of 91%TDE/9% VECTASHIELD resulted

in robust blinking of the dye and higher VECTASHIELD concentrations stabilized the dye,

preventing it from blinking. In $c(3)G^{cc\Delta1}$ homozygotes and heterozygotes, we had to increase the

concentration of VECTASHIELD to obtain the same robust blinking as WT, thus these samples

were mounted in 88%TDE/12% VECTASHIELD. All samples were mounted with NA1.5

coverslips on glass slides and sealed using clear nail polish for STORM imaging that same day.

Fluorescent BAC probes for fluorescent in situ hybridization (FISH)

FISH probes were designed from bacterial artificial chromosomes (BACs) obtained from the

Children's Hospital Oakland Research Institute (CHORI;

http://bacpacresources.org/library.php?id=30). The following BACs were used: for *2L* RP98-

28O9 (polytene band 22A2-22A4), RP98-43K24 (polytene band 32E2-32F2), RP98-7D17

(polytene band 38E4-38F4); for *3L* RP98-2N23 (polytene band 61D-61E), RP98-26C20

(polytene band 69B1-69C2), RP98-3J2 (polytene band 77F5-78B1); for the *X* RP98-3D13

(polytene band 3C3-3C7), RP98-9H1 (polytene band 15C1-15D6).

Bacteria containing each BAC were grown on LB+chloramphenicol and the BACs were

isolated using a modification of the QIAGEN Plasmid Midi Kit protocol (QIAGEN, 12143). A

single bacteria colony was grown in 5 mL LB+chloramphenicol for 6–8 hours, then that culture

was moved into 100 mL culture of LB+chloramphenicol and grown overnight. Next, cells were

spun down in two 50-mL bottles at 4500 g for 20 min at 4^{o}C and the supernatant was removed

(Note: Can stop here and freeze cells at 80^{o}C.) The pellets were resuspended in 10 mL Buffer

P1 followed by adding 10 mL Buffer P2 to each bottle and gently mixed by inverting 4–6 times.

Then, the tubes were incubated at room temp for 5 min. Chilled Buffer P3, 10 ml, was added to

each tube and immediately mixed by gently inverting 4–6 times. The tubes were incubated on

ice for 15 min. The lysed cells were centrifuged at ≥ 20,000 g for 30 min at 4^{o}C (JA-20 rotor) and

the supernatant was moved to a new tube followed by centrifugation at ≥ 20,000g for 15 min at

4^{o}C (JA-20 rotor). The QIAGEN-tip 100 were equilibrated by applying 4 mL Buffer QBT and

allowed to flow through the column by gravity. The two supernatants were pooled together and

added to the QIAGEN-tip column allowing gravity flow to run the liquid through the column. The

QIAGEN-tip column was washed twice with 10 mL Buffer QC and the BACs were eluted off the

column with 5 mL Buffer QF pre-warmed to 65^{o}C. Following elution, the DNA was precipitated

by adding 3.5 mL isopropanol, mixed well, and centrifuged at ≥ 15,000g for 30 min at 4^{o}C (JA-

20 rotor). The DNA pellet was washed with 2 mL 70% ethanol and centrifuged at ≥ 15,000g for

10 min, then allowed to air dry for 5–10 min. The DNA was dissolved in 50 µl of water, and

additional water may need to be added to solubilize all the DNA.

To make the FISH probes, the BACs were PCR amplified using the Illustra GenomiPhi

V2 DNA Amplification Kit (GE 25-6600-30). The concentration of the BAC DNA was determined

using the Quibit and 10 ng of BAC DNA was used for one amplification reaction. The

amplification reaction was performed as described in the kit protocol. To confirm that the

amplification reaction worked, each sample was run on a DNA agarose gel to display a bright

smear ranging from ≥10 kb–3 kb DNA sized fragments. Also, the Qubit dsDNA HS assay kit

protocol (ThermoFisher, Q32851) was used to confirm that each amplification reaction had ~4–

7 ng of DNA.

Next, the amplified BAC was restriction enzyme digested using AluI (NEB R137S), HaeII

(NEB R107S), MseI (NEB R0525S), RsaI (NEB R0167S), MboI (NEB R0147S) and MspI (NEB

R0106S). A 100-µl digestion reaction was set up adding in all 20 µl of the amplified BAC DNA,

64.5 µl of the New England BioLabs smart cut buffer, and 1 µl of each restriction enzyme (for

Msp1, only 0.5 µl was added). The digestion reaction was left overnight at 37°C in a

thermocycler. Complete digestion of the BAC DNA was checked on a DNA agarose gel where

most of the DNA fragments should be below the 500-bp DNA size marker. Following the

digestion, the DNA was precipitated using 2.5 volumes 100% ethanol, 1/10 volume 3M sodium

acetate, and 1 µl of 20 mg/mL glycogen (ThermoFisher, 10814010). The precipitated DNA was

washed using 70% ethanol and resuspended in 50–60 µl of the labeling buffer from the ULYSIS

Nucleic Acid Labeling Kits (ThermoFisher – AF647 kit, U21660; AF546 kit, U21652).

To label the DNA with AF647 or AF546, the ULYSIS Nucleic Acid Labeling Kits were

used. For labeling, we used 10 µl of the digested BAC DNA and followed the protocol provided

in the ULYSIS kits. We removed the unreacted dyes from the labeling reaction using Centri-Sep

Columns (Princeton Separation, CS-900). The protocol provided with the columns was used to

remove the unreacted dyes, thereby isolating the fluorescently labeled BAC probes. The

resulting labeled BAC DNA was stored at –20ºC.

Fluorescent in situ hybridization of ovaries

Ovaries were dissected in PBST and fixed in a 3.7% formaldehyde and sodium cacodylate

solution. A 2X fix buffer (200 mM sodium cacodylate pH7.2, 200 mM sucrose, 80 mM sodium

acetate, 20 mM EGTA) was made in advance and stored in 500-µl aliquots at –20ºC. The 1X fix

solution (500 µl 2X Fix buffer, 232 µl 16% formaldehyde, 268 µl sterile water) was made

immediately prior to dissecting the ovaries and each ovary sample was fixed in 500 µl of the 1X

fix solution. Ovaries were fixed for 4 min rocking on the nutator at room temperature. Then, the

ovaries were rinsed three times in 2X Saline Sodium Citrate buffer with 0.1% Tween-20 (SCCT)

for 10 min each. Next, the ovary tips were clipped off using forceps to help isolate the

germarium from the later stages of oogenesis. These tips were transferred into new thin-walled

0.5-mL tubes and stepped into 2X SCCT with 50% formamide by sequentially adding each

formamide solution for 10 min each while nutating. First, the tips were incubated in 500 µl of a

2X SCCT with 20% formamide, then 2X SCCT with 40% formamide, and finally 2X SCCT with

50% formamide. The previous 2X SCCT with 50% formamide was removed and replaced with

fresh 2X SCCT with 50% formamide that was then incubated in a thermocycler at 37ºC for 1–2

hr.

Following that incubation, the 2X SCCT with 50% formamide was removed and the

hybridization mixture was added to the ovary tips. For all the BAC probes, the hybridization

mixture contained 2 µl BAC probe labeled with AF647, 2 µl BAC probe labeled with AF546, and

36 µl 1.1X hybridization solution (1.0 g dextran sulfate, 1.5 mL 20X SSC, 5 mL formamide,

water up to 9 ml; made in advance and stored at 4ºC). In a thermocycler, the probes and

chromosomal DNA were denatured for 2 min at 91ºC, then hybridized overnight at 37ºC.

The following day 500 µl 2X SCCT with 50% formamide was added to the hybridization

mixture and left at 37ºC to settle. Then, the solution was removed and the ovaries were washed

with 2X SCCT with 50% formamide three times at 37ºC for 20 min each. Next, the ovary tips

were exchanged back into 2X SCCT without formamide while nutating at room temperature by

adding 2X SCCT with 40% formamide for 10 min and then adding 2X SCCT with 20%

formamide for 10 min. The ovaries were washed once with 2X SCCT for 10 min, then washed

two more times with PBST for 10 min each. Then, block [1% Fraction V BSA in PBST] was

added and ovary tips were incubated for 1 hr at room temperature on the nutator. The block was

removed and primary antibodies were added in PBST and incubated overnight at 4ºC on the

nutator. For all FISH samples, a cocktail of three mouse monoclonal anti-C(3)G C-terminal

antibodies (1A8–1G2, 5G4–1F1, 1G5–2F7; [23]) was added and each antibody was used at a

dilution of 1:500.

Ovary tips were washed three times for 20 min each in PBST and the Alexa Fluor 488

mouse secondary was added at a dilution of 1:500. The secondary antibody was incubated for 2

hr rocking at room temperature. Then, 5 µl of 100X DAPI was added and incubated for an

additional 10 min. Next, the ovary tips were washed three times for 20 min each in PBST and

mounted onto glass slides with NA1.5 coverslips in Prolong Gold. After 24 hr the slides were

sealed using clear nail polish and kept at 4ºC prior to imaging on the DeltaVison.

DeltaVision and STORM microscopy

Except for the STORM imaging (see below), all images were acquired on an inverted

DeltaVision microscopy system (GE Healthcare) with an Olympus 100x Objective (UPlanSApo

100x NA 1.40) and a high-resolution CCD camera. The images from the DeltaVision were

deconvolved using SoftWorRx v. 6.5 software following the GE healthcare protocol. Images

were cropped and brightness and contrast was slightly adjusted using ImageJ.

STORM Imaging was performed on an OMX V4 microscopy system (GE healthcare)

with an Olympus 60x TIRF Objective (APO N 60x NA 1.49). Alexa Fluor 647 was continuously

excited with a 642 nm laser without UV activation, and the emission photons were filtered using

a band-pass emission filter (679/41, Semrock) and collected by a PCO-Edge sCMOS camera

with an 80-nm pixel size. Laser intensity measured after the objective was 4–6 kW/cm-2.

16,000–20,000 frames were recorded for each STORM data with 12–15 ms exposure time

depending on signal intensity.

Meiotic recombination

The multiply-marked *X* chromosome (*y sc cv v f y$^+$*) used in all the *X* recombination assays

carries two versions of the yellow gene: (1) a mutant version of the yellow gene at the genomic

locus on the tip of the long arm of the *X* chromosome (represented as *y*) and (2) a wild type

copy of the yellow gene that was integrated onto the short arm of the *X* chromosome on other

side of the centromere (represented as *y$^+$*). To assay the frequency of recombination on the *X*

chromosome in *c(3)G^{ccΔ1}* homozygotes, males of the genotype *w$^+$/BsY; mm c(3)G^{ccΔ1}/TM3;*

sv$^{spa-pol}$/+ were crossed to *FM7w/y sc cv v f y$^+$; D/TM3* virgin females. The *y sc cv v f y$^+$/BsY; mm*

c(3)G^{ccΔ1}/+; sv$^{spa-pol}$/+ male progeny from this cross were then crossed to *w$^+$/y w; mm*

c(3)G^{ccΔ1}/TM3; sv$^{spa-pol}$/+ virgin females, which generated females carrying *y sc cv v f y$^+$/y w; mm*

c(3)G^{ccΔ1}; sv$^{spa-pol}$/+. These females were crossed individually to *y sc cv v f y$^+$/BsY* males and the

female progeny were scored for the presence of the *sc, cv, v, f* and *y$^+$* markers to determine the

X recombination frequency. *c(3)G^{ccΔ1}* heterozygotes were generated by crossing males of the

genotype *y w/BsY; mm c(3)G^{ccΔ1}/TM3; sv^{spa-pol}/+* to *FM7w/y sc cv v f y^{+}* virgin females. The

resulting female progeny of the genotype *y sc cv v f y^{+}/y w; mm c(3)G^{ccΔ1}/+; sv^{spa-pol}/+* were

individually crossed to *y sc cv v f y^{+}/BsY* males. The *X* recombination frequency was determined

by scoring the female progeny for the presence of *sc, cv, v, f* and *y^{+}* markers. To assay *X*

recombination in WT, *y w; sv^{spa-pol}* virgin females were crossed to *y sc cv v f y^{+}/BsY* males. The

virgin female progeny from this cross were individually crossed to *y sc cv v f y^{+}/BsY* males. The

X recombination frequency was determined by scoring the female progeny for the presence of

sc, cv, v, f and *y^{+}* markers.

To assay recombination frequency along the 2^{nd} chromosome in *c(3)G^{ccΔ1}* homozygotes,

virgin females of the genotype *y w; +/+; c(3)G^{ccΔ1} ca; sv^{spa-pol}* were crossed to *w/y; net dpp dpy b*

pr cn/cyo; mm c(3)G^{ccΔ1}/TM3; sv^{spa-pol}/+ males. Single virgin female progeny of the genotype

yw/w; net dpp dpy b pr cn/+; mm c(3)G^{ccΔ1}/ c(3)G^{ccΔ1} ca; sv^{spa-pol}/+ were crossed to *w^{+}/y; net dpp*

dpy b pr cn males. The recombination frequency was determined by scoring the female progeny

for the presence of *net, dpp, dpy, b, pr,* and *cn* markers. To assay recombination frequency

along the 2^{nd} chromosome in *c(3)G^{ccΔ1}* heterozygotes, virgin WT females were crossed to *w/y;*

net dpp dpy b pr cn/cyo; mm c(3)G^{ccΔ1}/TM3; sv^{spa-pol}/+ males. Single virgin female progeny of the

genotype *yw/w; net dpp dpy b pr cn/+; mm c(3)G^{ccΔ1}/+; sv^{spa-pol}/+* were crossed to *w^{+}/y; net dpp*

dpy b pr cn males. The recombination frequency was determined by scoring the female progeny

for the presence of *net, dpp, dpy, b, pr,* and *cn* markers. Lastly, to assay recombination

frequency along the 2^{nd} chromosome in wild type, virgin WT females (*y w; sv^{spa-pol}*) were crossed

to *w^{+}/y; net dpp dpy b pr cn* males. Single virgin female progeny of this cross were then crossed

to *w^{+}/y; net dpp dpy b pr cn* males. The 2^{nd} recombination frequency was determined by scoring

the female progeny for the presence of *net, dpp, dpy, b, pr,* and *cn* markers.

To assay recombination frequency along the 3^{rd} chromosome in $c(3)G^{ccΔ1}$ homozygotes,

virgin females of the genotype $y\ w/w^{+}$; $mm\ c(3)G^{ccΔ1}/TM3$; $sv^{spa-pol}/+$ were crossed to $y\ w$;

$c(3)G^{ccΔ1}\ ca$; $sv^{spa-pol}$ males. Single virgin female progeny of this cross that were homozygous for

the $c(3)G^{ccΔ1}$ deletion were then crossed to $ru\ h\ th\ st\ cu\ sr\ e\ ca$ males. The recombination

frequency was determined by scoring for the presence of $ru, h, th, st,$ and cu in female progeny.

To assay recombination frequency along the 3^{rd} chromosome in $c(3)G^{ccΔ1}$ heterozygotes, virgin

wild type females were crossed to $y\ w/w^{+}$; $mm\ c(3)G^{ccΔ1}$; $sv^{spa-pol}/+$ males. Single virgin female

progeny of this cross were then crossed to $ru\ h\ th\ st\ cu\ sr\ e\ ca$ males. The recombination

frequency was determined by scoring for the presence of $ru, h, th, st,$ and cu in female progeny.

Lastly, to assay recombination frequency along the 3^{rd} chromosome in WT, virgin WT females (y

w; $sv^{spa-pol}$) were crossed to $ru\ h\ th\ st\ cu\ sr\ e\ ca$ males. Single virgin female progeny of this

cross were then crossed to $ru\ h\ th\ st\ cu\ sr\ e\ ca$ males. The recombination frequency was

determined by scoring for the presence of $ru, h, th, st,$ and cu in female progeny.

Meiotic nondisjunction

For the X and 4^{th} chromosome nondisjunction assay, we introduced $y\ w$ marked X

chromosomes and $sv^{spa-pol}$ marked 4^{th} chromosomes into the $c(3)G^{ccΔ1}\ ca/TM3$ stock to create a

$y\ w/y^{+}Y$; $c(3)G^{ccΔ1}/TM3$; $sv^{spa-pol}$ stock. This was done by crossing males from $y\ w/y^{+}Y$; $Pr/TM3$;

$sv^{spa-pol}$ to the $c(3)G^{ccΔ1}/TM3$ stock. All nondisjunction assays were performed using the $y\ w/y^{+}Y$;

$c(3)G^{ccΔ1}/TM3$; $sv^{spa-pol}$ stock. X and 4^{th} chromosome nondisjunction was assayed by crossing

virgin $y\ w$; $c(3)G^{ccΔ1}$; $sv^{spa-pol}$ and $y\ w$; $c(3)G^{ccΔ1}/+$; $sv^{spa-pol}$ females to $attached-XY$, $y^{+}\ v\ f\ B$;

$C(4)RM,\ ci\ ey^{R}$ males, as described in Harris et al. 2003 (the $attached-XY$ symbol represents the

chromosome $C(1;Y), IN(1)EN$, which is arranged as such $Y^{S}X \cdot Y^{L}$) [53]. Virgin homozygous

females for $c(3)G^{ccΔ1}$ were collected from the $y\ w/y^{+}Y$; $c(3)G^{ccΔ1}/TM3$; $sv^{spa-pol}$ stock. Virgin

heterozygous $c(3)G^{ccΔ1}$ females were created by crossing $y\ w/\ y^{+}Y$; $c(3)G^{ccΔ1}/TM3$; $sv^{spa-pol}$

males to *y w; sv^spa-pol* virgin females and collecting the virgin female progeny that displayed the

correct genotype (*y w; c(3)G^ccΔ1/+; sv^spa-pol*).

In this assay, *X* chromosome nondisjunction at meiosis I will result in yellow white (*y w*)

females, which are created by having the two maternal *X* chromosomes segregate to the same

pole (also known as a diplo-*X* ova), or yellow-plus vermillion forked Bar (*y⁺ v f Bar*) males, which

are created by the segregation of none of the maternal *X* chromosomes into the egg, resulting in

an *attached-XY/0* male (also known as a nullo-*X* ova). Meiosis II nondisjunction is extremely

rare and using this assay as described above, the meiosis II nondisjunctional progeny will

phenocopy the meiosis I nondisjunctional progeny. Since *c(3)G^ccΔ1* mutants display no

nondisjunction, we did not assay for meiosis II nondisjunction.

4^th chromosome nondisjunction at meiosis I will result in sparkling poliert (*sv^spa-pol*)

progeny, which are created by having the two maternal *4^th* chromosomes segregate to the same

pole (also known as a diplo-*4* ova), or cubitus interruptus eyeless Russian (*ci ey*) progeny,

which are created by the segregation of none of the maternal *4^th* chromosomes into the egg,

resulting in a *C(4)RM/0* progeny (also known as a nullo-*4* ova). While rare, flies with four *4^th*

chromosomes are viable and indistinguishable from regular progeny.

Quantification and statistical analysis

γH2AV, CID, and SC intensity quantifications

Oocyte staging for the γH2AV and CID quantifications was done according to Matthies et al.

[54]. Briefly, the germarium was identified by DAPI staining and each region was defined by the

pattern of SC staining. Region 2A contains at least 4 cells of the 16-cell cyst with either partially

or fully assembled SC, and in region 2B only 2 cells contain full-length SC. Region 3 (stage 1) is

at the base of the germarium where only 1 cell has full-length SC and the other cell has either

fully disassembled or is in the process of disassembling the SC. The rest of the stages (stage

2–9) were defined by measuring the size of the oocyte. Stage 2 is 25x25 µm, stage 3 is 35x35

µm, stage 4 is 40x50 µm, stage 5 is 55x75 µm, stage 6 is 60x85 µm, and stage 7 is 70x115 µm.

Each oocyte was measured using the SoftWoRx software, then classified as the stage to which

it was the closest in size. For stages 8 and 9 there is no quantitative way to measure the egg to

determine these stages. Instead, stage 8 is defined by being bigger than a stage 7 with the

oocyte cytoplasm occupying less than one-third of the egg. Stage 9 is defined by the oocyte

cytoplasm occupying one-third of the egg and the follicle cells beginning to migrate over the

oocyte.

Prior to the quantification of γH2AV and CID foci, individual nuclei from each region and

stage were 3D-cropped in ImageJ (NIH open source software). The number of foci present for

both γH2AV and CID foci was determined by analyzing each cropped nucleus in 3D and

counting the number of foci. N value for both γH2AV and CID data sets represents the number

of nuclei scored in each stage. In the germarium, more nuclei can be scored in region 2A than in

region 2B and region 3, thus the number of germaria analyzed varies accordingly, such that the

total nuclei scored in each region was 30. The analysis of region 2A came from 4–7 individual

germaria. The analysis of region 2B came from 13–19 individual germarium. The analysis of

region 3 came from 30 individual germaria. A Wilcoxon rank sum test was used for the P-value

calculation of the γH2AV data.

To estimate protein density at different stages for the mutants acquired here, we

performed quantitative imaging of the SC antibody Corolla labeled with Alexa Fluor 488. For this

analysis, we assumed that the antibody penetration and epitope availability was the same

between each of the regions/stages and for each mutant analyzed. SC intensity measurements

were obtained by performing sum projections over SC regions and then measuring the

integrated intensity of the sum projected region after subtracting the average intensity from a

manually selected region near each individual SC. In some cases, the exposure time and laser

power were changed in order to obtain reasonable images of low-intensity SCs. In those cases,

the intensity values were corrected to account for these changes.

Euchromatic FISH pairing quantification

The same staging of the germarium as described above was done for the quantification of the

euchromatic FISH pairing. To measure the distance between the FISH probe foci, a custom

ImageJ plug-in was used. First, the two FISH probe foci were selected as points using the

points tool and the coordinates of each point were recorded in the ROI manager. Then, the

plugin measure "3D jru v1" was run to calculate the 3D distance (in μm) between ROI point one

and ROI point two (available at http://research.stowers.org/imagejplugins). A slice spacing of

0.20 and pixel spacing of 0.06370 were used in the plugin to calculate the 3D distance between

the foci. A locus was considered paired if the distance between the FISH probe foci was <0.75

μm and unpaired if the distanced between the FISH probe foci was ≥0.75 μm. The n value for

this data is the number of nuclei scored. Similar to the γH2AV and CID quantification, the

number of germaria scored for region 2A, region 2B, and region 3 vary because the number of

quantifiable nuclei in each region varies. For all the probes assayed, on average ~4 germaria

were scored for region 2A, ~8 germaria were scored for region 2B, and ~13 germaria were

scored for region 3.

STORM quantification

STORM data was analyzed through an open source ImageJ plug-in [55]. Briefly, the acquired

image was smoothed by a Gaussian filter, and then each blinking molecule was fitted to a

Gaussian function by maximum likelihood fitting. Individual molecule's position, photon counts

and fitting standard deviation were stored for future processing. To correct and optimize the

results of the analysis, only molecules having 1–16 nm localization precision and 80–160 nm

fitting standard deviation were chosen for rendering and quantification. Cross-correlation

between subsets (~2000 frames) of localized molecules was calculated to correct lateral drift.

Measurements were performed using ImageJ and the following custom plugins:

fit_multi_gaussian_jru_v4; set_multi_plot_offsets_jru_v1; resample_plot_jru_v1;

normalize_trajectories_jru_v1; average_trajectories_jru_v1 (available at

http://research.stowers.org/imagejplugins). Measurements were performed on raw molecule

position accumulation images. Intensity (i.e. molecule position accumulation) profiles were then

averaged over a five-pixel-wide stripe perpendicular to the paired C(3)G C-terminal tracks.

Profile fits to double Gaussian functions were used to determine the centers of each profile.

Profiles were then realigned to these centers, normalized, and averaged to obtain the final

averaged histograms as described in [21]. This averaged profile was then fit to a double

Gaussian function, by nonlinear least squares as described in [56], using the open source tools

described above. Error analysis of the fit parameters was performed using a Monte Carlo

approach with 100 random simulations (also described in [56]). For wild type, n=26 SC

measurements; for $c(3)G^{cc\Delta 1}$ heterozygotes, n = 34 SC measurements; and for $c(3)G^{cc\Delta 1}$

homozygotes, n = 32 SC measurements. Standard t-test was used for statistical comparisons

between the $c(3)G^{cc\Delta 1}$ mutants and WT.

Meiotic recombination quantification

N values for the recombination assay refer to the number of progeny scored in each genotype.

For all the chromosomes assayed, map length was determined by dividing the number of

progeny that had a crossover between a given set of markers by the total number of progeny

scored for one genotype on a specific chromosome. For example, the map length from *sc-cv*

was determined by dividing the number of progeny that had a crossover between *sc-cv* markers

by the total number of *X* chromosome recombinant progeny scored for a specific genotype.

Interference was calculated by: $1 - \dfrac{observed\ double\ crossovers\ between\ adjacent\ intervals}{expected\ double\ crossovers\ between\ adjacent\ intervals}$

The number of expected double crossovers occurring in adjacent intervals was determined from

the WT recombination data by multiplying the map length of one interval by the map length of

the adjacent interval and then multiplying by the total number of recombinants scored. For

example, on the *X* chromosome the expected number of double crossovers occurring between

the *sc-cv* and *cv-v* intervals was calculated by:

$\dfrac{sc-cv\ recombinants}{total\ X\ recombinants} \times \dfrac{cv-v\ recombinants}{total\ X\ recombinants} \times total\ X\ recombinants$

The number of observed double crossovers between adjacent intervals was empirically

determined from the raw data. The total number of noncrossovers (NCOs), single crossovers

(SCOs), double crossovers (DCOs), triple crossovers (TCOs) and quad crossovers (QCOs)

were empirically determined from the raw data. Exchange rank was determined according to the

method in Weinstein [36, 57].

Meiotic nondisjunction quantification

For both the *X* and *4th* chromosomes, half of the progeny from the nondisjunction assay are

inviable. Diplo-*X* ova are only viable when fertilized with sperm not carrying the *attached-XY*. If

these ova are fertilized with sperm carrying the *attached-XY,* the resulting progeny will be

XX/attached-XY, which is inviable. Nullo-*X* ova are only viable when fertilized with sperm

carrying the *attached-XY*. If these ova are fertilized with sperm not carrying the *attached-XY*, the

resulting progeny will not have any sex chromosomes, which is inviable. Since only half of the

nondisjunctional progeny are recovered, we adjusted the total by doubling the number of *X*

chromosome nondisjuctional progeny.

Nullo-*4* flies, from nullo-*4* ova fertilized by nullo-*4* sperm, are not able to survive.

Additionally, haplo-*4 minute* flies (or a single *4/0*) will occur as regular progeny from the cross,

but these flies are very poor and recorded but not scored in the nondisjunction tables. If both the

X and *4th* chromosomes nondisjoin together (*X–4*) only two of the four possible progeny are

viable. Thus, we adjusted the total by doubling the number of the *X–4* nondisjunctional progeny.

For both *X* and *4,* the frequency of nondisjunction was calculated as described in Hawley et al.

1992 [40].

Data and software availability

Primary data files for the figures in this paper are publicly accessible

at www.stowers.org/research/publications/odr. For data analysis, the custom ImageJ plugins

used are available at research.stowers.org/imagejplugins/zipped_plugins.html.

Key resources table

See document.

Acknowledgments

We thank Claudio Sunkel for antibodies; past and present members of the Hawley lab for

helpful discussion and comments on this manuscript especially Satomi Takeo, Cathy Lake,

Rachel Nielsen, and Elisabeth Bauerly; Diana Libuda for comments on this manuscript; and

Angela Miller for editorial and figure preparation assistance. R.S.H. is an American Cancer

Society Research Professor.

References

1.   Hassold T, Hall H, Hunt P. The origin of human aneuploidy: where we have been, where
we are going. Human molecular genetics. 2007;16 Spec No. 2:R203-8. Epub 2007/10/04. doi:
10.1093/hmg/ddm243. PubMed PMID: 17911163.

2. Keeney S, Giroux CN, Kleckner N. Meiosis-specific DNA double-strand breaks are
catalyzed by Spo11, a member of a widely conserved protein family. Cell. 1997;88(3):375-84.
Epub 1997/02/07. PubMed PMID: 9039264.
3. Gray S, Cohen PE. Control of Meiotic Crossovers: From Double-Strand Break Formation
to Designation. Annual review of genetics. 2016;50:175-210. doi: 10.1146/annurev-genet-
120215-035111. PubMed PMID: 27648641; PubMed Central PMCID: PMCPMC5319444.
4. Hunter N. Meiotic Recombination: The Essence of Heredity. Cold Spring Harb Perspect
Biol. 2015. doi: 10.1101/cshperspect.a016618. PubMed PMID: 26511629.
5. Lange J, Yamada S, Tischfield SE, Pan J, Kim S, Zhu X, et al. The Landscape of Mouse
Meiotic Double-Strand Break Formation, Processing, and Repair. Cell. 2016;167(3):695-708
e16. doi: 10.1016/j.cell.2016.09.035. PubMed PMID: 27745971; PubMed Central PMCID:
PMCPMC5117687.
6. Pan J, Sasaki M, Kniewel R, Murakami H, Blitzblau HG, Tischfield SE, et al. A
hierarchical combination of factors shapes the genome-wide topography of yeast meiotic
recombination initiation. Cell. 2011;144(5):719-31. doi: 10.1016/j.cell.2011.02.009. PubMed
PMID: 21376234; PubMed Central PMCID: PMCPMC3063416.
7. Lake CM, Hawley RS. Becoming a crossover-competent DSB. Semin Cell Dev Biol.
2016;54:117-25. doi: 10.1016/j.semcdb.2016.01.008. PubMed PMID: 26806636.
8. Xiang Y, Miller DE, Ross EJ, Sanchez Alvarado A, Hawley RS. Synaptonemal complex
extension from clustered telomeres mediates full-length chromosome pairing in Schmidtea
mediterranea. Proc Natl Acad Sci U S A. 2014;111(48):E5159-68. doi:
10.1073/pnas.1420287111. PubMed PMID: 25404302; PubMed Central PMCID:
PMCPMC4260563.
9. Miller DE, Smith CB, Kazemi NY, Cockrell AJ, Arvanitakas AV, Blumenstiel JP, et al.
Whole-Genome Analysis of Individual Meiotic Events in Drosophila melanogaster Reveals That
Noncrossover Gene Conversions Are Insensitive to Interference and the Centromere Effect.
Genetics. 2016;203(1):159-71. doi: 10.1534/genetics.115.186486. PubMed PMID: 26944917;
PubMed Central PMCID: PMCPMC4858771.
10. Lindsley DL, Sandler L. The genetic analysis of meiosis in female Drosophila
melanogaster. Philos Trans R Soc Lond B Biol Sci. 1977;277(955):295-312. PubMed PMID:
16292.
11. Hughes SE, Miller DE, Miller AL, Hawley RS. Female meiosis: synapsis, recombination
and segregation in *Drosophila melanogaster*. Genetics. 2017;*in press*.
12. Tanneti NS, Landy K, Joyce EF, McKim KS. A pathway for synapsis initiation during
zygotene in Drosophila oocytes. Current biology : CB. 2011;21(21):1852-7. doi:
10.1016/j.cub.2011.10.005. PubMed PMID: 22036181.
13. Christophorou N, Rubin T, Huynh J-R. Synaptonemal Complex Components Promote
Centromere Pairing in Pre-Meiotic Germ Cells. PLoS Genet. 2013.
14. Takeo S, Lake CM, Morais-de-Sa E, Sunkel CE, Hawley RS. Synaptonemal complex-
dependent centromeric clustering and the initiation of synapsis in Drosophila oocytes. Current
biology : CB. 2011;21(21):1845-51. doi: 10.1016/j.cub.2011.09.044. PubMed PMID: 22036182.
15. McKim KS, Green-Marroquin BL, Sekelsky JJ, Chin G, Steinberg C, Khodosh R, et al.
Meiotic synapsis in the absence of recombination. Science. 1998;279(5352):876-8. PubMed
PMID: 9452390.
16. Mehrotra S, McKim KS. Temporal analysis of meiotic DNA double-strand break
formation and repair in Drosophila females. PLoS Genet. 2006;2(11):e200. doi:
10.1371/journal.pgen.0020200. PubMed PMID: 17166055; PubMed Central PMCID:
PMCPMC1657055.
17. Libuda DE, Uzawa S, Meyer BJ, Villeneuve AM. Meiotic chromosome structures
constrain and respond to designation of crossover sites. Nature. 2013;502(7473):703-6. doi:
10.1038/nature12577. PubMed PMID: 24107990; PubMed Central PMCID: PMCPMC3920622.

18.   Hayashi M, Mlynarczyk-Evans S, Villeneuve AM. The synaptonemal complex shapes the
crossover landscape through cooperative assembly, crossover promotion and crossover
inhibition during Caenorhabditis elegans meiosis. Genetics. 2010;186(1):45-58. doi:
10.1534/genetics.110.115501. PubMed PMID: 20592266; PubMed Central PMCID:
PMCPMC2940310.
19.   Zickler D, Kleckner N. Recombination, Pairing, and Synapsis of Homologs during
Meiosis. Cold Spring Harb Perspect Biol. 2015;7(6). doi: 10.1101/cshperspect.a016626.
PubMed PMID: 25986558; PubMed Central PMCID: PMCPMC4448610.
20.   Zickler D, Kleckner N. Meiotic chromosomes: integrating structure and function. Annual
review of genetics. 1999;33:603-754. doi: 10.1146/annurev.genet.33.1.603. PubMed PMID:
10690419.
21.   Cahoon CK, Yu Z, Wang Y, Guo F, Unruh JR, Slaughter BD, et al. Superresolution
expansion microscopy reveals the three-dimensional organization of the Drosophila
synaptonemal complex. Proc Natl Acad Sci U S A. 2017;114(33):E6857-E66. doi:
10.1073/pnas.1705623114. PubMed PMID: 28760978; PubMed Central PMCID:
PMCPMC5565445.
22.   Page SL, Hawley RS. c(3)G encodes a Drosophila synaptonemal complex protein.
Genes & development. 2001;15(23):3130-43. doi: 10.1101/gad.935001. PubMed PMID:
11731477; PubMed Central PMCID: PMC312841.
23.   Anderson LK, Royer SM, Page SL, McKim KS, Lai A, Lilly MA, et al. Juxtaposition of
C(2)M and the transverse filament protein C(3)G within the central region of Drosophila
synaptonemal complex. Proc Natl Acad Sci U S A. 2005;102(12):4482-7. doi:
10.1073/pnas.0500172102. PubMed PMID: 15767569; PubMed Central PMCID: PMC555515.
24.   Collins KA, Unruh JR, Slaughter BD, Yu Z, Lake CM, Nielsen RJ, et al. Corolla Is a
Novel Protein that Contributes to the Architecture of the Synaptonemal Complex of Drosophila.
Genetics. 2014. doi: 10.1534/genetics.114.165290. PubMed PMID: 24913682.
25.   Page SL, Khetani RS, Lake CM, Nielsen RJ, Jeffress JK, Warren WD, et al. Corona is
required for higher-order assembly of transverse filaments into full-length synaptonemal
complex in Drosophila oocytes. PLoS Genet. 2008;4(9):e1000194. doi:
10.1371/journal.pgen.1000194. PubMed PMID: 18802461; PubMed Central PMCID:
PMC2529403.
26.   Lake CM, Hawley RS. The molecular control of meiotic chromosomal behavior: events in
early meiotic prophase in Drosophila oocytes. Annu Rev Physiol. 2012;74:425-51. doi:
10.1146/annurev-physiol-020911-153342. PubMed PMID: 22335798.
27.   Jeffress JK, Page SL, Royer SK, Belden ED, Blumenstiel JP, Anderson LK, et al. The
formation of the central element of the synaptonemal complex may occur by multiple
mechanisms: the roles of the N- and C-terminal domains of the Drosophila C(3)G protein in
mediating synapsis and recombination. Genetics. 2007;177(4):2445-56. doi:
10.1534/genetics.107.078717. PubMed PMID: 17947423; PubMed Central PMCID:
PMC2219479.
28.   Tung KS, Roeder GS. Meiotic chromosome morphology and behavior in zip1 mutants of
Saccharomyces cerevisiae. Genetics. 1998;149(2):817-32. PubMed PMID: 9611194; PubMed
Central PMCID: PMC1460213.
29.   Lake CM, Nielsen RJ, Guo F, Unruh JR, Slaughter BD, Hawley RS. Vilya, a component
of the recombination nodule, is required for meiotic double-strand break formation in Drosophila.
eLife. 2015;4:e08287. doi: 10.7554/eLife.08287. PubMed PMID: 26452093; PubMed Central
PMCID: PMCPMC4703084.
30.   Lake CM, Holsclaw JK, Bellendir SP, Sekelsky J, Hawley RS. The development of a
monoclonal antibody recognizing the Drosophila melanogaster phosphorylated histone H2A
variant (gamma-H2AV). G3 (Bethesda). 2013;3(9):1539-43. doi: 10.1534/g3.113.006833.
PubMed PMID: 23833215; PubMed Central PMCID: PMCPMC3755914.

31.   Zickler D, Kleckner N. A few of our favorite things: Pairing, the bouquet, crossover
interference and evolution of meiosis. Semin Cell Dev Biol. 2016;54:135-48. doi:
10.1016/j.semcdb.2016.02.024. PubMed PMID: 26927691; PubMed Central PMCID:
PMCPMC4867269.
32.   Gowen JW. Meiosis as a genetic character in *Drosophila melanogaster*. The Journal of
Experiment Zoology. 1933;65(1):83-106.
33.   Hall JC. Chromosome segregation influenced by two alleles of the meiotic mutant c(3)G
in Drosophila melanogaster. Genetics. 1972;71(3):367-400. PubMed PMID: 4624918.
34.   Miller DE. Genomic analysis of meiosis in *Drosophila melanogaster* [dissertation].
Kansas City, KS: University of Kansas Medical Center; 2016.
35.   Ashburner M, Golic KG, Hawley RS. Drosophila: A laboratory handbook. 2 ed. Cold
Spring Harbor, New York: Cold Spring Harbor Laboratory Press; 2005.
36.   Weinstein A. Coincidence of Crossing over in DROSOPHILA MELANOGASTER
(AMPELOPHILA). Genetics. 1918;3(2):135-72. PubMed PMID: 17245901; PubMed Central
PMCID: PMCPMC1200433.
37.   Yan R, McKee BD. The cohesion protein SOLO associates with SMC1 and is required
for synapsis, recombination, homolog bias and cohesion and pairing of centromeres in
Drosophila Meiosis. PLoS Genet. 2013;9(7):e1003637. doi: 10.1371/journal.pgen.1003637.
PubMed PMID: 23874232; PubMed Central PMCID: PMC3715423.
38.   Mason JM. Orientation disruptor (ord): a recombination-defective and disjunction-
defective meiotic mutant in Drosophila melanogaster. Genetics. 1976;84(3):545-72. PubMed
PMID: 826453; PubMed Central PMCID: PMCPMC1213594.
39.   Baker BS, Hall JC. Meiotic mutants: genie control of meiotic recombination and
chromosome segregation. In: Ashburner M, Novitski E, editors. The Genetics and Biology of
Drosophila. 1a. New York: Academic Press; 1976.
40.   Hawley RS, Irick H, Zitron AE, Haddox DA, Lohe A, New C, et al. There are two
mechanisms of achiasmate segregation in Drosophila females, one of which requires
heterochromatic homology. Dev Genet. 1992;13(6):440-67. doi: 10.1002/dvg.1020130608.
PubMed PMID: 1304424.
41.   Hawley RS, Theurkauf WE. Requiem for distributive segregation: achiasmate
segregation in Drosophila females. Trends Genet. 1993;9(9):310-7. PubMed PMID: 8236460.
42.   Balicky EM, Endres MW, Lai C, Bickel SE. Meiotic cohesion requires accumulation of
ORD on chromosomes before condensation. Molecular biology of the cell. 2002;13(11):3890-
900. doi: 10.1091/mbc.E02-06-0332. PubMed PMID: 12429833; PubMed Central PMCID:
PMC133601.
43.   Khetani RS, Bickel SE. Regulation of meiotic cohesion and chromosome core
morphogenesis during pachytene in Drosophila oocytes. Journal of cell science. 2007;120(Pt
17):3123-37. doi: 10.1242/jcs.009977. PubMed PMID: 17698920.
44.   Krishnan B, Thomas SE, Yan R, Yamada H, Zhulin IB, McKee BD. Sisters unbound is
required for meiotic centromeric cohesion in Drosophila melanogaster. Genetics.
2014;198(3):947-65. doi: 10.1534/genetics.114.166009. PubMed PMID: 25194162; PubMed
Central PMCID: PMCPMC4224182.
45.   Yan R, Thomas SE, Tsai JH, Yamada Y, McKee BD. SOLO: a meiotic protein required
for centromere cohesion, coorientation, and SMC1 localization in Drosophila melanogaster. The
Journal of cell biology. 2010;188(3):335-49. doi: 10.1083/jcb.200904040. PubMed PMID:
20142422; PubMed Central PMCID: PMC2819681.
46.   Gladstone MN, Obeso D, Chuong H, Dawson DS. The synaptonemal complex protein
Zip1 promotes bi-orientation of centromeres at meiosis I. PLoS Genet. 2009;5(12):e1000771.
doi: 10.1371/journal.pgen.1000771. PubMed PMID: 20011112; PubMed Central PMCID:
PMC2781170.

47.   Kurdzo EL, Obeso D, Chuong H, Dawson DS. Meiotic Centromere Coupling and Pairing
Function by Two Separate Mechanisms in Saccharomyces cerevisiae. Genetics.
2017;205(2):657-71. doi: 10.1534/genetics.116.190264. PubMed PMID: 27913618; PubMed
Central PMCID: PMCPMC5289843.
48.   Qiao H, Chen JK, Reynolds A, Hoog C, Paddy M, Hunter N. Interplay between
synaptonemal complex, homologous recombination, and centromeres during mammalian
meiosis. PLoS Genet. 2012;8(6):e1002790. doi: 10.1371/journal.pgen.1002790. PubMed PMID:
22761591; PubMed Central PMCID: PMC3386176.
49.   Carpenter A. Thoughts on recombination nodules, meiotic recombination, and
chiasmata. In: Kucherlapati R, Smith GR, editors. Genetic Recombination. Washington, DC:
American Society for Microbiology; 1988. p. 529-48.
50.   Mlynarczyk-Evans S, Villeneuve AM. Time-Course Analysis of Early Meiotic Prophase
Events Informs Mechanisms of Homolog Pairing and Synapsis in Caenorhabditis elegans.
Genetics. 2017;207(1):103-14. doi: 10.1534/genetics.117.204172. PubMed PMID: 28710064;
PubMed Central PMCID: PMCPMC5586365.
51.   Jaramillo-Lambert A, Ellefson M, Villeneuve AM, Engebrecht J. Differential timing of S
phases, X chromosome replication, and meiotic prophase in the C. elegans germ line. Dev Biol.
2007;308(1):206-21. doi: 10.1016/j.ydbio.2007.05.019. PubMed PMID: 17599823.
52.   Olivier N, Keller D, Rajan VS, Gonczy P, Manley S. Simple buffers for 3D STORM
microscopy. Biomed Opt Express. 2013;4(6):885-99. doi: 10.1364/BOE.4.000885. PubMed
PMID: 23761850; PubMed Central PMCID: PMCPMC3675867.
53.   Harris D, Orme C, Kramer J, Namba L, Champion M, Palladino MJ, et al. A deficiency
screen of the major autosomes identifies a gene (matrimony) that is haplo-insufficient for
achiasmate segregation in Drosophila oocytes. Genetics. 2003;165(2):637-52. PubMed PMID:
14573476; PubMed Central PMCID: PMCPMC1462769.
54.   Matthies HJG, Clarkson M, Saint RB, Namba R, Hawley RS. Analysis of meiosis in fixed
and live oocytes by light microscopy. In: Sullivan W, Ashburner M, Hawley RS, editors.
Drosophila protocols. Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press;
2000. p. 67-86.
55.   Ovesny M, Krizek P, Borkovec J, Svindrych Z, Hagen GM. ThunderSTORM: a
comprehensive ImageJ plug-in for PALM and STORM data analysis and super-resolution
imaging. Bioinformatics. 2014;30(16):2389-90. doi: 10.1093/bioinformatics/btu202. PubMed
PMID: 24771516; PubMed Central PMCID: PMCPMC4207427.
56.   Burns S, Avena JS, Unruh JR, Yu Z, Smith SE, Slaughter BD, et al. Structured
illumination with particle averaging reveals novel roles for yeast centrosome components during
duplication. eLife. 2015;4. doi: 10.7554/eLife.08586. PubMed PMID: 26371506; PubMed Central
PMCID: PMCPMC4564689.
57.   Hawley RS, Walker MY. Advanced Genetic Analysis. Malden, MA: Blackwell Publishing;
2009.

1116

1117

1118

1119

Figure Legends

Figure 1: Background on Drosophila female meiosis and the SC

(A) Diagram of Drosophila female meiosis, which is described in [11]. Briefly, at the anterior tip

of the germarium is a germline stem cell that divides asymmetrically to give rise to a cystoblast,

which undergoes four mitotic divisions with incomplete cytokinesis to give rise to a 16-cell cyst.

Homologous chromosome pairing and SC assembly begin at the centromeres (represented as

black dots on the chromosomes) during these mitotic divisions [13]. Then, four cells in the 16-

cell cyst officially enter meiosis in region 2A (zygotene/early pachytene) by assembling the SC

along the chromosome arms. During region 2A, DNA double-strand breaks (DSBs) are formed

within the context of the SC, starting meiotic recombination. By region 2B (early pachytene), the

fully synapsed cells have started to repair their DSBs and two of the four cells have backed out

of the meiotic program. By region 3 (mid pachytene), the oocyte nucleus has been selected and

is the only nucleus within the cyst that retains the SC—all other nuclei have backed out of

meiosis and have disassembled their SC to become nurse cells. Additionally, all the DSBs are

in the processes of being repaired into crossovers (COs) or noncrossovers by region 3. After

the germarium, the oocyte continues to develop and the SC is maintained along the

chromosome arms until stage 5. From stages 5–7 (mid/late pachytene), the SC is disassembled

from multiple regions along the chromosome arms, but the SC persists at the centromere into

1139   the later stages of prophase I (stages8–9) [12, 14]. (B) Model of the Drosophila SC with the

1140   transverse filament protein C(3)G in blue, the central region (CR) protein Corolla in green, the

1141   central element protein (CE) CONA in brown, and the lateral element (LE)/cohesin proteins

1142   represented by gray (adapted from [11]). Next to the SC model are diagrams of the gene and

1143   protein structures for wild type C(3)G$^+$ (blue) and $c(3)G^{cc\Delta1}$ (orange).

1144

1145

Figure 2: STORM analysis of the SC structure in $c(3)G^{cc\Delta1}$ mutants

STORM images of intact germaria with the C-terminus of C(3)G labeled in wild type (WT, grey),

$c(3)G^{cc\Delta1}$ heterozygotes ($c(3)G^{cc\Delta1}$/+, blue) and homozygotes ($c(3)G^{cc\Delta1}$, orange). The

quantification above the images displays the average distribution of the two C-terminal C(3)G

tracks based on the line profile analysis of each genotype (see Methods). The quantification

resulted an average width of 119nm ± SE 1.0nm in WT, 107nm ± SE 0.4nm in $c(3)G^{cc\Delta1}$

heterozygotes, and 78.6nm ± SE 0.3nm in $c(3)G^{cc\Delta1}$ homozygotes. The average distribution was

generated by averaging 26 line profiles from 8 WT nuclei, 34 line profiles from 17 $c(3)G^{cc\Delta1}$

heterozygous nuclei, and 32 line profiles from 13 $c(3)G^{cc\Delta1}$ heterozygous nuclei.

Figure 3: Euchromatic SC prematurely disassembles in both *c(3)G^{ccΔ1}* homozygotes and heterozygotes

(A) Images showing localization of the SC protein Corolla (green), the centromeric nucleosome CID (red) and DAPI-stained DNA (blue) in wild type (A), *c(3)G^{ccΔ1}*/+ heterozygotes (B), and *c(3)G^{ccΔ1}* homozygotes (C) from zygotene/early pachytene (region 2A) to mid/late pachytene (stages 8–9). Dotted lines indicate the location of the nucleus as defined by DAPI staining (blue). Scale bars, 2 µm.

1163

1164

Figure 4: *c(3)G^{ccΔ1}* **causes changes in recombination and euchromatic pairing on the *X*,**
***2^{nd}* and *3^{rd}* chromosomes**

The frequency of recombination (A, C, E) and the fraction of euchromatic pairing (B, D, F) in

wild type (WT, grey triangles), $c(3)G^{cc\Delta 1}$ heterozygotes [$c(3)G^{cc\Delta 1}$/+, blue squares] and

homozygotes [$c(3)G^{cc\Delta 1}$, orange circles] at distal (yellow), medial (green) and proximal (brown)

loci on the *X*, *2^{nd}* and *3^{rd}* chromosomes. The markers used to assay recombination on the *X*

chromosome are *scute* (*sc*), *crossveinless* (*cv*), *vermillion* (*v*), *forked* (*f*), and *yellow^{+}* (*y^{+}*,

integrated wild type allele). (The multiply marked *X* chromosome also carries a null mutant of

the *yellow* gene at the genomic locus, see Methods.) The markers used to assay recombination

on the *2^{nd}* chromosome are *net, decapentaplegic* (*dpp*), *dumpy* (*dpy*), *black* (*b*), *purple* (*pr*), and

cinnabar (*cn*). The markers used to assay recombination on the *3^{rd}* chromosome are *roughoid*

(*ru*), *hairy* (*h*), *thread* (*th*), *scarlet* (*st*), and *curled* (*cu*). For reference, below each set of charts is

a diagram of the corresponding chromosome being analyzed displaying the relative positions of

the recombination markers, locations of each FISH probe and the approximate amounts of

pericentromeric heterochromatin estimated from [35] (the black circle represents the

centromere; the chromosome diagrams do not include telomeric heterochromatin). The genomic

positions of all the markers and pairing probes within the figure are in Table S2 (base pair

numbers from flybase release FB2017_06).

1183

1184

1185

1186

1187

1188

1189

1190

1191   **Supplemental Figure Legends**

1192

Figure S1: SC intensity quantification

Average intensity of the Corolla antibody staining for wild type (grey), $c(3)G^{cc\Delta 1}$ heterozygotes

($cc\Delta 1/+$, blue) and homozygotes ($cc\Delta 1$, orange) from zygotene/early pachytene (region 2A) to

mid/late pachytene (stage 8–9). (See Methods for the detailed description of how this analysis

was performed). For WT, the average intensity was generated by averaging measurements

from 19 nuclei in region 2A, 18 nuclei in region 2B, 12 nuclei in region 3, 49 nuclei from stages

2–4, 39 nuclei from stages 5–7, and 18 nuclei from stages 8–9. For $c(3)G^{cc\Delta 1}$ heterozygotes, the

average intensity was generated by averaging measurements from 16 nuclei in region 2A, 13

nuclei in region 2B, 19 nuclei in region 3, 46 nuclei from stages 2–4, 47 nuclei from stages 5–7,

and 34 nuclei from stages 8–9. For $c(3)G^{cc\Delta 1}$ homozygotes, the average intensity was generated

by averaging measurements from 18 nuclei in region 2A, 11 nuclei in region 2B,16 nuclei in

region 3, 42 nuclei from stages 2–4, 47 nuclei from stages 5–7, and 31 nuclei from stages 8–9.

1206

Figure S2: The formation of double-strand breaks is normal in *c(3)G^{ccΔ1}* mutants

(A) Images of the germarium stained with the SC protein Corolla (green), γH2AV (magenta) and

DAPI (blue) in WT and *c(3)G^{ccΔ1}* heterozygotes (*c(3)G^{ccΔ1}/+*) and homozygotes (*c(3)G^{ccΔ1}*).

Region 1, 2A, 2B, and 3 are labeled on the images as a reference. (B) Quantification of the

number of DNA double-strand breaks as assayed by counting the number of γH2AV foci per

1212   nucleus in each region of the germarium for WT and $c(3)G^{ccΔ1}$ heterozygotes ($c(3)G^{ccΔ1}$/+) and

1213   homozygotes ($c(3)G^{ccΔ1}$).

1214

1215   **Figure S3: Centromeric SC is unaffected by the $c(3)G^{ccΔ1}$ mutation**

1216   Quantification of the percent of CID foci per nucleus in WT, $c(3)G^{ccΔ1}$ heterozygotes ($ccΔ1$/+),

1217   and homozygotes ($ccΔ1$) from zygotene/early pachytene (region 2A) to mid/late pachytene

1218   (stage 8–9). (See Methods for a description of how this quantification was performed.)

1219 **Tables**

Table 1. *X* Chromosome Recombination			
Maternal genotype	*yw; pol*	$c(3)G^{cc\Delta 1}$/+	$c(3)G^{cc\Delta 1}$
	(N = 1515)	(N = 1289)	(N = 1420)
Map Length			
sc–cv	8.8	7.1	0.4
cv–v	20.7	11.9	0.7
v–f	21.1	17.7	5.2
f–y⁺	12.4	14.4	5.4
Total	**63.0**	**51.2**	**11.8**
Interference			
sc/cv/v	0.7	0.9	1.0
cv/v/f	0.6	0.9	1.0
v/f/y⁺	0.7	0.9	0.9
Class			
NCO	688	702	1264
SCO	703	512	147
DCO	120	67	7
TCO	4	3	2
Exchange rank			
E_0	0.067	0.201	0.790
E_1	0.627	0.600	0.196
E_2	0.285	0.180	0.003
E_3	0.021	0.019	0.011
E_4	0	0	0

1220 Abbreviations: N, total number of flies scored; NCO, chromatids recovered exhibiting no
1221 crossovers; SCO, single-crossover chromatids; DCO, double-crossover chromatids; TCO, triple-
1222 crossover chromatids.

1223

1224

1225

1226

1227

1228

1229

Table 2. *2nd* Chromosome Recombination			
Maternal genotype	*yw; pol*	*c(3)G$^{cc\Delta 1}$/+*	*c(3)G$^{cc\Delta 1}$*
	(N = 2376)	(N = 2471)	(N = 1456)
Map Length			
net–dpp	5.7	7.2	3.4
dpp–dpy	8.0	9.5	5.8
dpy–b	28.5	29.5	26.9
b–pr	7.8	17.2	24.9
pr–cn	2.2	9.3	6.7
Total	**52.2**	**72.8**	**67.6**
Interference			
net/dpp/dpy	0.3	–0.7	0.8
dpp/dpy/b	0.6	–0.2	0.8
dpy/b/pr	0.6	–0.2	–0.2
b/pr/cn	–0.2	–5.3	–2.2
Class			
NCO	1249	1001	624
SCO	1021	1180	692
DCO	98	255	128
TCO	8	35	12
QCO	0	1	0
Exchange rank			
E_0	0.134	0.017	0.033
E_1	0.715	0.624	0.648
E_2	0.125	0.253	0.253
E_3	0.027	0.100	0.066
E_4	0	0.006	0

Abbreviations: N, total number of flies scored; NCO, chromatids recovered exhibiting no crossovers; SCO, single-crossover chromatids; DCO, double-crossover chromatids; TCO, triple-crossover chromatids; QCO, quadruple-crossover chromatid.

Table 3. *3rd* Chromosome Recombination

Maternal genotype	*yw; pol* (N = 1014)	*c(3)G$^{cc\Delta 1}$/+* (N = 1027)	*c(3)G$^{cc\Delta 1}$* (N = 1485)
Map Length			
ru–hu	22.4	18.4	9.5
h–th	21.8	21.5	21.7
th–st	0.6	1.1	2.0
st–cu	6.1	22.8	26.5
Total	**50.9**	**63.8**	**59.6**
Interference			
ru/h/th	0.7	0.9	0.5
h/th/st	1	1	0.2
th/st/cu	−4.5	−4.5	−12.6
Class			
NCO	574	465	746
SCO	420	470	586
DCO	45	91	146
TCO	2	1	6
Exchange rank			
E_0	0.168	0.083	0.202
E_1	0.663	0.567	0.420
E_2	0.154	0.343	0.345
E_3	0.016	0.008	0.032
E_4	0	0	0

1240 Abbreviations: N, total number of flies scored; NCO, chromatids recovered exhibiting no
1241 crossovers; SCO, single-crossover chromatids; DCO, double-crossover chromatids; TCO, triple-
1242 crossover chromatids.

1243

1244

1245

1246

1247

1248

1249

1250

Table S1: Average number of CID foci per stage of oogenesis*

	Stage 1			Stage 2-4	Stage 5-7	Stage 8-9
	Region 2a	**Region 2b**	**Region 3**	**Stage 2-4**	**Stage 5-7**	**Stage 8-9**
WT	2.0±SD 0.9 (30)	2.3±SD 0.9 (30)	1.9±SD 0.6 (30)	2.4±SD 0.9 (45)	2.7±SD 1.0 (45)	2.6±SD 0.9 (27)
c(3)G^{ccΔ1}/+	1.9±SD 0.6 (30)	1.7±SD 0.6 (30)	1.4±SD 0.5 (30)	2.4±SD 0.8 (45)	2.0± SD 0.9 (45)	2.0±SD 0.7 (30)
c(3)G^{ccΔ1}	2.5±SD 0.7 (30)	2.5±SD 0.6 (30)	2.4±SD 0.6 (30)	2.1±SD 0.7 (44)	1.8± SD 0.7 (45)	2.1±SD 0.7 (30)

* SD=standard deviation; (n)=number of nuclei score

Table S2: Genomic locations of recombination markers and pairing probes (from flybase release FB2017_06)

	Symbol/Band*	Genomic location**
X chromosome recombination markers	*sc*	*XL*: 396,060
	cv	*XL*: 5,690,002
	v	*XL*: 10,923,972
	f	*XL*: 17,232,942
	y+	*XR*: unknown
2nd chromosome recombination markers	*net*	2L: 82,421
	dpp	2L: 2,428,372
	dpy	2L: 4,477,462
	b	2L: 13,821,248
	pr	2L: 20,073,719
	cn	2R: 7,782,797
3rd chromosome recombination markers	*ru*	3L: 1,370,628
	h	3L: 8,675,759
	Diap1	3L: 16,038,410
	st	3L: 16,497,651
	cu	3R: 11,197,592
X chromosome pairing probes	3C3–3C7	*XL*: 2,907,702–3,141,427
	15C1–15D6	*XL*: 16,900,783–17,066,397
2nd chromosome pairing probes	22A2–22A4	2L: 1,428,615–1,645,199
	32E2–32F2	2L: 11,217,687–11,538,470
	38E4–38F4	2L: 20,722,927–20,910,490
3rd chromosome pairing probes	63C1–63D2	3L: 756,009–1,080,330
	69B1–69C2	3L: 12,298,256–12,491,589
	77F5–78B1	3L: 20,914,572–21,101,191

*Symbol = gene symbol of marker allele used to score recombination; band = location along polytene chromosome of probe used to score pairing
**Molecular location of each marker/probe is approximate

Table S3: Percent paired at each euchromatic locus assayed*

Chromosome *2L*

Genotype	Distal (2809:22A2-22A4)			Medial (43K24:32E2-32F2)			Proximal (7D17:38E4-38F4)		
	2a	2b	3	2a	2b	3	2a	2b	3
WT	100% (21)	100% (20)	100% (13)	100% (21)	100% (15)	100% (8)	100% (25)	100% (23)	92.3% (13)
$c(3)G^{cc\Delta 1}$/+	97.7% (44)	82.6% (23)	76.2% (21)	86.2% (29)	87.5% (24)	90.9% (11)	94.1% (34)	87.1% (31)	100% (16)
$c(3)G^{cc\Delta 1}$	87.5% (24)	25% (20)	1% (9)	89.7% (29)	88.2% (17)	100% (7)	100% (37)	85.7% (28)	86.7% (15)

Chromosome *3L*

Genotype	Distal (2N23:61D-61E)			Medial (26C20:69B1-69C2)			Proximal (3J2:77F5-78B1)		
	2a	2b	3	2a	2b	3	2a	2b	3
WT	100% (25)	100% (23)	92.3% (13)	100% (31)	100% (25)	100% (13)	97.1% (35)	100% (34)	100% (14)
$c(3)G^{cc\Delta 1}$/+	92.6% (27)	90.3% (21)	83.3% (18)	100% (32)	96.8% (31)	87.5% (16)	100% (36)	94.4% (36)	95.5% (22)
$c(3)G^{cc\Delta 1}$	45.8% (24)	40% (20)	15.4% (13)	100% (33)	73.1% (26)	73.68% (19)	98.1% (54)	90.9% (33)	100% (17)

Chromosome *X*

Genotype	Distal (3D13:3C3-3C7)			Proximal (9H1:15C1-15D6)		
	2a	2b	3	2a	2b	3
WT	97.4% (38)	93.8% (32)	100% (13)	100% (43)	100% (44)	100% (18)
$c(3)G^{cc\Delta 1}$/+	92.9% (42)	71.0% (31)	56.3% (16)	100% (35)	97.1% (35)	94.1% (17)
$c(3)G^{cc\Delta 1}$	67.9% (28)	10.7% (28)	5.3% (19)	90.6% (32)	69.0% (29)	38.5% (13)

*(n) = number of nuclei scored

Table S4: Nondisjuction of X and 4^{th} chromosomes

	Maternal genotype		
	yw; pol	*c(3)G$^{cc\Delta1}$/+*	*c(3)G$^{cc\Delta1}$*
Adj. total	1348	1655	954
% X	0.7	0.2	1.5
% 4	0.4	0.1	0.6