Logistic regression analysis of multiple interosseous hand muscle activation using surface electromyography during isometric finger tasks

Masayuki Yokoyama[*] and Masao Yanagisawa

Department of Computer Science and Communications Engineering, Waseda University

* Corresponding author

E-mail: masayuki.yokoyama@islab.cs.waseda.ac.jp

Abstract

Intrinsic hand muscles are densely located in the hand, and the myoelectric observation from the surface is sometimes unreliable because of some outside influences that may interfere with the signals. In the present study, we evaluated the activation of multiple interosseous hand muscles through the surface electromyographic signals during isometric finger-oriented tasks using univariate and multivariate logistic regression models. Ten healthy subjects participated in our experiment, and isometrically exercised each finger one by one in flexed form. The result of a univariate analysis with the power and amplitude domain predictor variables of the surface electromyographic signals showed a significant consistency between the activated finger and the inserted finger of the dorsal interosseous muscles to the proximal phalanx ($P < 0.001$). Meanwhile, the results of a multivariate analysis showed a higher correlation of the regression model of the fourth dorsal interosseous muscle during the action of the ring finger using frequency-domain variables (the Nagelkerke $R^2 = 0.716$ when the median frequency was used), compared to the model without the frequency-domain variables (the Nagelkerke $R^2 = 0.583$). We concluded that logistic regression models using surface electromyographic signals have a particular possibility to produce reliable results of an activation analysis of intrinsic hand muscles. Furthermore, the use of frequency- and amplitude-domain variables was shown to assist in improving data reliability.

Introduction

Surface electromyography (sEMG) is widely used to observe the state of muscle contraction in a non-invasive manner [1]. The characteristics of the sensor, which include safe use and less cost [2], help analyze human actions not only for clinical applications, such as hand rehabilitation systems [3-5], but also for non-clinical applications in information technology development, such as hand-oriented user interface [6, 7]. Hand and finger gesture recognition is a well-studied field as a key technology of the hand-oriented user interface. Compared to the significantly heavier mechanical gloves [8-12], the light-weighted sEMG sensors potentially save the user with long-term use from exhaustion [11, 12]. Therefore, the development of light-weighted hand wearable devices using sEMG sensors is encouraged and fostered to create promising products.

Muscles related to hand and finger activities are located in the forearm (i.e., extrinsic hand muscles) and hand (i.e., intrinsic hand muscles) [13]. sEMG electrodes are generally put on the forearm to analyze activities of the wrist [14] and the whole hand, such as grasping motion [15, 16]. In case of finger-related activities, intrinsic hand muscles work harmoniously with extrinsic hand muscles [17-20].

Johnston *et al*. [17] reported the co-activation of EMG signals of extrinsic and intrinsic hand muscles during two-digit grasping motions using the thumb and the index finger with various wrist angles. Al-Sukaini *et al*. [19] examined the dominance of extrinsic or intrinsic hand muscles in finger flexion using video motion tracking, and their results showed that 69.5% of the hands of the subjects had extrinsic dominance.

However, the extrinsic hand muscles themselves are not suitable to analyze the independent motion of each finger because the flexor digitorum superficialis muscle,

as a flexor, and the extensor digitorum muscle, as an extensor, manipulate multiple fingers (i.e., from the index to the little fingers). The boundaries of muscle fibers for each finger are not explicitly separated, except for some pollicis, indicis, and digiti minimi muscles. On the other hand, intrinsic hand muscles are located very close to the fingers and each muscle is inserted into a single phalanx of a finger.

It is not necessary to put the electrodes on all related muscles to analyze a motion, because according to the phenomenon "muscle synergies" [21], a behavior produced by contraction of many muscles can be represented by a small subset. This is important especially in the field of a hand-oriented user interface, where a small and light-weighted design is required for a wearable device. Intrinsic-domain sensing has a potential to help the designers because the electrodes may only be put on the hand and the design around the forearm may not be considered. Research to utilize this phenomenon for the analysis of finger activities was conducted by Choi *et al*. [22], who evaluated the finger force during pinching by the thumb and the index finger only from three intrinsic hand muscles.

However, there is a problem in measuring sEMG signals from intrinsic hand muscles because the area in contact with the electrodes is quite small, so the signals from the adjacent muscles can interfere. Therefore, relatively few studies of finger activities by sensing intrinsic hand muscles have been conducted so far [22]. Choi *et al*. [22] applied an artificial neural network (ANN) to predict the finger force, but the ANN classifiers are known as a black box [23], which finds suitable features automatically without knowing the characteristics of the input data. Due to an overfitting problem, the results predicted by ANN se, emed good. However, it was impossible to obtain clues to select an independent variable from the results, so the network had to be designed through trial-and-error.

In this study, we examined which independent variables obtained from sEMG signals would contribute to the analysis of isometric finger activities. Kaplanis *et al*. [24] evaluated 13 independent variables extracted in the time, frequency, and bispectrum domains from the Biceps Brachii muscle generated under isometric voluntary contraction. They analyzed the data univariately using the Wilcoxon signed rank test. Finger activities, to which many extrinsic and intrinsic muscles are related, are supposed to be more complex than upper-limb activities, so we adopted logistic regression models for the multivariate analysis of finger activities.

In this paper, we focused on the binary outcome, which represents the state of finger activities as active or non-active, to simplify the analysis. We chose four dorsal interosseous muscles (DI) in a hand as the measurement targets of our experiment by putting the electrodes to measure the sEMG signals of the DI muscles on the back of the hand, which did not prevent any flexion and extension activities of the fingers.

Materials and Methods

Subjects and Experimental Environment

Ten healthy male subjects in their early twenties participated in our experiment. This research was non-clinical, and the observed data were analyzed anonymously; therefore, verbal consent was obtained from all participants. The study was approved by the Human Ethics Committee of Waseda University (No.2016-HN035). Our data is available at https://doi.org/10.6084/m9.figshare.6467801.

We used wireless sEMG sensors OE-WES1224 (Oisaka Electronic Equipment Ltd.) with bipolar Ag/AgCl electrodes. The inter-electrode distance was approximately 20 mm, and the diameter of the electrodes was 10 mm. A disposal

sheet was cut into 10-mm squares to position the electrodes on the limited space of the back of the hand. The sampling frequency of the sensors was 1 kHz. A 16-bit A/D converter and a 4th-order band-pass filter from 30 to 700 Hz were used.

Experimental Procedure

The subjects tapped their fingers on a desk one after the other, and then released them. The tapping and releasing conditions were maintained for 3 s each, and iterated for 50 s for each finger. The subjects were told to tap on the desk with as much strength as possible. The experimental forms were kept the same between the tapping and releasing conditions: the tips of the fingers were always in contact with the desk, and the motion was quasi-static. There were two tapping forms: open and closed forms [25]. This time we adopted the open form tapping because the results were similar between the two forms in our previous research [26]. We adopted the electrode positions used in [13], at the back of the hand between the metacarpal bones.

Data Analysis

Mean with standard deviation (SD), 95% confidence interval (CI), and intra-class correlation (ICC) of active (i.e., tapped) and non-active (i.e., released) data were calculated. Statistical analyses were performed using SPSS Statistics version 24.0 and Matlab for Windows.

We applied both univariate and multivariate logistic regression analyses to the data. The logistic regression model with multiple predictor variables was defined by the following equation:

$$\ln\frac{p}{1-p} = b_0 + \sum_{k=1}^{n} b_k x_k \quad (1)$$

where p represents an outcome (or dependent variable) as a probability, $b_0 \ldots b_n$ represent the constant and the coefficients of the regression model, and $x_1 \ldots x_n$ represent the predictor (or independent) variables.

We chose the following six predictor variables: maximum amplitude (V_{max} [V]), mean amplitude (V_{mean} [V]), median frequency (f_{med} [Hz]), peak frequency (f_{peak} [Hz]), maximum power of the spectrum (P_{max} [V/Hz]), and total integrated power of the spectrum (P_{total} [V/Hz]).

P-value < 0.001 was considered to be statistically significant in our logistic regression analysis.

Results and Discussion

Univariate Analysis

Table 1 shows the results of our univariate logistic regression analysis. The results of each predictor variable are shown as columns, with the correlated muscles to the finger activity shown as rows. The results during the tapped state are shown as "active," while those during the released state are shown as "non-active." Note that only the connected muscles to each finger are shown in Table 1, which include the first DI muscle inserted into the radial side of the index finger, the second DI muscle inserted into the radial side of the middle finger, the third DI muscle inserted into the ulnar side of the middle finger, and the fourth DI muscle inserted into the ulnar side of the ring finger, as shown in Fig. 1 The ICC of the active state was computed from the first seven datasets of each session of the related finger activity among all subjects. Meanwhile, the ICC of the non-active state of each muscle was computed from the first seven datasets of all sessions of the related and non-related finger activities

among all subjects. The average values of signal-to-noise ratio (SNR) of each predictor variable are also shown in Table 1. The SNR of each subject as the linear scale was computed using the following equation:

$$SNR = \frac{VAR_{active}}{VAR_{non-active}} \quad (2)$$

where VAR_{active} represents the value of a predictor variable during the active state, and $VAR_{non-active}$ represents the value of a predictor variable during the non-active state.

The active and non-active states were significantly discriminated using the amplitude (V_{max} and V_{mean}) and power (P_{max} and P_{total}) of the sEMG signals, hence acting as the predictor variables. However, the two states were not significantly discriminated using the frequency-domain variables (f_{med} and f_{peak}). In fact, none of the f_{med} values was recorded as significant in our experiment, but only the two states of the third and fourth DI muscles were discriminated using f_{peak}, making it a potential predictor variable for our univariate logistic regression analysis as well.

Some of the ICCs using f_{peak} and P_{max} as the predictor variables showed fairly low values. The ICC of the third DI muscle using f_{peak} was 0.509 when the middle finger was active, and those of the first and third DI muscles were 0.355 and 0.668, respectively, when the index and middle fingers were not active. The results implied that f_{peak} and P_{max} are not reliable enough to be used as the predictor variables of our logistic regression analysis among the 10 subjects in our experiment.

The mean SNRs of the frequency-domain variables (f_{med} and f_{peak}) were lower than those of the other variables (e.g., 1.201 of f_{med} and 1.926 of f_{peak} of the first DI muscle). The results also indicated a subtle distinction between the active and non-active states.

Fig. 2 shows the box plots of the mean amplitudes (V_{mean}) of all DI muscles when one of the fingers was active. The results showed that the distribution of the

amplitude of the DI muscles inserted into the active fingers was higher than that of the non-active fingers. Table 2 also shows that the amplitude of the inserted muscles of the active fingers was significant, while that of the non-inserted muscles was not, even though some distributions of the amplitude seemed higher in Fig. 2 (e.g., the fourth DI muscle in Fig. 2 (d) when the little finger was active).

We also showed the two types of pseudo-R^2 coefficients proposed for the logistic regression analysis [27] [28]. Cox-Snell's R^2 coefficients [27] used in our analysis are shown in Table 3 and Nagelkerke R^2 coefficients [28] are shown in Table 4. Cox-Snell's R^2 is log-likelihood based and its maximum value is less than 1.0, while Nagelkerke R^2 solves the problem and its maximum value is 1.0 [29]. However, both R^2 coefficients showed the same trends comparing the results relatively among the different conditions in Tables 3 and 4. Both results represented a high correlation of DI muscles during the action of the inserted fingers, but a low correlation during the action of the non-inserted fingers. Meanwhile, the amplitudes of the signals of DI(I), which was inserted into the index finger, only indicated a high correlation during the action of the index finger (0.344 in Table 3 and 0.720 in Table 4). DI(II) and DI(III) were both inserted into the middle finger, and similarly, the results indicated a high correlation (0.372 and 0.243 in Table 3, 0.779 and 0.508 in Table 4) during the action of the middle finger. However, the results of DI(IV) showed only a relatively high correlation, but lower than the formers (0.257 in Table 3 and 0.537 in Table 4), during the action of the ring finger. Also, none of the muscles displayed any high correlation during the action of the little finger and the thumb, which were not inserted into any DI muscle, even though the position of the fingers were adjacent to the DI muscles (i.e., DI(I) to the thumb and DI(IV) to the little finger). Comparing the two types of coefficients, Nagelkerke R^2 could be easily used to understand the

correlation intuitively, particularly for the coefficients that imitated the range of the general non-pseudo ordinary least squares R^2 (i.e., from 0.0 to 1.0).

Multivariate Analysis

We analyzed a multivariate logistic regression model that consisted of two types of predictor variables in four ways: V_{mean}-V_{max}, P_{total}-f_{med}, P_{total}-f_{peak}, and P_{total}-P_{max}. The selections were made by combining the integrated power (or amplitude) with the peak power (V_{mean}-V_{max} and P_{total}-P_{max}) and the power domain with the frequency domain (P_{total}-f_{med}, P_{total}-f_{peak}, and P_{total}-P_{max}). V_{mean} was omitted from the combination with any frequency-domain variable because the non-normalized mean amplitude was equivalent to the integrated power (P_{total}). The results are shown in Tables 5 and 6 as Cox-Snell's R^2 and Nagelkerke R^2, respectively. The correlation values of univariate regression models using V_{mean} and P_{total} are also included in the tables as a comparison to those of multivariate regression models. Note that V_{max} and P_{max} had different characteristics, and thus, the results of the models were different.

Both of the results of Cox-Snell R^2 in Table 5 and Nagelkerke R^2 in Table 6 from the correlation of the univariate logistic regression models (i.e., V_{mean} and P_{total} as the predictor variables) were slightly higher than that of the multivariate logistic regression models (i.e., V_{mean}-V_{max}, P_{total}-f_{med}, P_{total}-f_{peak}, and P_{total}-P_{max} as the predictor variables). The models with two muscles are listed in the tables as DI(II,III) during the action of the middle finger because of the two insertions of the DI muscles into the middle finger. However, the correlation was not necessarily higher than the correlation of DI(II) both in Tables 5 and 6.

Only the multivariate regression models of the ring finger showed a higher increase (0.342 with P_{total}-f_{med}, 0.336 with P_{total}-f_{peak}, and 0.356 with P_{total}-P_{max} in

Table 5, and 0.716 with P_{total}-f_{med}, 0.702 with P_{total}-f_{peak}, and 0.745 with P_{total}-P_{max} in

Table 6) as compared to the univariate regression models (0.257 with V_{mean} and 0.278

with P_{total} in Table 5, and 0.537 with V_{mean} and 0.583 with P_{total} in Table 6). As a result,

we hypothesized that the regression models of the tapping action of the ring finger

had more complexity than the other two fingers with the insertions of the DI muscles

(i.e., the index finger and the middle finger), which was also contributed by some

muscle weaknesses and inexperienced actions. In our experiment, the regression

models of the ring finger with multiple frequency-domain predictor variables (P_{total}-

f_{med}, P_{total}-f_{peak}, and P_{total}-P_{max}) showed a similar trend to the models with multiple

amplitude-domain predictor variables (V_{mean}-V_{max}). Both f_{med} and f_{peak} enhanced the

correlations of the models, while neither univariate regression model was significant

for the discrimination. Therefore, further experiments with more participants might be

needed to verify the hypothesis of the multivariate models with the predictor variables

f_{peak} and P_{max} because some of the ICCs of the models shown in Table 1 were

reasonably low, as mentioned in the section of univariate analysis (i.e., 0.509 at

DI(III) with f_{peak}, 0.355 at DI(I), and 0.668 at DI(III) with P_{max}).

Conclusions

In this paper, we evaluated the activation of multiple interosseous hand

muscles through analyzing the surface electromyographic signals during isometric

finger-oriented tasks using univariate and multivariate logistic regression models. Ten

healthy subjects participated in our experiment. The results of the univariate analysis

with the power and amplitude domain predictor variables of the surface

electromyographic signals showed a significant consistency between the activated and

inserted fingers of the DI muscles into the proximal phalanx ($P < 0.001$). Meanwhile,

the results of the multivariate analysis showed a higher correlation of the regression model of the fourth DI muscle during the action of the ring finger using frequency-domain variables (Nagelkerke $R^2 = 0.716$ when the median frequency was used), compared to the model without the frequency-domain variables (Nagelkerke $R^2 = 0.583$). Conclusively, the logistic regression models using the surface electromyographic signals had a certain possibility to produce reliable results of an activation analysis of the intrinsic hand muscles. Moreover, the frequency-domain variables, in addition to the amplitude-domain variables, might assist in improving data reliability.

Acknowledgments

The authors thank Ryohei Koyama, Prof. N. Togawa, and Prof. Y. Shi for their contributions to this work.

References

1. Merletti R, Parker PA. Electromyography: physiology, engineering, and noninvasive applications. John Wiley & Sons, Inc; 2004.

2. Yokoyama M, Koyama R, Yanagisawa M. An evaluation of hand-force prediction using artificial neural-network regression models of surface EMG Signals for Handwear Devices. J Sensors 2017: 1-12.

3. Leonardis D, Barsotti M, Loconsole C, Solazzi M, Troncossi M, Mazzotti C, et al. An EMG-controlled robotic hand exoskeleton for bilateral rehabilitation. IEEE Trans Haptics 2015 Apr-Jun;8(2): 140-151.

4. Ho NS, Tong KY, Hu XL, Fung KL, Wei XJ, Rong W, et al. An EMG-driven exoskeleton hand robotic training device on chronic stroke subjects: task training system for stroke rehabilitation. IEEE Int Conf Rehabil Robot 2011;2011: 5975340.

5. Ang KY, Huang YY, Low KH. Electromyography analysis for pre-clinical trials of hand rehabilitation tasks using design of experiments. IEEE Int Conf Mech Auto. 2009 Aug: 915-920.

6. Saponas TS, Tan DS, Morris D, Balakrishnan R. Demonstrating the feasibility of using forearm electromyography for muscle-computer interfaces. SIGCHI Conf Human Factors Computing Systems. 2008 Apr: 515-524.

7. Zhang X, Chen X, Wang WH, Yang JH, Lantz V, Wang KQ. Hand gesture recognition and virtual game control based on 3D accelerometer and EMG sensors. Int Conf Intel User Int. 2009: 401-406.

8. Popescu VG, Burdea GC, Bouzit M, Hentz VR. A virtual-reality-based telerehabilitation system with force feedback. IEEE Trans Inf Technol Biomed. 2000 Mar;4(1): 45-51.

9. Jack D, Boian R, Merians AS, Tremaine M, Burdea GC, Adamovich SV, et al. Virtual reality-enhanced stroke rehabilitation. IEEE Trans Neural Syst Rehabil Eng. 2001 Sep;9(3): 308-318.

10. Dovat L, Lambercy O, Gassert R, Maeder T, Milner T, Leong TC, et al. HandCARE: a cable-actuated rehabilitation system to train hand function after stroke. IEEE Trans Neural Syst Rehabil Eng. 2008 Dec;16(6): 582-591.

11. Ma Z, Ben-Tzvi P, Danoff J. Hand rehabilitation learning system with an exoskeleton robotic glove. IEEE Trans Neural Syst Rehabil Eng. 2016 Dec;24(12): 1323-1332.

12. Demain S, Metcalf CD, Merrett GV, Zheng D, Cunningham S. A narrative review on haptic devices: relating the physiology and psychophysical properties of the hand to devices for rehabilitation in central nervous system disorders. Disabil Rehabil Assist Technol. 2013 May;8(3): 181-189.

13. Prerotto AO. Anatomical guide for the electromyographer: the limbs and trunk. 5th ed. Springfield, Illinois: Charles C Thomas Publisher Ltd; 2011.

14. Szeto GP, Lin JK. A study of forearm muscle activity and wrist kinematics in symptomatic office workers performing mouse-clicking tasks with different precision and speed demands. J Electromyogr Kinesiol. 2011 Feb;21(1): 59-66.

15. Green S, Stefanovic B, Warman J, Askew CD. Empirical modelling of the dynamic response of fatigue during intermittent submaximal contractions of human forearm and calf muscles. J Electromyogr Kinesiol. 2015 Feb;25(1): 20-27.

16. Hoozemans MJ, van Dieën JH. Prediction of handgrip forces using surface EMG of forearm muscles. J Electromyogr Kinesiol. 2005 Aug;15(4): 358-366.

17. Johnston JA, Bobich LR, Santello M. Coordination of intrinsic and extrinsic hand muscle activity as a function of wrist joint angle during two-digit grasping. Neurosci Lett. 2010 Apr 26;474(2): 104-108.

18. Buford WL Jr, Koh S, Andersen CR, Viegas SF. Analysis of intrinsic-extrinsic muscle function through interactive 3-dimensional kinematic simulation and cadaver studies. J Hand Surg Am. 2005 Nov;30(6): 1267-1275.

19. Al-Sukaini A, Singh HP, Dias JJ. Extrinsic versus intrinsic hand muscle dominance in finger flexion. J Hand Surg Eur Vol. 2016 May;41(4): 392-399.

20. Lee SK, Wisser JR. Restoration of pinch in intrinsic muscles of the hand. Hand Clin. 2012 Feb;28(1): 45-51.

21. d'Avella A, Saltiel P, Bizzi E. Combinations of muscle synergies in the construction of a natural motor behavior. Nat Neurosci. 2003 Mar;6(3): 300-308.

22. Choi C, Kwon S, Park W, Lee HD, Kim J. Real-time pinch force estimation by surface electromyography using an artificial neural network. Med Eng Phys. 2010 Jun;32(5): 429-436.

23. Ayer T, Chhatwal J, Alagoz O, Kahn CE Jr, Woods RW, Burnside ES. Comparison of logistic regression and artificial neural network models in breast cancer risk estimation. Radiographics 2010 Jan;30(1): 13-22.

24. Kaplanis PA, Pattichis CS, Hadjileontiadis LJ, Roberts VC. Surface EMG analysis on normal subjects based on isometric voluntary contraction. J Electromyogr Kines. 2009;19: 157-171.

25. Long C, Brown ME. Electromyographic kinesiology of the hand: muscles moving the long finger. J Bone Joint Surg Am. 1964 Dec;46: 1683-1706.

26. Yokoyama M, Koyama R, Yanagisawa M. Muscle analysis of hand and forearm during tapping using surface electromyography. IEEE Global Conf Consumer Electronics 2015: 595-598.

27. Cox D. R., and Snell E. J. The analysis of binary data, 2nd ed. London: Chapman and Hall; 1989.

28. Nagelkerke, N. J. D. 1991. A note on a general definition of the coefficient of determination. Biometrika, 78: 691-692.

29. Garson G. D. Logistic regression: binary & multinomial: 2016 Edition: Statistical Associates Publishers; 2016.

List of Figures:

Figure 1. Interosseous muscles of the dorsal side of the right hand.

Figure 2. Box plots of the mean amplitudes (mV) of the sEMG signals of the first DI(I) to the fourth DI(IV). (a) Index as active finger. (b) Middle as active finger. (c) Ring as active finger. (d) Little as active finger. (e) Thumb as active finger. (f) No active fingers.

List of Tables:

Table 1. Results of the univariate logistic regression analysis.

Finger	Variables					
Electrode Position	V_{max}	V_{mean}	f_{med}	f_{peak}	P_{max}	P_{total}
Index						
DI(I)						
Non-Active						
mean (SD)	0.037 ± 0.016	0.007 ± 0.002	181.838 ± 28.835	93.963 ± 48.003	1.265 ± 0.201	0.139 ± 0.054
CI	(0.0261 - 0.048)	(0.005 - 0.008)	(161.211 - 202.466)	(59.624 - 128.302)	(1.122 - 1.409)	(0.101 - 0.178)
ICC	0.904	0.927	0.977	0.963	0.355	0.918
Active						
mean (SD)	0.228 ± 0.111	0.041 ± 0.020	205.225 ± 34.333	164.636 ± 53.617	8.142 ± 3.891	1.072 ± 0.562
CI	(0.149 - 0.308)	(0.027 - 0.056)	(180.664 - 229.785)	(126.281 - 202.992)	(5.359 - 10.926)	(0.670 - 1.474)
ICC	0.982	0.995	0.993	0.918	0.984	0.995
SNR	5.423 ± 6.119	5.556 ± 5.040	1.201 ± 0.152	1.926 ± 0.930	5.134 ± 4.229	6.998 ± 7.744
P	<0.001	<0.001	0.054	0.003	<0.001	<0.001
Middle						
DI(II)						
Non-Active						
mean (SD)	0.043 ± 0.017	0.007 ± 0.002	199.839 ± 24.102	122.574 ± 56.240	1.230 ± 0.403	0.162 ± 0.067
CI	(0.031 - 0.055)	(0.005 - 0.008)	(182.598 - 217.080)	(82.342 - 162.806)	(0.941 - 1.518)	(0.114 - 0.210)
ICC	0.919	0.926	0.977	0.954	0.876	0.931
Active						
mean (SD)	0.202 ± 0.105	0.037 ± 0.015	233.447 ± 23.053	202.405 ± 28.271	7.015 ± 2.931	1.006 ± 0.414
CI	(0.127 - 0.277)	(0.026 - 0.048)	(216.956 - 249.939)	(182.181 - 222.629)	(4.918 - 9.112)	(0.710 - 1.302)
ICC	0.972	0.984	0.984	0.740	0.980	0.985
SNR	3.060 ± 1.204	3.949 ± 1.551	1.096 ± 0.068	1.603 ± 1.411	4.282 ± 1.857	4.468 ± 2.099
P	<0.001	<0.001	0.001	0.001	<0.001	<0.001
DI(III)						
Non-Active						
mean (SD)	0.026 ± 0.013	0.005 ± 0.002	195.459 ± 14.484	100.025 ± 48.403	0.852 ± 00.268	0.104 ± 0.046
CI	(0.017 - 0.035)	(0.004 - 0.006)	(185.098 - 205.820)	(65.400 - 134.651)	(0.660 - 1.044)	(0.071 - 0.138)
ICC	0.944	0.923	0.927	0.936	0.668	0.949

Active						
mean (SD)	0.215 ± 0.131	0.040 ± 0.025	225.525 ± 18.692	183.459 ± 29.035	7.558 ± 5.008	1.078 ± 0.663
CI	(0.121 - 0.309)	(0.022 - 0.058)	(212.154 - 238.896)	(162.689 - 204.230)	(3.975 - 11.140)	(0.604 - 1.552)
ICC	0.979	0.990	0.972	0.509	0.984	0.990
SNR	7.053 ± 4.126	7.321 ± 4.089	1.057 ± 0.103	1.615 ± 0.993	8.758 ± 4.858	8.965 ± 5.088
P	<0.001	<0.001	0.001	<0.001	<0.001	<0.001

Ring

DI(IV)

Non-Active						
mean (SD)	0.029 ± 0.009	0.006 ± 0.002	175.130 ± 24.321	81.499 ± 44.176	1.041 ± 0.321	0.116 ± 0.041
CI	(0.022 - 0.035)	(0.005 - 0.007)	(157.732 - 192.529)	(49.897 - 113.100)	(0.811 - 1.270)	(0.086 - 0.145)
ICC	0.901	0.960	0.970	0.962	0.904	0.943
Active						
mean (SD)	0.260 ± 0.183	0.047 ± 0.033	213.428 ± 20.391	173.584 ± 40.373	9.394 ± 7.444	1.234 ± 0.811
CI	(0.129 - 0.391)	(0.024 - 0.070)	(198.841 - 228.014)	(144.703 - 202.465)	(4.069 - 14.719)	(0.654 - 1.814)
ICC	0.979	0.990	0.977	0.748	0.983	0.988
SNR	7.817 ± 4.77	7.900 ± 5.906	1.244 ± 0.265	2.263 ± 0.934	8.821 ± 6.527	10.805 ± 8.052
P	<0.001	<0.001	0.001	<0.001	<0.001	<0.001

Table 2. *P*-values to predict the active and non-active states of each finger when the amplitude of sEMG signals of each DI muscle was used as the predictor variable.

Finger	*P*-value (V_{mean})			
	DI(I)	DI(II)	DI(III)	DI(IV)
Index	<0.001	0.373	0.735	0.069
Middle	0.326	<0.001	<0.001	0.153
Ring	0.512	0.133	0.01	<0.001
Little	0.056	0.276	0.826	0.106
Thumb	0.692	0.081	0.074	0.095

Table 3. Cox-Snell R^2 (V_{mean}) of the univariate analysis.

Finger	Cox-Snell R^2 (V_{mean})			
	DI(I)	DI(II)	DI(III)	DI(IV)
Index	0.344	0.008	0.001	0.032
Middle	0.010	0.372	0.243	0.020
Ring	0.004	0.022	0.064	0.257
Little	0.036	0.012	0.000	0.026
Thumb	0.002	0.030	0.031	0.027

Table 4. Nagelkerke R^2 (V_{mean}) of the univariate analysis.

Finger	Nagelkerke R^2 (V_{mean})			
	DI(I)	DI(II)	DI(III)	DI(IV)
Index	0.720	0.017	0.002	0.068
Middle	0.020	0.779	0.508	0.042
Ring	0.009	0.047	0.133	0.537
Little	0.075	0.025	0.000	0.054
Thumb	0.003	0.063	0.066	0.057

Table 5. Cox-Snell R^2 of the multivariate analysis.

Finger	Predictor Variables					
Electrode Position	V_{mean}	V_{mean} - V_{max}	P_{total}	P_{total} - f_{med}	P_{total} - f_{peak}	P_{total} - P_{max}
Index						
DI(I)	0.344	0.348	0.340	0.342	0.340	0.341
Middle						
DI(II)	0.372	0.380	0.370	0.371	0.370	0.375
DI(III)	0.243	0.253	0.246	0.265	0.272	0.248
DI(II,III)	0.374	0.398	0.371	0.382	0.374	0.381
Ring						
DI(IV)	0.257	0.274	0.278	0.342	0.336	0.356

Table 6. Nagelkerke R^2 of the multivariate analysis.

Finger	**Predictor Variables**					
Electrode Position	V_{mean}	$V_{mean} - V_{max}$	P_{total}	$P_{total} - f_{med}$	$P_{total} - f_{peak}$	$P_{total} - P_{max}$
Index						
DI(I)	0.720	0.729	0.712	0.716	0.712	0.713
Middle						
DI(II)	0.779	0.795	0.775	0.777	0.775	0.784
DI(III)	0.508	0.529	0.515	0.555	0.569	0.519
DI(II,III)	0.783	0.833	0.775	0.799	0.783	0.797
Ring						
DI(IV)	0.537	0.574	0.583	0.716	0.702	0.745

Fig. 1

DI(I)

DI(II)

DI(III)

DI(IV)

Fig. 2