Finding Nemo's Genes: A chromosome-scale reference assembly of the genome of the

orange clownfish *Amphiprion percula*

Robert Lehmann[1], Damien J. Lightfoot[1], Celia Schunter[1], Craig T. Michell[2], Hajime

Ohyanagi[3], Katsuhiko Mineta[3], Sylvain Foret[4,5], Michael L. Berumen[2], David J. Miller[4],

Manuel Aranda[2], Takashi Gojobori[3], Philip L. Munday[4] and Timothy Ravasi[1,*]

[1] KAUST Environmental Epigenetic Program, Division of Biological and Environmental

Sciences & Engineering, King Abdullah University of Science and Technology, Thuwal,

23955-6900, Kingdom of Saudi Arabia.

[2] Red Sea Research Center, Division of Biological and Environmental Sciences &

Engineering, King Abdullah University of Science and Technology, Thuwal, 23955-6900,

Kingdom of Saudi Arabia.

[3] Computational Bioscience Research Center, King Abdullah University of Science and

Technology, Thuwal, 23955-6900, Kingdom of Saudi Arabia.

[4] ARC Centre of Excellence for Coral Reef Studies, James Cook University, Townsville,

Queensland, 4811, Australia.

[5] Evolution, Ecology and Genetics, Research School of Biology, Australian

National University, Canberra, Australian Capital Territory, 2601, Australia.

Keywords:

Orange Clownfish, *Amphiprion percula*, Nemo, Functional Genomics, Chromosome-Scale

Assembly, Fish Genomics, Coral Reef Fish.

Running Title:

The Nemo Genome

[*]**Corresponding Author:**

Timothy Ravasi, Division of Biological and Environmental Sciences & Engineering, King

Abdullah University of Science and Technology, Thuwal, 23955-6900, Kingdom of Saudi

Arabia, timothy.ravasi@kaust.edu.sa

Abstract

The iconic orange clownfish, *Amphiprion percula*, is a model organism for studying the

ecology and evolution of reef fishes, including patterns of population connectivity, sex

change, social organization, habitat selection and adaptation to climate change. Notably, the

orange clownfish is the only reef fish for which a complete larval dispersal kernel has been

established and was the first fish species for which it was demonstrated that anti-predator

responses of reef fishes could be impaired by ocean acidification. Despite its importance,

molecular resources for this species remain scarce and until now it lacked a reference genome

assembly. Here we present a *de novo* chromosome-scale assembly of the genome of the

orange clownfish *Amphiprion percula*. We utilized single-molecule real-time sequencing

technology from Pacific Biosciences to produce an initial polished assembly comprised of

1,414 contigs, with a contig N50 length of 1.86 Mb. Using Hi-C based chromatin contact

maps, 98% of the genome assembly were placed into 24 chromosomes, resulting in a final

assembly of 908.8 Mb in length with contig and scaffold N50s of 3.12 and 38.4 Mb,

respectively. This makes it one of the most contiguous and complete fish genome assemblies

currently available. The genome was annotated with 26,597 protein coding genes and contains

96% of the core set of conserved actinopterygian orthologs. The availability of this reference

genome assembly as a community resource will further strengthen the role of the orange

clownfish as a model species for research on the ecology and evolution of reef fishes.

Introduction

The orange clownfish, *Amphiprion percula*, which was immortalized in the film "Finding Nemo", is arguably the most recognized fish on Earth. It is also one of the most important species for studying the ecology and evolution of coral reef fishes. The orange clownfish is used as a model species to study patterns and processes of social organization (Buston, Bogdanowicz, Wong, & Harrison, 2007; Buston & Wong, 2014; Wong, Uppaluri, Medina, Seymour, & Buston, 2016), sex change (Buston, 2003), mutualism (Schmiege, D'Aloia, & Buston, 2017), habitat selection (Dixson et al., 2008; Elliott & Mariscal, 2001; Scott & Dixson, 2016), lifespan (Buston & García, 2007) and predator-prey interactions (Dixson, 2012; Manassa, Dixson, McCormick, & Chivers, 2013). It has been central to ground-breaking research into the scale of larval dispersal and population connectivity in marine fishes (Almany et al., 2017; Pinsky et al., 2017; Planes, Jones, & Thorrold, 2009; Salles et al., 2016) and how this influences the efficacy of marine protected areas (Berumen et al., 2012; Planes et al., 2009). It is also used to study the ecological effects of environmental disturbances in marine ecosystems (Hess, Wenger, Ainsworth, & Rummer, 2015; Wenger et al., 2014), including climate change (McLeod et al., 2013; Saenz-Agudelo, Jones, Thorrold, & Planes, 2011) and ocean acidification (Dixson, Munday, & Jones, 2010; Jarrold, Humphrey, McCormick, & Munday, 2017; Munday et al., 2009; Simpson et al., 2011). Perhaps more than any other species, the orange clownfish has become a mainstay of research into the chemical, molecular, behavioral, population, conservation and climate-change ecology of marine fishes.

The orange clownfish is one of 30 species of anemonefishes belonging to the subfamily Amphiprioninae within the family Pomacentridae (damselfishes). The two clownfishes, *A. percula* (orange clownfish or clown anemonefish) and *A. ocellaris* (false clownfish or western

clown anemonefish) form a separate clade, alongside *Premnas biaculeatus*, within the

Amphiprioninae (J. Li, Chen, Kang, & Liu, 2015; Litsios, Pearman, Lanterbecq, Tolou, &

Salamin, 2014; Litsios & Salamin, 2014). The two species of clownfish are easily

distinguished from other anemonefishes by their bright orange body coloration and three

vertical white bars. The orange clownfish and the false clownfish have similar body

coloration, but largely distinct allopatric geographical distributions (Litsios & Salamin, 2014).

The orange clownfish occurs in northern Australia, including the Great Barrier Reef (GBR),

and in Papua New Guinea, Solomon Islands and Vanuatu, while the false clownfish occurs in

the Indo-Malaysian region, from the Ryukyu Islands of Japan, throughout south-east Asia and

south to north-western Australia (but not the GBR).

Like all anemonefishes, the orange clownfish has a mutualistic relationship with sea-

anemones. Wild adults and juveniles live exclusively in association with a sea anemone,

where they gain shelter from predators and benefit from food captured by the anemone

(Fautin, 1991; Fautin & Allen, 1997; Mebs, 2009). In return, the sea-anemone benefits by

gaining protection from predators (Fautin & Allen, 1997; Holbrook & Schmitt, 2005), from

supplemental nutrition from the clownfish's waste (Holbrook & Schmitt, 2005) and from

increased gas exchange as a result of increased water flow provided by clownfish movement

and activity (Herbert, Bröhl, Springer, & Kunzmann, 2017; Szczebak, Henry, Al-Horani, &

Chadwick, 2013). The orange clownfish associates with two species of anemone,

Stichodactyla gigantea and *Heteractis magnifica* (Fautin & Allen, 1997). Clownfish social

groups typically consist of an adult breeding pair and a variable number of smaller, size-

ranked juveniles that queue for breeding rights (Buston, 2003). The breeding female is larger

than the male. If the female disappears, the male changes sex to female and the largest non-

breeder matures into a breeding male. The breeding pair lays clutches of demersal eggs in

close proximity to their host anemone. Eggs hatch after 7-8 days and the larvae disperse into

the open ocean for a period of 11-12 days, at which time they return to the reef and settle to

an anemone.

The close association of clownfish and other anemonefishes with sea anemones makes them

excellent species for studying aspects of marine mutualisms and habitat selection. The easily

identified and delineated habitat they occupy, along with the ease with which the fish can be

observed in nature, makes them ideal candidates for behavioral and population ecology. The

unique capacity to collect juveniles immediately after they have settled to the reef from their

pelagic larval phase also makes them ideally suited to testing long-standing questions about

larval dispersal and population connectivity in reef fish populations. Using molecular

techniques to assign parentage between newly settled juveniles and adult anemonefishes,

recent studies have been able to describe for the first time the spatial scales of dispersal in reef

fish and its temporal consistency (Almany et al., 2017). The ability to map the connectivity of

clownfish populations in space and time has also opened the door to addressing challenging

questions about selection, fitness and adaptation in natural populations of marine fishes

(Pinsky et al., 2017; Salles et al., 2016). Finally, the orange clownfish is one of the relatively

few coral reef fishes that can easily be reared in captivity (Wittenrich, Turingan, & Creswell,

2007). Consequently, it has unrivalled potential for experimental manipulation to test

ecological and evolutionary questions in marine ecology (Dixson et al., 2014; Manassa et al.,

2013), including the impacts of climate change and ocean acidification (Nilsson et al., 2012).

Increasingly, genome-wide methods are being used to test ecological and evolutionary

questions and this is particularly true for coral reef species in the wake of anthropomorphic

climate change and its effects on these sensitive ecosystems (Stillman & Armstrong, 2015)

To date, genome assemblies of two anemonefish, *A. frenatus* (Marcionetti, Rossier, Bertrand,

Litsios, & Salamin, 2018) and *A. ocellaris* (Marcionetti et al., 2018), have been published.

Both of these were based on short-read Illumina technology with genome scaffolding

provided by shallow coverage of PacBio (Marcionetti et al., 2018) or Oxford Nanopore (Tan

et al., 2018) long reads. While the use of long reads to scaffold Illumina-based assemblies

improves contiguity, both genome assemblies are highly fragmented with respective contig

and scaffold N50s of 14.9 and 244.5 kb for *A. frenatus* and 323.6 and 401.7 kb for *A.*

ocellaris. Here we present a chromosome-scale genome assembly of the orange clownfish,

which was assembled using a primary PacBio long read strategy, followed by scaffolding

with Hi-C-based chromatin contact maps. The resulting final assembly is highly contiguous

with contig and scaffold N50 values of 3.12 and 38.4 Mb, respectively. This assembly will be

a valuable resource for the research community and will further establish the orange

clownfish as a model organism for genetic and genomic studies into ecological, evolutionary

and environmental aspects of reef fishes. To facilitate the use of this resource, we have

developed an integrated database, the Nemo Genome DB (http://nemogenome.org/), which

allows for the interrogation and mining of genomic and transcriptomic data described here.

Materials and Methods

Specimen collection and DNA extraction

Adult orange clownfish breeding pairs were collected on the northern GBR in Australia. Fish

were bred at the Experimental Aquarium Facility of James Cook University (JCU) and one

individual offspring was sacrificed at the age of 8 months. The whole brain was excised, snap

frozen and kept at -80°C until processing. High molecular weight DNA was extracted from

whole brain tissue using the Qiagen Genomic-tip 100/G extraction kit. The tissue was first

homogenized in lysis buffer G2 supplemented with 200 μg/mL RNase A using sterile beads

for 30 sec. After homogenization, proteinase K was added and the homogenate was incubated

at 50°C overnight. DNA extraction was then performed according to the manufacturer's

protocol with a final elution volume of 200 μl. DNA fragment size and quality was assessed

using pulsed-field gel electrophoresis. This study was completed under JCU animal ethics

permits A1961 and A2255.

PacBio library preparation and sequencing

For Pacific Biosciences (PacBio) long read sequencing, the extracted orange clownfish DNA

was first sheared using a g-TUBE (Covaris, MA, USA) (target size of 20 kb) and then

converted into SMRTbell template libraries according to the manufacturer's protocol (Pacific

Biosciences, CA, USA). Size selection was performed using BluePippin (Sage Science, MA,

USA) to generate two libraries with a minimum size of 10 and 15 kb, respectively.

Sequencing was performed using P6-C4 chemistry on the PacBio RS II instrument at the King

Abdullah University of Science and Technology (KAUST) Bioscience Core Laboratory

(BCL) with 360 mins movies. A total of 113 SMRT cells were sequenced.

Genome assembly

The genome sequence was assembled from the unprocessed PacBio reads (Table S1) using

the hierarchical diploid aware PacBio assembler FALCON v0.4.0 (Chin et al., 2016). To

obtain the optimal assembly, different parameters were tested (Table S2) to generate 12

candidate assemblies. The contiguity of these assemblies was assessed with QUAST v3.2

(Gurevich, Saveliev, Vyahhi, & Tesler, 2013), while assembly completeness was determined

with BUSCO v2.0 (Simão, Waterhouse, Ioannidis, Kriventseva, & Zdobnov, 2015).

Assembly "A7" exhibits the highest contiguity and single copy orthologous gene

completeness and was selected for further improvement. The FALCON_Unzip algorithm was

then applied to the initial A7 assembly obtain a haplotype-resolved, phased assembly, termed

"A7-phased". Contigs less than 20 kb in length were removed from the assembly. This phased

assembly was polished with Quiver to achieve final consensus sequence accuracies

comparable to Sanger sequencing (Chin et al., 2013) using default settings, which produced

the "A7-phased-polished" assembly.

<u>Genome assembly scaffolding with chromatin contact maps</u>

The flash-frozen brain tissue was sent to Phase Genomics (Seattle, WA, USA) for the

construction chromatin contact maps. Tissue fixation, chromatin isolation, library preparation

and 80-bp paired end sequencing were performed by Phase Genomics. The sequencing reads

were aligned to the A7-phased-polished version of the assembly with BWA (H. Li & Durbin,

2010) and uniquely mapping read pairs were retained. Contigs from the A7-phased-polished

assembly were clustered, ordered and then oriented using Proximo (Bickhart et al., 2017;

Burton et al., 2013), with settings as previously described (Peichel, Sullivan, Liachko, &

White, 2017). Briefly, contigs were clustered into chromosomal groups using a hierarchical

clustering algorithm based on the number of read pairs linking scaffolds, with the final

number of groups specified as the number of the haploid chromosomes. The haploid

chromosome number was set as 24, which is consistent with the observed haploid

chromosome number of the Amphiprioninae, as published for *A. ocellaris* (Arai, Inoue, & Ida,

1976), *A. frenatus*, (Molina & Galetti, 2004; Takai & Kosuga, 2007), *A. clarkii* (Arai &

Inoue, 1976; Takai & Kosuga, 2007), *A. perideraion* (Supiwong et al., 2015) and *A. polymnus*

(Tanomtong et al., 2012). After clustering into chromosomal groups, the scaffolds were

ordered based on Hi-C link densities and then oriented with respect to the adjacent scaffolds

using a weighted directed acyclic graph of all possible orientations based on the exact

locations of the Hi-C links between scaffolds. Gaps between contigs were represented with

100 Ns and the proximity-guided assembly was named "A7-PGA". Gaps in the scaffolded

assembly were subsequently closed using PBJelly from PBSuite v15.8.24 (English et al.,

2012) with the entire PacBio read dataset and Blasr (Chaisson & Tesler, 2012) (parameters: --

minMatch 8 --minPctIdentity 70 --bestn 1 --nCandidates 20 --maxScore -500 --nproc 32 --

noSplitSubreads), to give rise to the final version of the assembly, "Nemo v1".

<u>Genome assembly validation</u>

Genomic DNA was extracted from a second individual and Illumina sequencing libraries

were prepared using the NEBNext Ultra II DNA library prep kit for Illumina following the

manufacturer's protocol. Three cycles of PCR were used to enrich the library. The sequencing

libraries were sequenced on two lanes of a HiSeq 2500 at the KAUST BCL. A total of

1,199,533,204 paired reads were generated, covering approximately 181 Gb. The 151-bp

paired end reads were processed with Trimmomatic v0.33 to remove adapter sequences and

low-quality stretches of nucleotides (parameters: 2:30:10 LEADING:20 TRAILING:20

SLIDINGWINDOW:4:20 MINLEN:75) (Bolger, Lohse, & Usadel, 2014).

The genome assembly size was validated by comparison to a k-mer based estimate of genome

size. The first half of the paired-end reads of one sequencing lane (~ 25 Gb of data) was used

for the k-mer estimate of genome size. Firstly, KmerGenie (Chikhi & Medvedev, 2014) was

used to determine the optimal k-value for a k-mer based estimation. Following that, Jellyfish

v2.2.6 (Marçais & Kingsford, 2011) was used with k=71 to obtain the frequency distribution

of all k-mers with this length. The resulting distribution was analyzed with Genomescope

(Vurture et al., 2017) to estimate genome size, repeat content and the level of heterozygosity.

To further validate the assembly, we determined the proportion of trimmed Illumina short

reads that mapped to the Nemo v1 assembly with BWA v0.7.10 (H. Li & Durbin, 2010) and

SAMtools v1.1 (H. Li et al., 2009). Additionally, the completeness of the genome assembly

annotation as determined by the conservation of a core set of genes was measured using

BUSCO with default parameters.

Repeat annotation

A species-specific *de novo* repeat library was assembled by combining the results of three

distinct repeat annotation methods. Firstly, RepeatModeler v1.08 (Smit & Hubley, 2008) was

used to build an initial repeat library. Secondly, we used LtrHarvest (Ellinghaus, Kurtz, &

Willhoeft, 2008) and LTRdigest (Steinbiss, Willhoeft, Gremme, & Kurtz, 2009), both

accessed *via* genometools 1.5.6 (Gremme, Steinbiss, & Kurtz, 2013), with the following

parameters: -seed 76 -xdrop 7 -mat 2 -mis -2 -ins -3 -del -3 -mintsd 4 -maxtsd 20 -minlenltr

100 -maxlenltr 6000 -maxdistltr 25000 -mindistltr 1500 -similar 90. The resulting hits were

filtered with LTRdigest, accepting only sequences featuring a hit to one of the hidden markov

models in the GyDB 2.0 database. Thirdly, TransposonPSI v08222010 (Haas, 2018) was used

to detect sequences with similarities to known families of transposon open reading frames. To

remove duplicated sequences in the combined result from all three methods a clustering with

USEARCH (Edgar, 2010) was performed requiring at least 90% sequence identity, and only

cluster representatives were retained. The resulting representative sequences were classified

by RepeatClassifier (part of RepeatModeler), Censor v4.2.29 (Jurka, Klonowski, Dagman, &

Pelton, 1996) and Dfam v2.0 (Wheeler et al., 2012), and were then blasted against the

Uniprot/Swissprot database (release 2017_12) to obtain a unified classification. Furthermore,

these three classification methods and the blast result was used to filter out spurious matches

to protein-coding sequence. Specifically, putative repeat sequences were only retained when

at least one classification method recognized the sequence as a repeat and the best match in

Swissprot/Uniprot was not a protein-coding gene (default blastx settings). Furthermore,

sequences were retained if two of the three identification methods classified the sequence as

repeat, but the best blast hit was not a transposable element. This *de novo* library was

combined with the thoroughly curated zebrafish repeat library provided by Repbase v22.05

(Bao, Kojima, & Kohany, 2015) and this combined library was employed for repeat masking

in the Nemo v1 assembly using RepeatMasker (Smit, Hubley, & Green, 2010).

RNA extraction, library construction, sequencing and read processing

Tissues for RNA extraction were dissected from one eight-month old orange clownfish

individual. RNA was extracted from skin, eye, muscle, gill, liver, kidney, gallbladder,

stomach and fin tissues using the Qiagen AllPrep kit following manufacturer's instructions.

Sequencing libraries were prepared using the TruSeq Stranded mRNA Library Preparation kit

and 150 bp paired-end sequencing was performed on one lane of an Illumina HiSeq 4000

machine in the KAUST BCL. The RNA-seq reads were trimmed with Trimmomatic v0.33

(Bolger et al., 2014) (parameters: 2:30:10 LEADING:3 TRAILING:3

SLIDINGWINDOW:4:15 MINLEN:40) and contamination was removed with Kraken (Wood

& Salzberg, 2014) by retaining only unclassified reads.

<u>Genome assembly annotation</u>

After mapping the RNA-seq data with STAR v2.5.2b (Dobin et al., 2013) to the final

assembly, an *ab-initio* annotation with BRAKER1 v1.9 (Hoff, Lange, Lomsadze,

Borodovsky, & Stanke, 2016) was performed. This initial annotation identified 49,881 genes.

This annotation was then integrated with external evidence using the MAKER2 v2.31.8 (Holt

& Yandell, 2011) gene annotation pipeline. First, the transcriptome of the orange clownfish

was provided to MAKER2 as EST evidence in two forms, a *de novo* assembly of the

preprocessed RNA-seq reads obtained with Trinity v2.4.0 (Grabherr et al., 2011), and a

genome-guided assembly performed with the Hisat2 v2.1.0/Stringtie v1.3.3b workflow

(Pertea, Kim, Pertea, Leek, & Salzberg, 2016). Second, we combined the proteomes of

zebrafish (GCF_000002035.6_GRCz11), Nile tilapia (GCF_001858045.1_ASM185804v2)

and bicolor damselfish (*Stegastes partitus*) (GCA_000690725.1), together with the

Uniprot/Swissprot database (release 2017_12: 554,515 sequences) and the successfully

detected BUSCO genes to generate a reference protein set for homology based gene

prediction. In the initial MAKER2 run, the annotation edit distances (AED) were calculated

for the BRAKER1-obtained annotation, and only gene annotations with an AED of less than

0.1 and a corresponding protein length of greater than 50 amino acids were retained for

subsequent training of the gene prediction program SNAP v2013.11.29 (Korf, 2004).

Similarly, the AUGUSTUS v3.2.3 (Stanke et al., 2006) gene prediction program was trained

on 1,850 gene annotations that possessed: an AED score of less than 0.01; an initial start

codon, a terminal stop codon and no in-frame stop codons; more than one exon; and no

introns greater than 10 kb. The hidden markov gene model of GeneMark v4.32 (Ter-

Hovhannisyan, Lomsadze, Chernoff, & Borodovsky, 2008) was trained by BRAKER1. The

final annotation was then obtained in the second run of MAKER2 with the trained models for

SNAP, GeneMark, and AUGUSTUS. InterProScan 5 was then used to obtain the Pfam

protein domain annotations for all genes. The standard gene builds were then generated. The

output was filtered to include all annotated genes with evidence (AED less than 1) or with a

Pfam protein domain, as recommended (Campbell, Holt, Moore, & Yandell, 2014).

<u>Functional annotation</u>

The protein sequences produced from the genome assembly annotation were aligned to the

UniProtKB/Swiss-Prot database (release 2017_12) with blastp v2.2.29 (parameters: -outfmt 5

-evalue 1e-3 -word_size 3 -show_gis -num_alignments 20 -max_hsps 20) and protein

signatures were annotated with InterProScan 5. The results were then integrated with

Blast2GO v4.1.9 (Gotz et al., 2008).

<u>Genome assembly comparisons</u>

For genome assembly comparisons, we compared the Nemo v1 genome assembly to the 26

previously reported fish chromosome-scale genome assemblies (Table S3). Comparisons were

made for genome assembly contiguity and completeness. Contig N50 values are reported for

the scaffold-scale versions of each assembly and are taken from the indicated publication

(Table S3), database description (Table S3) or were generated with the Perl

assemblathon_stats_2.pl script (Bradnam et al., 2013). Genome assembly completeness was

assessed by determining the proportion of the genome size that is contained within the

chromosome content of each assembly. It should be noted that this comparison is relative to

the estimated genome size and not the published assembly size. The estimated genome size

was taken as either the published estimated genome size in the relevant paper (Table S3), or

from the Animal Genome Size Database (Gregory, 2018). Where possible, k-mer derived or

flow cytometry-based estimates of genome size were used. Before calculation, we remove

stretches of Ns from the genome assemblies as these are used to arbitrarily space scaffolds

and do not contain actual genome information. However, this step was not possible for the

Asian arowana, southern platyfish, yellowtail or croaker genomes as the chromosome-scale

assemblies have not been made publicly available. Genome assembly completeness was

determined with BUSCO (Simão et al., 2015) using the Actinopterygii set of 4,584 genes and

the AUGUSTUS zebrafish gene model provided with the software.

<u>Gene homology</u>

To investigate the gene space of the orange clownfish genome assembly, we used

OrthoFinder v1.1.4 (Emms & Kelly, 2015) to identify orthologous gene relationships between

the orange clownfish and four related fish species. The following four fish species were

utilized in addition to the orange clownfish: Asian seabass

GCF_001640805.1_ASM164080v1 (45,223 sequences), Nile tilapia

GCF_001858045.1_ASM185804v2 (58,087 sequences), southern platyfish

GCF_000241075.1_*Xiphophorus_maculatus*-4.4.2 (23,478 sequences), and zebrafish

GCF_000002035.6_GRCz11 (52,829 sequences). The longest isoform of each gene was

utilized in the analysis, which corresponded to 25,050, 28,497, 23,043, and 32,420 sequences,

respectively. 26,597 sequences were used for the orange clownfish. These protein sequences

were reciprocally blasted against each other and clusters of orthologous genes were then

defined using OrthoFinder with default parameters. As part of OrthoFinder, the concatenated

sequences of single-copy orthologs present in all species were then used to construct a

phylogenetic tree, which was rooted using STRIDE (Emms & Kelly, 2017).

Database system architecture and software

The Nemo Genome DB database (http://nemogenome.org) was implemented on a UNIX

server with CentOS version 7, Apache web server and MySQL Database server. JBrowse

(Buels et al., 2016) was employed to visualize the genome assembly and genomic features

graphically and interactively. JavaScript was adopted to implement client-side rich

applications. The JavaScript library, jQuery (http://jquery.com) was employed. Other

conventional utilities for UNIX computing were appropriately installed on the server if

necessary. All of the Nemo Genome DB resources are stored on the server and are available

through HTTP access.

Results and Discussion

Sequencing and assembly of the orange clownfish genome

Genomic DNA of an individual orange clownfish (Fig. 1A) was sequenced with the PacBio

RS II platform to generate 1,995,360 long reads, yielding 113.8 Gb, which corresponds to a

121-fold coverage of the genome (Table S1). After filtering with the read pre-assembly step

of the Falcon assembler, 5,764,748 reads, covering 54.3 Gb and representing a 58-fold

coverage of the genome, were available for assembly.

To optimize the assembly parameters, we performed 12 trial assemblies using a range of parameters for different stages of the Falcon assembler (Table S2). The assembly quality was assessed by considering assembly contiguity (contig N50 and L50), total assembly size, and also gene completeness (BUSCO) (Table 1). Assembly A7 exhibited the highest contig N50 (1.80 Mb), lowest contig L50 (138 contigs), lowest number of missing BUSCO genes (132) and is only slightly surpassed in the longest contig metric (15.8 Mb) by the highly similar assemblies A8 and A9 (16.5 Mb) (Table 1).

Genome assemblies represent a mixture of the two possible haplotypes of a diploid individual at each locus. This collapsing of haplotypes may result in a loss of important sequence information. However, diploid-aware assembly algorithms such as the Falcon_Unzip assembler are designed to detect single nucleotide polymorphisms (SNPs) as well as structural variations and to use this information to phase ("unzip") heterozygous regions into distinct haplotypes (Chin et al., 2016). This procedure results in a primary assembly and a set of associated haplotype contigs (haplotigs) capturing the divergent sequences. Having established the parameter set that gave the best assembly metrics with Falcon, we used Falcon_Unzip to produce a phased assembly ("A7-phased") of the orange clownfish (Table 2). The phased assembly was 905.0 Mb in length with a contig N50 of 1.85 Mb. As has been seen in previous genome assembly projects (Chin et al., 2016), Falcon_Unzip produced a smaller assembly with fewer contigs than the assembly produced by Falcon (Table 2). The phased primary assembly was then polished with Quiver, which yielded an assembly ("A7-phased-polished") with 1,414 contigs spanning 903.6 Mb with an N50 of 1.86 Mb (Table 2). This polishing step closed 91 gaps in the assembly and improved the N50 by approximately 14.3 kb. After polishing of the "unzipped" A7-phased-polished assembly, 9,971 secondary

contigs were resolved, covering 340.1 Mb of the genome assembly. The contig N50 of these

secondary contigs was 38.2 kb, with over 99% of them being longer than 10 kb in size.

Relative to the 903.6 Mb A7-phased-polished primary contig assembly, the secondary contigs

covered 38% of the assembly size. To the best of our knowledge, this is the first published

fish genome assembly that has been resolved to the haplotype level with Falcon_Unzip.

<u>Scaffolding of the orange clownfish genome assembly into chromosomes</u>

To build a chromosome-scale reference genome assembly of the orange clownfish, chromatin

contact maps were generated by Phase Genomics (Fig. S1). Scaffolding was performed by the

Proximo algorithm (Bickhart et al., 2017; Burton et al., 2013) on the A7-phased-polished

assembly using 231 million Hi-C-based paired-end reads to produce the proximity guided

assembly "A7-PGA" (Table 2). The contig clustering allowed the placement of 1,073 contigs

into 24 scaffolds (chromosomes) with lengths ranging from 23.4 to 45.8 Mb (Tables 2 and 3).

While only 76% of the contigs were assembled into chromosome clusters, this corresponds to

98% (885.4 Mb) of total assembly length and represents 95% of the estimated genome size of

938.9 Mb (Tables 2 and 3). This step substantially improved the overall assembly contiguity,

raising the N50 20-fold from 1.86 to 38.1 Mb.

A quality score for the order and orientation of contigs within the A7-PGA assembly was

determined. This metric is based on the differential log-likelihood of the contig orientation

having produced the observed log-likelihood, relative to its neighbors (Burton et al., 2013).

The orientation of a contig was deemed to be of high quality if its placement and orientation,

relative to neighbors, was 100 times more likely than alternatives (Burton et al., 2013). In A7-

PGA, the placements of 524 (37%) of the scaffolds were deemed to be of high quality,

accounting for 775.5 Mb (87%) of the scaffolded chromosomes, indicating the robustness of

the assembly.

A final polishing step was performed with PBJelly to generate the final Nemo v1 assembly.

This polishing step closed 369 gaps, thereby improving the contig N50 by 68% and increasing

the total assembly length by 5.21 Mb (Tables 2 and 3). The length of each chromosome was

increased, with a range of 23.7 to 46.1 Mb (Fig. 1B). Gaps were closed in each chromosome

except for chromosome 14, leaving an average of only 28 gaps per chromosome (Table 3).

The final assembly is 908.9 Mb in size and has contig and scaffold N50s of 3.12 and 38.4 Mb,

respectively. The assembly is highly contiguous as can be observed by the fact that 50% of

the genome length is contained within the largest 84 contigs. 890.2 Mb (98%) of the genome

assembly size was scaffolded into 24 chromosomes, with only 18.8 Mb of the assembly

failing to be grouped. The 18.8 Mb of unscaffolded assembly is comprised of 341 contigs

with a contig N50 of only 57.8 kb.

Validation of the orange clownfish genome assembly size

The final assembly size of 908.9 Mb is consistent with the results of a Feulgen image analysis

densitometry-based study, which determined a C-value of 0.96 pg and thus a genome size of

938.9 Mb for the orange clownfish (Hardie & Hebert, 2004). Furthermore, our assembly size

is in keeping with estimates of genome size for other fish of the *Amphiprion* genus, which

range from 792 to 1193 Mb (Gregory, 2018). We additionally validated the observed

assembly size by using a k-mer based approach. Specifically, the k-mer coverage and

frequency distribution were plotted and fitted with a four-component statistical model with

GenomeScope (Fig. S2A). This allowed us to generate an estimate of genome size as well as

the repeat content and level of heterozygosity. However, varying the k-value from the

recommended value of 21 up to 27 yielded a corresponding increase of the estimated genome

size. We therefore used KmerGenie to determine the optimal k-mer length of 71 to capture the

available sequence information. The utilization of small k-values might partially explain the

reported tendency of GenomeScope to underestimate the genome size (Vurture et al., 2017).

The final estimate of the haploid genome length by k-mer analysis was 906.6 Mb, with 732.8

Mb (80%) of unique sequence and a repeat content of 173.8 Mb (19%). Furthermore, the

estimated heterozygosity level of 0.12% is low considering that an F1 offspring of wild

caught fish was sequenced (Fig. S2B). While the short-read k-mer based genome size

estimate of 906.6 Mb matches the final assembly size of 908.9 Mb very well, the C-value

derived genome size estimate is slightly larger (938.9 Mb). As an additional validation of the

accuracy of the genome assembly, we mapped the trimmed Illumina short reads to the Nemo

v1 assembly and observed that 95% of the reads mapped to the assembly and that 84% of the

reads were properly paired.

Based on the C-value derived genome size estimate, there is approximately 29.9 Mb (3.3%)

of sequence length absent from our genome assembly. It seems likely that our assembly is

nearly complete for the euchromatic regions of the genome given our assessment of genome

size and gene content completeness. However, genomic regions such as the proximal and

distal boundaries of euchromatic regions contain heterochromatic and telomeric repeats,

respectively, are refractory to currently available sequencing techniques and are typically

absent from genome assemblies (Bickhart et al., 2017; Hoskins et al., 2007).

<u>Chromosome-scale fish genome assembly comparisons</u>

To date, chromosome-scale genome assemblies have been released for 26 other fish species

(Table S3). Here, we present the first chromosome-scale assembly of a tropical coral reef fish,

the orange clownfish. As a measure of genome assembly quality, we assessed the contiguity

and completeness of these 27 chromosome-scale genome assemblies. We investigated

genome contiguity with the contig N50 metric and characterized genome completeness for

each genome assembly by calculating the proportion of the estimated genome size that was

assigned to chromosomes. As shown in Fig. 1C, the orange clownfish genome assembly is

highly contiguous, with a scaffold-scale contig N50 of 1.86 Mb, which is only surpassed by

the contig N50 of the Nile tilapia genome assembly. Interestingly, even though different

assembler algorithms were utilized, the three genome assemblies based primarily on long read

PacBio technology were the most contiguous, with only Nile tilapia (3.09 Mb, Canu), orange

clownfish (1.86 Mb, Falcon) and Asian seabass (1.19 Mb, HGAP) genome assemblies

yielding contig N50s in excess of 1 Mb.

While the use of long read sequencing technologies facilitates the production of highly

contiguous genome assemblies, scaffold sizes are still much shorter than the length of the

underlying chromosomes. The use of further scaffolding technologies such as genetic linkage

maps, scaffolding based on synteny with genome assemblies from related organisms, as well

as *in vitro* and *in vivo* Hi-C based methods has allowed for the production of assemblies with

chromosome-sized scaffolds. Here, the use of Hi-C based chromatin contact maps allowed for

the placement of 98% of the Nemo v1 assembly length (890.2 of 908.9 Mb) into

chromosomes, yielding a final assembly with a scaffold N50 of 38.4 Mb. This corresponds to

95% of the estimated genome size (938.9 Mb), which suggests that the Nemo v1 assembly is

one of the most complete fish genome assemblies published to date (Fig. 1C). Only the

zebrafish (94%) and Atlantic cod (91%) genome assemblies had a comparably high

proportion of their estimated genome sizes scaffolded into chromosome-length scaffolds (Fig.

1C). It is likely that the use of both PacBio long reads and Hi-C based chromatin contact maps

contributed to the very high proportion of the orange clownfish genome that we were able to

both sequence and assemble into chromosomes.

While assembly contiguity is important, genome completeness with respect to gene content is

also vital for producing a genome assembly that will be utilized by the research community.

We evaluated the completeness of the 27 chromosome-scale assemblies with BUSCO and the

Actinopterygii lineage, which encompasses 4,584 highly-conserved genes. When ranked by

the total of complete (single copy and duplicate) genes, the orange clownfish assembly is the

second most complete, with 4,456 (97.2%) of the orthologs identified (Fig. 2). The top ranked

assembly, Nile tilapia, contains only 9 more of the core set of orthologs such that it contains

4,465 of the orthologs (97.4%). While the assemblies based on PacBio long read technology

are again amongst the most complete, it should also be noted that most of the assemblies

analyzed showed a very high level of completeness.

<u>Genome annotation</u>

To annotate repetitive sequences and transposable elements, we constructed an orange

clownfish-specific library by combining the results of Repeatmodeler, LTRharvest and

TransposonPSI. Duplicate sequences were removed and false positives were identified using

three classification protocols (Censor, Dfam, RepeatClassifier) as well as comparisons to

Uniprot/Swissprot databases. After these filtering steps, we identified 21,644 repetitive

sequences. These sequences, in combination with the zebrafish library of RepBase, were then

used for genome masking with RepeatMasker. This lead to a total of 28% of the assembly

being identified as repetitive (Fig. 3A and Table S4). It was observed that there is a general

trend for increased repeat density towards the ends of chromosome arms (Fig. 3B and S2).

The total fraction of repetitive genomic sequence is in good agreement with other related fish

species (Chalopin, Naville, Plard, Galiana, & Volff, 2015). Similarly, the high fraction of

DNA transposons (~10%) is in line with DNA transposon content in other fish species

(Chalopin et al., 2015) but is substantially higher that what has been reported in mammals

(~3%) (Chalopin et al., 2015; Lander et al., 2001).

Following the characterization of repetitive sequences in the Nemo v1 genome assembly,

gene annotation was performed with the BRAKER1 pipeline, which trained the AUGUSTUS

gene predictor with supplied RNA-seq data, and a successive refinement with the MAKER2

pipeline. We provided BRAKER1 with mapped RNA-seq data from 10 different tissues. This

initial annotation comprised 49,881 genes with 55,273 transcripts. The gene finder models of

SNAP and AUGUSTUS were refined based on the initial annotation, and MAKER2 was then

used to improve the annotation using the new models and the available protein homology and

RNA-seq evidence. The resulting annotation contained 26,606 genes and 35,498 transcripts,

which feature a low mean AED of 0.12, indicating a very good agreement with the provided

evidence. After retaining only genes with evidence support (AED of less than 1) or an

annotated Pfam protein domain, the filtered annotation was comprised of 26,597 genes,

corresponding to 35,478 transcripts (Table 4). This result is broadly consistent with the

average number of genes (23,475) found in the 22 diploid fish species considered in this study

(Table S3). Compared to the initial annotation, genes in the final annotation are 61% longer

(13,049 bp) and encode mRNAs that are 80% longer (17,727 bp). The proportion of the

genome that is covered by coding sequences also increased to 8.1% in the final annotation.

Together with the observed reduction in the gene number by 47%, this indicates a substantial

reduction of likely false positive gene annotations of short length and/or few exons. The gene

density across the 24 chromosomes of our assembly varied from 23.6 genes/Mb (chromosome

21) to 36.5 genes/Mb (chromosome 14), with a genome-wide average of one gene every 29.7

Mb (Table 3). The spatial distribution of genes across all 24 chromosomes is relatively even

(Fig. 1B), with regions of very low gene density presumably corresponding to centromeric

regions. We observed that the longest annotated gene was APERC1_00006329 (26.5 kb),

which encodes the extracellular matrix protein FRAS1, while the gene coding for the longest

protein sequence was APERC1_00011517, which codes for the 18,851 amino acid protein,

Titin. Functional annotation was carried out using Blast2GO and yielded annotations for

22,507 genes (85%) after aligning the protein sequences to the UniProt/Swissprot database

and annotating protein domains with InterProScan.

<u>Identification of orange clownfish-specific genes</u>

To investigate the gene space of the orange clownfish relative to other fishes, we used

OrthoFinder v1.1.4 (Emms & Kelly, 2015) to identify orthologous relationships between the

protein sequences of the orange clownfish and four other fish species (Asian seabass, Nile

tilapia, southern platyfish and zebrafish) from across the teleost phylogenetic tree (Betancur-

R. et al., 2013). The vast majority of sequences (89%) could be assigned to one of 19,838

orthogroups, with the remainder identified as "singlets" with no clear orthologs. We observed

a high degree of overlap of protein sequence sets between all five species, with 75% of all

orthogroups (14,783) shared amongst all species (Fig. 4A). The proteins within these

orthogroups presumably correspond to the core set of teleost genes. Of the 14,783

orthogroups with at least one sequence from each species, a subset of 8,905 orthogroups

contained only a single sequence from each species. The phylogeny obtained from these

single-copy orthologous gene sequences (Fig. 4B) is consistent with the known phylogenetic

tree of teleost fishes (Betancur-R. et al., 2013). Interestingly, we identified a total of 4,429

sequences that are specific to the orange clownfish, 2,293 (49%) of which possess functional

annotations (Fig. 4A). Future investigations will focus on the characterization of these unique

genes and what roles they may play in orange clownfish phenotypic traits.

Conclusion

Here, we present a reference-quality genome assembly of the iconic orange clownfish, *A.*

percula. We sequenced the genome to a depth of 121X with PacBio long reads and performed

a primary assembly with these reads utilizing the Falcon_Unzip algorithm. The primary

assembly was polished to yield an initial assembly of 903.6 Mb with a contig N50 value of

1.86 Mb. These contigs were then assembled into chromosome-sized scaffolds using Hi-C

chromatin contact maps, followed by gap-filling with the PacBio reads, to produce the final

reference assembly, Nemo v1. The Nemo v1 assembly is highly contiguous, with contig and

scaffold N50s of 3.12 and 38.4 Mb, respectively. The use of Hi-C chromatin contact maps

allowed us to scaffold 890.2 Mb (98%) of the 908.2 Mb final assembly into the 24

chromosomes of the orange clownfish. An analysis of the core set of Actinopterygii genes

suggests that our assembly is nearly complete, containing 97% of the core set of highly

conserved genes. The Nemo v1 assembly was annotated with 26,597 genes with an average

AED score of 0.12, suggesting that most gene models are highly supported.

The high-quality Nemo v1 reference genome assembly described here will facilitate the use of

this now genome-enabled model species to investigate ecological, environmental and

evolutionary aspects of reef fishes. To assist the research community, we have created the

Nemo Genome DB database, http://nemogenome.org/ (Fig. 5), where researchers can access,

mine and visualize the genomic and transcriptomic resources of the orange clownfish.

Acknowledgements

This study was supported by the Competitive Research Funds OCRF-2014-CRG3-62140408 from the King Abdullah University of Science and Technology (KAUST) to T.R., M.L.B. and P.L.M., as well as KAUST baseline support to M.L.B., M.A., T.G. and T.R. This project was completed under JCU Ethics A1233 and A1415. We thank Dr. Jennifer Donelson and staff at JCU's MARFU facility for assistance with animal husbandry, Dr. Susanne Sprungala for DNA extraction for Illumina library preparation, KAUST BCL for the PacBio sequencing, Dr. Hicham Mansour for sequencing advice and Dr. Rita Bartossek for the PacBio library preparations. We thank Dr. Salim Bougouffa for stimulating discussions. We also acknowledge Mr. Tane Sinclair-Taylor for providing the photograph of the orange clownfish (Fig. 1A). This paper is dedicated to our good friend and colleague, Dr. Sylvain Foret.

Author contributions

R.L. and D.J.L. designed and performed the computational analysis. R.L., T.R., C.S. and D.J.L. interpreted the results. H.O., K.M. and T.G. created the database. C.T.M. and S.F. produced sequencing libraries. R.L., D.J.L, T.R., P.L.M., M.L.B., M.A. and D.J.M. wrote the manuscript and all authors approved the final version. T.R. supervised the project.

References

Almany, G. R., Planes, S., Thorrold, S. R., Berumen, M. L., Bode, M., Saenz-Agudelo, P., …

Jones, G. P. (2017). Larval fish dispersal in a coral-reef seascape. *Nature Ecology &*

Evolution, *1*(6), 148.

Arai, R., & Inoue, M. (1976). Chromosomes of seven species of Pomacentridae and two

species of Acanthuridae from Japan. *Bulletin of the National Museum of Nature and*

Science, Series A, *2*, 73–78.

Arai, R., Inoue, M., & Ida, H. (1976). Chromosomes of four species of coral fishes from

Japan. *Bulletin of the National Museum of Nature and Science, Series A*, *2*, 137–141.

Bao, W., Kojima, K. K., & Kohany, O. (2015). Repbase Update, a database of repetitive

elements in eukaryotic genomes. *Mobile DNA*, *6*(1), 11.

Berumen, M. L., Almany, G. R., Planes, S., Jones, G. P., Saenz-Agudelo, P., & Thorrold, S.

R. (2012). Persistence of self-recruitment and patterns of larval connectivity in a marine

protected area network. *Ecology and Evolution*, *2*(2), 444–452.

Betancur-R., R., Broughton, R. E., Wiley, E. O., Carpenter, K., López, J. A., Li, C., … Ortí,

G. (2013). The tree of life and a new classification of bony fishes. *PLoS Currents*, *5*,

ecurrents.tol.53ba26640df0ccaee75bb165c8c26288.

Bickhart, D. M., Rosen, B. D., Koren, S., Sayre, B. L., Hastie, A. R., Chan, S., … Smith, T. P.

L. (2017). Single-molecule sequencing and chromatin conformation capture enable *de*

novo reference assembly of the domestic goat genome. *Nature Genetics*, *49*(4), 643–650.

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for

Illumina sequence data. *Bioinformatics*, *30*(15).

Bradnam, K. R., Fass, J. N., Alexandrov, A., Baranay, P., Bechner, M., Birol, I., … Korf, I. F.

(2013). Assemblathon 2: evaluating *de novo* methods of genome assembly in three

vertebrate species. *GigaScience*, *2*(1), 10.

Buels, R., Yao, E., Diesh, C. M., Hayes, R. D., Munoz-Torres, M., Helt, G., … Holmes, I. H.

(2016). JBrowse: a dynamic web platform for genome visualization and analysis.

Genome Biology, *17*(1), 66.

Burton, J. N., Adey, A., Patwardhan, R. P., Qiu, R., Kitzman, J. O., & Shendure, J. (2013).

Chromosome-scale scaffolding of de novo genome assemblies based on chromatin

interactions. *Nature Biotechnology*, *31*(12), 1119–1125.

Buston, P. M. (2003). Mortality is associated with social rank in the clown anemonefish

(*Amphiprion percula*). *Marine Biology*, *143*(4), 811–815.

Buston, P. M., Bogdanowicz, S. M., Wong, A., & Harrison, R. G. (2007). Are clownfish

groups composed of close relatives? An analysis of microsatellite DNA variation in

Amphiprion percula. Molecular Ecology, *16*(17), 3671–3678.

Buston, P. M., & García, M. B. (2007). An extraordinary life span estimate for the clown

anemonefish *Amphiprion percula. Journal of Fish Biology*, *70*(6), 1710–1719.

Buston, P. M., & Wong, M. (2014). Why some animals forgo reproduction in complex

societies. *American Scientist*, *102*(4), 290.

Campbell, M. S., Holt, C., Moore, B., & Yandell, M. (2014). Genome Annotation and

Curation Using MAKER and MAKER-P. In *Current Protocols in Bioinformatics* (Vol.

48, p. 4.11.1-4.11.39). Hoboken, NJ, USA: John Wiley & Sons, Inc.

Chaisson, M. J., & Tesler, G. (2012). Mapping single molecule sequencing reads using basic

local alignment with successive refinement (BLASR): application and theory. *BMC*

Bioinformatics, *13*(1), 238.

Chalopin, D., Naville, M., Plard, F., Galiana, D., & Volff, J.-N. (2015). Comparative analysis

of transposable elements highlights mobilome diversity and evolution in vertebrates.

Genome Biology and Evolution, *7*(2), 567–580.

Chikhi, R., & Medvedev, P. (2014). Informed and automated k-mer size selection for genome

assembly. *Bioinformatics*, *30*(1), 31–37.

Chin, C.-S., Alexander, D. H., Marks, P., Klammer, A. A., Drake, J., Heiner, C., … Korlach,

J. (2013). Nonhybrid, finished microbial genome assemblies from long-read SMRT

sequencing data. *Nature Methods*, *10*(6), 563–569.

Chin, C.-S., Peluso, P., Sedlazeck, F. J., Nattestad, M., Concepcion, G. T., Clum, A., …

Schatz, M. C. (2016). Phased diploid genome assembly with single-molecule real-time

sequencing. *Nature Methods*, *13*, 1050–1054.

Dixson, D. L. (2012). Predation risk assessment by larval reef fishes during settlement-site

selection. *Coral Reefs*, *31*(1), 255–261.

Dixson, D. L., Jones, G. P., Munday, P. L., Planes, S., Pratchett, M. S., Srinivasan, M., …

Thorrold, S. R. (2008). Coral reef fish smell leaves to find island homes. *Proceedings.*

Biological Sciences, *275*(1653), 2831–2839.

Dixson, D. L., Jones, G. P., Munday, P. L., Planes, S., Pratchett, M. S., & Thorrold, S. R.

(2014). Experimental evaluation of imprinting and the role innate preference plays in habitat selection in a coral reef fish. *Oecologia*, *174*(1), 99–107.

Dixson, D. L., Munday, P. L., & Jones, G. P. (2010). Ocean acidification disrupts the innate ability of fish to detect predator olfactory cues. *Ecology Letters*, *13*(1), 68–75.

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., … Gingeras, T. R. (2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*, *29*(1), 15–21.

Edgar, R. C. (2010). Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*, *26*(19), 2460–2461.

Ellinghaus, D., Kurtz, S., & Willhoeft, U. (2008). LTRharvest, an efficient and flexible software for *de novo* detection of LTR retrotransposons. *BMC Bioinformatics*, *9*(1), 18.

Elliott, J. K., & Mariscal, R. N. (2001). Coexistence of nine anemonefish species: differential host and habitat utilization, size and recruitment. *Marine Biology*, *138*(1), 23–36.

Emms, D. M., & Kelly, S. (2015). OrthoFinder: solving fundamental biases in whole genome comparisons dramatically improves orthogroup inference accuracy. *Genome Biology*, *16*(1), 157.

Emms, D. M., & Kelly, S. (2017). STRIDE: Species tree root inference from gene duplication events. *Molecular Biology and Evolution*, *34*(12), 3267–3278.

English, A. C., Richards, S., Han, Y., Wang, M., Vee, V., Qu, J., … Gibbs, R. A. (2012). Mind the Gap: Upgrading genomes with Pacific Biosciences RS long-read sequencing technology. *PLoS ONE*, *7*(11), e47768.

Fautin, D. G. (1991). The anemonefish symbiosis: what is known and what is not. *Symbiosis*, *10*, 23–46.

Fautin, D. G., & Allen, G. R. (1997). Life history of Anemonefishes. In: Anemone fishes and their host sea anemones. *Western Australian Museum, Perth*, 1–142.

Gotz, S., Garcia-Gomez, J. M., Terol, J., Williams, T. D., Nagaraj, S. H., Nueda, M. J., … Conesa, A. (2008). High-throughput functional annotation and data mining with the Blast2GO suite. *Nucleic Acids Research*, *36*(10), 3420–3435.

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., … Regev, A. (2011). Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nature Biotechnology*, *29*(7), 644–652.

Gregory, T. R. (2018). Animal Genome Size Database. Retrieved from http://www.genomesize.com

Gremme, G., Steinbiss, S., & Kurtz, S. (2013). GenomeTools: a comprehensive software

library for efficient processing of structured genome annotations. *IEEE/ACM*

Transactions on Computational Biology and Bioinformatics, *10*(3), 645–656.

Gurevich, A., Saveliev, V., Vyahhi, N., & Tesler, G. (2013). QUAST: quality assessment tool

for genome assemblies. *Bioinformatics*, *29*(8), 1072–1075.

Haas, B. J. (2018). TransposonPSI. Retrieved from http://transposonpsi.sourceforge.net/

Hardie, D. C., & Hebert, P. D. (2004). Genome-size evolution in fishes. *Canadian Journal of*

Fisheries and Aquatic Sciences, *61*(9), 1636–1646.

Herbert, N. A., Bröhl, S., Springer, K., & Kunzmann, A. (2017). Clownfish in hypoxic

anemones replenish host O_2 at only localised scales. *Scientific Reports*, *7*(1), 6547.

Hess, S., Wenger, A. S., Ainsworth, T. D., & Rummer, J. L. (2015). Exposure of clownfish

larvae to suspended sediment levels found on the Great Barrier Reef: Impacts on gill

structure and microbiome. *Scientific Reports*, *5*(1), 10561.

Hoff, K. J., Lange, S., Lomsadze, A., Borodovsky, M., & Stanke, M. (2016). BRAKER1:

Unsupervised RNA-Seq-based genome annotation with GeneMark-ET and

AUGUSTUS. *Bioinformatics*, *32*(5), 767–769.

Holbrook, S. J., & Schmitt, R. J. (2005). Growth, reproduction and survival of a tropical sea

anemone (Actiniaria): benefits of hosting anemonefish. *Coral Reefs*, *24*(1), 67–73.

Holt, C., & Yandell, M. (2011). MAKER2: an annotation pipeline and genome-database

management tool for second-generation genome projects. *BMC Bioinformatics*, *12*(1),

491.

Hoskins, R. A., Carlson, J. W., Kennedy, C., Acevedo, D., Evans-Holm, M., Frise, E., …

Celniker, S. E. (2007). Sequence finishing and mapping of *Drosophila melanogaster*

heterochromatin. *Science (New York, N.Y.)*, *316*(5831), 1625–1628.

Jarrold, M. D., Humphrey, C., McCormick, M. I., & Munday, P. L. (2017). Diel CO2 cycles

reduce severity of behavioural abnormalities in coral reef fish under ocean acidification.

Scientific Reports, *7*(1), 10153.

Jurka, J., Klonowski, P., Dagman, V., & Pelton, P. (1996). CENSOR--a program for

identification and elimination of repetitive elements from DNA sequences. *Computers &*

Chemistry, *20*(1), 119–121.

Korf, I. (2004). Gene finding in novel genomes. *BMC Bioinformatics*, *5*, 59.

Lander, E. S., Linton, L. M., Birren, B., Nusbaum, C., Zody, M. C., Baldwin, J., …

International Human Genome Sequencing Consortium. (2001). Initial sequencing and
analysis of the human genome. *Nature*, *409*(6822), 860–921.

Li, H., & Durbin, R. (2010). Fast and accurate long-read alignment with Burrows-Wheeler
transform. *Bioinformatics*, *26*(5), 589–595.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., … 1000 Genome
Project Data Processing Subgroup. (2009). The sequence alignment/map format and
SAMtools. *Bioinformatics*, *25*(16), 2078–2079.

Li, J., Chen, X., Kang, B., & Liu, M. (2015). Mitochondrial DNA Genomes Organization and
Phylogenetic Relationships Analysis of Eight Anemonefishes (Pomacentridae:
Amphiprioninae). *PLoS ONE*, *10*(4), e0123894.

Litsios, G., Pearman, P. B., Lanterbecq, D., Tolou, N., & Salamin, N. (2014). The radiation of
the clownfishes has two geographical replicates. *Journal of Biogeography*, *41*(11),
2140–2149.

Litsios, G., & Salamin, N. (2014). Hybridisation and diversification in the adaptive radiation
of clownfishes. *BMC Evolutionary Biology*, *14*(1), 245.

Manassa, R. P., Dixson, D. L., McCormick, M. I., & Chivers, D. P. (2013). Coral reef fish
incorporate multiple sources of visual and chemical information to mediate predation
risk. *Animal Behaviour*, *86*(4), 717–722.

Marçais, G., & Kingsford, C. (2011). A fast, lock-free approach for efficient parallel counting
of occurrences of k-mers. *Bioinformatics*, *27*(6), 764–770.

Marcionetti, A., Rossier, V., Bertrand, J. A. M., Litsios, G., & Salamin, N. (2018). First draft
genome of an iconic clownfish species (*Amphiprion frenatus*). *Molecular Ecology*
Resources.

McLeod, I. M., Rummer, J. L., Clark, T. D., Jones, G. P., McCormick, M. I., Wenger, A. S.,
& Munday, P. L. (2013). Climate change and the performance of larval coral reef fishes:
the interaction between temperature and food availability. *Conservation Physiology*,
1(1), cot024.

Mebs, D. (2009). Chemical biology of the mutualistic relationships of sea anemones with fish
and crustaceans. *Toxicon : Official Journal of the International Society on Toxinology*,
54(8), 1071–1074.

Molina, W. F., & Galetti, P. M. (2004). Karyotypic changes associated to the dispersive
potential on Pomacentridae (Pisces, Perciformes). *Journal of Experimental Marine*

Biology and Ecology, *309*(1), 109–119.

Munday, P. L., Dixson, D. L., Donelson, J. M., Jones, G. P., Pratchett, M. S., Devitsina, G. V,

& Døving, K. B. (2009). Ocean acidification impairs olfactory discrimination and

homing ability of a marine fish. *Proceedings of the National Academy of Sciences of the*

United States of America, *106*(6), 1848–1852.

Nilsson, G. E., Dixson, D. L., Domenici, P., McCormick, M. I., Sørensen, C., Watson, S.-A.,

& Munday, P. L. (2012). Near-future carbon dioxide levels alter fish behaviour by

interfering with neurotransmitter function. *Nature Climate Change*, *2*(3), 201–204.

Peichel, C. L., Sullivan, S. T., Liachko, I., & White, M. A. (2017). Improvement of the

Threespine Stickleback genome using a Hi-C-Based proximity-guided assembly. *The*

Journal of Heredity, *108*(6), 693–700.

Pertea, M., Kim, D., Pertea, G. M., Leek, J. T., & Salzberg, S. L. (2016). Transcript-level

expression analysis of RNA-seq experiments with HISAT, StringTie and Ballgown.

Nature Protocols, *11*(9), 1650–1667.

Pinsky, M. L., Saenz-Agudelo, P., Salles, O. C., Almany, G. R., Bode, M., Berumen, M. L.,

… Planes, S. (2017). Marine dispersal scales are congruent over evolutionary and

ecological time. *Current Biology*, *27*(1), 149–154.

Planes, S., Jones, G. P., & Thorrold, S. R. (2009). Larval dispersal connects fish populations

in a network of marine protected areas. *Proceedings of the National Academy of Sciences*

of the United States of America, *106*(14), 5693–5697.

Saenz-Agudelo, P., Jones, G. P., Thorrold, S. R., & Planes, S. (2011). Detrimental effects of

host anemone bleaching on anemonefish populations. *Coral Reefs*, *30*(2), 497–506.

Salles, O. C., Pujol, B., Maynard, J. A., Almany, G. R., Berumen, M. L., Jones, G. P., …

Planes, S. (2016). First genealogy for a wild marine fish population reveals

multigenerational philopatry. *Proceedings of the National Academy of Sciences*, *113*(46),

13245–13250.

Schmiege, P. F. P., D'Aloia, C. C., & Buston, P. M. (2017). Anemonefish personalities

influence the strength of mutualistic interactions with host sea anemones. *Marine*

Biology, *164*(1), 24.

Scott, A., & Dixson, D. L. (2016). Reef fishes can recognize bleached habitat during

settlement: sea anemone bleaching alters anemonefish host selection. *Proceedings.*

Biological Sciences, *283*(1831), 20152694.

Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V, & Zdobnov, E. M. (2015).

BUSCO: assessing genome assembly and annotation completeness with single-copy

orthologs. *Bioinformatics*, *31*(19), 3210–3212.

Simpson, S. D., Munday, P. L., Wittenrich, M. L., Manassa, R., Dixson, D. L., Gagliano, M.,

& Yan, H. Y. (2011). Ocean acidification erodes crucial auditory behaviour in a marine

fish. *Biology Letters*, *7*(6), 917–920.

Smit, A. F. A., & Hubley, R. (2008). RepeatModeler Open-1.0.

Smit, A. F. A., Hubley, R., & Green, P. (2010). RepeatMasker Open-4.0.

Stanke, M., Keller, O., Gunduz, I., Hayes, A., Waack, S., & Morgenstern, B. (2006).

AUGUSTUS: *ab initio* prediction of alternative transcripts. *Nucleic Acids Research*,

34(Web Server), W435–W439.

Steinbiss, S., Willhoeft, U., Gremme, G., & Kurtz, S. (2009). Fine-grained annotation and

classification of *de novo* predicted LTR retrotransposons. *Nucleic Acids Research*,

37(21), 7002–7013.

Stillman, J. H., & Armstrong, E. (2015). Genomics Are Transforming Our Understanding of

Responses to Climate Change. *BioScience*, *65*(3), 237–246.

Supiwong, W., Tanomtong, A., Pinthong, K., Kaewmad, P., Poungnak, P., & Jangsuwan, N.

(2015). The first chromosomal characteristics of nucleolar organizer regions and

karyological analysis of pink anemonefish, *Amphiprion perideraion* (Perciformes,

Amphiprioninae). *Cytologica*, *80*(3), 271–278.

Szczebak, J. T., Henry, R. P., Al-Horani, F. A., & Chadwick, N. E. (2013). Anemonefish

oxygenate their anemone hosts at night. *Journal of Experimental Biology*, *216*(6), 970–

976.

Takai, A., & Kosuga, S. (2007). Karyotypes and banded chromosomal features in two

anemonefishes (Pomacentridae, Perciformes). *Chromosome Science*, *10*(3), 71–74.

Tan, M. H., Austin, C. M., Hammer, M. P., Lee, Y. P., Croft, L. J., & Gan, H. M. (2018).

Finding Nemo: Hybrid assembly with Oxford Nanopore and Illumina reads greatly

improves the Clownfish (*Amphiprion ocellaris*) genome assembly. *GigaScience*, gix137.

Tanomtong, A., Supiwong, W., Chaveerach, A., Khakhong, S., Tanee, T., & Sanoamuang, L.

(2012). First report of chromosome analysis of saddleback anemonefish, *Amphiprion*

polymnus (*Perciformes*, *Amphiprioninae*), in Thailand. *Cytologica*, *77*(4), 441–446.

Ter-Hovhannisyan, V., Lomsadze, A., Chernoff, Y. O., & Borodovsky, M. (2008). Gene

832  prediction in novel fungal genomes using an *ab initio* algorithm with unsupervised

833  training. *Genome Research*, *18*(12), 1979–1990.

834 Vurture, G. W., Sedlazeck, F. J., Nattestad, M., Underwood, C. J., Fang, H., Gurtowski, J., &

835  Schatz, M. C. (2017). GenomeScope: fast reference-free genome profiling from short

836  reads. *Bioinformatics*, *33*(14), 2202–2204.

837 Wenger, A. S., McCormick, M. I., Endo, G. G. K., McLeod, I. M., Kroon, F. J., & Jones, G.

838  P. (2014). Suspended sediment prolongs larval development in a coral reef fish. *The*

839  *Journal of Experimental Biology*, *217*(Pt 7), 1122–1128.

840 Wheeler, T. J., Clements, J., Eddy, S. R., Hubley, R., Jones, T. A., Jurka, J., … Finn, R. D.

841  (2012). Dfam: a database of repetitive DNA based on profile hidden Markov models.

842  *Nucleic Acids Research*, *41*, D70–D82.

843 Wittenrich, M. L., Turingan, R. G., & Creswell, R. L. (2007). Spawning, early development

844  and first feeding in the gobiid fish *Priolepis nocturna*. *Aquaculture*, *270*(1–4), 132–141.

845 Wong, M., Uppaluri, C., Medina, A., Seymour, J., & Buston, P. M. (2016). The four elements

846  of within-group conflict in animal societies: an experimental test using the clown

847  anemonefish, *Amphiprion percula*. *Behavioral Ecology and Sociobiology*, *70*(9), 1467–

848  1475.

849 Wood, D. E., & Salzberg, S. L. (2014). Kraken: ultrafast metagenomic sequence classification

850  using exact alignments. *Genome Biology*, *15*(3), R46.

Data Accessibility

The assembled and annotated genome as well as the raw PacBio reads and Illumina reads are

available at the Nemo Genome DB (http://nemogenome.org). Furthermore, the assembled

genome will be available on GenBank as BioProject PRJNA436093 and BioSample accession

SAMN08615572. All raw sequencing data described in this study will be available via the

NCBI Sequencing Read Archive.

Tables and Figures

46.12 Mb

Fig. 1 (A) The iconic orange clownfish (*A. percula*). **(B)** Gene density on the 24

chromosomes, plotted in 100 kb windows. Chromosomes are ordered by size, as indicated on

the left axis in Mb. **(C)** Contiguity (x-axis) and genome assembly completeness (y-axis) of the

orange clownfish, and the 26 previously published, chromosome-scale fish genome

assemblies. Details and statistics of the 27 assemblies are presented in Table S3.

Fig. 2 Genome assembly completeness of all published chromosome-scale fish genome

assemblies, as measured by the proportion of the BUSCO set of core genes detected in each

assembly. Genome assemblies on the y-axis are sorted by the sum of single copy and

duplicated BUSCO genes.

Fig. 3 Repeat content of the orange clownfish genome assembly. **(A)** Repeat content of the whole genome as classified into transposable elements and repetitive sequences. **(B)** Spatial distribution of the four main identified classes of transposable elements on chromosome 1. Transposable element spatial distribution for chromosomes 2-24 is show in Fig S2. Detailed transposable element content is shown in Table S4.

Fig. 4 (A) The overlap of orthologous gene families of the orange clownfish, southern platyfish, Nile tilapia, zebrafish and Asian seabass. The total number of orthogroups (nOG) followed by the number of genes assigned to these groups is provided below the species name. The number of species-specific orthogroups (nSOG) and the respective number of genes is also indicated, followed by the number of genes not assigned to any orthogroups. **(B)** The inferred phylogenetic tree based on the ortholog groups that contain a single gene from each species.

Fig. 5 (A) Front page of the Nemo Genome DB database, which is a portal to access the data described in this manuscript and is accessible at www.nemogenome.org. **(B)** Genome viewer representation of the Titin gene.

916 **Table 1** Contig statistics for the preliminary candidate-assemblies

Assembly	Length (Mb)	Number	N50 (Mb)	L50	Longest (Mb)	Missing genes (number (%))
A1	950.4	4,874	1.024	254	9.59	148 (3.23)
A2	945.4	4,374	1.040	251	6.67	156 (3.30)
A3	926.5	3,629	1.070	236	7.21	140 (3.05)
A4	921.8	2,829	1.380	184	8.16	134 (2.92)
A5	883.9	1,017	1.469	167	10.24	146 (3.18)
A6	902.2	2,204	1.401	174	12.38	134 (2.92)
A7	920.7	2,473	1.801	138	15.84	132 (2.88)
A8	924.6	2,629	1.742	143	16.51	139 (3.03)
A9	924.9	2,638	1.742	143	16.51	140 (3.05)
A10	917.1	2,368	1.648	140	10.21	146 (3.18)
A11	899.9	2,049	1.571	160	9.07	151 (3.29)
A12	908.8	2,086	1.602	142	10.21	143 (3.12)

917
918
919
920

925 **Table 2** Assembly statistics of the orange clownfish genome assemblies
926

	A7	A7-phased	A7-phased-polished	A7-PGA	Nemo v1
Technology					
Falcon	✓	-	-	-	-
Falcon_Unzip	-	✓	✓	✓	✓
PacBio	✓	✓	✓	✓	✓
Quiver	-	-	✓	✓	✓
Hi-C maps	-	-	-	✓	✓
PBJelly	-	-	-	-	✓
Contigs					
Length (Mb)	920.7	905.0	903.6	903.6	908.9
Number	2,473	1,505	1,414	1,414	1,045
N50 length (Mb)	1.80	1.85	1.86	1.86	3.12
L50 count	138	135	134	134	84
Longest (Mb)	15.84	15.83	15.85	15.9	16.6
No. Scaffolded	-	-	-	1,073	704
Scaffolds					
Length (Mb)	-	-	-	903.7	908.9
Number	-	-	-	365	365
N50 length (Mb)	-	-	-	38.1	38.4
L50 count	-	-	-	12	12
Longest (Mb)	-	-	-	45.8	46.1
Ns	-	-	-	104,900	32,395
Number of gaps	-	-	-	1,049	680
Chromosomes					
Length in chr (Mb)	-	-	-	885.4	890.2
% assembly in chr	-	-	-	98.0%	97.9%
% assembly not in chr	-	-	-	2.0%	2.1%
% of predicted genome size in chr	-	-	-	94.3%	94.8%

928 * Predicted genome size is 938.88 Mb (Hardie & Hebert, 2004).

930 **Table 3** Chromosome metrics before and after polishing of the final assembly

Chromosome	A7-PGA assembly		Nemo v1 assembly			
	Contigs	Length (Mb)	Contigs	Length (Mb)	Genes	Gene density (genes/Mb)
1	57	45.8	31	46.1	1,091	23.8
2	41	43.3	31	43.4	1,132	26.1
3	55	43.2	28	43.4	1,395	32.3
4	47	42.0	29	42.2	1,259	30.0
5	32	40.5	31	40.6	1,303	32.2
6	44	40.4	24	40.6	1,337	33.1
7	37	40.2	32	40.4	1,324	32.9
8	42	39.3	26	39.4	1,276	32.5
9	47	39.0	25	39.2	1,083	27.8
10	55	38.3	38	38.6	1,339	35.0
11	40	38.3	23	38.5	1,037	27.1
12	48	38.1	23	38.4	1,067	28.0
13	30	37.6	20	37.7	1,014	27.0
14	33	37.3	33	37.4	1,362	36.5
15	45	37.3	22	37.4	1,091	29.2
16	77	36.3	50	36.6	1,018	28.0
17	35	35.2	23	35.4	987	28.0
18	40	34.9	32	35.1	1,126	32.3
19	53	34.0	35	34.2	1,062	31.2
20	46	33.4	31	33.7	1,132	33.9
21	40	30.7	21	30.8	725	23.6
22	29	29.6	20	29.8	786	26.6
23	32	27.2	23	27.4	904	33.2
24	68	23.4	53	23.7	723	30.9
In chr:	1,073	885.4	704	890.2	26,309	Ave: 29.7
Not in chr:	341	18.4	341	18.8	288	15.3
Total:	1,414	903.7	1,045	908.8	26,597	Ave: 29.3

933 **Table 4** Gene annotation statistics

	Initial BRAKER1	Final MAKER2
Genes	49,881	26,597
mRNAs	55,273	35,478
Exons	391,637	463,688
Introns	336,364	428,210
CDSs	55,273	35,478
Overlapping genes	2,407	1,852
Contained genes	744	463
Longest gene	264,684	264,684
Longest mRNA	264,684	264,684
Mean gene length	8,097	13,049
Mean mRNA length	9,841	17,727
% of genome covered by genes	44.4	38.2
% of genome covered by CDS	7.5	8.1
Exons per mRNA	7	13
Introns per mRNA	6	12
BUSCO		
Completeness	95.94%	96.25%
Complete	4,398	4,412
Single-Copy	3,588	3,888
Duplicated	810	524
Fragmented	138	96
Missing	48	76
Total	4,584	4,584