Ground-truthing environmental DNA (eDNA) metabarcoding for ecological hypothesis testing at the pondscape

Lynsey R. Harper[1]*, Lori Lawson Handley[1], Christoph Hahn[1,2], Neil Boonham[3,4], Helen C. Rees[5], Erin Lewis[3], Ian P. Adams[3], Peter Brotherton[6], Susanna Phillips[6] and Bernd Hänfling[1]

[1] School of Environmental Sciences, University of Hull, Hull, HU6 7RX, UK

E-mails: L.Harper@2015.hull.ac.uk, L.Lawson-Handley@hull.ac.uk, B.Haenfling@hull.ac.uk

[2] Institute of Zoology, University of Graz, Graz, Styria, Austria

E-mail: christoph.hahn@uni-graz.at

[3] Fera Science Ltd (Fera), Sand Hutton, York, YO14 1LZ, UK

E-mails: Erin.Lewis@fera.co.uk, Ian.Adams@fera.co.uk

[4] Newcastle University, Newcastle upon Tyne, NE1 7RU, UK

E-mail: neil.boonham@ncl.ac.uk

[5] ADAS, School of Veterinary Medicine and Science, The University of Nottingham, Sutton Bonington Campus, Leicestershire, LE12 5RD, UK

E-mail: Helen.Rees@adas.co.uk

[6] Natural England, Peterborough, PE1 1NG, UK

E-mails: Peter.Brotherton@naturalengland.org.uk, susanna.phillips@naturalengland.org.uk

Statement of authorship: B.H., L.R.H., L.L.H and N.B. conceived and designed the study. H.C.R. and N.B. contributed samples for processing. L.R.H. performed laboratory work and analysed the data. I.P.A. and E.L. offered advice on and supervised sequencing. C.H. assisted with bioinformatics analysis. P.B. and S.P. contributed datasets for analysis. L.R.H. wrote the manuscript, which all authors revised.

Running title: Hypothesis testing with eDNA metabarcoding

Article type: Letters

Abstract word count: 148 words

Main text word count: 5190 words

Number of references: 66

Number of figures, tables and boxes: 6

*Corresponding author: Lynsey R. Harper

School of Environmental Sciences, University of Hull, Hull, HU6 7RX, UK

E-mail: L.Harper@2015.hull.ac.uk

Tel: 01482 465804

Abstract

Environmental DNA (eDNA) metabarcoding is revolutionising biodiversity

monitoring but has unrealised potential for ecological hypothesis testing. Here, we

ground-truth eDNA metabarcoding for describing vertebrate communities from 532

UK ponds, examine associations between great crested newts (*Triturus cristatus*,

GCN) and other vertebrates, and investigate factors determining GCN occurrence and

vertebrate species richness at the pondscape. GCN occurrence was positively

correlated with amphibian and waterfowl species richness, where GCN had strong

associations with smooth newt (*Lissotriton vulgaris*) and common toad (*Bufo bufo*),

but negatively correlated to fish species richness, specifically stickleback presence.

Both GCN occupancy and species richness correlated with the GCN Habitat

Suitability Index score, supporting its application to GCN survey. We reaffirm

reported associations (e.g. GCN preference for deeper ponds) but also provide novel

insights, including a negative effect of pond outflow on GCN. Our findings

demonstrate the prospects of eDNA metabarcoding for hypothesis testing at

unprecedented scales.

INTRODUCTION

Environmental DNA (eDNA) analysis offers ecologists exceptional power to detect organisms within and across ecosystems. DNA secreted or excreted by organisms into their environment can be sampled and analysed using different approaches to reveal the distribution of single or multiple species (Rees *et al.* 2014; Lawson Handley 2015). When combined with high-throughput sequencing (i.e. eDNA metabarcoding), eDNA can yield efficient, comprehensive assessments of entire communities (Deiner *et al.* 2017), providing a step change in biodiversity monitoring (Hering *et al.* 2018). eDNA metabarcoding has untapped potential to test ecological hypotheses by enabling biodiversity monitoring at phenomenal scales with minimal impact to communities under investigation. Although this tool has been used to estimate species richness and assess diversity along environmental gradients (e.g. Hänfling *et al.* 2016; Kelly *et al.* 2016; Olds *et al.* 2016; Evans *et al.* 2017), its applications in community ecology are relatively unexplored.

Aquatic ecosystems are highly suited to eDNA studies (Muha *et al.* 2017) as eDNA exists in multiple states with rapid modes of transport and degradation, increasing detectability of contemporary biodiversity (Rees *et al.* 2014; Barnes & Turner 2015). Lentic systems provide further opportunities for eDNA research, being discrete water bodies with variable physicochemical properties that do not experience flow dynamics (Muha *et al.* 2017). Ponds in particular have enormous biodiversity and experimental virtue that has not been maximised in previous eDNA metabarcoding assessments of this habitat (Valentini *et al.* 2016; Bálint *et al.* 2017; Evans *et al.* 2017; Klymus *et al.* 2017; Ushio *et al.* 2017). These small and abundant water bodies span broad ecological gradients (De Meester *et al.* 2005) and comprise

pondscapes - a network of ponds and their surrounding terrestrial habitat (Hill *et al.*

2018). Pondscapes contribute substantially to aquatic and non-aquatic biodiversity

across spatial scales, with ponds supporting many rare and protected species in

fragmented landscapes (De Meester *et al.* 2005; Biggs *et al.* 2016; Hill *et al.* 2018).

Consequently, ponds are model systems for experimental validation and examination

of biogeographical patterns (De Meester *et al.* 2005). Habitat complexity and tools

required for different taxa with associated bias (Evans *et al.* 2017) and cost (Valentini

et al. 2016) once hindered exhaustive sampling of pond biodiversity (Hill *et al.* 2018),

but eDNA metabarcoding may overcome these barriers.

In the UK, the threatened great crested newt (*Triturus cristatus,* hereafter

GCN) is a flagship species for pond conservation. The extensive literature on the

GCN provides an excellent opportunity to ground-truth ecological patterns revealed

by eDNA metabarcoding. Both biotic (e.g. breeding substrate, prey, and predators)

and abiotic (e.g. pond area, depth, and temperature) factors are known to influence

GCN breeding success (Langton *et al.* 2001). The GCN Habitat Suitability Index

(HSI) (Oldham *et al.* 2000; ARG-UK 2010) accounts for these factors using 10

suitability indices that are scored and combined to calculate a decimal score between

0 and 1 (where 1 = excellent habitat). Larvae are susceptible to fish and waterfowl

predation (Rannap & Briggs 2006; Skei *et al.* 2006; Hartel *et al.* 2010), and adults

reportedly avoid ponds containing three-spined stickleback (*Gasterosteus aculeatus*)

(McLee & Scaife 1992), ninespine stickleback (*Pungitius pungitius*), crucian carp

(*Carassius carassius*), and common carp (*Carassius carpio*) (Rannap *et al.* 2009a, b).

Conversely, the GCN and smooth newt (*Lissotriton vulgaris*) prefer similar habitat

and often co-occur (Rannap & Briggs 2006; Skei *et al.* 2006; Rannap *et al.* 2009a;

Denoël *et al.* 2013). GCNs thrive in ponds with good water quality as indicated by

diverse macroinvertebrate communities (Oldham *et al.* 2000; Rannap *et al.* 2009a),

and water clarity is important for breeding displays, foraging success, and egg

survival (Rannap & Briggs 2006; Skei *et al.* 2006). Pond networks encourage GCN

occupancy (Joly *et al.* 2001; Rannap *et al.* 2009a; Hartel *et al.* 2010; Denoël *et al.*

2013), but larger area discourages presence (Joly *et al.* 2001). Ponds with heavy

shading (Vuorio *et al.* 2013) or dense macrophyte cover (Rannap & Briggs 2006; Skei

et al. 2006; Hartel *et al.* 2010) are unlikely to support viable populations. GCNs also

depend on terrestrial habitat, preferring open, semi-rural pondscapes (Denoël *et al.*

2013) containing pasture, extensively grazed and rough grassland, scrub, and

coniferous and deciduous woodland (Oldham *et al.* 2000; Rannap & Briggs 2006;

Rannap *et al.* 2009a; Gustafson *et al.* 2011; Vuorio *et al.* 2013).

We assessed vertebrate communities at the pondscape using a dataset

generated by eDNA metabarcoding for over 500 ponds with comprehensive

environmental metadata. We evaluated eDNA metabarcoding as a tool for ecological

hypothesis testing, and compared its outputs to previous results generated by

established methods. Specifically, we aimed to identify biotic (community presence-

absence data) and abiotic determinants (environmental metadata on ponds and

surrounding terrestrial habitat) of the GCN at an unparalleled scale, and determinants

of vertebrate species richness at the pondscape - an impractical task by conventional

means. Finally, we investigated applicability of the HSI to predict eDNA-based GCN

occupancy and vertebrate species richness of ponds.

MATERIALS AND METHODS

Samples

Samples from 508 ponds included in Natural England's Great Crested Newt Evidence Enhancement Programme were processed using eDNA metabarcoding alongside 24 privately surveyed ponds. Water samples were collected using established methodology (Biggs *et al.* 2015), detailed in Supporting Information. In brief, 20 x 30 mL water samples were collected from each pond and pooled. Six 15 mL subsamples were taken from the pooled sample and each added to 33.5 mL absolute ethanol and 1.5 mL sodium acetate 3 M (pH 5.2). Subsamples were pooled during DNA extraction to produce one eDNA sample per pond. Targeted qPCR detected GCN in 265 (49.81%) ponds (Harper *et al.* 2018).

Environmental metadata (Table S1) were collected for 504 of 532 ponds (Fig. S1), including: maximum depth; circumference; width; length; area; density; terrestrial overhang; shading; macrophyte cover; HSI score (Oldham *et al.* 2000); HSI band (categorical classification of HSI score by ARG-UK 2010); permanence; water quality; pond substrate; presence of inflow or outflow; presence of pollution; presence of other amphibians, fish and waterfowl; woodland; rough grass; scrub/hedge; ruderals; other good terrestrial habitat (i.e. good terrestrial habitat that did not conform to aforementioned habitat types); and overall terrestrial habitat quality.

DNA reference database construction

A custom, phylogenetically curated reference database of mitochondrial 12S ribosomal RNA (rRNA) sequences for UK fish species was previously constructed for eDNA metabarcoding of lake fish communities (Hänfling *et al.* 2016). In this study, additional reference databases for UK amphibians, reptiles, birds, and mammals were constructed (see Harper *et al.* 2018 and Supporting Information). Reference sequences available for species varied across vertebrate groups: amphibians 100.00% ($N = 21$), reptiles 90.00% ($N = 20$), mammals 83.93% ($N = 112$), and birds 55.88% ($N = 621$). Table S2 lists species without database representation, i.e. no records for any species in a genus. Sanger sequences were obtained from tissue of GCN, smooth newt, Alpine newt (*Mesotriton alpestris*), common toad (*Bufo bufo*), and common frog (*Rana temporaria*) to supplement the amphibian database (see Supporting Information). The complete reference databases compiled in GenBank format have been deposited in a dedicated GitHub repository for this study, permanently archived at: https://doi.org/10.5281/zenodo.1304107.

Primer validation

Reference databases were combined for *in silico* validation of published 12S rRNA primers 12S-V5-F (5'-ACTGGGATTAGATACCCC-3') and 12S-V5-R (5'-TAGAACAGGCTCCTCTAG-3') (Riaz *et al.* 2011) using ecoPCR software (Ficetola *et al.* 2010). Set parameters allowed a 50-250 bp fragment and three mismatches between each primer and reference sequence. Primers were validated *in vitro* for UK

fish by Hänfling *et al*. (2016) and here for six UK amphibian species (Fig. S2).

eDNA metabarcoding

Our two-step PCR workflow is fully described in Harper *et al*. (2018) and Supporting

Information. eDNA was first amplified with the aforementioned primers, where PCR

positive controls (six per PCR plate; N = 114) were cichlid (*Rhamphochromis esox*)

DNA and PCR negative controls (six per PCR plate; N = 114) were sterile molecular

grade water (Fisher Scientific UK Ltd, UK). PCR products were individually purified

using E.Z.N.A.® Cycle Pure V-Spin Clean-Up Kits (Omega Bio-tek, GA, USA)

following the manufacturer's protocol. The second PCR bound Multiplex

Identification (MID) tags to the purified products. PCR products were individually

purified using magnetic bead clean-up and quantified with a Quant-IT™ PicoGreen™

dsDNA Assay (Invitrogen, ThermoFisher Scientific, UK). Samples were normalised,

pooled, and libraries quantified using a Qubit™ dsDNA HS Assay Kit (Invitrogen,

ThermoFisher Scientific, UK). Libraries were sequenced on an Illumina MiSeq using

2 x 300 bp V3 chemistry (Illumina Inc., San Diego, CA, USA) and raw sequence

reads processed using metaBEAT (metaBarcoding and Environmental Analysis Tool)

v0.8 (https://github.com/HullUni-bioinformatics/metaBEAT). After quality trimming,

merging, chimera detection, and clustering, non-redundant query sequences were

compared against our reference database using BLAST (Zhang *et al.* 2000). Putative

taxonomic identity was assigned using a lowest common ancestor (LCA) approach

based on the top 10% BLAST matches for any query matching with at least 98%

identity to a reference sequence across more than 80% of its length. Unassigned

sequences were subjected to a separate BLAST against the complete NCBI nucleotide

(nt) database at 98% identity to determine the source via LCA as described above.

The bioinformatic analysis has been deposited in the GitHub repository for

reproducibility.

Data analysis

Analyses were performed in the statistical programming environment R v.3.4.2 (R

Core Team 2017). Data and R scripts have been deposited in the GitHub repository.

Assignments from different databases were merged, and spurious assignments (i.e.

non-UK species, invertebrates and bacteria) removed from the dataset. The genus

Strix was reassigned to tawny owl (*Strix aluco*) as it is the only UK representative of

this genus. Where family and genera assignments containing a single UK

representative had reads assigned to species, reads from all assignment levels were

merged and manually assigned to that species. Consequently, all taxonomic

assignments in the final dataset were of species resolution. The only misassignment

was Scottish wildcat (*Felis silvestris*), which does not occur at the sampling localities

(Kent, Lincolnshire and Cheshire) and was therefore reassigned to domestic cat (*Felis*

catus).

To minimise risk of false positives, species were only classed as present at

sites if their sequence frequency exceeded species-specific thresholds. Thresholds

were defined using the maximum sequence frequency of each species in PCR positive

controls ($N = 114$; Table S3). For example, the GCN threshold was 0.028% to omit

false positives in PCR positive controls. After applying thresholds, the read count data

were converted to a species presence-absence matrix. Analyses were based on

species-specific thresholds, but also performed for different blanket sequence

thresholds (0.05 - 30%, Tables S4-9). We tested biotic and abiotic determinants of

GCN occupancy and vertebrate species richness, and appropriateness of the HSI.

Hypotheses are summarised in Table 1.

All Generalised Linear Mixed Models (GLMMs) were executed using the R

package 'lme4' v1.1-12 (Bates *et al.* 2015). GCN occurrence relating to number of

other vertebrate species was investigated with a binomial GLMM, and species

associations identified using the R package 'cooccur' v1.3 (Griffith *et al.* 2016) ($N =$

532). Identified associations informed candidate biotic variables for inclusion with

abiotic variables (Table S1) in a binomial GLMM of GCN occurrence ($N = 504$).

Candidate explanatory variables were assessed for collinearity, relative importance,

and non-linearity (see Supporting Information). HSI score and band were collinear,

thus HSI score was analysed in a separate binomial GLMM. Using the R package

'ncf' v1.1-7 (Bjornstad 2009), spline correlograms of the Pearson residuals from a

binomial Generalised Linear Model (GLM) and GLMM were compared to assess

spatial autocorrelation potential in our dataset. Sample was treated as a random effect

in the GLMM to account for dependencies within ponds (Zuur *et al.* 2009). The

mixed model successfully accounted for spatial autocorrelation within sites. The same

steps were performed to identify explanatory variables and a modeling framework for

vertebrate species richness. A Poisson distribution was specified for all species

richness models as the response variable was integer count data.

Binomial and Poisson models considered respectively were nested thus best

models were chosen using stepwise backward deletion of terms based on Likelihood

Ratio Tests (LRTs). Akaike's Information Criteria (AIC) was employed to evaluate

model fit (Akaike 1973). Final models were tested for overdispersion using the R

package 'RVAideMemoire' v 0.9-69 (Hervé 2018) and a custom function testing

overdispersion of the Pearson residuals. Model fit was assessed using the Hosmer and

Lemeshow Goodness of Fit Test (Hosmer & Lemeshow 2000) within the R package

'ResourceSelection' v0.3-2 (Lele *et al.* 2016), quantile-quantile plots and partial

residual plots (Zuur *et al.* 2009). Model predictions were obtained using the

predictSE() function in the 'AICcmodavg' package v2.1-1 (Mazerolle 2016) and

upper and lower 95% CIs were calculated from the standard error of the predictions.

Results were plotted using the R package 'ggplot2' v 2.1.0 (Wickham 2009).

RESULTS

eDNA metabarcoding

A total of 532 eDNA samples and 228 PCR controls were processed across two

sequencing runs. The runs generated raw sequence read counts of 36,236,862 and

32,900,914 respectively. After trimming and merging of paired-end reads, 26,294,906

and 26,451,564 sequences remained. Following removal of chimeras and redundancy

via clustering, the libraries contained 14,141,237 and 14,081,939 sequences (average

read counts of 36,826 and 36,671 per sample respectively), of which 13,126,148 and

13,113,143 sequences were taxonomically assigned. The final dataset (thresholds

applied and assignments corrected) contained 60 vertebrate species (Table S10),

including six amphibians, 14 fish, 18 birds, and 22 mammals (Fig. S3).

Pondscape biodiversity

All native amphibians were found as well as the non-native marsh frog (*Pelophylax*

ridibundus). The GCN ($N = 149$), smooth newt ($N = 152$) and common frog ($N = 120$)

were widespread, but the common toad ($N = 42$), palmate newt (*Lissotriton*

helveticus, $N = 5$) and marsh frog were uncommon ($N = 1$). The threatened European

eel (*Anguilla anguilla*, $N = 15$), European bullhead (*Cottus gobio*, $N = 14$), and

crucian carp (*Carassius carassius,* $N = 2$) were detected alongside native fishes, such

as pike (*Esox lucius*, $N = 17$) and roach (*Rutilus rutilus*, $N = 72$), but also introduced

species, including common carp ($N = 41$), ruffe (*Gymnocephalus cernua*, $N = 1$), and

rainbow trout (*Oncorhynchus mykiss*, $N = 3$). Regular and irregular waterfowl were

identified, ranging from common moorhen (*Gallinula chloropus*, $N = 215$) to grey

heron (*Ardea cinerea*, $N = 1$) and Eurasian oystercatcher (*Haematopus ostralegus*, N

$= 1$). Terrestrial fauna were often detected in fewer than five ponds (Fig. S3c, d).

Buzzard (*Buteo buteo*, $N = 4$), Eurasian jay (*Garrulus glandarius*, $N = 7$), dunnock

(*Prunella modularis*, $N = 4$), and starling (*Sturnus vulgaris*, $N = 4$) were the most

frequently detected terrestrial birds. Domesticated, including cow (*Bos taurus*, $N =$

179) and pig (*Sus scrofa*, $N = 140$), or introduced mammals (Mathews *et al.* 2018),

such as grey squirrel (*Sciurus carolinensis*, $N = 57$) and Reeve's muntjac (*Muntiacus*

reevesi, $N = 3$), outweighed native mammals. Nonetheless, we detected several

mammals with Biodiversity Actions Plans and/or of conservation concern (Mathews

et al. 2018), including otter (*Lutra lutra*, $N = 1$), water vole (*Arvicola amphibius*, $N =$

16), European polecat (*Mustela putorius*, $N = 1$), brown hare (*Lepus europaeus*, $N =$

1) and water shrew (*Neomys fodiens*, *N* = 9). Notably, the invasive American mink

(*Neovison vison*) was absent despite widespread UK distribution (Mathews *et al.*

2018). All species and their detection frequencies are listed in Table S10.

Biotic determinants of GCN occurrence

GCN occupancy was negatively correlated with fish species richness, but positively

influenced by amphibian and waterfowl species richness (Fig. 1, GLMM:

overdispersion χ^2_{525} = 517.636, *P* = 0.582; fit χ^2_8 = 22.524, *P* = 0.004, R^2 = 9.43%).

The GCN had significant positive associations with four species (Fig. 2), including

the smooth newt, common coot (*Fulica atra*), common moorhen, and pig. However,

the GCN had significant negative associations with five species (Fig. 2), including the

common toad, ninespine stickleback, three-spined stickleback, grey squirrel, and

common pheasant (*Phasianus colchicus*). Non-significant negative associations were

also observed between the GCN and common carp, green-winged teal (*Anas*

carolinensis), and badger (*Meles meles*). Only presence-absence of the smooth newt,

common toad, three-spined stickleback, and grey squirrel were retained by model

selection as explanatory variables for the GLMM (Figs. 3a-d) with abiotic

determinants. Results of analyses are summarised and compared to previously

reported determinants in Table 1.

GCNs were more likely to occupy ponds with more amphibian species (Fig.

1a). GCNs were detected in 51.97% of ponds (*N* = 152) containing smooth newts, but

in only 11.91% of ponds (*N* = 42) with common toads (Fig. S3a). GCN occurrence

probability was lower in ponds with more fish species, and GCNs were absent from

ponds with more than four fish species (Fig. 1b). The GCN was only found in 14.29%

($N = 56$) and 6.67% ($N = 15$) of ponds inhabited by the three-spined stickleback and

ninespine stickleback respectively (Fig. S3b). In contrast, GCNs were more likely to

occur in ponds with more waterfowl species (Fig. 1c). GCNs occupied 41.67% ($N =$

48) and 35.81% ($N = 215$) of ponds with common coot and common moorhen

respectively (Fig. S3c). No significant relationships between terrestrial bird or

mammal species richness and GCN occupancy were found (Figs. 1d, e). Yet, species

associations between the GCN and pig, common pheasant, and grey squirrel were

identified. The GCN was detected in 37.14% of ponds ($N = 140$) where pig was

present (Fig. S3d) as opposed to 12.00% ($N = 25$) and 15.79% ($N = 57$) of ponds with

common pheasant (Fig. S3c) and grey squirrel (Fig. S3d) records respectively. GCN

occupancy positively correlated with overall vertebrate species richness (Fig. 3h),

irrespective of individual species associations.

Abiotic determinants of GCN occurrence

Five abiotic determinants were retained in the GLMM explaining GCN occupancy

(GLMM: overdispersion $\chi^2_{490} = 413.394$, $P = 0.995$; fit $\chi^2_8 = 11.794$, $P = 0.161$, $R^2 =$

38.58%). The probability of GCN occurrence increased with greater pond depth but

decreased in ponds with larger area, outflow, some other good terrestrial habitat, and

where ruderal habitat was absent (Table 1, Figs. 3e-g, i-j).

Abiotic determinants of vertebrate species richness

Our analysis (Table 2, GLMM: overdispersion $\chi^2_{494} = 431.959$, $P = 0.979$; fit $\chi^2_8 = -42.708$, $P = 1.000$, $R^2 = 8.94\%$) revealed species richness was greater in ponds with outflow (Fig. 4a), but reduced in those with some rough grass habitat compared to ponds with no or important rough grass habitat (Fig. 4b). Overall quality of terrestrial habitat was also influential (Fig. 4c), where ponds had higher species richness in areas considered to be poor or moderate habitat for GCNs. Species richness was reduced as percentages of terrestrial overhang (Fig. 4d) and macrophyte cover increased (Fig. 4e), but improved with pond density (Fig. 4f).

Applicability of the HSI to GCN and community eDNA surveys

HSI score positively correlated with both GCN occurrence (GLMM: overdispersion $\chi^2_{501} = 506.763$, $P = 0.4198$; fit $\chi^2_8 = 8.118$, $P = 0.422$, $R^2 = 4.99\%$) and vertebrate species richness (GLMM: overdispersion $\chi^2_{501} = 389.744$, $P = 0.999$; fit $\chi^2_8 = -145.12$, $P = 1.000$, $R^2 = 1.10\%$). GCN occupancy probability (Table 1, Fig. 3k) and vertebrate species richness (Table 2, Fig. 4g) were improved in ponds with higher HSI score.

DISCUSSION

We have ground-truthed eDNA metabarcoding as a tool for ecological hypothesis testing using the community data generated by this tool in combination with environmental metadata for ponds. We tested biotic and abiotic determinants of GCN occupancy and wider vertebrate biodiversity as well as the appropriateness of the GCN HSI for eDNA survey. GCN occupancy was higher in ponds containing smooth newts, but devoid of common toads, three-spined sticklebacks, and grey squirrels. Ponds inhabited by GCNs were typically small, deep, absent of outflow, and surrounded by ruderal and good quality terrestrial habitat. Vertebrate species richness was higher in ponds where outflow, some rough grass habitat, and poor or moderate terrestrial habitat for the GCN were present. Species richness was lower at higher percentages of terrestrial overhang and macrophyte cover, but greater at higher pond density. The GCN HSI was appropriate for predicting both GCN occupancy and vertebrate species richness. Our findings demonstrate the power of eDNA metabarcoding to reboot freshwater monitoring and research by providing biodiversity data en masse at low cost.

Pondscape biodiversity

eDNA metabarcoding detected six amphibian, 14 fish, 18 bird, and 22 mammal species across 532 UK ponds. This diverse species inventory emphasises the importance of ponds as habitat for aquatic taxa, but also as stepping stones for semi-aquatic and terrestrial taxa (De Meester *et al.* 2005; Hill *et al.* 2018) through

provision of drinking, foraging, dispersal, and reproductive opportunities (Biggs *et al.*

2016; Klymus *et al.* 2017). Some species detections may be the result of eDNA

transport from water bodies in the surrounding area (Hänfling *et al.* 2016) to ponds

via inflow; however, this signifies the capacity of ponds to provide natural samples of

freshwater biodiversity in the wider catchment.

Biotic determinants of GCN occurrence

GCN occurrence was positively associated with vertebrate species richness, which

may support its status as an indicator or umbrella species for pond biodiversity

(Gustafson *et al.* 2006). GCNs were more likely to occur in ponds with higher

amphibian species richness - particularly ponds containing smooth newts and absent

of common toads. GCNs and smooth newts have similar habitat requirements and

tend to breed in the same ponds (Skei *et al.* 2006; Rannap *et al.* 2009a; Denoël *et al.*

2013), with >60% overlap reported (Rannap & Briggs 2006). However, smooth newts

can inhabit a broader range of habitat (Rannap & Briggs 2006; Skei *et al.* 2006) than

GCNs, which depend on larger, deeper ponds with abundant macrophytes and no fish

located in open, semi-rural landscapes (Denoël *et al.* 2013). Common toads inhabit

fish-containing ponds (Manenti & Pennati 2016) which may explain their negative

association with GCNs, but newts may also predate toad eggs and larvae (Langton *et*

al. 2001).

GCNs were negatively associated with higher fish species richness, and

specifically the presence of three-spined and ninespine sticklebacks - fish that are

common in and typical of ponds. All GCN life stages may be predated by fishes

(Langton *et al.* 2001) and negative effects of fish presence-absence on GCN

occupancy, distribution, and abundance are repeatedly reported (Joly *et al.* 2001;

Rannap & Briggs 2006; Skei *et al.* 2006; Denoël & Ficetola 2008; Rannap *et al.*

2009a, b; Hartel *et al.* 2010; Denoël *et al.* 2013). Three-spined sticklebacks predate

GCN eggs and larvae (McLee & Scaife 1992; Jarvis 2010), and have non-

consumptive effects on GCN embryos (Jarvis 2010). GCN larvae were also found to

alter their behaviour when exposed to the predatory three-spined stickleback but not

the non-predatory crucian carp (Jarvis 2012), another fish characteristic of ponds.

In our study, we detected GCNs in 50% of ponds inhabited by crucian carp,

but <20% of ponds containing large and/or predatory fishes, e.g. pike and three-

spined stickleback. Although fewer ponds contained crucian carp than pike or

stickleback, previous research also indicates large and/or predatory fish are more

detrimental to GCN occurrence (Skei *et al.* 2006; Hartel *et al.* 2010; Chan 2011).

Crucian carp do not hinder GCN oviposition, larval behaviour, or recruitment success

(Chan 2011; Jarvis 2012), or pond invertebrate and macrophyte diversity (Stefanoudis

et al. 2017). In contrast, common carp foraging reduces invertebrate density and

macrophyte cover (Maceda-Veiga *et al.* 2017), which lowers GCN reproductive and

foraging success and heightens predator exposure (Gustafson *et al.* 2006; Rannap &

Briggs 2006; Chan 2011). The crucian carp and common carp are both included

among fish species assumed to negatively impact GCNs and whose presence-absence

is assessed for the GCN HSI (ARG-UK 2010). However, it is evident that crucian

carp do not directly predate GCNs or indirectly alter their behaviour, reproductive

success, or habitat. Therefore, we advocate a systematic re-evaluation of problematic

fish species for GCN conservation.

Unexpectedly, GCNs were positively associated with waterfowl species

richness, namely presence of common coot and common moorhen. These waterfowl

species share macrophytes and macroinvertebrates as resources with amphibians,

feeding on both directly (Perrow *et al.* 1997; Paillisson & Marion 2001; Wallau *et al.*

2010). Coots and moorhens crop emergent macrophytes to search for invertebrate

prey (Paillisson & Marion 2001; Wallau *et al.* 2010), which may indirectly benefit

GCN foraging. Although coots can also pull up submerged vegetation and damage

vegetation banks (Lauridsen *et al.* 1993), diet is macrophyte-dominated in late

summer and autumn (Perrow *et al.* 1997) and unlikely to impact GCN breeding in

spring (Langton *et al.* 2001). The positive association identified here between GCNs

and these waterfowl most likely reflects a shared preference for macrophyte-rich

ponds.

The GCN had negative associations with the common pheasant and grey

squirrel, but a positive association with pig. GCNs are at risk of predation during

terrestrial as well as aquatic phases of their life history (Langton *et al.* 2001;

Gustafson *et al.* 2011). There have been anecdotal records of pheasant predation on

herpetofauna, including the GCN (Rice 2016), which our results would support.

However, the terrestrial associations identified may instead reflect land-use and

indirect interaction. GCNs prefer ponds surrounded by deciduous forest and pasture

(Gustafson *et al.* 2011), where pheasants and domestic animals are commonplace,

over those in urban areas (Denoël & Ficetola 2008; Hartel *et al.* 2010), which support

dense populations of grey squirrels.

Abiotic determinants of GCN occurrence

GCNs were more likely to occupy deeper ponds, but less likely to inhabit larger ponds, with outflow present, no ruderal habitat, and some other good terrestrial habitat. Consistent with our results, pond depth was previously found to positively influence GCN occupancy (Denoël & Ficetola 2008). Shallow ponds can be inhospitable due to drying or freezing and may contain less prey, but detrimental effects are often observed in open farmland (Denoël & Ficetola 2008). Although our results indicate GCNs prefer larger ponds, pond area does not always influence occurrence (Maletzky *et al.* 2007; Denoël & Ficetola 2008; Gustafson *et al.* 2011) and was deemed a poor predictor of reproductive success (Vuorio *et al.* 2013). GCNs have been found to utilise small and large ponds (Rannap & Briggs 2006; Skei *et al.* 2006); however, very small ponds (less than 124 m^2) may be unable to support all life stages, and larger ponds may contain fish and experience eutrophication from agricultural or polluted run-off (Rannap & Briggs 2006). Effects of pond outflow (facilitated by drains, pipes or streams) are understudied, whereas inflow impacts biodiversity via polluted agricultural run-off and connections to streams and rivers containing large, predatory fish. Based on our findings that show outflow presence deters GCN occupation, we suggest outflow minimises fluctuations in pond depth (Freshwater Habitats Trust 2015) and alters patterns of colonisation and community structure to those of ponds which fluctuate with precipitation.

Our results support the importance of good terrestrial habitat to GCNs for shelter, foraging and dispersal (Langton *et al.* 2001). We observed higher GCN occurrence in ponds surrounded by ruderal or other good quality terrestrial habitat. However, the majority of terrestrial habitat variables were not retained by our model

selection. Hartel *et al.* (2010) also found landscape variables, excluding urbanisation, were inadequate predictors of GCN distribution, although their study area was mostly rural and thus optimal for amphibians. Conversely, research on more diverse landscapes found GCN occupancy was lower in coniferous forest, yet higher in deciduous or herb-rich forest and pasture (Gustafson *et al.* 2011; Vuorio *et al.* 2013). In our study, the metadata available were qualitative, preventing detailed analyses on terrestrial habitat quality in relation to GCN occupancy. Better understanding of occupancy and interactions with terrestrial species could be achieved with quantitative data on terrestrial habitat type, density, distance to ponds, and species utilisation, but this is a phenomenal task for large numbers of ponds across a vast landscape (Denoël & Ficetola 2008).

Abiotic determinants of vertebrate species richness

Vertebrate species richness was higher in ponds where outflow was present, with poor or moderate overall terrestrial habitat for GCNs, and more ponds nearby. Conversely, species richness was lower in ponds with higher percentages of macrophyte cover and terrestrial overhang, and absent of or surrounded by some rough grass habitat. We compare our results to previous studies of aquatic species richness but these largely focus on species assemblages or guilds, primarily macrophytes, macroinvertebrates, and amphibians. Outflow and inflow have been understudied in relation to pond biodiversity, although outflow may release harmful pollutants and pathogens (Beutel & Larson 2015) that would accumulate in a closed pond system. Species richness increased as pond density increased, which echoes the positive relationship between

pond density and macrophyte and macroinvertebrate richness observed by Gledhill *et*

al. (2008). These findings combined again highlight the importance of pond networks

for aquatic and non-aquatic taxa (Hill *et al.* 2018). Shade was identified as a principal

driver of macroinvertebrate and macrophyte diversity in ponds (Sayer *et al.* 2012).

Yet, canopy and macrophyte cover positively influence amphibian species richness

(Piha *et al.* 2007). Plentiful rough grass habitat may create more ecological niches and

foraging opportunity for different vertebrates, but quantitative data on type and

abundance of terrestrial habitat are needed to understand species preferences.

Applicability of the HSI to GCN and community eDNA surveys

We found the HSI can predict eDNA-based GCN occupancy at the pondscape. This

contradicts previous conventional studies which deemed the index inappropriate for

predicting GCN occupancy or survival probabilities (Unglaub *et al.* 2015). Our results

also suggest some indices comprising the GCN HSI, for example, outflow and

terrestrial habitat, represent suitability criteria for other biodiversity. An adapted HSI,

designed to predict species richness, could help select areas for management and

enhancement of aquatic and non-aquatic biodiversity. Nonetheless, the GCN HSI also

confers protection to pond biodiversity by identifying optimal habitat for pond

creation and restoration to encourage populations of this threatened amphibian. The

HSI is not without issue due to qualitative data used for score calculation and

subjective estimation of indices (Oldham *et al.* 2000). For future application of this

index in GCN eDNA survey, we recommend metabarcoding to quantify some

qualitatively assessed indices (e.g. water quality via macroinvertebrate diversity, fish

and waterfowl presence) alongside GCN detection. Provided rigorous spatial and

temporal sampling are undertaken, eDNA metabarcoding can also generate site

occupancy data to estimate relative species abundance (Hänfling *et al.* 2016;

Valentini *et al.* 2016).

Prospects of eDNA metabarcoding for conservation, management, and research

We have demonstrated the effectiveness of eDNA metabarcoding for landscape-scale

biodiversity monitoring and ecological hypothesis testing. We investigated

associations between aquatic and non-aquatic vertebrates, and combined

metabarcoding with environmental metadata to revisit important ecological

hypotheses at an unprecedented scale. Our findings indicate preferred habitat of a

threatened amphibian. GCNs were more likely to occupy ponds where smooth newts

were present, but common toads, three-spined sticklebacks, and grey squirrels were

absent. GCNs prefer small but deep ponds that are absent of outflow, and surrounded

by ruderal and good quality terrestrial habitat. These findings will guide management

in the face of increasing land-use and habitat fragmentation - a poignant issue as

protective legislation for the GCN in the UK is changing. Whilst conservation of

threatened species and their habitat should be a priority, the bigger picture should not

be ignored. eDNA metabarcoding could enhance our understanding of freshwater

networks - particularly pondscapes - to enable more effective monitoring, protection,

and management of aquatic and non-aquatic biodiversity. We are only now beginning

to realise and explore these opportunities.

ACKNOWLEDGEMENTS

This work was funded by the University of Hull. We would like to thank Jennifer

Hodgetts (Fera Science Ltd) for assisting with sample collection, and Jianlong Li

(University of Hull) for primer design and advice on laboratory protocols. Tissue

samples for primer validation and Sanger sequencing were provided by Andrew

Buxton and Richard Griffiths (DICE, University of Kent) under licence from Natural

England, and Barbara Mabel and Elizabeth Kilbride (University of Glasgow).

REFERENCES

Akaike, H. (1973). Maximum likelihood identification of Gaussian autoregressive moving average models. *Biometrika*, 60, 255–265.

ARG-UK (2010). *ARG UK Advice Note 5: Great Crested Newt Habitat Suitability Index*. Amphibian and Reptile Groups of the United Kingdom.

Bálint, M., Nowak, C., Márton, O., Pauls, S., Wittwer, C., Aramayo, J.L. *et al.* (2017). Twenty-five species of frogs in a liter of water: eDNA survey for exploring tropical frog diversity. *bioRxiv*, 176065. https://doi.org/10.1101/176065

Barnes, M.A. & Turner, C.R. (2015). The ecology of environmental DNA and implications for conservation genetics. *Conserv. Genet.*, 17, 1–17.

Bates, D., Mächler, M., Bolker, B. & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *J. Stat. Softw.*, 67, 1–48.

Beutel, M.W. & Larson, L. (2015). Pathogen removal from urban pond outflow using rock biofilters. *Ecol. Eng.*, 78, 72–78.

Biggs, J., Ewald, N., Valentini, A., Gaboriaud, C., Dejean, T., Griffiths, R.A., *et al.* (2015). Using eDNA to develop a national citizen science-based monitoring programme for the great crested newt (*Triturus cristatus*). *Biol. Conserv.*, 183, 19–28.

Biggs, J., von Fumetti, S. & Kelly-Quinn, M. (2016). The importance of small waterbodies for biodiversity and ecosystem services: implications for policy makers. *Hydrobiologia*, 793, 3–39.

Bjørnstad, O.N. (2016). ncf: spatial nonparametric covariance functions. R package version 1.1-7.

Chan, K. (2011). Can great crested newts (*Triturus cristatus*) coexist with fish?. Masters Thesis. University College London (UCL).

Deiner, K., Bik, H.M., Mächler, E., Seymour, M., Lacoursière-Roussel, A., Altermatt, F. *et al.* (2017). Environmental DNA metabarcoding: transforming how we survey animal and plant communities. *Mol. Ecol.*, 26, 5872–5895.

De Meester, L., Declerck, S., Stoks, R., Louette, G., Van De Meutter, F., De Bie, T. *et al.* (2005). Ponds and pools as model systems in conservation biology, ecology and evolutionary biology. *Aquat. Conserv.*, 15, 715–725.

Denoël, M. & Ficetola, G.F. (2008). Conservation of newt guilds in an agricultural landscape of Belgium: the importance of aquatic and terrestrial habitats. *Aquat. Conserv.*, 18, 714–728.

Denoël, M., Perez, A., Cornet, Y. & Ficetola, G.F. (2013). Similar local and landscape processes affect both a common and a rare newt species. *PLoS ONE*, 8, e62727.

Evans, N.T., Li, Y., Renshaw, M.A., Olds, B.P., Deiner, K., Turner, C.R. *et al.* (2017). Fish community assessment with eDNA metabarcoding: effects of sampling design and bioinformatic filtering. *Can. J. Fish. Aquat. Sci.*, 74, 1362-1374.

Ficetola, G.F., Coissac, E., Zundel, S., Riaz, T., Shehzad, W., Bessière, J. *et al.* (2010). An *In silico* approach for the evaluation of DNA barcodes. *BMC Genomics*, 11, 434.

Freshwater Habitats Trust (2015). Pond Habitat Survey. Survey Manual. Freshwater Habitats Trust, Oxford.

Gledhill, D.G., James, P. & Davies, D.H. (2008). Pond density as a determinant of aquatic species richness in an urban landscape. *Landsc. Ecol.*, 23, 1219–1230.

Griffith, D., Veech, J. & Marsh, C. (2016). cooccur: Probabilistic Species Co-Occurrence Analysis in R. *J. Stat. Softw.*, 69, 1–17.

Gustafson, D.H., Malmgren, J.C. & Mikusiński, G. (2011). Terrestrial Habitat Predicts use of Aquatic Habitat for Breeding Purposes — A Study on the Great Crested Newt (*Triturus cristatus*). *Ann. Zool. Fennici*, 48, 295–307.

Gustafson, D.H., Pettersson, C.J. & Malmgren, J.C. (2006). Great crested newts (*Triturus cristatus*) as indicators of aquatic plant diversity. *Herpetol. J.*, 16, 347–352.

Hänfling, B., Lawson Handley, L., Read, D.S., Hahn, C., Li, J., Nichols, P. *et al.* (2016). Environmental DNA metabarcoding of lake fish communities reflects long-term data from established survey methods. *Mol. Ecol.*, 25, 3101–3119.

Harper, L.R., Lawson Handley, L., Hahn, C., Boonham, N., Rees, H.C., Gough, K.C. *et al.* (2018). Needle in a haystack? A comparison of eDNA metabarcoding and targeted qPCR for detection of the great crested newt (*Triturus cristatus*). *Ecol. Evol.*, 8, 6330-6341.

Hartel, T., Nemes, S. & Oellerer, K. (2010). Using connectivity metrics and niche modelling to explore the occurrence of the northern crested newt *Triturus cristatus* (Amphibia, Caudata) in a traditionally managed landscape. *Environ. Conserv.*, 37, 195–200.

Hering, D., Borja, A., Jones, J.I., Pont, D., Boets, P., Bouchez, A., *et al.* (2018). Implementation options for DNA-based identification into ecological status assessment under the European Water Framework Directive. *Water Res.*, 138, 192–205.

Hervé, M. (2018) RVAideMemoire: Diverse basic statistical and graphical functions.
R package version 0.9-69.
Hill, M.J., Hassall, C., Oertli, B., Fahrig, L., Robson, B.J., Biggs, J. *et al.* (2018). New
policy directions for global pond conservation. *Conserv. Lett.*, e12447.
Hosmer, D.W. & Lemeshow, S. (2000). *Applied Logistic Regression*. Second Edition.
John Wiley & Sons, Inc., New York, USA, pp. 31–46.
Jarvis, L.E. (2010). Non-consumptive effects of predatory three-spined sticklebacks
(*Gasterosteus aculeatus*) on great crested newt (*Triturus cristatus*) embryos.
Herpetol. J., 20, 271–275.
Jarvis, L.E. (2012). Microhabitat preferences of the great crested newt *Triturus*
cristatus in a woodland area. PhD Thesis. Open University.
Joly, P., Miaud, C., Lehmann, A. & Grolet, O. (2001). Habitat Matrix Effects on Pond
Occupancy in Newts. *Conserv. Biol.*, 15, 239–248.
Kelly, R.P., O'Donnell, J.L., Lowell, N.C., Shelton, A.O., Samhouri, J.F., Hennessey,
S.M. *et al.* (2016). Genetic signatures of ecological diversity along an
urbanization gradient. *PeerJ*, 4, e2444.
Klymus, K.E., Richter, C.A., Thompson, N. & Hinck, J.E. (2017). Metabarcoding of
Environmental DNA Samples to Explore the Use of Uranium Mine Containment
Ponds as a Water Source for Wildlife. *Diversity*, 9, 54.
Langton, T., Beckett, C. & Foster, J. (2001). Great crested newt conservation
handbook. Froglife, Halesworth.
Lauridsen, T.L., Jeppesen, E. & Andersen, F.Ø. (1993). Colonization of submerged
macrophytes in shallow fish manipulated Lake Væng: impact of sediment
composition and waterfowl grazing. *Aquat. Bot.*, 46, 1–15.
Lawson Handley, L. (2015). How will the "molecular revolution" contribute to
biological recording? *Biol. J. Linn. Soc. Lond.*, 115, 750–766.
Lele, S.R., Keim, J.L. & Solymos, P. (2016). ResourceSelection: Resource Selection
(Probability) Functions for Use-Availability Data. R package version 0.3-2.
Maceda-Veiga, A., López, R. & Green, A.J. (2017). Dramatic impact of alien carp
Cyprinus carpio on globally threatened diving ducks and other waterbirds in
Mediterranean shallow lakes. *Biol. Conserv.*, 212, 74–85.
Maletzky, A., Kyek, M. & Goldschmid, A. (2007). Monitoring status, habitat features
and amphibian species richness of crested newt (*Triturus cristatus* superspecies)
ponds at the edge of the species range (Salzburg, Austria). *Ann. Limnol. – Int. J.*
of Lim., 43, 107–115.
Manenti, R. & Pennati, R. (2016). Environmental factors associated with amphibian
breeding in streams and springs: effects of habitat and fish occurrence. *Amphib-*
reptil., 37, 237–242.
Mathews, F., Kubasiewicz, L.M., Gurnell, J., Harrower, C.A., McDonald, R.A. &
Shore, R.F. (2018). A Review of the Population and Conservation Status of
British Mammals: Technical Summary. A report by the Mammal Society under
contract to Natural England, Natural Resources Wales and Scottish Natural
Heritage. Natural England, Peterborough.
Mazerolle, M.J. (2016). AICcmodavg: Model selection and multimodel inference
based on (Q)AIC(c). R package version 2.1-1.
McLee, A.G. & Scaife, R.W. (1992). The colonisation by great crested newts
(*Triturus cristatus*) of a water body following treatment with a piscicide to
remove a large population of sticklebacks (*Gasterosteus aculeatus*). *Herpetol.*
Bull., 42, 6–9.

Muha, T.P., Rodríguez-Rey, M., Rolla, M. & Tricarico, E. (2017). Using Environmental DNA to Improve Species Distribution Models for Freshwater Invaders. *Front. Ecol. Evol.*, 5, 158.

Oldham, R.S., Keeble, J., Swan, M.J.S. & and Jeffcote, M. (2000). Evaluating the suitability of habitat for the great crested newt (*Triturus cristatus*). *Herpetol. J.*, 10, 143–155.

Olds, B.P., Jerde, C.L., Renshaw, M.A., Li, Y., Evans, N.T., Turner, C.R. *et al.* (2016). Estimating species richness using environmental DNA. *Ecol. Evol.*, 6, 4214–4226.

Paillisson, J.-M. & Marion, L. (2001). Interaction between coot (*Fulica atra*) and waterlily (*Nymphaea alba*) in a lake: the indirect impact of foraging. *Aquat. Bot.*, 71, 209–216.

Perrow, M.R., Schutten, H.J., Howes, J.R., Holzer, T., Madgwick, F.J. & Jowitt, A. (1997). Interactions between coot (*Fulica atra*) and submerged macrophytes: the role of birds in the restoration process. In: *Shallow Lakes' 95*. Springer, Dordrecht, pp. 241–255.

Piha, H., Luoto, M. & Merilä, J. (2007). Amphibian occurrence is influenced by current and historic landscape characteristics. *Ecol. Appl.*, 17, 2298–2309.

Rannap, R. & Briggs, L. (2006). THE CHARACTERISTICS OF GREAT CRESTED NEWT TRITURUS CRISTATUS'BREEDING PONDS. Protection of the Great Crested Newt. Project Report. The experiences of LIFE-Nature project "Protection of *Triturus cristatus* in the Eastern Baltic Region" LIFE04NAT/EE/000070. Ministry of the Environment of the Republic of Estonia, Tallinn.

Rannap, R., Lõhmus, A. & Briggs, L. (2009a). Niche position, but not niche breadth, differs in two coexisting amphibians having contrasting trends in Europe. *Divers. Distrib.*, 15, 692–700.

Rannap, R., Lõhmus, A. & Briggs, L. (2009b). Restoring ponds for amphibians: a success story. *Hydrobiologia*, 634, 87–95.

Rees, H.C., Maddison, B.C., Middleditch, D.J., Patmore, J.R.M. & Gough, K.C. (2014). REVIEW: The detection of aquatic animal species using environmental DNA–a review of eDNA as a survey tool in ecology. *J. Appl. Ecol.*, 51, 1450–1459.

Riaz, T., Shehzad, W., Viari, A., Pompanon, F., Taberlet, P. & Coissac, E. (2011). ecoPrimers: inference of new DNA barcode markers from whole genome sequence analysis. *Nucleic Acids Res.*, 39, e145.

Rice, C.N. (2016). Abundance, impacts and resident perceptions of non-native common pheasants (*Phasianus colchicus*) in Jersey, UK Channel Islands. Masters Thesis. University of Kent.

Sayer, C., Andrews, K., Shilland, E., Edmonds, N., Edmonds-Brown, R., Patmore, I. *et al.* (2012). The role of pond management for biodiversity conservation in an agricultural landscape. *Aquat. Conserv.*, 22, 626–638.

Skei, J.K., Dolmen, D., Rønning, L. & Ringsby, T.H. (2006). Habitat use during the aquatic phase of the newts *Triturus vulgaris* (L.) and *T. cristatus* (Laurenti) in central Norway: proposition for a conservation and monitoring area. *Amphib-Reptil.*, 27, 309–324.

Stefanoudis, P.V., Sayer, C.D., Greaves, H.M., Davidson, T.A., Robson, H., Almeida, D. *et al.* (2017). Consequences of fish for cladoceran, water beetle and macrophyte communities in a farmland pond landscape: implications for conservation. *Fund. Appl. Limnol.*, 190, 141–156.

Unglaub, B., Steinfartz, S., Drechsler, A. & Schmidt, B.R. (2015). Linking habitat
suitability to demography in a pond-breeding amphibian. *Front. Zool.*, 12, 9.
Ushio, M., Fukuda, H., Inoue, T., Makoto, K., Kishida, O., Sato, K. *et al.* (2017).
Environmental DNA enables detection of terrestrial mammals from forest pond
water. *Mol. Ecol. Resour.*, 17, e63-e75.
Valentini, A., Taberlet, P., Miaud, C., Civade, R., Herder, J., Thomsen, P.F. *et al.*
(2016). Next-generation monitoring of aquatic biodiversity using environmental
DNA metabarcoding. *Mol. Ecol.*, 25, 929–942.
Vuorio, V., Heikkinen, R.K. & Tikkanen, O.-P. (2013). Breeding Success of the
Threatened Great Crested Newt in Boreal Forest Ponds. *Ann. Zool. Fennici*, 50,
158–169.
Wallau, G.L., Della-Flora, F., Bueno, A.S., Corso, J., Ortiz, M.F. & Cáceres, N.C.
(2010). Behaviour of the Common Moorhen in Rio Grande do Sul, Brazil. *Acta
Ethol.*, 13, 127–135.
Wickham, H. (2009). *ggplot2: Elegant Graphics for Data Analysis*. Springer New
York, USA.
Zhang, Z., Schwartz, S., Wagner, L. & Miller, W. (2000). A greedy algorithm for
aligning DNA sequences. *J. Comput. Biol.*, 7, 203–214.
Zuur, A.F., Ieno, E.N., Walker, N., Saveliev, A.A. & Smith, G.M. (2009). *Mixed
effects models and extensions in ecology with R*. Springer New York, USA.
793

Table 1 Summary of established and novel biotic and abiotic determinants of GCN occupancy. Reported effects on GCN occupancy in the literature and hypothesised effects on eDNA-based GCN occurrence are given for each determinant. Any determinants not reported in the literature are listed as UNK. Direction of observed effects on eDNA-based GCN occupancy determined by each analysis (GLMM assessing number of species in each vertebrate group, $N = 532$; co-occur analysis, $N = 532$; GLMM combining abiotic and biotic factors $N = 504$; and GLMM assessing HSI score, $N = 504$) are given. No, negative and positive effects are listed as 0, - and + respectively. For categorical variables with more than one level, effect size and standard error (SE) are only given for levels reported in the model summary. Test statistic is for LRT used and significant P-values (<0.05) are in bold. Variables included for model selection but not retained in the final model are listed as NR. Co-occur analysis was not applicable (NA) to abiotic factors.

Determinant	Effect reported	Hypothesised effect	Analysis					
			cooccur			GLMM		
			Effect	*P*	DF	Effect size (SE)	χ^2	*P*
Fish	-/0	-			1	-0.239 (0.124)	4.065	**0.044**
Three-spined stickleback	-	-	-	**0.009**	1	-1.432 (0.561)	9.453	**0.002**
Ninespine stickleback	-	-	-	**0.047**				
Common carp	-	-	-	0.070				
Crucian carp	-	-	0	NA				
Waterfowl	-	-			1	0.617 (0.181)	13.050	**<0.001**
Coot	UNK		+	**0.023**				
Moorhen	UNK		+	**<0.001**				
Green-winged teal	UNK		-	0.099				
Amphibians	UNK				1	0.558 (0.149)	16.641	**<0.001**
Smooth newt	+	+	+	**< 0.001**	1	1.081 (0.303)	7.434	**<0.001**
Common toad	UNK		-	**0.009**	1	-1.635 (0.696)	8.228	**0.004**
Terrestrial birds	UNK				1	-0.335 (0.291)	1.444	0.230
Common pheasant	UNK		-	**0.048**				
Terrestrial mammals	UNK				1	0.028 (0.091)	0.095	0.758
Grey squirrel	UNK		-	**0.018**	1	-1.591 (0.534)	12.432	**<0.001**
Badger	UNK		-	0.099				
Pig	UNK		+	**0.004**				
Cow	UNK		+	0.097				
Pond area	-/+	-	NA	NA	1	-0.0004 (0.0002)	6.453	**0.011**
Pond density	+	+	NA	NA		NR		

Pond depth	+	+	NA	NA	1	0.282 (0.139)	4.266	**0.039**
Water quality	+	+	NA	NA		NR		
Outflow	UNK		NA	NA	1	-0.713 (0.359)	4.467	**0.035**
Macrophyte cover	-/+	-	NA	NA		NR		
Shading	-/+	-	NA	NA		NR		
Woodland	+	+	NA	NA		NR		
Grassland	+	+	NA	NA		NR		
HSI	0/+	+	NA	NA	1	3.020 (0.791)	15.709	**<0.001**
Ruderal	UNK		NA	NA	2		6.507	**0.039**
None						-0.617 (0.527)		
Some						0.032 (0.528)		
Other good terrestrial habitat	UNK		NA	NA	2		7.918	**0.019**
None						0.428 (0.429)		
Some						-0.316 (0.424)		
Species richness	UNK		NA	NA	1	0.527 (0.105)	60.267	**<0.001**

807   **Table 2** Summary of analyses testing for variation in eDNA-based vertebrate species richness across

808   ponds (N = 504), attributable to aquatic and terrestrial habitat. Results of the separate GLMM assessing

809   variation explained by HSI score are italicised. For categorical variables with more than one level,

810   effect size and standard error (SE) are only given for levels reported in the model summary. Test

811   statistic is for LRT used and significant P-values (<0.05) are in bold.

Model variables	DF	Effect size (SE)	χ^2	*P*
Outflow	1		11.220	**<0.001**
Present		0.214 (0.063)		
Rough grass	2		16.715	**<0.001**
Some		-0.297 (0.074)		
None		-0.1402 (0.0795)		
Overall terrestrial habitat	2		8.244	**0.016**
Moderate		0.216 (0.078)		
Poor		0.115 (0.089)		
Pond density	1	0.006 (0.003)	4.564	**0.033**
Macrophyte cover	1	-0.002 (0.001)	4.117	**0.043**
Terrestrial overhang	1	-0.0026 (0.0008)	9.575	**0.002**
HSI score	*1*	*0.459 (0.002)*	*5.034*	***0.025***

Figure 1 GCN presence (orange) and absence (grey) in relation to number of species from different vertebrate groups detected by eDNA ($N = 532$ ponds): (a) other amphibians, (b) fish, (c) waterfowl, (d) terrestrial birds, and (e) mammals. Observed proportion of ponds with and without GCN (left) is plotted alongside predicted probability of GCN occurrence in ponds as determined by the binomial GLMM (right). Numbers on barplots of observed occupancy are the number of ponds for each category. In plots showing predicted GCN occupancy, the observed data is shown as points which have been jittered around 0 and 1 to clarify variation in point density. Blue points are outliers and boxes are the model predictions.

Figure 2 Heat map showing significant ($P < 0.05$) positive and negative species associations determined by the probabilistic co-occurrence model for the eDNA metabarcoding presence-absence data ($N = 532$ ponds). Species names are positioned to indicate the columns and rows that represent their pairwise relationships with other species. Species are ordered by those with the most negative interactions to those with the most positive interactions (left to right). Associations relevant to GCN are highlighted in red.

Figure 3 Biotic and abiotic determinants of GCN occurrence, as predicted by the binomial GLMMs (*N* = 504 ponds): (a) smooth newt occurrence, (b) common toad occurrence, (c) three-spined stickleback occurrence, (d) grey squirrel occurrence, (e) pond outflow, (f) ruderal habitat, (g) other good quality terrestrial habitat, (h) species richness, (i) pond area, (j) pond depth, and (k) HSI score. The 95% CIs, as calculated using the predicted GCN probability values and standard error for these predictions, are given for each relationship. The observed GCN presence (orange) and absence (grey) data are also displayed as points, which have been jittered around 0 and 1 to clarify variation in point density, against the predicted relationships (boxes/lines). Outliers are indicated by blue points.

842
843
Figure 4 Abiotic and biotic determinants of vertebrate species richness, as predicted by the Poisson

GLMMs (N = 504 ponds): (a) outflow, (b) rough grass habitat, (c) overall quality of terrestrial habitat,

(d) percentage of terrestrial overhang, (e) percentage of macrophyte cover, (f) pond density, and (g)

HSI score. The 95% CIs, as calculated using the predicted species richness values and standard error

for these predictions, are given for each relationship. The observed data are also displayed as points,

which have been jittered around 0 and 10 to clarify variation in point density, against the predicted

relationships (boxes/lines). Outliers are indicated by red points.