Generating and testing ecological hypotheses at the pondscape

with environmental DNA metabarcoding: a case study on a

threatened amphibian

Lynsey R. Harper[1,2*], Lori Lawson Handley[1], Christoph Hahn[1,3], Neil Boonham[4,5], Helen C.

Rees[6], Erin Lewis[4], Ian P. Adams[4], Peter Brotherton[7], Susanna Phillips[7] and Bernd Hänfling[1]

[1] Department of Biological and Marine Sciences, University of Hull, Hull, HU6 7RX, UK

[2] Illinois Natural History Survey, Prairie Research Institute, University of Illinois at Urbana-Champaign,

Champaign, Illinois, USA

[3] Institute of Zoology, University of Graz, Graz, Styria, Austria

[4] Fera Science Ltd (Fera), Sand Hutton, York, YO14 1LZ, UK

[5] Newcastle University, Newcastle upon Tyne, NE1 7RU, UK

[6] ADAS, School of Veterinary Medicine and Science, The University of Nottingham, Sutton Bonington Campus,

Leicestershire, LE12 5RD, UK

[7] Natural England, Peterborough, PE1 1NG, UK

*Corresponding author: Lynsey R. Harper

Illinois Natural History Survey, Prairie Research Institute, University of Illinois at Urbana-

Champaign, Champaign, Illinois, USA

E-mail: lynsey.harper2@gmail.com

Running title: Hypothesis testing with eDNA metabarcoding

Abstract

Environmental DNA (eDNA) metabarcoding is revolutionising biodiversity monitoring, but has unrealised potential for ecological hypothesis generation and testing. Here, we validate this potential in a large-scale analysis of vertebrate community data generated by eDNA metabarcoding of 532 UK ponds. We test biotic associations between the threatened great crested newt (*Triturus cristatus*) and other vertebrates as well as abiotic factors influencing *T. cristatus* detection at the pondscape. Furthermore, we test the status of *T. cristatus* as an umbrella species for pond conservation by assessing whether vertebrate species richness is greater in ponds with *T. cristatus* and higher *T. cristatus* Habitat Suitability Index (HSI) scores. *T. cristatus* detection was positively correlated with amphibian and waterfowl species richness. Specifically, *T. cristatus* was positively associated with smooth newt (*Lissotriton vulgaris*), common coot (*Fulica atra*), and common moorhen (*Gallinula chloropus*), but negatively associated with common toad (*Bufo bufo*). *T. cristatus* detection did not significantly decrease as fish species richness increased, but negative associations with common carp (*Cyprinus carpio*), three-spined stickleback (*Gasterosteus aculeatus*) and ninespine stickleback (*Pungitius pungitius*) were identified. *T. cristatus* detection was negatively correlated with mammal species richness, and *T. cristatus* was negatively associated with grey squirrel (*Sciurus carolinensis*). *T. cristatus* detection was negatively influenced by larger pond area, presence of inflow, and higher percentage of shading, but positively correlated with HSI score, supporting its application to *T. cristatus* survey. Vertebrate species richness was significantly higher in *T. cristatus* ponds and broadly increased as *T. cristatus* HSI scores increased. We reaffirm reported associations (e.g. *T. cristatus* preference for smaller ponds) but also provide novel insights, including a negative effect of pond inflow on *T. cristatus*. Our findings demonstrate the prospects of eDNA metabarcoding for ecological hypothesis generation and testing at landscape scale, and dramatic enhancement of freshwater conservation, management, monitoring and research.

Keywords: biodiversity assessment, environmental DNA (eDNA), hypothesis testing, metabarcoding, ponds, species associations, *Triturus cristatus*

1. Introduction

Environmental DNA (eDNA) analysis offers ecologists exceptional power to detect organisms within and across ecosystems. DNA released by organisms into their environment via secretions, excretions, gametes, blood, or decomposition, can be sampled and analysed using different approaches to reveal the distribution of single or multiple species (Rees et al., 2014; Lawson Handley, 2015). eDNA analysis combined with high-throughput sequencing (i.e. eDNA metabarcoding) can yield efficient, comprehensive assessments of entire communities (Deiner et al., 2017), providing a step change in biodiversity monitoring (Hering et al., 2018). eDNA metabarcoding has untapped potential to generate and test ecological hypotheses by enabling biodiversity monitoring at landscape scale with minimal impact to communities under investigation. This potential has already been demonstrated with targeted eDNA analysis by Wilcox et al. (2018), where climate-mediated responses of bull trout (*Salvelinus confluentus*) to biotic and abiotic factors were revealed using quantitative PCR (qPCR) on crowd-sourced eDNA samples. Although eDNA metabarcoding assessments of alpha and beta diversity along environmental gradients are increasing (e.g. Hänfling et al., 2016; Olds et al., 2016; Kelly et al., 2016; Evans et al., 2017; Li et al., 2018a; Nakagawa et al., 2018), this tool is less commonly used for ecological hypothesis testing, such as the impact of environmental stressors (Li et al., 2018b; Macher et al., 2018).

Aquatic ecosystems are highly suited to eDNA studies as eDNA exists in multiple states with rapid modes of transport and degradation, increasing detectability of contemporary biodiversity (Rees et al., 2014; Barnes & Turner, 2015). Lentic systems provide further opportunities for eDNA research, being discrete water bodies with variable physicochemical properties that do not experience flow dynamics (Harper et al., 2019).

Ponds in particular have enormous biodiversity and experimental virtue that has not been

maximised in previous eDNA metabarcoding assessments of this habitat (Valentini et al.,

2016; Evans et al., 2017; Klymus et al., 2017; Ushio et al., 2017; Bálint et al., 2018). These

small and abundant water bodies span broad ecological gradients (De Meester et al., 2005)

and comprise pondscapes - a network of ponds and their surrounding terrestrial habitat (Hill

et al., 2018). Pondscapes contribute substantially to aquatic and non-aquatic biodiversity

across spatial scales, with ponds supporting many rare and protected species in fragmented

landscapes (De Meester et al., 2005; Biggs et al., 2016; Hill et al., 2018). Consequently,

ponds are model systems for experimental validation and examination of biogeographical

patterns (De Meester et al., 2005). Habitat complexity and tools required for different taxa

with associated bias (Evans et al., 2017) and cost (Valentini et al., 2016) once hindered

exhaustive sampling of pond biodiversity (Hill et al., 2018), but eDNA metabarcoding may

overcome these barriers (Harper et al., 2019).

In the UK, the threatened great crested newt (*Triturus cristatus*) is an umbrella

species for pond conservation. The extensive literature on *T. cristatus* ecology provides an

excellent opportunity to validate ecological patterns revealed by eDNA metabarcoding. Both

biotic (e.g. breeding substrate, prey, and predators) and abiotic (e.g. pond area, depth, and

temperature) factors are known to influence *T. cristatus* breeding success (Langton, Beckett

& Foster, 2001). The *T. cristatus* Habitat Suitability Index (HSI [Oldham et al., 2000; ARG-UK,

2010; O'Brien et al., 2017]) accounts for these factors using 10 suitability indices that are

scored and combined to calculate a decimal score between 0 and 1 (where 1 = excellent

habitat). Larvae are susceptible to fish and waterfowl predation (Edgar & Bird, 2006; Rannap

& Briggs, 2006; Skei et al., 2006; Hartel, Nemes & Oellerer, 2010), and adults reportedly

avoid ponds containing three-spined stickleback (*Gasterosteus aculeatus*) (McLee & Scaife,

1992), ninespine stickleback (*Pungitius pungitius*), crucian carp (*Carassius carassius*), and

common carp (*Carassius carpio*) (Rannap, Lõhmus & Briggs, 2009a, b). Conversely, *T.*

cristatus and smooth newt (*Lissotriton vulgaris*) prefer similar habitat and often co-occur

(Rannap & Briggs, 2006; Skei et al., 2006; Rannap et al., 2009a; Denoël et al., 2013; Cayuela

et al., 2018). *T. cristatus* individuals thrive in ponds with good water quality as indicated by

diverse macroinvertebrate communities (Oldham et al., 2000; Rannap et al., 2009a). Pond

networks encourage *T. cristatus* occupancy (Joly et al., 2001; Rannap et al., 2009a; Hartel et

al., 2010; Denoël et al., 2013), but larger pond area discourages presence (Joly et al., 2001).

Ponds with heavy shading (Vuorio, Heikkinen & Tikkanen, 2013) or dense macrophyte cover

(Rannap & Briggs, 2006; Skei et al., 2006; Hartel et al., 2010) are unlikely to support viable

populations. *T. cristatus* individuals also depend on terrestrial habitat, preferring open,

semi-rural pondscapes (Denoël et al., 2013) containing pasture, extensively grazed and

rough grassland, scrub, and coniferous and deciduous woodland (Oldham et al., 2000;

Rannap & Briggs, 2006; Rannap et al., 2009a; Gustafson, Malmgren & Mikusiński, 2011;

Vuorio et al., 2013).

We assessed vertebrate communities at the pondscape using a dataset produced by

eDNA metabarcoding for over 500 ponds with comprehensive environmental metadata. We

validated eDNA metabarcoding as a tool for ecological hypothesis generation and testing,

and compared its outputs to previous results produced by established methods. Specifically,

we identified and tested biotic (community presence-absence data) and abiotic

(environmental metadata on ponds and surrounding terrestrial habitat) factors influencing

T. cristatus detection at the pondscape - an impractical task by conventional means.

Furthermore, we tested the applicability of the HSI to predict eDNA-based *T. cristatus*

detection. Finally, we assessed the umbrella species status of *T. cristatus* by investigating

whether *T. cristatus* detection and the *T. cristatus* HSI score can predict vertebrate species

richness of ponds.

2. Materials and methods

2.1 eDNA samples

Samples were previously collected from 508 ponds as part of Natural England's Great

Crested Newt Evidence Enhancement Programme and from 24 privately surveyed ponds

using established methodology (Biggs et al., 2015), detailed in Supporting Information:

Appendix 1. Briefly, 20 x 30 mL water samples were collected from each pond and pooled.

Six 15 mL subsamples were taken from the pooled sample and each added to 33.5 mL

absolute ethanol and 1.5 mL sodium acetate 3 M (pH 5.2). Subsamples were pooled during

DNA extraction to produce one eDNA sample per pond. Targeted qPCR detected *T. cristatus*

in 265 (49.81%) ponds (Harper et al., 2018).

Environmental metadata were collected for 504 of 532 ponds (Fig. S1) by

environmental consultants contracted for Natural England's Great Crested Newt Evidence

Enhancement Programme. Metadata included: maximum depth of ponds; pond

circumference; pond width; pond length; pond area; pond density (i.e. number of ponds per

km^2); terrestrial overhang; shading; macrophyte cover; HSI score (Oldham et al., 2000); HSI

band (categorical classification of HSI score [ARG-UK, 2010]); pond permanence; water

quality; pond substrate; presence of inflow or outflow; presence of pollution; presence of

other amphibians, fish and waterfowl; woodland; rough grass; scrub/hedge; ruderals; other

terrestrial habitat (i.e. good quality terrestrial habitat that did not conform to

aforementioned habitat types); and overall terrestrial habitat quality (see Table S1).

2.2. eDNA metabarcoding

We repurposed the taxonomically assigned sequence reads from Harper et al. (2018) that

were produced using eDNA metabarcoding of pond water to compare qPCR and eDNA

metabarcoding for *T. cristatus* detection. Here, we provide a summary of their eDNA

metabarcoding workflow (see Supporting Information: Appendix 1 for details).

A custom, phylogenetically curated reference database of mitochondrial 12S

ribosomal RNA (rRNA) sequences for UK fish species was previously constructed for eDNA

metabarcoding of lake fish communities (Hänfling et al., 2016). Harper et al. (2018)

constructed additional reference databases for UK amphibians, reptiles, birds, and

mammals. Reference sequences available for species varied across vertebrate groups:

amphibians 100.00% (N = 21), reptiles 90.00% (N = 20), mammals 83.93% (N = 112), and

birds 55.88% (N = 621). Table S2 lists species without database representation, i.e. no

records for any species in a genus. Sanger sequences were obtained from tissue of *T.*

cristatus, *L. vulgaris*, Alpine newt (*Ichthyosaura alpestris*), common toad (*Bufo bufo*), and

common frog (*Rana temporaria*) to supplement the amphibian database. The complete

reference databases compiled in GenBank format were deposited in a GitHub repository

and permanently archived (https://doi.org/10.5281/zenodo.1188710) by Harper et al.

(2018). Reference databases were combined for *in silico* validation of published 12S rRNA

primers 12S-V5-F (5'-ACTGGGATTAGATACCCC-3') and 12S-V5-R (5'-

TAGAACAGGCTCCTCTAG-3') (Riaz et al., 2011) using ecoPCR software (Ficetola et al., 2010).

Set parameters allowed a 50-250 bp fragment and three mismatches between each primer

and reference sequence. Primers were validated *in vitro* for UK fish by Hänfling et al. (2016)

and by Harper et al. (2018) for six UK amphibian species (Fig. S2).

eDNA samples were amplified in triplicate with the aforementioned primers, where

PCR positive controls (six per PCR plate; n = 114) were cichlid (*Rhamphochromis esox*) DNA

(0.284 ng/µL) and PCR negative controls (six per PCR plate; n = 114) were sterile molecular

grade water (Fisher Scientific UK Ltd, UK). PCR products were individually purified using

E.Z.N.A® Cycle Pure V-Spin Clean-Up Kits (Omega Bio-tek, GA, USA) following the

manufacturer's protocol. Purified product was pooled according to sample for the second

PCR which bound Multiplex Identification tags to the purified products. PCR products were

individually purified using magnetic bead clean-up and quantified with a Quant-IT™

PicoGreen™ dsDNA Assay (Invitrogen, UK). Samples were normalised, pooled, and libraries

quantified using a Qubit™ dsDNA HS Assay (Invitrogen, UK). Libraries were sequenced on an

Illumina® MiSeq using 2 x 300 bp V3 chemistry (Illumina, Inc, CA, USA) and raw sequence

reads processed using metaBEAT (metaBarcoding and Environmental Analysis Tool) v0.97.7

(https://github.com/HullUni-bioinformatics/metaBEAT).

After quality filtering (phred score Q30) and trimming (length filter of 90-110 bp)

using Trimmomatic v0.32 (Bolger, Lohse, & Usadel, 2014), merging (10 bp overlap minimum

and 10% mismatch maximum) using FLASH v1.2.11 (Magoč & Salzberg, 2011), chimera

detection using the uchime algorithm (Edgar et al., 2011) in vsearch v1.1 (Rognes et al.,

2016), and clustering (97% identity with 3 sequences minimum per cluster) with vsearch

v1.1 (Rognes et al., 2016), non-redundant query sequences were compared against the

reference database using BLAST (Zhang et al., 2000). Putative taxonomic identity was

assigned using a lowest common ancestor (LCA) approach based on the top 10% BLAST

matches for any query matching with at least 98% identity to a reference sequence across

more than 80% of its length. Unassigned sequences were subjected to a separate BLAST

against the complete NCBI nucleotide (nt) database at 98% identity to determine the source

via LCA as described above. The bioinformatic analysis was archived

(https://doi.org/10.5281/zenodo.1188710) by Harper et al. (2018) for reproducibility.

2.3 Data analysis

2.3.1 Dataset refinement

Analyses were performed in R v.3.4.3 (R Core Team, 2017). Data and R scripts have been

deposited in a dedicated GitHub repository for this study, which has been permanently

archived (https://doi.org/10.5281/zenodo.3365703). Assignments from different databases

were merged, and spurious assignments (i.e. non-UK species, invertebrates and bacteria)

removed from the dataset. The family Cichlidae was reassigned to *Rhamphochromis esox*.

The green-winged teal (*Anas carolinenisis*) was reassigned to *Anas* (dabbling ducks) because

this species is a rare migrant and available reference sequences were identical across our

metabarcode to those for mallard (*Anas platyrhynchos*) and Eurasian teal (*Anas crecca*),

which are widely distributed across the UK. Scottish wildcat (*Felis silvestris*) does not occur

at the sampling localities (Kent, Lincolnshire and Cheshire) and was therefore reassigned to

domestic cat (*Felis catus*). Wild boar (*Sus scrofa*) and grey wolf (*Canis lupus*) were reassigned

to domestic pig (*Sus scrofa domesticus*) and domestic dog (*Canis lupus familiaris*) given the

restricted distribution of *S. scrofa* and absence of *C. lupus* in the wild in the UK. The genus

Strix was reassigned to tawny owl (*Strix aluco*) as it is the only UK representative of this

genus. Where family and genera assignments containing a single UK representative had reads assigned to species, reads from all assignment levels were merged and manually assigned to that species.

Of the 114 PCR negative controls included, 50 produced no reads (Fig. S3). Reads generated for 64 of 114 PCR negative controls ranged from 0 to 49,227, and strength of each contaminant varied (mean = 0.021%, range = 0 - 100.0% of the total reads per PCR negative control). To minimise risk of false positives, we evaluated different sequence thresholds. These included the maximum sequence frequency of *R. esox* DNA in eDNA samples (100%), maximum sequence frequency of any DNA except *R. esox* in PCR positive controls (83.96%), and taxon-specific thresholds (maximum sequence frequency of each taxon in PCR positive controls). The different thresholds were applied to the eDNA samples and the results from each compared (Fig. S4). The taxon-specific thresholds (Table S3) retained the most biological information, thus these were selected for downstream analysis. Consequently, taxa were only classed as present at sites if their sequence frequency exceeded their threshold. After applying the taxon-specific thresholds, the PCR positive control (*R. esox*), human (*Homo sapiens*), and domestic species (Table S4) were removed from the dataset. Higher taxonomic assignments excluding the genus *Anas* were then removed, thus taxonomic assignments in the final dataset were predominantly of species resolution. The read count data were converted to a binary species matrix for downstream analyses using the package cooccur v1.3 (Griffith, Veech & Marsh, 2016) and Generalized Linear Mixed-effects Models (GLMMs), which were fit by maximum likelihood with the glmer function using the package lme4 v1.1-21 (Bates et al., 2015). County was included as a random effect in each GLMM to account for spatial dependencies between ponds (see Supporting Information: Appendix 1).

2.3.2 Hypothesis generation: biotic factors potentially influencing T. cristatus detection

We first mined the eDNA metabarcoding data and investigated the influence of vertebrate group richness on *T. cristatus* detection. We used a binomial GLMM with the logit link function that included species richness of other amphibians, fish, waterfowl, terrestrial birds, and mammals as fixed effects (model 1, N = 532). We then performed a preliminary analysis using the package cooccur v1.3 (Griffith, Veech & Marsh, 2016) to identify species associations between *T. cristatus* and other vertebrates (N = 532). Hypotheses generated by the eDNA metabarcoding data relating to factors influencing *T. cristatus* detection are summarised in Table 1.

2.3.3 Hypothesis testing: biotic and abiotic factors known to influence T. cristatus detection

Identified associations from the cooccur analysis in conjunction with the existing *T. cristatus* literature informed candidate biotic variables to be modelled against *T. cristatus* detection (model 2, n = 504). The existing *T. cristatus* literature informed candidate abiotic variables to be modelled against *T. cristatus* detection (model 3, n = 504). Hypotheses tested using the eDNA metabarcoding data and associated environmental metadata to examine biotic and biotic factors influencing *T. cristatus* detection are summarised in Table 2. Selection of a suitable set of explanatory variables and modelling framework is fully described in Supporting Information: Appendix 1. Briefly, candidate biotic and abiotic explanatory variables were assessed for: collinearity using a Spearman's rank pairwise correlation matrix and examination of variance inflation factors (Zuur et al., 2009), relative importance using a

classification tree within the package rpart v4.1-13 (Therneau, Atkinson & Ripley, 2014), and

non-linearity using Generalized Additive Models (GAMs) (Zuur et al., 2009). We constructed

separate binomial GLMMs with the logit link function for biotic (model 2) and abiotic

(model 3) explanatory variables that passed these assessments.

2.3.4 Hypothesis testing: T. cristatus HSI and umbrella status

We modelled HSI score (fixed effect) separately to prevent HSI score masking variation

caused by the individual biotic and abiotic variables it encompasses (model 4). Using a

Poisson GLMM, we tested the umbrella species status of *T. cristatus* by modelling

vertebrate species richness against *T. cristatus* detection and the *T. cristatus* HSI score

(model 5).

2.3.5 Model selection and validation

For each GLMM, we employed an information-theoretic approach using Akaike Information

Criterion (AIC) to determine the most parsimonious approximating model to make

predictions (Akaike, 1973). Biotic and abiotic models considered respectively were nested

thus the best models were chosen using stepwise backward deletion of terms based on

Likelihood Ratio Tests (LRTs). Models 1, 4 and 5 were compared to null GLMMs. The final

models were as follows:

Model 1: *T. cristatus* ~ (1|County) + amphibian richness + fish richness + waterfowl richness

+ terrestrial bird richness + mammal richness

Model 2: *T. cristatus* ~ (1|County) + *L. vulgaris* + *B. bufo* + *C. carpio* + *G. aculeatus* + *Gallinula*

chloropus + Waterfowl

Model 3: *T. cristatus* ~ (1|County) + inflow + pond area + shading

Model 4: *T. cristatus* ~ (1|County) + HSI score

Model 5: Vertebrate species richness ~ (1|County) + *T. cristatus* + HSI score

Final models were tested for overdispersion using a custom function testing overdispersion

of the Pearson residuals. Model fit was assessed using the Hosmer and Lemeshow Goodness

of Fit Test within the package ResourceSelection v0.3-2 (Lele, Keim & Solymos, 2016),

quantile-quantile plots, and partial residual plots (Zuur et al., 2009). Conditional R^2 was

calculated using the r.squaredGLMM function in the package MuMIn v1.42.1 (Bartoń, 2018).

The Benjamini-Hochberg correction was calculated for explanatory variable p-values from

the cooccur and GLMM analyses using the p.adjust function in the package stats v3.4.3, and

this alpha applied to the results to account for potential Type I error. The original p-values

and the adjusted Benjamini-Hochberg p-values are reported for comparison in Tables 1 and

2. Model predictions were obtained using the predictSE function in the package AICcmodavg

v2.1-1 (Mazerolle, 2016) and upper and lower 95% CIs were calculated from the standard

error of the predictions. Results were plotted using the package ggplot2 v3.1.0 (Wickham,

2016).

3. Results

3.1 eDNA metabarcoding

Harper et al. (2018) processed a total of 532 eDNA samples and 228 PCR controls across two sequencing runs. The runs generated raw sequence read counts of 36,236,862 and 32,900,914 respectively. After quality filtering, trimming, and merging of paired-end reads, 26,294,906 and 26,451,564 sequences remained. Following removal of chimeras and read dereplication via clustering, the libraries contained 14,141,237 and 14,081,939 sequences (average read counts of 36,826 and 36,671 per sample respectively), of which 13,126,148 and 13,113,143 sequences were taxonomically assigned. The final dataset used here (assignments corrected, thresholds applied, and assignments removed) contained 53 vertebrate species (Table S5), including six amphibians, 14 fish, 17 birds, and 16 mammals (Fig. 1).

3.2 Pondscape biodiversity

All native amphibians were detected as well as the non-native marsh frog (*Pelophylax ridibundus*). *T. cristatus* (n = 148), *L. vulgaris* (n = 151) and *R. temporaria* (n = 122) were widespread, but *B. bufo* (n = 42), palmate newt (*Lissotriton helveticus* [n = 5]) and *P. ridibundus* were uncommon (n = 1). The threatened European eel (*Anguilla anguilla* [n = 15]), European bullhead (*Cottus gobio* [n = 14]), and *C. carassius* (n = 2) were detected alongside native fishes, such as pike (*Esox Lucius* [n = 17]) and roach (*Rutilus rutilus* [n = 71]), but also introduced species, including *C. carpio* (n = 40), ruffe (*Gymnocephalus cernua* [n =

1]), and rainbow trout (*Oncorhynchus mykiss* [*n* = 3]). Some identified waterfowl were

ubiquitous, such as common moorhen (*Gallinula chloropus* [*n* = 211]), whereas others were

less common, e.g. grey heron (*Ardea cinerea* [*n* = 1]) and Eurasian oystercatcher

(*Haematopus ostralegus* [*n* = 1]). Terrestrial fauna were often detected in fewer than five

ponds (Figs. 1c, d). Buzzard (*Buteo buteo* [*n* = 4]), Eurasian jay (*Garrulus glandarius* [*n* = 7]),

dunnock (*Prunella modularis* [*n* = 4]), and starling (*Sturnus vulgaris* [*n* = 4]) were the most

frequently detected terrestrial birds. Introduced mammals (Mathews et al., 2018), such as

grey squirrel (*Sciurus carolinensis* [*n* = 57]) and Reeve's muntjac (*Muntiacus reevesi* [*n* = 3]),

outweighed native mammals. Nonetheless, we detected several mammals with Biodiversity

Actions Plans and/or of conservation concern (Mathews et al., 2018), including otter (*Lutra*

lutra [*n* = 1]), water vole (*Arvicola amphibius* [*n* = 16]), European polecat (*Mustela putorius*

[*n* = 1]), brown hare (*Lepus europaeus* [*n* = 1]) and water shrew (*Neomys fodiens* [*n* = 8]).

Notably, the invasive American mink (*Neovison vison*) was not detected despite widespread

UK distribution (Mathews et al., 2018). All species and their detection frequencies are listed

in Table S5.

3.3 Hypothesis generation: biotic factors potentially influencing *T. cristatus* detection

T. cristatus detection was positively influenced by amphibian and waterfowl species

richness, yet negatively influenced by mammal species richness. *T. cristatus* detection was

less probable as fish and terrestrial bird species richness increased, but these trends were

not significant (Fig. 2; GLMM: overdispersion θ = 0.995, χ^2_{525} = 522.276, *P* = 0.525; fit χ^2_8 =

1.568, *P* = 0.992, R^2 = 12.74%). With no correction for Type I error, *T. cristatus* had

significant positive associations with three species (Fig. S5), including *L. vulgaris*, common

coot (*Fulica atra*) and *G. chloropus*, and significant negative associations with six species

(Fig. S5), including *B. bufo*, *C. carpio*, *G. aculeatus*, *P. pungitius*, common pheasant

(*Phasianus colchicus*) and *S. carolinensis*. However, only *T. cristatus* associations with *L.*

vulgaris and *G. chloropus* were retained after applying the Benjamini-Hochberg correction.

T. cristatus detection was more likely in ponds with more amphibian species (Fig.

2a). *T. cristatus* was detected in 51.66% of ponds (*n* = 151) identified as containing *L.*

vulgaris, but in only 11.91% of ponds (*n* = 42) where *B. bufo* was detected (Fig. 1a).

Probability of *T. cristatus* detection was lower in ponds with more fish species, and *T.*

cristatus was not detected in ponds with more than four fish species (Fig. 2b). *T. cristatus*

was only detected in 15.00% (*n* = 40), 14.55% (*n* = 55) and 6.67% (*n* = 15) of ponds identified

as containing *C. carpio*, *G. aculeatus* and *P. pungitius* respectively (Fig. 1b). In contrast, *T.*

cristatus detection was more likely in ponds where more waterfowl species were detected

(Fig. 2c). *T. cristatus* was detected in 41.67% (*n* = 48) and 36.02% (*n* = 211) of ponds

identified as containing *F. atra* and *G. chloropus* respectively (Fig. 1c). *T. cristatus* detection

was negatively influenced by higher terrestrial bird species richness, but not significantly so

(Fig. 2d). However, *T. cristatus* was only detected in 12.00% (*n* = 25) ponds where *P.*

colchicus was detected (Fig. 1c). *T. cristatus* detection was less likely in ponds where more

mammal species were detected (Figs. 2e). Specifically, *T. cristatus* was only detected in

15.79% (*n* = 57) of ponds where *S. carolinensis* was detected (Fig. 1d).

3.4 Hypothesis testing: biotic and abiotic factors known to influence *T. cristatus* detection

Biotic variables put forward for model selection were eDNA-based detection/non-detection

of *L. vulgaris*, *B. bufo*, *C. carpio*, *G. aculeatus*, *P. pungitius*, *G. chloropus*, and *F. atra* as well

as presence/absence of other amphibians, fish, and waterfowl as recorded during HSI

assessment by environmental consultants. Only detection/non-detection of *L. vulgaris*, *B.*

bufo, *C. carpio*, *G. aculeatus*, and *G. chloropus* as well as waterfowl presence were retained

by model selection as explanatory variables for the biotic GLMM of *T. cristatus* detection

(Figs. 3a-f; GLMM: overdispersion θ = 1.013, χ^2_{495} = 495.000, *P* = 0.411; fit χ^2_8 = 4.437, *P* =

0.8157, R^2 = 27.77%). Abiotic variables put forward for model selection were max. depth,

pond density, presence of inflow, pond area, pond substrate, presence of outflow,

percentage of macrophyte cover, water quality, pond permanence, percentage of shading,

ruderals, scrub/hedge, and woodland. Only presence of inflow, pond area, and percentage

of shading were retained by model selection as explanatory variables for the abiotic GLMM

explaining *T. cristatus* detection (Figs. 3g-i; GLMM: overdispersion θ = 1.000, χ^2_{499} = 499.126,

P = 0.490; fit χ^2_8 = 4.898, *P* = 0.768, R^2 = 10.00%). Results are summarised and relationships

compared to those previously reported for *T. cristatus* in Table 2. Application of the

Benjamini-Hochberg correction for Type I error did not alter variable significance.

3.5 Hypothesis testing: *T. cristatus* HSI and umbrella status

HSI score positively correlated with *T. cristatus* detection probability (GLMM: overdispersion

θ = 0.995, χ^2_{501} = 498.372, *P* = 0.525; fit χ^2_8 = 5.395, *P* = 0.715, R^2 = 6.66%), where *T. cristatus*

detection was more likely in ponds with a higher HSI score (Fig. 3j; estimate ± standard error

= 3.094 ± 0.803, χ^2_1 = 16.020, *P* < 0.001). Vertebrate species richness was positively

associated with *T. cristatus* detection (GLMM: overdispersion θ = 1.112, χ^2_{500} = 555.800, *P* =

0.042; fit χ^2_8 = -11.91, *P* = 1.000, R^2 = 9.14%), with more species detected in ponds identified

as containing *T. cristatus* (Fig. 4a; estimate ± standard error = 0.408 ± 0.059, χ^2_1 = 46.265, *P*

< 0.001, adjusted *P* [Benjamini-Hochberg] < 0.001). However, vertebrate species richness

did not significantly increase with the *T. cristatus* HSI score (Fig. 4b; estimate ± standard

error = 0.228 ± 0.206, χ^2_1 = 1.227, *P* = 0.268, adjusted *P* [Benjamini-Hochberg] = 0.268).

4. Discussion

We have validated eDNA metabarcoding for ecological hypothesis generation and testing

using the community data produced by this tool in combination with environmental

metadata for ponds. We identified and tested biotic and abiotic factors influencing *T.*

cristatus detection, whether the HSI can be applied to *T. cristatus* eDNA survey, and

whether *T. cristatus* is truly an umbrella species for pond conservation. *T. cristatus*

detection was more likely in ponds where *L. vulgaris* and *G. chloropus* were detected, and

ponds where *B. bufo*, *C. carpio*, and *G. aculeatus* were not detected. *T. cristatus* detection

was also more likely in ponds where waterfowl presence was recorded during HSI

assessment. Ponds identified as containing *T. cristatus* were typically small, absent of inflow,

and not excessively shaded. The *T. cristatus* HSI correlated with *T. cristatus* detection, but

not vertebrate species richness. Nonetheless, more vertebrates were detected in ponds

where *T. cristatus* was detected thus presence of this amphibian may indicate good quality

habitat for other vertebrates. Our findings demonstrate the power of eDNA metabarcoding

to enhance freshwater monitoring and research by providing biodiversity data *en masse* at

low cost.

4.1 Pondscape biodiversity

eDNA metabarcoding detected six amphibian, 14 fish, 17 bird, and 16 mammal species across 532 UK ponds. This diverse species inventory emphasises the importance of ponds as habitat for aquatic taxa, but also as stepping stones for semi-aquatic and terrestrial taxa (De Meester et al., 2005; Hill et al., 2018) through provision of drinking, foraging, dispersive, and reproductive opportunities (Biggs et al., 2016; Klymus et al., 2017). Some species detections may be the result of eDNA transport from water bodies in the surrounding area (Hänfling et al., 2016) to ponds via inflow. However, this signifies the capacity of ponds to provide natural samples of freshwater and terrestrial biodiversity in the wider catchment (Deiner et al., 2017).

4.2 Biotic factors influencing *T. cristatus* detection

T. cristatus detection was more likely in ponds with higher amphibian species richness - particularly ponds identified as containing *L. vulgaris* but not *B. bufo*. *T. cristatus* and *L. vulgaris* have similar habitat requirements and tend to breed in the same ponds (Skei et al., 2006; Rannap et al., 2009a; Denoël et al., 2013; Cayuela et al., 2018), with >60% overlap reported (Rannap & Briggs, 2006). However, *L. vulgaris* can inhabit a broader range of habitat (Rannap & Briggs, 2006; Skei et al., 2006) than *T. cristatus*, which depends on larger, deeper ponds with abundant macrophytes and no fish located in open, semi-rural landscapes (Denoël et al., 2013). *B. bufo* can inhabit fish-containing ponds (Manenti & Pennati, 2016) and *T. cristatus* may predate *B. bufo* eggs and larvae (Langton et al., 2001). This may explain the negative association between *B. bufo* and *T. cristatus* as opposed to

the positively associated *T. cristatus* and *L. vulgaris*.

T. cristatus detection marginally decreased with higher fish species richness, and *T.*

cristatus was negatively associated with *C. carpio*, *G. aculeatus*, and *P. pungitius*. These

fishes are common in and typical of ponds. All *T. cristatus* life stages may be predated by

fishes (Langton et al., 2001) and negative effects of fish presence-absence on *T. cristatus*

occupancy, distribution, and abundance are repeatedly reported (Joly et al., 2001; Rannap &

Briggs, 2006; Skei et al., 2006; Denoël & Ficetola, 2008; Rannap et al., 2009a, b; Hartel et al.,

2010; Denoël et al., 2013). *G. aculeatus* predates *T. cristatus* eggs and larvae (McLee &

Scaife, 1992; Jarvis, 2010), and has non-consumptive effects on *T. cristatus* embryos (Jarvis,

2010). *T. cristatus* larvae were also found to alter their behaviour when exposed to

predatory *G. aculeatus* but not non-predatory *C. carassius* (Jarvis, 2012), another fish

characteristic of ponds.

In our study, we detected *T. cristatus* in 50% of ponds where *C. carassius* was

detected, but <20% of ponds where large and/or predatory fishes were detected, e.g. *C.*

carpio, *G. aculeatus*, *E. lucius*. Although we observed fewer detections from ponds for *C.*

carassius than *C. carpio*, *G. aculeatus* or *E. lucius*, previous research also indicates large

and/or predatory fish are more detrimental to *T. cristatus* occurrence (Skei et al., 2006;

Hartel et al., 2010; Chan, 2011). *C. carassius* does not hinder *T. cristatus* oviposition, larval

behaviour, or recruitment success (Chan, 2011; Jarvis, 2012), or pond invertebrate and

macrophyte diversity (Stefanoudis et al., 2017). In contrast, *C. carpio* foraging reduces

invertebrate density and macrophyte cover (Maceda-Veiga, López & Green, 2017), which

lowers *T. cristatus* reproductive and foraging success and heightens predator exposure

(Rannap & Briggs, 2006; Gustafson et al., 2006; Chan, 2011). *C. carassius* and *C. carpio* are

both included among fish species assumed to negatively impact *T. cristatus* and whose

presence-absence is assessed for the *T. cristatus* HSI (ARG-UK, 2010). However, given that *C.*

carassius does not directly predate *T. cristatus* or indirectly alter its behaviour, reproductive

success, or habitat, we advocate a systematic re-evaluation of problematic fish species for *T.*

cristatus conservation.

T. cristatus detection was positively associated with waterfowl species richness,

namely *F. atra* and *G. chloropus* detection. These waterfowl species share macrophytes and

macroinvertebrates as resources with amphibians, feeding on both directly (Perrow et al.,

1997; Paillisson & Marion, 2001; Wallau et al., 2010). *F. atra* and *G. chloropus* crop

emergent macrophytes to search for invertebrate prey (Paillisson & Marion, 2001; Wallau et

al., 2010), which may indirectly benefit *T. cristatus* foraging. Although *Fulica* spp. can also

pull up submerged vegetation and damage vegetation banks (Lauridsen, Jeppesen &

Andersen, 1993), diet is macrophyte-dominated in late summer and autumn (Perrow et al.,

1997) and unlikely to impact *T. cristatus* breeding in spring (Langton et al., 2001). The

positive association identified here between *T. cristatus* and these waterfowl most likely

reflects a shared preference for macrophyte-rich ponds.

T. cristatus detection was less frequent in ponds with higher mammal species

richness. Our preliminary cooccur analysis indicated *T. cristatus* had negative associations

with *P. colchicus* and *S. carolinensis,* but not when a correction for Type I error was applied.

T. cristatus associations with *F. atra*, *B. bufo*, *C. carpio*, *G. aculeatus*, and *P. pungitius* were

also non-significant when this correction was applied. These associations require further

investigation to determine whether they are real, and if so, whether they are direct and

reflect competition, predation or shared resources as discussed above, or indirect and

reflect land-use or effects of additional species.

4.3 Abiotic factors influencing *T. cristatus* detection

T. cristatus detection was less likely in large ponds with inflow present and a greater percentage of shading. Effects of pond area may depend on the size range of ponds studied. Although our results indicate *T. cristatus* prefers smaller ponds, pond area does not always influence occupancy (Maletzky, Kyek & Goldschmid, 2007; Denoël & Ficetola, 2008; Gustafson et al., 2011) and was deemed a poor predictor of reproductive success (Vuorio et al., 2013). *T. cristatus* has been found to utilise small and large ponds (Rannap & Briggs, 2006; Skei et al., 2006); however, very small ponds (<124 m^2) may be unable to support all life stages, and larger ponds may contain fish and experience eutrophication due to agricultural or polluted run-off (Rannap & Briggs, 2006). Inflow to ponds may exacerbate these problems by facilitating entry of agricultural or polluted run-off and connections to streams and rivers containing large, predatory fish (Freshwater Habitats Trust, 2015). Our results corroborate existing research where viable *T. cristatus* populations were unlikely in ponds that were shaded (Vuorio et al., 2013) or had dense macrophyte cover (Rannap & Briggs, 2006; Skei et al., 2006; Hartel et al., 2010).

In our study, most environmental metadata available were qualitative, preventing detailed analyses on pond properties and terrestrial habitat in relation to *T. cristatus* detection. Better understanding of *T. cristatus* detection in relation to species interactions and habitat quality could be achieved with quantitative data on pond properties (e.g. water chemistry), terrestrial habitat (e.g. type, density, distance to ponds), and aquatic and terrestrial habitat usage by different vertebrate species. Furthermore, given the metapopulation dynamics of *T. cristatus*, future research should investigate spatial drivers (e.g. pond density and other indices of connectivity, land cover, climate variables, roads,

rivers, elevation) of *T. cristatus* detection using innovative modelling approaches, such as

individual-based models (Messager & Olden, 2018). However, acquiring this data to perform

these models is a phenomenal task for large numbers of ponds across a vast landscape

(Denoël & Ficetola, 2008).

4.4 *T. cristatus* HSI and umbrella status

We found the HSI can predict eDNA-based *T. cristatus* detection at the UK pondscape. This

contradicts conventional studies which deemed the index inappropriate for predicting *T.*

cristatus occupancy or survival probabilities (Unglaub et al., 2015). We detected more

vertebrates in ponds identified as containing *T. cristatus*, which may support its status as an

umbrella species for pond biodiversity and conservation (Gustafson et al., 2006). We also

observed a non-significant increase in vertebrate species richness with increasing *T.*

cristatus HSI score. An adapted HSI, designed to predict species richness, could help select

areas for management and enhancement of aquatic and terrestrial biodiversity. Until then,

presence of *T. cristatus* and its HSI may confer protection to broader biodiversity by

identifying optimal habitat for pond creation and restoration to encourage populations of

this threatened amphibian. The HSI is not without issue due to qualitative data used for

score calculation and subjective estimation of indices (Oldham et al., 2000; O'Brien et al.,

2017). For future application of this index in *T. cristatus* eDNA survey, we recommend

metabarcoding to quantify some qualitatively assessed indices (e.g. water quality via

macroinvertebrate diversity, fish and waterfowl presence) alongside *T. cristatus* detection.

Provided rigorous spatial and temporal sampling are undertaken, eDNA metabarcoding can

also generate site occupancy data to estimate relative species abundance (Valentini et al.,

2016; Hänfling et al., 2016; Lawson Handley et al., 2019; Li et al., 2019a).

4.5 Limitations of repurposed eDNA samples for metabarcoding applications

This study was based on samples that were repurposed from targeted eDNA surveys for *T.*

cristatus. eDNA sampling, capture, and extraction was conducted in accordance with Biggs

et al. (2015), whose methods were chosen based on the eDNA literature at that time. Six 15

ml water samples were taken from a homogenised sample (600 ml). Ethanol precipitation

was used for eDNA capture, followed by DNA extraction on the combined lysate from the six

subsamples. Consequently, ponds were represented by a single eDNA sample, for which 12

qPCR replicates were performed (Biggs et al., 2015). However, eDNA metabarcoding may

require an entirely different workflow to enhance species detection (Harper et al., 2018;

Harper et al., 2019).

Independent biological replicates as opposed to pseudoreplicates from a single

water sample are key to account for spatial heterogeneity of eDNA from different species

(Hänfling et al., 2016; Bálint et al., 2018; Lawson Handley et al., 2019; Li et al., 2019b).

Larger volumes should be filtered instead of ethanol precipitation on small volumes to

maximise eDNA capture (Harper et al., 2019). Samples should be extracted individually to

maintain biological replication followed by independent technical replicates for PCR and

sequencing. These levels of replication are required for hierarchical occupancy modelling to

identify false positives and species detection probabilities as well as minimise false

negatives (Ficetola et al., 2015; Bálint et al., 2018; Dorazio & Erickson, 2018).

We performed three PCR replicates for each sample that were pooled prior to

sequencing. We included a large number of PCR controls and applied sequence thresholds

to control for any false positives in our dataset arising from potential laboratory

contamination. Pooling of technical replicates allowed us to screen vertebrate communities

from over 500 ponds, but prevented occupancy modelling to verify species detection.

Samples collected for targeted eDNA analysis can vary widely in terms of sampling strategy,

capture, extraction and storage protocols. Ultimately, these protocols may limit the data

generated by eDNA metabarcoding and the analyses that can be performed. This is an

important consideration for any future research where eDNA samples are repurposed to

address ecological questions.

4.6 Prospects of eDNA metabarcoding for freshwater conservation, management, and

research

We have demonstrated the effectiveness of eDNA metabarcoding for landscape-scale

biodiversity monitoring as well as ecological hypothesis generation and testing. We

combined metabarcoding with environmental metadata to identify new and revisit old

hypotheses relating to biotic and abiotic factors that influence a threatened amphibian at

the UK pondscape. Our findings will guide *T. cristatus* conservation in the face of increasing

land-use and habitat fragmentation - a poignant issue as protective legislation for this

species in the UK is changing. Whilst conservation of threatened species and their habitat

should be a priority, the bigger picture should not be ignored. eDNA metabarcoding could

enhance our understanding of freshwater networks, particularly pondscapes, to enable

more effective monitoring, protection, and management of aquatic and terrestrial

biodiversity. We are only now beginning to realise and explore these opportunities.

Acknowledgements

This work was funded by the University of Hull. We would like to thank Jennifer Hodgetts

(Fera Science Ltd) for assisting with sample collection, and Jianlong Li (University of Hull) for

primer design and advice on laboratory protocols. Tissue samples for primer validation and

Sanger sequencing were provided by Andrew Buxton and Richard Griffiths (DICE, University

of Kent) under licence from Natural England, and Barbara Mabel and Elizabeth Kilbride

(University of Glasgow).

Data Availability

The taxonomically assigned sequence reads used in this study were generated by Harper et

al. (2018). The raw sequence reads were archived on the NCBI Sequence Read Archive

(Bioproject: PRJNA417951; SRA accessions: SRR6285413 - SRR6285678). The bioinformatics

analysis was deposited in a GitHub repository and permanently archived

(https://doi.org/10.5281/zenodo.1188710). R scripts and corresponding data for this study

have been deposited in a separate GitHub repository, which has been permanently archived

(https://doi.org/10.5281/zenodo.2634427).

Author Contributions

B.H, L.R.H, L.L.H and N.B conceived and designed the study. H.C.R and N.B contributed samples for processing. L.R.H performed laboratory work and analysed the data. I.P.A and E.L offered advice on and supervised sequencing. C.H assisted with bioinformatics analysis. P.B and S.P contributed datasets for analysis. L.R.H wrote the manuscript, which all authors revised.

References

Akaike, H. (1973). Maximum likelihood identification of Gaussian autoregressive moving average models. *Biometrika*, **60**, 255–265. https://doi.org/10.2307/2334537

ARG-UK. (2010). *ARG UK Advice Note 5: Great Crested Newt Habitat Suitability Index*. Amphibian and Reptile Groups of the United Kingdom.

Bálint, M., Nowak, C., Márton, O., Pauls, S. U., Wittwer, C., Aramayo, J. L., Schulze, A., Chambert, T., Cocchiararo, B., & Jansen, M. (2018). Accuracy, limitations and cost efficiency of eDNA-based community survey in tropical frogs. *Molecular Ecology Resources*, **18**, 1415–1426. https://doi.org/10.1111/1755-0998.12934

Barnes, M. A., & Turner, C. R. (2015). The ecology of environmental DNA and implications for conservation genetics. *Conservation Genetics*, **17**, 1–17. https://doi.org/10.1007/s10592-015-775-4

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, **67**, 1–48. https://doi.org/10.18637/jss.v067.i01

Biggs, J., Ewald, N., Valentini, A., Gaboriaud, C., Dejean, T., Griffiths, R. A., Foster, J., Wilkinson, J. W., Arnell, A., Brotherton, P., Williams, P., & Dunn, F. (2015). Using eDNA to develop a national citizen science-based monitoring programme for the great crested newt (*Triturus cristatus*). *Biological Conservation*, **183**, 19–28. https://doi.org/10.1016/j.biocon.2014.11.029

Biggs, J., von Fumetti, S., & Kelly-Quinn, M. (2016). The importance of small waterbodies for biodiversity and ecosystem services: implications for policy makers. *Hydrobiologia*, **793**, 3–39. https://doi.org/10.1007/s10750-016-3007-0

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*, **30**, 2114–2120. https://doi.org/10.1093/bioinformatics/btu170

Cayuela, H., Grolet, O., & Joly, P. (2018). Context-dependent dispersal, public information, and heterospecific attraction in newts. *Oecologia*, **188**, 1069–1080. https://doi.org/10.1007/s00442-018-4267-3

Chan, K. (2011). Can great crested newts (*Triturus cristatus*) coexist with fish?. Masters Thesis. University College London (UCL).

Deiner, K., Bik, H. M., Mächler, E., Seymour, M., Lacoursière-Roussel, A., Altermatt, F., Creer, S., Bista, I., Lodge, D. M., de Vere, N., Pfrender, M. E., & Bernatchez, L. (2017). Environmental DNA metabarcoding: transforming how we survey animal and plant communities. *Molecular Ecology*, **26**, 5872–5895. https://doi.org/10.1111/mec.14350

De Meester, L., Declerck, S., Stoks, R., Louette, G., Van De Meutter, F., De Bie, T., Michels, E., & Brendonck, L. (2005). Ponds and pools as model systems in conservation biology, ecology and evolutionary biology. *Aquatic Conservation: Marine and Freshwater Ecosystems*, **15**, 715–725. https://doi.org/10.1002/aqc.748

Denoël, M., & Ficetola, G. F. (2008). Conservation of newt guilds in an agricultural landscape of Belgium: the importance of aquatic and terrestrial habitats. *Aquatic Conservation: Marine and Freshwater Ecosystems*, **18**, 714–728. https://doi.org/10.1002/aqc.853

Denoël, M., Perez, A., Cornet, Y., & Ficetola, G. F. (2013). Similar local and landscape processes affect both a common and a rare newt species. *PloS ONE*, **8**, e62727. https://doi.org/10.1371/journal.pone.0062727

Dorazio, R.M. & Erickson, R.A. (2018) ednaoccupancy: An r package for multiscale occupancy modelling of environmental DNA data. *Molecular Ecology Resources*, **18**, 368–380. https://doi.org/10.1111/1755-0998.12735

Edgar, P., & Bird, D. R. (2006). *Action Plan for the Conservation of the Crested Newt Triturus Cristatus Species Complex in Europe*. Council of the European Union, Strassbourg, Germany.

Edgar, R. C., Haas, B. J., Clemente, J. C., Quince, C., & Knight, R. (2011). UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics*, **27**, 2194–2200. https://doi.org/10.1093/bioinformatics/btr381

Evans, N. T., Li, Y., Renshaw, M. A., Olds, B. P., Deiner, K., Turner, C. R., Jerde, C. L., Lodge, D. M., Lamberti, G. A., & Pfrender, M. E. (2017). Fish community assessment with eDNA metabarcoding: effects of sampling design and bioinformatic filtering. *Canadian Journal of Fisheries and Aquatic Sciences*, **74**, 1362-1374. https://doi.org/10.1139/cjfas-2016-0306

691 Ficetola, G. F., Coissac, E., Zundel, S., Riaz, T., Shehzad, W., Bessière, J., Taberlet, P., & Pompanon, F. (2010). An *In silico* approach for the evaluation of DNA barcodes. *BMC Genomics*, **11**, 434. https://doi.org/10.1186/1471-2164-11-434

694 Ficetola, G.F., Pansu, J., Bonin, A., Coissac, E., Giguet-Covex, C., De Barba, M., Gielly, L., Lopes, C.M., Boyer, F., Pompanon, F., Rayé, G. & Taberlet, P. (2015) Replication levels, false presences and the estimation of the presence/absence from eDNA metabarcoding data. *Molecular Ecology Resources*, **15**, 543–556. https://doi.org/10.1111/1755-0998.12338

Freshwater Habitats Trust (2015). Pond Habitat Survey Method Booklet. Freshwater Habitats Trust, Oxford.

Griffith, D., Veech, J., & Marsh, C. (2016). cooccur: Probabilistic Species Co-Occurrence Analysis in R. *Journal of Statistical Software*, **69**, 1–17. https://doi.org/10.18637/jss.v069.c02

Gustafson, D. H., Pettersson, C. J., & Malmgren, J. C. (2006). Great crested newts (*Triturus*
cristatus) as indicators of aquatic plant diversity. *The Herpetological Journal*, **16**, 347–
352.

Gustafson, D. H., Malmgren, J. C., & Mikusiński, G. (2011). Terrestrial Habitat Predicts use of
Aquatic Habitat for Breeding Purposes — A Study on the Great Crested Newt (*Triturus*
cristatus). *Annales Zoologici Fennici*, **48**, 295–307. https://doi.org/10.5735/086.048.0504

Hänfling, B., Lawson Handley, L., Read, D. S., Hahn, C., Li, J., Nichols, P., Blackman, R. C.,
Oliver, A., & Winfield, I. J. (2016). Environmental DNA metabarcoding of lake fish
communities reflects long-term data from established survey methods. *Molecular*
Ecology, **25**, 3101–3119. https://doi.org/10.1111/mec.13660

Harper, L. R., Lawson Handley, L., Hahn, C., Boonham, N., Rees, H. C., Gough, K. C., Lewis, E.,
Adams, I. P., Brotherton, P., Phillips, S., & Hänfling, B. (2018). Needle in a haystack? A
comparison of eDNA metabarcoding and targeted qPCR for detection of the great crested
newt (*Triturus cristatus*). *Ecology and Evolution*, *8*, 6330–6341.
https://doi.org/10.1002/ece3.4013

Harper, L. R., Buxton, A. S., Rees, H. C., Bruce, K., Brys, R., Halfmaerten, D., Read, D. S.,
Watson, H. V., Sayer, C. D., Jones, E. P., Priestley, V., Mächler, E., Múrria, C., Garcés-
Pastor, S., Medupin, C., Burgess, K., Benson, G., Boonham, N., Griffiths, R. A., Lawson
Handley, L., & Hänfling, B. (2019). Prospects and challenges of environmental DNA
(eDNA) monitoring in freshwater ponds. *Hydrobiologia*, **826**, 25–41.
https://doi.org/10.1007/s10750-018-3750-5

Hartel, T., Nemes, S., & Oellerer, K. (2010). Using connectivity metrics and niche modelling
to explore the occurrence of the northern crested newt *Triturus cristatus* (Amphibia,
Caudata) in a traditionally managed landscape. *The Environmentalist*, **37**, 195–200.
https://doi.org/10.1017/S037689291000055X

Hering, D., Borja, A., Jones, J. I., Pont, D., Boets, P., Bouchez, A., Bruce, K., Drakare, S.,
Hänfling, B., Kahlert, M., Leese, F., Meissner, K., Mergen, P., Reyjol, Y., Segurado, P.,
Vogler, A., & Kelly, M. (2018). Implementation options for DNA-based identification into
ecological status assessment under the European Water Framework Directive. *Water*
Research, **138**, 192–205. https://doi.org/10.1016/j.watres.2018.03.003

Hill, M. J., Hassall, C., Oertli, B., Fahrig, L., Robson, B. J., Biggs, J., Samways, M. J., Usio, N.,
Takamura, N., Krishnaswamy, J., & Wood, P. J. (2018). New policy directions for global
pond conservation. *Conservation Letters*, **e12447**. https://doi.org/10.1111/conl.12447

Jarvis, L. E. (2010). Non-consumptive effects of predatory three-spined sticklebacks
(*Gasterosteus aculeatus*) on great crested newt (*Triturus cristatus*) embryos. *The*
Herpetological Journal, **20**, 271–275.

Jarvis, L. E. (2012). Microhabitat preferences of the great crested newt *Triturus cristatus* in a
woodland area. PhD Thesis. Open University.

Joly, P., Miaud, C., Lehmann, A., & Grolet, O. (2001). Habitat Matrix Effects on Pond
Occupancy in Newts. *Conservation Biology*, **15**, 239–248. https://doi.org/10.1111/j.1523-
1739.2001.99200.x

Bartoń, K. (2018). MuMIn: Multi-Model Inference. R package version 1.42.1.
https://CRAN.R-project.org/package=MuMIn

Kelly, R. P., O'Donnell, J. L., Lowell, N. C., Shelton, A. O., Samhouri, J. F., Hennessey, S. M.,
Feist, B. E., & Williams, G. D. (2016). Genetic signatures of ecological diversity along an
urbanization gradient. *PeerJ*, **4**, e2444. https://doi.org/10.7717/peerj.2444

Klymus, K. E., Richter, C. A., Thompson, N., & Hinck, J. E. (2017). Metabarcoding of Environmental DNA Samples to Explore the Use of Uranium Mine Containment Ponds as a Water Source for Wildlife. *Diversity*, **9**, 1-18. https://doi.org/10.3390/d9040054

Langton, T., Beckett, C., & Foster, J. (2001). Great crested newt conservation handbook. Froglife, Halesworth.

Lauridsen, T. L., Jeppesen, E., & Andersen, F. Ø. (1993). Colonization of submerged macrophytes in shallow fish manipulated Lake Væng: impact of sediment composition and waterfowl grazing. *Aquatic Botany*, **46**, 1–15. https://doi.org/10.1016/0304-3770(93)90061-Z

Lawson Handley, L. (2015). How will the "molecular revolution" contribute to biological recording? *Biological Journal of the Linnean Society,* 115, 750–766. https://doi.org/10.1111/bij.12516

Lawson Handley, L., Read, D.S., Winfield, I.J., Kimbell, H., Johnson, H., Li, J., Hahn, C., Blackman, R., Wilcox, R., Donnelly, R., Szitenberg, A. & Hänfling, B. (2019) Temporal and spatial variation in distribution of fish environmental DNA in England's largest lake. *Environmental DNA*, 12, e0176343. https://doi.org/10.1002/edn3.5

Lele, S. R., Keim, J. L., & Solymos, P. (2016). ResourceSelection: Resource Selection (Probability) Functions for Use-Availability Data.

Li, Y., Evans, N. T., Renshaw, M. A., Jerde, C. L., Olds, B. P., Shogren, A. J., Deiner, K., Lodge, D. M., Lamberti, G. A., & Pfrender, M. E. (2018a). Estimating fish alpha- and beta-diversity along a small stream with environmental DNA metabarcoding. *Metabarcoding and Metagenomics*, 2, e24262. https://doi.org/10.3897/mbmg.2.24262

Li, F., Peng, Y., Fang, W., Altermatt, F., Xie, Y., Yang, J., & Zhang, X. (2018b). Application of environmental DNA metabarcoding for predicting anthropogenic pollution in rivers. *Environmental Science & Technology*, 52, 11708–11719. https://doi.org/10.1021/acs.est.8b03869

Li, J., Hatton-Ellis, T.W., Lawson Handley, L.-J., Kimbell, H.S., Benucci, M., Peirson, G. & Hänfling, B. (2019a) Ground-truthing of a fish-based environmental DNA metabarcoding method for assessing the quality of lakes. *Journal of Applied Ecology*, **56**, 1232–1244. https://doi.org/10.1111/1365-2664.13352

Li, J., Lawson Handley, L.J., Harper, L.R., Brys, R., Watson, H.V., Di Muri, C., Zhang, X. & Hänfling, B. (2019b) Limited dispersion and quick degradation of environmental DNA in fish ponds inferred by metabarcoding. *Environmental DNA*, **5**, 133. https://doi.org/10.1002/edn3.24

Maceda-Veiga, A., López, R., & Green, A. J. (2017). Dramatic impact of alien carp *Cyprinus carpio* on globally threatened diving ducks and other waterbirds in Mediterranean shallow lakes. *Biological Conservation*, **212**, 74–85. https://doi.org/10.1016/j.biocon.2017.06.002

Macher, J. -N., Vivancos, A., Piggott, J. J., Centeno, F. C., Matthaei, C. D., & Leese, F. (2018). Comparison of environmental DNA and bulk-sample metabarcoding using highly degenerate cytochrome c oxidase I primers. *Molecular Ecology Resources*, **18**, 1456–1468. https://doi.org/10.1111/1755-0998.12940

Magoč, T., & Salzberg, S. L. (2011). FLASH: fast length adjustment of short reads to improve genome assemblies. *Bioinformatics*, **27**, 2957–2963. https://doi.org/10.1093/bioinformatics/btr507

Maletzky, A., Kyek, M., & Goldschmid, A. (2007). Monitoring status, habitat features and
amphibian species richness of crested newt (*Triturus cristatus* superspecies) ponds at the
edge of the species range (Salzburg, Austria). *Annales de Limnologie - International
Journal of Limnology*, **43**, 107–115. https://doi.org/10.1051/limn/2007017
Manenti, R., & Pennati, R. (2016). Environmental factors associated with amphibian
breeding in streams and springs: effects of habitat and fish occurrence. *Amphibia-
Reptilia*, **37**, 237–242. https://doi.org/10.1163/15685381-00003040
Mathews, F., Kubasiewicz, L. M., Gurnell, J., Harrower, C. A., McDonald, R. A., & Shore, R. F.
(2018). A Review of the Population and Conservation Status of British Mammals:
Technical Summary. A report by the Mammal Society under contract to Natural England,
Natural Resources Wales and Scottish Natural Heritage. Natural England, Peterborough.
Mazerolle, M. J. (2016). AICcmodavg: Model Selection and Multimodel Inference Based on
(Q)AIC(c).
McLee, A. G., & Scaife, R. W. (1992). The colonisation by great crested newts (*Triturus
cristatus*) of a water body following treatment with a piscicide to remove a large
population of sticklebacks (*Gasterosteus aculeatus*). *Herpetological Bulletin*, **42**, 6–9.
Messager, M. L., & Olden, J. D. (2018). Individual-based models forecast the spread and
inform the management of an emerging riverine invader. *Diversity & Distributions*, **24**,
1816–1829. https://doi.org/10.1111/ddi.12829
Nakagawa, H., Yamamoto, S., Sato, Y., Sado, T., Minamoto, T., & Miya, M. (2018). Comparing
local- and regional-scale estimations of the diversity of stream fish using eDNA
metabarcoding and conventional observation methods. *Freshwater Biology*, **63**, 569–580.
https://doi.org/10.1111/fwb.13094
O'Brien, D., Hall, J., Miró, A. & Wilkinson, J. (2017). Testing the validity of a commonly-used
habitat suitability index at the edge of a species' range: great crested newt *Triturus
cristatus* in Scotland. *Amphibia-Reptilia*, **38**, 265-273. https://doi.org/10.1163/15685381-
00003108
Oldham, R. S., Keeble, J., Swan, M. J. S., & Jeffcote, M. (2000). Evaluating the suitability of
habitat for the great crested newt (*Triturus cristatus*). *The Herpetological Journal*, **10**,
143–155.
Olds, B. P., Jerde, C. L., Renshaw, M. A., Li, Y., Evans, N. T., Turner, C. R., Deiner, K., Mahon,
A. R., Brueseke, M. A., Shirey, P. D., Pfrender, M. E., Lodge, D. M., & Lamberti, G. A.
(2016). Estimating species richness using environmental DNA. *Ecology and Evolution*, **6**,
4214–4226. https://doi.org/10.1002/ece3.2186
Paillisson, J. -M., & Marion, L. (2001). Interaction between coot (*Fulica atra*) and waterlily
(*Nymphaea alba*) in a lake: the indirect impact of foraging. *Aquatic Botany*, **71**, 209–216.
https://doi.org/10.1016/S0304-3770(01)00182-6
Perrow, M. R., Schutten, H. J., Howes, J. R., Holzer, T., Madgwick, F. J., & Jowitt, A. (1997).
Interactions between coot (*Fulica atra*) and submerged macrophytes: the role of birds in
the   restoration   process.   *Hydrobiologia*,   **342**,   241–255.
https://doi.org/10.1023/A:1017007911190
Rannap, R., & Briggs, L. (2006). THE CHARACTERISTICS OF GREAT CRESTED NEWT TRITURUS
CRISTATUS'BREEDING PONDS. Protection of the Great Crested Newt. Project Report. The
experiences of LIFE-Nature project "Protection of *Triturus cristatus* in the Eastern Baltic
Region" LIFE04NAT/EE/000070. Ministry of the Environment of the Republic of Estonia,
Tallinn.

Rannap, R., Lõhmus, A., & Briggs, L. (2009a). Niche position, but not niche breadth, differs in
two coexisting amphibians having contrasting trends in Europe. *Diversity & Distributions*,
15, 692–700. https://doi.org/10.1111/j.1472-4642.2009.00575.x

Rannap, R., Lõhmus, A., & Briggs, L. (2009b). Restoring ponds for amphibians: a success
story. *Hydrobiologia*, **634**, 87–95. https://doi.org/10.1007/s10750-009-9884-8

Rees, H. C., Maddison, B. C., Middleditch, D. J., Patmore, J. R. M., & Gough, K. C. (2014).
REVIEW: The detection of aquatic animal species using environmental DNA–a review of
eDNA as a survey tool in ecology. *Journal of Applied Ecology*, **51**, 1450–1459.
https://doi.org/10.1111/1365-2664.12306

Riaz, T., Shehzad, W., Viari, A., Pompanon, F., Taberlet, P., & Coissac, E. (2011). ecoPrimers:
inference of new DNA barcode markers from whole genome sequence analysis. *Nucleic
Acids Research*, **39**, e145. https://doi.org/10.1093/nar/gkr732

Rognes, T., Flouri, T., Nichols, B., Quince, C., & Mahé, F. (2016). VSEARCH: a versatile open
source tool for metagenomics. *PeerJ*, **4**, e2584. https://doi.org/10.7717/peerj.2584

Skei, J. K., Dolmen, D., Rønning, L., & Ringsby, T. H. (2006). Habitat use during the aquatic
phase of the newts *Triturus vulgaris* (L.) and *T. cristatus* (Laurenti) in central Norway:
proposition for a conservation and monitoring area. *Amphibia-Reptilia*, **27**, 309–324.
https://doi.org/10.1163/156853806778189972

Stefanoudis, P. V., Sayer, C. D., Greaves, H. M., Davidson, T. A., Robson, H., Almeida, D., &
Smith, E. (2017). Consequences of fish for cladoceran, water beetle and macrophyte
communities in a farmland pond landscape: implications for conservation. *Fundamental
and Applied Limnology*, **190**, 141–156. https://doi.org/10.1127/fal/2017/1004

Therneau, T., Atkinson, B. & Ripley, B. (2014). rpart: Recursive Partitioning and Regression
Trees. R package version 4.1-13.

Unglaub, B., Steinfartz, S., Drechsler, A., & Schmidt, B. R. (2015). Linking habitat suitability to
demography in a pond-breeding amphibian. *Frontiers in Zoology*, **12**.
https://doi.org/10.1186/s12983-015-0103-3

Ushio, M., Fukuda, H., Inoue, T., Makoto, K., Kishida, O., Sato, K., Murata, K., Nikaido, M.,
Sado, T., Sato, Y., Takeshita, M., Iwasaki, W., Yamanaka, H., Kondoh, M. & Miya, M.
(2017). Environmental DNA enables detection of terrestrial mammals from forest pond
water. *Molecular Ecology Resources*, **17**, e63-e65. https://doi.org/10.1111/1755-
0998.12690

Valentini, A., Taberlet, P., Miaud, C., Civade, R., Herder, J., Thomsen, P. F., Bellemain, E.,
Besnard, A., Coissac, E., Boyer, F., Gaboriaud, C., Jean, P., Poulet, N., Roset, N., Copp, G.
H., Geniez, P., Pont, D., Argillier, C., Baudoin, J. -M., Peroux, T., Crivelli, A. J., Olivier, A.,
Acqueberge, M., Le Brun, M., Møller, P. R., Willerslev, E. & Dejean, T. (2016). Next-
generation monitoring of aquatic biodiversity using environmental DNA metabarcoding.
Molecular Ecology, **25**, 929–942. https://doi.org/10.1111/mec.13428

Vuorio, V., Heikkinen, R. K., & Tikkanen, O. -P. (2013). Breeding Success of the Threatened
Great Crested Newt in Boreal Forest Ponds. *Annales Zoologici Fennici*, **50**, 158–169.
https://doi.org/10.5735/086.050.0303

Wallau, G. L., Della-Flora, F., Bueno, A. S., Corso, J., Ortiz, M. F., & Cáceres, N. C. (2010).
Behaviour of the Common Moorhen in Rio Grande do Sul, Brazil. *Acta Ethologica*, **13**,
127–135. https://doi.org/10.1007/s10211-010-0082-5

Wickham, H. (2009). *ggplot2: Elegant Graphics for Data Analysis*. Springer New York, USA.

Wilcox, T. M., Young, M. K., McKelvey, K. S., Isaak, D. J., Horan, D. L. & Schwartz, M. K.
(2018). Fine-scale environmental DNA sampling reveals climate-mediated interactions
between native and invasive trout species. *Ecosphere*, **9**, e02500.
https://doi.org/10.1002/ecs2.2500
Zhang, Z., Schwartz, S., Wagner, L., & Miller, W. (2000). A greedy algorithm for aligning DNA
sequences. *Journal of Computational Biology*, **7**, 203–214.
https://doi.org/10.1089/10665270050081478
Zuur, A. F., Ieno, E. N., Walker, N., Saveliev, A. A., & Smith, G. M. (2009). *Mixed effects
models and extensions in ecology with R*. Springer New York, USA.
895

Table 1. Summary of hypothesis generation using eDNA metabarcoding data, specifically
biotic factors that may influence *T. cristatus* detection probability. Direction of observed
effects on eDNA-based *T. cristatus* detection identified by the GLMM assessing species
richness in each vertebrate group (*N* = 532) and the preliminary cooccur analysis (*N* = 532)
are given. Negative and positive effects are listed as - and + respectively. Test statistic is for
LRT used and significant P-values (<0.05) are in bold.

Variable	Analysis							
	Cooccur			GLMM				
	Effect	*P*	Adjusted *P* (Benjamini-Hochberg)	DF	Effect size (SE)	χ²	*P*	Adjusted *P* (Benjamini-Hochberg)
Amphibians								
Species richness				1	0.526 (0.140)	14.195	**<0.001**	**<0.001**
L. vulgaris	+	**<0.001**	**<0.001**	1	1.616 (0.232)			
B. bufo	-	**0.010**	0.489	1	-1.793 (0.643)			
Fish								
Species richness				1	-0.163 (0.124)	1.798	0.180	0.225
G. aculeatus	-	**0.012**	0.489					
C. carpio	-	**0.040**	0.714					
P. pungitius	-	**0.049**	0.714					
C. carassius								
Waterfowl								
Species richness				1	0.491 (0.134)	13.530	**<0.001**	**<0.001**
G. chloropus	+	**0.001**	0.012					
F. atra	+	**0.022**	0.237					
Terrestrial birds								
Species richness				1	-0.044 (0.302)	0.021	0.884	0.884

| Variable | Analysis | | | | | | | |
| | Cooccur | | | GLMM | | | | |
	Effect	*P*	Adjusted *P* (Benjamini-Hochberg)	DF	Effect size (SE)	χ^2	*P*	Adjusted *P* (Benjamini-Hochberg)
P. colchicus	-	**0.050**	0.714					
Terrestrial mammals								
Species richness				1	-0.515 (0.188)	8.501	**0.004**	**0.006**
S. carolinensis	-	**0.020**	0.685					

Table 2. Summary of hypothesis testing using eDNA metabarcoding data and associated

environmental metadata, specifically biotic and abiotic factors known to influence *T.*

cristatus detection probability. Reported effects on *T. cristatus* detection in the literature

and hypothesised effects on eDNA-based *T. cristatus* detection are given for each variable.

Any variables not explicitly reported in the literature that were identified by the preliminary

cooccur analysis are listed as UNK. Direction of observed effects on eDNA-based *T. cristatus*

detection identified by each analysis (biotic GLMM, *n* = 504; abiotic GLMM, *n* = 504) are

given. Negative and positive effects are listed as - and + respectively. For categorical

variables with more than one level, effect size and standard error (SE) are only given for

levels reported in the model summary. Test statistic is for LRT used and significant P-values

(<0.05) are in bold. Variables included for model selection but not retained in the final

models are listed as NR.

Variable	Effect reported	Hypothesised effect	GLMM				
			DF	Effect size (SE)	χ^2	P	Adjusted P (Benjamini-Hochberg)
Amphibians							
L. vulgaris	+	+	1	1.616 (0.232)	50.024	**<0.001**	**<0.001**
B. bufo	UNK		1	-1.793 (0.643)	11.175	**<0.001**	**0.002**
Fish							
G. aculeatus	-	-	1	-1.048 (0.453)	6.300	**0.012**	**0.014**
C. carpio	-	-	1	-1.457 (0.586)	7.854	**0.005**	**0.008**
P. pungitius	-	-	NR				
C. carassius	-	-					
Waterfowl							
G. chloropus	UNK		1	0.893 (0.228)	15.703	**<0.001**	**<0.001**
F. atra	UNK		NR				
Waterfowl presence Minor Major	-	-	2	0.531 (0.234) 0.808 (0.495)	6.350	**0.042**	**0.042**
Pond area	-/+	-	1	-0.0003 (0.0002)	4.134	**0.042**	**0.042**

Variable	Effect reported	Hypothesised effect	GLMM				
			DF	Effect size (SE)	χ^2	*P*	Adjusted *P* (Benjamini-Hochberg)
Pond density	+	+			NR		
Pond depth	+	+			NR		
Pond substrate	+	+			NR		
Pond permanence	+	+			NR		
Water quality	+	+			NR		
Inflow Present	-	-	1	-0.820 (0.251)	11.437	**<0.001**	**0.001**
Outflow	UNK				NR		
Macrophyte cover	-/+	-			NR		
Shading	-/+	-	1	-0.010 (0.003)	12.266	**<0.001**	**0.001**
Woodland	+	+			NR		
Scrub/hedge	+	+			NR		
Ruderals	UNK				NR		

Figure 1. eDNA metabarcoding detection of *T. cristatus* in relation to other vertebrate species (*N* = 532 ponds): **(a)** other amphibians, **(b)** fish, **(c)** birds, and **(d)** mammals. Numbers on each bar are the number of ponds with (orange) and without (grey) *T. cristatus* in which a vertebrate species was detected.

Figure 2. *T. cristatus* detection (orange) and non-detection (grey) in relation to species richness of different vertebrate groups (*N* = 532 ponds): **(a)** other amphibians, **(b)** fish, **(c)** waterfowl, **(d)** terrestrial birds, and **(e)** mammals. Observed proportion of ponds with and without *T. cristatus* (left) is plotted alongside predicted probability of *T. cristatus* detection (right). Numbers on barplots of observed detection are the number of ponds for each category. In plots showing predicted *T. cristatus* detection, the observed data is shown as points (jittered around 0 and 1 to clarify variation in point density) and boxes are the model predictions.

Figure 3. Biotic and abiotic factors influencing *T. cristatus* detection (*n* = 504 ponds): **(a)** *L. vulgaris* detection, **(b)** *B. bufo* detection, **(c)** *C. carpio* detection, **(d)** *G. aculeatus* detection, **(e)** *G. chloropus* detection, **(f)** extent of waterfowl presence, **(g)** presence of inflow, **(h)** pond area, **(i)** percentage of shading, and **(j)** HSI score. The 95% CIs, as calculated using the predicted *T. cristatus* probability values and standard error for these predictions, are given for each relationship. The observed *T. cristatus* detection (orange) and non-detection (grey) data are displayed as points (jittered around 0 and 1 to clarify variation in point density) against the predicted relationships (boxes/lines).

943
944

945   **Figure 4.** Vertebrate species richness in ponds (*n* = 504) in relation to: **(a)** *T. cristatus*
946   detection and **(b)** the *T. cristatus* HSI score. The 95% CIs, as calculated using the predicted
947   species richness values and standard error for these predictions, are given for each
948   relationship. The observed data are displayed as points (jittered to clarify variation in point
949   density) against the predicted relationships (boxes/lines).