Understanding biodiversity at the pondscape using environmental DNA: a focus on great crested newts

Lynsey R. Harper[1]*, Lori Lawson Handley[1], Christoph Hahn[1,2], Neil Boonham[3,4], Helen C. Rees[5], Erin Lewis[3], Ian P. Adams[3], Peter Brotherton[6], Susanna Phillips[6] and Bernd Hänfling[1]

[1]School of Environmental Sciences, University of Hull, Hull, HU6 7RX, UK

[2]Institute of Zoology, University of Graz, Graz, Styria, Austria

[3]Fera, Sand Hutton, York, YO14 1LZ, UK

[4]Newcastle University, Newcastle upon Tyne, NE1 7RU, UK

[5]ADAS, School of Veterinary Medicine and Science, The University of Nottingham, Sutton Bonington Campus, Leicestershire, LE12 5RD, UK

[6] Natural England, Peterborough, PE1 1NG, UK

*Corresponding author:

Email: L.Harper@2015.hull.ac.uk

Word count: 9,563 words

eDNA metabarcoding represents a new tool for community biodiversity assessment in a broad range of aquatic and terrestrial habitats. However, much of the existing literature focuses on methodological development rather than testing of ecological hypotheses. Here, we use presence-absence data generated by eDNA metabarcoding of over 500 UK ponds to examine: 1) species associations between the great crested newt (*Triturus cristatus*) and other vertebrates, 2) determinants of great crested newt occurrence at the pondscape, and 3) determinants of vertebrate species richness at the pondscape. The great crested newt was significantly associated with nine vertebrate species. Occurrence in ponds was broadly reduced by more fish species, but enhanced by more waterfowl and other amphibian species. Abiotic determinants (including pond area, depth, and terrestrial habitat) were identified, which both corroborate and contradict existing literature on great crested newt ecology. Some of these abiotic factors (pond outflow) also determined species richness at the pondscape, but other factors were unique to great crested newt (pond area, depth, and ruderal habitat) or the wider biological community (pond density, macrophyte cover, terrestrial overhang, rough grass habitat, and overall terrestrial habitat quality) respectively. The great crested newt Habitat Suitability Index positively correlated with both eDNA-based great crested newt occupancy and vertebrate species richness. Our study is one of the first to use eDNA metabarcoding to test abiotic and biotic determinants of pond biodiversity. eDNA metabarcoding provided new insights at scales that were previously unattainable using established methods. This tool holds enormous potential for testing ecological hypotheses alongside biodiversity monitoring and pondscape management.

Freshwater ecosystems comprise <1% of the Earth's surface but provide vital

ecosystem services and are hotspots of biodiversity[1–3]. Nonetheless, freshwater

organisms are experiencing a greater rate of decline than marine or terrestrial

organisms[2,3]. Ponds especially represent critical habitat for biodiversity in a fragmented

landscape[4] and support many rare and protected species[5], such as the great crested newt

(*Triturus cristatus*) which is protected by UK and European legislation at all life

stages[5,6]. Ponds contribute substantially to regional- and landscape-scale aquatic

biodiversity[5,7–9] as well as non-aquatic biodiversity within pondscapes, i.e. a pond, its

immediate catchment, and the terrestrial matrix of land between ponds[5]. Until recently,

pondscapes were poorly understood[5] and neglected in research, scientific monitoring,

and policy[4,7,8]. Effective management of pondscapes requires knowledge of abiotic and

biotic factors that influence biodiversity, community structure and productivity.

Moreover, the biodiversity that ponds support individually and in combination must be

examined, but can only be maintained if stressors and threats to these systems are

understood[4,5,7,8,10]. Exhaustive sampling of pond biodiversity is impeded by the

complexity of these species-rich habitats, and numerous tools required for different taxa

with associated bias[11] and cost[12]. However, large-scale community-level monitoring,

encompassing alpha (site), beta (between-site) and gamma (landscape) diversity

analyses, is necessary to understand biodiversity in changing environments[13].

Analysis of environmental DNA (eDNA, i.e. DNA released by organisms via

skin cells, saliva, gametes, urine and faeces into the environment) is providing

ecologists with exceptional power to detect single species or describe whole

communities[14–18]. The great crested newt was the first and to date only UK protected

species to be routinely monitored using eDNA analysed with targeted real-time

quantitative PCR (qPCR)[19]. However, entire communities can be monitored using

High-Throughput Sequencing, i.e. eDNA metabarcoding[16–18]. This approach has been

used to estimate species richness and assess diversity along environmental

gradients[11,20–22], but studies have typically focused on species detection and

methodological improvement. eDNA metabarcoding has unprecedented diagnostic

power to test classic ecological hypotheses relating to the distribution of biodiversity

and its response to environmental pressures. Ponds are ideal model systems for

experimental validation and examination of biogeographical patterns as small,

abundant ecosystems that span broad ecological gradients[8]; however, few eDNA

metabarcoding studies to date have considered ponds[11,12,23–27].

Using ponds, we explore the potential of eDNA metabarcoding for hypothesis

testing. We focus on the threatened great crested newt as its ecology is well-understood.

Previous work established that both biotic (e.g. food availability, breeding substrate,

and predators) and abiotic (e.g. pond depth, area, permanence, and temperature)

variables strongly influence great crested newt breeding success[28]. These are

encompassed in the Habitat Suitability Index (HSI) used in species surveys[29,30]. The

HSI is comprised of 10 suitability indices (factors known to influence great crested

newts) which are scored and combined to calculate a decimal score between 0 and 1

representing habitat suitability (where 1 = excellent habitat); although some research

suggests HSI may not relate to great crested newt occupancy[31,32]. Fish species may

negatively impact great crested newt populations[28,33–40] or effects may be negligible[41].

Larvae tend to swim in open water, increasing susceptibility to fish and waterfowl

predation[34,36,38], and adults reportedly avoid ponds containing three-spined stickleback

(*Gasterosteus aculeatus*)[42], ninespine stickleback (*Pungitius pungitius*)[40], crucian carp

(*Carassius carassius*)[39,40], and common carp (*Carassius carpio*)[40]. Conversely, great

crested newts and smooth newts (*Lissotriton vulgaris*) are positively associated due to

shared habitat preferences[34,37,38,40]. Great crested newts are more likely in ponds with

better water quality (indicated by diverse macroinvertebrate communities)[29,40], higher nutrient content, and warmer temperature[43]. Water clarity is important for breeding displays, foraging success, and egg survival[34,38]. Higher density of ponds in a pondscape creates more opportunity for great crested newt occupation[33,36,37,40], but presence is negatively correlated with pond surface area[33]. Heavily shaded ponds[44], or those with high macrophyte cover[34,36,38], are less likely to support viable great crested newt populations. Great crested newts are also dependent on terrestrial habitat, preferring open, semi-rural pondscapes[37] containing pasture, extensively grazed and rough grassland, scrub, and coniferous and deciduous woodland[29,38,40,44,45].

The extensive literature on established determinants of the great crested newt provides an excellent opportunity to ground truth ecological patterns revealed by eDNA metabarcoding. We explore this tool's potential for biodiversity assessment at the pondscape using a dataset generated by eDNA metabarcoding of more than 500 ponds with comprehensive environmental metadata. We examined whether eDNA metabarcoding can test ecological hypotheses typically explored by established methods, and whether eDNA and established methods produce congruent results. Specifically, we sought to identify biotic determinants of great crested newt occurrence and species connections to the wider biological community. Using environmental metadata on pond properties and surrounding terrestrial habitat, we aimed to reaffirm abiotic determinants of great crested newts identified using established methods and revisit these important hypotheses at an unprecedented scale. We utilised eDNA metabarcoding for holistic biodiversity monitoring at the pondscape and uncovered abiotic determinants of vertebrate species richness - an impractical task by conventional means. Finally, we evaluated applicability of the great crested newt HSI[29,30] to eDNA-based great crested newt occupancy and vertebrate species richness of ponds.

Methods

Samples. 508 ponds, sampled as part of great crested newt surveys through Natural

England's Great Crested Newt Evidence Enhancement Programme, were processed

using eDNA metabarcoding alongside 24 ponds privately surveyed by ecological

consultants. All water samples were collected using methodology outlined by Biggs *et*

al. (2015)[19], detailed in Supplementary Methods. In brief, 20 x 30 mL water samples

were collected from each pond and pooled. Six 15 mL subsamples were taken from the

pooled sample and each added to 33.5 mL absolute ethanol and 1.5 mL sodium acetate

3 M (pH 5.2) for ethanol precipitation. Water subsamples from the same pond were

pooled during DNA extraction to produce one eDNA sample per pond. Targeted qPCR

detected great crested newt in 265 (49.81%) ponds. Egg searches performed at 506/508

ponds sampled for Natural England revealed great crested newt in 58 (11.46%) ponds[24].

Environmental metadata on pond characteristics and surrounding terrestrial

habitat was collected for 504/508 ponds sampled for Natural England (Supplementary

Fig. 1). Pond metadata included: maximum depth; circumference; width; length; area;

density; terrestrial overhang; shading; macrophyte cover; HSI score[29]; HSI band

(categorical classification of HSI score)[30]; permanence; water quality; pond substrate;

presence of inflow or outflow; presence of pollution; presence of other amphibians, fish

and waterfowl; woodland; rough grass; scrub/hedge; ruderals; other good terrestrial

habitat, i.e. good terrestrial habitat that did not conform to aforementioned habitat

types; and overall terrestrial habitat quality (see Supplementary Table 1 for details of

environmental variables).

DNA reference database construction. A custom, phylogenetically curated reference

database of mitochondrial 12S rRNA sequences for UK fish species was previously

created at University of Hull for an eDNA metabarcoding study of lake fish

communities[46]. Similar reference databases for UK amphibians, reptiles, birds, and

mammals were constructed using the ReproPhylo environment[47] in a Jupyter notebook

(Jupyter Team 2016). Full details of reference database construction are provided in

Harper *et al.* (in press)[24] and Supplementary Methods. Proportion of reference

sequences available for species varied within each vertebrate group: amphibians

100.00% (N = 21), reptiles 90.00% (N = 20), mammals 83.93% (N = 112), and birds

55.88% (N = 621). Species without any representation in these databases (i.e. no

records for that species or sister species within the same genus) are listed in

Supplementary Table 2. The amphibian database was supplemented by Sanger

sequences obtained from tissue of great crested newt, smooth newt, Alpine newt

(*Mesotriton alpestris*) and common toad (*Bufo bufo*) supplied by DICE, University of

Kent, under licence from Natural England, and common frog (*Rana temporaria*)

supplied by University of Glasgow (see Supplementary Methods). Databases for each

vertebrate group were combined and used for *in silico* validation of primers. The

complete reference databases compiled in GenBank format have been deposited in a

dedicated   GitHub   repository   for   this   study,   permanently   archived   at:

https://doi.org/10.5281/zenodo.1193609.

Primer validation. Published 12S ribosomal RNA (rRNA) primers 12S-V5-F (5'-

ACTGGGATTAGATACCCC-3') and 12S-V5-R (5'-TAGAACAGGCTCCTCTAG-

3')[48] were validated *in silico* using ecoPCR software[49] against our custom reference

database for UK vertebrates. Parameters set allowed a 50-250 bp fragment and

maximum of three mismatches between the primer pair and each sequence in the

reference database. Primers were previously validated *in vitro* for UK fish communities

by Hänfling *et al.* (2016)[46] and in the present study for six UK amphibian species

(Supplementary Fig. 2).

eDNA metabarcoding. Full details of the eDNA metabarcoding workflow performed

are provided in Harper *et al.* (in press)[24] and Supplementary Methods. eDNA was

amplified with a two-step PCR protocol, using the aforementioned 12S rRNA primers

in the first PCR. DNA from a cichlid (*Rhamphochromis esox*) was used for PCR

positive controls (six per PCR plate; $N = 114$), whilst sterile molecular grade water

(Fisher Scientific) substituted template DNA for negative controls (six per PCR plate;

$N = 114$). All PCR products were individually purified using E.Z.N.A. Cycle Pure V-

Spin Clean-Up Kits (VWR International) following manufacturer's protocol. A second

PCR was then used to bind Multiplex Identification (MID) tags to the amplified

product. PCR products were individually purified using a magnetic bead clean-up prior

to quantification with a Quant-IT™ PicoGreen™ dsDNA Assay. Using concentration

values, samples were normalised and pooled to create 4 nM pooled libraries, which

were quantified using a Qubit™ dsDNA HS Assay. Sequencing was performed on an

Illumina MiSeq using 2 x 300 bp V3 chemistry. Raw sequence reads were

taxonomically assigned against our UK vertebrate reference database using a custom

pipeline for reproducible analysis of metabarcoding data: metaBEAT (metaBarcoding

and Environmental Analysis Tool) v0.8 (https://github.com/HullUni-

bioinformatics/metaBEAT). After quality trimming, merging, chimera detection, and

clustering, non-redundant sets of query sequences were compared against our custom

reference database using BLAST[50]. Putative taxonomic identity was assigned using a

lowest common ancestor (LCA) approach based on the top 10% BLAST matches for

any query matching with at least 98% identity to a reference sequence across more than

80% of its length. Sequences that could not be assigned were subjected to a separate

BLAST search against the complete NCBI nucleotide (nt) database at 98% identity to

determine the source via LCA as described above. To ensure reproducibility, the

bioinformatic analysis has been deposited in the GitHub repository.

Data analysis. All downstream analyses were performed in the statistical programming

environment R v.3.4.2. (R Core Team 2017). Data and R scripts have been deposited

in the GitHub repository.

Non-target sequence assignments and original assignments at 98% identity were

merged. Any spurious assignments (i.e. non-UK species, invertebrates and bacteria)

were removed from the dataset. Assignments to genera or families which contained

only a single UK representative were manually assigned to that species. In our dataset,

only genus *Strix* was reassigned to tawny owl *Strix aluco*. Where family and genera

assignments containing a single UK representative had reads assigned to species, reads

from all assignment levels were merged and manually assigned to that species.

Consequently, all taxonomic assignments included in the final database were of species

resolution. Misassignments in our dataset were then corrected; again, only one instance

was identified. Scottish wildcat *Felis silvestris* was reassigned to domestic cat *Felis*

catus on the basis that Scottish wildcat does not occur where ponds were sampled

(Kent, Lincolnshire and Cheshire).

To reduce the potential for false positives, we applied species-specific

thresholds: a species was only classed as present at a given site if its sequence frequency

exceeded a species-specific threshold. Thresholds for each species were defined by

analysing sequence data from PCR positive controls ($N = 114$) and identifying the

maximum sequence frequency for a given species across all PCR positive controls

(Supplementary Table 3). For example, the great crested newt species-specific false

positive sequence threshold was 0.028% to omit all false detections in the PCR positive

controls. After thresholds were applied, the read count data for detected species were

converted to a presence-absence matrix for downstream analyses. In the main text, we

focus on the results inferred using the species-specific thresholds but all downstream

analyses were also performed across a variety of blanket sequence thresholds (0.05 -

30%, see Supplementary Tables 4-9). We tested the influence of fish and waterfowl

presence, pond characteristics and surrounding terrestrial habitat on great crested newt

occurrence as inferred by eDNA metabarcoding. We were particularly interested in the

appropriateness of HSI for eDNA-based great crested newt occupancy. Hypotheses are

summarised in Table 1.

  All Generalised Linear Mixed Models (GLMMs) were executed using the R

package 'lme4' v1.1-12[51]. First, correlations between great crested newt occurrence and

number of other vertebrate species were investigated using a binomial GLMM ($N =$

532). Individual species associations were then investigated using the method of Veech

(2013)[52] in the R package 'cooccur' v1.3[53] ($N = 532$). Identified associations informed

candidate biotic variables to be included with abiotic variables (Table S1) in a binomial

GLMM of great crested newt occurrence ($N = 504$). Collinearity and spatial

autocorrelation within the dataset were investigated before the most appropriate

regression model was determined. Collinearity between explanatory variables was

assessed using a Spearman's rank pairwise correlation matrix. After collinear variables

were removed, variance inflation factors (VIFs) of remaining variables were calculated

using the R package 'car' v2.1-6[54] to identify remnant multicollinearity. Variables

corresponding to HSI (HSI score, HSI band) were multicollinear and subsequently

removed prior to model selection (see Supplementary Methods), and HSI score

analysed separately in a binomial GLMM.

A large number of explanatory variables remained: max. depth; area; density,

overhang; macrophyte cover; permanence; water quality; pond substrate; inflow;

outflow; pollution; presence of amphibians, waterfowl and fish; woodland; rough grass;

scrub/hedge; ruderals; terrestrial other; and overall terrestrial habitat quality. The

relative importance of these for determining great crested newt occurrence was inferred

using a classification tree within the R package 'rpart' v4.1-13[55]. A pruning diagram

was applied to the data to cross-validate the classification tree and remove unimportant

explanatory variables (see Supplementary Methods). Many variables occurred more

than once in the classification tree, indicative of weak non-linear relationships with the

response variable. Generalised Additive Models (GAMs) were performed to deal with

non-linearity but several explanatory variables were in fact linear (estimated one degree

of freedom for smoother). A parametric, binomial Generalised Linear Model (GLM)

was applied and the potential for spatial autocorrelation assessed using spline

correlograms of the data using R package 'ncf' v1.1-7[56]. A binomial GLMM was

employed to account for dependencies within sites, handled with the introduction of

random effects[57–59]. Each eDNA sample represented a different pond and thus sample

was treated as a random effect. The mixed model successfully accounted for spatial

autocorrelation within sites when a spline correlogram of the Pearson residuals was

examined.

After identification of a suitable set of explanatory variables and modelling

framework, the variables which are the most important determinants of great crested

newt occurrence and a suitable, parsimonious approximating model to make predictions

were determined. An information-theoretic approach using Akaike's Information

Criteria (AIC) to evaluate model fit was employed[60]. A binomial distribution was

specified as the response variable was presence-absence data. After following a similar

workflow for identification of a suitable set of explanatory variables and modelling

framework (see Supplementary Methods), a set of variables that best explain vertebrate

species richness were constructed. A Poisson distribution was specified for all species

richness models as the response variable was integer count data. Model fit was

evaluated as above using AIC.

All binomial and Poisson models considered were nested and so the best models

of great crested newt occurrence and vertebrate species richness respectively were

chosen using stepwise backward deletion of terms based on Likelihood Ratio Tests

(LRTs). The final binomial and Poisson models were tested for overdispersion using

the R package 'RVAideMemoire' v 0.9-69[61] and custom functions to test

overdispersion of the Pearson residuals. Model fit was assessed using the Hosmer and

Lemeshow Goodness of Fit Test[62] within the R package 'ResourceSelection' v0.3-2[63],

quantile-quantile plots and partial residual plots[59,64]. Model predictions were obtained

using the predictSE() function in the 'AICcmodavg' package v2.1-1[65] and upper and

lower 95% CIs were calculated from the standard error of the predictions. All values

were bound in a new data frame and model results plotted for evaluation using the R

package 'ggplot2' v 2.2.1[66].

Data availability. Raw sequence reads have been archived on the NCBI Sequence

Read Archive (Bioproject: PRJNA417951; SRA accessions: SRR6285413 -

SRR6285678). Jupyter notebooks, R scripts and corresponding data are deposited in a

dedicated GitHub repository (https://github.com/HullUni-

bioinformatics/Harper_et_al_2018) which has been permanently archived

Results and discussion

Across two sequencing runs, 532 eDNA samples and 228 PCR controls were processed.

The runs generated raw sequence read counts of 36,236,862 and 32,900,914

respectively. After trimming and merging of paired-end reads, 26,294,906 and

26,451,564 sequences remained. Following removal of chimeras and redundancy via

clustering, the libraries contained 14,141,237 and 14,081,939 sequences (average read

counts of 36,826 and 36,671 per sample respectively), of which 13,126,148 and

13,113,143 sequences were taxonomically assigned. In the final dataset (thresholds

applied and assignments corrected), a total of 60 vertebrate species were detected by

eDNA metabarcoding across the 532 ponds surveyed (Supplementary Table 10). These

consisted of six amphibian species, 14 fish species, 18 bird species, and 22 mammal

species (Supplementary Fig. 3). Amphibian species detection ranged from 1 - 152

ponds (median 81 ponds) whilst fish species detection ranged from 1 - 72 ponds

(median 15 ponds). Bird species detection ranged between 1 and 215 ponds (median 3

ponds) whereas mammal species detection ranged between 1 and 179 ponds (median 9

ponds). The most common species detected across all vertebrate groups were common

moorhen (*Gallinula chloropus*, $N = 215$), cow (*Bos taurus*, $N = 179$), smooth newt (*N*

$= 152$), great crested newt ($N = 149$), pig (*Sus scrofa*, $N = 140$), and common frog ($N =$

120). All detected species and their frequency of detection are listed in Supplementary

Table 10.

We discuss great crested newt occupancy in the context of broad trends found

across vertebrate groups (GLMM: overdispersion $\chi^2_{525} = 517.636$, $P = 0.582$; fit $\chi^2_8 =$

22.524, $P = 0.004$, $R^2 = 9.43\%$) and individual species associations (Table 1).

Significant species associations with the great crested newt revealed by the co-

occurrence analysis were carried forward as candidate variables for analysis of biotic

and abiotic determinants (GLMM: overdispersion $\chi^2_{490} = 413.394$, $P = 0.995$; fit $\chi^2_8 =$

11.794, $P = 0.161$, $R^2 = 38.58\%$). Associations with support from both analyses are

summarised in Table 1. Great crested newt occupancy was best explained by smooth

newt occurrence (+), common toad occurrence (-), three-spined stickleback occurrence

(-), grey squirrel occurrence (-), vertebrate species richness (+), pond outflow (-),

ruderal habitat (-), other good terrestrial habitat (-), pond area (-), and max. depth (+).

Biotic determinants of great crested newt occurrence. We found a positive

correlation between great crested newt occurrence and increasing number of other

amphibian species (Table 1, Fig. 1a). Smooth newts were commonly detected in ponds

with great crested newts but palmate newts (*Lissotriton helveticus*) were not. Similarly,

common toad and common frog records were less frequent in ponds containing great

crested newts, and there was only one record of marsh frogs (*Pelophylax ridibundus*)

in a great crested newt pond (Supplementary Fig. 3a). Of these observations, a positive

association between the great crested newt and smooth newt (Table 1, Figs. 2, 3a) and

a negative association between the great crested newt and common toad were

significant (Table 1, Figs. 2, 3b). Great crested newts and smooth newts share similar

terrestrial and aquatic habitat requirements resulting in selection of the same ponds for

breeding[34,37,40], with more than 60% overlap in ponds reported[38]. Notably, research

suggests smooth newts are more versatile and capable of inhabiting a broader range of

habitat[34,38] whereas great crested newts may be associated with larger, deeper ponds

with an abundance of macrophytes and absence of fish located in open, semi-rural

landscapes[37]. Conversely, the negative association observed between the great crested

newt and common toad may be attributable to toads inhabiting fish-containing ponds[67]

or great crested newt predation on toad eggs and larvae[28].

Great crested newt occurrence was reduced in ponds containing a greater

number of fish species (Table 1), and newts were absent from ponds containing more

than four fish species (Fig. 1b). Nonetheless, all detected fish species were recorded in

great crested newt ponds to some extent, except Eurasian ruffe (*Gymnocephalus*

cernua) (Fig. S3b). It is important to note that some fish species detections may result

from eDNA transport into ponds via inflows from larger stream or river catchments,

when these species do not actually inhabit ponds. We discuss only associations with

fish species that are established inhabitants of ponds. The great crested newt had

significant negative associations with ninespine stickleback (Table 1, Fig. 2) and three-

spined stickleback (Table 1, Figs. 2, 3c), the latter of which was shared by the smooth

newt (Fig. 2). A non-significant negative co-occurrence was also observed between

great crested newts and common carp (Table 1). Common carp are ecosystem

engineers: their benthic foraging activity increases water turbidity and reduces

invertebrate density and macrophyte cover, affecting species that depend on these

groups[68,69]. Introduced fish species exerted a negative effect on site occupation of both

newt species in Belgium[37] and both species only colonised a site in England once three-

spined stickleback were removed[42]. Smooth newts are known to avoid fish occupied

sites, including ponds and wetlands[70,71], and negative effects of fish species on great

crested newt populations have been frequently reported[28,33–36,38,39]. Conversely, other

research suggests no or minimal negative interaction between fish and great crested

newts[41,72]. Fish species characteristic of ponds, such as crucian carp, are unlikely to be

damaging predators to amphibian populations[72,73]. Indeed, great crested newt detection

was equal in ponds containing or absent of crucian carp. However, consumption of

macroinvertebrates by fish can alter habitat suitability for great crested newts[35] as many

fish species share the same trophic status as newt species[70]. Fish also tend to be

associated with algal ponds where macrophyte diversity is impaired. Reduced

macrophyte availability imposes restrictions on egg-laying in great crested newts and

restricts the ecological niches that invertebrate prey may inhabit[38].

Unexpectedly, great crested newt occurrence was positively associated with

increasing number of waterfowl species (Table 1, Fig. 1c), despite absence of great

crested newts in ponds with certain waterfowl species (Supplementary Fig. 3c).

Furthermore, the great crested newt had significant positive associations with the

common coot (*Fulica atra*) and common moorhen (Table 1, Fig. 2), and a non-

significant negative association with the green-winged teal (Table 1). Great crested

newts are typically found in ponds with high macrophyte diversity as macrophyte

species dictate reproductive success and invertebrate prey availability[72,74]. Common

moorhen and common coot share macrophytes and macroinvertebrates as resources but

feed on both directly[75–77] thus competition between great crested newts and omnivorous

waterfowl may be reduced or indirect. Great crested newt breeding in April to June[28]

may be impacted by coots pulling up submerged vegetation, damaging vegetation

banks[78]. However, coot diet tends to be macrophyte-dominated in late summer and

autumn[77]. Both coot and moorhen also crop emergent macrophytes in their search for

invertebrate prey[75,76], but in doing so they may expose prey items to great crested newts

and confer indirect benefits.

The most common terrestrial detections in this study were domesticated or

introduced pest species, such as grey squirrel (*Sciurus carolinensis*) and muntjac deer

(*Muntiacus reevesi*)[69,79] (Supplementary Table 10). Nonetheless, we identified wild

species which emphasise the importance of ponds as stepping stones for both semi-

aquatic and terrestrial taxa[7,8], through provision of drinking, foraging, dispersal, and

reproductive opportunities[10,25,80]. The most frequent terrestrial bird detections included

buzzard (*Buteo buteo*), Eurasian jay (*Garrulus glandarius*), dunnock (*Prunella*

modularis), and starling (*Sturnus vulgaris*) (Supplementary Table 10), which utilise

different habitats. We detected several mammal species with Biodiversity Actions

Plans and/or of conservation concern, including otter (*Lutra lutra*), water vole (*Arvicola*

amphibius), European polecat (*Mustela putorius*), brown hare (*Lepus europaeus*) and

water shrew (*Neomys fodiens*)[79]. Notably, some mammals were only identified in one

pond (Supplementary Table 10) and American mink (*Neovison vison*) was absent

despite widespread UK distribution[79].

Records of great crested newts in relation to terrestrial species, and any

significant associations identified below, are unlikely to reflect direct species

interactions. Rather, these records and associations are a probable outcome of land-use

and indirect interaction. No significant relationships between the numbers of terrestrial

bird or mammal species and great crested newt occurrence were found (Table 1, Figs.

1d, e), but great crested newts were entirely absent from ponds where certain terrestrial

species were present e.g. great spotted woodpecker *(Dendrocopos major)*, tawny owl,

badger (*Meles meles*), and red deer (*Cervus elaphus*) (Supplementary Figs. 3c, d).The

great crested newt had a significant positive association with pig (Table 1, Fig. 2), but

significant negative associations with the grey squirrel (*Sciurus carolinensis*) (Table 1,

Figs. 2, 3d) and common pheasant (*Phasianus colchicus*) (Table 1, Fig. 2). A non-

significant negative co-occurrence between the great crested newt and badger was also

identified (Table 1). Excluding breeding, adult great crested newts live outside ponds

in terrestrial habitat for foraging, shelter, and hibernation[35,43]. Juveniles also spend two

to three years on land after emerging from ponds[70]. During time spent outside of ponds,

great crested newts may suffer dessication, or predation[28,45,70] from terrestrial species.

There have been anecdotal records of pheasant predation on herpetofauna, including

great crested newt[81], but existing literature on amphibian and terrestrial species

interactions is sparse with even less study of great crested newt interactions.

Critically, data points in all analyses of biotic determinants were not evenly

distributed across different species, or number of species within each vertebrate group.

Ponds containing a higher number of vertebrate species were much fewer than ponds

containing a lower number of vertebrate species (Fig. 1). Similarly, some species were

detected more frequently in ponds than others (Supplementary Table 10 and Fig. 3).

This uneven distribution is likely a natural outcome of species accumulation, but may

reduce capability of models to make accurate predictions.

Abiotic determinants of great crested newt occurrence. The probability of great

crested newt occurrence increased with greater pond depth but decreased in ponds with

larger area, outflow, without ruderal habitat, and with some other good terrestrial

habitat (Table 1, Figs. 3e-g, i-j). Previous work has shown great crested newts utilise

small and large ponds[34,38], although very small ponds (less than 124 m^2) were incapable

of supporting all life stages and larger ponds had greater occurrence of fish[38]. Large

ponds may also be more susceptible to eutrophication due to agricultural or polluted

run-off[38]. Yet, some studies found no effect of pond area on great crested newt

occurrence[35,41,45], and pond area has been deemed a poor predictor of great crested newt

reproductive success[44]. In contrast, past research showed a positive influence of pond

depth on great crested newt occupancy[35]. Conditions in shallow ponds may be too

unpredictable for great crested newt occupation, as they are susceptible to drying out

or freezing and may contain less prey. However, pond depth and surrounding terrestrial

habitat may be linked as detrimental effects of shallow water are more typically

observed in open farmland[33,35,71]. Temporary water bodies can be occupied provided

fish are absent[71] and drying may reduce predators such as fish and dragonfly larvae[28].

Unlike pond area and depth, there is little research on outflow to support an effect on

great crested newt occurrence. Pond inflow is known to affect biodiversity due to

polluted agricultural run-off and connections to streams and rivers containing large,

predatory fish. We suggest outflow (facilitated by drains, pipes or streams) may

stabilise maximum water level[82] and minimise fluctuations in pond depth, affecting

subsequent colonisation and structure of biological communities.

Our results support previous work demonstrating that good terrestrial habitat is

key to great crested newt success and serves multiple purposes, including daytime and

long-term shelter from extreme conditions in refugia, as well as foraging and dispersal

opportunities[28]. Previous research determined great crested newt occupancy and

breeding success was sub-optimal in coniferous forest yet enhanced in deciduous or

herb-rich forest and pasture[44,45]. Similarly, extensively grazed grassland and deadwood

positively influenced great crested newt presence whilst intensively grazed grasslands

were unoccupied[38]. Lower great crested newt abundance has been observed in

cultivated habitats[33], and modern forestry and increasing land use were deemed the

biggest great crested newt decline factors using a spatially explicit population model[44].

Conversely, others have found minimal effect of landscape context (excluding urban

areas) on great crested newts[36], suggesting terrestrial habitat may not restrict species

distribution; although, habitat degradation may increase isolation of ponds. Our results

indicate terrestrial habitat does influence great crested newt occupancy but without

quantitative data, these discrete effects cannot be teased apart. Data on type, density,

and great crested newt utilisation of terrestrial habitat, as well as distance of ponds to

terrestrial habitat, are necessary to fully understand great crested newt occupancy and

interactions with terrestrial species. However, this is a phenomenal task for large

numbers of ponds across a vast landscape[35].

HSI in relation to eDNA-based great crested newt occupancy. In a separate analysis

(GLMM: overdispersion $\chi^2_{501} = 506.763$, $P = 0.4198$; fit $\chi^2_8 = 8.118$, $P = 0.422$, $R^2 =$

4.99%), eDNA-based great crested newt occurrence positively correlated with HSI

score (Table 1), where probability of great crested newt occupancy was greater in ponds

with higher HSI score (Fig. 4). It has been suggested that HSI may be inappropriate for

predicting great crested newt occupancy or survival probabilities[32], but our finding

indicates HSI can be used to predict great crested newt occupancy at the pondscape.

HSI may help establish protection of ponds and the biodiversity they host by identifying

those which may be occupied by great crested newt. Optimal habitat can also be

identified for creation of new ponds or restoration of old ponds to encourage new

populations of this threatened amphibian species.

Nevertheless, issues remain with the HSI. Great crested newt occurrence may

indicate good quality habitat but may not reflect successful breeding and population

viability, albeit one study found ponds with higher HSI did have higher reproduction

probability[32]. Other issues include the use of qualitative data for score calculation, and

subjective, scorer-dependent estimation of indices[29]. For future application of HSI in

great crested newt eDNA survey, we recommend metabarcoding for quantification of

some indices which are qualitatively assessed (e.g. water quality via macroinvertebrate

diversity, fish and waterfowl presence) alongside detection of great crested newts.

Provided rigorous spatial and temporal sampling are undertaken, eDNA metabarcoding

can also generate site occupancy data to estimate relative abundance of species[12,46].

However, only conventional surveys can provide data on true great crested newt abundance to enable effective mitigation (e.g. translocation), understand population dynamics, and generate survival and reproduction probabilities.

Abiotic and biotic determinants of vertebrate species richness. In another analysis (GLMM: overdispersion $\chi^2_{494} = 431.959$, $P = 0.979$; fit $\chi^2_8 = -42.708$, $P = 1.000$, $R^2 = 8.94\%$), species richness was greater in ponds with outflow (0.214 ± 0.063, $\chi^2_1 = 11.220$, $P = 0.0008$, Fig. 5a), but reduced in those with some rough grass habitat (-0.297 ± 0.074) compared to ponds with no (-0.1402 ± 0.0795) or important rough grass habitat ($\chi^2_2 = 16.715$, $P = 0.0002$, Fig. 5b). Overall quality of terrestrial habitat was also influential ($\chi^2_2 = 8.244$, $P = 0.016$, Fig. 5c) where species richness was higher in ponds that were in areas considered to be poor (0.115 ± 0.089) or moderate (0.216 ± 0.078) habitat for great crested newts. Species richness was reduced as percentage of terrestrial overhang (-0.0026 ± 0.0008, $\chi^2_1 = 9.575$, $P = 0.002$, Fig. 5d) and percentage of macrophyte cover increased (-0.002 ± 0.001, $\chi^2_1 = 4.117$, $P = 0.043$, Fig. 5e) but improved with pond density (0.006 ± 0.003, $\chi^2_1 = 4.564$, $P = 0.033$, Fig. 5f). Many studies have focused on species richness of aquatic invertebrates as a range of invertebrate groups can be surveyed simultaneously using conventional tools. Until application of eDNA metabarcoding, this was not possible for aquatic and non-aquatic vertebrates. Instead indicator groups, such as amphibians, were chosen as representatives of pond biodiversity - although amphibians may in fact be poor surrogates of macroinvertebrate and macrophyte diversity[83,84]. Consequently, the literature on vertebrate species richness in aquatic ecosystems is sparse and we may only compare our results to studies which have investigated species richness of different vertebrate assemblages or species guilds, primarily amphibians and waterfowl.

Plentiful rough grass habitat can create more ecological niches and foraging

opportunity for a variety of vertebrates, but quantitative data on type and abundance of

terrestrial habitat surrounding ponds would be required to understand which species

prefer open or covered habitat. Pond outflow and inflow have received little

investigation in studies of freshwater biodiversity. Outflow may release harmful

pollutants and pathogens[85] that would otherwise accumulate and be retained in a closed

pond system. Outflow may benefit vertebrate biodiversity at risk of human health in

urban areas[85], but pollution was not identified as a candidate for model selection. Shade

has been identified as a principal driver of macroinvertebrate and macrophyte diversity

in freshwater ponds, negatively correlating with macrophyte cover, and can create

anoxic conditions in water bodies thereby decreasing productivity[9]. This can have

knock-on effects for consumers at higher trophic levels. For example, amphibians have

been observed to avoid ponds that are densely vegetated[86]. Yet, canopy and macrophyte

cover were also identified as positive drivers of amphibian species richness[86,87] and

abundance[88]. Our own results indicate highly shaded ponds are inconducive to high

vertebrate species richness but high densities of ponds support higher species richness,

providing further evidence of the importance of ponds for aquatic and non-aquatic

taxa[7,8].

In a separate analysis (GLMM: overdispersion $\chi^2_{501} = 389.744$, $P = 0.999$; fit

$\chi^2_8 = -145.12$, $P = 1.000$, $R^2 = 1.10\%$), vertebrate species richness positively correlated

with HSI score (0.459 ± 0.002, $\chi^2_1 = 5.034$, $P = 0.025$, $R^2 = 1.10\%$), where species

richness was improved in ponds with higher HSI score (Fig. 5g). HSI score also

positively correlated with probability of great crested newt occurrence (Fig. 4), and a

positive association was identified between vertebrate species richness and great

crested newt occurrence ($P < 0.0001$, Fig. 3b). Our results suggest some indices which

comprise the great crested newt HSI also represent key habitat criteria for broader

biodiversity, for example, outflow and terrestrial habitat. However, several indices

which affect great crested newts were not identified as determinants of vertebrate

species richness. Nonetheless, it may be possible to adapt HSI to more accurately

represent and predict vertebrate species richness in order to identify areas for pond

creation and management to enhance aquatic and non-aquatic biodiversity.

Implications for biodiversity assessment at the pondscape. Many species

associations were identified using eDNA metabarcoding (Fig. 2). However, there is no

literature available to confirm the nature, or even existence, of the majority of these

relationships. Lack of appropriate survey methods has caused freshwater research to

focus on single species or guilds and assemblages when studying predictors of species

diversity, richness, and abundance, or investigating impact of environmental change

and gradients. New methods are required for holistic biodiversity assessment in

response to ecosystem drivers and stressors. We have demonstrated how eDNA

metabarcoding can be used for landscape-scale biodiversity monitoring and ecological

study. Our results provide new insights and unparalleled biological understanding of

aquatic and non-aquatic biodiversity at the UK pondscape. Continued use of eDNA

metabarcoding could enhance our understanding of freshwater networks to enable more

effective protection and management for both aquatic and non-aquatic biodiversity.

Huge quantities of data can be generated to reduce the noise typically observed in

ecological datasets and at comparable cost to single-species eDNA monitoring[24]. We

investigated associations between aquatic and non-aquatic vertebrates and combined

metabarcoding with environmental metadata to revisit important ecological hypotheses

at an unprecedented scale, identifying determinants of great crested newts and broader

biodiversity. Our findings indicate preferred habitat of a threatened amphibian and will

guide management in the face of increasing land-use and habitat fragmentation - a

poignant issue as protective legislation for the great crested newt in the UK is under

review. Whilst conservation of threatened biodiversity and their habitat should be a

priority, the bigger picture should not be ignored. eDNA metabarcoding can create both

fine and broad-scale species inventories and allow researchers to examine the response

of entire communities' to environmental change, thereby allowing prioritisation of

regional- and landscape-scale conservation effort. eDNA metabarcoding holds great

promise for improved biodiversity monitoring and we are only beginning to realise and

explore these opportunities.

References

1. Strayer, D. L. & Dudgeon, D. Freshwater biodiversity conservation: recent progress and future challenges. *J. North Am. Benthol. Soc.* **29,** 344–358 (2010).
2. Collen, B. *et al.* Global patterns of freshwater species diversity, threat and endemism. *Glob. Ecol. Biogeogr.* **23,** 40–51 (2014).
3. Dudgeon, D. *et al.* Freshwater biodiversity: importance, threats, status and conservation challenges. *Biol. Rev. Camb. Philos. Soc.* **81,** 163–182 (2006).
4. Céréghino, R., Biggs, J., Oertli, B. & Declerck, S. The ecology of European ponds: defining the characteristics of a neglected freshwater habitat. *Hydrobiologia* **597,** 1–6 (2008).
5. Wood, P. J., Greenwood, M. T. & Agnew, M. D. Pond biodiversity and habitat loss in the UK. *Area* **35,** 206–216 (2003).
6. Rees, H. C. *et al.* The application of eDNA for monitoring of the Great Crested Newt in the UK. *Ecol. Evol.* **4,** 4023–4032 (2014).
7. Oertli, B., Céréghino, R., Hull, A. & Miracle, R. Pond conservation: from science to practice. *Hydrobiologia* **634,** 1–9 (2009).
8. De Meester, L. *et al.* Ponds and pools as model systems in conservation biology, ecology and evolutionary biology. *Aquat. Conserv.* **15,** 715–725 (2005).
9. Sayer, C. *et al.* The role of pond management for biodiversity conservation in an agricultural landscape. *Aquat. Conserv.* **22,** 626–638 (2012).
10. Biggs, J., von Fumetti, S. & Kelly-Quinn, M. The importance of small waterbodies for biodiversity and ecosystem services: implications for policy makers. *Hydrobiologia* **793,** 3–39 (2016).

11. Evans, D. N. T. *et al.* Fish community assessment with eDNA metabarcoding:
effects of sampling design and bioinformatic filtering. *Can. J. Fish. Aquat. Sci.*
74, 1362-1374 (2017).

12. Valentini, A. *et al.* Next-generation monitoring of aquatic biodiversity using
environmental DNA metabarcoding. *Mol. Ecol.* **25,** 929–942 (2016).

13. Hajibabaei, M., Baird, D. J., Fahner, N. A., Beiko, R. & Golding, G. B. A new
way to contemplate Darwin's tangled bank: how DNA barcodes are reconnecting
biodiversity science and biomonitoring. *Philos. Trans. R. Soc. Lond. B Biol. Sci.*
371, (2016).

14. Rees, H. C., Maddison, B. C., Middleditch, D. J., Patmore, J. R. M. & and Gough,
K. C. REVIEW: The detection of aquatic animal species using environmental
DNA – a review of eDNA as a survey tool in ecology. *J. Appl. Ecol.* **52,** 827-831
(2014).

15. Thomsen, P. F. & Willerslev, E. Environmental DNA – an emerging tool in
conservation for monitoring past and present biodiversity. *Biol. Conserv.* **183,** 4-
18 (2015).

16. Lawson Handley, L. How will the 'molecular revolution' contribute to biological
recording? *Biol. J. Linnaean. Soc.* **115,** 750–766 (2015).

17. Deiner, K. *et al.* Environmental DNA metabarcoding: transforming how we
survey animal and plant communities. *Mol. Ecol.* **26,** 5872-5895 (2017).

18. Taberlet, P., Coissac, E., Pompanon, F., Brochmann, C. & Willerslev, E. Towards
next-generation biodiversity assessment using DNA metabarcoding. *Mol. Ecol.*
21, 2045–2050 (2012).

19. Biggs, J. *et al.* Using eDNA to develop a national citizen science-based monitoring
programme for the great crested newt (*Triturus cristatus*). *Biol. Conserv.* **183,** 19–
28 (2015).

20. Kelly, R. P. *et al.* Genetic signatures of ecological diversity along an urbanization
gradient. *PeerJ* **4,** e2444 (2016).

21. Olds, B. P. *et al.* Estimating species richness using environmental DNA. *Ecol.*
Evol. **6,** 4214–4226 (2016).

22. Andruszkiewicz, E. A. *et al.* Biomonitoring of marine vertebrates in Monterey Bay
using eDNA metabarcoding. *PLoS One* **12,** e0176343 (2017).

23. Thomsen, P. F. *et al.* Monitoring endangered freshwater biodiversity using
environmental DNA. *Mol. Ecol.* **21,** 2565–2573 (2012).

24. Harper, L. R. *et al.* Needle in a haystack? A comparison of eDNA metabarcoding
and targeted qPCR for detection of the great crested newt (*Triturus cristatus*).
*Ecol.   Evol.   (in   press).   Pre-print   at
https://www.biorxiv.org/content/early/2018/03/05/215897.

25. Klymus, K. E., Richter, C. A., Thompson, N. & Hinck, J. E. Metabarcoding of
Environmental DNA Samples to Explore the Use of Uranium Mine Containment
Ponds as a Water Source for Wildlife. *Diversity* **9,** 54 (2017).

26. Bálint, M. *et al.* Twenty-five species of frogs in a liter of water: eDNA survey for
exploring   tropical   frog   diversity.   Pre-print   at
https://www.biorxiv.org/content/early/2017/08/15/176065 (2017).

27. Ushio, M. *et al.* Environmental DNA enables detection of terrestrial mammals
from forest pond water. *Mol. Ecol. Resour.* **17,** e63-e65 (2017).
28. Langton, T., Beckett, C. & Foster, J. *Great crested newt conservation handbook*.
(Froglife, Halesworth, 2001).
29. Oldham, R. S., Keeble, J., Swan, M. J. S. & and Jeffcote, M. Evaluating the
suitability of habitat for the great crested newt (*Triturus cristatus*). *Herpetol. J.* **10,**
143-155 (2000).
30. ARG-UK. *ARG UK Advice Note 5: Great Crested Newt Habitat Suitability Index*.
(Amphibian and Reptile Groups of the United Kingdom, 2010).
31. Sewell, D., Beebee, T. J. C. & Griffiths, R. A. Optimising biodiversity assessments
by volunteers: The application of occupancy modelling to large-scale amphibian
surveys. *Biol. Conserv.* **143,** 2102–2110 (2010).
32. Unglaub, B., Steinfartz, S., Drechsler, A. & Schmidt, B. R. Linking habitat
suitability to demography in a pond-breeding amphibian. *Front. Zool.* **12,** 9
(2015).
33. Joly, P., Miaud, C., Lehmann, A. & Grolet, O. Habitat Matrix Effects on Pond
Occupancy in Newts. *Conserv. Biol.* **15,** 239–248 (2001).
34. Skei, J. K., Dolmen, D., Rønning, L. & Ringsby, T. H. Habitat use during the
aquatic phase of the newts *Triturus vulgaris* (L.) and *T. cristatus* (Laurenti) in
central Norway: proposition for a conservation and monitoring area. *Amphib-*
Reptil. **27,** 309–324 (2006).
35. Denoël, M. & Ficetola, G. F. Conservation of newt guilds in an agricultural
landscape of Belgium: the importance of aquatic and terrestrial habitats. *Aquat.*
Conserv. **18,** 714–728 (2008).
36. Hartel, T., Nemes, S. & Oellerer, K. Using connectivity metrics and niche
modelling to explore the occurrence of the northern crested newt *Triturus cristatus*
(Amphibia, Caudata) in a traditionally managed landscape. *Environ. Conserv.* **37,**
195-200 (2010).
37. Denoël, M., Perez, A., Cornet, Y. & Ficetola, G. F. Similar local and landscape
processes affect both a common and a rare newt species. *PLoS One* **8,** e62727
(2013).
38. Rannap, R. & Briggs, L. THE CHARACTERISTICS OF GREAT CRESTED
NEWT TRITURUS CRISTATUS'BREEDING PONDS. (Project Report:
"Protection of the Great Crested Newt in the Eastern Baltic region",
LIFE2004NAT/EE/000070, Action A1, 2006).
39. Rannap, R., Lõhmus, A. & Briggs, L. Restoring ponds for amphibians: a success
story. *Hydrobiologia* **634,** 87–95 (2009).
40. Rannap, R., Lõhmus, A. & Briggs, L. Niche position, but not niche breadth, differs
in two coexisting amphibians having contrasting trends in Europe. *Divers.*
Distributions **15,** 692–700 (2009).
41. Maletzky, A., Kyek, M. & Goldschmid, A. Monitoring status, habitat features and
amphibian species richness of crested newt (*Triturus cristatus* superspecies) ponds
at the edge of the species range (Salzburg, Austria). *Ann. Limnol. – Int. J. Lim.* **43,**
107-115 (2007).

42. McLee, A. G. & Scaife, R. W. The colonisation by great crested newts (*Triturus*
cristatus) of a water body following treatment with a piscicide to remove a large
population of sticklebacks (*Gasterosteus aculeatus*). *Herpetol. Bull.* **42,** 6-9
(1992).
43. Gustafson, D. H., Andersen, A. S. L., Mikusiński, G. & Malmgren, J. C. Pond
Quality Determinants of Occurrence Patterns of Great Crested Newts (*Triturus*
cristatus). *J. Herpetol.* **43,** 300–310 (2009).
44. Vuorio, V., Heikkinen, R. K. & Tikkanen, O.-P. Breeding Success of the
Threatened Great Crested Newt in Boreal Forest Ponds. *Ann. Zool. Fennici* **50,**
158–169 (2013).
45. Gustafson, D. H., Malmgren, J. C. & Mikusiński, G. Terrestrial Habitat Predicts
use of Aquatic Habitat for Breeding Purposes — A Study on the Great Crested
Newt (*Triturus cristatus*). *Ann. Zool. Fennici* **48,** 295–307 (2011).
46. Hänfling, B. *et al.* Environmental DNA metabarcoding of lake fish communities
reflects long-term data from established survey methods. *Mol. Ecol.* **25**, 3101-
3119 (2016).
47. Szitenberg, A., John, M., Blaxter, M. L. & Lunt, D. H. ReproPhylo: An
Environment for Reproducible Phylogenomics. *PLoS Comput. Biol.* **11,** e1004447
(2015).
48. Riaz, T. *et al.* ecoPrimers: inference of new DNA barcode markers from whole
genome sequence analysis. *Nucleic Acids Res.* **39,** e145 (2011).
49. Ficetola, G. F. *et al.* An *In silico* approach for the evaluation of DNA barcodes.
BMC Genomics **11,** 434 (2010).
50. Zhang, Z., Schwartz, S., Wagner, L. & Miller, W. A greedy algorithm for aligning
DNA sequences. *J. Comput. Biol.* **7,** 203–214 (2000).
51. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects
Models Using lme4. *J. of Stat. Softw.* **67,** 1–48 (2015).
52. Veech, J. A. A probabilistic model for analysing species co-occurrence. *Glob.*
Ecol. Biogeogr. **22,** 252-260 (2013).
53. Griffith, D., Veech, J. & Marsh, C. cooccur: Probabilistic Species Co-Occurrence
Analysis in R. *J. Stat. Softw.* **69,** 1–17 (2016).
54. Fox, J. & Weisberg, S. *An R companion to applied regression, Second Edition.*
(Sage Publications, Thousand Oaks, CA, USA, 2011).
55. Therneau, T., Atkinson, B. & Ripley, B. rpart: Recursive Partitioning and
Regression Trees. R package version 4.1-13 (2018).
56. Bjørnstad, O. N. ncf: spatial nonparametric covariance functions. R package
version 1.1-7 (2016).
57. Pinheiro, J. C. & Bates, D. M. *Mixed-effects in S and S-Plus.* (Springer, New York,
USA, 2000).
58. McCulloch, C. E. & Searle, S. R. *Generalized, Linear, and Mixed Models* (John
Wiley & Sons, Inc., Hoboken, NJ, USA, 2001).
59. Zuur, A. F., Ieno, E. N., Walker, N., Saveliev, A. A. & Smith, G. M. *Mixed effects*
models and extensions in ecology with R (Springer, New York, USA, 2009).

60.   Akaike, H. Maximum likelihood identification of Gaussian autoregressive moving
average models. *Biometrika* **60,** 255-265 (1973).

61.   Hervé, M. RVAideMemoire: Diverse basic statistical and graphical functions. R
package version 0.9-69. (2018).

62.   Hosmer, D. W. & Lemeshow, S. *Applied Logistic Regression* (John Wiley & Sons,
Inc., Hoboken, NJ, USA, 2000).

63.   Lele, S. R., Keim, J. L. & Solymos, P. ResourceSelection: Resource Selection
(Probability) Functions for Use-Availability Data. R package version 0.3-2
(2017).

64.   Landwehr, J. M., Pregibon, D. & Shoemaker, A. C. Graphical Methods for
Assessing Logistic Regression Models. *J. Am. Stat. Assoc.* **79,** 61–71 (1984).

65.   Mazerolle, M. J. AICcmodavg: Model selection and multimodel inference based
on (Q)AIC(c). R package version 2.1-1 (2017).

66.   Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*. (Springer, New York,
USA, 2009).

67.   Manenti, R. & Pennati, R. Environmental factors associated with amphibian
breeding in streams and springs: effects of habitat and fish occurrence. *Amphib-*
Reptil. **37,** 237–242 (2016).

68.   Maceda-Veiga, A., López, R. & Green, A. J. Dramatic impact of alien carp
Cyprinus carpio on globally threatened diving ducks and other waterbirds in
Mediterranean shallow lakes. *Biol. Conserv.* **212,** 74–85 (2017).

69.   Manchester, S. J. & Bullock, J. M. The impacts of non-native species on UK
biodiversity and the effectiveness of control. *J. Appl. Ecol.* **37,** 845–864 (2000).

70.   Aronsson, S. & Stenson, J. A. E. Newt-fish interactions in a small forest lake.
Amphib-Reptil. **16,** 177–184 (1995).

71.   Ficetola, G. F. & De Bernardi, F. Amphibians in a human-dominated landscape:
the community structure is related to habitat features and isolation. *Biol. Conserv.*
119, 219–230 (2004).

72.   Chan, K. Can great crested newts (*Triturus cristatus*) coexist with fish?
(University College London, 2011).

73.   Hecnar, S. & M'Closkey, R. Amphibian species richness and distribution in
relation to pond water chemistry in south-western Ontario, Canada. *Freshw. Biol.*
(1996).

74.   Gustafson, D. H., Pettersson, C. J. & Malmgren, J. C. Great crested newts
(*Triturus cristatus*) as indicators of aquatic plant diversity. *Herpetol. J.* **16,** 347–
352 (2006).

75.   Paillisson, J.-M. & Marion, L. Interaction between coot (*Fulica atra*) and waterlily
(*Nymphaea alba*) in a lake: the indirect impact of foraging. *Aquat. Bot.* **71,** 209–
216 (2001).

76.   Wallau, G. L. *et al.* Behaviour of the Common Moorhen in Rio Grande do Sul,
Brazil. *Acta Ethol.* **13,** 127–135 (2010).

77.   Perrow, M. R. *et al.* Interactions between coot (*Fulica atra*) and submerged
macrophytes: the role of birds in the restoration process. *Hydrobiologia* **342/343,**
241–255 (1997).

78. Lauridsen, T. L., Jeppesen, E. & Andersen, F. Ø. Colonization of submerged
macrophytes in shallow fish manipulated Lake Væng: impact of sediment
composition and waterfowl grazing. *Aquat. Bot.* **46,** 1–15 (1993).
79. Battersby, J. & Partnership, T. M. *UK Mammals: Species Status and Population
Trends. First Report by the Tracking Mammals Partnership* (JNCC/Tracking
Mammals Partnership, Peterborough, 2005).
80. Almeida, D., Rodolfo, N., Sayer, C. D. & Copp, G. H. Seasonal use of ponds as
foraging habitat by Eurasian otter with description of an alternative handling
technique for common toad predation. *Folia Zool. Brno.* **62,** 214–221 (2013).
81. Rice, C. N. Abundance, impacts and resident perceptions of non-native common
pheasants (*Phasianus colchicus*) in Jersey, UK Channel Islands. (University of
Kent, 2016).
82. Freshwater Habitats Trust. *Pond Habitat Survey: Survey Manual* (Freshwater
Habitats Trust, Oxford, 2015).
83. Ilg, C. & Oertli, B. Effectiveness of amphibians as biodiversity surrogates in pond
conservation. *Conserv. Biol.* **31,** 437-445 (2016).
84. Sewell, D. & Griffiths, R. Can a Single Amphibian Species Be a Good
Biodiversity Indicator? *Diversity* **1,** 102–117 (2009).
85. Beutel, M. W. & Larson, L. Pathogen removal from urban pond outflow using
rock biofilters. *Ecol. Eng.* **78,** 72–78 (2015).
86. Piha, H., Luoto, M. & Merilä, J. Amphibian occurrence is influenced by current
and historic landscape characteristics. *Ecol. Appl.* **17,** 2298–2309 (2007).
87. Semlitsch, R. D., Peterman, W. E., Anderson, T. L., Drake, D. L. & Ousterhout,
B. H. Intermediate pond sizes contain the highest density, richness, and diversity
of pond-breeding amphibians. *PLoS One* **10,** e0123055 (2015).
88. Landi, M., Piazzini, S. & Saveri, C. The response of amphibian communities to
fish and habitat features in Mediterranean permanent ponds. *Biologia* **69,** (2014).
808

Acknowledgements

This work was funded by University of Hull. We would like to thank Jennifer Hodgetts (Fera) for assisting with sample collection, and Jianlong Li (University of Hull) for primer design and advice on laboratory protocols. Furthermore, Barbara Mabel (University of Glasgow), Andrew Buxton and Richard Griffiths (DICE, University of Kent) provided tissue samples for primer validation and Sanger sequencing to supplement the reference database.

Author contributions

B.H., L.R.H., L.L.H and N.B. conceived and designed the study. H.C.R. and N.B. contributed samples for processing. L.R.H. performed laboratory work and analysed the data. I.P.A. and E.L. offered advice on and supervised sequencing. C.H. assisted with bioinformatics analysis. P.B. and S.P. contributed datasets for analysis. L.R.H. wrote the manuscript, which all authors revised.

Competing interests

The authors declare no competing financial interests.

Materials and correspondence

All requests should be addressed to L.R.H., B.H. or L.L.H.

Table 1 | Summary of established and newly identified abiotic and biotic determinants of great crested newt occupancy. Reported effects on great crested newt occupancy in the literature and hypothesised effects on eDNA-based crested newt occurrence are given for each determinant. Any determinants not reported in the literature are listed as NR. Direction of observed effects on eDNA-based great crested newt occupancy determined by each analysis (GLMM assessing number of species in each vertebrate group, N = 532; co-occur analysis, N = 532; GLMM combining abiotic and biotic factors N = 504; and GLMM assessing HSI, N = 504) are given. No, negative and positive effects are listed as 0, - and + respectively. For categorical variables with more than one level, effect size and standard error (SE) are only given for levels reported in the model summary. Test statistic is for LRT used. Significant P-values (<0.05) are in bold.

Determinant	Effect reported	Hypothesised effect	Analysis					
			Co-occur		GLMM			
			Effect	P	DF	Effect size (SE)	χ^2	P
Fish	-/0	-			1	-0.239 (0.124)	4.065	**0.044**
Three-spined stickleback	-	-	-	**0.0091**	1	-1.432 (0.561)	9.453	**0.0021**
Ninespine stickleback	-	-	-	**0.0472**				
Common carp	-	-	-	0.0704				
Crucian carp	-	-						
Waterfowl	-	-			1	0.617 (0.181)	13.050	**0.0003**
Coot	NR		+	**0.0232**				
Moorhen	NR		+	**0.0007**				
Green-winged teal	NR		-	0.0987				
Amphibians	NR				1	0.558 (0.149)	16.640	**4.158x10^{-5}**
Smooth newt	+	+	+	**< 0.0001**	1	1.081 (0.303)	17.434	**2.975x10^{-5}**
Common toad	NR		-	**0.0088**	1	-1.635 (0.696)	8.228	**0.0041**
Terrestrial birds	NR				1	-0.335 (0.291)	1.444	0.2295
Common pheasant	NR		-	**0.0479**				
Mammals	NR				1	0.028 (0.091)	0.095	0.7583
Grey squirrel	NR		-	**0.0183**	1	-1.591 (0.534)	12.432	**0.0004**
Badger	NR		-	0.0987				
Pig	NR		+	**0.00395**				
Cow	NR		+	0.0971				

Pond area	-/+	-	1	-0.0004 (0.0002)	6.453	**0.0111**
Pond density	+	+				
Pond depth	+	+	1	0.282 (0.139)	4.266	**0.0389**
Water quality	+	+				
Outflow	NR		1	-0.713 (0.359)	4.467	**0.0346**
Macrophyte cover	-/+	-				
Shading	-/+	-				
Woodland	+	+				
Grassland	+	+				
HSI	0/+	+	1	3.0198 (0.7912)	15.709	**7.388x10^{-5}**
Ruderal	NR		2		6.507	**0.0387**
None				-0.617 (0.527)		
Some				0.032 (0.528)		
Other good terrestrial habitat	NR		2		7.918	**0.0191**
None				0.428 (0.429)		
Some				-0.316 (0.424)		
Species richness	NR		1	0.527 (0.105)	60.267	**8.281x10^{-15}**

Figure 1 | Great crested newt presence (orange) and absence (grey) in relation to number of species from different vertebrate groups detected by eDNA (*N* = 532 ponds). a, other amphibians; **b,** fish; **c,** waterfowl; **d,** terrestrial birds; **e,** mammals. Observed proportion of ponds with and without great crested newt (left) is plotted alongside predicted probability of great crested newt occurrence in ponds as determined by the binomial GLMM (right). Numbers on barplots of observed occupancy are the number of ponds for each category. In plots showing predicted crested newt occupancy, the observed data is shown as points which have been jittered around 0 and 1 to clarify variation in point density. Blue points are outliers and boxes are the model predictions.

Figure 2 | Heat map showing significant (P < 0.05) positive and negative species associations determined by the probabilistic co-occurrence model for the eDNA metabarcoding presence-absence data (N = 532 ponds). Species names are positioned to indicate the columns and rows that represent their pairwise relationships with other species. Species are ordered by those with the most negative interactions to those with the most positive interactions (left to right). Associations relevant to great crested newt are highlighted in red.

Figure 3 | Biotic and abiotic determinants of great crested newt occurrence, as predicted by the

binomial GLMM (*N* = 504 ponds). a, smooth newt occurrence, **b,** common toad occurrence, **c,** three-

spined stickleback occurrence, **d,** grey squirrel occurrence, **e,** pond outflow, **f,** ruderal habitat and **g,**

other good quality terrestrial habitat, **h,** species richness, **i,** pond area, **j,** pond depth. The 95% CIs, as

calculated using the predicted great crested newt probability values and standard error for these

predictions, are given for each relationship. The observed great crested newt presence (orange) and

absence (grey) data are also displayed as points, which have been jittered around 0 and 1 to clarify

variation in point density, against the predicted relationships (boxes/lines). Outliers are indicated by

blue points.

Figure 4 | Relationship between great crested newt occupancy and HSI score, as predicted by the

binomial GLMM (*N* = 504 ponds). The 95% CIs, as calculated using the predicted great crested newt

probability values and standard error for these predictions, are given. The observed great crested newt

presence (orange) and absence (grey) data are shown as points, which have been jittered around 0 and

1 to clarify variation in point density, against the predicted relationship (line).

Figure 5 | Abiotic and biotic determinants of vertebrate species richness, as predicted by the Poisson GLMM (*N* = 504 ponds). a, outflow, **b,** rough grass habitat, **c,** overall quality of terrestrial habitat, **d,** percentage of terrestrial overhang, **e,** percentage of macrophyte cover, **f,** pond density, and **g,** HSI score. The 95% CIs, as calculated using the predicted species richness values and standard error for these predictions, are given for each relationship. The observed data are also displayed as points, which have been jittered around 0 and 10 to clarify variation in point density, against the predicted relationships (boxes/lines). Outliers are indicated by red points.