Assessing the Relation between the Spatiotemporal Microhabitat Heterogeneity

and the Dispersion Patterns of Two Syntopic Small Mammals in a Midlatitude

Temperate Forest

De la Cruz-Arguello Ivan M.ᵃ (imda@xanum.uam.mx), Castro-Campillo Alondraᵃ (acc@xanum.uam.mx),

Zavala-Hurtado Alejandroᵃ (jazh@xanum.uam.mx), Salame-Méndez Arturoᵇ (asam@xanum.uam.mx), and

Ramírez-Pulido Joséᵃ (jrp@xanum.uam.mx).

Authors affiliations:

^{a} *Department of Biology (DB) and* ^{b} *Department of Reproduction Biology (DBR), Biological and Health*

Sciences Division (DCBS), Metropolitan Autonomous University campus Izatapalapa (UAM-I). AP 55-

535- Av. San Rafael Atlixco 186, Vicentina, 09340 Iztapalapa, CDMX, México.

Running title: Ecological Dispersion of Syntopic Small Mammals

Abstract

One of the basics problems in ecology is to understand the factors that shape the spatial

patterns in the distribution of the species and the coexistence of close relatives species.

Among the most critical factors governing the distributions and the coexistence of

species are the spatiotemporal changes occurring in the microhabitat heterogeneity.

Here, we assessed the heterogeneity of microhabitats and how they affect the spatial

segregation of two species of small mammals (*i. e.*, *Peromyscus difficilis* and *P.*

melanotis), which coexist in a temperate, mixed forest. We evaluated the microhabitat

heterogeneity through multivariate statistics, using onto 23 habitat variables describing

horizontal-vertical habitat structure along pluvial seasons. To detect specific

microdistribution changes and habitat preferences by two species of small mammals, we

used second-order spatial statistics and general linear models. According to their

respective morphology and locomotive adaptations, the middle sized, midscansorial *P.*

difficilis was resident all year long and preferred microhabitats with a high log ground

cover, while the opportunistic, small sized, cursorial *P. melanotis* changed its

occupancy area, depending on the density of herbaceous and woody plants cover. Under

the more benign microhabitat conditions of the rainy season (denser plant coverage,

milder temperature), both species showed closer microdistribution patterns; while these

became repulsive at the less favorable conditions of the dry season (scarcer plant cover,

colder temperature). Thus, we could confirm that seasonal changes of microhabitat

heterogeneity promoted *Peromyscus* coexistence, through dispersion patterns reflecting

partition of microhabitat resources.

Keywords: Microhabitat selection, Use of space, Small mammals, Dispersion, Resource

partition.

Introduction

Spatiotemporal habitat heterogeneity has a substantial impact on coexistence among

species since they become spatially segregated, according to niche preferences such as

resource requirements (Valladares *et al*. 2015). Model simulations reveal the potentially

important role of spatial heterogeneity and its complex and delicate interplay with

dispersal in mediating long-term outcomes of species coexistence (Schreiber and

Killingback 2013; Valladares *et al*. 2015). For example, an increase in the number of

habitat types, resources, and structural complexity should increase the available niche

space and thus allow more species to coexist (Currie 1991). Equally crucial for the

maintenance of coexistence is the heterogeneity in time, with an influence on natural

communities also variable depending upon the temporal scales (Valladares *et al*. 2015).

Temporal fluctuations on habitat structure can stabilize coexistence via storage effect

(Chesson 2000), when inter-annual variation in climate or resource availability favors

alternatively one group of species over the others (Zavaleta *et al*. 2003). For instance,

Peromyscus mice have been shown to respond to spatiotemporal habitat heterogeneity

regarding species coexistence (M'Closkey and Fieldwick 1975; M'Closkey and Lajoie

1975), spatial segregation (Monamy and Fox 1999), and competition (Morris 1984;

Seagle 1985).

Heterogeneous habitats contain patches of varying size, distribution, resources,

environmental conditions, and species composition (Fahrig and Merriam 1994) that

vary spatiotemporally, depending on observation scale and habitat type under study

(Wiens 2000). It is highly likely that small size species such as small mammals perceive

spatial habitat heterogeneity at a fine scale (microhabitat). Therefore, both movement

and foraging by these kinds of mammals are affected by habitat heterogeneity (Bowne

et al. 1999), having different responses among species or demographic groups within

them (Dooley and Bowers 1996). Here we focus on whether small mammals are

capable of perceiving and to respond to spatiotemporal microhabitat heterogeneity.

Small mammals use resources selectively, based on their requirements for

growth, survival, and reproduction (Johnson and Gaines 1980). Several studies have

quantified variation in resource use in heterogeneous environments either among

congeneric and co-family mice species within a community (Kaufman and Kaufman

1989 and references therein). Indeed, syntopic related species are especially valuable for

studies of spatial and resource partitioning, since they are most likely to be current or

past competitors (Kalcounis-Rüppell and Millar 2002). Since habitat use varies

spatiotemporally (Haim and Rozenfeld 1993) due to distribution and availability of

resources, we expect that such variation also affects the use of resources and

relationships among non-closely related species. In this study, the primary objective is

to analyze how heterogeneity of microhabitats is structured and change through pluvial

seasons in a temperate, mixed forest, and how such changes affect the coexistence of

two syntopic species of *Peromyscus* (*i. e., P. difficilis* and *P. melanotis*). Therefore, 1)

We assess the presence of the spatiotemporal heterogeneity at a fine scale

(microhabitat) according to vertical and horizontal structure indicators. 2) We evaluate

the ecological dispersion patterns of two species of *Peromyscus* and is tested if the use

of space of the two small mammals is affected by the spatiotemporal changes in

microhabitat heterogeneity. 3) We determine what kind of spatial relationships occur

between both species, depending on seasonal changes in microhabitats heterogeneity (*i.*

e., attraction or positive: sharing of same microhabitats and resources; repulsion or

negative: not sharing). Finally, we assessed which specific structural elements of each

one of the microhabitats better explain the capture frequency of the two *Peromyscus*

species between seasons.

105   **Material and methods**

Ethics statement

Animal capture and handling followed guidelines of the American Society of

Mammalogists (Sikes *et al*. 2011). If animals died, corpses were prepared as voucher

specimens to house them at the Mammal Collection of the Universidad Autónoma

Metropolitana-Iztapalapa (UAMI, Ramírez-Pulido *et al*. 1989). Collecting permit,

SEMARNAT-08-049-B, was issued by DGVS, SGPA-09712/13, SEMARNAT, to

Alondra Castro-Campillo (ACC).

Study area

The study area was located in a temperate forest of conifers and broad-leaved trees at

Parque Nacional Desierto de Los Leones (PNDL, CONANP 2006), Mexico City, within

the Transmexican Neovolcanic Range. Temperature averages 12.6 ± 6 °C from April to

July, while it descends to 8.1 ± 2 °C from December to February. Rainfalls occur from

late spring throughout summer (May to August), with average precipitation of 235 ± 30

mm, though this period can go until mid-fall (October). From winter to early spring

(December to March), precipitation average becomes 12 ± 4 mm, introducing a dry and

cold climate (PNDL, CONANP 2006). Dominating vegetation by strata includes

canopy, *Abies religiosa, Pinus hartwegii, P. leiophylla, P. montezumae, Prunus*

serotina, Garrya laurifolia, Salix paradoxa, and Buddleia cordata; understory shrubs,

Baccharis conferta and *Senecio spp.*, and herbs, *Acaena elongata, Sigesbeckia*

jorullensis, and *Alchemilla procumbens*; ground level, a rich variety of mosses and

fungi.

Mice sampling

We captured mice along ten months (October 2013- August 2014), including the dry

and wet seasons within a 2,475 m² surface plot (Fig. 1) at 2289 m. The plot was gridded

(12 vertical lines, A-L; 10 horizontal lines, 1-10) marking intersections each 5 m with

buried wood stakes (150x2.5x2.5 cm) to construct a coordinate system for independent

sampling stations, where we set 120 Sherman live traps (Tallahassee, FL 32303, USA),

baited with oat flakes and vanilla scent (Fig. 1). Traps were set for two continues nights

(2400 trap-nights), shortly before dusk, left open overnight, and checked at next day

dawn. To avoid bias in the abundance and use of space of both *Peromyscus* during each

sampling session, we used a temporal mark (gentian violet applied in the abdomen of

the mice) to identify the captures and recaptures. If we saw signals that the gentian

violet was disappearing in the abdomen of the mice, we applied it again. This temporal

mark enabled us to count the net frequency of the individuals in each microhabitat. We

carried out an initial analysis with the recaptures, and we found the same patterns that

we obtained with the first frequency (without recaptures). Thus, for the analyses, we

used only the first capture data for each of the 120 sampling stations during all trapping

sessions. All the traps were cleaned, and bedding was changed at each trap check event.

Also, all the traps were set randomly in each sampling station to avoid bias in frequency

quantification due to recurrence behavior or shyness of the mice to the trapping.

Coordinates of all trapped mice were recorded and the species and sex determined

through visual inspection.

Microhabitat features

To assess spatiotemporal heterogeneity of microhabitat within the entire plot, we

delimited an influence zone (sub-plot, Fig. 1; close-up, Fig. 1), drawing a rectangle of

2.5 m² around each sampling station. 23 variables were sampled (Table 1) once in each

influence zone in both seasons. The sampling of the entire plot took ten days, we

sampled one transect per day, starting at 9 am in the second week of the most

represented month of rains (July) and in February for the dry season (PNDL, CONANP

2006). These variables have been proved to be components of the vertical and

horizontal structure of the microhabitat. Also as factors affecting the distribution of

small mammals at fine scales (Morris 1984, 1987; Stapp 1997; Jorgensen 2004,

Coppeto *et al*. 2006; Villanueva-Hernández *et al*. 2017). Environmental variables

include vegetation coverage at different heights (VC10-100 cm), number of herbaceous

forms (FH11-15), and number of woody forms (VF21-25) at five different heights: 10,

25, 35, 50, and 100 cm, respectively. Rock coverage (RC), logs coverage which include

stumps and fallen trunks (CLO); coverage of organic matter (OM); coverage of bare soil

(BS); vegetation species richness (VR); total herbaceous plants (TH); total woody plants

(TW) and the number of burrows (BW) also were sampled (Table 1). Overall, these

variables provide information about possible refuges for the small mammals against

predators, some elements of microhabitat also imply spaces for resting, breeding, for

food resources and safe roads to move from one place to another, avoiding predation

risk (Jorgensen 2004). Detailed information about ecological meanings for each variable

appears in Table 1.

We adapted Canfield's (1941) method of Line Intercept (LI) to measure all

variables in each pluvial station, for it allows sampling within-plot variation and

quantifies changes, both in plant species cover and height over time, by using transects

located within a plot. We drew eight graduated transects (cm) from the respective

influence zone of each station to cover the 2.5 m² sub-plots (Fig. 1); four 2.5 m transects

were cardinally oriented (N, S, E, W), while the other four 3.30 m ones were diagonally

oriented (NE, NW, SE, SW). Only plants intercepted by diagonal transects were

counted for vegetation species richness, number of life forms at different heights

(herbaceous and woody plants), and total herbaceous and woody plants; while all

transects were used to measure plant coverage at different heights, as well as coverage

of fixed elements (*i. e.,* rocks, logs, bare soil, dead organic matter). We calculated

coverage percentage of each variable within the sub-plots with the formula $\sum(Xi)\frac{100}{2400}$,

where *Xi* is centimeters occupied by every plant in each transect, and 2400 equals total

longitude (sum of the eight transects). We counted the number of woody and

herbaceous plant life forms at different heights and burrows within each influence zone

of every sampling station. We took each hole in the ground ca. 8-12 cm of diameter as a

potential burrow for both *Peromyscus* (Álvarez-Castañeda 2005; Fernandez *et al*. 2010).

Statistical data analysis

We standardized all variables and run statistical tests at $p \leq 0.05$. We did not found

multicollinearity among the 23 variables which was assessed by examining the Variance

Inflation Factor (*i. e.,* all values were in a threshold <6) (Zar 1999). To identify and

categorized microhabitat heterogeneity within the grid, we performed two independent

Hierarchical Cluster Analysis (HCA) in JMP® (ver. 9, SAS Institute Inc., Cary, NC,

1989-2007) for each season. These analyses enabled us to cluster the sample stations

with similar characteristics according to the sampled variables for both seasons. We

used Ward´s method (1963) where the distance between two clusters is the ANOVA'

sum of squares between them, added up over all variables. Visual inspection of the

dendrogram and no drastic changes in the variance scree plot gave us three general

groups of stations for both seasons, which we interpreted as distinct microhabitat types

(M1-3). We further statistically validated such microhabitats carrying out independent

Discriminant Analyses for both seasons (DA, Addinsoft SARL's XLSTAT 2013; F =

9.99 dry; F = 9.64, rains; Wilk's Lambda = 0.0001 in both) (Fig. A.1). These analyses

allow us to know what type of microhabitat corresponded to each sampling station.

Thus, we constructed two different maps, assigning to each sampling station a color

mark that represented the three different microhabitats for both seasons (M1: red, M2:

green, M3: blue) (Fig. 2, Fig. A.1). To create typologies that describe at a fine scale the

behavior of each one of the 23 variables within three microhabitats for both seasons, we

built a plot whereas "Y" axis comprises of the 23 standardized variables and in the "X"

axis corresponds to the three microhabitats constructed with the same 23 variables (Fig.

A.1).

Due to not all 23 variables did not reveal a normal distribution, we applied two

different tests to analyze the distribution changes on each microhabitat variable between

the rainy and dry seasons. First, distribution normality was checked for every variable

with Kolmogorov-Smirnov tests (Zar 1999). Thus, we used Student's t-parametric tests

for variables normally distributed and non-parametric Wilcoxon test for non-normal

variables (Table 2). To know whether the number of stations structuring each type of

microhabitat changed between seasons (seasonal microhabitat heterogeneity), we used a

contingency table and x^2 test. All these analyses were performed with XLSTAT 13

(Addinsoft SARL). To examine the association between changes in spatial

heterogeneity of microhabitats and dispersion patterns of *Peromyscus*, along with the

entire grid, we conducted a Contingency Tables Analysis. Then, for visualizing

associations between the *Peromyscus* and microhabitats, we performed a

Correspondence Analysis with the constructed Contingency Tables for both seasons,

including the abundance of each species in the three microhabitats (Table A.2).

Spatial analysis

To map variations in point density captures of small mammals, and to find density

gradients across the plot area, we used the Kernel function (PAST, ver. 3.14, Hammer *et*

al. 2001). To test the ecological dispersion patterns of both species within the plot (*i.e.*,

clustering or overdispersion) for both seasons, we used a "Nearest Neighbor Analysis"

(Clark and Evans, 1954) using the PAST software (ver. 3.14, Hammer *et al*. 2001). We

applied the Wrap-around edge effect adjustment which is only appropriate for

rectangular domains as our plot sampling (Hammer *et al*. 2001). In general, the

"Nearest Neighbor Analysis" compares the mean distance of each main individual from

its nearest neighbor of the same species with the mean distance expected for a set of

points randomly dispersed at the same density (Vázquez and Álvarez-Castañeda 2011).

The ratio of the observed mean distance to the expected distance (*R*) indicates how the

observed distribution deviates from random. Clustered points give *R<1*, Poisson

patterns give *R~1*, while overdispersed points give *R>1*.

To assess whether both species were associated or disassociated, we compared

dispersion patterns between the two mice, using Ripley's K bivariate function (Ripley

1977), since the method considers all distances among individuals located under a

Cartesian scheme (X, Y) in a quadrat plot (Ripley 1976, Dale 1999, Zavala-Hurtado *et*

al. 2000). We used PASSaGE (*ver*. *2*, Rosenberg and Anderson 2011) to carry out

Ripley's bivariate K analysis. We used the option to test the associations conditional on

current locations hypothesis in PASSaGE. In this test, the point locations remain fixed,

and only the types are randomized (the two *Peromyscus*). The number of each type

remains fixed, but the association of each type with a specific, fixed location is

randomized. In this case, one is not testing whether the points are themselves random or

clustered, but rather whether the association of type A with type B (*P. difficilis vs. P.*

melanotis) is what one would expect given the locations of the points as fixed

(Rosenberg and Anderson 2011). We assigned coordinates to each sampling stations

considering 5 m distance between them. Therefore, because each sampling station had a

Cartesian coordinate system, we used as data input each station were we capture mice.

Thereby, our null hypothesis involved independence between our two species *P.*

difficilis and *P. melanotis* (Ripley 1976, Diggle 1983, Dale 1999), since we were

interested in whether points of a specific type (*P. difficilis*) were associated, or

disassociated, with points of the other specific type (*P. melanotis*). In other words,

Ripley's $K_{12}(d)$ allowed us to assess spatial attraction or repulsion between the two

species, among the stations of the plot. To evaluate the statistical significance of $K_{12}(d)$,

we estimated 95% confidence envelops (95% CI), using a Monte Carlo procedure,

based on 1000 stochastic relocation simulations of the sampling stations in the plot

(Upton and Fingleton 1985; Bailey and Gatrell 1995; Zavala-Hurtado *et al*. 2000).

When L(*d*) was positive and took values over the upper limit of the confidence region,

we inferred dissociation or repulsion between the *Peromyscus* at the corresponding (*d*)

scale; whereas a significant negative deviation indicated us a pattern of association or

attraction between the two mice (Dale 1999). If L(*d*) remains within the limits of the

95% CI for a given value of *d*, the null hypothesis of independence between the two

contrasts cannot be rejected (Dale 1999). The height of the L(*d*) function (peak height)

indicates the intensity of the association or repulsion. Control for edge effect was

carried out for analysis by rescaling the counts based overlap with the study boundary

setting the options area/volume which has been recommended for quadrats plots

(Rosenberg and Anderson 2011).

Prediction of microhabitat elements affecting the use of space by each species

The next *s*tep was to assess which specific environmental variables best explained the

frequency of both species within the plot at each season. For this, we performed

Generalized Linear Models (GLMs), using JMP® (ver. 9, SAS Institute Inc., 1989-

2007, Cary, NC,). First, we conducted a Principal Component Analysis (PCA) on the

means of the 23 variables to obtain functions summarizing the most significant variance

at each season. Because in both seasons the three first functions summarized a good

portion of the variance (38.61 for the dry season and 45.92 for rains), we used these

functions to construct the GLMs. Therefore, we assessed different models using the PCs

as effect variables to explain the frequency of the individuals of both species for both

seasons. Then, to visualize the fit of the model, the predicted values from the GLMs

were plotting against the Principal Components of the best model chosen. For each

species, the response variable in the models was the capture frequency at each trapping

station of the grid, and we assumed a Poisson distribution with a log link function.

301   **Results**

Frequency of captures

The total number of individuals captured from both species during the dry season was

111, while for the rainy season was 168. For *P*. *difficilis* the total number of frequency

capture in the dry season was 64. For *P*. *melanotis* during the dry season, we captured 47

individuals. On the other hand, during the rainy season, the frequency of capture of *P*.

difficilis was 87. While the total captures for *P*. *melanotis* were 81 (Table A.2).

The Behavior of 23 variables of microhabitat between seasons

When considering the mean value over all stations, 16 out of the 23 environmental

variables showed a significant difference between the two seasons (Table 2, Fig. 3, Fig.

A.1). The main changes were observed for variables related to herbaceous vegetation,

according to plant coverage at different heights, and vegetation species richness (VR).

As expected in overall, woody life forms were more stable between seasons (Table 2,

Fig. 3, Fig. A.1). Components of microhabitat showing no statistical change between

seasons were frequency of herbaceous plants at 10 cm (FH11), the frequency of both 35

cm (FW23) and 50 cm (FW24) woody plants, as well as coverage of logs (CLO), rocks

(RC), and bare soil (BS). Therefore, space configuration given by these structural

features remained stable in both seasons (Fig. 3).

Microhabitat heterogeneity

The dendrogram from cluster analysis revealed three different classes of microhabitats

(M1, M2, and M3) within the sampling plot in each season (Fig. A.1). M1 included

more sampling stations during the dry season, while M2 covered the majority of

sampling stations during the rainy season (Fig. 2, Fig. A.1, Table A.1). Indeed, Fisher's

exact test of Contingency Tables Analysis revealed significant changes in the number of

sampling stations ($p = 0.004$) from dry to rains. The main changes occurred between

M1 ($x^2 = 2.123, p < 0.05$) and M2 ($x^2 = 3.348, p < 0.05$), especially in the latter, since it

augmented 24 sampling stations from the dry season to the rainy season (Fig. 2, Table

A.1). Conversely, the number of sampling stations remained quite stable in M3 ($x^2 =$

0.011) throughout the study (Fig. 2, Fig. A.1, Table A.1).

Discriminant analyses validated heterogeneity of the three microhabitats (Fig. 2)

within and between pluvial seasons. the percentage of variance associated to each

discriminant function was clearer during the rainy season (dry season: F1 = 58.67 %, F2

= 42.32 %; rainy season: F1 = 72.581 %, F2 = 27.419 %) (Fig. 2, Fig. A.1).

Discriminant functions explained variation due to woody plants in the understory in the

dry season, while herbaceous vegetation together with decayed matter and logs became

more relevant at rains (Table A.3). Wilk's Lambda showed that at least the mean of one

microhabitat differed statistically from the others either in the dry and rainy season ($p =$

0.0001; $p = 0.0001$, respectively) (Fig. A.1). These results corroborated spatial and

temporal heterogeneity drawn from cluster analysis of individual station features in the

grid plot. During the dry season, 93.33 % of individual sampling stations remained

correctly classified, and 94.17 % of them during the rainy season (Fig. 2).

Description of microhabitats within the plot

Microhabitat 1. M1 (Fig. 2, Fig. A.1) was characterized by low amounts, or absence, of

vegetation cover at different heights (VC) and by a low frequency of herbaceous and

woody plants (FH and FW, respectively). In contrast, organic matter on the ground

(OM) was the most frequent component. This microhabitat was also distinctive for

having extensive coverage of logs (CLO) on the ground and for being the microhabitat

with the significant presence of burrows (BW).

Microhabitat 2. M2 (Fig. 2, Fig. A.1) was the largest area covered by herbaceous life

forms at different heights; vegetation species richness (VR) and the total number of

herbaceous plants (TH) also remained very high. Woody life forms at 25 cm (FW22)

were present but at low frequencies, while FW24 and FW25 remained at higher

frequencies. Coverage on the ground of small logs or dead wood (CLO), of rocks (RC),

and organic matter (OM) were shallow during the study. Bare soil surface (BS)

increased to higher amounts during the dry season, while it decreased during the rainy

season. There were no burrows (BW) in this microhabitat.

Microhabitat 3. In M3 (Fig. 2, Fig. A.1), VC25, VC35, VC50, and VC100 showed high

values, while vegetation species richness (VR) averaged the highest compared to other

microhabitats. The total number of woody plants (TW) and frequency of woody plants

at different heights (FW21-25) remained very high, while the total number of

herbaceous plants (TH) was low. Vegetation coverage at 10 cm (VC10), rocks (RC),

logs (CLO), and bare soil surface (BS) were low during the dry season; however, VC10

increased in the rainy season. Conversely, OM was present in the dry season but

decreased in the rainy season. The number of burrows (BW) remained quite scarce.

Association between the microhabitats and species

Kernel Graphs (Fig. 4) indicated that the two species used space distinctive since they

were differentially scattered along the plot. *Peromyscus difficilis* was more abundant

during the dry season, occupying a large portion of the plot. Instead, *P. melanotis*

increased its distribution during the rains when it seemed to displace *P. difficilis* into

other sampling stations (Fig. 4). Distribution rearrangements between seasons implied

that one species occupied some stations more frequently than the other, and vice versa

(Fig. 4). Indeed, the two x^2 tests yielded statistical abundance differences in each species

at both microhabitats and seasons (Table A.2: dry season; R^2 (U) = 0.09, $n = 111$, df = 2,

Likelihood Ratio $x^2 = 15.07$, $p = 0.00005$; rainy season; R^2 (U) = 0.02, $n = 168$, df = 2,

Likelihood Ratio $x^2 = 6.40$, $p = 0.0406$).

During the dry season (Fig. 5 and Table A.2), *Peromyscus difficilis* was highly

related to M2 (80 % captured mice) and also related to M1 (64 % captured mice);

conversely, *P. melanotis* was captured more frequently in M3 (66 % captured mice).

Spatial use of microhabitats changed for both species in the rains (Fig. 5 and Table

A.2), showing a microhabitat partition again, though less visible: in the Correspondence

Analysis, 61 % of mice caught in M1 were *P. difficilis*. This species was also related to

M3, while 60% of mice caught of *P. melanotis* were captured in M2.

Spatial Patterns

The Nearest Neighbor Analysis revealed that in both seasons both species showed a

significant clustered pattern within the plot (Table 3). In the dry season, the mean

distance between individuals of *P. difficilis* displayed to be more clustered than the

individuals of *P. melanotis*. However, the mean distance between individuals of *P.*

difficilis increases in the rainy season, while individuals of *P. melanotis* showed a

reduction in their area occupied (Table 3).

On the other hand, during the dry season, Ripley's bivariate K showed a

statistically significant pattern of repulsion for both *Peromyscus* in almost all analyzed

distances of the entire plot, excused at 2 m where it shows an attraction pattern (Fig.

6a). Such repulsion pattern turned over in almost all distances during the rainy season

(Fig. 6b), since the two species became more associated, sharing microhabitats in

almost all capture stations. However, during the rainy season, significant statistical

peaks of repulsion reappeared between the species at distances of five, ten, and 14 m

(Fig. 6b). Analyses also revealed the intensity of these patterns; *e.g.*, the likelihood for

finding individuals of *P. difficilis* and *P. melanotis* together at the same capture station

was very low during the dry season (Fig. 6a), while this probability increased in the

rainy season (Fig. 6b).

Structural components of microhabitats affecting the distribution of small mammals

Criteria for the selecting models that most explained the frequency of mice for each

species at each season in the sampling stations included: model significance of $p <$

0.05; lowest Akaike Information Criterion (AIC), *i. e.,* a measure of goodness-of-fit

penalized by the number of variables (Posada and Buckley 2004); Pearson goodness-of-

fit ($p < 0.05$) and its deviation (Deviance; $p < 0.05$); as well as the biological and

ecological meaning of results. Therefore, during the dry season, PC1 and PC3 predicted

the frequency of *P. difficilis* according to the best Generalized Linear Model for this

season (AICc 212.55) (Table 5, Fig. 5c). Environmental variables with higher scores in

the PC1 ($p = 0.0186$) were the total number of herbaceous plants (TH), vegetation

species richness (VR), and vegetation cover at 10 cm (VC10). In PC3 ($p = 0.0001$),

characterizing variables were the number of logs on the ground (CLO), as well as the

presence of burrows (BW) (Table 4, 5, Fig. 5c). These five variables were also the main

elements characterizing M2 (PC1) and M1 (PC3), the two habitats where the frequency

of capture of *P. difficils* was the higher. As for *P. melanotis* (Fig. 5c), the frequency of

capture of *P. melanotis* was positively related to PC2 (AICc = 204.47, $p = 0.0001$).

Here, among the four variables with higher scores, were the total number of woody

vegetation (TW) and frequency of woody life forms characterizing 10 cm height

(FW21), the two main elements at M3 for the dry season (Table 4, 5, Fig. 5c).

During the rainy season, a similar pattern of microhabitat partition occurred

between the two species. The best GLM model for *P. difficilis* in this season (Table 4, 5,

Fig. 5c) was mainly associated with the total number of woody plants (TW),

herbaceous forms at 50 cm (FH14), woody forms at 10 and 100 cm (FW21, FW25,

respectively) and with the number of logs on the ground (CLO), in PC2; and with the

number of logs on the ground (CLO), and organic matter (OM) in PC3 (AICc = 286.76,

$p = 0.0001$) (Table 4, 5). The latterly involved variables with a high presence at M1

(PC3) and M3 (PC2) during the wet season, where *P. difficilis* was more frequently

captured. On the other hand, *P. melanotis* (Fig. 5c) was only associated with PC1 (AICc

= 241.01, $p = 0.0343$). The primary variables characterizing this component were the

number of total herbaceous plants (TH), vegetation species richness (VR), and the plant

cover at almost all analyzed heights (VC10-100 cm) (Table 4).

450   **Discussion**

Our results of Contingency Tables Analysis, Correspondence Analysis, and GLMs

models showed that the two syntopic species of *Peromyscus* switched and split

resources in space, as environmental conditions changed. Also, our findings on

microhabitat relations of rodents in a patch of a mid-latitude temperate forest, concur

with the habitat heterogeneity coexistence hypothesis (Cramer and Willig 2002;

Valladares *et al*. 2015). We found that both *Peromyscus* were sensitive to slight changes

in microhabitat structure and that the three microhabitats provided different resources

for each one. Our results also suggest that coexistence between these syntopic

Peromyscus is facilitated by temporal differences in space structure, and probably also

by changes on the availability of both food and shelter that go with it (Pianka 1973;

Schoener 1974). For instance, *P. difficilis* was more related to M1 and M2 by the dry

season, while *P. melanotis* was highly related to M3. Conversely, *P. difficilis* was more

related to M1 and frequently captured to M3 by the rainy season, while *P. melanotis*

was highly related to M2. Partition in space use has already been documented for

coexisting species of *Peromyscus* within a community; *i.e.,* extensive studies of niche

partitioning by *Peromyscus leucopus* and *P. maniculatus* (Kaufman and Kaufman

1989). Moreover, several studies have demonstrated variation in spatial resource use by

congeneric species (Barry *et al*. 1990; Dooley and Dueser 1996; Kalcounis-Rüppell and

Millar 2002).

In our study, *P. difficilis* remained related to M1 in both seasons, especially to

the former during the wet season. The number of fallen logs was very high at M1. Such

microhabitat represents small patches of food for small mammals, offering invertebrates

(Bellows *et al*. 2001), refuges and shelters (Bowman *et al*. 2001). In this microhabitat,

individuals of *P. difficilis* can also use the long and big fallen logs as pathways for quick

and straight locomotion inside the forest (Bellows *et al.* 2001; Dewalt *et al.* 2003).

Indeed, fallen logs promote structural complexity of forests and may enhance positive

interaction among species of small mammals (Bowman and Facelli 2013). GLM's

analyses showed that PC1 and PC3 explained the abundance of *P. difficilis* during the

dry season and by PC2 and PC3 in the rainy season. In the dry season, variables

contributing most to PCA ordination in PC1 and PC3 were the total number of

herbaceous vegetation (TH), vegetation species richness (VR), vegetation coverage at

10 cm (VC10), herbaceous forms at 10 cm (FH11) and the number of logs on the

ground (CLO), as well as burrows (BW). While in the dry season, PC2 and PC3

represented woody vegetation (TW), the number of logs on the ground, herbaceous

forms at 50 cm (FH14), woody forms at 25 and 100 cm (FW21 and FW25, respectively)

and organic matter (OM). Bellows *et al.* (2001) found a similar result in a high latitude

temperate forest (Virginia, USA), where distribution of a small mammal, generalist

species was associated with the diameter of fallen logs, the frequency of shrubs, and

degree of canopy closure. Association between rodents and fallen logs was also

documented for *Nectomys squamipes* that builds up its nests inside decomposed, fallen

logs (Briani *et al.* 2004); this has also been recorded for other rodent species in several

biomes with different vegetation types (*e. g.*, *Rattus rattus*, *Nesomys audeberti* in

Lehtonen *et al.* 2001; *Oligoryzomys nigripes* in Dalmagro and Vieira 2005). In contrast,

P. melanotis showed a close relation with only one PC over each pluvial season. During

the dry season, the frequency of this species was only related to PC2, which represented

the total number of woody vegetation (TW) and FW21. Then by the rainy season, the

species became only related to PC1, which accounted for understory dominated by

herbaceous life forms (TH). Relationships between frequency of rodents and habitats

with increased understory density have been interpreted as a protection against aerial

predators (Dalmagro and Vieira 2005) or as a preference for more complex habitats that

enable for vertical stratifications and, thus for extended coexistence inside fragments

(Grelle 2003; Johnson 2007).

Intrinsic factors as sex and mating system must also be related to the coexistence

of both *Peromyscus*. For instance, the Nearest Neighbor Analysis revealed that in both

seasons both species showed a significant clustered pattern within the plot. However,

the mean distance between individuals of *P. difficilis* displayed to be more clustered

than the individuals of *P. melanotis*, especially during the dry season. Indeed, the mean

distance between individuals of *P. difficilis* increases in the rainy season, while

individuals of *P. melanotis* showed a reduction in their area occupied. These patterns

are related with the breeding season of both species. Our data (not published) revealed

that during the dry and rainy seasons were captured individuals of *P. difficilis* with signs

of reproductive activity, especially during the dry season (males: inguinal vs. scrotal

testicles, females: alopecia, the presence of milk, the appearance of the vagina,

gestation). While for *P. melanotis* the primary breeding season occurs during the rains.

Individuals need different resources requirements to fulfill the breeding season and this

impact on its dispersion and coexistence. We believe that *P. melanotis* may be moving

to other areas with more beneficial conditions (*i. e*. more availability of resources).

Thus, the coexistence between both *Peromyscus* may occur because the geographic

segregation of *P. melanotis* to search for better environmental conditions to survive.

Indeed, in this study, we define to *P. melanotis* as an opportunistic species throughout

the study area because the capture frequencies of the individuals only increased (as we

pointed out early) when the environmental conditions became more beneficial for the

survival of these. That is, throughout the time of capture except for the rainy season, its

frequency of capture remained at low levels compared to those of *P. difficilis* (resident

species), increasing drastically in the rainy season when there is greater accessibility for

the use of resources and competition between the two species decreases. Therefore, *P.*

melanotis may be coming back to the area during the rainy season to carry out the

breeding season. However, Chupp (2002) showed an opportunistic habitat use by small-

sized *Peromyscus leucopus*, which he related to predation risk. Further studies about the

geographic segregation and the influence of predation on the opportunistic behavior of

P. melanotis are needed. Also, the increase in the number of captures of both species

from the dry to the rains could be leaded for the breeding season and resources

availability. One possible consequence of such intrinsic and extrinsic operating factors

is that *P. difficilis* remained more abundant and always present at the microhabitats with

more stable elements in the study area (our unpublished data), while *P. melanotis* only

increased its abundance during the rainy season, when environmental conditions

became more benign, and plant cover became denser. Since rains promote an increase

of primary productivity, allowing more resource availability (mainly food) and

enhancing microhabitat carrying capacity, such habitat changes facilitate coexistence

between both *Peromyscus*, and with other small land mammals in the area (Castro-

Campillo *et al*. 2008, 2012). Ripley's bivariate K supported and shed light into such

outcome since the species showed repulsion between them in almost all analyzed

distances during the dry season; *i.e*., due to resources availability are more scarce during

this season, is not so common that a mixed pair of mice used the same microhabitat.

Moreover, as in this season is the primary breeding season of *P. difficilis* is probably

that territoriality behavior can occur. Conversely, the intensity of such repulsion

decreased substantially during the rainy season; indeed, at some analyzed distances, the

two deer mice showed an association pattern. Holding our results that the increase of

resources available during the rainy season can allow that the species can share the

habitat. Thus, our results indicate that in these two syntopic *Peromyscus*, both

microhabitat use and spatial organization are seasonally variable (Brown and Zeng

1989; Cramer and Willig 2002). It is important to point out that during both seasons

Ripley's bivariate K indicated an interesting attraction pattern between 2 and 3 m. This

patterns could be due to sampling design. Ergo we set only one Sherman trap in each

sampling station for two days, therefore if one individual of *P. difficilis* was captured in

the first day and then one individual of *P. melanotis* was captured in the same station

where we captured the *P. difficilis*, the analysis counts this pattern as an association

because they were trapped in the same station at a very short distance.

Also, the partition of space by this syntopic deer mice must be facilitated by

their respective locomotive habits as semiarboreal (*P. difficilis*) and cursorial (*P.*

melanotis). The long tail of *P. difficilis* must enable it to rush and climb along shrubs or

trees (Álvarez-Castañeda 2005), thus easing its preference for habitats where it can

escape from predators. In fact, adult color coat changed in *P. difficilis* between pluvial

seasons, becoming more similar to ground litter; *e. g.*, in this resident deer mouse, color

phenotype plasticity must be a cryptic response elicited against predators. On the other

hand, a smaller sized body, together with a shorter tail and narrower soles (Fernandez *et*

al. 2010), should enable the cursorial *P. melanotis* to occupy such zones as M2 and M3,

where predators cannot easily spot it through a dense vegetation cover, so it can scape

very quickly. Indeed, high shrubs cover provide both protection from predators and

food sources, since seeds may be concentrated under shrub canopies (Thompson 1982;

Mohammadi 2010). In fact, rodents usually avoid foraging in unsheltered microhabitats

and forest edges where they are more likely to be spotted by avian (Kotler *et al*. 1991)

and other vertebrate predators (Morris and Davidson 2000; Mohammadi 2010).

Finally, we want to highlight and recommend our sampling of environmental

features and quantitative analyses, especially for Ripley's bivariate K, as neither one has

been extensively used in studies focusing on the spatial dynamics of dispersion in small

rodents, despite its important advantages. The methodological approach used here,

together with the gathered evidence became relevant for providing us with necessary

information about the ability of this two *Peromyscus* to split resources in a mid-latitude

temperate forest at the edge of a megalopolis, which continuous growth produces

fragmentation and patching of natural microhabitats. Therefore, our local scope was

explicitly designed at this initial point, to focus on small-sized rodents with low vagility.

Besides, we successfully aimed to reconstruct vegetation structure, to provide us with

clues as to how to eventually manage such perturbed ecosystems for conservation

purposes. We also think that our methodology can be helpful in other scenarios for

understanding small rodents, such as ethological dynamics, activity patterns, and

reproductive systems at either or both intra or interpopulation levels.

590   **Acknowledgments**

We thank Juan Luis Patiño Ortega and Claudia Peralta Juarez for their fieldwork

assistance and friendly logistic support. The staff at UAMI mammal scientific collection

kindly provided all the facilities for specimen curation and housing. Authorities in

charge of DLNP (CORENA, SEMARNAT), as well as the forest rangers and

administration staff at the National Park, buoyed us with information, security, and

logistics to complete our studies. This paper contains partial information from the thesis

of Master in Biology (DCBS, UAM-I) of IMDA; IMDA received financial support

through a fellowship (283799, CVU 479479) from the National Council of Science and

Technology (CONACyT). This study was supported by annual grants (143.***.46 IB-

DCBS-UAMI, 2013-2017) to AACC.

603   **References**

Álvarez-Castañeda, S.T. (2005). *Peromyscus melanotis*. Mamm. Spec. 764, 1-4.

Barry Jr, R.E., Heft, A.A., Baummer, TE. (1990). Spatial relationships of syntopic white-
footed mice, *Peromyscus leucopus*, deer mice, *Peromyscus maniculatus*, and red-
backed voles, *Clethrionomys gapperi*. Can Field-Nat. 104, 87-393.

Bellows, A.S., Pagels, J.F., Mitchell, J.C. (2001). Macrohabitat and microhabitat
affinities of small mammals in a fragmented landscape on the upper coastal plain of
Virginia. Amer. Midl. Nat. 146(2), 345-360.

Briani, D.C., Palma, A.R., Vieira, E.M., Henriques, R.P. (2004). Post-fire succession of
small mammals in the Cerrado of central Brazil. Biodivers. Conserv. 13(5), 1023-
1037.

Brown J.H, Zeng, Z. (1989). Comparative population ecology of eleven species of rodents
in the Chihuahuan Desert. Ecol. 70(5), 1507-1525.

Bowers, M.A. (1982). Foraging behavior of heteromyid rodents: field evidence of
resource partitioning. J. Mamm. 63(3), 361-367.

Bowers, J.H., Baker, RJ., Smith, M.H. (1973). Chromosomal, electrophoretic, and
breeding studies of selected populations of deer mice (*Peromyscus maniculatus*) and
Black-eared mice (*P. melanotis*). Evol. 27(3), 378-386.

Bowman, J., Forbes, G., Dilworth, T. (2001). Landscape context and small-mammal
abundance in a managed forest. For. Ecol. Manag. 140(2), 249-255.

Bowman, A.S., Facelli, J.M. (2013). Fallen logs as sources of patchiness in chenopod
shrublands of South Australia. J. Arid Environ. 97, 66-72.

Bowne, D.R., Peles, J.D., Barrett, GW. (1999). Effects of landscape spatial structure on
movement patterns of the hispid cotton rat (*Sigmodon hispidus*). Landsc. Ecol. 14(1),
53-65.

Canfield, R.H. (1941). Application of the line interception method in sampling range
vegetation. J. For. 39(4), 388-394.

Carey, A.B., Harrington, C.A. (2001). Small mammals in young forests: implications for
management for sustainability. For. Ecol. Manag. 154(1), 289-309.

Carleton, M.D. (1989). Systematic and evolution. Pp. 7-141 in Advances in the study of
Peromyscus (Rodentia) (G. L. Kirkland, Jr., and J. N. Layne, eds.). Texas Tech
University, Lubbock.

Castro-Campillo, A., León-Altamirano, L., Herrera-Muñoz, J., Salgado-Ugarte, I.,
Mendieta Márquez, E., Contreras-Montiel, JL., Serrano, H., Ramírez-Pulido, J.,
Salame-Méndez, A. (2012). Is there a difference between T contents in two
populations of the Black-eared Mouse, living under similar conditions but with
differences in two population patterns? Acta Zool. Mex. (n. s.) 28(3), 525-539.

Castro-Campillo, A., Salame-Méndez, A., Vergara-Huerta, A., Castillo, A., Ramírez-
Pulido, J. (2008). Fluctuaciones de micromamíferos terrestres en bosques templados
aledaños a la ciudad de México. Distrito Federal. Pp. 391-410 in Avances en el
Estudio de los Mamíferos de México (Lorenzo, C, Espinoza E, Ortega J, eds.).

Publicaciones especiales. Vol. II, Asociación Mexicana de Mastozoología, A. C.,
México, D. F. 726 pp.

Cramer, M.J., Willig, M.R. (2002). Habitat heterogeneity, habitat associations, and rodent
species diversity in a sand–shinnery-oak landscape. J. Mamm. 83(3), 743-753.

Coppeto, S.A., Kelt, D.A., Van Vuren, D.H., Wilson, J.A., Bigelow, S. (2006). Habitat
associations of small mammals at two spatial scales in the northern Sierra Nevada. J.
Mamm. 87, 402-413.

Currie, D.J. (1991). Energy and large-scale patterns of animal and plant species richness.
American Naturalist 137: 27-49.

Chesson, P. (2000). Mechanisms of maintenance of species diversity. Annu. Rev. Ecol.
Syst. 31, 343–366. doi: 10.1146/annurev.ecolsys.31.1.343

Chupp, AD. (2002). Habitat selection in four sympatric small mammal species and the
effects of potential predators on *Peromyscus leucopus*. Master of Science Thesis.
Department of Biology, Ohio University, 61 pp.

Dale, MRT. (1999). Spatial pattern analysis in plant ecology. Cambridge University
Press.

Dalmagro, A.D., Vieira, E.M. 2005. Patterns of habitat utilization of small rodents in an
area of Araucaria forest in Southern Brazil. Austral Ecol. 30, 353-362.

De la Cruz-Argüello, I.M. (2015). Patrones de microdistribución en los mamíferos
pequeños en vegetación secundaria en el Parque Nacional Desierto de los Leones.
Tesis de Maestría en Ciencias. División de Ciencias Biológicas y de la Salud,
Universidad Autónoma Metropolitana, Unidad Iztapalapa, pp. 256.

DeWalt, S.J., Maliakal, S.K., Denslow, J.S. (2003). Changes in vegetation structure and
composition along a tropical forest chronosequence: implications for wildlife. For.
Ecol. Manag. 182(1), 139-151.

Dooley, J.L., Dueser, R.D. (1996). Experimental tests of nest site competition in two
Peromyscus species. Oecologia 105, 81-86

Dooley, J.L., Bowers, M.A. (1998). Demographic responses to habitat fragmentation:
experimental tests at the landscape and patch scale. Ecol. 79(3), 969-980.

Fahrig, L., Merriam, G. (1994). Conservation of fragmented populations. Conserv. Biol.
8(1), 50-59.

Fernández, J.A., García-Campusano, F., Hafner, M.S. (2010). *Peromyscus difficilis*
(Rodentia: Cricetidae). Mamm. Spec. 42(1), 220-229.

Forsman, E.D., Meslow, E.C., Wight, H.M. (1984). Distribution and biology of the
spotted owl in Oregon, USA. Wildl. Monographs. 87, 1-64.

Galindo, C., Krebs, C.J. (1985). Habitat use and abundance of Deer Mice: interactions
with Meadow Voles and Red-backed Voles. Can. J. Zool. 63, 1870-1879.

Gunther, P.M., Horn, B.S., Babb, G.D. (1983). Small mammal populations and food
selection in relation to timber harvest practices in the western Cascade Mountains.
Northwest Sci. 57, 32-44.

Grelle, C.E.V. (2003). Forest structure and vertical stratification of small mammals in a
secondary Atlantic forest, southeastern Brazil. Stud. Neotrop. Fauna Environ. 38(2),
81-85.

Haim, A., Rozenfeld, F.M. (1993). Temporal segregation in coexisting *Acomys* species:
the role of odour. Physiol. Behav. 54(6), 1159-1161.

Hammer, Ø., Harper, D.A.T., Ryan, P.D. (2001). PAST: PAleontological STatistics
software package for education and data analysis. Palaeontol. Electronica 4, 1-9.

Johnson, M.L., Gaines, M.S. (1990). Evolution of dispersal: theoretical models and
empirical tests using birds and mammals. Annu. Rev. Ecol. Syst. 21(1), 449-480.

693 Johnson, M.D. (2007). Measuring habitat quality: a review. Condor 109(3), 89-504.

694 Jorgensen, E.E. (2004). Small mammals use of microhabitat reviewed. J. Mamm. 85(3),
695  531-539.

696 Kalcounis-Rüppell, M.C., Millar, J.S. (2002). Partitioning of space, food, and time by
697  syntopic *Peromyscus boylii* and *P. californicus*. J. Mamm. 83, 614-625.

698 Kaufman, D.W., Kaufman, G.A. (1989). Population biology. pp. 233-271 *in* Advances in
699  the study of *Peromyscus* (Rodentia) (G. Kirkland, JR. and J. Layne, eds.). Texas Tech
700  University Press, Lubbock.

701 Kotler, B.P., Ayal, Y., Subach, A. (1994). Effects of predatory risk and resource renewal
702  on the timing of foraging activity in a gerbil community. Oecologia 100(4), 391-396.

703 Kotler, B.P., Brown, J.S., Hasson, O. 1991. Factors affecting gerbil foraging behavior
704  and rates of owl predation. Ecol. 72, 2249-2260.

705 Lehtonen, J.T., Mustonen, O., Ramiarinjanahary, H., Niemelä, J., Rita, H. (2001). Habitat
706  use by endemic and introduced rodents along a gradient of forest disturbance in
707  Madagascar. Biodivers. Conserv. 10, 1185-1202.

708 Maser, C., Maser, Z. (1988). Interactions among squirrels, mycorrhizal fungi, and
709  coniferous forests in Oregon. Great Basin Nat. 48, 358-369.

710 M'Closkey, R.T., Fieldwick, B. (1975). Ecological separation of sympatric rodents
711  (*Peromyscus* and *Microtus*). J. Mamm. 56(1), 119-129.

712 M'Closkey, R.T., Lajoie, D.T. (1975). Determinants of local distribution and abundance
713  in White-Footed Mice. Ecol. 56(2), 467-472.

714 Mohammadi, S. (2010). Microhabitat selection by small mammals. Adv. Biol. Res. 4(5),
715  283-287.

716 Monamy, V., Fox, B.J. (1999). Habitat selection by female swamp rats (*Rattus lutreolus*)
717  drives asymmetric competition and coexistence with long-tailed mice (*Pseudomys*
718  *higginsi*). J. Mamm. 80, 232-242.

719 Morris, D.W. (1987). Ecological scale and habitat use. Ecol. 68(2), 362-369.

720 Morris, D.W. (1984a). Sexual differences in habitat use by small mammals: evolutionary
721  strategy or reproductive constraint? Oecologia 65(1), 51-57.

722 Morris, D.W. (1984b). Patterns and scale of habitat use in two temperate-zone, small
723  mammal faunas. Can. J. Zool. 62(8), 1540-1547.

724 Pianka, E.R. (1973). The structure of lizard communities. Annu. Rev. Ecol. Syst. 4(1),
725  53-74.

726 Pyare, S., Longland, W.S. (2001). Patterns of ectomycorrhizal fungi consumption by
727  small mammals in remnant old-growth forests of the Sierra Nevada. J. Mamm. 82,
728  681-689.

729 Posada, D. and Buckley, T.R. (2004). Model selection and model averaging in
730  phylogenetics: advantages of Akaike Information Criterion and Bayesian
731  Approaches Over Likelihood Ratio Tests, Systematic Biology 53(5), 793-808,

732 Ramírez-Pulido, J., Lira, I., Gaona, S., Müdespacher, C., Castro-Campillo, A. (1989).
733  Manejo y mantenimiento de colecciones mastozoológicas. Universidad Autónoma
734  Metropolitana, Unidad Iztapalapa, México, D.F.

735 Ripley, B.D. (1976). The second-order analysis of stationary point processes. J. Appl.
736  Probab. 13(2), 255-266.

737 Rosenberg, M.S., Anderson, C.D. (2011). PASSaGE: pattern analysis, spatial statistics,
738  and geographic exegesis. Version 2. Methods Ecol. Evol. 2(3), 229-232.

739 Schoener, T.W. (1974). Resource partitioning in ecological communities. Science
740  185(4145), 27-39.

Schreiber, S. and Killingback P.T. (2013). Spatial heterogeneity promotes coexistence of rock-paper-scissor metacommunities. Theoretical population biology. 86. 10.1016/j.tpb.2013.02.004.

Seagle, S.W. (1985). Patterns of small mammal microhabitat utilization in cedar glade and deciduous forest habitats. J. Mamm. 66(1), 22-35.

Schreiber, S. J., and Killingback, T. P. 2013. Spatial heterogeneity promotes coexistence of rock-paper-scissors metacommunities. Theor. Popul. Biol. 86, 1–11. doi: 10.1016/j.tpb.2013.02.004

Smartt, R.A. (1978). A comparison of ecological and morphological overlap in a *Peromyscus* community. Ecol., 59:21.

Thompson, S.D. (1982). Structure and species composition of desert heteromyid rodent species assemblages: effects of a simple habitat manipulation. Ecol. 63(5),1313-1321.

Valladares, F., Bastias, C.C., Godoy, O., Granda, E., Escudero, A. (2015). Species coexistence in a changing world. Front. Plant Sci. 6, art. 866.

Vander Wall, S.B., Thayer, T.C., Hogbe, S.B., Beck, M.J., Roth, J.K. (2001). Scatter-hoarding behavior of deer mice (*Peromyscus maniculatus*). West. N. Am. Nat. 61, 109-113.

Villanueva-Hernández, A.I., Delgado-Zamora, D.A., Heynes-Silerio, S.A., Ruacho-González, L., López-González, C. (2017). Habitat selection by rodents at the transition between the Sierra Madre Occidental and the Mexican Plateau, México. J. Mamm. 98(1), 293-301.

Ward, J.H. (1963). Hierarchical grouping to optimize an objective function. J. Am. Stat. Assoc. 58(301), 236-244.

Wiens, J.A. (2000). Ecological heterogeneity: an ontogeny of concepts and approaches. *In*: The Ecological Consequences of Environmental Heterogeneity (Stewart AJA, John EA, Hutchings MJ, eds). Blackwell, Oxford, Pp. 9-31.

Zar, JH. (1999). Biostatistical Analysis. Prentice Hall, Princeton.

Zavala-Hurtado, JA., Valverde, PL., Herrera-Fuentes, MC., Diáz-Solis, A. (2000). Influence of leaf-cutting ants (*Atta mexicana*) on performance and dispersion patterns of perennial desert shrubs in an inter-tropical region of Central Mexico. J. Arid Environ. 46, 93-102.

Zavaleta, E. S., Shaw, M. R., Chiariello, N. R., Mooney, H. A., and Field, C. B. (2003). Additive effects of simulated climate changes, elevated CO_2, and nitrogen deposition on grassland diversity. Proc. Nat. Acad. Sci. 100, 7650–7654. doi: 10.1073/pnas.0932734100

Zielinski, W.J., Spencer, W.D., Barrett, R.H. (1983). Relationship between food habits and activity patterns of Pine Martens. J. Mamm. 64, 387-396.

 Figure legends

Fig. 1. Sampling plot design. a) Total plot area was 2475 m²; each intersection (n = 120)

was an independent sampling station for capturing mice. An influence area of 2.5 m²

around each station was constructed for sampling 24 microhabitat environmental

variables. b) Close up to the influence area showing the eight transects used for drawing

environmental variables; note subdivisions into three subtransects.

Fig. 2. Map of the distribution of the microhabitats (M1 = red, M2 = green, M3 = blue)

in the study quadrant for dry (A) and rainfall (B). The bar graph below shows the change

in the number of sampling stations between dry and rainy seasons. ** $p = 0.001$

Fig. 3. Box-plots are showing the seasonal heterogeneity of the 23 variables sampled in

the study plot between dry and rainy seasons. Overall, vegetation variables show an

increase in the rainy season, while the organic matter (OM) decrease. Structural elements

as logs on the ground do not shot differentiation between seasons. Some variables were

not present in high quantities along the study plot. Therefore we were unable to build the

box-plot. The red cross represents the mean of each variable.

Fig. 4. Spatial analysis with Kernel density points showing capture frequencies for *P.*

difficilis at dry (a) and rainy (b) seasons, and for *P. melanotis* at dry (c) dry and rainy (d).

Dots depict actual capture points, while hot spots color indicates higher density.

Fig. 5. Correspondence analysis from contingency tables of frequency capture data for

each deermouse (*P. difficilis* and *P. melanotis*) in each microhabitat (M1-3), during the

dry (a) and rainy (b) seasons, respectively. Axis c1 and c2 indicate the coordinates

resulted from the ordination of each variable from the analysis. Figure c shows the plots

for the GLMs models, the response variable for each graph are the predicted values of the

model to that principal component (the model effect). All of the PCs fitted in the model

show a positive relationship, indicating that the probability of capture frequency of the

individuals of two mice, increases with the presence of that variables. Therefore, the

figure also shows in the "X" axis in parenthesis the variables with the higher loading

scores in the ordination of this PCs.

Fig. 6. Ripley's bivariate K Analysis for interactions between the two *Peromyscus*. (a)

Repulsion pattern between both species in all analyzed distances during the dry season.

(b) Attraction from 6 - 12 m, and repulsion at 5, 10, and 14 m during the rainy season.

Black solid line represent the Ripleys K bivariate pattern; red dashed lines represent the

95% CI; blue solid line shows the null hypothesis.

844 **Fig. 1**

Fig. 3

930
931 **Fig. 4**
932
933
934
935

979
980 **Fig. 5**
981
982

998
999

1005
1006
1007

1014
1015
1016

1028

Fig. 6

Table 1. Environmental variables used to measure the horizontal and vertical structure of microhabitat at each sampling station, showing its ecological interpretation (kind or resource type) for both species of *Peromyscus*.

Variable code	Predation		Resources				
	Diminished risk	Refuges	Breeding Space	Food	Temporal food and refuges	Permanent	Resting places
Vegetation Cover (cm)							
VC10	X			X	X		X
VC25	X		X	X	X		X
VC35	X		X	X	X		X
VC50	X		X	X	X		X
VC100	X		X	X	X		X
Frequency of Herbaceous Plants (cm)*							
FH11	X		X	X	X		X
FH12	X		X	X	X		X
FH13	X		X	X	X		X
FH14	X		X	X	X		X
FH15	X		X	X	X		X
Frequency of Woody Plants (cm)*							
FW21	X	X	X	X	X	X	X
FW22	X	X	X	X	X	X	X
FW23	X	X	X	X	X	X	X
FW24	X	X	X	X	X	X	X
FW25	X	X	X	X	X	X	X
Species Vegetation Richness							
VR	X	X	X	X	X	X	X
Total of Herbaceous Plants							
TH	X		X	X	X		X
Total of Woody Plants							
TW	X	X	X	X	X	X	X
Rocks							
RC	X	X	X			X	X
Organic Matter (litter, twigs, small trunks)							
OM				X	X		
Bare Soil							
BS					X		
Coverage of Logs (stumps, fallen trunks)							
CLO	X	X	X			X	X
Numbers of Burrows							
BW	X	X	X			X	X

1079

* = Frequencies of plant types (FH, FW) involve the same heights as in VC.

1081
1082
1083
1084

1085 **Table 2.** Normality (**NOR**) tests for 22 variables (**VAR**) of microhabitat, and

1086 comparisons[abc] (**COM**) for average values between the dry (**DRY**) and rainy (**RAINS**)

1087 seasons, respectively. Variable names in Table 1.

1088

VAR	NOR[·]		COM	VAR	NOR[·]		COM	VAR	NOR[·]		COM
	DRY	RAINS			DRY	RAINS			DRY	RAINS	
BS	0.0001	0.0001	*1.0*[·]	**VC100**	0.002	0.003	0.0001[·]	**TW**	*0.143*	*0.08*	0.0004[b]
CLO	0.0001	0.0001	*0.972*[·]	**VR**	0.006	*0.082*	0.0001[b]	**FW21**	0.0001	0.0001	0.004[·]
OM	0.010	0.034	0.0001[·]	**TH**	0.0001	*0.072*	0.0001[b]	**FW22**	0.0001	0.0001	0.026[·]
RC	0.0001	0.0001	*0.432*[·]	**FH11**	0.015	0.028	*0.658*[·]	**FW23**	0.0001	0.0001	*0.875*[·]
VC10	0.002	0.041	0.0001[·]	**FH12**	0.0001	0.0001	0.001[·]	**FW24**	0.0001	0.0001	*0.378*[·]
VC25	0.009	0.003	0.0001[·]	**FH13**	0.0001	0.0001	0.0001[·]	**FW25**	0.0001	0.001	0.0001[·]
VC35	0.0001	0.002	0.0001[·]	**FH14**	0.0001	0.0001	0.0001[·]	BW	0.0001	0.0001	1
VC50	0.0001	0.019	0.0001[·]	**FH15**	0.0001	0.0001	0.0001[·]				

1089

1090 Parametric: [·]Kolmogorov-Smirnov; [b]Student for two samples; non-parametric: [·]Wilcoxon). The *p*-values in italics do

1091 not reject normality (*p* = 0.05), or do not indicate significant differences between pluvial seasons.

1092

1093

1094

1095

1096

1097

1098

1099

1100

1101

1102

1103

1104

1105

1106

1107 **Table 3.** Nearest Neighbor Analysis for both species of *Peromyscus*, it shows the mean

1108 distance in which individuals from both *Peromyscus* are segregated. R-value indicates

1109 how the observed distribution deviates from random. Clustered points give $R<1$, Poisson

1110 patterns give $R\sim1$, while overdispersed points give $R>1$.

1111
1112

1113

1114

1115

1116

1117

	Dry season		Rain season	
	P. difficilis	*P. melanotis*	*P. difficilis*	*P. melanotis*
Points	57	39	58	52
Mean distance	1.40	2.39	1.67	2.09
R	0.42	0.77	0.55	0.64
P value	1.1263×10^{-16}	0.006	8.6577×10^{-11}	8.7708×10^{-7}
Point pattern	Clustering	Clustering	Clustering	Clustering

1118

1119

1120

1121

1122

1123

1124

1125

1126

1127

1128

1129

1130

1131

1132 **Table 4.** Principal components used as effect model in the GLMs analysis for both

1133 seasons. Red colors represent the variables that have the major eigenfactors to the

1134 ordination of the components.

1135

Variables	Dry season			Rainy season		
	PC1	PC2	PC3	PC1	PC2	PC3
VC10	0.713	-0.388	-0.134	0.659	-0.179	-0.374
RC	-0.145	-0.055	-0.110	0.014	0.062	0.034
CLO	0.030	0.118	0.828	-0.153	0.411	-0.583
OM	-0.604	0.303	-0.487	-0.499	-0.122	0.775
BS	0.130	-0.187	0.112	-0.028	0.154	-0.328
VC25	0.415	0.280	0.034	0.658	-0.061	0.237
VC32	0.287	0.184	0.052	0.580	-0.003	-0.126
VC50	0.523	0.107	0.078	0.669	-0.173	-0.030
VC100	0.335	0.408	0.052	0.656	-0.053	0.107
VR	0.762	0.251	-0.269	0.776	0.380	0.058
TH	0.837	-0.430	-0.131	0.874	-0.319	-0.076
TW	0.520	0.684	-0.168	0.615	0.711	0.270
FH11	0.784	-0.394	-0.214	0.678	0.007	-0.228
FH12	0.573	-0.345	0.176	0.548	-0.278	0.256
FH13	0.247	0.073	0.113	0.463	-0.260	0.067
FH14	0.149	-0.186	-0.023	0.580	-0.530	-0.043
FH14	0.080	0.117	-0.236	0.481	-0.267	-0.019
FW21	0.127	0.510	-0.394	0.342	0.603	0.311
FW22	0.015	0.064	-0.105	0.266	0.212	0.416
FW23	0.199	0.256	-0.228	0.196	0.366	-0.241
FW24	0.424	0.359	0.195	0.337	0.165	0.159
FW25	0.515	0.366	0.380	0.494	0.469	0.026
BW	-0.105	0.249	0.557	-0.170	0.332	-0.218

1148

1149

1150

1151

1152

1153

1154

1155

Table 5. Generalized Linear Models constructed with the Principal Components and the

frequency of P. *difficils* and *P. melanotis* for both seasons. The partition of resources by

both species is observed at both seasons of the year. Different main components within

each season explain the frequency of capture of each of the species. Also, the table is

related to what type of microhabitat the new variables belong to (M1-M3). The goodness

of fit tests are shown; Pearson and Deviance.

1163

Term	Estimate	Std Error	L-R x^2	P value	Pearson	Deviance
				P. difficilis		
				Dry season		
Intercept	-0.97	0.15	55.73	0.0001	$x^2 = 155.80\ (p = 0.0096)$	$x^2 = 123.56\ (p = 0.3210)$
PC1 = M2	0.11	0.05	5.53	0.0186	AICc = 212.55	
PC3 = M1	0.45	0.06	44.73	0.0001		
				Rainy season		
Term	Estimate	Std Error	L-R x^2	P value	Pearson	Deviance
Intercept	-0.40	0.11	14.37	0.0002	$x^2 = 168.56\ (p = 0.0008)$	$x^2 = 156.43\ (p = 0.0062)$
PC2 = M3	0.23	0.06	10.92	0.0009	AICc = 286.76	
PC3 = M1	0.26	0.07	10.95	0.0009		
				P. melanotis		
				Dry season		
Term	Estimate	Std Error	L-R x^2	P value	Pearson	Deviance
Intercept	-1.21	0.18	74.28	0.0001	$x^2 = 237.65\ (p = 0.0001)$	$x^2 = 138.70\ (p = 0.0935)$
PC2 = M3	0.46	0.09	23.34	0.0001	AICc = 204.47	
				Rainy season		
Term	Estimate	Std Error	L-R x^2	P value	Pearson	Deviance
Intercept	0.60	0.12	28.15	0.0001	$x^2 = 132.71\ (p = 0.13)$	$x^2 = 132.72\ (p = 0.13)$
PC1 = M2	0.10	0.04	4.47	0.0343	AICc = 241.01	

1164

1165

1166

1167

1168

1169