Assessing the Relation between the Spatiotemporal Microhabitat Heterogeneity

and the Dispersion Patterns of Two Syntopic Small Mammals in a Midlatitude

Temperate Forest

De la Cruz-Arguello Ivan M.[a], Castro-Campillo Alondra[a], Zavala-Hurtado Alejandro[a],

Salame-Méndez Arturo[b], and Ramírez-Pulido José[a]

[a] Department of Biology (DB) and [b] Department of Reproduction Biology (DBR),

Biological and Health Sciences Division (DCBS), Metropolitan Autonomous University-

Iztapalapa (UAM-I). AP 55-535- Av. San Rafael Atlixco 186, Vicentina, 09340

Iztapalapa, CDMX, México.

E-mail of corresponding author: Alondra Castro-Campillo, acc@xanum.uam.mx

Laboratory of Mammals

Edificio "Anexo S" AS-012 PB

Biology, DCBS, UAM-I, AP 55-535

Av. San Rafael Atlixco 186, Vicentina, Iztapalapa

PC 09340, CDMX, México

Abstract

One of the basics problems in ecology is to understand the factors that shape the spatial patterns in the distribution of the species and the coexistence of close relatives species. Among the most critical factors governing the distributions and the coexistence of species are the spatiotemporal changes occurring in the microhabitat heterogeneity. Here, we assessed the heterogeneity of microhabitats and how they affect the spatial segregation of two species of small mammals (*i. e.*, *Peromyscus difficilis* and *P. melanotis*), which coexist in a temperate, mixed forest. We evaluated the microhabitat heterogeneity through multivariate statistics, using onto 23 habitat variables describing horizontal-vertical habitat structure along pluvial seasons. To detect specific microdistribution changes and habitat preferences by two species of small mammals, we used second-order spatial statistics and general linear models. According to their respective morphology and locomotive adaptations, the middle sized, midscansorial *P. difficilis* was resident all year long and preferred microhabitats with a high log ground cover, while the opportunistic, small sized, cursorial *P. melanotis* changed its occupancy area, depending on the density of herbaceous and woody plants cover. Under the more benign microhabitat conditions of the rainy season (denser plant coverage, milder temperature), both species showed closer microdistribution patterns; while these became repulsive at the less favorable conditions of the dry season (scarcer plant cover, colder temperature). Thus, we could confirm that seasonal changes of microhabitat heterogeneity promoted *Peromyscus* coexistence, through dispersion patterns reflecting partition of microhabitat resources.

Key words: Microhabitat selection, Use of space, Small mammals, Dispersion, Resource partition.

Introduction

51
52

Spatiotemporal habitat heterogeneity has a strong impact on coexistence among species

since they become spatially segregated, according to niche preferences such as resource

requirements (Valladares *et al.* 2015). Model simulations reveal the potentially important

role of spatial heterogeneity and its complex and delicate interplay with dispersal in

mediating long-term outcomes of species coexistence (Schreiber and Killingback 2013;

Valladares *et al.* 2015). For example, an increase in the number of habitat types,

resources, and structural complexity should increase the available niche space and thus

allow more species to coexist (Currie 1991). Equally crucial for the maintenance of

coexistence is the heterogeneity in time, with an influence on natural communities also

variable depending upon the temporal scales (Valladares *et al.* 2015). Temporal

fluctuations on habitat structure can stabilize coexistence via storage effect (Chesson

2000), when inter-annual variation in climate or resource availability favors alternatively

one group of species over the others (Zavaleta *et al.* 2003). For instance, *Peromyscus*

mice have been shown to respond to spatiotemporal habitat heterogeneity regarding

species coexistence (M'Closkey and Fieldwick 1975; M'Closkey and Lajoie 1975), spatial

segregation (Monamy and Fox 1999), and competition (Morris 1984; Seagle 1985).

Heterogeneous habitats contain patches of varying size, distribution, resources,

environmental conditions, and species composition (Fahrig and Merriam 1994) that vary

spatiotemporally, depending on observation scale and habitat type under study (Wiens

2000). It is highly likely that small size species such as small mammals perceive spatial

habitat heterogeneity at a fine scale (microhabitat). Therefore, both movement and

foraging by these kinds of mammals are affected by habitat heterogeneity (Bowne *et al.*

1999), having different responses among species or demographic groups within them

(Dooley and Bowers 1996). Here we focus on whether small mammals are capable of

perceiving and to respond to spatiotemporal microhabitat heterogeneity.

Small mammals use resources selectively, based on their requirements for growth,

survival, and reproduction, (Johnson and Gaines 1980). Several studies have quantified

variation in resource use in heterogeneous environments either among congeneric and co-

family mice species within a community (Kaufman and Kaufman 1989 and references

therein). Indeed, syntopic related species are especially valuable for studies of spatial and

resource partitioning, since they are most likely to be current or past competitors

(Kalcounis-Rüppell and Millar 2002). Since habitat use varies spatiotemporally (Haim

and Rozenfeld 1993) due to distribution and availability of resources, we expect that such

variation also affects the use of resources and relationships among non-closely related

species. In this study, the primary objective is to analyze how heterogeneity of

microhabitats is structured and change through pluvial seasons in a temperate, mixed

forest, and how such changes affect the coexistence of two syntopic species of

Peromyscus (*i. e., P. difficilis* and *P. melanotis*). Therefore, 1) We assess the presence of

the spatiotemporal heterogeneity at a fine scale (microhabitat) according to vertical and

horizontal structure indicators. 2) We evaluate the ecological dispersion patterns of two

species of *Peromyscus*, and is tested if the use of space of the two small mammals is

affected by the spatiotemporal changes in microhabitat heterogeneity. 3) We determine

what kind of spatial relationships occur between both species, depending on seasonal

changes in microhabitats heterogeneity (*i. e*., attraction or positive: sharing of same

microhabitats and resources; repulsion or negative: not sharing). Finally, we assessed

which specific structural elements of each one of the microhabitats better explain the

capture frequency of the two *Peromyscus* species between seasons.

 Material and methods

Ethics statement

Animal capture and handling followed guidelines of the American Society of

Mammalogists (Sikes *et al.* 2011). If animals died, corpses were prepared as voucher

specimens to house them at the Mammal Collection of the Universidad Autónoma

Metropolitana-Iztapalapa (UAMI, Ramírez-Pulido *et al*. 1989). Collecting permit,

SEMARNAT-08-049-B, was issued by DGVS, SGPA-09712/13, SEMARNAT, to

Alondra Castro-Campillo (ACC).

Study area

The study area was located in a temperate forest of conifers and broad-leaved trees at

Parque Nacional Desierto de los Leones (PNDL, CONANP 2006), Mexico City, within

the Transmexican Neovolcanic Range. Temperature averages 12.6 ± 6 ℃ from April to

July, while it descends to 8.1 ± 2 °C from December to February. Rainfalls occur from

late spring throughout summer (May to August), with average precipitation of 235 ± 30

mm, though this period can go until mid-fall (October). From winter to early spring

(December to March), precipitation average becomes 12 ± 4 mm, introducing a dry and

cold climate (PNDL, CONANP 2006). Dominating vegetation by strata includes canopy,

Abies religiosa, Pinus hartwegii, P. leiophylla, P. montezumae, Prunus serotina, Garrya

laurifolia, Salix paradoxa, and Buddleia cordata; understory shrubs, Baccharis conferta

and *Senecio spp.,* and herbs*, Acaena elongata, Sigesbeckia jorullensis,* and *Alchemilla*

procumbens; ground level, a rich variety of mosses and fungi.

Mice sampling

We captured mice along ten months (October 2013- August 2014), including the dry and wet seasons within a 2,475 m2 surface plot (Fig. 1) at 2289 m. The plot was gridded (12 vertical lines, A-L; 10 horizontal lines, 1-10) marking intersections each 5 m with buried wood stakes (150x2.5x2.5 cm) to construct a coordinate system for independent sampling stations, where we set 120 Sherman live traps (Tallahassee, FL 32303, USA), baited with oat flakes and vanilla scent (Fig. 1). Traps were set for two continues nights (2400 trap-nights), shortly before dusk, left open overnight, and checked at next day dawn. We recorded the frequency of captures of each of the two *Peromyscus* in the 120 sampling stations during all trapping sessions. All the traps were cleaned, and bedding was changed at each trap check event (Brouard 2015). All the traps were set randomly in each sampling station to avoid bias in frequency quantification due to recurrence behavior or shyness of the mice to the trapping. Coordinates of all trapped mice were recorded and the species and sex determined through visual inspection.

Microhabitat features

To assess spatiotemporal heterogeneity of microhabitat within the entire plot, we delimited an influence zone (sub-plot, Fig. 1; close-up, Fig. 1), drawing a rectangle of 2.5 m^2 around each sampling station. 23 variables were sampled (Table 1) once in each influence zone in both seasons. The sampling of the entire plot took ten days, we sampled one transect per day, starting at 9 am in the second week of the most represented month

of rains (July) and in February for the dry season (PNDL, CONANP 2006). These

variables have been proved to be components of the vertical and horizontal structure of

the microhabitat. Also as factors affecting the distribution of small mammals at fine scales

(Morris 1984, 1987; Stapp 1997; Jorgensen 2004, Coppeto *et al.* 2006; Villanueva-

Hernández *et al.* 2017). Environmental variables include vegetation coverage at different

heights (VC10-100 cm), number of herbaceous forms (FH11-15), and number of woody

forms (VF21-25) at five different heights: 10, 25, 35, 50, and 100 cm, respectively. Rock

coverage (RC), logs coverage which include stumps and fallen trunks (CLO); coverage

of organic matter (OM); coverage of bare soil (BS); vegetation species richness (VR);

total herbaceous plants (TH); total woody plants (TW) and the number of burrows (BW)

also were sampled (Table 1). Overall, these variables provide information about possible

refuges for the small mammals against predators, some elements of microhabitat also

imply spaces for resting, breeding, for food resources and safe roads to move from one

place to another, avoiding predation risk (Jorgensen 2004). Detailed information about

ecological meanings for each variable appears in Table 1.

We adapted Canfield's (1941) method of Line Intercept (LI) to measure all

variables in each pluvial station, for it allows sampling within-plot variation and

quantifies changes, both in plant species cover and height over time, by using transects

located within a plot. We drew eight graduated transects (cm) from the respective

influence zone of each station to cover the 2.5 m^2 sub-plots (Fig. 1); four 2.5 m transects

were cardinally oriented (N, S, E, W), while other four 3.30 m ones were diagonally

oriented (NE, NW, SE, SW). Only plants intercepted by diagonal transects were counted

for vegetation species richness, number of life forms at different heights (herbaceous and

woody plants), and total herbaceous and woody plants; while all transects were used to

measure plant coverage at different heights, as well as coverage of fixed elements (*i. e.,*

rocks, logs, bare soil, dead organic matter). We calculated coverage percentage of each

variable within the sub-plots with the formula $\sum(Xi)\frac{100}{2400}$, where Xi is centimeters

occupied by every plant in each transect, and 2400 equals total longitude (sum of the eight

transects). We counted the number of woody and herbaceous plant life forms at different

heights and burrows within each influence zone of every sampling station. We took each

hole in the ground ca. 8-12 cm of diameter as a potential burrow for both *Peromyscus*

(Álvarez-Castañeda 2005; Fernandez *et al.* 2010).

Statistical data analysis

We standardized all variables and run statistical tests at $p \leq 0.05$. We did not found

multicollinearity among the 23 variables which was assessed by examining the Variance

Inflation Factor (*i. e.,* all values were in a threshold <6) (Zar 1999). To identify and

categorized microhabitat heterogeneity within the grid, we performed two independent

Hierarchical Cluster Analysis (HCA) in JMP® (ver. 9, SAS Institute Inc., Cary, NC,

1989-2007) for each season. These analyses allowed us to cluster the sample stations with

similar characteristics according to with the sampled variables for both seasons. We used

Ward´s method (1963) where the distance between two clusters is the ANOVA' sum of

squares between them, added up over all variables. Visual inspection of the dendrogram

and no drastic changes in the variance screeplot, gave us three general groups of stations

for both seasons, which we interpreted as distinct microhabitat types (M1-3). We further

statistically validated such microhabitats carrying out independent Discriminant Analyses

for both seasons (DA, Addinsoft SARL's XLSTAT 2013; F = 9.99 dry; F = 9.64, rains;

Wilk's Lambda = 0.0001 in both) (Fig. A.1). These analyses allow us to know what type

of microhabitat corresponded to each sampling station. Thus, we constructed two

different maps, assigning to each sampling station a color mark that represented the three

different microhabitats for both seasons (M1: red, M2: green, M3: blue) (Fig. 2, Fig. A.1).

To create typologies that describe at a fine scale the behavior of each one of the 23

variables within three microhabitats for both seasons, we built a plot whereas "Y" axis

comprises of the 23 standardized variables and in the "X" axis corresponds to the three

microhabitats constructed with the same 23 variables (Fig. A.1).

Due to not all 23 variables did not reveal a normal distribution, We applied two

different tests to analyze the distribution changes on each microhabitat variable between

the rainy and dry seasons. First, distribution normality was checked of every variable with

Kolmogorov-Smirnov tests (Zar 1999). Thus, We used Student's t-parametric tests for

variables normally distributed and non-parametric Wilcoxon test for non-normal

variables (Table 2). To know whether the number of stations structuring each type of

microhabitat changed between seasons (seasonal microhabitat heterogeneity), we used a

contingency table and x^2 test. All these analyses were performed with XLSTAT 13

(Addinsoft SARL). To examine the association between changes in spatial heterogeneity

of microhabitats and dispersion patterns of *Peromyscus*, along with the entire grid, we

conducted a Contingency Tables Analysis. Then, for visualizing associations between the

Peromyscus and microhabitats, we performed a Correspondence Analysis with the

constructed Contingency Tables for both seasons, including the abundance of each

species in the three microhabitats (Table A.2).

Spatial analysis

To map variations in point density captures of small mammals, and to find density gradients across the plot area, we used the Kernel function (PAST, ver. 3.14, Hammer *et al.* 2001). To test the ecological dispersion patterns of both species within the plot (*i. e.,* clustering or overdispersion) for both seasons, we used a "Nearest Neighbor Analysis" (Clark and Evans, 1954) using the PAST software (ver. 3.14, Hammer *et al.* 2001). We applied the Wrap-around edge effect adjustment which is only appropriate for rectangular domains as our plot sampling (Hammer *et al.* 2001). In general, the "Nearest Neighbor Analysis" compares the mean distance of each main individual from its nearest neighbor of the same species with the mean distance expected for a set of points randomly dispersed at the same density (Vázquez and Álvarez-Castañeda 2011). The ratio of the observed mean distance to the expected distance (*R*) indicates how the observed distribution deviates from random. Clustered points give *R<1*, Poisson patterns give *R~1*, while overdispersed points give *R>1*.

To assess whether both species were associated or disassociated, we compared dispersion patterns between the two mice, using Ripley's K bivariate function (Ripley 1977), since the method considers all distances among individuals located under a Cartesian scheme (X,Y) in a quadrat plot (Ripley 1976, Dale 1999, Zavala-Hurtado *et al.* 2000). We used PASSaGE (*ver. 2,* Rosenberg and Anderson 2011) to carry out Ripley's bivariate K analysis. We used the option to test the associations conditional on current locations hypothesis in PASSaGE. In this test, the point locations remain fixed, and only the types are randomized (the two *Peromyscus*). The number of each type remains fixed, but the association of each type with a specific, fixed location is randomized. In this case, one is not testing whether the points are themselves random or clustered, but rather

whether the association of type A with type A (*P. difficilis vs. P. diffiiclis*) (or *P. difficilis*

vs. P. melanotis) is what one would expect given the locations of the points as fixed

(Rosenberg and Anderson 2011). We assigned coordinates to each sampling stations

considering 5 m distance between them. Therefore, because each sampling station had a

Cartesian coordinate system, we used as data input each station were we capture mice.

Thereby, our null hypothesis involved independence between our two species *P. difficilis*

and *P. melanotis* (Ripley 1976, Diggle 1983, Dale 1999), since we were interested in

whether points of a specific type (*P. difficilis*) were associated, or disassociated, with

points of the other specific type (*P. melanotis*). In other words, Ripley's $K_{12}(d)$ allowed

us to assess spatial attraction or repulsion between the two species, among the stations of

the plot. To evaluate the statistical significance of $K_{12}(d)$, we estimated 95% confidence

envelops (95% CI), using a Monte Carlo procedure, based on 1000 stochastic relocation

simulations of the sampling stations in the plot (Upton and Fingleton 1985; Bailey and

Gatrell 1995; Zavala-Hurtado *et al.* 2000). When $L(d)$ was positive and took values over

the upper limit of the confidence region, we inferred dissociation or repulsion between

the *Peromyscus* at the corresponding (d) scale; whereas a significant negative deviation

indicated us a pattern of association or attraction between the two mice (Dale 1999). If

$L(d)$ remains within the limits of the 95% CI for a given value of d, the null hypothesis

of independence between the two contrasts cannot be rejected (Dale 1999). The height of

the $L(d)$ function (peak height) indicates the intensity of the association or repulsion.

Control for edge effect was carried out for analysis by rescaling the counts based overlap

with the study boundary setting the options area/volume which has been recommended

for quadrats plots (Rosenberg and Anderson 2011).

Prediction of microhabitat elements affecting the use of space by each species

To determine which specific environmental variables best explained the frequency of

both species at each season, we performed Generalized Linear Models (GLMs), using

JMP® (ver. 9, SAS Institute Inc., 1989-2007, Cary, NC,). First, we conducted a Principal

Component Analysis (PCA) onto the means of the 23 variables to obtain functions

summarizing the most significant variance at each season. Because in both seasons the

three first functions summarized a good portion of the variance (38.61 for the dry season

and 45.92 for rains), we used these functions to construct the GLMs. Therefore, we

assessed different models using the PCs as effect variables to explain the frequency of

the individuals of both species for both seasons. Then, to visualize the fit of the model,

the predicted values from the GLMs were plotting against the Principal Components of

the best model chosen. For each species, the response variable in the models was the

capture frequency at each trapping station of the grid, and we assumed a Poisson

distribution with a log link function. Criteria for the selecting models that most explained

the frequency of mice for each species at each season in the sampling stations included:

model significance of $p < 0.05$; lowest Akaike Information Criterion (AIC), *i. e.,* a

measure of variance explained by the model (Posada and Buckley 2004); Pearson

goodness-of-fit ($p < 0.05$) and its deviation (Deviance; $p < 0.05$); as well as the biological

and ecological meaning of results.

299   **Results**

The Behavior of 23 variables of microhabitat between seasons

When considering the mean value over all stations, 16 out of the 23 environmental

variables showed a significant difference between the two seasons (Table 2, Fig. 3, Fig.

A.1). The main changes were observed for variables related to herbaceous vegetation,

according to plant coverage at different heights, and vegetation species richness (VR). As

expected in overall, woody life forms were more stable between seasons (Table 2, Fi. 3,

Fig. A.1). Components of microhabitat showing no statistical change between seasons

were frequency of 10 cm height herbaceous plants (FH11), the frequency of both 35 cm

(FW23) and 50 cm (FW24) woody plants, as well as coverage of logs (CLO), rocks (RC),

and bare soil (BS). Therefore, space configuration given by these structural features

remained stable in both seasons (Fig. 3).

Microhabitat heterogeneity

The dendrogram from cluster analysis revealed three different classes of microhabitats

(M1, M2, and M3) within the sampling plot in each season (Fig. A.1). M1 included more

sampling stations during the dry season, while M2 covered the majority of sampling

stations during the rainy season (Fig. 2, Fig. A.1, Table A.1). Indeed, Fisher's exact test

of Contingency Tables Analysis revealed significant changes in the number of sampling

stations ($p = 0.004$) from dry to rainy seasons. The main changes occurred between M1

($x^2 = 2.123$, $p < 0.05$) and M2 ($x^2 = 3.348$, $p < 0.05$), especially in the latter, since it

augmented 24 sampling stations from the dry season to the rainy season (Fig. 2, Table

A.1). Conversely, the number of sampling stations remained quite stable in M3 ($x^2 =$

0.011) throughout the study (Fig. 2, Fig. A.1, Table A.1).

Discriminant analyses validated heterogeneity of the three microhabitats (Fig. 2)

within and between pluvial seasons. the percentage of variance associated to each

discriminant function was clearer during the rainy season (dry season: F1 = 58.67 %, F2

= 42.32 %; rainy season: F1 = 72.581 %, F2 = 27.419 %) (Fig. 2, Fig. A.1). Discriminant

functions explained variation due to woody plants in the understory in the dry season,

while herbaceous vegetation together with decayed matter and logs became more relevant

at rains (Table A.3). Wilk's Lambda showed that at least the mean of one microhabitat

differed statistically from the others either in the dry and rainy season (p = 0.0001; p =

0.0001, respectively) (Fig. A.1). These results corroborated spatial and temporal

heterogeneity drawn from cluster analysis of individual station features in the grid plot.

During the dry season, 93.33 % of individual sampling stations remained correctly

classified, and 94.17 % of them during the rainy season (Fig. 2).

Description of microhabitats within the plot

Microhabitat 1. M1 (Fig. 2, Fig. A.1) was characterized by low amounts, or absence, of

vegetation cover at different heights (VC) and by a low frequency of herbaceous and

woody plants (FH and FW, respectively). In contrast, organic matter on the ground (OM)

was the most frequent component. This microhabitat was also distinctive for having

extensive coverage of logs (CLO) on the ground and for being the microhabitat with the

significant presence of burrows (BW).

Microhabitat 2. M2 (Fig. 2, Fig. A.1) was the largest area covered by herbaceous life

forms at different heights; vegetation species richness (VR) and the total number of

herbaceous plants (TH) also remained very high. Woody life forms at 25 cm (FW22)

were present but at low frequencies, while FW24 and FW25 remained at higher

frequencies. Coverage on the ground of small logs or dead wood (CLO), of rocks (RC),

and organic matter (OM) were shallow during the study. Bare soil surface (BS) increased

to higher amounts during the dry season, while it decreased during the rainy season. There

were no burrows (BW) in this microhabitat.

Microhabitat 3. In M3 (Fig. 2, Fig. A.1), VC25, VC35, VC50, and VC100 showed high

values, while vegetation species richness (VR) averaged the highest compared to other

microhabitats. The total number of woody plants (TW) and frequency of woody plants at

different heights (FW21-25) remained very high, while the total number of herbaceous

plants (TH) was low. Vegetation coverage at 10 cm (VC10), rocks (RC), logs (CLO), and

bare soil surface (BS) were low during the dry season; however, VC10 increased in the

rainy season. Conversely, OM was present in the dry season but decreased in the rainy

season. The number of burrows (BW) remained quite scarce.

Association between the microhabitats and species

Kernel Graphs (Fig. 4) indicated that the two species used space distinctivel, since they

were differentially scattered along the plot. *Peromyscus difficilis* was more abundant

during the dry season, occupying a large portion of the plot. Instead, *P. melanotis*

increased its distribution during the rains when it seemed to displace *P. difficilis* into other

sampling stations (Fig. 4). Distribution rearrangements between seasons implied that one

species occupied some stations more frequently than the other, and vice versa (Fig. 4).

Indeed, the two x^2 tests yielded statistical abundance differences in each species at both

microhabitats and seasons (Table A.2: dry season; R^2 (U) = 0.09, n = 111, df = 2,

Likelihood Ratio $x^2 = 15.07$, $p = 0.00005$; rainy season; R^2 (U) = 0.02, $n = 168$, df = 2,

Likelihood Ratio $x^2 = 6.40$, $p = 0.0406$).

During the dry season (Fig. 5 and Table A.2), *Peromyscus difficilis* was highly

related to M2 (80 % captured mice) and frequently related to M1 (64 % captured mice);

conversely, *P. melanotis* was captured more frequently in M3 (66 % captured mice).

Spatial use of microhabitats changed for both species in the rains (Fig. 5 and Table A.2),

showing a microhabitat partition again, though less visible: in the Correspondence

Analysis, 61 % of mice caught in M1 were *P. difficilis*. This species was also related to

M3, while 60% of mice caught of *P. melanotis* were captured in M2.

Spatial Patterns

The Nearest Neighbor Analysis revealed that in both seasons both species showed a

significant clustered pattern within the plot (Table 3). In the dry season, the mean distance

between individuals of *P. difficilis* displayed to be more clustered than the individuals of

P. melanotis. However, the mean distance between individuals of *P. difficilis* increases in

the rainy season, while individuals of *P. melanotis* showed a reduction in their area

occupied (Table 3).

On the other hand, during the dry season, Ripley's bivariate K showed a

statistically significant pattern of repulsion for both *Peromyscus* in almost all analyzed

distances of the entire plot, excused at 2 m where it shows an attraction pattern (Fig. 6a).

Such repulsion pattern turned over in almost all distances during the rainy season (Fig.

6b), since the two species became more associated, sharing microhabitats in almost all

capture stations. However, during the rainy season, significant statistical peaks of

repulsion reappeared between the species at distances of five, ten, and 14 m (Fig. 6b).

Analyses also revealed the intensity of these patterns; *e.g.,* the likelihood for finding

individuals of *P. difficilis* and *P. melanotis* together at the same capture station was very

low during the dry season (Fig. 6a), while this probability increased in the rainy season

(Fig. 6b).

Structural components of microhabitats affecting the distribution of small mammals

During the dry season, PC1 and PC3 predicted the frequency of *P. difficilis* according to

the best Generalized Linear Model for this season (AICc 212.55) (Table 5, Fig. 5c).

Environmental variables with higher scores in the PC1 ($p = 0.0186$) were the total number

of herbaceous plants (TH), vegetation species richness (VR), and vegetation cover at 10

cm height (VC10). In PC3 ($p = 0.0001$), characterizing variables were the number of logs

on the ground (CLO), as well as the presence of burrows (BW) (Table 4, 5, Fig. 5c). These

five variables were also the main elements characterizing M2 (PC1) and M3 (PC3), the

two habitats where the frequency of capture of *P. difficils* was the higher. As for *P.*

melanotis (Fig. 5c), the frequency of capture of *P. melanotis* was positively related to

PC2 (AICc = 204.47, $p = 0.0001$). Here, among the four variables with higher scores,

were the total number of woody vegetation (TW) and frequency of woody life forms

characterizing 10 cm height (FW21), the two main elements at M3 for the dry season

(Table 4, 5, Fig. 5c).

During the rainy season, a similar pattern of microhabitat partition occurred

between the two species. The best GLM model for *P. difficilis* in this season was (Table

4, 5, Fig. 5c) was mainly associated with the total number of woody plants (TW),

herbaceous forms at 50 cm (FH14), woody forms at 10 and 100 cm (FW21, FW25,

respectively) and with the number of logs on the ground (CLO), in PC2; and with were

number of logs on the ground (CLO), and organic matter (OM) in PC3 (AICc = 286.76,

$p = 0.0001$) (Table 4, 5). The latterly involved variables with a high presence at M1 (PC3)

and M3 (PC2) during the wet season, where *P. difficilis* was more frequently captured.

On the other hand, *P. melanotis* (Fig. 5c) was only associated to PC1 (AICc = 241.01, *p*

= 0.0343). The primary variables characterizing this component were the number of total

herbaceous plants (TH), vegetation species richness (VR), and the plant cover at almost

all analyzed heights (VC10-100 cm) (Table 4). Therefore, results of Contingency Tables

Analysis, Correspondence Analysis, and GLMs models showed that the two syntopic

species of *Peromyscus* switched and split resources in space, as environmental conditions

changed.

435   **Discussion**

Our findings on microhabitat relations of rodents in a patch of a mid-latitude temperate

forest, concur with the habitat heterogeneity coexistence hypothesis (Cramer and Willig

2002; Valladares *et al.* 2015). We found that both *Peromyscus* were sensitive to slight

changes in microhabitat structure and that the three microhabitats provided different

resources for each one. Our results also suggest that coexistence between these syntopic

Peromyscus is facilitated by temporal differences in space structure, and probably also by

changes on the availability of both food and shelter that go with it (Pianka 1973; Schoener

1974). For instance, *P. difficilis* was more related to M1 and M2 by the dry season, while

P. melanotis was highly related to M3. Conversely, *P. difficilis* was more related to M1

and slightly captured to M3 by the rainy season, while *P. melanotis* was highly related to

M2. Partition in space use has already been documented for coexisting species of

Peromyscus within a community; *i.e.,* extensive studies of niche partitioning by

Peromyscus leucopus and *P. maniculatus* (Kaufman and Kaufman 1989). Moreover,

several studies have demonstrated variation in spatial resource use by congeneric species

(Barry *et al.* 1990; Dooley and Dueser 1996; Kalcounis-Rüppell and Millar 2002).

In our study, *P. difficilis* remained related to M1 and M3 in both seasons,

especially to the former during the wet season. The number of fallen logs was very high

at M1, while understory complexity, together with woody vegetation, were more

representative in M3. Such microhabitats represent small patches of food for small

mammals, offering invertebrates (Bellows *et al.* 2001), flowers, grains, and fungi items

(Bowman *et al.* 2001). In these microhabitats, individuals of *P. difficilis* can also use

fallen logs as shelter, as well as pathways for quick and straight locomotion inside the

forest (Bellows *et al.* 2001; Dewalt *et al.* 2003). Indeed, fallen logs promote structural

complexity of forests and may enhance positive interaction among species of small

mammals (Bowman and Facelli 2013). GLM's analyses showed that abundance of *P.*

difficilis was explained by PC1 and PC3 during the dry season and by PC2-4 in the rainy

season. In the dry season, variables contributing most to PCA ordination in PC1 and PC3

were the total number of herbaceous vegetation (TH) and the number of logs on the

ground (CLO), as well as burrows (BW). While in the dry season, PC2-4 represented

woody vegetation (TW), number of logs, burrows, and understory complexity (VR).

Bellows *et al.* (2001) found a similar result in a high latitude temperate forest (Virginia,

USA), where distribution of a small mammal, generalist species was associated with the

diameter of fallen logs, the frequency of shrubs, and degree of canopy closure.

Association between rodents and fallen logs was also documented for *Nectomys*

squamipes that builds up its nests inside decomposed, fallen logs (Briani *et al.* 2004); this

has also been recorded for other rodent species in several biomes with different vegetation

types (*e. g*., *Rattus rattus*, *Nesomys audeberti* in Lehtonen *et al.* 2001; *Oligoryzomys*

nigripes in Dalmagro and Vieira 2005). In contrast, *P. melanotis* showed a close relation

with only one PC over each pluvial season. During the dry season, the frequency of this

species was only related to PC2, which represented the total number of woody vegetation

(TW). Then by the rainy season, the species became only related to PC1, which accounted

for understory complexity dominated by herbaceous life forms (TH). Relationships

between frequency of rodents and habitats with increased understory density has been

interpreted as protection against aerial predators (Dalmagro and Vieira 2005), or as

preference for more complex habitats that enable for vertical stratifications and, thus for

extended coexistence inside fragments (Grelle 2003; Johnson 2007).

Intrinsic factors as sex and mating system must also be related to the coexistence

of both *Peromyscus*. For instance, the Nearest Neighbor Analysis revealed that in both

seasons both species showed a significant clustered pattern within the plot (Table x).

However, the mean distance between individuals of *P. difficilis* displayed to be more

clustered than the individuals of *P. melanotis* especially during the dry season. Indeed,

the mean distance between individuals of *P. difficilis* increases in the rainy season, while

individuals of *P. melanotis* showed a reduction in their area occupied (Table x). This

patterns are related with the breeding season of both species. Our data (not published)

revealed that during the dry season were captured more individuals with signs of

reproductive activity (males: inguinal vs. scrotal testicles, females: alopecia, the presence

of milk, the appearance of the vagina, gestation). While for *P. melanotis* occurs during

the rainy season. Individuals need different resources requirements to fulfill the breeding

season and this impact on its dispersion and coexistence. Likewise, the partition of space

by these syntopic deermice must be facilitated by their respective locomotive habits as

semiarboreal (*P. difficilis*) and cursorial (*P. melanotis*). The long tail of *P. difficilis*, must

enable it to rush and climb along shrubs or trees (Álvarez-Castañeda 2005), thus easing

its preference for habitats where it can escape from predators. Indeed, adult color coat

changed in *P. difficilis* between pluvial seasons, becoming more similar to ground litter;

e. g., in this resident deermouse, color phenotype plasticity must be a cryptic response

elicited against predators. On the other hand, a smaller sized body, together with a shorter

tail and narrower soles (Fernandez *et al*. 2010), should enable the cursorial *P. melanotis*

to occupy such zones as M2 and M3, where predators cannot easily spot it through a

dense vegetation cover, so it can scape very quickly. Indeed, high shrubs cover provide

both protection from predators and food sources, since seeds may be concentrated under

shrub canopies (Thompson 1982; Mohammadi 2010). In fact, rodents usually avoid

foraging in unsheltered microhabitats and forest edges where they are more likely to be

spotted by avian (Kotler *et al*. 1991) and other vertebrate predators (Morris and Davidson

2000; Mohammadi 2010). In this study, we define to *P. melanotis* as an opportunistic

species throughout the study area because the capture frequencies of the individuals only

increased when the environmental conditions (*i. e*. availability of resources) became more

beneficial for the survival of these. That is, throughout the time of capture except for the

rainy season, its frequency of capture remained at low levels compared to those of *P.*

difficilis (resident species), increasing drastically in the rainy season when there is greater

accessibility for the use of resources and competition between the two species decreases.

Therefore, *P. melanotis* generally showed a more opportunistic pattern of habitat

occupation as compared to the larger sized *P. diffiilis*. Chupp (2002) also showed an

opportunistic habitat by small-sized *Peromyscus leucopus*, which he related to predation

risk. Further studies about the influence of predation on the opportunistic behavior of *P.*

melanotis are needed.

One possible consequence of such extrinsic and intrinsic operating factors is that

P. difficilis remained more abundant and always present at the microhabitats with more

stable elements in the study area (our unpublished data), while *P. melanotis* only

increased its abundance during the rainy season, when environmental conditions became

more benign, and plant cover became denser. Since rains promote an increase of primary

productivity, allowing more resource availability (mainly food) and enhancing

microhabitat carrying capacity, such habitat changes facilitate coexistence between both

Peromyscus, and with other small land mammals in the area (Castro-Campillo *et al.* 2008,

2012). Ripley's bivariate K supported and shed light into such outcome since the species

showed repulsion between them in almost all analyzed distances during the dry season;

i.e., due to resources availability are more scarce during this season, is almost null that a

mixed pair of mice used the same microhabitat. Moreover, as in this season is the primary

breeding season of *P. difficilis* is probably that territoriality behavior can occur.

Conversely, the intensity of such repulsion decreased substantially during the rainy

season; moreover, at some analyzed distances, the two deermice showed an association

pattern. Holding our results that the increase of resources available during the rainy

season can allow that the species can share the habitat. Thus, our results indicate that in

these two syntopic *Peromyscus*, both microhabitat use and spatial organization are

seasonally variable (Brown and Zeng 1989; Cramer and Willig 2002).

Finally, we want to highlight and recommend our sampling of environmental

features and quantitative analyses, especially for Ripley's bivariate K, as neither one has

been extensively used in studies focusing on the spatial dynamics of dispersion in small

rodents, despite its important advantages. The methodological approach used here,

together with the gathered evidence became relevant for providing us with necessary

information about the ability of this two *Peromyscus* to split resources in a mid-latitude

temperate forest at the edge of a megalopolis, which continuous growth produces

fragmentation and patching of natural microhabitats. Therefore, our local scope was

explicitly designed at this initial point, to focus on small-sized rodents with low vagility.

Besides, we successfully aimed to reconstruct vegetation structure, to provide us with

clues as to how to eventually manage such perturbed ecosystems for conservation

purposes. We also think that our methodology can be helpful in other scenarios for

understanding small rodents, such as ethological dynamics, activity patterns, and

reproductive systems at either or both intra or interpopulation levels.

Acknowledgments

We thank Juan Luis Patiño Ortega and Claudia Peralta Juarez for their fieldwork

assistance and friendly logistic support. The staff at UAMI mammal scientific collection

kindly provided all the facilities for specimen curation and housing. Authorities in charge

of DLNP (CORENA, SEMARNAT), as well as the forest rangers and administration staff

at the National Park, buoyed us with information, security, and logistics to complete our

studies. This paper contains partial information from the thesis of Master in Biology

(DCBS, UAM-I) of IMDA; IMDA received financial support through a fellowship

(283799, CVU 479479) from the National Council of Science and Technology

(CONACyT). This study was supported by annual grants (143.***.46 IB-DCBS-UAMI,

2013-2017) to AACC.

References

Álvarez-Castañeda, S.T. (2005). *Peromyscus melanotis*. Mamm. Spec. 764, 1-4.
Barry Jr, R.E., Heft, A.A., Baummer, TE. (1990). Spatial relationships of syntopic white-
footed mice, *Peromyscus leucopus*, deer mice, *Peromyscus maniculatus*, and red-
backed voles, *Clethrionomys gapperi*. Can Field-Nat. 104, 87-393.
Bellows, A.S., Pagels, J.F., Mitchell, J.C. (2001). Macrohabitat and microhabitat
affinities of small mammals in a fragmented landscape on the upper coastal plain of
Virginia. Amer. Midl. Nat. 146(2), 345-360.

Briani, D.C., Palma, A.R., Vieira, E.M., Henriques, R.P. (2004). Post-fire succession of
small mammals in the Cerrado of central Brazil. Biodivers. Conserv. 13(5), 1023-
1037.
Brown J.H, Zeng, Z. (1989). Comparative population ecology of eleven species of rodents
in the Chihuahuan Desert. Ecol. 70(5), 1507-1525.
Bowers, M.A. (1982). Foraging behavior of heteromyid rodents: field evidence of
resource partitioning. J. Mamm. 63(3), 361-367.
Bowers, J.H., Baker, RJ., Smith, M.H. (1973). Chromosomal, electrophoretic, and
breeding studies of selected populations of deer mice (*Peromyscus maniculatus*) and
black-eared mice (*P. melanotis*). Evol. 27(3), 378-386.
Bowman, J., Forbes, G., Dilworth, T. (2001). Landscape context and small-mammal
abundance in a managed forest. For. Ecol. Manag. 140(2), 249-255.
Bowman, A.S., Facelli, J.M. (2013). Fallen logs as sources of patchiness in chenopod
shrublands of South Australia. J. Arid Environ. 97, 66-72.
Bowne, D.R., Peles, J.D., Barrett, GW. (1999). Effects of landscape spatial structure on
movement patterns of the hispid cotton rat (*Sigmodon hispidus*). Landsc. Ecol. 14(1),
53-65.
Canfield, R.H. (1941). Application of the line interception method in sampling range
vegetation. J. For. 39(4), 388-394.
Carey, A.B., Harrington, C.A. (2001). Small mammals in young forests: implications for
management for sustainability. For. Ecol. Manag. 154(1), 289-309.
Carleton, M.D. (1989). Systematic and evolution. pp. 7-141 in Advances in the study of
Peromyscus (Rodentia) (G. L. Kirkland, Jr., and J. N. Layne, eds.). Texas Tech
University, Lubbock.
Castro-Campillo, A., León-Altamirano, L., Herrera-Muñoz, J., Salgado-Ugarte, I.,
Mendieta Márquez, E., Contreras-Montiel, JL., Serrano, H., Ramírez-Pulido, J.,
Salame-Méndez, A. (2012). Is there difference between T contents in two
populations of the Black eared Mouse, living under similar conditions but with
differences in two population patterns? Acta Zool. Mex. (n. s.) 28(3), 525-539.
Castro-Campillo, A., Salame-Méndez, A., Vergara-Huerta, A., Castillo, A., Ramírez-
Pulido, J. (2008). Fluctuaciones de micromamíferos terrestres en bosques templados
aledaños a la ciudad de México. Distrito Federal. Pp. 391-410 in Avances en el
Estudio de los Mamíferos de México (Lorenzo, C, Espinoza E, Ortega J, eds.).
Publicaciones especiales. Vol. II, Asociación Mexicana de Mastozoología, A. C.,
México, D. F. 726 pp.
Cramer, M.J., Willig, M.R. (2002). Habitat heterogeneity, habitat associations, and rodent
species diversity in a sand–shinnery-oak landscape. J. Mamm. 83(3), 743-753.
Coppeto, S.A., Kelt, D.A., Van Vuren, D.H., Wilson, J.A., Bigelow, S. (2006). Habitat
associations of small mammals at two spatial scales in the northern Sierra Nevada. J.
Mamm. 87, 402-413.
Currie, D.J. (1991). Energy and large-scale patterns of animal and plant species richness.
American Naturalist 137: 27-49.
Chesson, P. (2000). Mechanisms of maintenance of species diversity. Annu. Rev. Ecol.
Syst. 31, 343–366. doi: 10.1146/annurev.ecolsys.31.1.343
Chupp, AD. (2002). Habitat selection in four sympatric small mammal species and the
effects of potential predators on *Peromyscus leucopus*. Master of Science Thesis.
Department of Biology, Ohio University, 61 pp.
Dale, MRT. (1999). Spatial pattern analysis in plant ecology. Cambridge University
Press.

Dalmagro, A.D., Vieira, E.M. 2005. Patterns of habitat utilization of small rodents in an area of Araucaria forest in Southern Brazil. Austral Ecol. 30, 353-362.

De la Cruz-Argüello, I.M. (2015). Patrones de microdistribución en los mamíferos pequeños en vegetación secundaria en el Parque Nacional Desierto de los Leones. Tesis de Maestría en Ciencias. División de Ciencias Biológicas y de la Salud, Universidad Autónoma Metropolitana, Unidad Iztapalapa, pp. 256.

DeWalt, S.J., Maliakal, S.K., Denslow, J.S. (2003). Changes in vegetation structure and composition along a tropical forest chronosequence: implications for wildlife. For. Ecol. Manag. 182(1), 139-151.

Dooley, J.L., Dueser, R.D. (1996). Experimental tests of nest site competition in two *Peromyscus* species. Oecologia 105, 81-86

Dooley, J.L., Bowers, M.A. (1998). Demographic responses to habitat fragmentation: experimental tests at the landscape and patch scale. Ecol. 79(3), 969-980.

Fahrig, L., Merriam, G. (1994). Conservation of fragmented populations. Conserv. Biol. 8(1), 50-59.

Fernández, J.A., García-Campusano, F., Hafner, M.S. (2010). *Peromyscus difficilis* (Rodentia: Cricetidae). Mamm. Spec. 42(1), 220-229.

Forsman, E.D., Meslow, E.C., Wight, H.M. (1984). Distribution and biology of the spotted owl in Oregon, USA. Wildl. Monogrsaphs. 87, 1-64.

Galindo, C., Krebs, C.J. (1985). Habitat use and abundance of Deer Mice: interactions with Meadow Voles and Red-backed Voles. Can. J. Zool. 63, 1870-1879.

Gunther, P.M., Horn, B.S., Babb, G.D. (1983). Small mammal populations and food selection in relation to timber harvest practices in the western Cascade Mountains. Northwest Sci. 57, 32-44.

Grelle, C.E.V. (2003). Forest structure and vertical stratification of small mammals in a secondary Atlantic forest, southeastern Brazil. Stud. Neotrop. Fauna Environ. 38(2), 81-85.

Haim, A., Rozenfeld, F.M. (1993). Temporal segregation in coexisting *Acomys* species: the role of odour. Physiol. Behav. 54(6), 1159-1161.

Hammer, Ø., Harper, D.A.T., Ryan, P.D. (2001). PAST: PAleontological STatistics software package for education and data analysis. Palaeontol. Electronica 4, 1-9.

Johnson, M.L., Gaines, M.S. (1990). Evolution of dispersal: theoretical models and empirical tests using birds and mammals. Annu. Rev. Ecol. Syst. 21(1), 449-480.

Johnson, M.D. (2007). Measuring habitat quality: a review. Condor 109(3), 89-504.

Jorgensen, E.E. (2004). Small mammals use of microhabitat reviewed. J. Mamm. 85(3), 531-539.

Kalcounis-Rüppell, M.C., Millar, J.S. (2002). Partitioning of space, food, and time by syntopic *Peromyscus boylii* and *P. californicus*. J. Mamm. 83, 614-625.

Kaufman, D.W., Kaufman, G.A. (1989). Population biology. pp. 233-271 *in* Advances in the study of *Peromyscus* (Rodentia) (G. Kirkland, JR. and J. Layne, eds.). Texas Tech University Press, Lubbock.

Kotler, B.P., Ayal, Y., Subach, A. (1994). Effects of predatory risk and resource renewal on the timing of foraging activity in a gerbil community. Oecologia 100(4), 391-396.

Kotler, B.P., Brown, J.S., Hasson, O. 1991. Factors affecting gerbil foraging behavior and rates of owl predation. Ecol. 72, 2249-2260.

Lehtonen, J.T., Mustonen, O., Ramiarinjanahary, H., Niemelä, J., Rita, H. (2001). Habitat use by endemic and introduced rodents along a gradient of forest disturbance in Madagascar. Biodivers. Conserv. 10, 1185-1202.

Maser, C., Maser, Z. (1988). Interactions among squirrels, mycorrhizal fungi, and
coniferous forests in Oregon. Great Basin Nat. 48, 358-369.
M'Closkey, R.T., Fieldwick, B. (1975). Ecological separation of sympatric rodents
(*Peromyscus* and *Microtus*). J. Mamm. 56(1), 119-129.
M'Closkey, R.T., Lajoie, D.T. (1975). Determinants of local distribution and abundance
in White-Footed Mice. Ecol. 56(2), 467-472.
Mohammadi, S. (2010). Microhabitat selection by small mammals. Adv. Biol. Res. 4(5),
283-287.
Monamy, V., Fox, B.J. (1999). Habitat selection by female swamp rats (*Rattus lutreolus*)
drives asymmetric competition and coexistence with long-tailed mice (*Pseudomys*
higginsi). J. Mamm. 80, 232-242.
Morris, D.W. (1987). Ecological scale and habitat use. Ecol. 68(2), 362-369.
Morris, D.W. (1984a). Sexual differences in habitat use by small mammals: evolutionary
strategy or reproductive constraint? Oecologia 65(1), 51-57.
Morris, D.W. (1984b). Patterns and scale of habitat use in two temperate-zone, small
mammal faunas. Can. J. Zool. 62(8), 1540-1547.
Pianka, E.R. (1973). The structure of lizard communities. Annu. Rev. Ecol. Syst. 4(1),
53-74.
Pyare, S., Longland, W.S. (2001). Patterns of ectomycorrhizal fungi consumption by
small mammals in remnant old-growth forests of the Sierra Nevada. J. Mamm. 82,
681-689.
Ramírez-Pulido, J., Lira, I., Gaona, S., Müdespacher, C., Castro-Campillo, A. (1989).
Manejo y mantenimiento de colecciones mastozoológicas. Universidad Autónoma
Metropolitana, Unidad Iztapalapa, México, D.F.
Ripley, B.D. (1976). The second-order analysis of stationary point processes. J. Appl.
Probab. 13(2), 255-266.
Rosenberg, M.S., Anderson, C.D. (2011). PASSaGE: pattern analysis, spatial statistics
and geographic exegesis. Version 2. Methods Ecol. Evol. 2(3), 229-232.
Schoener, T.W. (1974). Resource partitioning in ecological communities. Science
185(4145), 27-39.
Seagle, S.W. (1985). Patterns of small mammal microhabitat utilization in cedar glade
and deciduous forest habitats. J. Mamm. 66(1), 22-35.
Schreiber, S. J., and Killingback, T. P. 2013. Spatial heterogeneity promotes coexistence
of rock-paper-scissors metacommunities. Theor. Popul. Biol. 86, 1–11. doi:
10.1016/j.tpb.2013.02.004
Smartt, R.A. (1978). A comparison of ecological and morphological overlap in a
Peromyscus community. Ecol., 59:21.
Thompson, S.D. (1982). Structure and species composition of desert heteromyid rodent
species assemblages: effects of a simple habitat manipulation. Ecol. 63(5),1313-
1321.
Valladares, F., Bastias, C.C., Godoy, O., Granda, E., Escudero, A. (2015). Species
coexistence in a changing world. Front. Plant Sci. 6, art. 866.
Vander Wall, S.B., Thayer, T.C., Hogbe, S.B., Beck, M.J., Roth, J.K. (2001). Scatter-
hoarding behavior of deer mice (*Peromyscus maniculatus*). West. N. Am. Nat. 61,
109-113.
Villanueva-Hernández, A.I., Delgado-Zamora, D.A., Heynes-Silerio, S.A., Ruacho-
González, L., López-González, C. (2017). Habitat selection by rodents at the
transition between the Sierra Madre Occidental and the Mexican Plateau, México. J.
Mamm. 98(1), 293-301.

Ward, J.H. (1963). Hierarchical grouping to optimize an objective function. J. Am. Stat.
Assoc. 58(301), 236-244.
Wiens, J.A. (2000). Ecological heterogeneity: an ontogeny of concepts and approaches.
In: The Ecological Consequences of Environmental Heterogeneity (Stewart AJA,
John EA, Hutchings MJ, eds). Blackwell, Oxford, Pp. 9-31.
Zar, JH. (1999). Biostatistical Analysis. Prentice Hall, Princeton.
Zavala-Hurtado, JA., Valverde, PL., Herrera-Fuentes, MC., Diáz-Solis, A. (2000).
Influence of leaf-cutting ants (*Atta mexicana*) on performance and dispersion
patterns of perennial desert shrubs in an inter-tropical region of Central Mexico. J.
Arid Environ. 46, 93-102.
Zavaleta, E. S., Shaw, M. R., Chiariello, N. R., Mooney, H. A., and Field, C. B. (2003).
Additive effects of simulated climate changes, elevated CO_2, and nitrogen deposition
on grassland diversity. Proc. Nat. Acad. Sci. 100, 7650–7654. doi:
10.1073/pnas.0932734100
Zielinski, W.J., Spencer, W.D., Barrett, R.H. (1983). Relationship between food habits
and activity patterns of Pine Martens. J. Mamm. 64, 387-396.

Table 1. Environmental variables used to measure the horizontal and vertical structure of microhabitat at each sampling station, showing its ecological interpretation (kind or resource type) for both species of *Peromyscus*.

Variable code	Predation		Resources				
	Diminished risk	Refuges	Breeding Space	Food	Temporal food and refuges	Permanent	Resting places
Vegetation Cover (cm)							
VC10	X			X	X		X
VC25	X		X	X	X		X
VC35	X		X	X	X		X
VC50	X		X	X	X		X
VC100	X		X	X	X		X
Frequency of Herbaceous Plants (cm)*							
FH11	X		X	X	X		X
FH12	X		X	X	X		X
FH13	X		X	X	X		X
FH14	X		X	X	X		X
FH15	X		X	X	X		X
Frequency of Woody Plants (cm)*							
FW21	X	X	X	X	X	X	X
FW22	X	X	X	X	X	X	X
FW23	X	X	X	X	X	X	X
FW24	X	X	X	X	X	X	X
FW25	X	X	X	X	X	X	X
Species Vegetation Richness							
VR	X	X	X	X	X	X	X
Total of Herbaceous Plants							
TH	X		X	X	X		X
Total of Woody Plants							
TW	X	X	X	X	X	X	X
Rocks							
RC	X	X	X			X	X
Organic Matter (litter, twigs, small trunks)							
OM				X	X		
Bare Soil							
BS					X		
Coverage of Logs (stumps, fallen trunks)							
CLO	X	X	X			X	X
Numbers of Burrows							
BW	X	X	X			X	X

* = Frequencies of plant types (FH, FW) involve the same heights as in VC.

Table 2. Normality (**NOR**) tests for 22 variables (**VAR**) of microhabitat, and comparisons[abc] (**COM**) for average values between the dry (**DRY**) and rainy (**RAINS**) seasons, respectively. Variable names in Table 1.

VAR	NOR[a]		COM	VAR	NOR[a]		COM	VAR	NOR[a]		COM
	DRY	RAINS			DRY	RAINS			DRY	RAINS	
BS	0.0001	0.0001	*1.0[c]*	**VC100**	0.002	0.003	0.0001[c]	**TW**	*0.143*	*0.08*	0.0004[b]
CLO	0.0001	0.0001	*0.972[c]*	**VR**	0.006	*0.082*	0.0001[b]	**FW21**	0.0001	0.0001	0.004[c]
OM	0.010	0.034	0.0001[c]	**TH**	0.0001	*0.072*	0.0001[b]	**FW22**	0.0001	0.0001	0.026[c]
RC	0.0001	0.0001	*0.432[c]*	**FH11**	0.015	0.028	*0.658[c]*	**FW23**	0.0001	0.0001	*0.875[a]*
VC10	0.002	0.041	0.0001[c]	**FH12**	0.0001	0.0001	0.001[c]	**FW24**	0.0001	0.0001	*0.378[c]*
VC25	0.009	0.003	0.0001[c]	**FH13**	0.0001	0.0001	0.0001[c]	**FW25**	0.0001	0.001	0.0001[c]
VC35	0.0001	0.002	0.0001[c]	**FH14**	0.0001	0.0001	0.0001[c]	BW	0.0001	0.0001	1
VC50	0.0001	0.019	0.0001[c]	**FH15**	0.0001	0.0001	0.0001[a]				

Parametric: [a]Kolmogorov-Smirnov; [b]Student for two samples; non-parametric: [c]Wilcoxon). The *p*-values in italics do not reject normality ($p \geq 0.05$), or do not indicate significant differences between pluvial seasons.

 Table 3. Nearest Neighbor Analysis for both species of *Peromyscus*, it shows the mean

788 distance in which individuals from both *Peromyscus* are segregated. R-value indicates

789 how the observed distribution deviates from random. Clustered points give $R<1$, Poisson

790 patterns give $R\sim1$, while overdispersed points give $R>1$.

791
792

	Dry season		Rain season	
	P. difficilis	*P. melanotis*	*P. difficilis*	*P. melanotis*
Points	57	39	58	52
Mean distance	1.40	2.39	1.67	2.09
R	0.42	0.77	0.55	0.64
P value	1.1263×10^{-16}	0.006	8.6577×10^{11}	8.7708×10^{7}
Point pattern	Clustering	Clustering	Clustering	Clustering

812 **Table 4.** Principal components used as effect model in the GLMs analysis for both

813 seasons. Red colors represent the variables that have the major eigenfactors to the

814 ordination of the components.

Variables	Dry season			Rainy season		
	PC1	PC2	PC3	PC1	PC2	PC3
VC10	0.713	-0.388	-0.134	0.659	-0.179	-0.374
RC	-0.145	-0.055	-0.110	0.014	0.062	0.034
CLO	0.030	0.118	0.828	-0.153	0.411	-0.583
OM	-0.604	0.303	-0.487	-0.499	-0.122	0.775
BS	0.130	-0.187	0.112	-0.028	0.154	-0.328
VC25	0.415	0.280	0.034	0.658	-0.061	0.237
VC32	0.287	0.184	0.052	0.580	-0.003	-0.126
VC50	0.523	0.107	0.078	0.669	-0.173	-0.030
VC100	0.335	0.408	0.052	0.656	-0.053	0.107
VR	0.762	0.251	-0.269	0.776	0.380	0.058
TH	0.837	-0.430	-0.131	0.874	-0.319	-0.076
TW	0.520	0.684	-0.168	0.615	0.711	0.270
FH11	0.784	-0.394	-0.214	0.678	0.007	-0.228
FH12	0.573	-0.345	0.176	0.548	-0.278	0.256
FH13	0.247	0.073	0.113	0.463	-0.260	0.067
FH14	0.149	-0.186	-0.023	0.580	-0.530	-0.043
FH14	0.080	0.117	-0.236	0.481	-0.267	-0.019
FW21	0.127	0.510	-0.394	0.342	0.603	0.311
FW22	0.015	0.064	-0.105	0.266	0.212	0.416
FW23	0.199	0.256	-0.228	0.196	0.366	-0.241
FW24	0.424	0.359	0.195	0.337	0.165	0.159
FW25	0.515	0.366	0.380	0.494	0.469	0.026
BW	-0.105	0.249	0.557	-0.170	0.332	-0.218

Table 5. Generalized Linear Models constructed with the Principal Components and the

frequency of P. *difficils* and *P. melanotis* for both seasons. The partition of resources by

both species is observed at both seasons of the year. Different main components within

each season explain the frequency of capture of each of the species. Also, the table is

related to what type of microhabitat the new variables belong to (M1-M3). The goodness

of fit tests are shown; Pearson and Deviance.

Term	Estimate	Std Error	L-R x^2	P value	Pearson	Deviance
P. difficilis						
Dry season						
Intercept	-0.97	0.15	55.73	0.0001	$x^2 = 155.80$ ($p = 0.0096$)	$x^2 = 123.56$ ($p = 0.3210$)
PC1 = M2	0.11	0.05	5.53	0.0186	AICc = 212.55	
PC3 = M1	0.45	0.06	44.73	0.0001		
Rainy season						
Intercept	-0.40	0.11	14.37	0.0002	$x^2 = 168.56$ ($p = 0.0008$)	$x^2 = 156.43$ ($p = 0.0062$)
PC2 = M3	0.23	0.06	10.92	0.0009	AICc = 286.76	
PC3 = M1	0.26	0.07	10.95	0.0009		
P. melanotis						
Dry season						
Intercept	-1.21	0.18	74.28	0.0001	$x^2 = 237.65$ ($p = 0.0001$)	$x^2 = 138.70$ ($p = 0.0935$)
PC2 = M3	0.46	0.09	23.34	0.0001	AICc = 204.47	
Rainy season						
Intercept	0.60	0.12	28.15	0.0001	$x^2 = 132.71$ ($p = 0.13$)	$x^2 = 132.72$ ($p = 0.13$)
PC1 = M2	0.10	0.04	4.47	0.0343	AICc = 241.01	

Fig. 1. Sampling plot design. a) Total plot area was 2475 m^2; each intersection (n = 120)

was an independent sampling station for capturing mice. An influence area of 2.5 m^2

around each station was constructed for sampling 24 microhabitat environmental

variables. b) Close up to the influence area showing the eight transects used for drawing

environmental variables; note subdivisions into three subtransects.

Fig. 2. Map of the distribution of the microhabitats (M1 = network, M2 = green, M3 =

blue) in the study quadrant for dry (A) and rainfall (B). The bar graph below shows the

change in the number of sampling stations between dry and rainy seasons. ** $p = 0.001$

Fig. 3. Box-plots are showing the seasonal heterogeneity of the 23 variables sampled in

the study plot between dry and rainy seasons. Overall, vegetation variables show an

increase in the rainy season, while the organic matter (OM) decrease. Structural elements

as logs on the ground do not shot differentiation between seasons. Some variables were

not present in high quantities along the study plot. Therefore we were unable to build the

box-plot. The red cross represents the mean of each variable.

Fig. 4. Spatial analysis with Kernel density points showing capture frequencies for *P.*

difficilis at dry (a) and rainy (b) seasons, and for *P. melanotis* at dry (c) dry and rainy (d).

Dots depict actual capture points, while hot spots color indicates higher density.

Fig. 5. Correspondence analysis from contingency tables of frequency capture data for

each deermouse (*P. difficilis* and *P. melanotis*) in each microhabitat (M1-3), during the

dry (a) and rainy (b) seasons, respectively. Axis c1 and c2 indicate the coordinates

resulted from the ordination of each variable from the analysis. Figure c shows the plots

for the GLMs models, the response variable for each graph are the predicted values of the

model to that principal component (the model effect). All of the PCs fitted in the model

show a positive relationship, indicating that the probability of capture frequency of the

individuals of two mice, increases with the presence of that variables. Therefore, the

figure also shows in the "X" axis in parenthesis the variables with the higher loading

scores in the ordination of this PCs.

Fig. 6. Ripley's bivariate K Analysis for interactions between the two *Peromyscus*. (a)

Repulsion pattern between both species in all analyzed distances during the dry season.

(b) Attraction from 6 - 12 m, and repulsion at 5, 11, and 14 m during the rainy season.

Black solid line represent the Ripleys K bivariate pattern; red dashed lines represent the

95% CI; blue solid line shows the null hypothesis.

905   **Fig. 1**

936
937 **Fig. 2**
938
939
940
941

961
962 **Fig. 3**
963
964
965

991
992 **Fig. 4**

1041 **Fig. 5**

1042

1043

1044

1059

1060

1067

1068

1075

1076

1077

Fig. 6