3   Spatiotemporal microhabitat heterogeneity and dispersion patterns of two small

4   mammals in a temperate forest

5   Heterogeneidad espaciotemporal del microhábitat y los patrones de dispersión de dos

6   pequeños mamíferos en un bosque templado

8   Ivan Mijail De-la-Cruz[1,3]*, Alondra Castro-Campillo[1], Alejandro Zavala-Hurtado[1],

9   Arturo Salame-Méndez[2], and José Ramírez-Pulido[1]

[1]Department of Biology, Metropolitan Autonomous University, Iztapalapa Unit (UAM-I).

Av. San Rafael Atlixco 186, CP. 09340, Ciudad de México, México. Email:

imda@xanum.uam.mx (IMDC)*, acc@xanum.uam.mx (ACC), jazh@xanum.uam.mx (AZH),

jrp@xanum.uam.mx (JRP).

[2]Department of Reproduction Biology, Metropolitan Autonomous University, Iztapalapa Unit

(UAM-I). Av. San Rafael Atlixco 186, CP. 09340, Ciudad de México, México. Email:

asam@xanum.uam.mx (ASM).

[3]Department of Evolutionary Ecology, Institute of Ecology, National Autonomous University

of Mexico. Av. Circuito Exterior S/N anexo Jardín Botánico exterior, Ciudad Universitaria,

C.P. 04500, Ciudad de México, México. Email: imda@xanum.uam.mx (IMDC)*.

* Corresponding author

Abstract. One basic issue in ecology is to understand how seasonal spatiotemporal shifts in

habitat heterogeneity shape spatial patterns in the distribution and coexistence of organisms.

For example, how do spatial dispersion of syntopic species reflects their respective resource

preferences and partition, during microhabitat seasonal shifts? To address this, we assessed

changes in microhabitat structure heterogeneity, between the two annual seasons (dry and

rainy seasons) in a midlatitude temperate forest by analyzing 23 habitat variables using

multivariate statistics. Furthermore, to determine how such microhabitat changes affected the

spatial segregation of two congeneric species of small mammals (*Peromyscus difficilis*, *P.*

melanotis), we used second-order spatial statistics to detect changes in their microdistribution

changes and general linear models to evaluate their habitat preferences. In accordance with

their respective morphology and ambulatory habits, the medium-sized, semi-scansorial *P.*

difficilis was resident year-round and preferred microhabitats with high amounts of log

ground cover, while the small, opportunistic, cursorial *P. melanotis* varied its occupancy area

depending on the density of herbaceous and cover of woody plants. Both species showed

more similar microdistribution patterns under the more benign microhabitat conditions of the

rainy season (denser plant coverage, humid-cool weather), but became more segregated form

each other during the less favorable conditions of the dry season (scarcer plant coverage, dry-

cold weather). Therefore, seasonal changes in microhabitat heterogeneity influenced the

dispersion patterns of both *Peromyscus* species and revealed their temporal preferences for

microhabitat resources.

Keywords: ecological dispersion; microhabitat selection; *Peromyscus*; resource partition;

use of space.

Resumen. Un tema básico de la ecología es comprender cómo los cambios espacio-

temporales de la heterogeneidad del microhábitat, dan forma a los patrones espaciales de

distribución y coexistencia de los organismos. En otras palabras, cómo es que la dispersión

espacial de especies sintópicas refleja sus preferencias y partición de recursos durante los

cambios estacionales del microhábitat? Para abordar esto, evaluamos los cambios en la

estructura vertical-horizontal del microhábitat (heterogeneidad) a lo largo de las dos

estaciones anuales (temporadas seca y lluviosa) de un bosque templado, analizando 23

variables con estadística multivariada. Además, para determinar cómo tales cambios del

microhábitat afectaban la segregación espacial de dos especies congenéricas de pequeños

mamíferos (*Peromyscus difficilis*, *P. melanotis*), utilizamos estadística espacial de segundo

orden y modelos lineales generalizados para detectar sus cambios específicos de

microdistribución y sus preferencias de hábitat, respectivamente. De acuerdo con su

morfología y hábitos ambulatorios respectivos, el mediano y semiescansorial *P. difficilis* fue

residente durante todo el año y prefirió los microhábitats con una alta cobertura de troncos

sobre el suelo, mientras que el oportunista, pequeño y cursorial *P. melanotis*, varió su área de

ocupación, dependiendo de la densidad de plantas herbáceas y la cobertura de plantas leñosas.

Bajo condiciones de microhábitats más benignas en la época de lluvias (mayor densidad de

cobertura vegetal, clima húmedo-fresco), ambas especies mostraron patrones de

microdistribución más próximos, pero se volvieron más segregadas entre sí durante las

condiciones menos favorables de la temporada seca (cobertura vegetal más escasa, clima

seco-frío). Por lo tanto, los cambios estacionales en la heterogeneidad del microhábitat

influyeron en los patrones de dispersión de ambas especies de *Peromyscus* y revelaron sus

preferencias temporales por los recursos de microhábitat.

Palabras clave: dispersión ecológica; *Peromyscus*; selección del microhábitat; partición de

recursos; uso del espacio.

Introduction

Spatiotemporal habitat heterogeneity has a substantial impact on species coexistence. For

example, species may become spatially segregated according to niche preferences, such as

resource requirements (Valladares *et al.* 2015). Model simulations reveal the potentially

important role of spatial heterogeneity and its complex and delicate interplay with dispersal in

mediating long-term outcomes of species coexistence (Schreiber and Killingback 2013;

Valladares *et al.* 2015). For instance, an increase in the number of habitat types, resources,

and structural complexity should increase available niche spaces, thus allowing more species

to coexist (Currie 1991). Equally crucial for the maintenance of species coexistence is habitat

heterogeneity over time, whose influence on natural communities also varies depending on

the temporal scale (Valladares *et al.* 2015). Temporal fluctuations in habitat structure can

stabilize coexistence via the "storage effect" (Chesson 2000), when inter-annual variation in

climate or resource availability favors one group of species over others (Zavaleta *et al*. 2003).

Heterogeneous habitats contain patches of varying size, distribution, resources,

environmental conditions, and species composition (Fahrig and Merriam 1994) that vary

spatiotemporally, depending on observation scale and habitat type (Wiens 2000). It is highly

likely that small body size species such as small mice perceive spatial habitat heterogeneity at

a fine scale (microhabitat). Therefore, both movement and foraging by this kind of small

mammals are affected by habitat heterogeneity (Bowne *et al.* 1999), having different

responses among species or demographic groups within them (Dooley and Bowers 1998).

Here we focus on whether small mammals are capable of perceiving and responding to

spatiotemporal microhabitat heterogeneity.

Small mammals use resources selectively, based on their requirements for growth,

survival, and reproduction (Johnson and Gaines 1980). For instance, spatiotemporal habitat

heterogeneity affects *Peromyscus* mice in terms of species coexistence (M'Closkey and

Fieldwick 1975; M'Closkey and Lajoie 1975), spatial segregation (Monamy and Fox 1999),

and competition (Morris 1984a; Seagle 1985). Indeed, several studies have quantified

variation in resource use in heterogeneous environments within a community among mice

from the same family or genus (Kaufman and Kaufman 1989 and references therein). Indeed,

closely related species are especially valuable for studies of spatial and resource partitioning,

since they are most likely to be current or past competitors (Kalcounis-Rüppell and Millar

2002). Therefore, since habitat use varies spatiotemporally (Haim and Rozenfeld 1993) due

to distribution and availability of resources, we expect this variation to affect the use of

resources and relationships among closely related species.

In this study, the primary objective was to analyze how microhabitat heterogeneity is

structured between the dry and rainy seasons of a temperate mixed forest, and how these

changes affect the coexistence of *Peromyscus difficilis* and *P. melanotis*. To do so, we

addressed four main aspects; first, we assessed the presence of spatiotemporal heterogeneity

at a fine scale (microhabitat) according to vertical and horizontal structure indicators.

Second, we determined the ecological dispersion patterns of *P. difficilis* and *P. melanotis* and

tested whether the use of space by these two *Peromyscus* was affected by spatiotemporal

changes in microhabitat heterogeneity. Third, we determined what kind of spatial

relationships occurred between the two species depending on seasonal changes in

microhabitat heterogeneity (*i. e*., attraction/positive: sharing microhabitats and resources, or

repulsion/negative: not sharing). Finally, we assessed what particular structural elements of

each one of the microhabitats best explained capture frequency for each *Peromyscus* species

in each season.

Materials and methods

Study area. The study area was located in a temperate mixed forest (coniferous and broad-

leaved trees) at Desierto de Los Leones National Park (DLNP, CONANP 2006), Mexico City,

within the Transmexican Neovolcanic Range (Universal Transverse Mercator, UTM,

boundary coordinates: 465261.25 m E, 2137029.52 m N; 468996.54 m E, 2129839.47 m N).

The isothermal temperate climate belongs to the most humid of the subhumid type with

summer rainfalls (C(W2) W (b´)ig); winter rainfall accounts for < 5% of total annual rainfall

(CONANP 2006). There is an annual accumulated precipitation of 1340.6 mm (monthly

mean 111.72 ± 109.1; minimum 10.3 in December, maximum 275.9 in August). The rainy

season (CONANP 2006) is from summer through early fall (June - September) with humid-

cool weather and a monthly average precipitation of 252.92 ± 28.01 (218.8, September -

275.9, August) and an average monthly temperature (°C) of 11.72 ± 0.53 (11.3, September -

12.5, June). The dry season (CONANP 2006) occurs from late fall through winter

(November - February) with cold, dry weather: monthly average precipitation (mm) from

November - January is 13.2 ± 3.11 (10.3, December - 16.5, November), and average monthly

temperature (°C) is 8.97 ± 0.68 (8.3, January - 9.7, November). The dominant vegetation in

most of DLNP is *Abies-Pinus-Quercus* forest, at 2,800 - 3,000 m (CONANP 2006).

According to official sources (CONANP 2006) and our unpublished data, the most

conspicuous plant species by stratum are as follows. The canopy includes Christmas fir

(*Abies religiosa*), several species of pine trees, including *Pinus patula, P. hartwegii, P.*

leiophylla, and *P. montezumae*, and oak trees *Quercus laurina, Q. castanea, Q. laeta*, together

with other broad-leaved trees (*e. g., Prunus serotina, Salix paradoxa*, and *Buddleia cordata*).

Among understory shrubs are *Senecio barba-johannis, Symphoricarpos microphyllus,*

Cestrum anagyris, Solanum cervantesii, Physalis viscosa, Fuchsia microphylla, *Baccharis*

conferta; there are also isolated elements of *Berberis moranensis, Garrya laurifolia, Arbutus*

xalapensis, Buddleia cordata, and *Clethra mexicana* within the oak trees. Among the most

frequent herbs are *Acaena elongata, Sigesbeckia jorullensis, Alchemilla procumbens*,

Geranium seemannii, Valeriana clematitis, and *Archibaccharis hirtella*. The ground level

bears a rich variety of mosses (*e. g., Aloinella catenula, Anacolia laevisphaera, Bartramia*

spp., *Bryum procerum, Grimmia* spp., *Herzogiella cylindricarpa, Mironia* spp.) and fungi,

whose fruiting bodies are conspicuous during the rainy season (*e. g.,Cantharellus cibarius,*

Phlogiotis helvelloides, Amanita ccf. *citrina, Geastrum triplex, Ramaria* sp.).

Microhabitat features. We set a 2,475 m^2 (55 x 45 m) surface plot at 2,289 m (UTM 2144958

m N, 14477883 m E) (Figure 1). The plot was gridded with 12 vertical lines (A - L) by 10

horizontal lines (1 - 10), with lines placed every 5 m. Intersections between

vertical/horizontal lines were marked with buried wooden stakes (150 x 2.5 x 2.5 cm) to

construct a coordinate system for 120 independent sampling stations (Figure 1). To assess

spatiotemporal heterogeneity of microhabitats within the entire plot, we delimited an

influence zone (Figure 1), drawing a 2.5 m^2 square around each sampling station. We used

these influence zones to sample microhabitat characteristics around each sampling station,

using 23 environmental variables (Table 1) for ten days: once in the dry season (July) and

once during the rainy season (February; CONANP 2006). All selected variables qualify as

components of the vertical and horizontal structure of microhabitats, as well as factors

affecting rodent distribution at this spatial scale (Morris 1984b, 1987; Jorgensen 2004;

Coppeto *et al.* 2006; Villanueva-Hernández *et al.* 2017). Further, taken together, these 23

variables (Table 1) provide information about possible refuges for the small mammals from

predators, as well as spaces for resting, breeding, food resources, and safe routes which can be

used to move from one place to another to avoid predation risk (Jorgensen 2004). More

detailed information about the ecological significance of each variable appears in Table 1.

The variables we sampled include the following: Vegetation Cover at five different

heights at the understory in cm: *i. e*., 10, 25, 35, 50, and 100 cm (VC10 - VC100; *e. g*., VC10

is vegetation coverage at 10 cm height). Number of herbaceous forms at five different

heights: *i. e*., 10, 25, 35, 50, and 100 cm (HF1-5; *e. g*., HF3 is number of herbaceous plants at

level 3, or 25 cm height). Number of woody forms (WF1-5; *e. g.,* WF5 is the number of

woody forms at level 5, or 100 cm height). Total Herbaceous plants (TH) and Total Woody

plants (TW) are the total number of each type counted within each influence zone.

Vegetation species richness (VR) is total number of plant species. Other variables included

measurements of the coverage of fixed microhabitat elements (rocks = RC, logs = CLO,

stumps and fallen branches), organic matter (OM, mostly leaf litter and small twigs), and bare

soil (BS). Finally, we counted the number of mouse burrows (BW), recording each hole in

the ground *ca.* 8 - 12 cm of diameter within the influence zone (Álvarez-Castañeda 2005;

Fernandez *et al.* 2010).

To measure all variables for each season in all influence zones, we adapted Canfield's

(1941) Line Intercept (LI) method, which allowed us to sample the within-plot variation and

quantify microhabitat changes in plant species cover and height over time. We drew eight

graduated transects (cm) within the respective influence zone of each station to cover the

entire surface area (Figure 1): four 2.5 - m transects were cardinally oriented (N, S, E, W),

while the other four (3.30 m long) were diagonally oriented (NE, NW, SE, SW). We

measured plant coverage at the five different heights (VC10 - 100), as well as coverage of

fixed elements; *i. e.,* rocks (RC), logs (CLO), bare soil (BS), dead organic matter (OM) for all

transects. We only counted plants intercepted by diagonal transects for vegetation species

richness (VR), for number of plant life forms at the different height levels (herbaceous, HF1 -

5; woody, WF1 - 5), and for total herbaceous (TH) and woody plants (TW). We calculated

the percent coverage of each variable within the influence zones, using the formula

$\sum(Xi)\frac{100}{2400}$, where Xi is the length along the transect occupied by each plant in centimeters,

and 2,400 is the total length (sum of the eight transects).

Mouse sampling. We captured *Peromyscus* mice alive over ten months to include data for the

dry (October 2013 to February 2014) and rainy (March 2014 to July 2014) seasons. We set a

single live trap (H. B. Sherman, Inc., Tallahassee, FL 32303, USA), baited with oat flakes and

vanilla extract (Figure 1) at each independent sampling station of the plot (n = 120) (Figure

1). Traps were set for two consecutive nights, for a total of 20 capture events and 2, 400 trap-

nights. To record the capture frequency of mice in all the stations for every capture event, we

recorded the sampling station coordinates at which mice were caught. We also recorded the

species, sex, age, reproductive condition, and any other conspicuous trait that allowed us for

individual recognition of every mouse by visual inspection; *e. g.*, acquired traits, such as scars

and missing parts in the ears, tail or a missing eye. In addition, to discriminate between new

captured individuals and recaptured individuals between capture events, we stained the

abdomen of each captured mouse with gentian violet. Since temporary marking with gentian

violet lasted in average 75 ± 7 days in the rainy season and up to 180 ± 8 days in the dry

season, it proved to be a non-intrusive, affordable, and suitable method for our study

objectives. When the gentian violet mark began to fade and a mouse was recaptured, the

mark was reapplied. This non-intrusive marking allowed us to avoid overestimating the

capture frequency by ensuring we never counted the same individual twice during the same

capture event. Moreover, our analyses showed the same dispersion patterns regardless of

whether we used both the frequency of newly-captured individuals and recapture frequencies

during all capture events, or only new captures in our analyses. In addition, to avoid

recurrence behavior (*e. g.,* mice coming back to the traps for bait) or shyness (*e. g.,* mice

avoiding traps due to other mice odors), all the traps were thoroughly cleaned and randomly

oriented within the influence zone in each capture event. To prevent hypothermia during

capture, we placed 3 - 5 cotton balls inside the trap and put the traps inside open plastic bags.

Mice capture and handling followed the guidelines of the American Society of Mammalogists

(Sikes 2016). Any mice that died were prepared as voucher specimens and incorporated into

the Mammal Collection of the Universidad Autónoma Metropolitana-Iztapalapa (UAMI,

Ramírez-Pulido *et al*. 1989). A scientific collecting permit, SEMARNAT – 08 – 049 - B, was

issued to Alondra Castro-Campillo (ACC) by DGVS, SGPA - 09712/13, SEMARNAT.

Statistical analysis. We standardized all variables and run statistical tests with an alpha level

of 0.05. Examining the Variance Inflation Factor revealed showed no multicollinearity

among the 23 habitat variables (*i. e.,* all values were at a threshold < 6, Zar 1999). To identify

and categorize microhabitat heterogeneity within the plot grid, we performed two independent

Hierarchical Cluster Analysis (HCA) in JMP® (ver. 9, SAS Institute Inc., Cary, NC, 1989 -

2007), one for each season. These analyses enabled us to cluster sampling stations with

similar microhabitat characteristics, according to the 23 environmental variables, in each

season. We used Ward's method (1963) to construct the respective dendrograms, where the

distance between two clusters is the ANOVA sum of squares between them, added up over all

quantitative variables. From visual inspection of each dendrogram and given that there were

no drastic changes in the variance scree plot, we obtained three groups of sampling stations

for each season, which we interpreted as distinct microhabitat types (M1 - 3). We further

statistically validated such microhabitats (Figure A.1) by carrying out independent

Discriminant Analyses in each season (DA, Addinsoft SARL's XLSTAT 2013; F = 9.99 dry;

F = 9.64, rains; Wilk's Lambda = 0.0001 in both). We constructed a different map for each

season, assigning each sampling station one of three colors, according to its microhabitat type

(M1: red, M2: green, M3: blue, Figure 2). To construct the respective descriptive typologies

for the fine scale behavior of each one of the environmental variables, we plotted the 23

standardized variables in the "Y" axis and the microhabitat types in the "X" axis in each

season (Figure A.1).

We calculated descriptive statistics for each variable (*e. g.*, mean, median, standard

deviation, minimum and maximum values), and tested for normal distribution with

Kolmogorov-Smirnov tests (Zar 1999). Then, we analyze distribution changes in each

variable between the rainy and dry seasons using Student's t-test for normally distributed

variables and the non-parametric Wilcoxon test for non-normal variables (Table 2). We used

a contingency table and x^2 test to determine whether the number of stations belonging to each

type of microhabitat changed between seasons (seasonal microhabitat heterogeneity). To

examine the association between changes in spatial heterogeneity of microhabitats and

dispersion patterns of the *Peromyscus* along the entire grid, we conducted a Contingency

Table Analysis in each season. In each contingency table, microhabitats were the rows and

capture frequency of each *Peromyscus* species were the columns (Table A.2). Then, in order

to visualize associations between the *Peromyscus* and microhabitats, we performed a

correspondence analysis using the contingency tables for each season (Figure 5). We

performed all of these analyses with XLSTAT 13 (Addinsoft SARL).

Spatial analysis. To map variations in point density captures of small mammals, and to find

density gradients across the plot area, we used the Kernel function (PAST, ver. 3.14, Hammer

et al. 2001). To test the ecological dispersion patterns of each *Peromyscus* species within the

plot (*i. e.,* clustering or overdispersion) in each season, we used a "Nearest Neighbor

Analysis" (NNA, Clark and Evans 1954) using PAST software (ver. 3.14, Hammer *et al.*

2001). We applied the wrap-around edge effect adjustment, which is only appropriate for

rectangular domains (like our sampling plot; Hammer *et al.* 2001). In general, the NNA

compares the mean distance of each focal individual from its nearest conspecific neighbor,

with the mean distance expected for a set of points randomly dispersed at the same density

(Vázquez and Álvarez-Castañeda 2011). The ratio of the observed mean distance to the

expected distance (R) indicates how the observed distribution deviates from random; i.e.,

clustered points give $R < 1$, Poisson patterns give $R \sim 1$, and overdispersed points give $R > 1$.

273   To assess whether the two species were associated (attracted) or disassociated

(repelled) with each other, we compared dispersion patterns between the two mice, using

Ripley's K bivariate function (Ripley 1976), since the method considers all distances among

individuals located under a Cartesian scheme (X, Y) in a quadrat plot (Ripley 1976; Dale

1999; Zavala-Hurtado *et al.* 2000). We used PASSaGE (ver. 2, Rosenberg and Anderson

2011) to carry out Ripley's bivariate K analysis. We used the option to test the associations

conditional on the current locations hypothesis in PASSaGE, where the point locations

(sampling stations) remain fixed, and only the point types (each *Peromyscus* species as type

points A or B) are randomized; once the number of points of a determined type remains fixed

(*Peromyscus* species A), the association of the another points type (*Peromyscus* species B)

with a specific fixed location is randomized. In this case, the test does not address whether

the *Peromyscus* species are themselves at random or clustered, but rather whether the

association of type A points (*P. difficilis*) with type B points (*P. melanotis*) is what one would

expect, given the locations of the points as fixed (Rosenberg and Anderson 2011). We

assigned coordinates to each sampling stations considering 5 m distance between them, using

the Cartesian coordinates of each sampling station where we captured mice as data input. Our

null hypothesis was independence between *P. difficilis* and *P. melanotis* (Ripley 1976; Diggle

1983; Dale 1999); *i. e.,* we wanted to assess whether type A points (*P. difficilis*) were

associated, or disassociated, with type B points (*P. melanotis*). In other words, Ripley's

$K_{12}(d)$ allowed us to assess spatial attraction or repulsion between the two species, among the

stations of the plot. To evaluate the statistical significance of $K_{12}(d)$, we estimated the 95 %

confidence interval (95 % CI), using a Monte Carlo procedure, based on 1000 stochastic

relocation simulations of the sampling stations in the plot (Upton and Fingleton 1985; Zavala-

Hurtado *et al.* 2000). When L(d) was positive and above the upper limit of the 95% CI, we

inferred dissociation or repulsion between the two *Peromyscus* at the corresponding (d) scale;

a significant negative deviation indicated a pattern of association or attraction between them

(Dale 1999). If L(d) was within the 95 % CI for a given value of d, the null hypothesis of

independence between the two contrasts could not be rejected (Dale 1999). The height of the

L(d) function (peak height) indicates the intensity of the association or repulsion. We carried

out a control for edge effect for the Ripley's K analysis, rescaling the base counts overlapped

with the study boundary (Rosenberg and Anderson 2011).

Prediction of microhabitat elements affecting the use of space by each species. The next *s*tep

was to assess which specific environmental variables best explained the frequency of each

species within the complete grid at each season. For this, we performed Generalized Linear

Models (GLMs), using JMP ® (ver. 9, SAS Institute Inc., 1989 - 2007, Cary, NC,). First, we

calculated the mean for each one of the 23 variables in each microhabitat type, and then

conducted a Principal Component Analysis (PCA) on the microhabitat means to obtain

functions summarizing the most variance within each season. Because in both seasons the

first three principal components (PC) summarized a good portion of the variance (38.61 for

the dry season and 45.92 for rains), we used them to construct the GLMs. Since we found no

multicollinearity between variables, we assessed different models using the PCs as

explanatory variables to explain the frequency of individuals of each species for every season.

Then, to visualize the fit of the model, the predicted values from the GLMs were plotted

against the PC of the best model chosen. For each species, the response variable in the

models was the capture frequency at each trapping station of the grid, and we assumed a

Poisson distribution with a log link function.

Criteria for selecting the best models to explain the frequency of mice of each species

in every season within the sampling stations included: a) model significance of $p < 0.05$; b)

lowest Akaike Information Criterion (AIC), *i. e.,* a measure of goodness-of-fit penalized by

the number of variables (Posada and Buckley 2004); c) Pearson goodness-of-fit ($p < 0.05$)

and its deviation (Deviance; $p < 0.05$). If the models met all the criteria and showed very

similar results, we selected the final model also taking into consideration the ecological and

biological significance of the results.

Results

Frequency of captures. During the dry season, the total number of individuals was 111: 64

individuals for *P. difficilis* and 47 individuals for *P. melanotis*. During the rainy season, the

total number of individuals increased to 168: 87 individuals for *P. difficilis* and 81 individuals

for *P. melanotis* (Table A.2).

Behavior of the 23 microhabitat variables between seasons. When considering the mean

value over all stations, 16 out of the 23 environmental variables showed a significant

difference between the two seasons (Table 2, Figure 3). Main observed changes involved

variables related to herbaceous vegetation, according to plant coverage at different heights,

and vegetation species richness (VR). As expected, overall, woody life forms were more

stable between seasons (Table 2, Figure 3). Components of microhabitat showing no

statistical change between seasons were frequency of herbaceous plants at 10 cm (HF11),

frequency of woody plants at 35 cm (WF23) and 50 cm (WF24), as well as coverage of logs

(CLO), rocks (RC), and bare soil (BS); therefore, space configuration given by these

structural features remained stable in both seasons (Figure 3).

Microhabitat heterogeneity. The three different microhabitat types (M1, M2, and M3) within

the sampling plot in each season, as revealed by cluster analyses, revealed temporal habitat

heterogeneity in the entire plot (Figure A.1). In addition, Fisher's exact test of Contingency

Tables Analysis revealed significant changes ($p = 0.004$) in the number of sampling stations

assigned to each microhabitat type from the dry to the rainy seasons with main changes

occurring within M1 and M2. That is, M1 included more sampling stations during the dry

season $x^2 = 2.123$, $p < 0.05$), while M2 covered the majority of sampling stations during the

rainy season ($x^2 = 3.348$, $p < 0.05$), as its area increased by 24 sampling stations from the dry

to the rainy seasons (Figure 2, Figure A.1). Conversely, M3 remained quite stable ($x^2 =$

0.011), with almost the same number of sampling stations throughout the study (Figure 2).

Discriminant analyses validated heterogeneity of the three microhabitats (Figure 2)

within and between seasons. The percentage of variance associated with each discriminant

function was higher during the rainy season (dry season: F1 = 58.67 %, F2 = 42.32 %; rainy

season: F1 = 72.581 %, F2 = 27.419 %, Figure A.1). Discriminant functions explained

variation due to woody plants in the understory during the dry season, while herbaceous

vegetation together with decayed matter and logs became more relevant during the rainy

season (Table A.3). Wilk's Lambda showed that at least the mean of one microhabitat

differed statistically from the others either in the dry and rainy season ($p = 0.0001$ each,

respectively, Figure A.1). These results corroborated spatial and temporal heterogeneity

drawn from cluster analysis of individual station features in the grid plot. During the dry

season and rainy season, 93.33 % and 94.17% of individual sampling stations, respectively,

remained correctly classified (Figure 2).

Description of microhabitats within the plot

Microhabitat 1. M1 (Figure 2, Figure A.1) was characterized by little to no vegetation cover

at different heights (VC) and by a low frequency of herbaceous (HF) and woody (WF) plants.

In contrast, organic matter on the ground (OM) was the most frequent component. This

microhabitat was also distinctive for having extensive coverage of logs (CLO) on the ground

and for being the microhabitat with the most conspicuous presence of burrows (BW).

Microhabitat 2. M2 (Figure 2, Figure A.1) was the largest area covered by herbaceous life

forms at different heights (WF11-15); vegetation species richness (VR) and the total number

of herbaceous plants (TH) also remained very high. Woody life forms at 25 cm (WF22) were

present only at low frequencies, while WF24 (35 cm) and WF25 (100 cm) remained at higher

frequencies. Coverages on the ground of small logs or dead wood (CLO), rocks (RC), and

organic matter (OM) remained low during the study. Bare soil surface (BS) increased to

higher amounts during the dry season and decreased during the rainy season. There were no

burrows (BW) in this microhabitat.

Microhabitat 3. In M3 (Figure 2, Figure A.1), vegetation coverage showed high values at all

height levels (VC25, VC35, VC50, and VC100), except for the first (VC10), while vegetation

species richness (VR) averaged the highest, compared to other microhabitats. Total number

of woody plants (TW) and frequency of woody plants at different heights (WF21 - 25)

remained very high, while total number of herbaceous plants (TH) was low. Coverage of

rocks (RC), logs (CLO), and bare soil surface (BS) were low during the dry season.

Vegetation coverage at 10 cm (VC10) was also low during that season but increased in the

rainy season. Conversely, OM was present in the dry season but decreased in the rainy

season. There was a low number of burrows (BW).

Association between the microhabitats and species. Kernel Graphs (Figure 4) indicated that

the two species had a distinctive use of space, since they were differentially scattered along

the plot. *Peromyscus difficilis* was more abundant during the dry season, occupying a large

portion of the plot. *P. melanotis*, on the other hand, increased its distribution during the rains,

when it seemed to displace *P. difficilis* into other sampling stations (Figure 4). Distribution

rearrangements between seasons implied that one species occupied some stations more

frequently than the other, and vice versa (Figure 4). Indeed, the two x^2 tests yielded

statistically significant differences in abundance of each species among microhabitats and

between seasons (Table A.2: dry season; R^2 (U) = 0.09, n = 111, df = 2, Likelihood Ratio x^2 =

15.07, p = 0.00005; rainy season; R^2 (U) = 0.02, n = 168, df = 2, Likelihood Ratio x^2 = 6.40, p

= 0.0406).

During the dry season (Figure 5 and Table A.2), *Peromyscus difficilis* was highly

associated with M2 (80 % of captured mice) and somewhat less with M1 (64 % of captured

mice); conversely, *P. melanotis* was captured more frequently in M3 (66 % captured mice).

Spatial use of microhabitats changed for both species in the rainy season (Figure 5 and Table

A.2), also showing a microhabitat partition, though less conspicuous; *i. e.*, in the

Correspondence Analysis, 61 % of mice caught in M1 were *P. difficilis*. This species was

also related to M3, while 60 % of mice caught of *P. melanotis* were captured in M2.

Spatial Patterns. The Nearest Neighbor Analysis revealed a significantly clustered pattern

within the plot for each species in both seasons (Table 3). During the dry season, the mean

distance between individuals of *P. difficilis* was more clustered than for individuals of *P.*

melanotis. However, during the rainy season, the mean distance between individuals

increased in *P. difficilis*, but decreased in *P. melanotis* (Table 3).

During the dry season, Ripley's bivariate K showed a statistically significant pattern of

repulsion between the *Peromyscus* species in almost all analyzed distances of the entire plot,

except at 2 m where it showed an attraction pattern (Figure 6A). This repulsion pattern

reversed in almost all distances during the rainy season (Figure 6B), with the two species

becoming more associated, sharing microhabitats in almost all sampling stations. However,

during the rainy season, statistically significant peaks of repulsion reappeared between the

species at distances of five, ten, and 14 m (Figure 6B). These analyses also revealed the

intensity of these patterns; *e.g.,* the likelihood of finding individuals of *P. difficilis* and *P.*

melanotis together at the same capture station was very low during the dry season (Figure

6A), while this probability increased in the rainy season (Figure 6B).

Effects of microhabitat structural components on distribution. According to the best

Generalized Linear Model (AICc = 212.55, Table 5, Figure 5C), during the dry season PC1

and PC3 were the best predictors of capture frequency of *P. difficilis*. Environmental

variables with the highest scores in PC1 ($p = 0.0186$) were the total number of herbaceous

plants (TH), vegetation species richness (VR), and vegetation cover at 10 cm (VC10). In PC3

($p = 0.0001$, Table 4, 5, Figure 5C), characterizing variables were the number of logs on the

ground (CLO), and the presence of burrows (BW). These five variables were also the main

elements characterizing M2 (PC1) and M1 (PC3), the two habitats with the highest capture

frequencies of *P. difficils*. As for *P. melanotis* (Figure 5C), capture frequencies were

positively related to PC2 (AICc = 204.47, *p* = 0.0001), and among the four variables with the

highest PC scores were total number of woody plants (TW) and frequency of woody life

forms at 10 cm height (WF21), the two main elements of M3 during the dry season (Table 5,

Figure 5C).

During the rainy season, a similar pattern of microhabitat partition occurred between

the two species. The best GLM model for *P. difficilis* in this season (Table 5) showed effects

of PC2 and PC3. The most important environmental variables (AICc = 286.76, *p* = 0.0001,

Table 4, 5) in PC2 were total number of woody plants (TW), herbaceous forms at 50 cm

(HF14), woody forms at 10 and 100 cm (WF21, WF25, respectively), and number of logs on

the ground (CLO). In PC3, the variables were number of logs on the ground (CLO) and

organic matter (OM), which were associated with a higher capture frequency of *P. difficilis* in

M1 (PC3) and M3 (PC2) during the rainy season. On the other hand, *P. melanotis* (Figure

5C) was only associated with PC1 (AICc = 241.01, *p* = 0.0343), and the main characterizing

variables in this component were number of total herbaceous plants (TH), vegetation species

richness (VR), and plant cover at almost all analyzed heights (VC10 - 100 cm, Table 4).

446   **Discussion**

The results of our analyses show that as environmental conditions changed, the two syntopic

species of *Peromyscus* switched and split space resources. Our findings on the microhabitat

relationships between these two *Peromyscus*, dwelling in a patch of mid-latitude temperate

forest, concur with the habitat heterogeneity coexistence hypothesis (Cramer and Willig 2002;

Valladares *et al.* 2015). We found that both *Peromyscus* were sensitive to slight changes in

microhabitat structure and that each one of the three microhabitat types provided different

resources for each species. Our results also suggest that temporal differences in spatial

structure, and the associated shifts in food and shelter availability, facilitate coexistence

between these two syntopic *Peromyscus* (Pianka 1973; Schoener 1974). For instance, during

the dry season, *P. difficilis* was most associatedwith M1 and M2, while *P. melanotis* was

highly associated with M3. Conversely, in the rainy season, *P. difficilis* was more frequently

captured in M1 and M3, while *P. melanotis* was highly associated with M2. Partition in space

use has already been documented for coexisting species of *Peromyscus* within a community

(*i. e.,* niche partitioning by *Peromyscus leucopus* and *P. maniculatus*, Kaufman and Kaufman

1989), and several studies have demonstrated variation in spatial resource use by congeneric

species (Barry *et al.* 1990; Dooley and Dueser 1996; Kalcounis-Rüppell and Millar 2002).

In our study, *P. difficilis* remained highly associated with M1 in both seasons, but

especially so during the rainy season. There was a very high number of fallen logs at M1,

which represents small patches of microhabitat for small mammals with food sources, such as

plant items and invertebrates (Bellows *et al.* 2001), as well as sources for refuge and shelter

(Bowman *et al.* 2001). In this microhabitat, individuals of *P. difficilis* can also use the large

fallen logs as pathways for quick and straight locomotion within the forest (Bellows *et al.*

2001; Dewalt *et al.* 2003). Indeed, fallen logs promote structural complexity of forests and

may enhance positive interactions among species of small mammals (Bowman and Facelli

2013). GLM analyses showed that the main variables contributing to the abundance of *P.*

difficilis during the rainy season were herbaceous vegetation (TH), vegetation species richness

(VR), and the number of logs on the ground (CLO), among others. However, in the dry

season, the main variables contributing to the abundance of *P. difficilis* were woody

vegetation (TW), the number of logs on the ground, and organic matter (OM), among others.

Bellows *et al.* (2001) found a similar result in a high latitude temperate forest (Virginia,

USA), where distribution of a generalist small mammal was associated with the diameter of

fallen logs, the frequency of shrubs, and degree of canopy closure. Association between

rodents and fallen logs was also documented for *Nectomys squamipes*, which builds its nests

inside decomposed fallen logs (Briani *et al.* 2004). This has also been recorded for other

rodent species in several biomes with different vegetation types (*e. g.*, *Rattus rattus*, *Nesomys*

audeberti in Lehtonen *et al.* 2001; *Oligoryzomys nigripes* in Dalmagro and Vieira 2005). In

contrast, *P. melanotis* was only related with one PC in each season. For instance, during the

dry season, *P. melanotis* was associated with only PC2, a component representing mostly the

total number of woody plants (TW) and small woody forms (WF21, 10 cm height). During

the rainy season, the presence of *P. melanotis* was only related to PC1, which is characterized

by an understory dominated by herbaceous life forms (TH). Higher rodent frequency in

higher-density understory habitats has been interpreted as a strategy to avoid predation by

aerial hunters (Grelle 2003; Dalmagro and Vieira 2005; Johnson 2007).

Other intrinsic factors, such as the breeding seasons, are likely to be related to

coexistence of these two syntopic *Peromyscus*. For instance, the Nearest Neighbor Analysis

revealed that in both seasons each species showed a distinctive, significant clustered pattern

within the plot. Individuals of *P. difficilis* were more clustered together by mean distances

than individuals of *P. melanotis*, particularly during the dry season. Conversely, during the

rainy season, the mean distance between individuals increased in *P. difficilis* but decreased in

P. melanotis. These shifting dispersion patterns are species-specific and relate with their

respective breeding seasons (De-la-Cruz *et al*. 2019). The main breeding activity in *P.*

difficilis occurs during the dry season, while it is during the rains for *P. melanotis*.

In addition, species-specific needs for different resource requirements to fulfill the

breeding season, could affect the dispersion and coexistence between these two *Peromyscus*

in the study area. For instance, coexistence between the two *Peromyscus* may be improved

by temporal segregation of *P. melanotis*, compared to the resident behavior of *P. difficilis*.

(species). Indeed, *P. melanotis* behaved as an opportunistic species throughout our study area

since its capture frequencies increased when environmental conditions were more beneficial

during the rainy season but remained at low levels during the harsher dry season. This

smaller *Peromyscus* may migrate to areas with more availability of resources and better

environmental conditions to survive during the dry season, coming back to the area during the

rainy season to carry out the breeding season, when there is greater accessibility of food and

spatial resources, and when competition with *P. difficilis* may decrease. On the other hand, as

showed by Mohammadi (2010) in another small member of this genus, *Peromyscus leucopus*,

opportunistic habitat use may be related to predation risk, which in our study is also supported

for *P. melanotis* given its affinity for microhabitats that offer greater coverage against aerial

predators. Further studies of the spatial segregation and predation influence on the

opportunistic behavior of *P. melanotis* are needed.

Increase in number of captures for both species from the dry to the rainy seasons is

due to resource availability, but they are also due to the type of resources. For instance, *P.*

difficilis remained more abundant and always present at microhabitats with more stable

elements in the study area throughout the study, while the increase in abundance of *P.*

melanotis depended on denser plant cover during the rainy season. Since rains promote an

increase of primary productivity, allowing more resource availability (mainly food) and

enhancing microhabitat carrying capacity, such habitat changes facilitated coexistence

between both *Peromyscus*, and with other small land mammals in the area (Castro-Campillo

et al. 2008, 2012). Ripley's bivariate K supported and shed light on this result, since the two

species showed repulsion between them in almost all analyzed distances during the dry

season; indeed, during the dry season when resources were scarce, incidents of both species

using the same microhabitat were uncommon. Since the dry season is the primary breeding

season for the medium-sized *P. difficilis*, it is likely that this species displays territorial

behavior against its smaller-sized congeneric. Conversely, the intensity of this repulsion

decreased substantially during the rainy season, with the two species showing an association

pattern at some analyzed distances. This supports the idea that increased resource availability

during the rainy season allows the two species to share the habitat. Moreover, our results also

indicate that spatial dispersion patterns are seasonally variable in both *Peromyscus* (Brown

and Zeng 1989; Cramer and Willig 2002). In this sense, it is important to point out that in

both seasons Ripley's bivariate K indicated an attraction pattern between the two species

within 2 and 3 m, which is more likely an artefact of our sample design than a biological

result. This is because if an individual of *P. difficilis* was captured during the first of our two

sampling nights, followed by capture of one individual of *P. melanotis* in that same station the

next day, the analysis counted the pattern as an association because they were trapped in the

same station at a very short distance.

Also, partition of space by these syntopic deermice must be facilitated by their

respective locomotive habits- semiarboreal in *P. difficilis* and cursorial in *P. melanotis*. The

long tail of *P. difficilis* enables it to rush and climb along shrubs or trees (Fernandez *et al*.

2010), thus increasing its preference for habitats with these and other fixed (*e. g.,* fallen logs

and twigs) elements where it can escape from predators. In fact, adult coat color changes in

P. difficilis between seasons, becoming more similar to ground litter, suggesting that in this

resident deermouse, phenotype plasticity in color is a cryptic response elicited by predation

risk. In contrast, smaller body size, together with a shorter tail and narrower soles (Álvarez-

Castañeda 2005), should enable the cursorial *P. melanotis* to occupy such zones as M2 and

M3, where predators cannot easily spot it through dense vegetation cover, so it can scape very

quickly. Indeed, coverage by high shrubs provides both protection from predators and food

sources, since seeds may be concentrated under shrub canopies (Thompson 1982;

Mohammadi 2010). In fact, rodents usually avoid foraging in unsheltered microhabitats and

forest edges where they are more likely to be spotted by avian (Kotler *et al*. 1991) and other

vertebrate predators (Morris and Davidson 2000; Mohammadi 2010).

Finally, several aspects of our methods for sampling the environmental features of

microhabitat, as well as the quantitative analyses used here, especially Ripley's bivariate K,

make it particularly informative in studies of spatial dynamics of dispersion in small rodents

at a microhabitat scale. Our explicitly designed methodological approach, focused on small-

sized rodents with low vagility, provided us with necessary information about the ability of

these two *Peromyscus* species to split resources in the complex understory of a mid-latitude

temperate forest. Our initial aim to reconstruct microhabitat structure also successfully

provided us with clues as to how to eventually manage perturbed ecosystems for conservation

purposes, such as those at the edge of a megalopolis, whose continuous growth produces

fragmentation of natural microhabitats. We think that our methodology will also be helpful in

other scenarios for understanding the ecology of small rodents, such as behavioral dynamics,

activity patterns, and reproductive systems at intra- and/or inter-population levels.

568   **Acknowledgments**

We thank C. Peralta-Juarez and J. Patiño-Ortega for their friendly fieldwork assistance and

logistic support. The staff at UAMI mammal scientific collection kindly provided all the

facilities for specimen curation and housing. Authorities in charge of DLNP (CORENA,

SEMARNAT), as well as the forest rangers and administration staff at the National Park gave

us necessary information, security, and logistics to complete our studies in the field. This

paper contains partial information from the Masters in Biology thesis (DCBS, UAM-I) of

IMDA; IMDA received financial support through a fellowship (283799, CVU 479479) from

the Consejo Nacional de Ciencia y Tecnología (CONACyT). This study was supported by

annual grants (143.***.46 IB-DCBS-UAMI, 2013-2017) to AACC.

579   **References**

Álvarez-Castañeda, S. T. 2005. *Peromyscus melanotis*. Mammalian Species 764:1-4.

Barry Jr, R. E., A. A. Heft., and T. E. Baummer. 1990. Spatial relationships of syntopic

white-footed mice, *Peromyscus leucopus*, deer mice, *Peromyscus maniculatus*, and red-

backed voles, *Clethrionomys gapperi*. The Canadian Field-Naturalist 104:87-393.

Bellows, A. S., J. F. Pagels., and J. C. Mitchell. 2001. Macrohabitat and microhabitat

affinities of small mammals in a fragmented landscape on the upper coastal plain of

Virginia. The American Midland Naturalist 146:345-360.

Briani, D. C., A. R., Palma, E. M. Vieira., and R. P. Henriques. 2004. Post-fire

succession of small mammals in the Cerrado of central Brazil. Biodiversity and

Conservation 13:1023-1037.

Brown J. H and Z. Zeng. 1989. Comparative population ecology of eleven species of

rodents in the Chihuahuan Desert. Ecology 70:1507-1525.

Bowman, J., G. Forbes., and T. Dilworth. 2001. Landscape context and small-mammal

abundance in a managed forest. Forest Ecology and Management 140:249-255.

Bowman, A. S., and J. M. Facelli. 2013. Fallen logs as sources of patchiness in chenopod

shrublands of South Australia. Journal of Arid Environment 97:66-72.

Bowne, D. R., J. D. Peles., and G. W. Barrett. 1999. Effects of landscape spatial structure

on movement patterns of the hispid cotton rat (*Sigmodon hispidus*). Landscape Ecology

14: 53-65.

Canfield, R. H. 1941. Application of the line interception method in sampling range

vegetation. Journal of Forestry 39:388-394.

Castro-Campillo, A., A. Salame-Méndez., A. Vergara-Huerta., A. Castillo., and J.

Ramírez-Pulido. 2008. Fluctuaciones de micromamíferos terrestres en bosques

templados aledaños a la ciudad de México. pp. 391-410 in Avances en el Estudio de los

Mamíferos de México (Lorenzo C., E. Espinoza and J. Orteg, eds.). Publicaciones

especiales. Vol. II, Asociación Mexicana de Mastozoología, A. C., México, CDMX.

Castro-Campillo, A., L. León-Altamirano., J. Herrera-Muñoz., I. Salgado-Ugarte., E.

Mendieta Márquez., J. L. Contreras-Montiel., H. Serrano., J. Ramírez-Pulido., and

A. Salame-Méndez. 2012. Is there a difference between T contents in two populations

of the Black-eared Mouse, living under similar conditions but with differences in two

population patterns? Acta Zoológica Mexicana (n. s.) 28:525-539.

Clark, P. J., and F. C. Evans. 1954. Distance to nearest neighbor as a measure of spatial

relationships in populations. Ecology 35:445-453

Cramer, M. J., M. R. Willig. 2002. Habitat heterogeneity, habitat associations, and rodent

species diversity in a sand-shinnery-oak landscape. Journal of Mammalogy 83:743-753.

CONANP. 2006. Programa de Conservación y Manejo Parque Nacional Desierto de los

Leones. Dirección General de Manejo para la Conservación, Comisión Nacional de

Áreas Naturales Protegidas, Secretaria de Medio Ambiente del Gobierno del Distrito

Federal, Secretaria del Medio Ambiente y Recursos Naturales.

http://www.sma.df.gob.mx/

Coppeto, S. A., D. A. Kelt., D. H. Van Vuren., J. A. Wilson., S. Bigelow. 2006. Habitat

associations of small mammals at two spatial scales in the northern Sierra Nevada.

Journal of Mammalogy 87:402-413.

Currie, D. J. 1991. Energy and large-scale patterns of animal and plant species richness.

American Naturalist 137:27-49.

Chesson, P. 2000. Mechanisms of maintenance of species diversity. Annual Review of

Ecology, Evolution, and Systematics 31:343-366.

Dale, M. R. T. 1999. Spatial pattern analysis in plant ecology. Cambridge University Press.

Dalmagro, A. D., and E. M. Vieira. 2005. Patterns of habitat utilization of small rodents in

an area of Araucaria forest in Southern Brazil. Austral Ecology 30:353-362.

De-la-Cruz I. M., A. Castro-Campillo., A. Zavala-Hurtado., A. Salame-Méndez., and J.

Ramírez-Pulido. 2019. Differentiation pattern in the use of space by males and females

of two species of small mammals (*Peromyscus difficilis* and *P. melanotis*) in a temperate

forest. Therya 10:3-10.

DeWalt, S. J., S. K. Maliakal., J. S. Denslow. 2003. Changes in vegetation structure and

composition along a tropical forest chronosequence: implications for wildlife. Forest

Ecology and Management 182:139-151.

Diggle, P. J. 1983. Statistical Analysis of Spatial Point Pattern. London: Academic Press.

Dooley, J. L., and R. D. Dueser. 1996. Experimental tests of nest site competition in two

Peromyscus species. Oecologia 105:81-86

Dooley, J. L., and M. A. Bowers. 1998. Demographic responses to habitat fragmentation:

experimental tests at the landscape and patch scale. Ecology 79:969-980.

Fahrig, L., and G. Merriam. 1994. Conservation of fragmented populations. Conservation

Biology 8:50-59.

Fernández, J. A., F. García-Campusano., and M. S. Hafner. 2010. *Peromyscus difficilis*

(Rodentia: Cricetidae). Mammalian Species 42:220-229.

Grelle, C. E. V. 2003. Forest structure and vertical stratification of small mammals in a

secondary Atlantic forest, southeastern Brazil. Studies on Neotropical Fauna and

Environment 38:81-85.

Haim, A., and F. M. Rozenfeld. 1993. Temporal segregation in coexisting *Acomys* species:

the role of odour. Physiology & Behavior 54:1159-1161.

Hammer, Ø., D. A. T. Harper., and P. D. Ryan. 2001. PAST: PAleontological STatistics

software package for education and data analysis. Palaeontologia Electronica 4:1-9.

Johnson, M. L., and M. S. Gaines. 1990. Evolution of dispersal: theoretical models and

empirical tests using birds and mammals. Annual Review of Ecology, Evolution, and

Systematics 21:449-480.

Johnson, M. D. 2007. Measuring habitat quality: a review. Condor 109:89-504.

Jorgensen, E. E. 2004. Small mammals' use of microhabitat reviewed. Journal of

Mammalogy 85:531-539.

Kalcounis-Rüppell, M. C., and J. S. Millar. 2002. Partitioning of space, food, and time by

syntopic *Peromyscus boylii* and *P. californicus*. Journal of Mammalogy 83:614-625.

Kaufman, D. W., and G. A. Kaufman. 1989. Population biology. pp. 233-271 *in*

Advances in the study of *Peromyscus* (Rodentia) (Kirkland G., and J. Layne, eds.).

Texas Tech University Press, Lubbock.

Kotler, B. P., J. S. Brown and O. Hasson. 1991. Factors affecting gerbil foraging behavior

and rates of owl predation. Ecology 72:2249-2260.

Lehtonen, J. T., O. Mustonen., H. Ramiarinjanahary., J. Niemelä., and H. Rita. 2001.

Habitat use by endemic and introduced rodents along a gradient of forest disturbance in

Madagascar. Biodiversity and Conservation 10:1185-1202.

M'Closkey, R. T., and B. Fieldwick. 1975. Ecological separation of sympatric rodents

(*Peromyscus* and *Microtus*). Journal of Mammalogy 56:119-129.

M'Closkey, R. T., and D. T. Lajoie. 1975. Determinants of local distribution and

abundance in White-Footed Mice. Ecology 56:467-472.

Mohammadi, S. 2010. Microhabitat selection by small mammals. Advances in Biological

Research 4:283-287.

Monamy, V., and B. J. Fox. 1999. Habitat selection by female swamp rats (*Rattus*

lutreolus) drives asymmetric competition and coexistence with long-tailed mice

(*Pseudomys higginsi*). Journal of Mammalogy 80:232-242.

Morris, D. W. 1984a. Sexual differences in habitat use by small mammals: evolutionary

strategy or reproductive constraint? Oecologia 65:51-57.

Morris, D. W. 1984b. Patterns and scale of habitat use in two temperate-zone, small

mammal faunas. Canadian Journal of Zoology 62:1540-1547.

Morris, D. W. 1987. Ecological scale and habitat use. Ecology 68:362-369.

Morris, D. W., and L. D. Davidson. 2000. Optimally foraging mice match patch use with

habitat differences in fitness. Ecology 81:2061-2066.

Pianka, E. R. 1973. The structure of lizard communities. Annual Review of Ecology,

Evolution, and Systematics 4:53-74.

Posada, D., and T. R. Buckley. 2004. Model selection and model averaging in

phylogenetics: advantages of Akaike Information Criterion and Bayesian Approaches

Over Likelihood Ratio Tests. Systematic Biology 53:793-808,

Ramírez-Pulido, J., I. Lira., S. Gaona., C. Müdespacher., and A. Castro-Campillo. 1989.

Manejo y mantenimiento de colecciones mastozoológicas. Universidad Autónoma

Metropolitana, Unidad Iztapalapa, México, D.F.

Ripley, B. D. 1976. The second-order analysis of stationary point processes. Journal of

Applied Probability 13:255-266.

Rosenberg, M. S., and C. D. Anderson. 2011. PASSaGE: pattern analysis, spatial

statistics, and geographic exegesis. Version 2. Methods in Ecology and Evolution 2:229-

232.

Schoener, T. W. 1974. Resource partitioning in ecological communities. Science 185:27-

39.

Schreiber, S., and P. T. Killingback. 2013. Spatial heterogeneity promotes coexistence of

rock-paper-scissor metacommunities. Theoretical population biology 86:1-11

Seagle, S. W. 1985. Patterns of small mammal microhabitat utilization in cedar glade and

deciduous forest habitats. Journal of Mammalogy 66:22-35.

Sikes, R. S. 2016. Guidelines of the American Society of Mammalogists for the use of wild

mammals in research and education. Journal of Mammalogy 97:663-688,

Thompson, S. D. 1982. Structure and species composition of desert heteromyid rodent

species assemblages: effects of a simple habitat manipulation. Ecology 63:1313-1321.

Upton, G. J. G., and B. Fingleton. 1985. Spatial data analysis by example. Wiley,

Chichester; New York.

Valladares, F., C. C. Bastias., O. Godoy., E. Granda., and A. Escudero. 2015. Species

coexistence in a changing world. Frontiers in Plant Science 6, art. 866.

Vázquez, J., and S. T. Álvarez-Castañeda. 2011. Spatial relationships between burrows of

an insular population of *Dipodomys merriami*. Mammalian Biology 76:577-582.

V**illanueva-Hernández, A. I., D. A. Delgado-Zamora., S. A. Heynes-Silerio., L. Ruacho-**

González., and C. López-González. 2017. Habitat selection by rodents at the transition

between the Sierra Madre Occidental and the Mexican Plateau, México. Journal of

Mammalogy 98:293-301.

Ward, J. H. 1963. Hierarchical grouping to optimize an objective function. Journal of the

American Statistical Association 58:236-244.

Wiens, J. A. 2000. Ecological heterogeneity: an ontogeny of concepts and approaches. pp

9-31 in The Ecological Consequences of Environmental Heterogeneity (Stewart A. J. A.,

E. A. John., and M. J. Hutchings, eds). Blackwell, Oxford.

Zar, J. H. 1999. Biostatistical Analysis. Prentice Hall, Princeton.

Zavala-Hurtado, J. A., P. L. Valverde., M. C. Herrera-Fuentes., and A. Diáz-Solis.

2000. Influence of leaf-cutting ants (*Atta mexicana*) on performance and dispersion

patterns of perennial desert shrubs in an inter-tropical region of Central Mexico. Journal

of Arid Environments 46:93-102.

Zavaleta, E. S., M. R. Shaw., N. R. Chiariello., H. A. Mooney., and C. B. Field. 2003.

Additive effects of simulated climate changes, elevated CO2, and nitrogen deposition on

grassland diversity. Proceedings of the National Academy of Science 100:7650-7654.

 Figure legends

Figure 1. Sampling plot design (A). Total plot area was 2475 m^2; each intersection (dots, n

= 120) was an independent sampling station for capturing mice. Influence area of 2.5 m^2

around each station sampling (central green triangle) (B), where 23 environmental variables

of microhabitat were sampled along eight transects (see text).

Figure 2. Distribution maps for microhabitats (M1 = red, M2 = green, M3 = blue) within the

study quadrant for the dry (A) and rainy (B) seasons. Bar graph showing number shifts of

microhabitat types in the dry and rainy seasons (C). Asterisks in M1* and M2* refer to p =

0.001.

Figure 3. Seasonal heterogeneity between the dry and rainy seasons as evaluated using 23

environmental variables sampled in the study plot. Overall, vegetation variables increased

during the rainy season, while organic matter (OM) decreased. Fixed elements, such as logs

on the ground (CLO), showed no shifts between seasons. Variables with low values in the

study plot (*e. g*., RC) are depicted with red crosses (mean value).

Figure 4. Spatial analysis with Kernel density points showing capture frequencies during the

dry and rainy seasons for *P. difficilis* (A and B respectively) and *P. melanotis* (C and D,

respectively). Dots depict actual capture points, while the color refers to the density of

captures, with red color indicating the highest frequencies (hot spots). Numbers on each axis

indicate actual distances (m).

Figure 5. Correspondence analysis from contingency tables of capture frequency data for

each deermouse (*P. difficilis* and *P. melanotis*) in each microhabitat (M1 - 3), during the dry

(A) and rainy (B) seasons. Axes c1 and c2 indicate coordinates resulting from the ordination

of each variable in the analysis. Plots for the GLMs models (C), showing predicted

frequencies for each species, according to a determined principal component (model effect).

All fitted PCs showed a positive relationship in the model, indicating that the probability of

capture frequency of individuals of certain species, increased with the presence of the

environmental variables within parenthesis (those having the highest loading scores).

Figure 6. Ripley's bivariate K Analysis for the interaction between the two *Peromyscus*. (A)

shows a repulsion pattern between both species in all analyzed distances during the dry

season. (B) Shows attraction from 6 - 12 m and repulsion at 5, 10, and 14 m, between the two

species during the rainy season. The solid black line represents Ripley's K bivariate pattern;

red dashed lines depict a 95 % CI; solid blue line shows the null hypothesis. See Methods.

Variable code	Predation		Resources				
	Diminished risk	Refuges	Breeding Space	Food	Temporal food and refuges	Permanent	Resting Places
Vegetation Cover (cm)							
VC10	X			X	X		X
VC25	X		X	X	X		X
VC35	X		X	X	X		X
VC50	X		X	X	X		X
VC100	X		X	X	X		X
Frequency of Herbaceous Plants (cm)*							
HF11	X		X	X	X		X
HF12	X		X	X	X		X
HF13	X		X	X	X		X
HF14	X		X	X	X		X
HF15	X		X	X	X		X
Frequency of Woody Plants (cm)*							
WF21	X	X	X	X	X	X	X
WF22	X	X	X	X	X	X	X
WF23	X	X	X	X	X	X	X
WF24	X	X	X	X	X	X	X
WF25	X	X	X	X	X	X	X
Species Vegetation Richness							
VR	X	X	X	X	X	X	X
Total of Herbaceous Plants							
TH	X		X	X	X		X
Total of Woody Plants							
TW	X	X	X	X	X	X	X
Rocks							
RC	X	X	X			X	X
Organic Matter (litter, twigs, small trunks)							
OM				X	X		
Bare Soil							
BS					X		
Coverage of Logs (stumps, fallen trunks)							
CLO	X	X	X			X	X
Numbers of Burrows							
BW	X	X	X			X	X

* = Frequencies of plant forms (HF, WF) involve the same heights as in VC.

831 **Table 2.** Normality (**NOR**) tests for 23 variables (**VAR**) of microhabitat, and comparisons[abc]

832 (**COM**) for average values between the dry (**DRY**) and rainy (**RAINS**) seasons, respectively.

833 Variable names as in Table 1.

VAR	NOR[a]		COM	VAR	NOR[a]		COM	VAR	NOR[a]		COM
	DRY	RAINS			DRY	RAINS			DRY	RAINS	
BS	0.0001	0.0001	*1.0*[c]	VC100	0.002	0.003	0.0001[c]	TW	*0.143*	*0.08*	0.0004[b]
CLO	0.0001	0.0001	*0.972*[c]	VR	0.006	*0.082*	0.0001[b]	WF21	0.0001	0.0001	0.004[c]
OM	0.010	0.034	0.0001[c]	TH	0.0001	*0.072*	0.0001[b]	WF22	0.0001	0.0001	0.026[c]
RC	0.0001	0.0001	*0.432*[c]	HF11	0.015	0.028	*0.658*[c]	WF23	0.0001	0.0001	*0.875*[a]
VC10	0.002	0.041	0.0001[c]	HF12	0.0001	0.0001	0.001[c]	WF24	0.0001	0.0001	*0.378*[c]
VC25	0.009	0.003	0.0001[c]	HF13	0.0001	0.0001	0.0001[c]	WF25	0.0001	0.001	0.0001[c]
VC35	0.0001	0.002	0.0001[c]	HF14	0.0001	0.0001	0.0001[c]	BW	0.0001	0.0001	1
VC50	0.0001	0.019	0.0001[c]	HF15	0.0001	0.0001	0.0001[a]				

836 Parametric: [a]Kolmogorov-Smirnov; [b]Student for two samples; non-parametric: [c]Wilcoxon). The *p*-values in italics do not

837 reject normality ($p = 0.05$), or do not indicate significant differences between pluvial seasons.

850 **Table 3.** Nearest Neighbor Analysis for the species of *Peromyscus*, showing the mean

851 distance at what individuals of each species are dispersed among each other. In all instances,

852 individuals of either species had a clustered point pattern. R - values indicate how the

853 observed distribution deviates from random. Clustered points give $R < 1$, Poisson patterns

854 give $R \sim 1$, while overdispersed points give $R > 1$.

	Dry season		Rain season	
	P. difficilis	*P. melanotis*	*P. difficilis*	*P. melanotis*
Points	57	39	58	52
Mean distance	1.40	2.39	1.67	2.09
R	0.42	0.77	0.55	0.64
P value	1.1263×10^{-16}	0.006	8.6577×10^{11}	8.7708×10^{7}
Point pattern	Clustered	Clustered	Clustered	Clustered

869   **Table 4.** Principal components (PC1 - 3) used as effect model in the GLMs analysis for both

870   seasons. Variable names as in Table 1.

Variables	Dry season (38.61 % of explained variance)			Rainy season (45.92 % of explained variance)		
	PC1	PC2	PC3	PC1	PC2	PC3
VC10	0.713	-0.388	-0.134	0.659	-0.179	-0.374
RC	-0.145	-0.055	-0.110	0.014	0.062	0.034
CLO	0.030	0.118	0.828	-0.153	0.411	-0.583
OM	-0.604	0.303	-0.487	-0.499	-0.122	0.775
BS	0.130	-0.187	0.112	-0.028	0.154	-0.328
VC25	0.415	0.280	0.034	0.658	-0.061	0.237
VC32	0.287	0.184	0.052	0.580	-0.003	-0.126
VC50	0.523	0.107	0.078	0.669	-0.173	-0.030
VC100	0.335	0.408	0.052	0.656	-0.053	0.107
VR	0.762	0.251	-0.269	0.776	0.380	0.058
TH	0.837	-0.430	-0.131	0.874	-0.319	-0.076
TW	0.520	0.684	-0.168	0.615	0.711	0.270
HF11	0.784	-0.394	-0.214	0.678	0.007	-0.228
HF12	0.573	-0.345	0.176	0.548	-0.278	0.256
HF13	0.247	0.073	0.113	0.463	-0.260	0.067
HF14	0.149	-0.186	-0.023	0.580	-0.530	-0.043
HF14	0.080	0.117	-0.236	0.481	-0.267	-0.019
WF21	0.127	0.510	-0.394	0.342	0.603	0.311
WF22	0.015	0.064	-0.105	0.266	0.212	0.416
WF23	0.199	0.256	-0.228	0.196	0.366	-0.241
WF24	0.424	0.359	0.195	0.337	0.165	0.159
WF25	0.515	0.366	0.380	0.494	0.469	0.026
BW	-0.105	0.249	0.557	-0.170	0.332	-0.218

Table 5. Generalized Linear Models constructed with the Principal Components (PC1 - 3,
habitat resources) and frequencies of *P. difficils* and *P. melanotis* for each pluvial season.
Resource partitions are observed in both species and seasons, since each PC relates to one
microhabitat (M1 - M3); note how PC1 - 3 explain capture frequencies for each species
within each season. Tests of Pearson and Deviance for fit goodness of are also shown.

Term	Estimate	Std Error	L-R x^2	P value	Pearson	Deviance
			Peromyscus difficilis			
			Dry season			
Intercept	-0.97	0.15	55.73	0.0001	$x^2 = 155.80$ ($p = 0.0096$)	$x^2 = 123.56$ ($p = 0.3210$)
PC1 = M2	0.11	0.05	5.53	0.0186	AICc = 212.55	
PC3 = M1	0.45	0.06	44.73	0.0001		
			Rainy season			
Intercept	-0.4	0.11	14.37	0.0002	$x^2 = 168.56$ ($p = 0.0008$)	$x^2 = 156.43$ ($p = 0.0062$)
PC2 = M3	0.23	0.06	10.92	0.0009	AICc = 286.76	
PC3 = M1	0.26	0.07	10.95	0.0009		
			Peromyscus melanotis			
			Dry season			
Intercept	-1.21	0.18	74.28	0.0001	$x^2 = 237.65$ ($p = 0.0001$)	$x^2 = 138.70$ ($p = 0.0935$)
PC2 = M3	0.46	0.09	23.34	0.0001	AICc = 204.47	
			Rainy season			
Intercept	0.6	0.12	28.15	0.0001	$x^2 = 132.71$ ($p = 0.13$)	$x^2 = 132.72$ ($p = 0.13$)
PC1 = M2	0.1	0.04	4.47	0.0343	AICc = 241.01	

906
907

915   **Figure 1**

   Figure 2

Figure 3

Figure 4

Figure 5

Figure 6