TITLE PAGE

Intracellular FMRpolyG-HSP70 complex: Possible use as biochemical marker of Fragile X Tremor Ataxia Syndrome

Authors: Giuseppe Bonapace[1*], Rosa Gullace[1], Daniela Concolino,[1] Grazia Iannello,[2,3] Radha Procopio,[2,3] Monica Gagliardi, [2] Gennarina Arabia,[3] Gaetano Barbagallo,[3] Angela Lupo,[3] Lucia Ilaria Manfredini,[3] Grazia Annesi,[2] Aldo Quattrone.[2,3]

[1] University Magna Graecia, Faculty of Medicine, Pediatrics

[2] Neuroimaging Research Unit, IBFM/CNR; Institute of Neurology,

[3] Institute of Neurology, University "Magna Graecia" of Catanzaro, Department of Medical Sciences

[*] **Corresponding author**:

Giuseppe Bonapace

Dept. of Pediatrics

University Magna Graecia

Campus S. Venuta

Catanzaro Italy 88100

Phone: +39 09613694127

bonapace@unicz.it

KEYWORDS

Toxic aggregates

FXTAS

Heat Shock proteins

Biochemical Marker

HSP70

INTRODUCTION

Fragile X-associated tremor/ataxia syndrome (FXTAS) is a late-onset neurodegenerative disorder that affects about 25% of carriers of non-coding CGG-repeat expansions (55–200 repeats) in the premutation range within the fragile X gene (*FMR1*). Main clinical features include intention tremor, cerebellar ataxia, and Parkinsonism [1]. Recently, great emphasis on the pathogenetic role of toxic aggregates, produced by a RAN translation process on the 5' expanded CGG region, has been given [2,3]. These aggregates contain a small peptide with a polyglycin stretch on the aminoterminal ends, (FMRpolyG), and so far, have been isolated and characterized in drosophila, mouse model and post mortem FXTAS patients biopsies, but never in living patients [4]. The early diagnosis of this syndrome, in adult carriers of premutations, remains difficult due to the absence of specific markers and, the complexity of the procedure aimed at producing human cellular models, makes the study of the role of these aggregates in the pathogenesis of FXTAS difficult as well.

Here we demonstrate, for the first time, using immunocystochemistry (ICC) and Western Blot procedure (WB), that it is possible to easily assess the presence of aggregates containing FMRpolyG protein, in fibroblasts isolated from a clinical and molecular characterized FXTAS living patient, and discuss some possible practical implications.

MATERIALS AND METHODS

Case report

The patient was a 50-year male with a history of tremor and slowness since age 43. He also complained of occasional syncopal episodes. His past and family histories were unremarkable. Neurological examination revealed broad-based gait, facial hypomimia and mild asymmetric (right greater than left) resting tremor in the distal upper and lower extremities, in association with postural and action tremor in the upper limbs. Tone, strength, and sensation were normal. There were also a slight bradykinesia and increased deep tendon reflexes. General physical examination was normal, without sexual or urinary disturbances, except for orthostatic hypotension with a decrease in systolic and diastolic blood pressure of 34 and 14 mmHg respectively, changing position from supine to standing upright. A generalized cerebral atrophy with severe involvement of the cerebellum and of the middle cerebellar peduncles were present in 3T MRI.

Gene Scan analysis

Amplicons depending on the size of the CGG repeat region, were produced using a specific FRAXA Triple Primed PCR Kit (Abbott Molecular). Hi-Di formamide, (Applied Biosystems), and 2μl of ROX 350 size standard (Celera Corporation) were combined with 2μl of PCR product. Samples were denatured at 95°C for 2 minutes before loading onto an ABI 310 with POP-6 polymer on a 36-cm or 50-cm array. Data were analyzed using GeneMapper v.3.7 software.

Immunocytochemistry

Patient's and control fibroblasts from skin biopsies, at confluence, were plated in chamber slide (Nunc Rochester NY) and incubated 48 h in RPMI 10% FBS. After paraformaldehyde fixation, permeabilization with 0.2% Triton X-100 followed by blocking in 0,2% gelatin at 4°C, cells were incubated for 3h at room temperature with a anti FMRpolyG monoclonal Antibody (clone 2J7 Millipore USA) and a goat anti Human HsP70 1:200 (Santa Cruz). Anti-mouse FITC conjugated and anti-goat TRITC conjugated specifc secondary antibody 1:500 were used for the detection. Hundred microscopic fields were analyzed and cells containing aggresomes (positive) were counted and evaluated as percentage of the total cells.

Immunoprecipitation studies

After precleaning with 60 μl of a Staphylococcus A membrane, the fibroblast' cytosolic fraction, was isolated and incubated overnight at 4°C with an anti FMRpolyG monoclonal antibody. The immunocomplexes were separated by SDS page and Western blot and analyzed by using a goat anti human Hsp70 and a rabbit antigoat HRP conjugated. The same anti Hsp70 antibody was used to assess the amount of free Hsp70 (without immunoprecipitation) in fibroblasts from normal control and patients.

RESULTS

The molecular analysis of the FMR1CGG repeats shows in our patient a mosaicism for a

44 and 116 CGG expansions in the premuation range for FRAXA

(Fig 1A).

To assess the production of a toxic FMRpolyG protein we performed

immunocytochemistry experiments both on FXTAS patient's and normal control's

fibroblasts.

Fig 1B shows that the FMRpolyG accumulates in patient's fibroblasts both in the

cytoplasm and nucleus (yellow arrow). As expetcted in presence of toxic soluble

aggregates, there is a clear colocalization of FMRpolyG with the Heat Shock protein 70

(HSP70), a well known cellular chaperone involved in the protection against the toxic

effect triggered by aggregosomes. (S.I.)

To demonstrate that the HSP70 does not merely colocalize with FMRpolyG, but it is a

component of the same cellular complex and involved in a defensive reaction against the

toxic aggregates, we performed immunoprecipitation experiment on the cytosolic

fibroblast fraction.

Fig 1C left, shows that the anti FMRpolyG antibody is able to immunoprecipitate the

HSP70 protein, confirming that there is a specific physical interaction with FMRpolyG

as expected in the first step of proteosomal mediated degradation pathway. [5]

Western blot only for the HSP70 protein on the same amount of cellular extract is shown

on the right side of the figure. According to the defense response, the total HSP70 levels

are increased in the FXTAS fibroblasts in comparison to the steady state levels in normal

controls.

DISCUSSION

Features of FXTAS overlap many other neurodegenerative disorders, including Parkinson, Alzheimer diseases and frontotemporal dementia (e.g., progressive cognitive impairment, altered mood and behavior). [6] These observations underscore the need to develop a new and reliable diagnostic tool, to confirm the diagnosis of FXTAS. Todd and colleagues recently demonstrated as a translation of CGG repeats without ATG (RAN) occurs in two out of the three frames, giving rise to short proteins containing either a polyalanine or a polyglycine stretch and that polyglycine protein resulted in the formation of heterogeneous protein inclusions, which were toxic both in neuronal transfected cells and in Drosophila . [3]

Based on this observation we speculated that would have been possible to exploit these FMRpolyG aggregate as possible diagnostic marker for FXTAS in living patients. Starting from the observation that the FMR1 protein is expressed in a wide range of cells, [7] we demonstrate, the presence *in vivo*, of FMRpolyG aggregates in fibroblasts from a patient carrier of a premutated FMR1 allele associated to clinical signs of FXTAS. In our knowledge this is the first observation in a living patient.

Because of the presence of the HsP70, a protective cellular chaperone against toxic aggregates, has been described as marker of toxicity, in the inclusions found in neurons and astrocytes from FXTAS post mortem brain, [8], we also studied such reaction in the fibroblasts of our patient. Our data demonstrating an increase in the amount of total Hsp70, along with an evident molecular interaction between the expanded peptide and

the HsP70, confirm in the patient, the cellular toxicity of the aggregates and the presence of a protective cellular response. Even though the validation of this procedure needs larger research to assess its predictive value for the clinical outcome of FMR1 premutation carriers without signs of FXTAS, these data demonstrate that it is possible to assess the presence of toxic HSP70-FMRpolyG aggregates responsible for FXTAS, in skin biopsies from living patients, establish a new simple, fast and reproducible human cellular model to study in *vivo* the progression of the disease and suggest that in a close future this procedure can be exploited to confirm the clinical diagnosis in patients showing neurological signs of FXTAS associated to a CGG premutated allele.

AKNOWLEDGMENTS

Author Contributions:

Giuseppe Bonapace, Study concept and design, immunohistochemistry and WB analysis

Rosa Gullace, Cell Colture and WB experiments

Daniela Concolino, Analysis and interpretation of data

Grazia Iannello, Molecular Biology data

Radha Procopio, Molecular Biology data

Monica Gagliardi, Fragment analysis

Grazia Annesi, Study concept and design, critical revision of manuscript

Gennarina Arabia, Draft paper, clinical data collection

Gaetano Barbagallo, Draft paper and skin biopsy

Angela Lupo, Clinical data collection

Lucia Ilaria Manfredini, Clinical data collection

Aldo Quattrone, Study supervision, Critical revision of manuscript for intellectual content

CONFLICT OF INTEREST

All the authors listed above report no financial disclosure /conflict of interest related to the manuscript.

REFERENCES

[1] Hall DA, Birch RC, Anheim MA et al. Emerging topics in FXTAS. J Neurodev Dis, 2014; 6: 31-40

[2] Usdin K, Hayward BE, Kumari D. et al. Repeat mediated genetic and epigenetic changes at FMR1 locus in Fragile X-related disorders. Front Genet 2014 5: 226-242.

[3]Kearse GM, Todd K. Repeat-Associated non-AUG translation and its impact in neurodegenerative disease. Neurotherapeutics .2014; 11: 721-731.

[4] Buijsen AMR, Sellier C, Severijnen LA,et al. FMR-polyG-positive inclusions in CNS and non-CNS organs of fragile X premutation carrier with fragile X-associated tremor/ataxia syndrome. Acta Neuropathologica Communications 2014; 2: 162-163

[5] Shiber A, Ravid A. Chaperoning protein for destruction: Diverse roles of HSP70 chaperones and their cochaperones in targeting misfolded proteins to the proteasome. Biomol. 2014; 4: 704-724

[6] Fraint A, Vittal P, Szewka A, et al. New observations in the fragile X-associated tremor/ataxia syndrome (FXTAS) phenotype. Front Genet. 2014; 5:365. [PubMed: 25368631]

[7] Thul PJ, Åkesson L, Wiking M, et al. A subcellular map of the human proteome. Science. 2017; DOI: 10.1126/science.aal3321

[8] Iwahashi C.K., Yasui D.H., An H.J et al. Protein composition of intranuclear inclusions of FXTAS. Brain 2006; 129: 256-271

FIGURE LEGEND

Fig 1

A

Genescan analysis of the CGG repeats within FMR1 gene from Normal male control (Left) and from FXTAS patient (Right):

Alleles size are shown in the upper right corner. The patient was a mosaic for a 44 allele 1(red arrow) and for a 116 allele (green arrow)

B

Recruitment of HSP70 cellular chaperone to FMRpolyG:

Fibroblasts from FXTAS patient and control were processed for indirect immunofluorescence using antibodies against FMRpolyG and against HSP70. HSP70 and FMRpolyG colocalize in about 45% of FXTAS fibroblasts

No association is evident in normal control fibroblasts

C

Western Blot Analysis of HSP70 FMRpolyG interaction:

Cellular extracts from patient's fibroblast and normal control, immunoprecipitated with anti FMRpolyG (left) were separated on 12% polyacrilammide gel. After transfer, the filter was incubated with an anti hHsP70 Ab and developped with a secondary HRP conjugated antibody. Western blot of HSP70 protein on the same amount of cellular

extracts , without IP is shown in the right side of the panel. Strikingly ,the recruitment of

HSP70 by FMRpolyG (left w/ IP) and the increased levels of total HSP70 (both ATP

and free forms,) in the patient's fibroblasts are confirmed

W/IP. Immunoprecipitation

W/0 IP : No immunoprecipitation

A   37   44,116

| DAPI | FMRP | HSP70 | MERGE |

C