Fasting protects hypoxia-induced defects in proteostasis in *C. elegans*

Short title: Fasting prevents hypoxia-induced defects in proteostasis

Nicole N Iranon[1,2], Bailey E Jochim[2], and Dana L Miller[2]

[1] Graduate Program in Molecular and Cellular Biology, University of Washington School of Medicine, Seattle WA 98195-7350, USA
[2] Department of Biochemistry, University of Washington School of Medicine, Seattle WA 98195-7350, USA

Address correspondence to DLM: dlm16@uw.edu

Abstract

Low oxygen conditions (hypoxia) can impair essential physiological processes and cause cellular
damage and death. We have shown that specific hypoxic conditions disrupt protein
homeostasis in *C. elegans*, leading to protein aggregation and proteotoxicity. Here, we show
that nutritional cues regulate the effect of hypoxia on proteostasis. Animals that are fasted
prior to hypoxic exposure develop dramatically fewer protein aggregates compared to their fed
counterparts. Our results suggest an important role for the nutritional environment
experienced at the onset of hypoxia in mediating hypoxia's effect on proteostasis, as fasting
protection can be both induced and reversed rapidly. Fasting is effective at protecting against
hypoxia-induced proteostasis defects in multiple developmental stages, tissues, and in different
models of misfolded or aggregation prone proteins. We further demonstrate that the
insulin/IGF-like signaling (IIS) pathway plays a role in mediating the protective effects of fasting
in hypoxia. Animals with mutations in *daf-2*, the *C. elegans* insulin-like receptor, display wild-
type levels of hypoxia-induced protein aggregation upon exposure to hypoxia when fed, but are
not protected by fasting. Our data further show that DAF-2 acts independently of the FOXO
transcription factor, DAF-16, to mediate the protective effects of fasting. These results suggest
a non-canonical role for the IIS pathway in coordinating the effects of hypoxia and nutritional
state on proteostasis.

Author Summary

When blood flow to various parts of the body becomes restricted, those tissues suffer from a
lack of oxygen, a condition called hypoxia. Hypoxia can cause cellular damage and death, such
as is observed as a result of stroke and cardiovascular disease. We have found that in the model
organism *C. elegans* (a roundworm) specific concentrations of hypoxia cause aggregation of
polyglutamine proteins – the same kind of proteins that are found in an aggregated state in the
neurodegenerative disorder Huntington's disease. Here, we show that that worms can be
protected from hypoxia-induced protein aggregation if they are fasted, or removed from their
food source prior to experiencing hypoxia. Furthermore, we show that the insulin receptor is
required for this protection. The insulin receptor is responsible for detecting insulin, a hormone
that is released after feeding. Worms with a nonfunctional version of the insulin receptor
displayed hypoxia-induced protein aggregation despite being fasted before the hypoxic
exposure. Our results highlight a new role for the insulin signaling pathway in coordinating the
effects of both hypoxia and nutritional state on protein aggregation.

Introduction

In order to survive in changing conditions, organisms need to successfully integrate a

number of environmental signals and respond appropriately in order to maintain homeostasis.

Aerobic heterotrophs must meet their requirements for food and oxygen by taking in these

resources from the environment. An inadequate response to low levels of oxygen (hypoxia) can

lead to cellular damage or death, an unsurprising outcome given oxygen's central role in

cellular metabolism. Like hypoxia, food deprivation presents an obstacle to homeostasis by

impinging on cellular metabolism and disturbing anabolic pathways. However, in many cases

food restriction can have beneficial effects, such as extending lifespan and delaying the onset of

neurodegenerative diseases and their associated pathologies [1]. In a mouse model of

Alzheimer's disease, 12 weeks of caloric restriction reduces Aβ plaque burden [2], and mice

expressing human mutant huntingtin maintained on an alternate-day-feeding diet have

reduced brain atrophy and decreased huntingtin aggregate formation [3]. Depriving *C. elegans*

of their bacterial food source reduces damage associated with expressing polyglutamine

proteins [4].

The protective effect of fasting is not limited to symptoms of neurodegeneration – there

are many studies that show fasting can protect against damage associated with hypoxia in

mammals. For example, mice on an alternate-day feeding regimen have higher survival rates

after myocardial ischemia induced via coronary occlusion [5]. Similar results have been

obtained with ischemic damage to the liver. Mice on a calorically restricted diet have reduced

infarct damage compared to ad-libitum fed controls [6], and mice that have been fasted for 3

days display reduced hepatocellular apoptosis and damage [7]. Calorie restriction can also

improve outcomes after cerebral ischemic injury by protecting cortical and striatal neurons [8],

and reducing neurological deficits and infarct volume [9]. These observations suggest that

understanding the mechanistic basis underlying the protective effects of fasting in hypoxia

could provide novel insight into therapeutic strategies to treat pathological conditions

associated with ischemia and reperfusion injury.

We have previously shown that in *C. elegans* the cellular response to specific hypoxic

conditions involves a disruption of proteostasis – the coordination of protein synthesis, folding,

degradation, and quality control required to maintain a functional proteome [10]. Here we

show that fasting prevents the hypoxia-induced disruption of proteostasis. Our data indicate

that the nutritional context of an animal at the onset of hypoxia has the power to alter

hypoxia's effect on proteostasis and that the insulin-like signaling (IIS) pathway plays a role in

fasting's ability to protect against proteostasis decline independently of the canonical

downstream transcription factor DAF-16/FOXO.

Results

In order to investigate the effect of nutritional status on proteostasis in hypoxia, we first

used transgenic *C. elegans* that express yellow fluorescent protein (YFP) fused to a

polyglutamine tract in the body wall muscles [11]. We refer to these animals as QX::YFP, where

X refers to the number of glutamine residues fused to YFP, such that Q35::YFP animals express

YFP with 35 glutamine residues. In these animals, the number of YFP foci, which correspond to

large protein aggregates, can be used as an *in vivo* measure of cellular proteostasis [12].

Exposing animals to 0.1% oxygen for 24 hours while fed resulted in an increase in the

number of YFP foci (Fig. 1B-1D), consistent with a decrease in proteostasis as has been

demonstrated previously [10]. However, we found that the number of YFP foci that formed in

hypoxia was dramatically reduced if the animals were removed from food for six hours before

the hypoxic exposure and remained off of food for the duration of hypoxia (Fig. 1A). Hypoxia-

induced protein aggregation (HIPA) was prevented by fasting in fourth-stage larvae (L4)

Q35::YFP animals (Fig. 1C) as well as in first-stage larvae (L1) Q40::YFP (Fig. 1D). We conclude

that fasting prevents HIPA and that this effect persists across development.

Figure 1. Fasting protects against hypoxia-induced protein aggregation. A.
Experimental Schematic. Cohorts of age-synchronized animals were split into three groups: the
first was maintained on food in room air, the second was maintained on food before and during
exposure to hypoxia, and the third was removed from food before exposure to hypoxia. Fasting
is indicated by white plates, yellow plates indicate animals on food. F= the duration of fasting
(h) before hypoxia; H = duration of hypoxia (h). Unless otherwise noted, aggregates were
counted immediately upon removal from hypoxia. **B.** Representative images of Q40::YFP
animals from cohorts of animals maintained in room air, exposed to hypoxia on food (hypoxia
fed), or exposed to hypoxia while fasted (hypoxia fasted). F=6h, H=24h. Scale bars = 100μm**. C-**
D. Aggregation measurements for L4 Q35::YFP (**C**) and L1 Q40::YFP (**D**) animals exposed to
hypoxia on food (fed, magenta) or after removal from food (fasted, blue). Controls remained in
room air (green). Data from one representative experiment is shown. Each experiment was
repeated at least 3 times. Each circle is the number of YFP foci in a single animal, the mean is
indicated by the line, and error bars are the standard deviation. Statistical comparisons were
made between animals exposed to hypoxia and controls maintained in room air. Significance:
*** $p < 0.001$; ns, not significant.

Figure 2: Fasting protection against HIPA is quickly induced and reversed. A. Effect of fasting
occurs rapidly in hypoxic conditions. Cohorts of L1 Q40::YFP animals were removed from food
before exposure to hypoxia (F = 0, 2, or 4 h; H=24 h). All animals were off of food when exposed
to hypoxia and the number of foci was scored immediately upon removal from hypoxia (cyan).
Controls remained in room air (green), were continuously on food (fed, magenta), or were
fasted for a full 6 h before hypoxia (fasted, blue). Data from one representative experiment is
shown. Each experiment was repeated at least 3 times. Each circle is the number of YFP foci in a
single animal, the mean is indicated by the line, error bars are the standard deviation.
Significance was calculated using a Kruskal-Wallis test and Dunn's multiple comparisons post
hoc analysis. Significant differences ($p < 0.05$) in aggregation between conditions are indicated
by letters above each group as follows: a - significantly different from room air controls; b -
significantly different from fed hypoxic controls; c - significantly different from fasted hypoxic
controls. **B.** Fasting before exposure to hypoxia improves proteostasis. As shown in the
schematic above the graph, cohorts of L1 Q40::YFP animals were removed from food 6h before
exposure to hypoxia, and fasted for 2, 4, or 6 h before being returned to food. All cohorts were
on food when exposed to hypoxia (H=24 h). The number of foci was scored immediately upon
return to room air (gray). Controls remained in room air (green), were continuously on food and
exposed to hypoxia (fed, magenta), or were not returned to food before hypoxia (fasted, blue).
Data from one representative experiment is shown. Each experiment was repeated at least 3
times. Each circle is the number of YFP foci in a single animal, the mean is indicated by the line,

error bars are the standard deviation. Statistical comparisons were made between animals
fasted for the indicated amount of time and controls maintained in room air, fed controls
exposed to hypoxia after being continuously on food, and fasted controls that were not
returned to food before hypoxia. Significance was calculated using a Kruskal-Wallis test and
Dunn's multiple comparisons post hoc analysis. Significant differences ($p < 0.05$) in aggregation
between conditions are indicated by letters above each group as follows: a - significantly
different from room air controls; b - significantly different from fed hypoxic controls; c -
significantly different from fasted hypoxic controls.

We originally chose to fast animals for 6h before exposure to hypoxia to allow animals

time to alter gene expression [13], and this period of time off of food is sufficient to deplete

stored glycogen as measured by iodine staining (DLM unpublished). However, there is no

evidence to suggest that the protective effects of fasting in hypoxia requires changes in gene

expression or glycogen stores. Therefore we next measured how long of a fasting period was

required to mitigate the effects of hypoxia on aggregation of polyglutamine proteins.

To determine the pre-hypoxia fasting duration required to protect against HIPA, we

removed Q35::YFP animals from food for varying lengths of time before being exposed to

hypoxia (as diagrammed in Fig. 2A). We found that animals removed from food immediately

before exposure to hypoxia developed significantly fewer YFP foci in hypoxia as compared to

controls that remained on food in hypoxia (Fig. 2A, 6h fed compared to fed). We conclude that

extended fasting before exposure to hypoxia is not required to prevent HIPA. Instead, our data

show that the protective effects of fasting occur very rapidly. In fact, the full protection against

HIPA is realized with only 2h fasting before exposure to hypoxia (Fig. 2A). These results suggest

that at least some of the protective effects of fasting are due to the absence of food directly,

rather than metabolic changes or alterations in gene expression that occur during fasting prior

to the hypoxic insult.

144   Work in other systems has shown that fasting can have a protective effect that persists

145 even after animals are returned to food (Robertson and Mitchell 2014). To further explore the

146 requirements for fasting to protect against HIPA we next asked whether the protective effects

147 of fasting against HIPA could be reversed. In these experiments (Fig. 2B), we began fasting

148 animals 6h before exposure to hypoxia but then returned the animals to food prior to initiation

149 of hypoxia. We observed that animals fasted for a full 6h and then returned to food

150 immediately before exposure to hypoxia (Fig. 2B, 6h fasted) developed significantly more YFP

151 foci than animals that were fasted for 6h and then exposed to hypoxia in the absence of food

152 (Fig 2B, fasted), suggesting that the nutritional context of an animal as it experiences hypoxia is

153 able to mediate the effect of hypoxia on proteostasis. Furthermore, we found no protection

154 from HIPA if animals were fasted for 4h, but then fed for 2 h before exposure to hypoxia (Fig.

155 2B, 4h fasted), even though 4h of fasting was sufficient for complete protection against HIPA in

156 the absence of food (Fig. 2A, 2h fed). This result indicates that the protective effects of fasting

157 are fully reversed within 2h of return to food. We conclude that the protective effects of fasting

158 in hypoxia are rapidly reversed.

Figure 3: Fasting protects against long-term effects of hypoxia on proteostasis.
Cohorts of L4 Q35::YFP animals were exposed to hypoxia (H=10 h) on food (magenta) or fasted
(blue, F=6h). Controls remained in room air on food (green). The number of YFP foci was scored
after return to room air as indicated. Data from one representative experiment is shown. The
experiment was repeated at least 3 times. Each cohort included at least 20 animals per time
point.

Shorter exposures to hypoxia that do not immediately increase the number of

polyglutamine protein aggregates still disrupt long-term proteostasis, as evidenced by the

increased rate of age-associated protein aggregation after return to room air [10]. We

therefore asked whether fasting could protect against these long-term proteostasis deficits in

addition to HIPA. We exposed Q35::YFP L4 animals to hypoxia for only 10h either in the fed

state or after fasting for 6h (F = 6 hours, H = 10 hours as per Fig. 1A). Control animals remained

on food in room air. Immediately after this short hypoxic exposure, there was no observed

increase in the number of YFP foci in animals exposed to hypoxia regardless of whether food

was present (Fig. 3, 0 hours post-hypoxia). As expected, the animals exposed to hypoxia in the

fed state accumulate aggregates faster than control animals. In contrast, animals exposed to

178   hypoxia while fasted accumulate YFP foci at the same rate as control animals. These data

179   indicate that fasting both prevents HIPA and protects against the long-term effects on

180   proteostasis induced by a short exposure to hypoxia.

Figure 4. Fasting has general protective effects against hypoxia-induced defects in
proteostasis. A. Fasting protects against toxicity of Q40::YFP. Cohorts of L1 animals expressing
Q40::YFP were exposed to hypoxia on food (magenta), or fasted (blue) before exposure to
hypoxia (F=6h, H=24 h). Paralysis was scored after return to room air, beginning the first day of
adulthood. Controls remained on food in room air (green). Data from one representative
experiment is shown, each cohort included at least 70 animals. Each experiment was repeated
at least 3 times. Significance was calculated using a Log-rank (Mantel-Cox) test with a
Bonferroni correction for multiple comparisons. Statistical comparisons were made between
animals exposed to hypoxia and animals maintained in room air. **** $p < 0.0001$; ** $p < 0.01$.
B. Fasting protects against toxicity of Aβ_{1-42}. Cohorts of L4 animals expressing Aβ_{1-42} were
exposed to hypoxia on food (magenta) or fasted (blue) before exposure to hypoxia (F=6h,
H=24h). Paralysis was scored after return to room air, beginning at the first day of adulthood.

Controls remained on food in room air (green). Data from one representative experiment is
shown, each cohort included at least 70 animals. Each experiment was repeated at least 3
times. Significance was calculated using a Log-rank (Mantel-Cox) test with a Bonferroni
correction for multiple comparisons. Statistical comparisons were made between animals
exposed to hypoxia and animals maintained in room air. *** $p < 0.001$. **C.** Fasting protects
against hypoxia effects on metastable DYN-1. Temperature-sensitive *dyn-1(ky51)* mutant
animals were exposed to hypoxia at the permissive temperature on food (magenta), or after
fasting (blue). Controls remained on food in room air at the permissive temperature (green) or
on food at the non-permissive temperature (28°C, gray). Paralysis was scored 1h after return to
room air. Average data from 3 independent experiments is shown, each cohort included 10
animals. Significance was calculated using a repeated measures two-way ANOVA and Dunnett's
multiple comparisons test. Statistical comparisons were made between animals exposed to
hypoxia or animals maintained at the restricted temperature and animals maintained in room
air. Significance: *** $p < 0.001$; ** $p < 0.01$
209

The cellular role of protein aggregates is controversial, with some reports finding a

protective role and others suggesting a cytotoxic effect [14]. We have previously shown that

aggregates induced by hypoxia are cytotoxic, resulting in accelerated paralysis after animals are

returned to room air [10]. We therefore next asked if fasting would protect against increased

proteotoxicity in addition to HIPA. To address this, we exposed cohorts of L1 Q40::YFP animals

to hypoxia for 24 hours while fed or fasted, then returned the animals to room air and

measured the onset of paralysis in each cohort. We found that fasting slowed the rate at which

paralysis developed relative to animals exposed to hypoxia while fed (Fig. 4A). This result

indicates that fasting protects against hypoxic effects of increased protein aggregation and

proteotoxicity.

We next sought to determine whether fasting's protective effects on proteostasis

extend to other models of proteotoxicity. Human amyloid β (Aβ)$_{1-42}$ peptide expressed in the

body wall muscles of *C. elegans* results in cytoplasmic plaque formation, with a subsequent

phenotype of progressive paralysis [15]. *C. elegans* expressing Aβ$_{1-42}$ in their body wall muscles

224 become paralyzed more quickly when they are exposed to hypoxia [10]. We found that this

225 effect of hypoxia was reversed by fasting, as the rate that paralysis develops is slowed if animals

226 expressing $A\beta_{1-42}$ are exposed to hypoxia while fasting (Fig. 4B). Because $A\beta_{1-42}$ and Q40::YFP are

227 both expressed in body wall muscles, we also evaluated if fasting protected animals expressing

228 a metastable version of the neuronal dynamin protein DYN-1 from the effects of hypoxia. The

229 *dyn-1(ky51)* mutant contains a temperature-sensitive (ts) mutation, such that the DYN-1

230 protein is functional and *dyn-1(ky51)* mutant animals exhibit wild-type motility at the

231 permissive temperature (20°C), but become uncoordinated at the restrictive temperature

232 (28°C) due to improper folding of the DYN-1 protein [16]. Genetic and environmental factors

233 that disrupt proteostasis, including hypoxia, prevent the proper folding of the DYN-1 protein at

234 the permissive temperature, thereby rendering the *dyn-1(ky51)* animals uncoordinated [17,

235 10]. Similar to our experiments with Q40::YFP and $A\beta_{1-42}$, we found that fasting *dyn-1(ky51)*

236 mutant animals before exposure to hypoxia results in a partial rescue of hypoxia-induced

237 uncoordination at the permissive temperature (Fig. 4C). Together, our results suggest that

238 fasting has a general protective effect against proteostasis defects induced by hypoxia, and that

239 this protective effect is not specific to a particular tissue, developmental stage, or

240 misfolded/aggregation prone model.

241   Dysregulation of insulin-like signaling (IIS) has been tied to protein aggregation and

242 neurodegeneration in a number of model organisms [18]. As the IIS pathway links food

243 availability to growth, development, stress resistance, and aging, we hypothesized that changes

244 in IIS could explain how fasting modulates the effect of hypoxia on proteostasis. The IIS

pathway is widely conserved in metazoans [19]. We therefore explored the hypothesis that IIS

would mediate the effects of fasting to prevent HIPA.

We first looked at the localization of DAF-16::GFP in animals exposed to hypoxia to

determine if IIS is active in hypoxia. DAF-16 is the *C. elegans* orthologue of the FOXO

transcription factor. When active, the insulin/IGF-like receptor DAF-2 initiates a

phosphorylation cascade that results in the phosphorylation and nuclear exclusion of DAF-16

protein [20, 21]. Conversely, when nutrients are scarce, DAF-16 remains unphosphorylated by

upstream kinases and is able to enter the nucleus and bind to its target genes [20, 22]. We

found that DAF-16::GFP remained diffuse and cytoplasmic in control worms maintained in room

air on food (Fig 5B, 5C), but accumulated in the nucleus of animals that were removed from

food in room air (Fig. 5B, 5C) or were exposed to hypoxia on food (Fig. 5B, 5C). These results

suggest that IIS activity is reduced by fasting and hypoxia, consistent with previous reports [23,

24]. Surprisingly, DAF-16::GFP did not accumulate in the nuclei of animals exposed to hypoxia

after fasting (Fig 5B, 5C), despite hypoxia and fasting both individually resulting in nuclear

accumulation.

These DAF-16::GFP localization patterns led us to interrogate requirements for DAF-16

and the upstream IIS receptor DAF-2 in mediating fasted and fed responses to hypoxia. To this

end, we crossed the Q35::YFP transgene into *daf-2(e1370)* and *daf-16(mu86)* backgrounds. The

fact that DAF-16::GFP is localized to the nucleus in fed animals exposed to hypoxia suggests the

possibility that DAF-16 facilitates HIPA. We found that *Q35::YFP; daf-16(mu86)* mutant animals

exhibit robust HIPA on food (Fig 5D), indicating that DAF-16 is not required for HIPA despite its

nuclear accumulation in fed hypoxic animals. We also asked if there was a genetic requirement

for the IIS receptor DAF-2. Our data indicate that IIS does not mediate the effects of hypoxia on

proteostasis in fed animals, as *Q35::YFP; daf-2(e1370)* mutant animals exhibit robust HIPA

when fed (Fig. 5D). Thus, neither DAF-16 nor DAF-2 activities are required for HIPA in fed

animals.

Given the IIS-independent nature of HIPA in fed animals, we next investigated whether

fasting protection requires IIS. We discovered that DAF-2, but not DAF-16 is required for fasting

protection against HIPA. Fasting protects the *Q35::YFP; daf-16(mu86)* similar to wild-type (Fig

5D); however, we observe significant HIPA when *Q35; daf-2(e1370)* and *Q35; daf-2(e1368)*

mutant animals are exposed to hypoxia when fasted (Fig 5D and Supplemental Fig. 1). These

results show that protective effects of fasting in hypoxia require DAF-2, but not DAF-16. This is

consistent with our observation that DAF-16::GFP is not localized to the nucleus in fasted

animals exposed to hypoxia (Fig 5B, 5C).

We found that the insulin/IGF-like receptor DAF-2 mediates the protective effects of

fasting on HIPA, while the the FOXO transcription factor DAF-16 is not required for protection.

Given this finding, we also checked the DAF-16::GFP localization pattern in worms with a *daf-*

2(e1370) mutation. These mutants have constitutively nuclear DAF-16 in the fed state due to

decreased signaling through the IIS pathway [20]. Since DAF-16::GFP is not localized to the

nucleus in fasting-protected wild-type animals exposed to hypoxia, we sought to investigate

whether the nuclear localization of DAF-16 in *daf-2(e1370)* mutants, which are not protected by

fasting, would be altered by hypoxia. We found that DAF-16::GFP is fully nuclear in all

conditions, including fasted hypoxia, in these animals (Supplemental Fig. 2).

In *C. elegans*, DAF-16 mediates the effects of decreased signaling through DAF-2.

Mutations in *daf-16* suppress most *daf-2* mutant phenotypes including increased lifespan,

enhanced dauer formation, increased fat storage, reproductive delays, and increased resistance

to heat and oxidative stress. [25, 26]. This coupled with the nuclear localization of DAF-16::GFP

in *daf-2* mutants led us to hypothesize that *daf-16* would be required for the HIPA in fasted

Q35; daf-2(e1370) mutant animals. While *Q35; daf-16(mu86)* mutant animals were protected

from HIPA by fasting similar to wild-type controls, *Q35; daf-2(e1370);daf-16(mu86)* animals still

exhibit significant HIPA when fasted (Fig. 5D). These results indicate that DAF-2 mediates the

effects of fasting to prevent HIPA at least partly independently of DAF-16.

Figure 5. The insulin-like signaling pathway is required for fasting protection. A Schematic of
key insulin-signaling pathway members in *C. elegans*. Under nutrient-rich conditions, insulin-like
peptides bind to the insulin receptor DAF-2, initiating a phosphorylation cascades that
ultimately leads to the phosphorylation of the FoxO transcription factor DAF-16, excluding it
from the nucleus. Conversely, when nutrients are scarce, DAF-16 remains unphosphorylated
and is able to enter the nucleus and bind to its target genes. **B.** DAF-16 is not localized to the
nucleus in fasted animals exposed to hypoxia. Cohorts of 20 DAF-16::GFP animals were
maintained in room air on food for 24 hrs (fed + room air), fasted in room air for 24 hrs (fasted
+ room air), exposed to hypoxia for 24 hrs on food (fed + hypoxia), or exposed to hypoxia after
fasting (fasted + hypoxia; F=6h, H=24hr). Scale bars = 100μm. **C** Quantification of DAF-16::GFP
nuclear accumulation. The percent of animals with nuclear GFP was scored immediately post
hypoxia. Average data from 3 independent experiments is shown. The bar height indicates the
mean. Error bars are the standard deviation. **D** Fasting does not protect *daf-2* mutants against
HIPA. Aggregation measurements (F=6h, H=24h) for L4 Q35::YFP animals with mutations in *daf-*
2(e1370), *daf-16(mu86)*, and the *daf-2(e1370); daf-16(mu86)* double mutant. Animals were
maintained on food in room air (room air, green), were exposed to hypoxia on food (fed,
magenta), or were exposed to hypoxia after removal from food (hypoxia fasted, blue). Each
circle is the number of YFP foci in a single animal, the mean is indicated by the line, error bars
are the standard deviation. Data from one representative experiment is shown. Each cohort
included at least 20 animals, and each experiment was repeated at least 3 times. Significance
was calculated using a Kruskal-Wallis test and Dunn's multiple comparisons post hoc analysis.
Significant differences ($p < 0.05$) in aggregation for a given strain between conditions are
indicated by letters above each group as follows: a - significantly different from room air
controls; b - significantly different from fed hypoxic controls; c - significantly different from
fasted hypoxic controls.
323

Discussion

This study illustrates the power of fasting to ameliorate the deleterious effects of

hypoxia on proteostasis. These findings are consistent with phenomena that have been

observed in mammals – fasting mice for a single day increases survival after kidney ischemia

and also reduces ischemic damage to the liver [27]. Our results suggest that the nutritional

milieu present at the onset of hypoxia can dictate the effect of hypoxia on proteostasis, as

fasting protection against hypoxia can be induced quite quickly. Animals that are removed from

food immediately before hypoxia are protected against HIPA to a significant degree, even after

being maintained on food for the entire pre-hypoxic period. This implies that worms are

integrating information about their environment, including nutrient availability, right as they

sense hypoxia. The importance of the nutritional environment of the animal as it experiences

hypoxia is further supported by the fact that we also see a rapid reversal of fasting protection.

Worms fasted for six hours but that are moved onto food immediately preceding hypoxia are

not as protected against HIPA compared to worms that were fasted and remained off of food

for the duration of hypoxia. The speed with which fasting protection can be induced and

reversed indicates that protection cannot be explained solely by changes in gene expression

resulting in a hypoxia-resistant pre-adapted state. Furthermore, the rapidity with which fasting

protection can be reversed suggests that altered gene expression or metabolism resulting from

the fasting period is alone insufficient to protect against HIPA. Although *C. elegans* enter a

reproductive and developmental diapause in 0.1% oxygen [28], the protection conferred by

fasting does not represent a simple delay in the onset of proteostasis decline due to the time

spent in hypoxia. Rather, fasting provides long-term protection against the accrual of protein

aggregates and toxicity even after the return to room air.

We found that IIS mediates fasting protection against HIPA. Notably, IIS is not required

for the fed response to hypoxia, as fed IIS mutants show increased aggregate levels comparable

to wild-type animals. In worms and flies, mutations in the insulin receptor are generally

considered protective against hypoxia. In *C. elegans, daf-2* mutants have a hypoxia-resistant

phenotype, displaying reduced muscle and neuronal cell death following hypoxia [29, 30], while

flies with defective insulin signaling (mutations in the insulin receptor *InR*, or *Chico*, the insulin

receptor substrate) are protected against anoxia/reoxygenation injury [31]. The *daf-2*

phenotype uncovered here is therefore distinct in that these mutants are sensitive to hypoxia in

the fasted state, with fasted *daf-2* mutant animals exhibiting increased HIPA compared to wild-

type controls. These results contradict the *a priori* expectation that *daf-2* mutants might be

resistant to hypoxia even in the fed state due to their inability to detect insulin-like peptides.

Mammalian systems offer precedents of insulin receptor mutations causing sensitivity

to hypoxic stress. Knockdown of neuronal insulin-like growth factor 1 receptor (IGF-1R)

exacerbates hypoxic injury and increases mortality in mice [32], and IGF-1R is required in order

for IGF-1 to protect myocardial cell exposed to ischemia [33]. However, data on the role of

mammalian IIS in response to hypoxia are mixed, and are complicated by the fact that different

types of insulin receptors mediate distinct cellular functions [34]. As such, the simplified *C.*

elegans IIS system may be useful for understanding contextual inputs that alter IIS outputs.

DAF-16 is believed to be the main nexus of IIS [20, 35-37], which makes the DAF-2-

dependent, but DAF-16-independent nature of the protective effect of fasting described here

unusual in *C. elegans*. Decreased DAF-2 activity results in phenotypes such as increased

lifespan, reproductive delays, and increased resistance to heat and oxidative stress, all of which

require DAF-16 [26]. However, a few other examples exist in the literature of DAF-2 dependent,

DAF-16 independent phenomena: dauer formation at 27°, meiotic progression of oocytes, salt

chemotaxis learning, and regulation of the dao-3 and hsp-90 genes [38- 42]. In chemotaxis

learning, DAf-2 acts on learning through phosphatidylinositol 3,4,5-triphosphate (PIP_3), but not

DAF-16. Similar to these studies, fasting-mediated protection against HIPA supports the

existence of downstream targets of DAF-2 separate from DAF-16 that are capable of influencing

stress responses and proteostasis.

Materials and methods

C. elegans strains and methods

Animals were maintained on nematode growth media (NGM) with OP50 *E. coli* at 20°C

(Brenner, 1974). See Supplementary Table S6 for worm strains. Strains were obtained from the

Caenhorabditis Genetics Center at the University of Minnesota. Double and triple mutants were

generated using standard genetic techniques, and genotypes were verified using PCR.

Construction of hypoxic chambers

Hypoxic conditions were maintained using continuous flow chambers, as described in

Fawcett et al. 2012. Compressed gas tanks (1000 ppm O_2 balanced with N_2) were Certified

Standard (within 2% of target concentration) from Airgas (Seattle, WA). Oxygen flow was

regulating using Aalborg rotameters (Aalborg Intruments and Controls, Inc., Orangeburg, NY,

USA). Hypoxic chambers (and room air controls?) were maintained in a 20°C incubator for the

duration of the experiments.

YFP::polyQ aggregation assays

Synchronous cohorts of L1 YFP::polyQ$_{40}$ animals were generated by either bleaching

first-day adult animals in a 20% alkaline hypochlorite solution or allowing first-day adult animals

to lay eggs for 1-2 hrs on seeded NGM plates. The adults were then removed, and the plates

were incubated at 20°C. The next morning, cohorts of hatched L1 larvae were suspended in M9

and mouth-pipetted to new NGM plates for hypoxic exposure. Synchronous cohorts of L4

YFP::polyQ$_{35}$ animals were generated by picking L4 animals from well-fed, logarithmically

growing populations.

Cohorts of 25-35 YFP::polyQ animals were exposed to hypoxia for approximately 24 h at

20°C on unseeded 3 cm NGM plates with 40mg/mL carbenicillin or NGM plates seeded with live

OP50 food. Plates were ringed with palmitic acid (10mg/mL in ethanol), creating a physical

barrier around the edge of each plate to discourage animals from leaving the surface of the

agar.

To quantify the number of YFP foci, worms were mounted a 2% agar pad in a drop of

50mM sodium azide as anesthetic. Control experiments showed that azide did not affect the

aggregation of YFP::polyQ$_{35}$ or YFP::polyQ4$_{0}$ (Moronetti Mazzeo et al. 2012). YFP foci were

identified and quantified as described in Morley et al. (2002) and Silva et al. (2011). A Nikon 90i

fluorescence microscope with the YFP filter and 10x objective (Nikon Instruments Inc., Melville,

NY, USA) was used to visualize and quantify aggregates. In all experiments, the number of

aggregates was counted blind to treatment and genotype. Statistical significance was evaluated

by calculating P-values between conditions using a Kruskal-Wallis test and Dunn's multiple

comparisons post hoc analysis in GraphPad Prism version 7.0c for Mac OSX (GraphPad Softare,

San Diego, California, USA) In all cases, $P < 0.05$ was considered statistically significant.

Paralysis and uncoordination assays of proteotoxicity

Animals expressing A$\beta_{1\text{-}42}$ or YFP::polyQ$_{40}$ were exposed to 1000 ppm O$_2$ for 24 at 20°C

as L4 or L1, respectively. For both, animals were grown on seeded NGM plates until 6 hrs before

hypoxic exposure, at which point fasted animals were transferred to unseeded NGM plates,

where they remained until the end of the hypoxic exposure. Fed animals were transferred to

new seeded NGM plates. After hypoxic exposure, all animals were returned to food and

normoxia, and incubated at 20°C. Paralysis was scored daily. Worms were considered paralyzed

if they failed to respond, other than with movement of the nose or pharyngeal pumping, when

tapped with a platinum wire pick 3 consecutive times. Dead or bagged worms were censored

from the experiment on the day of death/bagging. Paralyzed worms were removed from the

plate on the day of paralysis. Live worms that were not paralyzed were moved to a new plate

each day until all worms were scored as either paralyzed or dead. Statistical significance was

calculated using Kaplan-Meier log-rank (Mantel-Cox) tests and a Bonferroni correction for

multiple comparisons using GraphPad Prism version 7.0c for Mac OSX (GraphPad Softare, San

Diego, California, USA).

431
DAF-16::GFP localization

Synchronous cohorts of L2 animals expressing DAF-16::GFP were exposed to hypoxia for

24 h at 20°C on unseeded unseeded 3 cm NGM plates with 40mg/mL carbenicillin or NGM

plates seeded with live OP50 food. Plates were ringed with palmitic acid (10mg/mL in ethanol),

creating a physical barrier around the edge of each plate to discourage animals from leaving the

surface of the agar. To visualize the localization of DAF-16::GFP, worms were mounted a 2%

agar pad in a drop of 10mM levamisole as anesthetic. A Nikon 90i fluorescence microscope with

the GFP filter and 10x objective (Nikon Instruments Inc., Melville, NY, USA) was used to visualize

DAF-16::GFP. For quantification, percent of animals with nuclear GFP was scored immediately

after removal from hypoxia. In all experiments, the GFP localization was scored blind to

treatment and genotype. Statistical significance was evaluated by calculating P-values between

conditions using a Kruskal-Wallis test and Dunn's multiple comparisons post hoc analysis in

GraphPad Prism version 7.0c for Mac OSX (GraphPad Softare, San Diego, California, USA). P <

0.05 was considered statistically significant.

Acknowledgements

We

References:

1. Contestabile A. Benefits of caloric restriction on brain aging and related pathological states: understanding mechanisms to devise novel therapies. Curr Med Chem. 2009;16:350-361.

2. Patel NV, Gordon MN, Connor KE et al. Caloric restriction attenuates Abeta-deposition in Alzheimer transgenic models. Neurobiol Aging. 2005;26:995-1000.

3. Duan W, Guo Z, Jiang H, Ware M, Li XJ, Mattson MP. Dietary restriction normalizes glucose metabolism and BDNF levels, slows disease progression, and increases survival in huntingtin mutant mice. Proc Natl Acad Sci U S A. 2003;100:2911-2916.

4. Steinkraus KA, Smith ED, Davis C et al. Dietary restriction suppresses proteotoxicity and enhances longevity by an hsf-1-dependent mechanism in Caenorhabditis elegans. Aging Cell. 2008;7:394-404.

5. Katare RG, Kakinuma Y, Arikawa M, Yamasaki F, Sato T. Chronic intermittent fasting improves the survival following large myocardial ischemia by activation of BDNF/VEGF/PI3K signaling pathway. J Mol Cell Cardiol. 2009;46:405-412.

6. Menezes-Filho SL, Amigo I, Prado FM et al. Caloric restriction protects livers from ischemia/reperfusion damage by preventing Ca^{2+}-induced mitochondrial permeability transition. Free Radic Biol Med. 2017;110:219-227.

7. Qin J, Zhou J, Dai X et al. Short-term starvation attenuates liver ischemia-reperfusion injury (IRI) by Sirt1-autophagy signaling in mice. Am J Transl Res. 2016;8:3364-3375.

8. Duan W, Guo Z, Mattson MP. Brain-derived neurotrophic factor mediates an excitoprotective effect of dietary restriction in mice. J Neurochem. 2001;76:619-626.

9. Ran M, Li Z, Yang L, Tong L, Zhang L, Dong H. Calorie restriction attenuates cerebral ischemic injury via increasing SIRT1 synthesis in the rat. Brain Res. 2015;1610:61-68.

10. Fawcett EM, Hoyt JM, Johnson JK, Miller DL. Hypoxia disrupts proteostasis in Caenorhabditis elegans. Aging Cell. 2015;14:92-101.

11. Morley JF, Brignull HR, Weyers JJ, Morimoto RI. The threshold for polyglutamine-expansion protein aggregation and cellular toxicity is dynamic and influenced by aging in Caenorhabditis elegans. Proc Natl Acad Sci U S A. 2002;99:10417-10422.

12. Satyal SH, Schmidt E, Kitagawa K et al. Polyglutamine aggregates alter protein folding homeostasis in Caenorhabditis elegans. Proc Natl Acad Sci U S A. 2000;97:5750-5755.

13. Van Gilst MR, Hadjivassiliou H, Yamamoto KR. A Caenorhabditis elegans nutrient response system partially dependent on nuclear receptor NHR-49. Proc Natl Acad Sci U S A. 2005;102:13496-13501.

14. Soto C. Unfolding the role of protein misfolding in neurodegenerative diseases. Nat Rev Neurosci. 2003;4:49-60.

15. Link CD. Expression of human beta-amyloid peptide in transgenic Caenorhabditis elegans. Proc Natl Acad Sci USA. 1995;92:9368.

16. Clark SG, Shurland DL, Meyerowitz EM, Bargmann CI, van der Bliek AM. A dynamin GTPase mutation causes a rapid and reversible temperature-inducible locomotion defect in C. elegans. Proc Natl Acad Sci U S A. 1997;94:10438-10443.

17. Ben-Zvi A, Miller EA, Morimoto RI. Collapse of proteostasis represents an early molecular event in Caenorhabditis elegans aging. Proc Natl Acad Sci U S A. 2009;106:14914-14919.

18. Cohen E, Bieschke J, Perciavalle RM, Kelly JW, Dillin A. Opposing activities protect against age-onset proteotoxicity. Science. 2006;313:1604-1610.

19. Piñero González J, Carrillo Farnés O, Vasconcelos AT, González Pérez A. Conservation of key members in the course of the evolution of the insulin signaling pathway. Biosystems. 2009;95:7-16.

20. Lin K, Hsin H, Libina N, Kenyon C. Regulation of the Caenorhabditis elegans longevity protein DAF-16 by insulin/IGF-1 and germline signaling. Nat Genet. 2001;28:139-145.

21. Henderson ST, Johnson TE. daf-16 integrates developmental and environmental inputs to mediate aging in the nematode Caenorhabditis elegans. Curr Biol. 2001;11:1975-1980.

22. Murphy CT, McCarroll SA, Bargmann CI et al. Genes that act downstream of DAF-16 to influence the lifespan of Caenorhabditis elegans. Nature. 2003;424:277-283.

23. Honjoh S, Yamamoto T, Uno M, Nishida E. Signalling through RHEB-1 mediates intermittent fasting-induced longevity in C. elegans. Nature. 2009;457:726-730.

24. Leiser SF, Fletcher M, Begun A, Kaeberlein M. Life-span extension from hypoxia in Caenorhabditis elegans requires both HIF-1 and DAF-16 and is antagonized by SKN-1. J Gerontol A Biol Sci Med Sci. 2013;68:1135-1144.

25. Ogg S, Paradis S, Gottlieb S et al. The Fork head transcription factor DAF-16 transduces insulin-like metabolic and longevity signals in C. elegans. Nature. 1997;389:994-999.

26. Zhou KI, Pincus Z, Slack FJ. Longevity and stress in Caenorhabditis elegans. Aging (Albany NY). 2011;3:733-753.

27. Mitchell JR, Verweij M, Brand K et al. Short-term dietary restriction and fasting precondition against ischemia reperfusion injury in mice. Aging Cell. 2010;9:40-53.

28. Miller DL, Roth MB. C. elegans are protected from lethal hypoxia by an embryonic diapause. Curr Biol. 2009;19:1233-1237.

29. Scott BA, Avidan MS, Crowder CM. Regulation of hypoxic death in C. elegans by the insulin/IGF receptor homolog DAF-2. Science. 2002;296:2388-2391.

30. Mabon ME, Scott BA, Crowder CM. Divergent mechanisms controlling hypoxic sensitivity and lifespan by the DAF-2/insulin/IGF-receptor pathway. PLoS One. 2009;4:e7937.

31. Vigne P, Tauc M, Frelin C. Strong dietary restrictions protect Drosophila against anoxia/reoxygenation injuries. PLoS One. 2009;4:e5422.

32. Liu W, D'Ercole JA, Ye P. Blunting type 1 insulin-like growth factor receptor expression exacerbates neuronal apoptosis following hypoxic/ischemic injury. BMC Neurosci. 2011;12:64.

33. Liu A, Zhang X, Gu H, Li P, Yu T. Insulin-like growth factor-1 protects ischemic myocardial cells via PI3K/AKT pathway. Int J Clin Exp Pathol. 2016;9:12574-12580.

34. Cai W, Sakaguchi M, Kleinridders A et al. Domain-dependent effects of insulin and IGF-1 receptors on signalling and gene expression. Nat Commun. 2017;8:14892.

35. Dillin A, Crawford DK, Kenyon C. Timing requirements for insulin/IGF-1 signaling in C. elegans. Science. 2002;298:830-834.

36. Hsu AL, Murphy CT, Kenyon C. Regulation of aging and age-related disease by DAF-16 and heat-shock factor. Science. 2003;300:1142-1145.

37. Honda Y, Honda S. The daf-2 gene network for longevity regulates oxidative stress resistance and Mn-superoxide dismutase gene expression in Caenorhabditis elegans. FASEB J. 1999;13:1385-1393.

38. Ailion M, Thomas JH. Dauer formation induced by high temperatures in Caenorhabditis elegans. Genetics. 2000;156:1047-1067.

39. Lopez AL, Chen J, Joo HJ et al. DAF-2 and ERK couple nutrient availability to meiotic progression during Caenorhabditis elegans oogenesis. Dev Cell. 2013;27:227-240.

40. Tomioka M, Adachi T, Suzuki H, Kunitomo H, Schafer WR, Iino Y. The insulin/PI 3-kinase pathway regulates salt chemotaxis learning in Caenorhabditis elegans. Neuron. 2006;51:613-625.

41. Vellai T, McCulloch D, Gems D, Kovács AL. Effects of sex and insulin/insulin-like growth factor-1 signaling on performance in an associative learning paradigm in Caenorhabditis elegans. Genetics. 2006;174:309-316.

42. Yu H, Larsen PL. DAF-16-dependent and independent expression targets of DAF-2 insulin receptor-like pathway in Caenorhabditis elegans include FKBPs. J Mol Biol. 2001;314:1017-1028.