Decomposing the subclonal structure of tumors with two-way mixture models on copy number aberrations

models on copy number aberrations

An-Shun Tai[1], Chien-Hua Peng[1,*], Shih-Chi Peng[1], and Wen-Ping Hsieh[1,*]

1. Institute of Statistics, National Tsing Hua University, Hsinchu, Taiwan, R.O.C.

* Corresponding author

Wen-Ping Hsieh[1] Institute of Statistics, National Tsing Hua University 101 sec. 2, Kwang-Fu Rd. , Hsin-Chu City, Taiwan, R.O.C. 30013; TEL: 886-35715131-33188; FAX:886-35728318;

E-mail:wphsieh@stat.nthu.edu.tw

Chien-Hua Peng[1] Institute of Statistics, National Tsing Hua University 101 sec. 2, Kwang-Fu Rd. , Hsin-Chu City, Taiwan, R.O.C. 30013; TEL: 886-35715131-33195; FAX:886-35728318;

E-mail: ariel0207@gmail.com

Abstract

Multistage tumorigenesis is a dynamic process characterized by the accumulation of mutations. Thus, a tumor mass is composed of genetically divergent cell subclones. With the advancement of next-generation sequencing (NGS), mathematical models have been recently developed to decompose tumor subclonal architecture from a collective genome sequencing data. Most of the methods focused on single-nucleotide variants (SNVs). However, somatic copy number aberrations (CNAs) also play critical roles in carcinogenesis. Therefore, further modeling subclonal CNAs composition would hold the promise to improve the analysis of tumor heterogeneity and cancer evolution. To address this issue, we developed a two-way mixture Poisson model, named CloneDeMix for the deconvolution of read-depth information. It can infer the subclonal copy number, mutational cellular prevalence (MCP), subclone composition, and the order in which mutations occurred in the evolutionary hierarchy. The performance of CloneDeMix was systematically assessed in simulations. As a result, the accuracy of CNA inference was nearly 93% and the MCP was also accurately restored. Furthermore, we also demonstrated its applicability using head and neck cancer samples from TCGA. Our results inform about the extent of subclonal CNA diversity, and a group of candidate genes that probably initiate lymph node metastasis during tumor evolution was also discovered. Most importantly, these driver genes are located at 11q13.3 which is highly susceptible to copy number change in head and neck cancer genomes. This study successfully estimates subclonal CNAs and exhibit the evolutionary relationships of mutation events. By doing so, we can track tumor heterogeneity and identify crucial mutations during evolution process. Hence, it

43 facilitates not only understanding the cancer development but finding potential

44 therapeutic targets. Briefly, this framework has implications for improved modeling of

45 tumor evolution and the importance of inclusion of subclonal CNAs.

Introduction

Cancer, a dynamic disease, is characterized by unusual cells with somatic

mutations. These mutations are caused by environmental factors accumulated during

an individual's lifetime; this accumulation of mutational events results in a large degree

of genetic heterogeneity among cancer cells. The intratumor heterogeneity causes

difficulties in devising personalized treatment strategies.

To decipher intratumor heterogeneity, understanding how cancer evolves is a key

step. The hypothesis for the somatic evolution of cancer was proposed in the 1970s [1].

It states that all tumor cells descend from a single founder cell, and cells with some

advantageous mutations become more competitive than normal cells for growth and

clonal expansion. This hypothesis could also be formed through random drift. Gradually,

subsequent clonal expansion occurs, and the tumor evolves into an organization of

multiple cell subpopulations. Understanding clonal evolution in cancer is one of the

goals of cancer medicine [2]. Presently, sequencing technology enables performing a

large-scale molecular profiling of tumors to comprehend cancer development and

determine disease progression. However, the process of evolution is not directly

observed because tissues for measuring somatic mutations are typically obtained from

patients at a single time point. Thus, the ancestral relationship among tumor subclones

have to be inferred, and this is closely related to a well-studied problem, phylogenetic

tree reconstruction. To construct a phylogenetic tree, the mutations in each cancer cell

should be measured to infer evolutionary relationships among various cells. For

addressing this concern, the current technology of single-cell sequencing seems

appropriate [3, 4]. However, this technology is not widely used because of some

technical limitations and financial considerations [5]. Most studies on tumor evolution

rely on DNA sequencing technology with a bulk tumor containing genetically different

cells. Therefore, the cellular prevalence of each subclone have to be measured through

the relative read count information of the variants.

Single-nucleotide variants (SNVs) and copy number aberrations (CNAs) are

widely used data types to study tumor evolution. Recently, studies inferring the

population structure and clonal architecture have either focused on SNVs according to

variant allele frequencies (VAFs) or on CNAs with read counts obtained through DNA

sequencing [6, 7]. Methods for either type of data can adopt the other type of data to

improve their reconstruction, and most methods have developed corresponding

computational tools.

The first category of method reconstructs models with only SNV data. AncesTree

and clonality inference in tumors using phylogeny (CITUP) are the representatives of

this category, and they build models based on heterozygous SNV to study cancer

progression, assuming that the copy number is two [8, 9]. To relax the assumption of

the normal copy number status, many studies have included CNAs to correct the

baseline [10-13]. For instance, Pyclone is one of the clonal inference approaches, and

it applies a hierarchical Bayes binomial distribution to model allelic counts [13]. This

approach applies a Dirichlet process prior on group mutations and infers the posterior

distribution to estimate the cellular prevalence, which is the fraction of cancer cells

harboring a mutation.

Unfortunately, the aforementioned algorithms only considered abnormal copy

number states but do not infer the clonal structure of copy number changes. If we do

not account for clonal evolutionary architecture, the estimation of CNAs would be

inaccurate and just reported as an average of the CNAs of all tumor subclones. Hence,

in contrast to the SNV-based models, some studies focus on subclonal CNA

heterogeneity [7, 14-18]. They recognize that subclonal CNAs could technically

improve the analysis accuracy. THetA is one of the most popular tools for subclonal

copy number decomposition [7]; it searches all possible combinations of copy numbers

across all segments and applies the maximum likelihood approach to infer the most

likely subclonal structures. However, THetA has an identifiability concern, such that

several solutions of subclone structures and copy number status levels can explain the

read-depth information equally well [15, 16].

Integrating other data, such as single-nucleotide polymorphisms, to jointly analyze

tumor progression is a solution to the identification problem. The methods developed

on the basis of these integrated data types constitute another category of cancer

subclone reconstruction approaches [14-18]. In 2014, Oesper et al. modified THetA to

THetA2, which designs a probabilistic model of B-allele frequencies (BAFs) to solve

the identification problem and simultaneously improves the efficiency of the algorithm

[14]. Furthermore, PyLOH resolves the identifiability problem by integrating CNAs

and loss of heterozygosity (LOH) within a unified probabilistic model [15]. PyLOH

aims at determining the contamination from normal cells and evaluating tumor purity,

which is the fraction of tumor cells within a tumor tissue. Instead of tumor purity,

MixClone improves PyLOH with a more delicate measurement of tumor progression,

the subclonal cellular prevalence (SCP) [16]. The major concept of PyLOH and

MixClone is to use the Poisson and binomial models simultaneously to analyze the read

depth and BAFs.

Most of the above mentioned methods that reconstruct the process of copy number

evolution assume heterozygous SNV sites within chromosome segments. This

assumption facilitates the decomposition of clonal CNAs, but it ignores segments

without any somatic SNVs. Therefore, to more effectively address this concern, we

developed a new algorithm, called CloneDeMix, which considers subclonal copy

number changes when inferring the clonal evolutionary structures. It requires only the

read-depth information of loci of any sizes no matter SNVs are included or not. The

input can be a predefined segment of the chromosome or simply a single nucleotide

locus. CloneDeMix is a two-way clustering model that clusters each locus into an

appropriate copy number state and a most likely clonal group. The procedure can

simultaneously evaluate all loci and regions. The algorithm uses information from all

samples and loci simultaneously to infer clone progression and can efficiently reduce

the identification bias. The flowchart of CloneDeMix is demonstrated in Fig 1.

In this study, we demonstrated the performance of the algorithm with simulation

data and applied it to a head and neck cancer dataset from The Cancer Genome Atlas

(TCGA) and primary esophageal squamous cell carcinoma (ESCC) [19]. The

simulation demonstrated the accuracy of clone identification and subclonal copy

number change detection, particularly in early mutational events, which could be the candidate of driver mutations. The specificity of the copy number detection exceeded 98%, and the sensitivity was nearly 93.5%. These simulations support that our approach can successfully identify the copy number mutation and deconvolute its amplification or deletion state from the clonal architecture.

Our results obtained for 75 paired normal–tumor samples recapitulated most of the findings reported in head and neck cancer [20-23]. The novel subclonal CNAs have also been identified, and their subclonal structure has been shown to facilitate the discovery of driver mutations for advanced tumor progression. Furthermore, we provide evidence for the association between tumor heterogeneity and metastasis. A large heterogeneity tends to promote tumor metastasis. To sum up, CloneDeMix demonstrated ability to accurately identify subclonal CNAs and clarify intratumor heterogeneity. It is useful complement to other methods for cancer evolution studies.

Fig 1. Flowchart of CloneDeMix

Our approach includes three main steps, data preparation, running CloneDeMix, and inference of tumor heterogeneity.

Methods

Two-way Poisson mixture model

We delineated the structure of cellular evolution based on two concepts: SCP and mutational cellular prevalence (MCP), as shown in Fig 2. The SCP is defined as the fraction of cells that are relatively homogeneous and carry the same set of mutations. The MCP is defined as the fraction of cells that carry a certain mutation. The SCPs can be added to match the MCPs according to the evolutionary structure of subclones (Fig 3A). The evolution matrix, an upper triangular matrix, in Fig 3A provides information on the ancestral relationship among the subclones. There are five subclones in this toy example and their relationship is shown in the evolution tree in Fig 3A. The percentages indicate the corresponding SCPs. In this evolutionary structure, six mutations create five subclones. For example, locus A exists in every tumor subclone because of its presence at the top of the tree. Hence, the MCP of this locus can be calculated as the sum of all SCPs. By contrast, locus G is a later mutation and only exists in the leaf subclone C4. The corresponding MCP is equal to the SCP of C4.

Fig 2. Illustration of SCP and MCP

A tissue has two decompositions. Panel (A) provides an overhead view that divides the cells into several disjoint groups according to their mutations. The cells in the same group are relatively homogeneous and carry the same set of mutations. The size of a group or the fraction of cells is called the SCP. In contrast to the SCP, panel (B) demonstrates the MCP, which is defined as the fraction of cells carrying a certain mutation.

Fig 3. Two-way mixture model for inferring tumor progression by using copy numbers

(A) A toy example for tumor progression of five distinct subclones. Six of the ten loci (A, B, E, F, G, and J) have gained or lost copy numbers, and the remaining loci (C, D, H, and I) show no copy number change. The mutation in each locus forms a new subclone. MCPs can be determined by multiplying the SCP and evolution matrix. (B) The copy number status of each locus is listed in the table, and the MCP of each locus is listed under the table. (C) Each locus belongs to one of the 21 clusters in CloneDeMix. The columns represent five MCP levels, and the rows represent five copy number states considered in the example.

169   The read depth of each locus is proportional to the copy number and MCP. To

170 delineate the read depth of each somatic copy number variant into its copy number state

171 and MCP, this study proposes a two-way mixture model (CloneDeMix). Any locus in a

172 sample has only two states, namely normal and mutated states; the proportion of both

173 types differs across different loci. For example, locus E shows copy number changes in

174 subclone C2 but not in the other subclones (Fig 3B). Hence, all other subclones

175 comprise the normal allele for locus E. Furthermore, locus F has copy number changes

176 in C3, C4, and C5; hence, it is classified as normal in subclones C1 and C2.

177 CloneDeMix clusters all loci according to their copy number state and MCP. As shown

178 in Fig 3C, all loci in this case are classified into five copy number states and

179 simultaneously into five MCP levels. This results in 21 groups because we could not

180 distinguish the MCP levels for the loci of two copies. The MCPs for the five MCP

181 groups are unknown and have to be estimated. Thus, CloneDeMix provides the copy

182 number and MCP for each locus.

183   The input in CloneDeMix is the read depth of each analyzed locus. When the locus

184 represents a segment, such as an exon or a predefined amplicon, the average read depth

185 is adopted. Let X_i be the read depth of locus i or the average read depth rounded to the

closest integer in region i, and assume that it follows a two-way mixture Poisson

distribution.

$$P\big(X_i\big|\{r_h\},\{\pi_{kh}\},a_{base,i}\big) = \sum_{k=1}^{m_1} \sum_{h=1}^{m_2} \pi_{kh} f_{kh}(X_i|r_h, a_{base,i}) \ \forall \ i$$

Each component $f_{kh}(X_i)$ in the model represents the distribution of read depths

sampled from the k-th and h-th groups of the copy number state and MCPs, respectively.

The read count for each combined group is specified as a Poisson distribution; the mean

of this distribution is proportional to a function of the mutated copy number and the

MCP. It is specified as

$$\mu_{hk} = a_{base,i} \times \big(\ 2(1 - r_h) + c_k r_h \ \big),$$

where r_h is the MCP for the h-th group, c_k is the copy number of the k-th copy number

group, and $a_{base,i}$ is a normalization number for locus i. The corresponding mixture

weight is denoted as π_{kh}. Without further evidence, the copy number of the normal cells

can be considered to be two in CloneDeMix. The number of groups for copy numbers

and cellular proportions are pre-specified as m_1 and m_2, respectively; we select m_1 and

m_2 according to the Akaike information criterion (AIC).

Estimating MCPs and copy number by using expectation–maximization algorithm

The parameters of CloneDeMix include the normalization constants $a_{base,i}$, MCPs

$\mathbf{r} = \{r_h\}$, and weights $\Pi = \{ \pi_{kh} \}$. The plug-in estimator of $a_{base,i}$ is estimated from the

paired normal sample of each tumor sample. Because all samples are assumed to be

globally normalized, and the sample-specific variation is removed before the analysis,

the read depth of locus i in the normal sample represents an unbiased estimator of the

mean read depth in the tumor sample when the copy number is two. Hence, we use half

of the read depth of locus i in the paired normal sample as the estimator of $a_{base,i}$. In

case of no paired normal sample, we suggest taking half of the sample mean across all

existing normal samples to estimate $a_{base,i}$. All other parameters are estimated using the

expectation–maximization (EM) algorithm to approximate the maximum likelihood

estimation (MLE).

We introduce a sequence of latent binary variables for locus i. Variables $Y_i =$

$\{Y_{ikh}\}_{k=1,\dots,m_1;h=1,\dots,m_2}$ take the value of 0 or 1, indicating the memberships of the copy

number and MCP groups for locus i. If $Y_{ikh} = 1$, then $X_i|Y_{ikh} = 1, c_k, r_h, \hat{a}_{base,i}$ has

the following distribution

$$f_{kh}(X_i) = Poisson\left(X_i|\mu_{hk} = \hat{a}_{base,i} \times \left(2(1-r_h) + c_k r_h\right)\right). \tag{2}$$

219   A complete form of the conditional distribution is

$$P\left(X_i|Z_i, Y_i, \tilde{r}, \hat{a}_{base,i}\right) = \prod_h \prod_k f_{kh}\left(X_i|\hat{a}_{base,i}\right)^{Y_{ikh}}. \tag{3}$$

220   According to the mixture model construction, the probability of $Y_{ikh} = 1$ is π_{kh}.

221   Specifically,

$$P(Y_{ikh} = 1) = \pi_{kh} \text{ for each locus } i. \tag{4}$$

222   Hence, the density functions of $Y_i = \{Y_{ikh}\}_{k=1,\dots,m_1;h=1,\dots,m_2}$ follow multinomial

223   distributions with probability functions

$$P(Y_i|\Pi) = \prod_h \prod_k \pi_{kh}{}^{Y_{ikh}}. \tag{5}$$

224   According to the definition of conditional probability, the joint density function of

225   X_i and Y_i can be written as follows:

$$
\begin{aligned}
P(X_i, Y_i | \Pi, \tilde{r}, \hat{a}_{base,i}) \ &= P(X_i | Y_i, \tilde{r}, \hat{a}_{base,i}) P(Y_i | \Pi) \\
&= \prod_h \prod_k f_{kh}(X_i | \hat{a}_{base,i})^{Y_{ikh}} \prod_h \prod_k \pi_{kh}{}^{Y_{ikh}} \\
&= \prod_h \prod_k [\pi_{kh} f_{kh}(X_i | \hat{a}_{base,i})]^{Y_{ikh}}. \quad (6)
\end{aligned}
$$

226   The log likelihood of Π and $\tilde{r}$ is

$$
\begin{aligned}
l(\Pi, \tilde{r} | X, Y) \ &= log \prod_i P(X_i, Y_i | \Pi, \tilde{r}, \hat{a}_{base,i}) \\
&= \sum_i \sum_h \sum_k Y_{ikh} log(f_{kh}(X_i | \hat{a}_{base,i}) \pi_{kh}) \quad (7)
\end{aligned}
$$

Because there is no closed form for the maximum likelihood estimator of Π and

$\tilde{r}$, we adopted the EM algorithm to determine the MLE. The EM algorithm iteratively

maximizes the expected log likelihood in two steps: E and M steps.

The E step of the EM algorithm determines the expected value of the log likelihood

over the value of the latent variable Y, given the observed data X and current parameter

value $\Pi = \Pi^0$ and $\tilde{r} = \tilde{r}^0$. Thus, we derive the following equation:

$$
\begin{aligned}
&E_{Y|\Pi^0, c, \tilde{r}^0, X} [l(\Pi, \tilde{r} | X, Z, Y)] \\
&= E_{Y|\Pi^0, c, \tilde{r}^0, X} \left[\sum_i \sum_h \sum_k Y_{ikh} log(f_{kh}(X_i | \hat{a}_{base,i}) \pi_{kh}) \right] \\
&= \sum_i \sum_h \sum_k E_{Y|\Pi^0, c, \tilde{r}^0, X} \left[Y_{ikh} log(f_{kh}(X_i | \hat{a}_{base,i}) \pi_{kh}) \right] \\
&= \sum_i \sum_h \sum_k E_{Y|\Pi^0, c, \tilde{r}^0, X} [Y_{ikh}] \times log(f_{kh}(X_i | \hat{a}_{base,i}) \pi_{kh}) \quad (8)
\end{aligned}
$$

233   According to the definition of Y_{ikh},

$$
\begin{aligned}
E_{Y|\Pi^0, X}[Y_{ikh}] \ &= 1 \times P(Y_{ikh} = 1 | \Pi^0, X) + 0 \times P(Z_{ikh} = 0 | \Pi^0, X) \\
&= \frac{P(X_i | \Pi^0, c, \tilde{r}^0, Y_{ikh} = 1) \times P(Y_{ikh} = 1 | \Pi^0)}{P(X_i | \Pi^0)} \\
&= \frac{P(X_i | \Pi^0, c, \tilde{r}^0, Y_{ikh} = 1) \times P(Y_{ikh} = 1 | \Pi^0)}{\sum_k \sum_h [P(X_i | c, \tilde{r}^0, \Pi^0, Y_{ikh}) P(Y_{ikh} | \Pi^0)]}
\end{aligned}
$$

$$= \frac{f_{kh}(X_i \mid \hat{a}_{base,i}) \times \pi_{kh}^0}{\sum_k \sum_h [f_{kh}(X_i \mid \hat{a}_{base,i}) \times \pi_{kh}^0]} \tag{9}$$

234   Let $E_{Y \mid \Pi^0, X}[Y_{ikh}] = Y_{ikh}^0$ and substitute it into equation (8); with some

235   arrangement, we obtain

$$E_{Y \mid \Pi^0, c, \tilde{r}^0, X} \left[l(\Pi, \tilde{r} \mid X, Z, Y) \right]$$

$$= \sum_i \sum_h \sum_k E_{Y \mid \Pi^0, c, \tilde{r}^0, X}[Y_{ikh}] \times \log(f_{kh}(X_i \mid \hat{a}_{base,i}) \pi_{kh})$$

$$= \sum_i \sum_h \sum_k Y_{ikh}^0 \times \log(f_{kh}(X_i \mid \hat{a}_{base,i}) \pi_{kh})$$

$$= \sum_i \sum_h \sum_k Y_{ikh}^0 \times [\log(f_{kh}(X_i \mid \hat{a}_{base,i})) + \log(\pi_{kh})] \tag{10}$$

236   The M step of the EM algorithm maximizes equation (10) over $\Pi, \tilde{r}$ to determine

237   the next estimates (e.g., Π^1 and $\tilde{r}^1$). The maximization over Π involves only the

238   second term in equation (10):

$$\pi_{kh}^1 = argmax_{\pi_{kh}} (\sum_i \sum_h Y_{ikh}^0 \times \log(\pi_{kh})) \text{ under } \sum \pi_{kh} = 1 \tag{11}$$

239   The solution is $\pi_{kh}^1 = \sum_{i=1}^n Y_{ikh}^0 / n$. The maximization of $\tilde{r}$ concerns the first

240   term of equation (10), and the solution has no closed form. Numeric algorithms, such

241   as the Newton–Raphson method, are required to solve the equation. We used the

242   Newton–Raphson method with the R function *optim*(), and the iterative algorithm for

243    $\tilde{r}$ is

$$\tilde{r}^1 = argmax_{\tilde{r}} \left(\sum_i \sum_h Y_{ikh}^0 \times log \left(f_{kh}(X_i \mid \hat{a}_{base,i}) \right) \right)$$

$$= argmax_{\tilde{r}} \left(\sum_i \sum_h Y_{ikh}^0 \times [-\hat{a}_{base,i} \times (2(1 - r_h) + c_k r_h) + X_i \log(\hat{a}_{base,i} \times (2(1 - r_h) + c_k r_h))] \right) \tag{12}$$

244   The solutions $(\Pi^1, \tilde{r}^1)$ are substituted into equation (10) to replace $(\Pi^0, \tilde{r}^0)$. The

245   expectation is then rewritten as $E_{Y \mid \Pi^1, c, \tilde{r}^1, X}[l(\Pi, \tilde{r} \mid X, Y)]$. The algorithm continues

246   iteratively to maximize the expectation of the log likelihood.

Determining the order of copy number variants

Based on the subclone size inferred using two-way cluster modeling, we can

determine the order of any pairs of recurrent mutations existing in multiple samples.

Herein, we use the notation MCP $\hat{r}_{ij}$ to indicate the estimated MCP of mutation i from

the model of sample j. If a pair of mutations is recurrent in tumors with a fixed order,

the relative size of their estimated MCPs should be consistent. For any two loci a and

b with somatic mutations, the MCP profiles across p samples are $(\hat{r}_{a1}, \dots \hat{r}_{ap})$ and

$(\hat{r}_{b1}, \dots \hat{r}_{bp})$. To determine whether the two mutations are highly related, the Wilcoxon

signed-rank test can be applied to the profiles of the two mutations. In the event of

significant inequality, when one mutation is more common in cells than the other

mutation, it indicates a recurrent evolutionary order between the two mutations.

Results

In this study, we first evaluated the prediction accuracy of CloneDeMix by

simulated data. Simulation study is useful to verify how well an algorithm behaves with

data generated from the theory, but it cannot inform us how well the theory fits reality.

To that end, we collected normal RNA sequences from TCGA and applied down-

sampling to these normal data to create artificial copy number changes. We used the

data to compare CloneDeMix with THetA by evaluating weighted root mean square

error of MCP estimation and positive rate of copy number prediction. We also applied

CloneDeMix on head and neck cancer data from TCGA and serial biopsies of

esophageal cancer [19] to infer genomic evolution based on copy number change.

Simulation

The simulation considered four variant states of copy numbers, namely 0, 1, 3, and

4 copies. Four MCPs were included: 0.1, 0.3, 0.5, 0.7, and 0.9. Each combination was

repeated three times, thus resulting in 60 regions with copy number changes.

Furthermore, each region was assigned 20 bases generated with a Poisson distribution

whose mean value was determined by its assigned copy number state and MCP. In

addition to the mutated regions, 100 normal regions were scattered among the mutated

regions; their copy number state was two. The simulation generated depths for a long

sequence with 3,200 bases for each of the 60 samples. CloneDeMix was subsequently

applied to the simulation data to reconstruct respective copy number states and MCP

groups. The entire simulation process was repeated 10,000 times to obtain a conclusion.

The simulation was performed to evaluate the model estimation accuracy. Table 1

shows the mean and standard deviation (SD) of simulation results for MCP estimation,

and Fig 4 demonstrates the accuracy of assignments for copy number states. As

presented in Table 1, the MCP estimates were very close to the underlying truth,

indicating high performance for MCP estimation. Notably, the bias decreased as the

ground value of the MCP increased. Detecting mutations of low cellular prevalence was

relatively difficult because the signal was not adequately strong.

As illustrated in Fig 4, the accuracy of the copy number assignment under each

condition was calculated from 10,000 simulations. The specificity of CloneDeMix was

found to be 99.58%, and the sensitivity for amplification and deletion were 93.65% and

93.89%, respectively. Thus, CloneDeMix represents a high specificity and efficiently

controls false positive results. As mentioned in the discussion on MCP estimation,

estimating mutations of low cellular prevalence was biased. These biased MCPs

directly caused the misclassification of the copy number state and reduced the model

sensitivity. In conclusion, these simulations support that CloneDeMix can successfully

identify the potential copy number mutations and deconvolute its amplification or

deletion state from the clonal architecture.

Table 1. Mean and SD of MCP estimation

True value	Summary statistics	
	Mean	**SD**
0.1	0.100	0.0067
0.3	0.300	0.0036
0.5	0.499	0.0022
0.7	0.699	0.0014
0.9	0.900	0.0008

Fig 4.   Result of copy number estimation

The size of the circle is proportional to the number of loci assigned to each estimated status from 10,000 simulations. The CNA status is divided into three conditions: deletion, amplification, and normal conditions.

Comparison with THetA2

In this section, we evaluated CloneDeMix on a more realistic simulation scenario

and compared it with THetA2. The core concept of this simulation scenario is the use

of down-sampling technique to resample reads of real normal sequencing data with

artificial copy number changes.

To that end, we first collected 75 normal samples from TCGA and then performed

standard quintile normalization to reduce noise. For simplicity, we only used

chromosome 1 for validation, and chromosome 1 was first cut into 200 different regions.

According to the raw data, we have the raw read counts of each region per sample. The

75 samples were equally divided into case and control. In the control group, the

resampled read count of each region was generated from a binomial distribution. For

the parameter setting of a binomial distribution, the number of trials is set as two times

raw read count and the success probability is 0.5. This procedure is called down-

sampling and it guarantees the mean of resampled count is the same as the mean of raw

count. In the case group, we need to randomly assign 20 regions to have copy number

change. The resampled read count of CNA region also followed the binomial

distribution with the number of trials equal to two times of the raw read count, but the

success probability is set as $0.5\times(2\times(1\text{-MCP})+C\times\text{MCP})/2$ which is determined by a

predefined copy number C and MCP. The predefined copy number of a variant was set

to be 0, 1, 3, and 4. The MCP was set to be 15 different values ranging from 0.1 to 0.9

as shown in Fig 5.

Most studies integrate CNAs and single nucleotide change to improve the accuracy

of copy number identification and to reduce the bias of cellular prevalence estimation.

However, those approaches only study the regions that contain single nucleotide change,

and this constraint apparently limits our understanding of the chromosome structure

change. It has been reported that CNAs affect a larger fraction of the genome in cancers

than any other type of somatic genetic mutation does [23]. For example, a large-scale

study of somatic CNAs across different cancers shows that in a typical cancer sample,

17% of the genome was amplified and 16% genome was deleted on average [24]. Hence,

for a fair comparison, we only compared CloneDeMix with THetA2 because THetA2

is also a subclonal copy number decomposition method and supports direct tumor

heterogeneity inference without considering SNVs.

Both of CloneDeMix and THetA2 are developed for multiple clone identification,

but THetA2 tends to identify single clone in our experience. Therefore, we designed

the resampled data as a mixture of normal cells and one subclone of tumor cells. In this

simple case, the MCP is equal to the tumor purity and we explored the model

performance in different purity. In Fig 5A, we measured the performance of purity

estimation by weighted root mean square error (WRMSE) which is a type of adjusted

RMSE. WRMSE adopts the inverse of true purity as the weight for adjustment because

the variance of purity estimation is a function of the true purity. The variation of purity

estimate increases when the purity increases. Across the 15 different purity settings,

CloneDeMix outperforms THetA2 on measuring purity as demonstrated in Fig 5A. It

is notable that the WRMSEs of THetA2 are missing zero in Fig 5 at low purity settings

(0.1, and 0.16) because THetA2 cannot identify tumor population at low tumor purity.

We calculated the true positive rate (TPR) and false positive rate (FPR) of copy number

assignment at different purity levels in Fig 5B and Fig 5C. We found that both of them

performed well when tumor purity was larger than 0.5. CloneDeMix outperformed

THetA2 in the low purity. It indicates CloneDeMix and THetA2 are equally well at

exploring large subclones while CloneDeMix has better detection power for small

subclones.

Fig 5. Comparison of CloneDeMix and THetA2 with resampled data
(A) The Y-axis is the weighted root mean square error (WRMSE) for measuring the performance of MCP (or purity) estimate, and X-axis represents the true purity setting. (B) The true positive rate (TPR) of copy number detection. (C) The false positive rate of copy number detection.

Preprocessing of TCGA data

We analyzed the whole-exon sequencing data of 75 head and neck tumor samples

with their paired normal samples from TCGA (http://cancergenome.nih.gov/). This

dataset includes a total of 20,846 genes with 180,243 exons. We assumed the copy

number state of a single exon to be homogeneous. Each exon was represented by the

mean read depth. The read-depth profile of a tumor sample was normalized with loess

transformation against its paired normal sample. The baseline parameter $a_{base,i}$ for

exon i was estimated from the paired normal sample by using half of the read depth of

the normal sample at the same locus. Because the normal sample could also have an

abnormal copy number status, we checked it against all other normal samples. The

target normal sample was first normalized against all other normal samples by using

the cyclic loess method and was subsequently processed through CloneDeMix to

identify the copy number status at each locus. In this step, the average profile of all

other normal samples was treated as the baseline. If, for example, an abnormal copy

number is found to be k, the raw read depth of this locus would be divided by k to

provide the estimate of $a_{base,i}$ for tumor modeling.

Copy number distribution and clone structure

We applied CloneDeMix to each normalized sample and estimated the copy

number state of each locus as well as the corresponding MCPs. Fig 6A shows the

chromosomes that were mutated most frequently, and the results of all other

chromosomes are shown in S1 Fig. This figure presents the copy number events across

180,243 exons for each of the 75 tumor–control sample pair. The proportion of exons

with a normal copy number was high in all samples, and it was close to 100% in the

control samples. The proportion was significantly decreased in the tumor samples,

indicating considerable structural variations during cancer development.

Fig 6.   **Copy number estimation of chromosomes with high mutation rates**

(A) The estimated copy number states for exons across the genome are presented by different colors. Light blue and red represent the deletion and amplification events, respectively. Black indicates no copy number changes. (B) The black dots indicate the estimated MCPs with respect to the left axis. The red bars represent the number of MCPs with respect to the right axis.

On average, 4.7% and 8.7% of exons were estimated to have deletion and

amplification, respectively. We also found that the exons located at 3p, 21p, and 18q

were deleted most frequently, and the average proportions of deletion within these

chromosomal arms were 19%, 17%, and 13%, respectively. Conversely, the estimated

amplification frequently occurred at 3q, 8q, and 5p, with average frequency levels of

29%, 24%, and 23%, respectively. Previous studies have reported a loss of 3p and 8p

as well as gains of 3q, 5p, and 8q not only in head and neck cancer but also in most

tumors [20-23]; these results are concordant with our findings. Other novel subclonal

CNA regions that were not reported in pan-cancer data analysis [20-23] were identified

as multiple tumor subpopulations were considered (e.g. Deletion in 21p, S1 Fig). These

subclonal CNA signals may be diluted in the previous studies that assumed only one

homogeneous tumor clone and inferred CNAs from the average of whole tumor

information. Our results confirm the identification strength of large-scale structural

variations based on clonal evolution.

Fig 6B presents a summary of MCP estimation. The number of MCPs was

determined using the model selection criterion AIC. We associated the number of

subclones in the tumors with clinical outcomes because this number is closely related

to tumor heterogeneity. The target phenotype included tumor invasion and metastasis,

which are particularly ominous signs of poor prognosis in head and neck cancer. The

association analysis was applied to only 68 samples because the clinical records of the

other samples were incomplete in TCGA. Fig 7A illustrates the box plot of the number

of MCP groups under each clinical group. In this figure, a sample is denoted as "NO"

if no record of either invasion or metastasis exists; otherwise, it is denoted as "YES."

There appeared to be a tendency of increased tumor heterogeneity for tumors with

invasion or metastasis. The variation of numbers of MCPs was larger for this group. To

more comprehensively clarify this factor, we dichotomized the number of MCPs into

two groups. The number of MCPs exceeding 4 indicated strong tumor heterogeneity,

whereas a lower number indicated less heterogeneity. The contingency table (Fig 7B)

shows the dichotomization of tumor heterogeneity associated with the clinical

outcomes. The corresponding odds ratio was 3.64, and the p value evaluated with

logistic regression was 0.029. For the samples with higher tumor heterogeneity, the

odds of invasion and metastasis were 3.64 times higher than those for the samples with

lower tumor heterogeneity. In recent studies of head and neck cancer, this association

between tumor heterogeneity and metastasis was explored by whole exome sequencing

and single cell RNA sequencing [25-27]. These studies also found the difference in

tumor heterogeneity between primary and matched lymph node metastases samples.

We further investigated the association of overall patient survival and tumor

heterogeneity by survival analysis, and used two different ways to demonstrate this

association. First, we directly considered the subclone number as a covariate of survival

analysis, and then applied Cox model to analyze the effect of subclones. We got a p-

value, 0.036, by Wald's test, and apparently tumor heterogeneity is a risk factor for

survival. Next, we considered three different tumor heterogeneity levels of samples and

performed Kaplan-Meier (KM) curve for different levels. To this end, all of the samples

are divided into three classes by its subclones number, low-heterogeneity (less than 5

subclones), median-heterogeneity ($5 \leq$ subclone number ≤ 8), and high-heterogeneity

(large than 8 subclones). The sample sizes of the three classes are 20, 36, and 19,

respectively. Fig 8 showed the survival curves of the three classes with different colors,

and the survival curve of high-heterogeneity samples is worse than the others. Hence,

high-heterogeneity is associated with poor overall survival. It indicates the tumor

behavior varies with its heterogeneity. The heterogeneity and mortality in head and neck

cancer was also investigated by a different approach [26], and it also concluded that

high-heterogeneity in tumors had doubled the hazard of death.

Fig 7. Comparison for the number of MCPs in different clinical groups
(A) The box plot for the number of MCPs with and without invasion or metastasis. The number of MCPs in each sample is represented by a black point jittered around the box. (B) Contingency table for dichotomization of tumor heterogeneity and clinical outcomes.

Fig 8. Survival curves between different classes of heterogeneity levels
There are three Kaplan-Meier (KM) curves. The blue, yellow, and green represent the group of low, median, and high heterogeneity, respectively.

Inference of evolutionary order of mutations

As stated in the Methods section, we inferred the evolutionary order of recurrent

variants with multiple samples. For easy comprehension, we demonstrated the result at

the gene level through a series of summary steps. We first selected the genes with

consistent amplification or deletion states in more than 25% of the exons within at least

one sample. A total of 3,244 genes were included in this demonstration, and this set is

called the background gene set. For each sample, the MCP of a gene was represented

by the mean MCP of its exons. We then performed the Wilcoxon signed-rank test using

the gene-level MCP of any two genes across the samples to derive all pairwise

evolutionary relationships. For example, if the MCP of gene i was larger than that of

gene *j* (p = 0.05), the mutation on gene *i* was more likely to be an earlier event than that

on gene *j*. This relationship was marked as 1; otherwise, it was marked as 0. The 0–1

matrices of pairwise evolutionary relationships were separately calculated for samples

with and without nodal metastasis, and they could be denoted as a matrix M_{neg} and

M_{pos}. The element of the matrix could be denoted as $M_{E,ij}$, representing the

evolutionary order of mutations on gene *i* against mutations on gene *j* inferred with

samples under the E condition, which could be *neg* or *pos*.

The evolutionary order matrix can be used to construct an evolutionary tree of all

mutations. However, a tree of 3,244 genes is highly complicated, rendering the

comparison of different clinical traits difficult. Therefore, for simplification, we

proposed a progression score to summarize the relative position of a mutation on the

evolutionary tree of tumor formation. The scores of a gene in advanced tumors can be

compared with those of genes in newly developed tumors to select the ones that

recurrently occur in the early stage of tumor development. The P score of gene *i* under

condition E is thus defined as a summary statistic from the evolution matrix and is

formulated as follows:

$$\text{P score (gene } i \,|\text{E)} \; = \; \sum_{j \neq i} M_{E,ij} / (\sum_{j \neq i} M_{E,ij} + \sum_{k \neq i} M_{E,ki}). \qquad (13)$$

Among all relations of gene *i* with other genes, the P score provides the number of times

the mutation in gene *i* is more likely to occur before that in other genes. If a gene is

close to the root of an evolutionary tree, its corresponding P score must be higher than

that of its descending gene.

We first investigated the P-score behavior of prevalent genes which have been

discussed in head and neck cancer [22], and the results are listed in Table 2. The P-score

of PIK3CA is consistently larger than 0.9 across different clinical traits. That is, the

mutation of PIK3CA occurs early in the tumor progression. In contrast, patients with

perineural invasion acquire early mutation of CDKN2A gene more often. Some of the

well-known cancer genes are not powerful in our P-score analysis. For example, we

identified structure variation of TP53 only in a few patients, and these few MCPs are

not enough to construct a powerful P-score.

We compared the P score between the samples with and without nodal metastasis

by plotting a scatter plot (Fig 9). Most background genes tend to mutate in a random

order not related to tumor progression. According to our P score definition, we

postulated that the driving genes of lymph node metastasis would be scattered above

the diagonal line. The genes above the diagonal line of the plot are more likely to

acquire mutations at an earlier stage of tumor formation and occupy a significant

proportion of the tumor at its advanced stage. This would yield higher P scores when

only the samples with lymph node metastasis are considered. By contrast, the

prevalence of mutations in those genes might be low in the samples without lymph node

metastasis and hence yield lower P scores.

Table 2. The P-score of CDKN2A, E2F1, and PIK3CA in different clinical outcomes

	All patients	**Margin status**		**Vital status**		**ECS**		**Invasion**	
		Positive	**Negative**	**Dead**	**Alive**	**Positive**	**Negative**	**Yes**	**No**
CDKN2A	0.665	0.676	0.703	0.747	0.511	0.593	0.625	0.875	0.292
E2F1	0.726	0.092	0.840	0.697	0.889	0.618	0.899	0.078	0.997
FAT1	0.776	0.937	0.714	0.714	0.931	0.683	0.657	0.687	0.866
HAS2	0.081	0.041	0.078	0.016	0.371	0.757	0.064	0.351	0.006
TGFBR2	0.591	0.443	0.629	0.697	0.412	0.646	0.643	0.684	0.394
PIK3CA	0.964	0.998	0.955	0.978	0.947	0.990	0.906	0.991	0.924

Fig 9.   Scatter plot of P scores between nodal positive and nodal negative samples

The red points indicate the background gene set. The red curve indicates the loess smoothing curve constructed using all points in the figure. Genes related to cell migration are marked in black. The genes from 11q13.3 are marked in blue. The literature supporting genes are labeled.

To confirm our conjecture, we selected the genes by their biological functions using

ConsensusPathDB web (http://cpdb.molgen.mpg.de/) and investigated whether genes

related to metastasis in the literature are more likely to be distributed above the diagonal

line. Because cell migration is a crucial step in the metastatic cascade, we selected cell-

migration-related genes, which are marked as black in Fig 9. Consequently, we found

that 43 genes had the function of cell migration. Most of these genes were distributed

above the diagonal line of the P score scatter plot, whereas some were distributed below

the diagonal line. Recurrent mutations in these cell migration genes are expected to be

the driving forces for the initiation of lymph node metastasis, consistent with our

observations. For example, HAS2 is a member of the gene family encoding putative

hyaluronan synthases, which control the biosynthesis of hyaluronan and critically

modulate the tumor microenvironment. Several studies have shown that the inhibition

of HAS2 reduced the invasion of oral squamous cell carcinoma [28-30]. Similar to

HAS2, ANGPT1 is located in the upper left corner and has been recently investigated

for the mechanism of lymph node metastasis [31-34]. ANGPT1 plays an important role

in the regulation of vascular development and maintenance of vessel integrity. A study

showed that the activity of ANGPT1 induced the enlargement of tumor blood vessels

to facilitate tumor cell dissemination and increased the ability of metastasis in tumors

[34]. Fibroblast growth factor (FGF)-4 is another notable example. The P score of FGF4

significantly differs in nodal positive and negative patients. FGF4 is a member of the

FGF family and possesses broad mitogenic and cell survival activities. It has been

proposed to be involved in tumor growth, cell proliferation, and lymph node metastasis

[35-37]. In contrast to the black genes located in the upper left corner of the plot in Fig

9, few studies have reported any relationship between the black genes located in the

lower right corner and lymph node metastasis, although they have the same biological

function. A complete literature review of the genes associated with cell migration and

tumor metastasis is presented in S1 Table. The observations suggest that our inference

of the clonal evolutionary order is relevant and can be applied for identifying causal

drivers.

Another notable observation is about the neighboring genes of FGF4. As

mentioned, FGF4 is an important gene for driving lymph node metastasis. It is located

in 11q13.3, which is frequently amplified in head and neck squamous cell carcinoma

[35]. Sugahara also listed several other genes in 11q13.3 that are related to cancer

development, namely *TPCN2*, *MYEOV*, *CCND1*, *ORAOV1*, *TMEM16A*, *FADD*,

PPFIA1, *CTTN*, *SHANK2*, and *DHCR7*. We also assessed their status by using the P

score analysis; the genes are indicated in blue in Fig 9. All these genes were above the

diagonal line. Their corresponding P scores showed considerably significant differences

between patients with and without nodal metastasis. Hence, we postulated that those

genes in 11q13.3 are possibly related to lymph node metastasis in head and neck cancer.

Several previous studies have confirmed this observation, as reported in S2 Table.

Application on serial biopsies of esophageal cancer

We next applied CloneDeMix on multiregional whole-exome sequencing data from 13 primary esophageal squamous cell carcinoma (ESCC) patients [19]. There are 51 tumor regions and 13 matched morphologically normal esophageal tissues sequenced with the mean coverage of 150×. For fair comparison, we selected 11 of 13 patients based on its platform. We also removed patient ESCC07 because we only got two regions successfully aligned to the reference genome. In total, we included 10 patients in this application, and, for each patient, we have four different tumor regions with one matched esophageal tissue. As preprocess of TCGA data, the read-depth profiles of ESCC tumors are normalized with loess transformation against its paired normal sample. For each individual, the paired normal tissue is also used to calculate the estimates of baseline, and then applied CloneDeMix to tumors for gene-specific CNVs and MCPs.

In this application, we aim to explore the variability of evolutionary structure among multiregional tumors by inferring the order of copy number change. For the purpose of studying variability between regions, we only focused on the frequently mutated genes which are informative about tumor evolution. Although the construction through these genes is not able to resolve completely the entire evolutionary structure, the inferred structure between regions can still facilitate the understanding of tumor progression. To that end, we collected the target gene list from the Ion AmpliSeq Comprehensive Cancer Panel which includes 7,044 exons of 409 tumor suppressor genes and oncogenes. The estimated CNVs and MCPs of the ESCC biopsies for this

gene set were summarized and interpreted as follows.

We first investigated genomic heterogeneity of ECSS through MCP comparison.

MCP is a gene-specific measurement of fraction of cells that carry a certain mutation,

and we can study the overall structure of MCPs across whole genome to reveal the

genomic heterogeneity of a given sample. We calculated the correlation matrix of MCP

between samples, and this correlation matrix is presented in Fig 10. The diagonal blocks

of this correlation matrix are tissues of the same sample and are slightly higher than the

others. The average correlation of diagonal block is 0.5 and the average of off-diagonal

cells 0.3. It shows that the MCP structure within each patient is more consistent than

between patients.

Next, we identified the evolution-related genes for each individual. In ESCC study,

each tumor was dissected into four regions, and this kind of serial biopsies has a natural

assumption that the size of MCPs is comparable within a given tumor. This

characteristic can facilitate the individual-specific heterogeneity study. In order to

explore individual-specific heterogeneity, we first identified genes on the trunk and on

the branch of the evolutionary tree separately. The trunk of the tree refers to the CNAs

consisted in all regions while the branch refers to those only in some regions. We can

identify these genes according to the MCP across regions. A gene is located on the trunk

of a tree if its average MCP across four regions is larger than 0.8, and a gene is located

on the branch of a tree if the MCP of one region is larger than the average MCP of all

the remaining regions by 0.7. Instead of tree comparison, we directly compare the MCP

matrix of selected genes (Fig 11). In Fig 11, genes in red rectangles are selected to be

the trunk genes, and the remaining genes are on the branch of a tree.

The two types of genes defined above reveals huge variability of evolution

structure across tumors. The genes on the trunk of a given tree represent the genes

changed in copy number at an earlier stage of tumor formation, and these genes have

potential ability to drive tumor growth. For example, most of the genes on the trunk of

sample ESCC12 (CCND1, EGFR, APC, TGFBR2, XPC, XPA, FLI1, and NUMA1)

have been identified and initially reported on the esophageal cancer [38-42]. Although

the genes on the trunks of trees vary among different individuals, there are still genes

repeatedly identified in multiple individuals such as CCND1, JAK2, UGT1A1, FLI1,

NFE2L2, SOX2, CDKN2B, and MYC. Specifically, CCND1 was identified in six

individuals as the trunk gene and is a well-known cancer oncogene located on 11q13.

Its amplification has been reported in several human neoplasias [43].

Fig 10.　Correlation matrix of MCP between samples
Each cell indicates the correlation of MCPs between the corresponding ESCC samples.

Fig 11.　MCP matrices of selected genes among 10 samples
There are six MCP matrices. The color of each cell represents the MCP quantity of a gene for a given
sample. The labels of rows indicate the gene symbols, and the labels of columns are region index A
gene within the red rectangle is identified as the gene located on the trunk of an evolutionary tree.

Discussion

In this study, we developed CloneDeMix for the deconvolution of tumor

progression through high-throughput DNA sequencing data. The features of

CloneDeMix are as follows. First, it reconstructs an evolutionary structure of copy

number changes during tumorigenesis. Most existing methods for cancer evolution

discuss the history of single-nucleotide changes and derive the potential driver genes.

However, the importance of CNAs is growing and its influence on disease and cancer

development is clearly established [44]. Therefore, the reconstruction of copy number

evolution in tumor progression is in demand. Second, CloneDeMix provides the MCP

as a measure of the evolutionary structure. This measurement is used to estimate the

fraction of cells containing a specific set of mutational events. According to the

definition of the MCP, it provides a more direct evolutionary reconstruction than does

the SCP, which is defined as the size of a subclone in a tumor. For instance, the MCPs

of early mutations in cancer must exceed those of other mutations, but no such structural

relationship exists for SCPs. Although MCPs of a tumor is related to its phylogenetic

tree, we do not have DNA haplotypes to resolve the tree architecture from many

possibilities for each individual tumor. Hence, in this study, we only borrow the strength

of multiple samples to understand potential evolutionary orders using the P score. Third,

our model exhibits high flexibility. CloneDeMix can identify the copy number state of

any type of variant, from a single nucleotide to a moderate size of regions. Furthermore,

the model facilitates the simultaneous analysis of multiple types of targets because it

depends on only the summary information of each locus.

The simulation study revealed that CloneDeMix can identify the current clonal

structures of a tumor. The accuracy of copy number states was nearly 93%, and the

MCP was also accurately restored (Table 1). Furthermore, the application of

CloneDeMix to head and neck cancer data from TCGA yielded promising putative

CNAs. The deletions observed on chromosomes 3p, 18q, and 21p and the

amplifications on chromosomes 3q, 5p, and 8q are consistent with most cancer studies

on copy number identification [20-23]. This observation strongly supports our CNA

inference procedure.

When the estimation accuracy reaches a certain level, the most important concern

is to understand the relationship between tumor heterogeneity and disease progression.

Tumor clone dynamics have been associated with clinical outcomes for different types

of cancer [45-47]. Our method provides a quantitative measurement of clonality, and it

is associated with tumor invasion and metastasis development in TCGA database.

Tumors with more subclones are a result of complex branched evolution, implying a

series of adaptations to a new environment. These newly emerged subclones may

contribute to metastatic initiation or acquire a new ability to invade the lymphatic or

vascular system. Thus, the strong prognostic association of the number of MCPs with

invasion or metastasis reinforces its clinical relevance; this index appears to be a novel

feature for further exploration.

We established a novel score, the P score, for evaluating the order of a recurrent

mutation in the evolutionary hierarchy by analyzing multiple samples. By comparing

the P scores of a somatic variant between different clinical groups, we could identify

the copy number mutations that occur early in the tumor stage and expand the

accompanied subclones with time. The utility of P scores was also demonstrated in the

head and neck cancer data according to the sample status of metastasis. Furthermore,

we identified a group of genes that matched this condition. Specifically, the genes

located at 11q13.3 are well known to be frequently amplified in head and neck

squamous cell carcinomas. Their P scores in our analysis were particularly high for the

samples with lymph node metastasis and relatively low for those without metastasis.

Accordingly, those gene amplifications are potential causal mutations to drive

metastatic cascade in head and neck cancer. Hence, screening for genes that differ

considerably in their P scores is meaningful for driver gene detection.

The success of our approach highly depends on the coverage of DNA sequencing.

A higher read depth can more efficiently reflect the clonal structure and copy number

changes of different loci. Currently, CloneDeMix makes an independent assumption

without considering the dependency among closely located loci. Hence, the

neighboring segments are not grouped into the same copy number events. This can be

an advantage as well as a disadvantage because there is no clear understanding about

the range covered by a copy number event. Technically, we can still integrate the

correlation structure into CloneDeMix to improve the flexibility; this is an ongoing

project for our next version of the R package.

CloneDeMix can easily integrate different types of somatic mutations detected in

sequencing data. For example, the well-studied SNVs carry extensive information

about the clonal expansion in tumors. CloneDeMix can consider the copy number status

of two alleles individually if the detection of each allele is optimized. Therefore, we

expect CloneDeMix to be useful in understanding tumor heterogeneity and how it

evolves to the current status. Moreover, CloneDeMix has high specificity for detecting

early mutations in tumor progression; these early mutations would be good candidates

for disease driver genes and targeted therapies.

Acknowledgments

669 This work was supported by the Ministry of Science and Technology [MOST 105-2118-

670 M-007-001].

Reference

1.   Nowell PC. The clonal evolution of tumor cell populations. Science.
1976;194(4260):23-8.

2.   Kreso A, O'Brien CA, van Galen P, Gan OI, Notta F, Brown AM, et al. Variable clonal
repopulation dynamics influence chemotherapy response in colorectal cancer. Science.
2013;339(6119):543-8.

3.   Hou Y, Song L, Zhu P, Zhang B, Tao Y, Xu X, et al. Single-cell exome sequencing and
monoclonal evolution of a JAK2-negative myeloproliferative neoplasm. Cell.
2012;148(5):873-85.

4.   Xu X, Hou Y, Yin X, Bao L, Tang A, Song L, et al. Single-cell exome sequencing
reveals single-nucleotide mutation characteristics of a kidney tumor. Cell.
2012;148(5):886-95.

5.   Wang Y, Waters J, Leung ML, Unruh A, Roh W, Shi X, et al. Clonal evolution in
breast cancer revealed by single nucleus genome sequencing. Nature.
2014;512(7513):155-60.

6.   Ding L, Raphael BJ, Chen F, Wendl MC. Advances for studying clonal evolution in
cancer. Cancer letters. 2013;340(2):212-9.

7.   Oesper L, Mahmoody A, Raphael BJ. THetA: inferring intra-tumor heterogeneity
from high-throughput DNA sequencing data. Genome Biol. 2013;14(7):R80.

8.   Malikic S, McPherson AW, Donmez N, Sahinalp CS. Clonality inference in multiple
tumor samples using phylogeny. Bioinformatics. 2015;31(9):1349-56.

9.   El-Kebir M, Oesper L, Acheson-Field H, Raphael BJ. Reconstruction of clonal trees
and tumor composition from multi-sample sequencing data. Bioinformatics.
2015;31(12):i62-i70.

10.   Deshwar AG, Vembu S, Yung CK, Jang GH, Stein L, Morris Q. PhyloWGS:
Reconstructing subclonal composition and evolution from whole-genome sequencing
of tumors. Genome biology. 2015;16(1):35.

11.   Jiao W, Vembu S, Deshwar AG, Stein L, Morris Q. Inferring clonal evolution of
tumors from single nucleotide somatic mutations. BMC bioinformatics. 2014;15(1):35.

12.   Miller CA, White BS, Dees ND, Griffith M, Welch JS, Griffith OL, et al. SciClone:
inferring clonal architecture and tracking the spatial and temporal patterns of tumor
evolution. 2014.

13.   Roth A, Khattra J, Yap D, Wan A, Laks E, Biele J, et al. PyClone: statistical inference
of clonal population structure in cancer. Nature methods. 2014;11(4):396-8.

14.   Oesper L, Satas G, Raphael BJ. Quantifying tumor heterogeneity in whole-genome
and whole-exome sequencing data. Bioinformatics. 2014;30(24):3532-40.

15.   Li Y, Xie X. Deconvolving tumor purity and ploidy by integrating copy number
alterations and loss of heterozygosity. Bioinformatics. 2014:btu174.

16.   Li Y, Xie X. MixClone: a mixture model for inferring tumor subclonal populations.
BMC genomics. 2015;16(Suppl 2):S1.

17.   Yu Z, Li A, Wang M. CloneCNA: detecting subclonal somatic copy number
alterations in heterogeneous tumor samples from whole-exome sequencing data.
BMC bioinformatics. 2016;17(1):310.

18.   Fischer A, Vázquez-García I, Illingworth CJ, Mustonen V. High-definition
reconstruction of clonal composition in cancer. Cell reports. 2014;7(5):1740-52.

19.   Hao J-J, Lin D-C, Dinh HQ, Mayakonda A, Jiang Y-Y, Chang C, et al. Spatial
intratumoral heterogeneity and temporal clonal evolution in esophageal squamous
cell carcinoma. Nature genetics. 2016;48(12):1500.

20.   Koboldt DC, Zhang Q, Larson DE, Shen D, McLellan MD, Lin L, et al. VarScan 2:
somatic mutation and copy number alteration discovery in cancer by exome
sequencing. Genome research. 2012;22(3):568-76.

21.   Krijgsman O, Carvalho B, Meijer GA, Steenbergen RD, Ylstra B. Focal chromosomal
copy number aberrations in cancer—Needles in a genome haystack. Biochimica et
Biophysica Acta (BBA)-Molecular Cell Research. 2014;1843(11):2698-704.

22. Network CGA. Comprehensive genomic characterization of head and neck
squamous cell carcinomas. Nature. 2015;517(7536):576-82.

23. Zack TI, Schumacher SE, Carter SL, Cherniack AD, Saksena G, Tabak B, et al. Pan-
cancer patterns of somatic copy number alteration. Nature genetics.
2013;45(10):1134-40.

24. Beroukhim R, Mermel CH, Porter D, Wei G, Raychaudhuri S, Donovan J, et al. The
landscape of somatic copy-number alteration across human cancers. Nature.
2010;463(7283):899.

25. Puram SV, Tirosh I, Parikh AS, Patel AP, Yizhak K, Gillespie S, et al. Single-Cell
Transcriptomic Analysis of Primary and Metastatic Tumor Ecosystems in Head and
Neck Cancer. Cell. 2017;171(7):1611-24. e24.

26. Mroz EA, Tward AM, Hammon RJ, Ren Y, Rocco JW. Intra-tumor genetic
heterogeneity and mortality in head and neck cancer: analysis of data from the Cancer
Genome Atlas. PLoS medicine. 2015;12(2):e1001786.

27. Mroz EA, Tward AD, Pickering CR, Myers JN, Ferris RL, Rocco JW. High intratumor
genetic heterogeneity is related to worse outcome in patients with head and neck
squamous cell carcinoma. Cancer. 2013;119(16):3034-42.

28. Bernert B, Porsch H, Heldin P. Hyaluronan synthase 2 (HAS2) promotes breast
cancer cell invasion by suppression of tissue metalloproteinase inhibitor 1 (TIMP-1).
Journal of Biological Chemistry. 2011;286(49):42349-59.

29. Zhang Z, Tao D, Zhang P, Liu X, Zhang Y, Cheng J, et al. Hyaluronan synthase 2
expressed by cancer-associated fibroblasts promotes oral cancer invasion. Journal of
Experimental & Clinical Cancer Research. 2016;35(1):181.

30. Li P, Xiang T, Li H, Li Q, Yang B, Huang J, et al. Hyaluronan synthase 2
overexpression is correlated with the tumorigenesis and metastasis of human breast
cancer. International journal of clinical and experimental pathology. 2015;8(10):12101.

31. Fagiani E, Lorentz P, Kopfstein L, Christofori G. Angiopoietin-1 and-2 exert
antagonistic functions in tumor angiogenesis, yet both induce lymphangiogenesis.
Cancer research. 2011;71(17):5717-27.

32. Holopainen T, Huang H, Chen C, Kim KE, Zhang L, Zhou F, et al. Angiopoietin-1
overexpression modulates vascular endothelium to facilitate tumor cell dissemination
and metastasis establishment. Cancer research. 2009;69(11):4656-64.

33. Mangiola S, Hong MK, Cmero M, Kurganovs N, Ryan A, Costello AJ, et al.
Comparing nodal versus bony metastatic spread using tumour phylogenies. Scientific
Reports. 2016;6.

34. Li T, Yang J, Zhou Q, He Y. Molecular regulation of lymphangiogenesis in

development and tumor microenvironment. Cancer Microenvironment.
2012;5(3):249-60.

35. Sugahara K, Michikawa Y, Ishikawa K, Shoji Y, Iwakawa M, Shibahara T, et al.
Combination effects of distinct cores in 11q13 amplification region on cervical lymph
node metastasis of oral squamous cell carcinoma. International journal of oncology.
2011;39(4):761.

36. Muller D, Millon R, Lidereau R, Engelmann A, Bronner G, Flesch H, et al. Frequent
amplification of 11q13 DNA markers is associated with lymph node involvement in
human head and neck squamous cell carcinomas. European Journal of Cancer Part B:
Oral Oncology. 1994;30(2):113-20.

37. Qi L, Song W, Li L, Cao L, Yu Y, Song C, et al. FGF4 induces epithelial-mesenchymal
transition by inducing store-operated calcium entry in lung adenocarcinoma.
Oncotarget. 2016;7(45).

38. Hao J-J, Shi Z-Z, Zhao Z-X, Zhang Y, Gong T, Li C-X, et al. Characterization of genetic
rearrangements in esophageal squamous carcinoma cell lines by a combination of M-
FISH and array-CGH: further confirmation of some split genomic regions in primary
tumors. BMC cancer. 2012;12(1):367.

39. Yang P-W, Hsieh C-Y, Kuo F-T, Huang P-M, Hsu H-H, Kuo S-W, et al. The survival
impact of XPA and XPC genetic polymorphisms on patients with esophageal squamous
cell carcinoma. Annals of surgical oncology. 2013;20(2):562-71.

40. Nomura T, Miyashita M, Makino H, Maruyama H, Katsuta M, Kashiwabara M, et
al. Argon plasma coagulation for the treatment of superficial esophageal carcinoma.
Journal of Nippon Medical School. 2007;74(2):163-7.

41. Petty RD, Dahle-Smith A, Stevenson DA, Osborne A, Massie D, Clark C, et al.
Gefitinib and EGFR gene copy number aberrations in esophageal cancer. Journal of
Clinical Oncology. 2017;35(20):2279-87.

42. Hu X, Moon JW, Li S, Xu W, Wang X, Liu Y, et al. Amplification and overexpression
of CTTN and CCND1 at chromosome 11q13 in Esophagus squamous cell carcinoma
(ESCC) of North Eastern Chinese Population. International journal of medical sciences.
2016;13(11):868.

43. Santarius T, Shipley J, Brewer D, Stratton MR, Cooper CS. A census of amplified
and overexpressed human cancer genes. Nature Reviews Cancer. 2010;10(1):59.

44. Valsesia A, Macé A, Jacquemont S, Beckmann JS, Kutalik Z. The growing
importance of CNVs: new insights for detection and clinical interpretation. Frontiers in
genetics. 2013;4:92.

45. McGranahan N, Furness AJ, Rosenthal R, Ramskov S, Lyngaa R, Saini SK, et al.
Clonal neoantigens elicit T cell immunoreactivity and sensitivity to immune checkpoint

blockade. Science. 2016;351(6280):1463-9.

46. Park SY, Gönen M, Kim HJ, Michor F, Polyak K. Cellular and genetic diversity in the
progression of in situ human breast carcinomas to an invasive phenotype. The Journal
of clinical investigation. 2010;120(2):636-44.

47. Maley CC, Galipeau PC, Finley JC, Wongsurawat VJ, Li X, Sanchez CA, et al. Genetic
clonal diversity predicts progression to esophageal adenocarcinoma. Nature genetics.
2006;38(4):468-73.

Supporting Information

S1 Table. Reference list of cell-migration-related genes

S2 Table. List of reference genes in 11q13.3

S1 Fig. Copy number estimation results

The estimated copy number states for the exons across the genome are presented in different colors. Light blue and red represent the deletion and amplification events, respectively. Black indicates no copy number changes.

S1 Software.

Software S1 is an R package called "CloneDeMix" that implements subclonal copy number decomposition and it is available at https://github.com/AshTai/CloneDeMix.

Data Preparation

DNA sequencing

Normal tissue

Read count

Position

Tumors

···

➢ Define regions of interest.
  (It can be a single base, exon or gene.)

➢ Calculate average read count of regions.

Input

Average read count

Order of region

CloneDeMix

Output

Tumor No.1   *Tumor* No.2   *Tumor* No.3

✓ **Number of subclones**

✓ **Target-specific copy number**

✓ **MCP**

Inference of Heterogeneity

➢ Use MCPs to infer tumor evolution by comparing P-score.

Early event

P-score in case 2

P-score in case 1

Compare P-score

MCP matrix of an individual

A

Evolution matrix

$$\begin{matrix} c1 \\ c2 \\ c3 \\ c4 \\ c5 \end{matrix} \begin{bmatrix} 1 & 1 & 1 & 1 & 1 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix} \times \begin{matrix} \text{SCP} \\ \begin{bmatrix} 5\% \\ 20\% \\ 10\% \\ 25\% \\ 15\% \end{bmatrix} \end{matrix} = \begin{matrix} \text{MCP} \\ \begin{bmatrix} 75\% \\ 20\% \\ 50\% \\ 25\% \\ 15\% \end{bmatrix} \end{matrix}$$

B

Clone (SCP)	Locus									
	A	**B**	**C**	**D**	**E**	**F**	**G**	**H**	**I**	**J**
C1 (5%)	3	1	2	2	2	2	2	2	2	2
C2 (20%)	3	1	2	2	0	2	2	2	2	2
C3 (10%)	3	1	2	2	2	4	2	2	2	2
C4 (25%)	3	1	2	2	2	4	0	2	2	2
C5 (15%)	3	1	2	2	2	4	2	2	2	3

MCP	75%	75%	0%	0%	20%	50%	25%	0%	0%	15%

C

Copy number state	MCP				
	r1 (75%)	**r2 (50%)**	**r3 (25%)**	**r4 (20%)**	**r5 (15%)**
0-copy				E	J
1-copy	B				
2-copy (Normal State)	C, D, H, I				
3-copy	A		G		
4-copy		F			

A

B

A

B

	Clinical trait (Invasion/Metastasis)	
	NO	YES
Numb. of MCPs ≤ 4	16	22
Numb. of MCPs > 5	5	25
Odds Ratio (P-value)	3.64 (p=0.029*)	

Color Key

Correlation of MCPs

ECSS03 - t1
ECSS03 - t2
ECSS03 - t3
ECSS03 - t4
ECSS04 - t1
ECSS04 - t2
ECSS04 - t3
ECSS04 - t4
ECSS05 - t1
ECSS05 - t2
ECSS05 - t3
ECSS05 - t4
ECSS06 - t1
ECSS06 - t2
ECSS06 - t3
ECSS06 - t4
ECSS09 - t1
ECSS09 - t2
ECSS09 - t3
ECSS09 - t4
ECSS10 - t1
ECSS10 - t2
ECSS10 - t3
ECSS10 - t4
ECSS08 - t1
ECSS08 - t2
ECSS08 - t3
ECSS08 - t4
ECSS11 - t1
ECSS11 - t2
ECSS11 - t3
ECSS11 - t4
ECSS12 - t1
ECSS12 - t2
ECSS12 - t3
ECSS12 - t4
ECSS13 - t1
ECSS13 - t2
ECSS13 - t3
ECSS13 - t4

ECSS03 - t1
ECSS03 - t2
ECSS03 - t3
ECSS03 - t4
ECSS04 - t1
ECSS04 - t2
ECSS04 - t3
ECSS04 - t4
ECSS05 - t1
ECSS05 - t2
ECSS05 - t3
ECSS05 - t4
ECSS06 - t1
ECSS06 - t2
ECSS06 - t3
ECSS06 - t4
ECSS09 - t1
ECSS09 - t2
ECSS09 - t3
ECSS09 - t4
ECSS10 - t1
ECSS10 - t2
ECSS10 - t3
ECSS10 - t4
ECSS08 - t1
ECSS08 - t2
ECSS08 - t3
ECSS08 - t4
ECSS11 - t1
ECSS11 - t2
ECSS11 - t3
ECSS11 - t4
ECSS12 - t1
ECSS12 - t2
ECSS12 - t3
ECSS12 - t4
ECSS13 - t1
ECSS13 - t2
ECSS13 - t3
ECSS13 - t4

ESCC-03

ESCC-04

ESCC-05

ESCC-06

ESCC-08

ESCC-09

ESCC-10

ESCC-11

ESCC-12

ESCC-13