Article:

The neuronal ceroid lipofuscinosis protein Cln7 regulates neural development from the
post-synaptic cell.

Kyle J. Connolly* [1], Megan B. O'Hare* [2], Alamin Mohammed* [1], Katelyn M. Aitchison[1], Niki
C. Anthoney[1], Amy L. Roberts[1], Matthew J. Taylor [1], Bryan A. Stewart[3,4], Richard I. Tuxworth#
[1,2] and Guy Tear# [2]

8
9   1. Institute of Cancer and Genomic Sciences
10   University of Birmingham
11   Birmingham
12   B15 2TT
13   UK

14
2: Department of Developmental Neurobiology
King's College London
New Hunt's House
London
SE1 1UL
UK

21
22   3: Department of Biology
23   University of Toronto Mississauga
24   Mississauga, ON
25   L5L 1C6
26   Canada

27
28   4. Department of Cell and Systems Biology
29   University of Toronto
30   Toronto, ON
31   M5S 3G5
32   Canada

33
34
35   * Equal contribution

36
37   # co-senior and co-corresponding authors:
38   Guy.Tear@kcl.ac.uk
39   +44 (0)20 7848 6539

40
R.I.Tuxworth@bham.ac.uk
+44 (0)121 414 7046

ORCIDs:

MJT: 0000-0003-3055-4453
NCA: 0000-0003-3311-6328
KJC: 0000-0003-2425-3910
BAS: 0000-0003-2520-3632
RIT: 0000-0001-5697-6254
GT: 0000-0002-3114-7537

Running Title:
Cln7 regulates synapse development

Abstract

The development and maintenance of synapses requires co-ordination between the

pre- and post-synaptic partners. We demonstrate that in *Drosophila,* Cln7, a MFS-domain

transporter protein, is required to regulate normal synapse development. Cln7 is restricted

to the post-synaptic muscle suggesting it is a novel component of a retrograde pathway

regulating trans-synaptic communication. Loss of Cln7 leads to dysregulation of TORC

activity, including excessive organismal growth and increased lipid storage phenotypes

which are both dependent on TORC2 activity. Cln7 is mutated in late-infantile neuronal

ceroid lipofuscinosis, an early-onset monogenetic neurodegenerative disorder, indicating

that dysregulated TORC activity resulting in developmental synapse pathology may underpin

the disease.

Introduction

The development and maintenance of synapses need to be precisely regulated to

allow accurate flow of signals from the neuron to its target. Following circuit formation,

synapse size continues to be adjusted and maintained through homeostatic mechanisms[1,2].

This plasticity of the synapse allows strengthening or weakening of connections and the

accurate matching of presynaptic signal strength to the size of the postsynaptic target.

Synapse homeostasis relies on anterograde and retrograde signaling across the synapse,

which is mediated by a variety of signals[3,4]. Failure to form, maintain or adjust the synapse

can lead to synapse failure or inefficiency leading to loss of neural function and

degeneration. As such the synapse is a common target for neurodegenerative disease[4,5].

The *Drosophila* neuromuscular junction (NMJ) is a widely used genetically tractable

model system to study synapse formation and homeostasis[6-8]. The *Drosophila* NMJ forms

during embryonic development where the size of the pre-synaptic compartment is aligned

with muscle size. During subsequent larval stages the muscle grows rapidly and synapse size

expands to match this growth by increasing the number of neurotransmitter release sites

(boutons), the amount of neurotransmitter release and the expression level of glutamate

receptor subunits[9]. Retrograde signals such as the BMP ligand, glass bottom boat (gbb), are

released from the post-synaptic side, activating a SMAD pathway in the neuron to influence

growth, reorganization of the cytoskeleton and the secretion of membrane[9]. The expansion

of the pre-synapse is also modulated via a variety of factors acting within neurons including

JNK signaling[10,11], levels of reactive oxygen species[12] autophagic pathways and regulators of

membrane traffic[4]. TORC-based mechanisms also act to co-ordinate postsynaptic glutamate

receptor levels in response to unknown trans-synaptic signals[13].

Many genes associated with human neurological conditions, including

developmental disorders and neurodegenerative diseases, have roles regulating NMJ

development[14,15]. Here we have investigated a role for the gene, *Cln7*, which is mutated in a

form of the childhood neurodegenerative disorder, neuronal ceroid lipofuscinosis (NCL)[16].

Evidence has accumulated in several models that loss of the *CLN* genes impacts on synapse

function, suggesting that they play a common role either within neurons or non-neuronal

cells to maintain function[17-19]. Recently, *Cln7* has been implicated in the activation of mTORC

signaling[20] and given the importance of TORC and autophagy for neuronal growth and

synapse homeostasis, we generated a *Drosophila Cln7* mutant to ask how these processes

might be affected in its absence. We identified a requirement for *Cln7* for normal neural

growth but, surprisingly, *Cln7* is restricted to the post-synaptic side of the NMJ where it

concentrates at the post-synaptic density. Autophagy and retrograde BMP signaling are

largely intact in *Cln7* mutants and instead the neurodevelopmental defect is likely due to

changes in TORC activity in the post-synaptic cell via interactions between Cln7 and Rheb.

This places Cln7 as a novel regulator of a retrograde TORC activity at the synapse and

highlights the importance of this pathway in neural development and disease.

Results

Cln7 is localized in the basal sub-synaptic reticulum

Disruption of the *Cln7* gene has been identified as the cause of a childhood

neurodegenerative disorder, late-infantile neuronal ceroid lipofuscinosis (LINCL/Cln7

disease)[16]. We had previously identified that the *Drosophila* Cln7 protein is expressed in the

body wall muscles that form the post-synaptic cells of the neuromuscular junction (NMJ),

where it concentrates at the post-synaptic density (PSD) (Fig. 1a)[21]. This distribution

suggests the protein may function to regulate the synapse and that LINCL may be a result of

synapse failure. We further examined the localization of Cln7 using a CRIPSR-generated

Venus-YFP-Cln7 knock-in and higher-resolution Airyscan 3D confocal imaging. We find that

Cln7 is restricted to a region overlapping with the basal sub-synaptic reticulum (SSR) (Fig. 1c-

f). The SSR is a convoluted membranous structure membrane in the post synaptic cell that

surrounds the pre-synaptic membrane swellings (the boutons) and contains the post-

synaptic glutamate receptors. Electron microscopy studies have revealed the SSR fully

surrounds the pre-synaptic bouton and anti-Discs large staining (DLG, the *Drosophila*

homologue of PSD-95), which marks the SSR, shows the same distribution (Fig. 1e and f).

Interestingly, the Cln7 protein appears to be associated with vesicles or the SSR membrane

network only on the basal side; Cln7 is not present surrounding the upper surface of the

bouton (Fig. 1c-f). Since retrograde signaling from the post synaptic cell is known to

influence synapse structure[9] we investigated the consequence of a loss of *Cln7* function.

Cln7 is required for normal development of the NMJ

We generated two loss-of-function alleles of *Cln7* by imprecise excision of the P-

element NP0345 inserted in the 5' UTR upstream of the start codon. Two alleles were

generated: exons 1-5 were excised in the *Cln7*[84D] allele, with some P-element sequence

remaining and exons 1 and 2 and part of exon 3 were excised in *Cln7*[36H]. Both mutant alleles

are viable and fertile when homozygous and show no developmental delay. The peripheral

NMJs in late stage *Drosophila* larvae are simple, with 2-4 individual motoneurons contacting

each muscle cell. The type I motoneurons that innervate the muscle can be further

subdivided into two classes called type Ib and type Is classified according to the size of the

presynaptic boutons (Fig. 2a). We find that in *Cln7*[-/-] animals the NMJ is reduced in size. At

muscle 4 we see a significant reduction of the type Ib motoneuron terminals in the *Cln7*

mutant larvae when compared to isogenic control larvae (Fig. 2a). We used a non-subjective

method of measuring terminal volume from 3D-rendered confocal images to confirm the

type Ib terminals were significantly smaller than controls while the type Is terminals appear

unaffected (Fig. 2b). As a further measure of synapse size, we counted the number of active

zones at the NMJ. Active zones are visualized with an antibody to Bruchpilot (ELKS), an

active zone component[22]. Bruchpilot positive puncta were counted at muscle 4 and this

confirmed that the number of active zones within Ib boutons is reduced in *Cln7*[-/-] animals

concomitant with the reduction in volume but is unchanged in type Is junctions (Fig. 2c).

Both alleles of Cln7 had an identical phenotype as does the trans-heterozygous

combination. Both alleles were also haploinsufficient, suggesting some processes important

for NMJ development are critically dependent on the levels of the Cln7 protein: potentially

the stoichiometry of a membrane complex. These results reveal a requirement for Cln7 for

normal development of the NMJ.

160   To investigate whether Cln7 is required in the post synaptic cell we used RNAi to

reduce *Cln7* expression in various cell types at the synapse. Standard GAL4 drivers were

used to express dsRNA specifically targeted at *Cln7* either pre-synaptically (Elav-Gal4), post-

synaptically in the muscle (Mef2-Gal4), in all glia (Repo-Gal4) and in only the perineurial glia

(PG-Gal4) that form the blood-brain-barrier, where *Cln7* is also highly expressed[21]. Only

knockdown of *Cln7* in the muscle replicated the under-development phenotype at the NMJ

(Fig. 2d), consistent with the expression of *Cln7*. We also performed the converse

experiment of re-expressing YFP-Cln7 in the *Cln7*[-/-] background, again with ubiquitous

expression (actin-Gal4) or expression restricted to neurons (Elav-Gal4), to muscle (Mef2-

Gal4), in both neurons and muscle together (Spin-Gal4) or to glia (Repo-Gal4). Muscle

expression with Mef2- or Spin-Gal4 was only capable of partial rescue of function which

failed to reach significance. Only ubiquitous expression fully rescued the phenotype (Fig. 2e)

indicating that some Cln7 function is required elsewhere in addition to the muscle.

Loss of Cln7 leads to synapse dysfunction
176
To examine the functional consequence of the abnormal development of the

synapse we characterized synaptic physiology and larval motility. An electrophysiological

analysis of the NMJ was carried out using intracellular recordings. We recorded the

spontaneous transmitter release at the larval NMJs for two minutes to calculate miniature

excitatory junction potential (mEJP) frequency and amplitude in 0.5, 0.75, 1.0 and 2.0 mM

extracellular Ca^{2+} concentration but there was no significant difference due to genotype in

either case (Fig. 3a). Taken together, we concluded there was likely to be little overt change

in spontaneous synaptic vesicle release or post-synaptic receptor density.

Nerve-evoked excitatory junction potentials (EJPs), were recorded at 1 Hz frequency

at 1 mM extracellular Ca^{2+} to evaluate synapse function. We found that EJPs at *Cln7*

neuromuscular junctions were similar to those produced by control animals (Fig. 3b). We

next applied continuous 5 Hz stimulation to the motor neurons to assess changes in the

ability to recruit from the reserve pool of neurotransmitter vesicles. In the control animals,

the evoked amplitudes followed a trend towards mild depression as vesicles in the reserve

pool become depleted. However, a much more accentuated decline of EJP amplitude was

observed in *Cln7* mutant animals under the same conditions (Fig. 3c). This suggests a

potential defect in the ability of *Cln7* neurons to activate vesicles from the reserve pool or a

defect in synaptic vesicle recycling.

The NMJs regulate body-wall muscle contraction. To identify if the

electrophysiological changes we detected led to a quantifiable change in locomotion, we

measured instantaneous speed of larval crawling. To address large variations in behavior

within a population of larvae, and to overcome the tendency of larvae to avoid open fields,

we used a custom 3D printed chamber to record the movement of multiple larvae

simultaneously in oval "racetracks" over periods of 5 min (see methods). *Cln7* mutant larvae

move significantly more slowly than isogenic control larvae (Fig. 3d).

Autophagy and BMP signaling is unaffected in Cln7 animals
204
Cln7 is required in post synaptic cells for normal development and function of the

NMJ we investigated potential pathways of action. Impairment of autophagy is known to

cause a reduction in synapse size[24], similar to that seen in the *Cln7* mutants. Furthermore, in

cultured cells Cln7 is associated predominantly with lysosomes[15], which are essential

mediators of autophagy and mutations affecting lysosomal function lead to a block in

autophagic flux. We examined accumulation of p62, a multiple domain protein that acts as a

receptor to activate autophagy. Under nutrient-rich conditions, p62 is usually degraded in

lysosomes but in starvation conditions autophagic flux is increased and some non-degraded

p62 accumulates in lysosomes (Fig. 4a; see w^{1118} control). Thus, p62 levels are often

monitored as one readout of autophagic flux. In *Cln7* larvae no p62 accumulates under

growth conditions. However, a small but not statistically significant amount of p62 did

accumulate under starvation conditions in the *Cln7* larvae, suggested a minor impairment in

autophagy or lysosomal function (Fig. 4a). This contrasted markedly with *atg18* mutants.

Here, p62 clearly accumulated during growth and did so very dramatically under starvation

conditions (Fig. 4a).

We also examined whether increasing autophagic flux could rescue the *Cln7*

phenotype by inhibiting the TOR complex activity using rapamycin or torin. TORC1 is a

central regulator of growth and autophagy and homeostasis in cells and TORC2 is required

for normal NMJ development[15,23]. Inhibition of TORC1 increases autophagic flux which, in

the *Drosophila* NMJ, leads to an overgrowth phenotype[24]. Previously, it was reported that

inhibiting TORC1 with rapamycin was unable to rescue the smaller NMJ phenotype seen in

atg18 mutants, presumably because autophagy is severely compromised in these animals

and this cannot be circumvented[24]. We reasoned that rapamycin treatment would similarly

have no effect on the *Cln7* phenotype if it was due to defective autophagy. However, both

rapamycin treatment or treatment with the TORC1 and TORC2 inhibitor, torin, caused a very

robust increase in NMJ size in both control and *Cln7* larvae indicating that autophagic flux

can be increased (Fig. 4b). Taken together with the lack of p62 accumulation, this suggests

that loss of Cln7 does not lead to an overt block in autophagy.

Given enrichment of Cln7 at the PSD we proceeded to investigate whether Cln7

functions in the classical retrograde signaling mediated by the post-synaptic BMP ligand

Glass-bottomed boat (Gbb). Loss of the type I BMP receptor Saxophone (Sax) leads to a

reduced NMJ of a size almost identical to that in *Cln7* larvae (Fig. 4c). However, a

s*ax*$^{4/5}$*;Cln7*^{84D} double mutant combination was lethal before late larval stage suggesting an

additive effect and indicating that Cln7 and Sax operate in different pathways. Conversely,

mutations in *Dad*, which encodes an inhibitory SMAD, lead to NMJ synapse overgrowth and

a profusion of small, satellite boutons (Fig. 4c). A *Dad*[J1e4];*Cln7*[84D] double mutant combination

showed a small reduction in overgrowth resulting in an intermediate phenotype. This

suggests that phosphorylation of Mad in response to a Gbb signal is intact in the *Cln7*

mutant larvae. To further confirm this, we stained the CNS of wild-type and *Cln7* larvae with

anti-pMad. The nuclei of motor neurons clearly showed pMad present at the synapse in

both genotypes (Fig. 4d) indicating that BMP signaling is unaffected in the absence of Cln7.

Dysregulated TORC signaling in Cln7 mutants

During our study, we noticed that the *Cln7* flies are considerably larger than their

isogenic control lines (Fig. 5a). *Cln7* flies are not only heavier than controls, their body parts

are also larger. We measured wing blade area from detached wings and identified a

significant increase in *Cln7* animals *vs.* controls (Fig. 5b). Increased wing size could be due to

additional cell divisions or larger cells. Additional divisions would suggest dysregulation of

cell cycle control, but larger cells could indicate a potential hyper-activation of TORC. We

measured the average area of cells within the wing and found a significant increase in cell

size in flies lacking Cln7 (Fig. 5b). Hyper-activation of TORC may also lead to excessive

storage of lipids[25]. We measured triacylgylcerides, the principal lipid stored in *Drosophila*

and identified that Cln7 flies have more than 2-fold greater lipid storage than control flies

(Fig. 5c).

Since TORC2 complex activity is necessary to restrict cell growth and loss of the

TORC2-specific component, Rictor, leads to a large expansion of the NMJ[23], we wondered

whether Cln7 may function in a pathway regulated by TORC2. We generated a

rictor[Δ42];*Cln7*[84D] double mutant and found that these animals have a significantly reduced

mass identical to that of *rictor*[Δ42] single mutants (Fig. 5d). Similarly, the *rictor*[Δ42];*Cln7*[84D]

double mutant show an expanded NMJ indistinguishable in size from the *rictor*[Δ42] single

mutant (Fig. 5e). This places *Cln7* genetically upstream of TORC2 but, as with autophagic

flux, TORC2 activity appears to be function principally in the neuron downstream of Tsc2[23].

Given the restriction of Cln7 to the post-synaptic side, it is not likely to be directly acting on

TORC2 in the neuron but could feasibly regulate TORC2 indirectly via a retrograde signal.

We looked for further supporting evidence that TORC is dysregulated by loss of Cln7.

We stained NMJs with an antibody to phosphorylated pS6, which is a marker of TORC1

activity. Phosphorylated S6 was seen clearly to localise at active zones in the pre-synaptic

boutons, suggesting localised activation of TORC1 at these release sites (Fig. 5f). This pattern

was retained in the Cln7 mutant NMJ (Fig. 5g), consistent with the restriction of Cln7 to the

post-synaptic side and acting through TORC2 rather than TORC1 in neural development.

Interestingly, some pS6 staining was seen in a vesicular pattern throughout the muscle and

this was reduced in the Cln7 mutant (Fig. 5g).

Next, we looked for more direct biochemical support that Cln7 regulates TORC

activity. Rheb is a small GTPase located primarily in the lysosomal membrane and

potentially co-located with Cln7, so we investigated whether Cln7 might be in a protein

complex with Rheb. We expressed tagged forms of Cln7 and Rheb in *Drosophila* S2 cells and

were able to co-immunoprecipitate the two proteins (Fig. 5g).

Finally, we switched to mammalian cells to confirm that the involvement of Cln7 in

the regulation of TORC signalling is conserved. We starved control and *Cln7*[−/−] mouse

embryonic fibroblasts cells to reduce basal TORC activity and then returned insulin, amino

acids or glucose to the cells. In each case, a robust of activation of TORC1 leading to

activation phosphorylation of p70 S6 kinase (Fig. 5h). In contrast, TORC1 fails to be activated

after starvation in *Cln7*[−/−] cells, consistent with a previous report[20].

Discussion

We have uncovered a role for Cln7 in the regulation of synapse growth and

development through a retrograde signaling process from the post-synaptic cell. TORC

signaling is dysregulated by loss of Cln7, potentially though disruption of a Cln7 and Rheb-

containing membrane complex. Interestingly, Cln7 mutant flies show classic indications of

excessive TORC1 activity leading to larger cell size, increased mass and excessive lipid

storage yet blocking TORC2 activity reverses these phenotypes. TORC1 activity fails to

reactivate in *Cln7*[−/−] cells after starvation indicating Cln7 function is conserved across species.

TOR has a conserved postsynaptic role to influence retrograde regulation of synaptic

activity at central and peripheral synapses in both *Drosophila* and vertebrates[13,26]. Several

activating mutations upstream of mTORC1, including loss of PTEN or Tsc1/2, result in

overgrowth of neurons and increase in density of dendritic spines in the mammalian CNS[27].

Less is understood about the role of the mTORC2 complex in the CNS although deletion of

the TORC2-specific component, Rictor, in all neurons or specifically in Purkinje cells causes

deficits in neural growth and development, including morphological effects with

consequential early-onset motor deficits[28,29].

Studies in *Drosophila* have revealed that TOR signaling is required for normal

synapse function and regulates multiple pathways controlling diverse mechanisms in both

the pre- and post-synapse partners. Reduced TORC1 activity in the pre-synapse can limit JNK

activity required for presynaptic membrane growth[11] where TORC2 also stimulates Akt

activity necessary to limit presynaptic membrane addition[23]. The TOR target Akt is also

required in the postsynaptic cell for the growth and assembly of the sub-synaptic reticulum

and targeting of glutamate receptor subunits to the postsynaptic cell surface necessary for

synapse expansion[30]. Additionally, TOR-mediated regulation of S6 kinase and presence of

the translation initiation factor eIF4E level is needed in the postsynaptic cell to produce an

unknown retrograde signal that regulates pre-synaptic neurotransmitter release[13]. It

remains to be discovered how Cln7 precisely influences TOR signaling to mediate

appropriate synapse growth and homeostasis.

The conserved role for TORC in regulating synaptic development has implications for

human CLN7 disease (also known as late-infantile neuronal ceroid lipofuscinosis). Our data

suggest dysregulated mTORC activity leading to developmental synaptic pathology may

underpin the disease. It is also significant that Cln7 protein was recently found to co-localise

with the post-synaptic marker, PSD-95, in the outer plexiform layer of the murine retina[31];

loss of visual acuity is an early symptom in CLN7 disease in humans.

Methods

***Drosophila* lines.**

Drosophila lines were obtained from the Bloomington Stock Center at Indiana University except for

an isogenic *w^{1118}* control strain from the Vienna *Drosophila* Resource Center; PG-Gal4 (*NP6293) from

the Kyoto DGRC Stock Center; Spin-Gal4 and *Dad^{j1e4}* from Dr. Sean Sweeney (University of York, UK);

and *rictor$^{\Delta 42}$* from Dr. Joseph Bateman (King's College London). Gal4 lines used with stock numbers

as of March 2018 were as follows: actin (BL3954); Elav^{c155} (BL458); Mef2 (BL27930); vGlut (BL26160);

Repo (BL7415); PG (Kyoto: 105188). The Cln7 RNAi line used was yv;TRiP.HMC03819 (BL55664). To

generate UAS-Venus-Cln7, the full-length Cln7 ORF was amplified by proofreading PCR using clone

GH22722 as a template and cloned into pENTR (Thermo). After sequence verification it was

recombined into the pTVW destination vector obtained from the *Drosophila* Genetics Resource

Center, University of Indiana and transformants generated by BestGene Inc.

Generation of Cln7 mutants.

The P-element NP0345 inserted in the 5'UTR of the *Cln7* locus (CG8596) was excised by mating with

to the Δ2-3 transposase line. Two independent alleles, 36H and 84D, were backcrossed over 6

generations to the isogenic w^{1118} control.

Visualization of the larval NMJ

The pre-synaptic membrane was visulaised with Alexa-594 goat anti-HRP (1;1000; Jackson Immuno)

except for volume measurements from confocal z-sections when a membrane-localized GFP was

used (OK371-Gal4, UAS-CD8::GFP) with rabbit anti-GFP (AbCam, 1:1000). To ensure constant culture

density and conditions between samples, cohorts of 100 1[st] instar larvae were picked from the agar

plates and transferred to vials of freshly made food and incubated for 3-4 days until wandering 3[rd]

instar stage. Larval flatmounts were immunostained using standard procedures, mounted in Prolong

Gold (Thermo) and visualized on a Zeiss LSM510 Meta, LSM780 or LSM880 Airsycan confocal

microscopes using using 40X N.A. 1.4 oil immersion, 40X N.A. 1.2 water immersion or 100x N.A. 1.42

oil immersion PlanApo lenses, respectively. 3D Renders were processed in Volocity (Perkin Elmer).

Active zones were stained with mouse anti-Bruchpilot (clone NC82, 1:25; Developmental Studies

Hybridoma Bank) and post-synaptic densities with anti-Dlg (clone 4F3, 1:20; DSHB). Rabbit anti-

pMAD (1:500) was a gift of Prof. Carl-Henrik Heldin (Ludwig Institute for Cancer Research, Uppsala).

YFP-Cln7 was visualized with rabbit anti-GFP (1:5000; AbCam ab290). Rabbit anti-pS6 (1:400) was a

kind gift of Dr. Jongkyeong Chung, Seoul National University[32].

Quantification of NMJ size

Volumes were acquired with an optical slice width of 1.0 µM and steps of 0.5 µM using a 40x N.A. 1.4

PlanApo objective and zoom = 2.0. Volocity was used to render the volumes then to measure the

volume of the pre-synaptic membrane and to count active zones. Boutons were counted manually

for the type Ib junction on muscle 4 of segments A3-A5. In each case a transmitted light image of the

corresponding muscle was obtained at 10x and the muscle surface area calculated by outlining in

ImageJ. Bouton number was then normalized to the muscle surface area.

Quantification of larval movement

Wandering 3[rd] instar stage larvae were transferred to a custom 3D-printed chamber comprising 12

oval troughs each 3 mm deep and with a track length of 75 mm. Each trough was lined with a thin

coating of 3% w/v agar. The chamber was lit from below by a LED light table and filmed from above

with a Basler GigE camera. Cohorts of 12 larvae crawling were filmed for 5 min each and their

position tracked at 2.5 fps using Ethovision software (Noldus).

Electrophysiology

Wandering 3rd instar larvae were dissected in HL3.1 supplemented with Ca^{2+} concentration at 0.5,

0.75, 1.0 or 2.0 mM depending on the paradigm[33]. Recordings were made from muscle 4 of

segments A3-A5 using standard protocols[34]. Baseline synaptic transmission was characterized by

recording mEJPs for 2 min and 16 EJPs stimulated at 1 Hz. For high frequency stimulation, 16 EPSPs

at 1 Hz were recorded followed by 5 Hz stimulation for 5 min. In all experiments a minimum of 8

larvae were used with maximum of 2 recordings per larva. In each case the starting resting potential

of the muscle was -60 to -75 mV and resistance of 5-10 mΩ. Analysis of electrophysiological

recordings was performed in Clampfit 10.0 (Molecular Devices) and data exported for further

analysis in Prism 7.0 (Graphpad).

***Drosophila* starvation assay**

First instar larvae of control *w^{1118}*, *Cln7^{84D}* and Atg18a^{KG03090}/Df(3L)Exel6112 were transferred in

cohorts of 20 from grape juice agar plates to standard food vials smeared with fresh yeast paste and

allowed to develop for a further 24 h at 25 °C. Larvae were then removed from the food and protein

samples prepared for each genotype. The remaining 10 larvae for each genotype were starved of

amino acids for 4 h at 25 °C. Lysates of total protein were prepared by manually crushing the larvae

with a mini-pestle directly into 200 μl Laemmli buffer containing 8 mM DTT and protease inhibitors.

Cell transfection and immunoprecipitation.

To generate expression plasmids, the Cln7 pENTR clone was recombined into pAMW. The full-length

Rheb ORF was amplified from clone GH10361 and cloned in to pENTR. After sequence verification it

was recombined in pAFW. Tagged proteins were immunoprecipitated with Myc-Trap_MA beads

(ChromoTek) or anti-FLAG M2 Affinity gel (Sigma) overnight at 4°C.

Serum withdrawal experiments

Mouse embryonic fibroblasts were seeded at 20% confluence then incubated in DMEM + 10 % FCS

(both Thermo) for 24 h. Cells were then washed once in EBSS (Thermo) and then starved for 18 h in

1 ml of EBSS. For re-feeding, the EBBS was replaced with complete medium (EBSS plus 10% dialysed

FBS (USB Biologicals), 1x amino acid mix (Thermo), 5.5 mM D-glucose, 4 µg/mL insulin, pH 7.95) or

partial recovery medium lacking either the amino acids, the glucose or the insulin. For the non-

starved control, basal DMEM + 10% FCS was used. Cells were lysed directly in the culture dishes after

1 h in 1X RIPA buffer (New England Biolabs) then clarified by centrifugation at 15,000 x g for 10 min

at 4°C. The protein concentration of the supernatants was determined by Pierce 660 nm protein

assay before SDS-PAGE and Western blotting.

Western blotting

Protein were separated by standard SDS-PAGE and transferred to PVDF membranes by wet transfer.

Primary antibodies were: mouse anti-Flag M2 (1:5000; Sigma); goat anti-myc (1:1000; AbCam

ab9132); rabbit anti-Ref(2)P (*Drosophila* homologue of p62, 1:500; a gift of Dr. Gabor Juhasz, Eötvös

Loránd University, Budapest); rabbit anti-p70 S6K pThr389 (1:500; Cell Signaling Technologies) and

mouse anti-actin (clone JLA-20, 1:500; DSHB). Secondary antibodies were HRP-conjugated goat anti-

mouse or anti-rabbit IgG (both 1:4000, Cell Signaling Technologies). Detection of HRP was with

SuperSignal West femto ECL reagent (Pierce) in conjunction with a Vilber Fusion FX system. p62

levels were normalized to actin from 16-bit images using ImageJ.

Statistical analysis

All statistical tests were performed within Prism 7 (GraphPad). Normality of datasets was

determined using a D'Agostino-Pearson test and datasets not conforming to Gaussian distributions

were transformed with logarithmic or reciprocal transformations. Non-parametric tests were used

where distributions remained non-Gaussian after transformation. Samples sizes and tests used are

described in the figure legends. Significance in all cases was set at $p < 0.05$.

Acknowledgements

The authors are very grateful to Dr. Stephan Storch for sharing the *Cln7$^{tm1d/tm1d}$* mouse embryonic

fibroblasts ahead of publication; to Drs. Sean Sweeney and Joe Bateman for sending *Drosophila*

lines; and to Dr. Gabor Juhasz, Prof. Carl-Henrik Heldin and Dr. Jongkyeong Chung for the anti-

Ref(2)P, anti-pMad and anti-pS6 antibodies, respectively. Research was funded by The Wellcome

Trust grant number 082004 to GT and by a UK Biotechnology and Biological Sciences Research

Council New Investigator award BB/N008472/1 to RIT. BAS was supported by a Natural Sciences and

Engineering Research Council of Canada grant number 250078. MO'H was supported by a King's

College, London Quota BBSRC PhD studentship and a KCL Peter Baker Travelling Fellowship. AM and

KJC were supported by University of Birmingham College of Medical and Dental Sciences PhD

References

Davis, G. W. Homeostatic control of neural activity: from phenomenology to molecular design. *Annu Rev Neurosci* **29**, 307-323, doi:10.1146/annurev.neuro.28.061604.135751 (2006).

Turrigiano, G. G. The self-tuning neuron: synaptic scaling of excitatory synapses. *Cell* **135**, 422-435, doi:10.1016/j.cell.2008.10.008 (2008).

Marques, G. Morphogens and synaptogenesis in Drosophila. *J Neurobiol* **64**, 417-434, doi:10.1002/neu.20165 (2005).

Deshpande, M. & Rodal, A. A. The Crossroads of Synaptic Growth Signaling, Membrane Traffic and Neurological Disease: Insights from Drosophila. *Traffic* **17**, 87-101, doi:10.1111/tra.12345 (2016).

Bayat, V., Jaiswal, M. & Bellen, H. J. The BMP signaling pathway at the Drosophila neuromuscular junction and its links to neurodegenerative diseases. *Curr Opin Neurobiol* **21**, 182-188, doi:S0959-4388(10)00134-0 [pii]10.1016/j.conb.2010.08.014 (2011).

Davis, G. W. & Goodman, C. S. Synapse-specific control of synaptic efficacy at the terminals of a single neuron. *Nature* **392**, 82-86 (1998).

Collins, C. A. & DiAntonio, A. Synaptic development: insights from Drosophila. *Current opinion in neurobiology* **17**, 35-42, doi:10.1016/j.conb.2007.01.001 (2007).

Frank, C. A. *et al.* New approaches for studying synaptic development, function, and plasticity using Drosophila as a model system. *J Neurosci* **33**, 17560-17568, doi:10.1523/JNEUROSCI.3261-13.2013 (2013).

Harris, K. P. & Littleton, J. T. Transmission, Development, and Plasticity of Synapses. *Genetics* **201**, 345-375, doi:10.1534/genetics.115.176529 (2015).

Collins, C. A., Wairkar, Y. P., Johnson, S. L. & DiAntonio, A. Highwire restrains synaptic growth by attenuating a MAP kinase signal. *Neuron* **51**, 57-69, doi:S0896-6273(06)00439-9 [pii]
10.1016/j.neuron.2006.05.026 (2006).

11	Wong, C. O. *et al.* Diminished MTORC1-Dependent JNK Activation Underlies the Neurodevelopmental Defects Associated with Lysosomal Dysfunction. *Cell reports* **12**, 2009-2020, doi:10.1016/j.celrep.2015.08.047 (2015).

12	Milton, V. J. & Sweeney, S. T. Oxidative stress in synapse development and function. *Dev Neurobiol* **72**, 100-110, doi:10.1002/dneu.20957 (2012).

13	Penney, J. *et al.* TOR is required for the retrograde regulation of synaptic homeostasis at the Drosophila neuromuscular junction. *Neuron* **74**, 166-178, doi:10.1016/j.neuron.2012.01.030 (2012).

14	Charng, W. L., Yamamoto, S. & Bellen, H. J. Shared mechanisms between Drosophila peripheral nervous system development and human neurodegenerative diseases. *Curr Opin Neurobiol* **27**, 158-164, doi:10.1016/j.conb.2014.03.001 (2014).

15	Sharifi, A. *et al.* Expression and lysosomal targeting of CLN7, a major facilitator superfamily transporter associated with variant late-infantile neuronal ceroid lipofuscinosis. *Hum Mol Genet*, doi:ddq381 [pii]
10.1093/hmg/ddq381 (2010).

16	Kousi, M. *et al.* Mutations in CLN7/MFSD8 are a common cause of variant late-infantile neuronal ceroid lipofuscinosis. *Brain* **132**, 810-819, doi:awn366 [pii]
10.1093/brain/awn366 (2009).

17   Aby, E. *et al.* Mutations in palmitoyl-protein thioesterase 1 alter exocytosis and
endocytosis at synapses in Drosophila larvae. *Fly* **7**, 267-279, doi:10.4161/fly.26630 (2013).
18   Kielar, C. *et al.* Molecular correlates of axonal and synaptic pathology in mouse
models of Batten disease. *Human molecular genetics* **18**, 4066-4080,
doi:10.1093/hmg/ddp355 (2009).
19   Grunewald, B. *et al.* Defective synaptic transmission causes disease signs in a mouse
model of juvenile neuronal ceroid lipofuscinosis. *Elife* **6**, doi:10.7554/eLife.28685 (2017).
20   Danyukova, T. *et al.* Loss of CLN7 results in depletion of soluble lysosomal proteins
and impaired mTOR reactivation. *Hum Mol Genet* **27**, 1711-1722, doi:10.1093/hmg/ddy076
(2018).
21   Mohammed, A., O'Hare, M. B., Warley, A., Tear, G. & Tuxworth, R. I. in vivo
localization of the neuronal ceroid lipofuscinosis proteins, CLN3 and CLN7, at endogenous
expression levels. *Neurobiology of disease* **103**, 123-132, doi:10.1016/j.nbd.2017.03.015
(2017).
22   Wagh, D. A. *et al.* Bruchpilot, a protein with homology to ELKS/CAST, is required for
structural integrity and function of synaptic active zones in Drosophila. *Neuron* **49**, 833-844,
doi:10.1016/j.neuron.2006.02.008 (2006).
23   Natarajan, R., Trivedi-Vyas, D. & Wairkar, Y. P. Tuberous sclerosis complex regulates
Drosophila neuromuscular junction growth via the TORC2/Akt pathway. *Human molecular*
genetics **22**, 2010-2023, doi:10.1093/hmg/ddt053 (2013).
24   Shen, W. & Ganetzky, B. Autophagy promotes synapse development in Drosophila. *J*
Cell Biol **187**, 71-79, doi:10.1083/jcb.200907109 (2009).
25   Caron, A., Richard, D. & Laplante, M. The Roles of mTOR Complexes in Lipid
Metabolism. *Annu Rev Nutr* **35**, 321-348, doi:10.1146/annurev-nutr-071714-034355 (2015).
26   Henry, F. E. *et al.* A Unique Homeostatic Signaling Pathway Links Synaptic Inactivity
to Postsynaptic mTORC1. *The Journal of neuroscience : the official journal of the Society for*
Neuroscience **38**, 2207-2225, doi:10.1523/JNEUROSCI.1843-17.2017 (2018).
27   Tavazoie, S. F., Alvarez, V. A., Ridenour, D. A., Kwiatkowski, D. J. & Sabatini, B. L.
Regulation of neuronal morphology and function by the tumor suppressors Tsc1 and Tsc2.
Nat Neurosci **8**, 1727-1734, doi:10.1038/nn1566 (2005).
28   Angliker, N., Burri, M., Zaichuk, M., Fritschy, J. M. & Ruegg, M. A. mTORC1 and
mTORC2 have largely distinct functions in Purkinje cells. *Eur J Neurosci* **42**, 2595-2612,
doi:10.1111/ejn.13051 (2015).
29   Thomanetz, V. *et al.* Ablation of the mTORC2 component rictor in brain or Purkinje
cells affects size and neuron morphology. *J Cell Biol* **201**, 293-308,
doi:10.1083/jcb.201205030 (2013).
30   Lee, H. G., Zhao, N., Campion, B. K., Nguyen, M. M. & Selleck, S. B. Akt regulates
glutamate receptor trafficking and postsynaptic membrane elaboration at the Drosophila
neuromuscular junction. *Dev Neurobiol* **73**, 723-743, doi:10.1002/dneu.22086 (2013).
31   Khan, K. N. *et al.* Specific Alleles of CLN7/MFSD8, a Protein That Localizes to
Photoreceptor Synaptic Terminals, Cause a Spectrum of Nonsyndromic Retinal Dystrophy.
Investigative ophthalmology & visual science **58**, 2906-2914, doi:10.1167/iovs.16-20608
(2017).
32   Kim, W., Jang, Y. G., Yang, J. & Chung, J. Spatial Activation of TORC1 Is Regulated by
Hedgehog and E2F1 Signaling in the Drosophila Eye. *Dev Cell* **42**, 363-375 e364,
doi:10.1016/j.devcel.2017.07.020 (2017).

33   Stewart, B. A., Atwood, H. L., Renger, J. J., Wang, J. & Wu, C. F. Improved stability of
Drosophila larval neuromuscular preparations in haemolymph-like physiological solutions. *J*
Comp Physiol A **175**, 179-191 (1994).
34   Zhang, B. & Stewart, B. Electrophysiological recording from Drosophila larval body-
wall muscles. *Cold Spring Harbor protocols* **2010**, pdb prot5487, doi:10.1101/pdb.prot5487
(2010).
537
538

Figure 1. Cln7 localizes to the basal sub-synaptic reticulum at the neuromuscular junction.

A-B): Immunofluorescence staining of the NMJ at muscle 6/7 stained with anti-HRP to

visualize the neuronal membrane (red in A and B), anti-DLG for the sub-synaptic reticulum

(SSR; blue in A) and anti-GFP to show YFP-Cln7 localization (green in B). YFP-Cln7 is

concentrates at the SSR. Scale bar = 10 μm. C-E): Higher Airyscan resolution imaging of

boutons at NMJ3, rendered into 3D and rotated using Volocity. Note the four panels are

from the same acquired image and show anti-GFP (green) with either anti-HRP (red; C&D) or

anti-DLG (blue; E&F). YFP-Cln7 localizes in a vesicular pattern around the boutons but is

restricted to the basal part of SSR (view from muscle side, D). GFP staining is largely absent

from the SSR above the boutons (view from side, E; GFP and DLG overlap below the boutons

only).

Figure 2. Loss of Cln7 leads to neurodevelopmental defects.

A). Anti-HRP staining of the late larval NMJ of w^{1118} control and $Cln7^{84D}$ flies to visualize the

neuronal membrane. Swellings of the pre-synaptic membrane (boutons) are indicated with

arrowheads. There are fewer boutons in $Cln7^{84D}$ larvae but these are occasionally swollen.

Scale bar = 10 µm. B). The volume of type Ib junction is reduced in $Cln7^{84D}$ flies but the type

Is junction is unchanged. C). Active zones number is reduced in $Cln7^{84D}$ type Ib but not in

type Is junctions. (B&C; 2-tailed t-tests, n is indicated). D). The number of boutons can be

used as a proxy for junction size. Type Ib junctions are smaller when $Cln7$ expression is

knocked down post-synaptically in muscle (Mef2-gal4) but not with pre-synaptic (Elav), pan-

glial (Repo) or perineurial glial knockdown (PG). E). Normal NMJ development is restored

only when Cln7 is re-expressed ubiquitously (actin-Gal4) in the mutant background. (D&E:

ANOVA with Dunnett's post-hoc comparisons, n is indicated).

Figure 3. Altered neural function in *Cln7* mutant flies

A-C): Electrophysiological recordings from muscle 4 in *w1118* control and *Cln7*[84D] larvae. A. The frequency (A) and amplitude (B) of miniature excitatory junction potentials (mEJP) are not significantly changed in *Cln7* larvae. mEJP frequency (left) was calculated over 120 s for each larva and amplitude (right) from 201-835 events using four different concentrations of extraceullular Ca^{2+}. Mean is indicated by a solid bar; error bars indicate SD. 2-way ANVOA with Sidak's multiple comparisons. F = 0.6086; DFn = 1; DFd = 66; p=0.4381 for both. B. Excitatory junction potentials (EJPs) are unchanged in Cln7 larvae. The mean of 16 EJPs at 1 Hz was calculated from n=13 (*w*[1118]) or 14 (*Cln7*[84D]) in 1 mM Ca^{2+}. Mean is indicated by a solid line; error bars show SD. p=0.274; Mann-Whitney. C). *Cln7* mutant synapses display an increased rate of fatigue at 5 Hz stimulation. EJP amplitude at 5 Hz was normalised to the mean amplitude at 1 Hz stimulation immediately beforehand. n=4 NMJs for *w*[1118] and =6 for *Cln7*[84D]. Trend lines fitted by linear regression are shown with solid lines with 95% confidence intervals indicated by dashes. R^2: *w*[1118] = 0.149; *Cln7*[84D] = 0.383. The slopes are significantly different: F= 4.786; DFn = 1; DFd = 96; p=0.0311, t-test. D). Crawling speed is reduced in *Cln7*[84D] larvae. Mean speed of *w*[1118] control (n=109) and *Cln7*[84D] (n=86) larvae was determined at 2.5 fps over 5 mins. **** p<0.0001; Mann-Whitney.

Figure 4. Autophagy and retrograde BMP signaling are unaffected by loss of Cln7.

A-B). The lysosomal substrate, p62, does not accumulate in control of *Cln7*[84D] larvae when feeding but does in *atg18* mutants. A. After 4 h starvation, p62 accumulates in control animals. Accumulation is slightly but not significantly increased in *Cln7*[84D] larvae. In contrast, p62 accumulates to very high levels in starved *atg18* mutants. Bars show mean p62 band intensity +/- SEM normalised to actin, n=3 for each. ** p<0.01, ANOVA. C). Inhibition of TORC1 with Rapamycin or TORC1 and 2 by Torin drives NMJ overgrowth by increasing autophagic flux. *Cln7* mutant larvae respond similarly to control *w*[1118] larvae suggesting no block in autophagy. *** p=0.01; **** p<0.001; ANOVA with Dunnett's post-hoc comparisons, n is indicated. C). Mutations affecting retrograde signaling from the muscle alter neural development. Mutations in the inhibitory SMAD, *Dad*, drive overgrowth of the NMJ; mutations in the *saxophone* type I TGFβ receptor result in underdevelopment at a similar level to *Cln7* mutants. A double *Dad;Cln7* mutant displays a small but significant moderation of the *Dad* phenotype (2-tailed t-test: t=2.181, Df=21, p=0.0407). A *sax;Cln7* double mutant is lethal early in development. Control *w*[1118] and *Cln7*[84D] data from Fig.2 are

included for comparison. D). Mad is phosphorylated after binding of the Gbb ligand to Tkv,

Sax and Wit receptors in the neuron. Characteristic pMad staining is clearly visible in

columns of motor neurons in the control and *Cln7^{84D}* CNS but is absent in a *wit* mutant CNS

(anti-pMad, top row, arrowheads). pMad staining in boutons is also present in boutons in

control and *Cln7^{84D}* NMJs but absent in a *wit* mutant (anti-pMad, bottom column; anti-HRP

marks the neuronal plasma membrane, middle column). Similarly, pMad is localised within

muscle nuclei but absent in *wit* mutants (dashed circles). Scale bar = 20 μM.

Figure 5. Loss of Cln7 leads to dysregulation of TORC

A). $Cln7^{84D}$ flies are significantly larger than isogenic w^{1118} controls. The mean mass of adult flies is shown. B). $Cln7^{84D}$ female flies have larger wings than w^{1118} controls due to increased cell size, C). $Cln7^{84D}$ female flies higher levels of triacylgylceride (TAG) storage than w^{1118} controls. TAG concentration normalized to protein concentration is shown for n=3. (A-C: 2-tailed t-tests, n is indicated for A&B). D-E). Cln7 is genetically upstream of TORC2 signaling. D). Flies mutant for the TORC2 component, *rictor,* are smaller than controls and a *rictor;;Cln7* double mutant displays a *rictor* phenotype E). $rictor^{\Delta42}$ larvae have overgrown type 1b junctions. A *rictor;;Cln7* double mutant also shows an overgrowth phenotype (D&E: ANOVA with Dunn's post-hoc comparisons, n is indicated). F). Immunofluorescence of NMJ6/7 stained for pre-synaptic membrane (anti-HRP, red) and TORC1 activity (anti-pS6, green). pRpS6 is phosphorylated by TORC1. Distinct foci of pS6 are present at the active zones in pre-synaptic boutons. These are not altered by loss of Cln7, indicating no change in pre-synaptic TORC1 activity. Scale bar = 10 μm. The vesicular anti-pS6 staining pattern in the post-synaptic muscle cells is partially reduced in $Cln7^{84D}$ mutants. H). Cln7 can be

immunoprecipitated in a complex with the TOR activating protein, Rheb. Myc-tagged Cln7

and Flag-tagged Rheb proteins were expressed in *Drosophila* S2 cells. I). Western blots for

anti-p70 S6K to indicate mTORC1 activity in control of *Cln7*[-/-] mouse embryonic fibroblasts.

Cells were starved of serum, amino acids and glucose for 18 h then insulin, glucose or amino

acids returned. Insulin is sufficient to activate mTORC1 but all three are required for full

activation. In *Cln7*[-/-] cells, mTORC1 activity is not activated. Anti-actin was used as a loading

control.