Dendritic NMDA receptors in parvalbumin neurons enable strong and stable neuronal assemblies

Jonathan Cornford[1][*], Marion S Mercier[1][*], Marco Leite[1], Vincent Magloire[1], Michael Häusser[2], Dimitri M Kullmann[1]

Affiliations:

[1]UCL Institute of Neurology, University College London,

[2]Wolfson Institute for Biomedical Research, University College London,

*These authors contributed equally

Correspondence to: jonathan.cornford@gmail.com or d.kullmann@ucl.ac.uk

Abstract: Parvalbumin-expressing (PV+) GABAergic interneurons mediate feedforward and feedback inhibition and have a key role in gamma oscillations and information processing. The importance of fast synaptic recruitment, action potential initiation and repolarization, and rapid synchronous GABA release by PV+ cells is well established. In contrast, the functional significance of PV+ cell NMDA receptors (NMDARs), which generate relatively slow postsynaptic currents, is unclear. Underlining their importance, several studies implicate PV+ cell NMDAR disruption in impaired network function and circuit pathologies. Here, we show that dendritic NMDARs underlie supralinear integration of feedback excitation from local pyramidal neurons onto mouse CA1 PV+ cells. Furthermore, by incorporating NMDARs at feedback connections onto PV+ cells in spiking networks, we show that these receptors enable cooperative recruitment of PV+ interneurons, strengthening and stabilising principal cell assemblies. Failure of this phenomenon provides a parsimonious explanation for cognitive and sensory gating deficits in pathologies with impaired PV+ NMDAR signalling.

Introduction

Interactions among cell assemblies underlie information representation and processing in the brain (Buzsáki, 2010). Inhibitory interneurons, including fast-spiking PV+ cells, which mediate feedforward and feedback inhibition and are central to gamma oscillations, have a major role in segregating excitatory principal cells into functional groups. PV+ cells have broad receptive fields inherited from multiple converging heterogeneously tuned principal neurons (Kerlin et al., 2010) and coupled with their powerful somatic inhibition of principal cells, they are positioned to mediate a "winner-takes-all" scheme in which neuronal assemblies inhibit each other (Agetsuma et al., 2018; Trouche et al., 2016).

The biophysical properties of PV+ cells that make them suited to fast inhibition of target neurons are well established (Jonas et al., 2004). These properties are critical for functions such as the enforcement of narrow temporal integration, input normalization, and sparsification of neuronal assemblies (de Almeida et al., 2009; Pouille et al., 2009; Pouille and Scanziani, 2001). However, PV+ interneurons are also equipped with NMDARs whose slow kinetics and nonlinear voltage dependence do not appear well-aligned with fast inhibition of principal cells. Although NMDARs contribute relatively less to synaptic excitation of PV+ cells than principal neurons (Geiger et al., 1997; Lamsa et al., 2007; Matta et al., 2013), several sources of evidence suggest that they are important for the normal operation of cell assemblies. In particular, genetic deletion of NMDARs in PV+ interneurons disrupts both gamma rhythms (Carlén et al., 2012) and spatial representations (Korotkova et al., 2010). Moreover, impaired NMDAR-mediated signalling in PV+ interneurons has been suggested to be a core feature of schizophrenia (Coyle, 2012; Lisman et al., 2008). Indeed, genetic manipulation of the schizophrenia risk genes encoding neuregulin and ErbB4, which amongst other functions regulate NMDARs, impairs recruitment of PV+ interneurons and recapitulates some features of the disease (delPino et al., 2013; Kotzadimitriou et al., 2018).

A recent study investigating plasticity rules of glutamatergic inputs onto CA1 PV+ interneurons reported NMDAR-dependent long-term potentiation (LTP) at feedback synapses from local pyramidal neurons but not at feedforward connections made by Schaffer collaterals (Le Roux et al., 2013), and attributed the difference to a larger

NMDAR conductance at feedback synapses. A natural question prompted by these

findings is the degree to which NMDARs contribute towards synaptic integration of

glutamatergic feedback inputs on PV+ cells. In principal neurons, NMDAR-mediated

dendritic nonlinearities enhance the computing capacity of individual cells (Gasparini,

2006; Losonczy and Magee, 2006; Poirazi and Mel, 2001; Stuart and Spruston, 2015).

Do NMDARs have an analogous function in PV+ interneurons? Furthermore, given

the importance of excitatory feedback connections on interneurons for microcircuit

motifs, how do NMDARs in PV+ interneurons affect interactions between neuronal

assemblies?

Here we combine *in vitro* optogenetic stimulation and two-photon glutamate uncaging

with modelling to assess the role of NMDARs at excitatory feedback connections

onto mouse hippocampal CA1 PV+ interneurons. We show that NMDARs at

feedback synapses mediate integrative dendritic nonlinearities in PV+ interneurons.

Importantly, this mechanism can be exploited to promote the formation of robust cell

assemblies that are stable in the face of distracting noise.

83
Results

Differential input integration at stratum oriens and stratum radiatum dendrites
of PV+ interneurons.

Experiments were performed in acute hippocampal slices from mice obtained by
crossing PV-Cre mice with Ai9 mice, and tdTomato expression was used to target
fast-spiking PV+ interneurons in CA1 stratum pyramidale. Such neurons, which
mainly comprise basket cells in addition to axo-axonic and bistratified cells (Bezaire
and Soltesz, 2013), receive excitatory feedforward inputs across the full extent of
their dendritic trees, in both strata radiatum and oriens. In contrast, feedback inputs
from axon collaterals of local pyramidal cells are confined to dendrites in the stratum
oriens (Amaral et al., 1991). In order to compare the contribution of NMDARs to
dendritic integration of feedforward and feedback excitatory postsynaptic potentials
(EPSPs), we took advantage of the anatomical restriction of feedback inputs onto
oriens dendrites, and recorded somatic responses to two-photon glutamate uncaging at
multiple sites within a 15 µm dendritic segment in either stratum oriens or stratum
radiatum (**Figure 1A**). Activation of individual uncaging locations in either stratum
evoked uncaging-evoked EPSPs (uEPSPs) that were comparable in amplitude and
kinetics to spontaneous EPSPs (**Figure 1B** and Figure 1 – Figure Supplement 1),
consistent with a high density of excitatory synapses innervating PV+ interneuron
dendrites (Gulyás et al., 1999). To quantify the degree of nonlinearity of dendritic
integration, we compared compound uEPSPs elicited by near-synchronous activation
of increasing numbers of uncaging locations to the arithmetic sum of individual
uEPSPs at the same sites (**Figure 1C**). Activation of sites on dendrites in stratum
oriens revealed supralinear uEPSP summation (peak amplitude nonlinearity: $24.0 \pm$
4.5%, mean ± SEM, n = 14; **Figure 1D,E**; unscaled responses in Figure 1 – Figure
Supplement 2). This nonlinearity was even larger when measured using the time-
integral of uEPSPs measured between 0 and 50 ms from onset (time-integral
nonlinearity: $54.0 \pm 10.1\%$; **Figure 1F**). In contrast, when glutamate was uncaged
along dendritic segments in stratum radiatum, uEPSPs summated in an approximately
linear fashion (peak amplitude nonlinearity: $3.8 \pm 5.0\%$, time-integral nonlinearity:
$6.3 \pm 7.6\%$, n = 9; oriens vs. radiatum $P = 0.0083$ and $P = 0.0028$ for peak amplitude

and time-integral comparisons respectively, unpaired *t*-tests, **Figure 1D–F**). The

difference between strata was also observed in a subset of paired recordings in which

dendrites in both strata were tested (Figure 1 – Figure Supplement 3). There was no

significant relationship between integration nonlinearity and either uncaging distance

from soma or the size of the arithmetic sum of the uEPSPs (Figure 1 – Figure

Supplement 4).

125
126

Figure 1. Differential input integration at stratum oriens and stratum
radiatum dendrites of PV+ interneurons.
(**A**) Two-photon z-projection image of a PV+ interneuron recorded via a
patch pipette in stratum pyramidale (SP) and filled with Alexa-594 (left),
with two dendritic regions of interest at higher magnification (right: top,
stratum radiatum, SR; bottom, stratum oriens, SO), showing glutamate
uncaging locations (numbered).
(**B**) Individual uEPSP responses from radiatum dendritic locations shown
in (A).
(**C**) Comparison of arithmetic sum of individual uEPSPs and recorded
uEPSPs evoked by near-synchronous uncaging at multiple locations in
stratum radiatum (blue) and oriens (red).
(**D**) Peak amplitudes of recorded uEPSPs plotted against arithmetically
summed waveforms for the two regions shown in (A). Dashed line shows
line of identity. Right: bar chart showing percentage amplitude
nonlinearity. Red: oriens, blue: radiatum.
(**E**) Summary of scaled peak amplitude comparisons for all cells (oriens
locations: n = 14, radiatum locations: n = 9). Filled circles and error bars
indicate mean ± SEM. Right: bar chart showing quantification of
amplitude nonlinearity.
(**F**) Time-integral nonlinearity plotted as for (E). **: $p < 0.01$.
148

Given that synapses in stratum oriens are innervated by both local pyramidal neurons

and Schaffer collaterals, the striking supralinear summation of uEPSPs uncovered

here may underestimate the true extent of dendritic nonlinearity at feedback

connections.

NMDAR expression and dendrite morphology underlie stratum-dependent

differences in synaptic integration.

Supralinear dendritic integration in pyramidal neurons depends on the recruitment of

voltage-dependent conductances. We therefore investigated the role of such

conductances in PV+ interneurons. In line with previous evidence for a substantial

NMDAR component at feedback inputs onto PV+ cells (Le Roux et al., 2013),

supralinear dendritic summation in stratum oriens was abolished when NMDARs

were blocked by D-AP5 (100 µM) (time-integral nonlinearity: 2.5 ± 3.0 %, vs control

without the drug $P = 0.0004$, n = 10; **Figure 2A–C**). Dendritic integration in stratum

radiatum was unchanged from control conditions (time-integral nonlinearity: 3.3 ± 2.6

%, vs control $P = 0.88$, n = 4; **Figure 2A–C**). In contrast to D-AP5, the sodium

channel blocker tetrodotoxin (TTX, 100 nM) did not significantly affect integration in

either stratum oriens or radiatum (oriens time-integral nonlinearity 40.1 ± 5.6 %, vs

control $P = 0.23$, n = 16; radiatum time-integral nonlinearity 9.4 ± 3.3 %, vs control P

= 0.71, n = 9; **Figure 2D**). The failure of TTX to affect uEPSP integration is

consistent with the view that PV+ interneuron dendrites generally do not support

regenerative events (Hu et al., 2010) (although see (Chiovini et al., 2014)). The

effects of pharmacological manipulations were consistent whether measuring time-

integrals or peak uEPSP amplitudes (Figure 2 – Figure Supplement 1). Furthermore,

uncaging distances from soma were comparable across all conditions, as were somatic

uEPSP amplitudes (Figure 2 – Figure Supplement 2). Dendrites of PV+ interneurons

mediating feedback inhibition, but not feedforward inhibition, thus exhibit NMDAR-

dependent supralinear input integration. These findings imply that clusters of coactive

synapses supplied by local pyramidal neurons cooperate via depolarization-dependent

relief of NMDARs from Mg^{2+} blockade.

Figure 2. NMDARs mediate stratum oriens dendrite synaptic
integration supralinearity.
(**A**) Two-photon z-stack of PV+ interneuron in CA1 region of
hippocampus. Red box marks glutamate uncaging location.
(**B**) Comparison of arithmetic and recorded uEPSP summation waveforms
in the presence of D-AP5. Right: peak recorded amplitude vs peak
arithmetic amplitude.

(**C**) Summary data of time-integrals plotted against arithmetic sum time-
integrals for 14 dendritic locations recorded in D-AP5 (oriens locations: n
= 10, radiatum locations: n = 4). Right: quantified synaptic integration
nonlinearity. The dashed line marks the average magnitude of oriens
nonlinearity from Figure 1F.
(**D**) Summary data for 25 dendritic locations recorded in TTX (oriens
locations: n = 16, radiatum locations: n = 9). Right: quantification of
synaptic integration nonlinearity. Filled circles and error bars indicate
mean ± SEM.
198

The results above are consistent with previous evidence of a larger NMDAR/AMPAR

conductance ratio at feedback than feedforward synapses, estimated by electrically

stimulating axons in the alveus/stratum oriens or stratum radiatum respectively, while

clamping PV+ interneurons at positive and negative potentials to separate AMPAR

and NMDAR components (Le Roux et al., 2013; Pouille and Scanziani, 2004).

However, the low input resistance of PV+ interneurons, together with different

dendritic morphologies in strata oriens and radiatum (Hu et al., 2010), potentially

confounds the comparison of excitatory postsynaptic currents (EPSCs) originating

from the two locations and recorded at positive holding potentials (Williams and

Mitchell, 2008). We therefore used an alternative experimental design to estimate the

relative contribution of AMPARs and NMDARs. Specifically, we recorded EPSCs in

a low (0.1 mM) extracellular $[Mg^{2+}]$ solution to partially unblock NMDARs while

holding PV+ interneurons at −60 mV, and used sequential addition of AMPAR and

NMDAR blockers to separate the two components of transmission (**Figure 3A**).

Pharmacological dissection of EPSCs in 0.1 mM $[Mg^{2+}]$ revealed a >2-fold greater

NMDAR/AMPAR charge ratio when stimulating in the alveus than when stimulating

in stratum radiatum (charge ratio: 3.5 ± 0.7 vs 1.3 ± 0.3, P = 0.0017, n = 10, paired *t*-

test; **Figure 3A**). The decay time constant of the NMDAR-mediated EPSCs was

similar for the two inputs (Schaffer collaterals: 154.6 ± 25.9 ms vs alveus: 148.1 ±

13.4 ms, P = 0.8, n = 10, paired *t*-test; **Figure 3B**) providing no evidence for

differences in NR2B subunit inclusion, again consistent with previous work that

showed similar effects of selective blockade of NR2B-containing receptors (Le Roux

et al., 2013) (although see (Matta et al., 2013)).

Dendrites of PV+ interneurons in stratum oriens are generally thinner and shorter than

those in stratum radiatum (Gulyás et al., 1999) suggestive of a higher effective local

input impedance. This raises the possibility that, in addition to enriched NMDAR

expression, oriens dendrites may be depolarized more effectively by glutamate

uncaging resulting in an enhanced relief of NMDARs from voltage-dependent Mg^{2+}

block (Branco et al., 2010). To investigate the relationship between synaptic

integration and dendritic geometry we used a detailed compartmental model of a CA1

PV+ interneuron (**Figure 3C**). Voltage-dependent conductance densities and

membrane properties were implemented according to previously published models

(Hu and Jonas, 2014; Nörenberg et al., 2010), and the relative densities of synaptic

AMPARs and NMDARs were initially assumed to be the same on oriens and

radiatum dendrites. Simulation parameters closely followed the uncaging

experiments, with clusters of synapses activated across the range of experimentally

measured locations. These simulations revealed supralinear summation of EPSPs

recorded at the soma that was more pronounced for stratum oriens than for stratum

radiatum dendrites (oriens vs radiatum time-integral nonlinearity: 42.5 ± 3.5 % vs

23.9 ± 2.3 %, $P = 0.0005$, **Figure 3D**), supporting a role for dendritic morphology in

mediating the difference between strata. The simulation results were very similar

whether AMPARs were assumed to show polyamine-dependent inward rectification

or to have a fixed open-channel conductance (Figure 3 – Figure Supplement 1). The ~

2-fold difference in supralinearity between strata was, however, smaller than the >8-

fold difference observed experimentally (oriens vs radiatum time-integral

nonlinearity: 54.0 ± 10.1 % vs 6.3 ± 7.6 %; **Figure 1**).

Reducing the simulated NMDAR/AMPAR conductance ratio at radiatum dendrites to

half that of the oriens dendrites, in line with results from experiments in **Figure 3A**,

improved agreement with the glutamate uncaging data (time-integral supralinearity in

simulations: 42.5 ± 3.5 % vs 8.5 ± 0.8 % for stratum oriens vs stratum radiatum;

Figure 3E). The difference in dendritic integration in oriens and radiatum dendrites

observed experimentally (**Figure 1**) may therefore be accounted for by a combination

of differential NMDAR expression (greater in stratum oriens) and dendritic

morphology (greater impedance in stratum oriens, thus facilitating depolarization and

NMDAR opening).

Figure 3 Differential NMDAR expression and dendrite morphology explain stratum-dependent synaptic integration difference.
(**A**) Schematic describing stimulation of feedforward (S1, blue) and antidromic stimulation of feedback (S2, red) axons. Middle: example paired AMPAR and NMDAR EPSC components in low [Mg^{2+}]. Right: NMDAR/AMPAR charge ratios (n = 10).
(**B**) Decay time constants of NMDAR-mediated EPSCs recorded in the same PV+ neurons in response to stimulation of feedforward (blue) and feedback (red) axons (n = 10).
(**C**) Top: reconstruction of a PV+ interneuron (axon not shown). Simulated synaptic locations are shown in gray. Bottom: example simulated uncaging experiment at the synapses marked with red circles; graph shows recorded EPSP amplitudes vs arithmetic sum of EPSP amplitudes. Inset: red solid lines, recorded summation; dashed black lines,

arithmetic summation; waveforms calculated from individual synaptic

responses.

(**D**) Scaled recorded time-integrals vs scaled arithmetic sum of time-

integrals at all locations with equal NMDAR conductance (oriens

locations: n=28, radiatum locations: n=16). Right: quantified synaptic

integration nonlinearity.

(**E**) As (D), but with reduced NMDAR/AMPAR conductance ratio at

radiatum dendrites. Oriens data replotted from (D).

NMDAR recruitment at CA1 pyramidal cell feedback connections onto PV+

interneurons.

While electrical stimulation of the alveus recruits local pyramidal cell axon

collaterals, it may also recruit other extrinsic afferents that could contribute to the

observed NMDAR currents. In order to isolate the feedback input from CA1

pyramidal cells to PV+ interneurons, and to measure the magnitude of the NMDAR

component at these synapses, we combined voltage clamp using a Cs^+-based pipette

solution with optogenetic stimulation of feedback fibers. ChR2 was selectively

expressed in CA1 pyramidal cells by injecting an adeno-associated virus (AAV)

encoding ChR2-EYFP under the control of the CaMKII promoter in CA1 of the

dorsal hippocampus. We routinely verified that expression was confined to CA1 and

did not spread to CA3 (**Figure 4A**). Wide-field illumination pulses (1ms) of 470 nm

light elicited monophasic EPSCs, in agreement with the low associative connectivity

among CA1 pyramidal neurons (Amaral et al., 1991; Deuchars and Thomson, 1996)

(**Figure 4B**). AMPAR- and NMDAR-mediated light-evoked feedback EPSCs were

recorded at -60 mV and +60 mV, respectively, and NBQX was added to isolate the

NMDAR component (**Figure 4B**). This revealed large NMDAR currents (amplitude:

459.1 ± 89.2 pA, integral: 39.1 ± 6.8 nA ms) and NMDAR/AMPAR ratios

(amplitude: 0.7 ± 0.2, integral: 2.7 ± 0.8, **Figure 4C**), confirming abundant expression

of functional NMDARs at feedback excitatory synapses on PV+ interneurons.

In order to assess whether feedback inputs from local CA1 pyramidal cells depolarize

PV+ interneurons sufficiently to recruit NMDARs, we employed the same

optogenetic strategy, and measured the contribution of NMDARs to EPSPs recorded

in current clamp and elicited by cycling the light intensity through a duty cycle

(**Figure 4D**). Perfusion of the NMDAR blocker D-AP5 significantly reduced both the

average time-integral and the ratio of integral to peak of EPSPs evoked at the

maximal light intensity ($P = 0.033$; $P = 0.014$; n=5), but not the peak EPSPs (**Figure**

4E,F). Furthermore, in line with cooperative postsynaptic voltage-dependent relief of

Mg^{2+} blockade, we observed a larger NMDAR contribution to EPSPs elicited by

stronger light pulses (**Figure 4G**). Together, these experiments confirm that synaptic

release at feedback inputs from CA1 pyramidal cells can elicit NMDAR-mediated

depolarization of PV+ interneurons.

Figure 4. NMDAR recruitment at CA1 pyramidal cell feedback connections onto PV+ interneurons.

(**A**) Schematic of viral injections into dorsal CA1 of PV-tdTomato mice (left), and confocal image of a sagittal hippocampal slice showing selective ChR2-EYFP expression in CA1 pyramidal cells (middle, inset scale: 100 µm). Right: schematic of optogenetic patch clamp experiments.

(**B**) Example traces of light-evoked feedback EPSCs in a PV+ interneuron held at -60 mV (black), +60 mV (black dashed), +60mV with application of NBQX (red) or +60mV with NBQX and D-AP5 in voltage clamp.

(**C**) Quantification of absolute NMDAR-mediated feedback EPSCs (amplitude and integral, left, n = 7) and NMDAR/AMPAR ratios (amplitude and integral, right, n = 6), measured from voltage clamp experiments as in (B). Black bars indicate mean ± SEM.

(**D**) Schematic of optogenetic stimulation protocol for current clamp experiments: light power was cycled from 20% to 100% of power for maximal response (see methods).

(**E**) EPSP integral of maximal response over time, with 20 min application of D-AP5.

(**F**) EPSP amplitude, EPSP integral and integral/amplitude ratio in the presence (red) or absence (black) of D-AP5 (n = 5, one-tailed *t*-tests; control = average of baseline and wash). Filled circles and error bars indicate mean ± SEM.

(**G**) Normalised EPSP integrals (black example traces) vs normalised EPSP integrals in the presence of D-AP5 (red example traces), for all stimulation intensities (n = 5).

NMDAR recruitment at feedback connections onto PV+ interneurons strengthens and stabilises neuronal assemblies.

NMDAR-dependent supralinear dendritic integration increases the computational capacity of principal neurons (Mel, 1992). In an extreme case cooperativity among synapses could implement an AND gate where a somatic depolarization is conditional on more than one near-simultaneous excitatory input impinging on a dendritic branch. We asked how supralinear summation of feedback excitation of PV+ interneurons could affect local circuit behaviour. Feedback recruitment of interneurons has been implicated in lateral inhibition, implementing a winner-takes-all mechanism (de Almeida et al., 2009). We therefore simulated a network of 250 excitatory point neurons (Izhikevich, 2003) reciprocally connected to a single fast-spiking inhibitory neuron (Ferguson et al., 2014, 2013) (**Figure 5A,B**). The inhibitory neuron received dual-component (AMPAR and NMDAR) synaptic conductances from the excitatory neurons (**Figure 5C**), and synapses located close to one another were allowed to interact cooperatively and engage the non-ohmic behaviour of NMDARs. The strength of interaction between individual excitatory synapses on the interneuron fell

off with distance in an abstract input space (Figure 5 – Figure Supplement 1), inspired
by the experimental evidence for clustering of homotopic inputs on dendritic
segments in principal neurons (Iacaruso et al., 2017; Wilson et al., 2016). Excitatory
neurons were assumed to be driven by a Poisson process whose rate across the
population was defined to be either clustered or dispersed in the input space. The
overall intensity was such that the network entered a sparsely firing oscillatory state
akin to a cortical gamma rhythm, with the simulated PV+ interneuron firing at around
40 Hz. Excitatory neurons driven by a compact 'hump' of excitation in input space
cooperated in recruiting NMDARs on the interneuron to a greater extent than
equivalent excitation shuffled randomly in input space (**Figure 5D**). The
disproportionate NMDAR activation by compact versus distributed excitation
recapitulates multiple co-active synapses within a small region of the dendritic tree
cooperating to relieve NMDARs from Mg^{2+} blockade. Although individual pyramidal
neurons fired sparsely, an effect of the gamma oscillation was to synchronize them so
that the local depolarization was maximized.

Recruitment of NMDAR conductances in the interneuron also maintained sparse
principal cell firing over several oscillatory cycles (Figure 5 – Figure Supplement 2).
In contrast, without NMDARs, the hump of active principal cells broadened as the
oscillation stabilized. Principal cells at the core of the hump of activity (as defined by
synaptic location on the interneuron dendritic tree) thus influence the firing of the
interneuron disproportionately as a result of the recruitment of NMDARs. We
propose, therefore, that NMDARs on PV+ interneurons contribute to maintaining a
sharp assembly representation, dependent on the spatial arrangement of active
synapses on the dendritic tree of the interneuron.

Next, we simulated two similar networks mutually inhibiting one another (Geisler et
al., 2007; Trouche et al., 2016) to understand how NMDARs in the inhibitory neurons
could affect competition among cell assemblies. When one network received a stable
and compact hump of excitation (again, with input space defined by location on the
interneuron dendritic tree) it was much more likely to 'win' than a competing network
receiving an equal amount of excitation that was dispersed (**Figure 6A**). This effect
results from the additional interneuron depolarization mediated by NMDARs which
were recruited by clustered synapses. The tendency for the network receiving a hump

of excitation to win disappeared when NMDARs were removed from the inhibitory

neurons (**Figure 6B,C**).

Finally, we explored the ability of the combined network to 'lock' onto one of two

inputs of similar strength and compactness presented to the two sub-networks (**Figure**

7). An ethologically relevant analogous task in humans is the ability to stabilize

perception of a Necker cube (**Figure 7A**). Although we make no claim as to how this

task is solved, it exemplifies a situation where two sensory or cognitive

representations compete for recruitment of a network. The net excitatory drive to each

of the sub-networks was allowed to fluctuate independently with time **(Figure 7B)**.

As a result of this stochastic variability in the input strength, and neuronal

accommodation, the combined network intermittently 'flipped' between the two

inputs. However, the frequency of flipping increased steeply when the normalized

conductance of NMDARs was decreased in the inhibitory neurons, resulting in a

flickering of the dominant assemblies (**Figure 7C**). In contrast, the frequency of

flipping was relatively unaffected by reducing the AMPAR conductance (**Figure 7D**).

We thus conclude that NMDAR-mediated cooperative interactions among clustered

synapses on an interneuron stabilize cell assemblies.

Figure 5. Network architecture and NMDAR recruitment at feedback connections.
(**A**) Schematic of network structure.
(**B**) Voltage traces of interneuron (black) and principal cells (blue, cell # at right) during network simulation. The network was driven by an asynchronous barrage of spikes, maximal in cell #125 ('clustered' input).
(**C**) Corresponding currents in interneuron.
(**D**) Left: schematic showing cell assemblies receiving clustered (top) or dispersed (bottom) external inputs, and middle: corresponding summary plots of network simulation showing external input distribution (grey), pyramidal cell firing (blue, circles), and interneuron firing (black, and vertical dashed lines). Right: average NMDAR and AMPAR charge in interneuron per principal neuron spike. (Autaptic and feedback connections from PV+ cells are omitted from the schematic for clarity.)

433
434

Figure 6. The role of NMDARs at feedback connections in cell assembly competition.
(**A**) Left: schematic showing competing cell assemblies with clustered (blue) or dispersed (orange) inputs; right: example simulation of lateral inhibition between these subnetworks with NMDARs at feedback connections to interneuron (input distribution: grey; pyramidal cell firing: blue/orange, circles; interneuron firing: blue/orange spikes and vertical dashed lines). The network receiving the clustered input out-competed the network receiving the dispersed input.
(**B**) Same as (A) but without NMDARs at feedback connections, showing, in one case the network receiving clustered input firing more than the network receiving dispersed input (top), and in the other case the network receiving dispersed input winning (bottom).
(**C**) Summary of 250 simulations showing ratio of principal cell spikes for each subnetwork with and without NMDARs at feedback inputs onto interneuron. Numbers correspond to simulations illustrated in (A) and (B).

451
452

Figure 7. The role of NMDARs at feedback connections in cell assembly stability.

(**A**) Cartoon illustrating a bistable neural representation.

(**B**) Schematic of competing subnetworks both receiving clustered inputs (left) with random fluctuations in external input strength (plotted right).

(**C**) Example simulation of network activity with NMDARs at synapses on interneurons (top), and with NMDARs scaled down to 1/8 of baseline (bottom). Input distribution: grey; pyramidal cell firing: blue/orange, circles; interneuron firing: blue/orange spikes. Although the dominant networks flipped spontaneously in both cases, the frequency of flipping was much higher with the down-scaled NMDARs.

(**D**) Plot of network flip rate vs NMDAR and AMPAR conductance. White box labelled 1: baseline NMDAR simulation parameters; black box labelled 2: NMDARs down-scaled to 12.5%.

Discussion

The present study shows that clustered excitatory synapses on stratum oriens dendrites of CA1 PV+ interneurons interact supralinearly. The high impedance of stratum oriens dendrites, which are innervated by axon collaterals of local pyramidal neurons, facilitates the cooperative recruitment of NMDARs. We place these results in the context of cell assembly competition by including nonlinear feedback integration in a spiking neural network model. NMDARs at synapses on a simulated feedback interneuron allow multiple principal cells co-innervating a subset of its input space, equivalent to a dendritic branch, to interact cooperatively in recruiting the interneuron. We show that cooperative assemblies are more able to efficiently engage interneurons in the inhibition of rival competing assembly representations. Furthermore, we report that inclusion of NMDAR conductances at feedback synapses stabilises cell assemblies, allowing the network to 'lock' on to an input. An adaptive role of NMDARs in feedback excitation of PV+ interneurons can therefore be to facilitate the emergence of strong and stable cell assemblies.

Extrapolating from the behaviour of a spiking neural network model to information processing clearly depends on a number of assumptions, not least that the principles underlying NMDAR-dependent input integration observed in CA1 PV+ interneurons apply generally throughout the brain, and that clustering of homotopic inputs on inhibitory dendritic segments obeys the same rules as in excitatory neurons (Iacaruso et al., 2017; Wilson et al., 2016). In addition, while PV+ cells predominantly show NMDAR-independent anti-Hebbian LTP (Lamsa et al., 2007), dependent on calcium-permeable AMPARs, feedback synapses also display NMDAR-dependent Hebbian plasticity rules (Le Roux et al., 2013). In this study we isolate the integrative properties of NMDARs from their role in synaptic plasticity, but it is expected that plasticity mechanisms may also contribute to the network phenomena described here. For example, NMDAR-dependent plasticity, which could be induced by cooperative feedback synaptic integration, would allow PV+ interneurons to be wired into stimulus-specific ensembles (Khan et al., 2018). Overall, therefore, NMDAR-dependent supralinear integration in the feedback inhibitory loop potentially expands the computational power of a canonical cortical motif.

Selective knockdown of NMDARs in PV+ interneurons has previously been shown to

cause a range of functional impairments, including working-memory deficits and a

reduction in the precision of hippocampal spatial representations (Korotkova et al.,

2010). The network simulations presented here provide a mechanistic explanation for

some of these results, in particular, a reduction in spatial information conveyed by

principal cell spiking (Korotkova et al., 2010). Other studies highlight a role for PV+

cells and inhibition in cell-assembly competition: in the visual cortex PV+ inhibition

increases assembly similarity (Agetsuma et al., 2018) and in the hippocampus

silencing of a dominant assembly was shown to uncover an alternative previously

inhibited assembly (Trouche et al., 2016). The present study indicates that NMDARs

may be integral to these functions.

Our network simulation results also resonate with multiple convergent findings that

implicate PV+ cell NMDAR hypofunction at the centre of schizophrenia

pathophysiology (Bygrave et al., 2016; Lisman et al., 2008; Nakazawa et al., 2012).

For instance, NMDAR blockers have been shown to recapitulate some features of

schizophrenia in healthy individuals (Krystal et al., 1994). When coupled with the

observation that NMDAR antagonists cause a net disinhibition of principal cell

activity (Homayoun and Moghaddam, 2007; Jackson et al., 2004), it has been

suggested that cortical circuits are especially vulnerable to failure of NMDAR-

mediated signalling in PV+ interneurons. Moreover, post-mortem studies have

revealed a selective loss of PV+ interneurons in people with schizophrenia (Lewis et

al., 2012) and overexpression of Neuregulin 1, a leading schizophrenia susceptibility

gene, is associated with a reduction of NMDARs on PV+ cells (Kotzadimitriou et al.,

2018). Destabilisation of dominant neuronal assemblies, as shown here to result from

impaired NMDAR signalling on PV+ interneurons, may thus explain a failure of

sensory gating (Javitt and Freedman, 2015) and evidence for reduced cognitive

control, for instance in the Necker cube test (McBain et al., 2011), in schizophrenia.

Feedback inhibition, such as that mediated by PV+ interneurons, is thought to be

critical for preventing runaway excitation. Indeed, a failure of feedback inhibitory

restraint is a major factor in the emergence of pathological states such as epileptic

seizures. Given the importance of inhibition, how then do PV+ interneurons

participate in assemblies that are composed of cells that they also regulate? One

possibility is that PV+ cells are "transient allies" (Buzsáki, 2010), only aligning with

principal cell assemblies over short time windows and at specific times. For example,

in a recent study of circuit changes in the visual cortex during learning, PV+ cells

were found to become more selective to task-relevant stimuli by increasing their

coupling to stimuli selective principal cells and becoming less influenced by the

general activity of the remaining surrounding network (Khan et al., 2018).

It is worth noting that feedback interneurons operating over longer time scales, for

example somatostatin-positive oriens-lacunosum/moleculare interneurons, or

cholecystokinin-expressing basket cells, may also be suited to mediate assembly

competition. However, once recruited they would disengage more slowly because

they integrate principal cell firing over a longer time frame. In contrast, PV+ cell

firing is thought to define the period for gamma oscillations. As a result, NMDARs on

PV+ interneurons are ideally suited to allow for stable assemblies while the stimulus

is constant, but also allow for quick switching if the nature of the stimulus changes.

Recurrent connections between PV+ interneurons and local pyramidal cell circuits are

found throughout the brain. A challenge for future studies will be to establish whether

feedback connections outside of hippocampal area CA1 also display NMDAR-

dependent supralinear feedback integration. Furthermore, experiments acutely and

specifically blocking NMDARs at feedback synapses onto PV+ cells, as opposed to

nonspecific knockdown of all PV+ cell NMDARs during development, will be

necessary in order to fully characterize the importance of PV+ cell NMDAR

processing *in vivo*. Taken together, our results expand the computational role of

NMDARs on PV+ cells, providing a parsimonious mechanism uniting a number of

hitherto unexplained observations, relating to both basic neuronal network function

and pathophysiology.

Acknowledgements

We are grateful to members of the Experimental Epilepsy Group at the UCL Institute

of Neurology, Peter Latham and Arnd Roth for advice and technical assistance. This

work was supported by the Wellcome Trust, the Medical Research Council and the

Brain Research Trust.

Competing interests

The authors declare no competing interests.

574
575 **References**
576

Agetsuma M, Hamm JP, Tao K, Fujisawa S, Yuste R. 2018. Parvalbumin-Positive
Interneurons Regulate Neuronal Ensembles in Visual Cortex. *Cereb Cortex N*
Y N 1991 **28**:1831–1845. doi:10.1093/cercor/bhx169
Amaral DG, Dolorfo C, Alvarez-Royo P. 1991. Organization of CA1 projections to
the subiculum: a PHA-L analysis in the rat. *Hippocampus* **1**:415–435.
doi:10.1002/hipo.450010410
Bartos M, Vida I, Frotscher M, Geiger JRP, Jonas P. 2001. Rapid Signaling at
Inhibitory Synapses in a Dentate Gyrus Interneuron Network. *J Neurosci*
21:2687–2698. doi:10.1523/JNEUROSCI.21-08-02687.2001
Bezaire MJ, Soltesz I. 2013. Quantitative assessment of CA1 local circuits:
knowledge base for interneuron-pyramidal cell connectivity. *Hippocampus*
23:751–785. doi:10.1002/hipo.22141
Branco T, Clark BA, Häusser M. 2010. Dendritic discrimination of temporal input
sequences in cortical neurons. *Science* **329**:1671–1675.
Buzsáki G. 2010. Neural Syntax: Cell Assemblies, Synapsembles, and Readers.
Neuron **68**:362–385. doi:10.1016/j.neuron.2010.09.023
Bygrave AM, Masiulis S, Nicholson E, Berkemann M, Barkus C, Sprengel R,
Harrison PJ, Kullmann DM, Bannerman DM, Kätzel D. 2016. Knockout of
NMDA-receptors from parvalbumin interneurons sensitizes to schizophrenia-
related deficits induced by MK-801. *Transl Psychiatry* **6**:e778.
doi:10.1038/tp.2016.44
Carlén M, Meletis K, Siegle JH, Cardin JA, Futai K, Vierling-Claassen D, Rühlmann
C, Jones SR, Deisseroth K, Sheng M, Moore CI, Tsai L-H. 2012. A critical
role for NMDA receptors in parvalbumin interneurons for gamma rhythm
induction and behavior. *Mol Psychiatry* **17**:537–548. doi:10.1038/mp.2011.31
Carnevale NT, Hines ML. 2009. The NEURON Book. Cambridge University Press.
Chiovini B, Turi G, Katona G, Kaszás A, Pálfi D, Maák P, Szalay G, Szabó M, Szabó
G, Szadai Z, Káli S, Rózsa B. 2014. Dendritic Spikes Induce Ripples in
Parvalbumin Interneurons during Hippocampal Sharp Waves. *Neuron* **82**:908–
924. doi:10.1016/j.neuron.2014.04.004
Coyle JT. 2012. NMDA receptor and schizophrenia: A brief history. *Schizophr Bull*
38:920–926. doi:10.1093/schbul/sbs076
Cuntz H, Forstner F, Borst A, Häusser M. 2010. One rule to grow them all: a general
theory of neuronal branching and its practical application. *PLoS Comput Biol*
6. doi:10.1371/journal.pcbi.1000877
de Almeida L, Idiart M, Lisman JE. 2009. A second function of gamma frequency
oscillations: an E%-max winner-take-all mechanism selects which cells fire. *J*
Neurosci Off J Soc Neurosci **29**:7497–7503. doi:10.1523/JNEUROSCI.6044-
08.2009
delPino I, García-Frigola C, Dehorter N, Brotons-Mas JR, Alvarez-Salvado E,
MartínezdeLagrán M, Ciceri G, Gabaldón MV, Moratal D, Dierssen M,
Canals S, Marín O, Rico B. 2013. Erbb4 Deletion from Fast-Spiking
Interneurons Causes Schizophrenia-like Phenotypes. *Neuron* **79**:1152–1168.
doi:10.1016/j.neuron.2013.07.010

Deuchars J, Thomson AM. 1996. CA1 pyramid-pyramid connections in rat
hippocampus in vitro: Dual intracellular recordings with biocytin filling.
Neuroscience **74**:1009–1018. doi:10.1016/S0306-4522(96)00251-5
Ferguson KA, Huh CYL, Amilhon B, Williams S, Skinner FK. 2014. Simple,
biologically-constrained CA1 pyramidal cell models using an intact, whole
hippocampus context. *F1000Research* **3**:104.
doi:10.12688/f1000research.3894.1
Ferguson KA, Huh CYL, Amilhon B, Williams S, Skinner FK. 2013. Experimentally
constrained CA1 fast-firing parvalbumin-positive interneuron network models
exhibit sharp transitions into coherent high frequency rhythms. *Front Comput*
Neurosci **7**:144. doi:10.3389/fncom.2013.00144
Gasparini S. 2006. State-Dependent Dendritic Computation in Hippocampal CA1
Pyramidal Neurons. *J Neurosci* **26**:2088–2100.
doi:10.1523/JNEUROSCI.4428-05.2006
Geiger JR, Lübke J, Roth A, Frotscher M, Jonas P. 1997. Submillisecond AMPA
receptor-mediated signaling at a principal neuron-interneuron synapse. *Neuron*
18:1009–1023.
Geisler C, Robbe D, Zugaro M, Sirota A, Buzsáki G. 2007. Hippocampal place cell
assemblies are speed-controlled oscillators. *Proc Natl Acad Sci* **104**:8149–
8154.
Gulyás AI, Megías M, Emri Z, Freund TF. 1999. Total number and ratio of excitatory
and inhibitory synapses converging onto single interneurons of different types
in the CA1 area of the rat hippocampus. *J Neurosci Off J Soc Neurosci*
19:10082–10097.
Hines ML, Carnevale NT. 1997. The NEURON simulation environment. *Neural*
Comput **9**:1179–1209. doi:10.1162/neco.1997.9.6.1179
Homayoun H, Moghaddam B. 2007. NMDA receptor hypofunction produces opposite
effects on prefrontal cortex interneurons and pyramidal neurons. *J Neurosci*
Off J Soc Neurosci **27**:11496–11500. doi:10.1523/JNEUROSCI.2213-07.2007
Hu H, Jonas P. 2014. A supercritical density of Na(+) channels ensures fast signaling
in GABAergic interneuron axons. *Nat Neurosci* **17**:686–693.
Hu H, Martina M, Jonas P. 2010. Dendritic mechanisms underlying rapid synaptic
activation of fast-spiking hippocampal interneurons. *Science* **327**:52–58.
doi:10.1126/science.1177876
Iacaruso MF, Gasler IT, Hofer SB. 2017. Synaptic organization of visual space in
primary visual cortex. *Nature* **547**:449–452. doi:10.1038/nature23019
Izhikevich EM. 2003. Simple model of spiking neurons. *IEEE Trans Neural Netw*
14:1569–1572. doi:10.1109/TNN.2003.820440
Jackson ME, Homayoun H, Moghaddam B. 2004. NMDA receptor hypofunction
produces concomitant firing rate potentiation and burst activity reduction in
the prefrontal cortex. *Proc Natl Acad Sci U S A* **101**:8467–8472.
doi:10.1073/pnas.0308455101
Javitt DC, Freedman R. 2015. Sensory processing dysfunction in the personal
experience and neuronal machinery of schizophrenia. *Am J Psychiatry*
172:17–31. doi:10.1176/appi.ajp.2014.13121691
Jonas P, Bischofberger J, Fricker D, Miles R. 2004. Interneuron Diversity series: Fast
in, fast out – temporal and spatial signal processing in hippocampal
interneurons. *Trends Neurosci* **27**:30–40.

Kerlin AM, Andermann ML, Berezovskii VK, Reid RC. 2010. Broadly tuned
response properties of diverse inhibitory neuron subtypes in mouse visual
cortex. *Neuron* **67**:858–871. doi:10.1016/j.neuron.2010.08.002

Khan AG, Poort J, Chadwick A, Blot A, Sahani M, Mrsic-flogel TD, Hofer SB. 2018.
selectivity and interactions of GABAergic interneuron classes in visual cortex.
Nat Neurosci **21**. doi:10.1038/s41593-018-0143-z

Korotkova T, Fuchs EC, Ponomarenko A, von Engelhardt J, Monyer H. 2010. NMDA
receptor ablation on parvalbumin-positive interneurons impairs hippocampal
synchrony, spatial representations, and working memory. *Neuron* **68**:557–569.
doi:10.1016/j.neuron.2010.09.017

Kotzadimitriou D, Nissen W, Paizs M, Newton K, Harrison PJ, Paulsen O, Lamsa K.
2018. Neuregulin 1 Type-I over-Expression Is Associated with Reduced
NMDA Receptor-Mediated Synaptic Signaling in Hippocampal Interneurons
Expressing PV or CCK. *eNeuro* **5**. doi:10.1523/ENEURO.0418-17.2018

Krystal J, Karper L, Seibyl J, Freeman G, Delaney R, Bremner D, Heninger G,
Bowers M, Charney D. 1994. Subanesthetic effects of the NMDA antagonist,
ketamine, in humans: psychotomimetic, perceptual, cognitive, and
neuroendocrine effects. *Arch Gen Psychiatry* **51**:199–214.
doi:10.1017/CBO9781107415324.004

Lamsa KP, Heeroma JH, Somogyi P, Rusakov DA, Kullmann DM. 2007. Anti-
hebbian long-term potentiation in the hippocampal feedback inhibitory circuit.
Science **315**:1262–1266. doi:10.1126/science.1137450

Le Roux N, Cabezas C, Böhm UL, Poncer JC. 2013. Input-specific learning rules at
excitatory synapses onto hippocampal parvalbumin-expressing interneurons. *J
Physiol* **591**:1809–22. doi:10.1113/jphysiol.2012.245852

Lewis DA, Curley AA, Glausier JR, Volk DW. 2012. Cortical parvalbumin
interneurons and cognitive dysfunction in schizophrenia. *Trends Neurosci*
35:57–67. doi:10.1016/j.tins.2011.10.004

Lisman JE, Coyle JT, Green RW, Javitt DC, Benes FM, Heckers S, Grace AA. 2008.
Circuit-based framework for understanding neurotransmitter and risk gene
interactions in schizophrenia. *Trends Neurosci* **31**:234–242.
doi:10.1016/j.tins.2008.02.005

Losonczy A, Magee JC. 2006. Integrative properties of radial oblique dendrites in
hippocampal CA1 pyramidal neurons. *Neuron* **50**:291–307.
doi:10.1016/j.neuron.2006.03.016

Matta JA, Pelkey KA, Craig MT, Chittajallu R, Jeffries BW, McBain CJ. 2013.
Developmental origin dictates interneuron AMPA and NMDA receptor
subunit composition and plasticity. *Nat Neurosci* **16**:1032–1041.
doi:10.1038/nn.3459

McBain R, Norton DJ, Kim J, Chen Y. 2011. Reduced cognitive control of a visually
bistable image in schizophrenia. *J Int Neuropsychol Soc* **17**:551–556.
doi:10.1017/S1355617711000245

Mel BW. 1992. NMDA-Based Pattern Discrimination in a Modeled Cortical Neuron.
Neural Comput **4**:502–517. doi:10.1162/neco.1992.4.4.502

Nakazawa K, Zsiros V, Jiang Z, Nakao K, Kolata S, Zhang S, Belforte JE. 2012.
GABAergic interneuron origin of schizophrenia pathophysiology.
Neuropharmacology **62**:1574–1583. doi:10.1016/j.neuropharm.2011.01.022

Nörenberg A, Hu H, Vida I, Bartos M, Jonas P. 2010. Distinct nonuniform cable
properties optimize rapid and efficient activation of fast-spiking GABAergic

interneurons. *Proc Natl Acad Sci U S A* **107**:894–899. doi:10.1073/pnas.0910716107

Poirazi P, Mel BW. 2001. Impact of active dendrites and structural plasticity on the memory capacity of neural tissue. *Neuron* **29**:779–796. doi:10.1016/S0896-6273(01)00252-5

Pouille F, Marin-Burgin A, Adesnik H, Atallah B V, Scanziani M. 2009. Input normalization by global feedforward inhibition expands cortical dynamic range. *Nat Neurosci* **12**:1577–1585. doi:10.1038/nn.2441

Pouille F, Scanziani M. 2004. Routing of spike series by dynamic circuits in the hippocampus. *Nature* **429**:717–723. doi:10.1038/nature02615

Pouille F, Scanziani M. 2001. Enforcement of temporal fidelity in pyramidal cells by somatic feed-forward inhibition. *Science* **293**:1159–63. doi:10.1126/science.1060342

Roth A, van Rossum MCW. 2009. Modeling Synapses. *Comput Model Methods Neurosci* **06**:139–160. doi:10.7551/mitpress/9780262013277.003.0007

Stuart GJ, Spruston N. 2015. Dendritic integration: 60 years of progress. *Nat Neurosci* **18**:1713–1721. doi:10.1038/nn.4157

Trouche S, Perestenko P V, van de Ven GM, Bratley CT, McNamara CG, Campo-Urriza N, Black SL, Reijmers LG, Dupret D. 2016. Recoding a cocaine-place memory engram to a neutral engram in the hippocampus. *Nat Neurosci* **19**:564–567. doi:10.1038/nn.4250

Wang XJ, Buzsáki G. 1996. Gamma oscillation by synaptic inhibition in a hippocampal interneuronal network model. *J Neurosci* **16**:6402–6413.

Williams SR, Mitchell SJ. 2008. Direct measurement of somatic voltage clamp errors in central neurons. *Nat Neurosci* **11**:790–798.

Wilson DE, Whitney DE, Scholl B, Fitzpatrick D. 2016. Orientation selectivity and the functional clustering of synaptic inputs in primary visual cortex. *Nat Neurosci* **19**:1003–1009. doi:10.1038/nn.4323

Materials and Methods

<u>Animals</u>

Hippocampal slices were obtained from postnatal day 14 – 24 male and female mice, or from 2-3 month old male and female mice (optogenetic experiments), expressing tdTomato in PV+ interneurons. Experimental mice were obtained by crossing homozygous mice expressing Cre under the PV promoter (Jackson Labs: B6;129P2-Pvalb$^{tm1(cre)Arbr}$) with homozygous Ai9 Cre reporter mice (Jackson Labs: B6.Cg-$Gt(ROSA)26Sor^{tm9(CAG-tdTomato)Hze}$). Animals were group-housed under a non-reversed 12 h light/dark cycle, and allowed access to food and water *ad libitum*. All procedures were carried out in accordance with the UK Animals (Scientific Procedures) Act, 1986.

<u>Surgery for viral injections</u>

Mice (minimum age: 6 weeks) were anaesthetized with isoflurane and viruses were stereotaxically injected into the dorsal CA1 region of both hippocampi using a Hamilton syringe. The injection coordinates were 2.15 mm caudal and 1.4 mm lateral of Bregma, and 1.2 and 1.0 mm deep from the pia. 50 nl of virus was injected at each site at a rate of 100 nl/min, and the needle was left in place for 5 minutes following injections before withdrawal. Slices were prepared for experiments after a minimum of three weeks post-surgery.

<u>Slice preparation and electrophysiology</u>

Acute sagittal brain slices (300 µm) were prepared using a Vibratome (Leica VT1200 S). Slices were cut in an ice-cold artificial cerebrospinal fluid (ACSF) solution, containing (in mM): NaCl (119), KCl (2.5), NaH$_2$PO$_4$ (1.25), NaHCO$_3$ (25), glucose (20), CaCl$_2$ (1.5), MgSO$_4$ (1.3), and saturated with 95% O$_2$, 5% CO$_2$. Slices were allowed to recover at 32°C for 15 min after slicing, before subsequent storage in ACSF at room temperature. Older mice (>1 month) were transcardially perfused with ice-cold sucrose-based ACSF solution, containing (in mM): sucrose (75), NaCl (87), KCl (2.5), NaH$_2$PO$_4$ (1.25), NaHCO$_3$ (25), glucose (25), CaCl$_2$ (0.5), MgCl$_2$ (7), and saturated with 95% O$_2$, 5% CO$_2$. Slices were cut in the same solution, and left to recover at 32°C for 15 min, before being

transferred to normal ACSF (same as above but with 2.5 mM $CaCl_2$) for storage at

room temperature. All experiments were carried out in ACSF maintained at 30° –

32°C and perfused at 2–3 ml/min. For dissection of AMPAR and NMDAR

components at feedforward and feedback synapses (**Figure 3A,B**) a modified

ACSF containing 0.1 mM $MgSO_4$ was used to partially relieve Mg^{2+} blockade of

NMDARs at rest. NMDARs and AMPARs were sequentially blocked with D-AP5

(100 μM) and 2,3-dihydroxy-6-nitro-7-sulfamoyl-benzo[f]quinoxaline-2,3-dione

(NBQX, 10 μM), respectively. Picrotoxin (100 μM) and CGP 55845 (1 μM) were

included throughout these experiments, as well as the HCN channel blocker ZD

7288 (30 μM), which was included in order to hyperpolarize pyramidal cells and

decrease network excitability. Picrotoxin (100 μM) and CGP 55845 (1 μM) were

also included throughout the optogenetic experiments (**Figure 4**). For blockade

of NMDARs or Na^+ channels during uncaging experiments (**Figure 2**) D-AP5 (100

μM) or TTX (0.1 μM) were used, respectively.

Fluorescence-guided somatic whole-cell recordings were obtained from PV+

interneurons and pyramidal cells using a Multiclamp 700B amplifier (Molecular

Devices), filtered at 5 kHz, and digitized at 20 kHz (National Instruments PCI-

6221), with custom software written in LabVIEW. Patch pipettes of 3 – 4 MΩ

resistance were filled with KGluconate- or CsGluconate-based internal solution

for current clamp or voltage clamp experiments, respectively. These solutions

contained (in mM): KGluconate (140), KOH-HEPES (10), EGTA (0.2), NaCl (8),

Mg-ATP (2), Na-GTP (0.3) or CsGluconate (125), HCsO-HEPES (10), EGTA (0.2),

NaCl (8), Mg-ATP (4), Na-GTP (0.33), Na-Phosphocreatine (10), TEA-Cl (5)

QX314 (5). Cells were held at -60 mV or +60 mV during voltage-clamp

experiments (**Figure 3A,B**, and **Figure 4B,C**). All other experiments were

performed in current-clamp mode, and current was continuously injected to

maintain cell membrane between -65 and -70 mV. The series resistance during

voltage clamp recordings was <15 MΩ and during current clamp recordings was

< 25 MΩ. For field stimulation experiments, concentric bipolar stimulating

electrodes (FHC) coupled to constant current stimulators (Digitimer) were

placed in the alveus and stratum radiatum in order to evoke responses from

feedback and feedforward inputs, respectively (Pouille and Scanziani, 2004).

Stimuli were delivered to each pathway at 0.05 Hz, and alternated between the

pathways. Optogenetic responses were elicited using 470 nm light pulses (1 ms,

1-15 mW) at 0.2 Hz, generated by an LED light-source (ThorLabs) and delivered

through a 40X objective lens (Olympus). The light power necessary to elicit

maximal and minimal responses was identified for each cell, and the difference

in power was divided by 5 to define the 5 stimulation strengths (20%, 40%,

60%, 80% and 100%) used in each cycle. Maximal responses were identified as

the maximum response elicited without generating an action potential (typically

≤ 15 mV), and minimal responses were the smallest response visible.

<u>Two-photon imaging and uncaging experiments</u>

Slices were submerged in a perfusion chamber on an upright microscope (FV1000

BX61, Olympus). Simultaneous two-photon imaging and uncaging of MNI-caged

glutamate was performed with two Ti-sapphire lasers tuned to 810 nm and 720 nm, for

imaging and uncaging respectively (Mai-Tai, Spectra Physics; Chameleon, Coherent).

MNI-caged-glutamate-TFA (3 mM; Femtonics) dissolved in the recording ACSF

solution was perfused in a closed system.

Uncaging locations (range: 8 – 12, mean: 8.4 vs 8.5 sites in radiatum and oriens

respectively) were selected either side of a dendritic region of interest, separated

by 2 – 3 μm and within 1 μm of the dendrite. Uncaging-evoked EPSPs (uEPSPs)

were evoked using 0.5 ms-long pulses of 720 nm laser-light. To account for

differing depths of dendritic segments, uncaging-laser intensity was adjusted

using a Pockels Cell (Conoptics). uEPSPs were first evoked by sequential

stimulation of individual uncaging spots with an inter-stimulus interval of 200

ms. Uncaging locations (order chosen at random) were then stimulated at

intervals of 1 ms, with 10 s delays between trials. Beginning with a single

location, the number of uncaging locations was increased on successive trials

until all locations were activated. The entire sequence was repeated between 2

and 5 times, without changing the order of uncaging locations, and the responses

to each combination of uncaging locations were averaged. Arithmetic compound

uEPSPs were constructed from individual spot recordings offline, including a 1

ms waveform shift to match the experimental protocol, and compared to

recorded uEPSPs. Uncaging times and locations were controlled by scanning

software (Fluoview 1000V) and a pulse generator (Berkeley Nucleonics) coupled

to the Pockels cell. Experiments were discontinued and excluded from analysis if

photo-damage to PV+ cells was observed, or if physical drift occurred.

Quantification and statistical analysis

Data analysis was performed using custom code written in Python. The nonlinearity of

responses recorded from uncaging at each dendritic location was quantified using the

following equation:

$$nonlinearity\ \% = \sum_{i=2}^{n} \frac{\frac{Mi}{Ai} - 1}{n - 1} \cdot 100\% \tag{1}$$

where Mi is the amplitude of the ith measured uEPSP (composed of i individual

uncaging spots), Ai is the amplitude of the ith constructed arithmetic summed

uEPSP, and n is the total number of uncaging locations. For uEPSP integral

analysis a Savitzky-Golay filter was applied to traces.

For analysis of NMDAR/AMPAR ratios in **Figure 3A**, the AMPAR-mediated

response was calculated by subtracting the NMDAR-mediated response

(recorded in NBQX) from the baseline EPSC. NMDAR and AMPAR charge were

then calculated by integrating the first 500 ms or 20 ms, respectively, of these

isolated traces. NMDAR/AMPAR ratios in **Figure 4C** were calculated as the ratio

between the NMDAR-mediated response recorded at +60 mV in the presence of

NBQX (with the +60 mV response in the presence of D-AP5 subtracted), and the

AMPAR-mediated response recorded at -60 mV.

Statistical significance was assessed using Student's paired or unpaired t-tests.

Data are presented as mean ± SEM, unless stated otherwise. Sample sizes were

estimated to obtain 80% power to detect effects at p<0.05. n values are for cells.

Multi-compartmental modelling

Multi-compartmental modelling was performed with the NEURON 7.5 simulation

environment (Hines and Carnevale, 1997). The soma and dendrites of a PV+

interneuron were reconstructed using the TREES toolbox in MATLAB (Cuntz et

al., 2010). The axon was not included in the reconstruction. As PV+ interneuron

dendrites are generally smooth, addition of spines or correction of synaptic

responses for spines was deemed unnecessary. The number of segments per

section was constrained to odd numbers and set according to the d-lambda rule

(Carnevale and Hines, 2009) to have a length no more than 10% of the

alternating current length constant at 1 kHz. The model contained 500 segments

in total with a maximal segment length of 8.7 μm.

The biophysical parameters were based on previously published models of

dentate gyrus PV+ basket cells (Hu and Jonas, 2014; Nörenberg et al., 2010). The

specific membrane capacitance (C_m) and intracellular resistance (R_i) were

assumed to be spatially uniform (for values see Table 2.1). In contrast, the

specific membrane resistance (R_m) was assumed to vary as a step function with

distance from the soma. R_m at distal dendrites was 10 times larger than at

proximal dendrites, and R_m was chosen so as to make the model cell's input

resistance 78 MΩ, the mean experimentally recorded input resistance (78.6 ± 5.2

MΩ). The border between proximal and distal dendrites was defined to be 120

μm from the soma.

Table 1: NEURON model parameters

Parameter	Proximal	Distal	Units
C_m	0.9	0.9	$\mu F\ cm^{-2}$
R_{axial}	170	170	$\Omega\ cm$
R_m	5.55	55.5	$k\Omega$
e_{leak}	-65	-	mV
e_{gK}	-90	-	mV
e_{gNa}	55	-	mV
v shift	-12	-12	mV
g_k dend	300	300	$pS\ \mu m^{-2}$
g_{Na} dend	200	100	$pS\ \mu m^{-2}$
g_{Na} soma	2000	-	$pS\ \mu m^{-2}$

Wang and Buzsaki (WB) Na⁺ and K⁺ channels were inserted in the model neuron

to confer a fast-spiking action potential phenotype (Wang and Buzsáki, 1996).

However, in order to produce a realistic firing frequency - current injection

relationship a hyperpolarizing voltage shift was included in the WB

implementation. The depolarized threshold of the WB mechanism has been

discussed previously (Ferguson et al., 2013).

Subthreshold synaptic integration curves were produced by first finding all sites

on the dendritic tree that were located between 40 and 190 μm from the soma.

Simulations then closely followed the experimental protocol detailed above. At

each dendritic site, 15 synapses were placed within a distance of 30 μm. Each

synapse was activated individually and the arithmetic sum calculated from the

somatic membrane potential. Synapses were then activated in increasing

numbers, with an interval of 1 ms between activations, and the integral and

amplitude of these measured responses compared to the calculated arithmetic

responses. Quantification of dendritic nonlinearity was identical to that applied

to experimental data.

<u>Network modelling</u>

Single cell modelling

For network simulations, PV+ interneurons and CA1 pyramidal cells were represented

by two-dimensional Izhikevich model neurons (Izhikevich, 2003). Izhikevich models

for these neurons have previously been parameterized from experimental data

(Ferguson et al., 2014, 2013). In line with this previous work, the models were slightly

modified to reproduce the narrow PV+ interneuron spike width, and had the following

form:

$$C_m \frac{dv}{dt} = k(v - v_r)(v - v_t) - u + I_{applied}$$

$$\frac{du}{dt} = a[b(v - v_r) - u] \tag{2}$$

$$if\ v \geq v_{peak}, \text{set } v = c,\ u = u + d$$

$$\text{Where } k = k_{low} \text{ if } v \leq v_t;\ k = k_{high} \text{ if } v > v_t$$

The variable v represents the membrane potential, and u represents a slow

"refractory" current, that, aside from subthreshold effects governed by b, is

increased by d when the neuron fires, and decays at a rate determined by a. The

parameter C_m represents the membrane capacitance; k is a scalar; v_r is the

resting membrane potential; v_t is the instantaneous spiking threshold potential;

v_{peak} is the peak action potential voltage; $I_{applied}$ is the applied current,

comprised of the sum of all synaptic inputs to the cell; a is the recovery inverse

time constant of the refractory current, u; b is the sensitivity of u to subthreshold

voltage fluctuations; c is the voltage reset value; and d is the amount of current

generated by the after-spike behaviour. The values for all parameters of the

network simulations are presented in Table 2.2

Synaptic modelling

Synaptic connections between neurons were modelled as bi-exponential,

conductance-based synapses (Roth and van Rossum, 2009), which can be

written in the following differential form:

$$g^{chan}(v^{syn}) = \frac{G^{chan}(v^{syn})}{\tau_1^{chan} - \tau_2^{chan}}(g_1^{chan} - g_2^{chan})$$

$$\frac{dg_1^{chan}}{dt} = \left(\delta^{chan}(t) - \frac{1}{\tau_1^{chan}}g_1^{chan}\right)$$

$$\frac{dg_2^{chan}}{dt} = \left(\delta^{chan}(t) - \frac{1}{\tau_2^{chan}}g_2^{chan}\right) \tag{3}$$

$$G^{NMDA}(v^{syn}) = \left(\frac{1}{2}tanh\left[\frac{v_i^{syn} + 50mV}{10mV}\right] + \frac{1}{2}\right)$$

$$G^{AMPA}(v^{syn}) = G^{ext}(v^{syn}) = G^{GABA}(v^{syn}) = 1$$

Where g^{chan} is the total synaptic conductance of a given channel family

(composed of a rise term g_1^{chan} and a decay term g_2^{chan}), $G^{chan}(v^{syn})$ is an

eventual instantaneous voltage-gating term dependent on a local synaptic

voltage v^{syn} (only relevant for NMDA channels; detailed in the next section),

τ_1^{chan} is the rising exponential time constant, τ_2^{chan} is the decay exponential time

constant, the variable t is time, and $\delta^{chan}(t)$ represents the input spike train of

Dirac deltas.

NMDA receptor modelling

NMDARs, present at the feedback connections from principal cells onto the

interneuron, and the cooperative relief of NMDAR Mg²⁺ block by co-active

synaptic inputs, were modelled in an abstract manner. We assumed that all co-

active inputs from the population of n_{pyr} principal cells had a degree of

cooperation, or functional clustering (Wilson et al., 2016), which was weighted

by a distance matrix, $D^{n_{pyr} \times n_{pyr}}$. D was defined as a Toeplitz matrix, with the i_{th}

element of the row vector $D_{\frac{n_{pyr}}{2}}$ equal to:

$$D_{\frac{n_{pyr}}{2},i} = \frac{1}{\sqrt{2\pi \, \sigma_D^2}} e^{-\frac{(i-n_{pyr}/2)^2}{2 \, \sigma_D^2}} \qquad (4)$$

967 Where σ_D^2 controlled the specificity of local cooperation. We then modelled the

968 time evolution of the voltage of the local synaptic membrane patch, v_i^{syn}, with

969 the following equation:

$$C_{syn} \frac{dv_i^{syn}}{dt} = k^{syn} I_i^{syn} + g^{leak}\left(e^{leak} - v_i^{syn}\right) \qquad (5)$$

$$I_i^{syn} = \sum_{j=1}^{n_{pyr}} D_{i,j}\left(k_{pv+}^{AMPA} g_j^{AMPA} + k_{pv+}^{NMDA} g_j^{NMDA}(v_j^{syn})\right)\left(e^{glu} - v_i^{syn}\right)$$

971 Where C_{syn} is the local patch membrane capacitance, k^{syn} is a gain applied to the

972 input current I_i^{syn}, k_{pv+}^{chan} are gain constants defining each channel family

973 synaptic strength onto the interneuron, g^{leak} is a leak conductance with reversal

974 potential e^{leak}.

975 Finally, the total current, $I_{applied}$, an interneuron receives is given by:

$$I_{applied} = \left(\sum_{i=1}^{n_{pyr}} k_{pv+}^{AMPA} \, g_i^{AMPA} + k_{pv+}^{NMDA} g_i^{NMDA}(v_i^{syn}) + k_{pv+}^{ext} \, g_i^{ext}\right)(e^{glu} - v)$$

$$+ \, k_{pv+}^{GABA}\left(\sum_{i=1}^{n_{pv+}} g_i^{GABA}\right)(e^{GABA} - v) \qquad (6)$$

977 Accordingly, for the *ith* pyramidal cell the input current is more simply given by:

$$I_{Applied}^i = k_{pyr}^{ext} \, g_i^{ext}(e^{glu} - v) + k_{pyr}^{GABA}\left(\sum_{i=1}^{n_{pv+}} g_i^{GABA}\right)(e^{GABA} - v) \qquad (7)$$

External inputs

The external input to the i_{th} pyramidal neuron was modelled as a non-

homogeneous Poisson process, in which the instantaneous rate of incoming

spikes was given by an Ornstein–Uhlenbeck process with mean μ_{OU}, volatility

σ_{OU} and time constant τ_{OU}, multiplied by a Gaussian gain function representing

the neuron position in the receptive field. The input of the PV+ interneurons was

the mean of all inputs to its afferent pyramidal cells. The peak mean μ_{OU} was

5000 spikes per second, τ_{OU} was 50 ms and σ_{OU} was $\frac{1}{6}\mu_{OU}\sqrt{2\,\tau_{OU}}$ for the

simulations in **Figure 7** and zero everywhere else (i.e. the Poisson rates were

constant).

Table 2: model neuron parameters for network modelling (Bartos et al., 2001; Geiger et al., 1997; Roth and van Rossum, 2009)

Parameter	FS PV+	Pyramidal	Units
C_m	90	115	pF
k_{low}	14	3.3	nS/mV
k_{high}	1.7	0.1	nS/mV
v_r	-60.6	-65.8	mV
v_{peak}	2.5	22.6	mV
v_t	-43.1	-57	mV
a	0.1	0.0012	ms^{-1}
b	-0.1	3	nS
c	-67	-65.8	mV
d	0.1	10	pA
τ_1^{AMPA}	0.25	0.2	ms
τ_2^{AMPA}	0.77	1.2	ms
τ_1^{GABA}	0.3	0.26	ms
τ_2^{GABA}	2.5	8.5	ms
τ_1^{NMDA}	2	-	ms
τ_2^{NMDA}	60	-	ms
e^{glu}	0	0	mV
e^{GABA}	-70	-70	mV
$\sigma_D{}^2$	$0.015/(n_{pyr}{}^2)$	-	-
C_{syn}	9	-	pF
k^{syn}	$3/n_{pyr}$	-	-
g^{leak}	5	-	nS
e^{leak}	-60.6	-	mV
k^{AMPA}	2^9	-	-
k^{NMDA}	2^{13}	-	-
k^{GABA}	80	80	-
k^{ext}	5	1	-

993
994

995 **Figure supplements**
996
997

998
999
1000

Figure 1 – Figure Supplement 1.

Somatic glutamate uncaging-evoked membrane responses.

(A) Comparison of uEPSPs to spontaneous EPSPs. Half-maximum width (ms) plotted

against peak amplitude (mV) for n = 210 spontaneous events (grey) and n = 78

uEPSPs (red); data from 4 cells. Inset: peak amplitude (i) and half-width (ii) for one

event.

(B) Distribution of arithmetic sum of peak amplitudes for oriens and radiatum

uncaging locations.

(C) As (B) but for arithmetic sum of time-integrals.

1010
1011

1012

1013

Figure 1 – Figure Supplement 2.

Unscaled uEPSP integration location-dependent nonlinearity.

Comparisons of the arithmetic sum of individual uEPSPs and corresponding recorded

uEPSPs evoked at locations in stratum radiatum (blue, n = 9) and stratum oriens (red,

n = 14).

(A, B) Unscaled uEPSP amplitudes, average (A) and raw data (B).

(C, D) Unscaled uEPSP integrals, average (C) and raw data (D).

For panels (A) and (C) linearly interpolated averages are shown ± SEM.

1022

1023

1024

1025

1026

Figure 1 – Figure Supplement 3.

uEPSP integration location-dependent nonlinearity by cell.

(A) Peak amplitude nonlinearity comparison by dendrite location. Statistics for

all data, oriens vs radiatum: 26.0 ± 6.0 % (n = 9) vs 5.4 ± 5.4 % (n = 8), *P* = 0.02.

Statistics for paired data oriens vs radiatum: 29.6 ± 8.7 % vs 6.4 ± 6.7 %, *P* = 0.01,

n = 6.

(B) Time-integral nonlinearity comparison by dendrite location. Statistics for all

data, oriens vs radiatum: 55.9 ± 11.0 % (n = 9) vs 9.9 ± 7.6 % (n = 8), *P* = 0.004.

Statistics for paired data, oriens vs radiatum: 59.3 ± 16.4 % vs 13.1 ± 9.6 %, *P* =

0.035, n = 6.

Closed circles: paired recordings, open circles: unpaired. Bars show mean ± SEM

of all recordings.

1039

1040

Figure 1 – Figure Supplement 4.

uEPSP nonlinearity does not depend on uncaging location distance from

soma or on the size of the arithmetic sum of uEPSPs.

(A) uEPSP amplitude nonlinearity plotted against glutamate uncaging site

distance from soma.

(B) uEPSP amplitude nonlinearity plotted against maximal arithmetic sum of

uEPSP amplitudes.

(C) As (A) but for time-integrals.

(D) As (B) but for time-integrals.

Statistics: *r* = Pearson correlation coefficient, *P* = associated two-tailed

significance.

1062
1063
1064

Figure 2 – Figure Supplement 1.

Peak amplitude nonlinearity in oriens dendrites is abolished by blocking

NMDARs with D-AP5 but not by blocking sodium channels with TTX.

(A) Summary scaled peak amplitude uEPSP data for 14 dendritic locations

recorded in D-AP5, n = 10 oriens; n = 4 radiatum. Right: quantified synaptic

integration nonlinearity. Oriens vs radiatum: 1.9 ± 2.7 % vs 7.3 ± 2.5 %, P = 0.27.

Oriens D-AP5 vs oriens control, P = 0.001. Radiatum D-AP5 vs radiatum control,

P = 0.66.

(B) Summary scaled peak amplitude uEPSP data for 25 dendritic locations

recorded with TTX in the ACSF, n = 16 oriens; n = 9 radiatum. Right: quantified

synaptic integration nonlinearity. Oriens vs radiatum: 22.6 ± 5.2% vs 2.9 ± 3.9%,

P = 0.016. Oriens TTX vs oriens control, P = 0.84. Radiatum TTX vs radiatum

control, P = 0.9.

Figure 2 – Figure Supplement 2.

Arithmetic sum maximum uEPSP amplitudes, and integration nonlinearity

vs uncaging location distances, across pharmacological conditions.

(A – C) Peak arithmetic uEPSP amplitudes by uncaging dendrite location in

control conditions (A, replotted from Figure 1 – Figure Supplement 1B for

comparison), D-AP5 (B), and TTX (C).

(D – F) uEPSP peak amplitude nonlinearity vs uncaging location distance from

soma in control conditions (D, replotted from Figure 1 – Figure Supplement 4 for

comparison), D-AP5 (E), and TTX (F).

Figure 3 – Figure Supplement 1.

Simulations including polyamine modulation of AMPA receptors show synaptic integration differences between strata oriens and radiatum dendrite locations.

(A) Mean percentage integral nonlinearity difference between oriens and radiatum dendrites as peak synaptic AMPA and NMDA receptor conductance is varied.

(B) As (A), but with 100 µM intracellular spermine. Polyamine modulation was simulated using a kinetic scheme based on (Bowie et al., 1998).

Values for peak synaptic conductance were calculated at -60 mV and +60 mV for AMPA and NMDA receptors respectively. n=16 radiatum, n=28 oriens.

1124

Figure 5 – Figure Supplement 1.

Modelling principal cell input cooperation onto feedback interneurons. (A)

Network schematic showing postsynaptic membrane locations of Izhikevich-

model PV+ interneuron.

(B) Steady state transfer resistance for a PV+ cell dendritic tree reconstructed

using the TREES toolbox (Cuntz et al., 2010). The matrix shows the electrotonic

'proximity' of neighbouring patches of membrane on the dendritic tree (colour

scale: μV/pA). The main branches of the dendritic tree are indicated at the

bottom.

(C) Input cooperativity matrix used in simulations. Inset shows zoomed view of

cooperativity matrix for the first 50 principal neurons.

1136
1137

1138
1139
1140

Figure 5 – Figure Supplement 2.

NMDARs help to maintain a sparse and sharp representation of a 'hump' of

excitation to the feedback circuit shown in Figures 5D.

(A) Mean firing rate by neuron during the first 75 ms (solid line) and last 75 ms

(dashed line) averaged over one hundred 300 ms-long simulations. The hump of

excitation was centred on neuron #125. Black: without NMDARs; red: with

NMDARs.

(B) Mean network firing rates during first and last 75 ms of network simulation runs.

Black: without NMDARs; red: with NMDARs.

(C) Standard deviation of average network "receptive field" during first and last 75

ms of simulation runs. Black: without NMDARs; red: with NMDARs.

1152
1153

1154