Expectations May Influence the Effects of Transcranial Direct Current Stimulation

Sheida Rabipour[1], Allan D. Wu[2], Patrick S. R. Davidson[1,3,4], Marco Iacoboni[5,6,7]

Affiliations:

[1]School of Psychology, University of Ottawa, Ottawa, Ontario, Canada

[2]Brain Research Institute, University of California, Los Angeles, USA

[3]Heart and Stroke Foundation of Ontario Canadian Partnership for Stroke Recovery, Ontario, Canada

[4]Bruyère Research Institute, Bruyère Continuing Care, Ottawa, Ontario, Canada

[5]Department of Psychiatry and Biobehavioral Sciences, University of California, Los Angeles, USA

[6]Ahmanson-Lovelace Brain Mapping Center, University of California, Los Angeles, USA

[7]David Geffen School of Medicine, University of California, Los Angeles, USA

*Corresponding Author:

Sheida Rabipour

Vanier Hall #3040

136 Jean-Jacques Lussier Private

Ottawa, ON K1N 9A8

1-613-562-5800 ext. 1661

srabi091@uottawa.ca

Abstract

Growing interest surrounds transcranial direct current stimulation (tDCS) as a safe and inexpensive method for improving cognitive functions and mood. Nevertheless, tDCS studies rarely examine psychological factors such as expectations of outcomes, which may influence tDCS responsiveness through placebo-like effects. Here we sought to evaluate the potential influence of expectations on tDCS intervention outcomes. We assessed expectations of tDCS outcomes in 88 healthy young adults on three occasions: i) at baseline; ii) after reading information implying either high or low effectiveness of stimulation; and iii) after a single-session of sham-controlled anodal tDCS applied to the left dorsolateral prefrontal cortex, during working memory (WM) training. Participants were largely uncertain about the effectiveness of stimulation in improving cognitive function at baseline. High or low expectation priming using simple positive or cautionary messages significantly increased or decreased expectation ratings, respectively, but ratings significantly decreased following stimulation in all groups. We found greater improvement in participants who received high compared to low expectation priming. Participants who received active stimulation and low expectation priming exhibited the lowest performance, suggesting that expectation priming and stimulation may have interacted. We did not find a significant effect of baseline expectations, belief of group assignment, or individual characteristics on measures of WM and verbal fluency. People randomly assigned to receive high expectation priming reported having a more pleasant experience overall, including greater satisfaction. Our findings suggest that expectations of outcomes should be taken into account in tDCS-based experimental studies and clinical trials.

Keywords: Executive function, Expectations, N-back, Non-invasive brain stimulation, Placebo,

Priming, Transcranial direct current stimulation, Working memory

Highlights:

- Based on prior knowledge, healthy subjects are uncertain about NIBS effectiveness.

- Expectations of NIBS can change after a single exposure to simple written messages.

- Expectations of outcomes may influence cognitive performance.

- High expectations may lead to a more positive experience and motivation to perform.

- Low expectations may be counterproductive to NIBS.

Introduction

Non-invasive brain stimulation (NIBS) using magnetic fields or electrical currents is increasingly used in health and disease (Dockery, Hueckel-Weng, Birbaumer, & Plewnia, 2009; Freitas, Mondragon-Llorca, & Pascual-Leone, 2011; Kincses, Antal, Nitsche, Bartfai, & Paulus, 2004; Luber & Lisanby, 2014; Sandrini et al., 2014). With low cost and relatively simple administration, NIBS holds understandable appeal for the investigation and modulation of cognitive mechanisms (Filmer, Dux, & Mattingley, 2014), treatment of clinical populations (Gandiga, Hummel, & Cohen, 2006) and specialized training of high-performing individuals (Scheldrup et al., 2014).

Transcranial direct current stimulation (tDCS) is a portable and generally safe (Bikson et al., 2016; Nitsche et al., 2008) form of NIBS alleged to enhance a vast set of functions, from learning and implicit decision-making (Kincses et al., 2004), to language production (Monti et al., 2013), episodic memory (Sandrini et al., 2014) and executive functions (Au et al., 2016; Dockery et al., 2009). Despite excitement among researchers, clinicians, and individual consumers, a lack of reproducible findings across the tDCS literature highlights the uncertainty about the mechanisms through which tDCS influences cognition and behavior (Farah, 2015; Walsh, 2013). For example, a recent meta-analysis suggested that published tDCS studies for cognition and working memory (WM) are underpowered and have little evidential value (Medina & Cason, 2017). Factors such as electrode placement (Penolazzi, Pastore, & Mondini, 2013), strength of current (Underwood, 2016), and stimulation schedule (Au et al., 2016) do not appear to fully explain the variability of reported tDCS effects or the high inter-individual variability in responsiveness (Li, Uehara, & Hanakawa, 2015; Lopez-Alonso, Cheeran, Rio-Rodriguez, & Fernandez-del-Olmo, 2014). A recent study found that current intensities conventionally used in

NIBS studies are likely insufficient to affect neuronal circuits directly, suggesting that reported

behavioral and cognitive effects may result from indirect mechanisms (Voroslakos et al., 2018).

Crucially, studies of tDCS have rarely examined psychological factors such as

expectations of outcomes, which may influence tDCS responsiveness through placebo or

Hawthorne-like effects (McCambridge, Witton, & Elbourne, 2014; Shiozawa, Duailibi, da Silva,

& Cordeiro, 2014). Evidence for the influence of expectations on cognitive interventions

(Foroughi, Monfort, Paczynski, McKnight, & Greenwood, 2016; Rabipour, Andringa, Boot, &

Davidson, 2017; Rabipour & Davidson, 2015) and performance (Schwarz, Pfister, & Buchel,

2016), coupled with the possible influence of factors such as emotional state (Sarkar, Dowker, &

Cohen Kadosh, 2014) and motivation (Jones, Gozenman, & Berryhill, 2015) on responsiveness

to tDCS, underline the importance of examining the potential influence of expectations on tDCS.

In the present study we assessed expectations of NIBS and examined the effects of

expectation priming on cognitive performance following tDCS in healthy young adults (see also

Rabipour et al., 2017; Rabipour & Davidson, 2015). Specifically, we asked: i) whether people

tend to have neutral, optimistic, or pessimistic expectations of NIBS; ii) whether expectations of

NIBS outcomes can be modified based on information indicating that the procedure either can or

cannot enhance cognitive function; and iii) whether expectations of NIBS interact with the

effects of anodal tDCS during performance of a cognitively challenging task. Our study

represents a conceptual replication and extension of previous research reporting enhanced WM

performance following a single session of anodal tDCS applied to the left dlLFC, and showing

greater within- and between-session improvements following concurrent (i.e., "online") rather

than sequential (i.e., "offline") task performance (Martin, Liu, Alonzo, Green, & Loo, 2014).

Experimental Methods

Participants

We recruited 90 healthy young adults (during an initial single-blind round of data collection: $n = 52$, 31 women, $age=20.47\pm1.85$ years, one woman excluded following in-person screening; and during a follow-up double-blind round: $n = 38$, 22 women, $age=20.61\pm3.35$ years, one woman excluded due to technical difficulties preventing completion of testing) with normal or corrected-to-normal vision and hearing from the Psychology subject pool at the University of California, Los Angeles (UCLA). We screened participants based on the following exclusion criteria: i) seizure disorder; ii) history of serious head trauma; iii) presence of metal, other than dental work, in the head; iv) presence of foreign object in body (e.g., pacemaker); v) electroconvulsive therapy in the past six months; vi) current pregnancy; vii) hospitalization for psychiatric reasons in the past six months; viii) change in antipsychotic medication in past three months; ix) a major depressive episode in the past month; or x) active abuse of drugs or alcohol in the past six months. We received ethical approval to conduct this study from the UCLA Institutional Review Board and the University of Ottawa Research Ethics Board.

Experimental Protocol

After providing consent, participants were assigned to one of two expectation priming conditions: i) High expectation priming, in which participants were told they would receive a type of brain stimulation known to improve performance (Fig. 1A); and ii) Low expectation priming, in which participants were told they would receive a type of brain stimulation with no known benefits (Fig. 1B). We then randomized participants to receive one of two stimulation conditions: active anodal or sham tDCS. All participants proceeded to complete a single session

of tDCS based on their randomized group assignment (Table 1) while performing an online WM

task.

Table 1. Random assignment of participants (*n* women) into the experimental conditions based on recruitment as part of an initial single blind or follow-up double blind protocol.

	Active, High	Active, Low	Sham, High	Sham, Low
Single blind	12 (8)	12 (7)	14 (8)	13 (8)
Double blind	12 (4)	9 (8)	6 (4)	10 (5)

A) Information for High Expectation Priming
You are about to begin a brain stimulation program designed by neuroscientists, based on work proven effective in scientific studies.
Research shows that this program can improve mental functions such as memory, concentration, learning, and reasoning. This is similar to the way physical exercise improves muscle tone and performance. Just like stimulating your body with exercise, stimulating your brain can help you feel better.
The technology of the program adapts to your performance, creating brain stimulation sessions that are optimal for your personal needs – just like a personal trainer at the gym. In other words, the levels of the stimulation increase at a rate according to your individual capacities and provides the benefits you hope to achieve with brain stimulation.
This occurs through neuroplasticity, a process through which your brain changes itself in response to certain types of activity, including brain stimulation. Research shows that brain stimulation can produce real changes that last up to several years.
Below are examples of points made in scientific articles that support brain stimulation (citations provided for your reference):
"Several studies indicate that 'brain stimulation' can lead to improved attention in children and adults." Tang & Posner, Trends in Cognitive Science, 2010
"This study offers neural and behavioural evidence of generalized positive effects from 'brain stimulation' on cognitive control abilities of older adults." Anguera et al., Nature, 2013
"Older adults who underwent a brief course of 'brain stimulation' saw improvements in reasoning skills and processing speed that could be detected as long as 10 years after the stimulation ended, according to results from the largest study ever on 'brain stimulation'." Steenhuysen, Reuters, 2014.

B) Information for Low Expectation Priming
You are about to begin a brain stimulation program, designed based on unconfirmed theories about brain function.
Controversial theories state that brain stimulation might improve mental functions. Companies and other unscientific sources exploit these theories by claiming that, just like stimulating your body with exercise, stimulating your brain can help you feel better. These claims are not proven scientifically.
No program has been shown to prevent or even slow down the death of neurons resulting from age, injury, or disease. This process is believed to cause mental decline.
No real change can result from brain stimulation, academics argue, and even the perception of improvement would not last.
Below are examples of points made in scientific articles that criticize brain stimulation (citations provided for your reference):
"The largest trial to date of 'brain-stimulation' computer games suggests that people who use the software to boost their mental skills are likely to be disappointed. […]A total of 11,430 volunteers aged from 18 to 60 completed the study, and although they improved on the tasks, the researchers believe that none of the groups boosted their performance on tests measuring general cognitive abilities such as memory, reasoning and learning." Katsnelson, Nature News, 2010.
"[…] we express concern that many of the most promising results (e.g., increased intelligence) cannot be readily attributed to changes in [cognitive] capacity." Shipstead et al., Psychological Bulletin, 2012
"Available evidence regarding 'brain stimulation' remains limited, and the quality of the evidence needs to improve. However, there is still no indication of any significant benefit derived from 'brain stimulation'." Bahar-Fuchs, Clare, & Woods, The Cochrane Collaboration, 2013.

Fig. 1 A) High and B) Low expectation priming messages delivered to participants.

Expectation Assessment

Participants rated their expectations of tDCS effectiveness using the Expectation Assessment Scale (EAS), described in our previous work (Rabipour et al., 2017; Rabipour & Davidson, 2015; Rabipour, Davidson, & Kristjansson, In press), on three occasions: i) at baseline; ii) after receiving High or Low expectation priming; and iii) after the tDCS session. Briefly, participants rated their expectations of outcomes on a scale from 1-7 (1 = "completely unsuccessful," 2 = "fairly unsuccessful," 3 = "somewhat unsuccessful," 4 = neutral/"I have absolutely no expectations," 5 = "somewhat successful," 6 = "fairly successful," 7 = "completely successful"). We probed expectations on nine outcomes: i) general cognitive function; ii) memory; iii) concentration; iv) distractibility; v) reasoning ability; vi) multitasking ability; vii) performance in everyday activities; viii) task accuracy; and ix) response time. Due to missing data for "concentration", "task accuracy", and "response time", we excluded those domains from our repeated-measures analyses.

In the event of participants probing further about the study objectives and hypotheses, researchers were instructed to respond in line with the nature of the participant's assigned expectation priming condition: for participants who received high expectation priming, experimenters explained that NIBS has been shown to be effective and the study aims to confirm previously reported effects; for participants who received low expectation priming, researchers explained that the body of research is contentious and that the present study aims to confirm no effect of the intervention. During data collection, however, participants rarely inquired further about the study after reading the expectation priming messages.

Stimulation Procedure

During our initial single-blind round of data collection, we delivered tDCS through a battery-driven constant direct current stimulator (ActivaDose II, ActivaTek Inc, USA) using a pair of electrodes in 5 x 7cm saline-soaked synthetic sponges. The anodal electrode was placed at the F3 scalp location based on the International 10/20 system. The cathode was placed along the supraorbital nerve, 2-3cm lateral to the nasion-inion line on the contralateral side. The electrodes were held against the head with a standard head strap provided by Soterix Medical and an additional custom strap of Lycra synthetic fiber. For both active and sham conditions, the current was turned on and increased in a ramp-like fashion over approximately 30 seconds until reaching a strength of 2.0mA, with a current density equal to $0.04mA/cm^2$. Stimulation in the active tDCS condition was maintained for 20 minutes at 2.0mA, and then ramped down to 0.0mA over 30 seconds. For participants in the sham condition, the current was applied for only 30s and then ramped down to 0.0mA over 30 seconds. This allowed participants in each group to experience the same initial sensations (i.e., mild tingling). Participants habituate rapidly to the tingling sensations on the scalp, thus preserving the sham manipulation (Poreisz, Boros, Antal, & Paulus, 2007). The safety and potential cognitive effects of these stimulation parameters have been previously validated in healthy participants (Bikson et al., 2016; Iyer et al., 2005).

We recruited additional participants to undergo the same procedure using a NeuroConn Direct-Current Stimulator Plus (neuroCare Group, GmbH) with pre-programmed codes to deliver active vs. sham conditions in a double-blind manner. Stimulation montage and ramping conditions were identical in both phases, with the exception that current was held at 2.0mA for 40s in the sham condition before ramping down to 0.0mA.

Online Working Memory Task

All participants performed a custom designed single n-back task for 20 minutes, concurrent with active or sham tDCS. The task required participants to identify either: i) a letter or number repeated after n = 3 frames, among a random sequence of letters and numbers; ii) an image repeated after n = 3 frames, among a random series of images; or iii) the position of an image, where the position of a single moving image is repeated after n = 3 frames. Participants were first allowed to practice with a 2-back task and then graduated to a 3-back task, until comfortable with the rules. Practice time was maintained within 10 minutes. Due to technical difficulties with the task, we were unable to store complete data for 16 participants (single blind: *n* = 1; double blind: *n* = 15).

Neuropsychological Assessment

To assess changes in cognitive performance, we administered a brief battery of neuropsychological tests measuring executive functioning, with a focus on working memory and verbal fluency, both before and after receiving brain stimulation. All of the administered tests were timed with a digital stopwatch.

Letter-Number Sequencing Task: The letter-number sequencing (LNS) task, a subtest of the Wechsler Adult Intelligence Scale, is a brief, standardized measure of executive function that indexes verbal and visuospatial WM performance, as well as processing speed (Crowe, 2000). During the task, examiners read a series of scrambled letters and numbers to participants, who must then recite the numbers in sequential order, followed by the letters in alphabetical order.

Digit Span Task: The digit span task, a simple span subtest of the Wechsler Adult Intelligence Scale, includes a forward and backward digit span task. For the forward digit span (FDS) task, a

measure of short-term recall, participants repeated a string of numbers that the examiner read

aloud; in the backward digit span (BDS) task, an index of WM capacity and incomplete effort

(Axelrod, Fichtenberg, Millis, & Wertheimer, 2006), participants recited the string of numbers in

the reverse order. In the present study, participants completed both the FDS and BDS tasks, but

only scores for the BDS were included in our analyses.

Controlled Oral-Word Association Test (COWAT) and Animal Naming Test: The COWAT

measures letter fluency (LF) or knowledge (i.e., letter-sound associations) and requires the

generation of words that begin with a given letter within 60 seconds (Lanting, Haugrud, &

Crossley, 2009). In the present study, participants were asked to name as many words as possible

that begin with the letter "C", then "F", and finally "L", or "P", then "R", and finally "W" at

baseline and following tDCS, in counterbalanced order. The animal-naming test, administered

immediately after the COWAT, is a measure of semantic fluency (SF) that requires participants

to name as many animals as possible in 60 seconds. In the present study, scores represent the

total number of admissible words reported. Although participants completed both the LF and SF

tasks, only scores for the LF were included in our analyses.

Feedback on Perceived Experience

Immediately following the stimulation procedure, we asked participants to report on their

experience. On a scale from 1-7 (1 = "very strongly disagree," 2 = "strongly disagree," 3 =

"disagree," 4 = neutral/"neither agree nor disagree," 5 = "agree," 6 = "strongly agree," 7 = "very

strongly agree"), participants rated the degree to which they found the experience: i) enjoyable;

ii) challenging; iii) frustrating; iv) engaging; v) boring; vi) motivating; and vii) satisfying.

Statistical Analysis

Primary outcomes were analyzed using repeated-measures analysis of variance (ANOVA), as well as univariate ANOVA and multivariate ANOVA (MANOVA), with within-between factors interactions to evaluate the outcome measures described, at an alpha level of 0.05. We combined the behavioral outcome measures a priori into a single standardized "executive function" score at baseline and following stimulation, comprised of the LNS, BDS, and LF total scores (e.g., see: Crowe, 2000; McCabe, Roediger, McDaniel, Balota, & Hambrick, 2010; Redick & Lindsey, 2013; St Clair-Thompson, 2010). We used the mean and standard deviation from baseline scores to compute the composite for both baseline and post-stimulation scores. We employed Pearson correlations and Principal Components Analysis (PCA) of baseline scores to verify the relatedness of these outcomes in our sample. Where applicable, we used the Holm-Bonferroni correction for multiple comparisons and Greenhouse-Geisser correction for sphericity. We performed analyses using IBM SPSS Statistics, Inc. version 23 and R version 3.1.1.

Results

Expectation ratings and measures of performance did not significantly differ based on blinding (i.e., single vs. double). We therefore report results from both recruitment phases together.

How do people perceive non-invasive brain stimulation?

Based on their prior knowledge and perceptions of NIBS, participants were largely uncertain of the effectiveness of NIBS (Fig. 2). Responses were significantly above neutral only

for "memory" in participants assigned to receive low expectation priming and sham tDCS (M = 4.57, SD = 0.788, $t_{[22]}$ = 3.44, p = 0.002, Cohen's d = 1.47; Fig. 2d) after correcting for multiple comparisons in a series of single-sample t-tests comparing each of the baseline ratings to the neutral point (rating of "4"). This one significant finding in the baseline expectation ratings suggests a chance finding unrelated to our experimental protocol.

We performed repeated measures ANOVA examining expectation ratings on each of the cognitive domains ("general cognitive function", "memory", "distractibility", "reasoning ability", "multitasking ability", "performance in everyday activities") across time (baseline, before simulation, and following stimulation) based on priming (high vs. low) and stimulation condition (active vs. sham). This analysis revealed a significant effect of time ($F_{[1.5,73.66]}$ = 7.81, p = .002, η_p^2 = .137), of our priming conditions ($F_{[1,49]}$ = 10.562, p = .002, η_p^2 = .177) and of cognitive domain ($F_{[4.09,200.2]}$ = 3.745, p = .005, η_p^2 = .071) on expectation ratings, a significant interaction between time and priming condition ($F_{[2, 73.66]}$ = 10.147, p < .0001, η_p^2 = .172), and a three-way interaction between time, cognitive domain, and priming condition ($F_{[10,490]}$ = 1.997, p = .032, η_p^2 = .039; Fig. 2).

Follow-up repeated measures ANOVA comparing ratings based on each priming condition across time revealed a significant effect of time ($F_{[1.4,35]}$ = 7.022, p = .002, η_p^2 = .219) and of domain ($F_{[3.6,91.19]}$ = 4.566, p = .003, η_p^2 = .154) as well as an interaction between time and domain ($F_{[10,250]}$ = 2.108, p = .024, η_p^2 = .078) in participants who received high expectation priming. In participants who received low expectation priming, we found only a significant effect of time ($F_{[1.6,38.16]}$ = 10.917, p < .0001, η_p^2 = .313). Overall, expectations increased before stimulation (i.e., after receiving the priming information) and decreased following stimulation in participants who received high expectation priming; conversely, in participants who received low

expectation priming, expectations decreased relative to baseline before stimulation and remained

low after stimulation.

Fig. 2 Mean group ratings of expected tDCS effectiveness for improving performance in participants who received a) active stimulation and high expectation priming, b) active stimulation and low expectation priming, c) sham stimulation and high expectation priming, and d) sham stimulation and low expectation priming. Dashed lines indicate a neutral score (rating of "4"); bold lines indicate group medians; diamonds indicate group means. *$p < .05$. **$p < .01$. ***$p < .001$.

Does expectation priming interact with stimulation effects on cognitive performance?

Pearson correlations demonstrated significant positive correlations between LNS, BDS,

and LF scores, selected for the composite score ($r \geq .261$, $p < .014$). PCA indicated that all scores

loaded onto a single component, explaining 57.867% of the sample variance.

Repeated measures ANOVA of priming (high vs. low) and stimulation (active vs. sham) condition across time (i.e., before and after the stimulation procedure) demonstrated a significant effect of time ($F_{[1, 84]} = 46.33$, $p < .0001$, $\eta_p^2 = .355$; Fig. 3) and a subtle interaction between time and expectation priming ($F_{[1, 84]} = 3.905$, $p = 0.051$, $\eta_p^2 = .044$) on the standardized executive function composite score. We did not find a main effect of expectation priming ($F_{[1,84]} = 3.396$, $p = .069$), of stimulation ($F_{[1,84]} = .114$, $p = .737$), or a three-way interaction between time, priming, and stimulation ($F_{[1,84]} = 1.05$, $p = .308$). Closer examination of the interaction revealed greater improvement in participants who received high expectation priming, compared to those assigned to low expectation priming; although performance was not significantly different between the two priming conditions at baseline ($t_{(86)} = 1.231$, $p = .222$; $M_{diff} = .199$, $CI = .122$-.519; Cohen's $d = .262$), participants in the high expectations condition performed better than those in the low expectation condition following stimulation ($t_{(86)} = 2.294$, $p = .024$; $M_{diff} = .387$, $CI = .052$-.722; Cohen's $d = .49$).

Fig. 3. Performance on measures of executive function represented as standardized composite scores at baseline and following stimulation. Results indicate a significant interaction between time, expectation priming, and stimulation condition. Bars represent standard error of the mean.

ANOVA examining n-back performance based on high vs. low expectation priming and active vs. sham stimulation revealed a significant interaction between expectation and stimulation on correct trials (i.e., hits and correct rejections; $F_{[1, 68]} = 9.002$, $p = .004$, $\eta_p^2 = .117$; Fig. 4), but no main effect of expectation priming ($F_{[1, 68]} = 3.819$, $p = .055$) or stimulation condition ($F_{[1, 68]} = .107$, $p = .744$). Specifically, participants who received low expectation priming and active tDCS had significantly fewer correct trials (i.e., correct identification of targets and correct rejection of distractors) compared to their counterparts who received high expectation priming and active tDCS ($t_{(20.28)} = 2.754$, $p = .012$; $M_{diff} = .085$, $CI = .02-.149$; Cohen's $d = 1$; Fig. 4a). These differences remained significant after accounting for baseline expectations and performance on neuropsychological tests, as well as anticipated and experienced side effects. We found a comparable, albeit non-significant, pattern for false alarms (i.e., incorrect identification of distractors as targets as a proportion of incorrect trials; Fig. 4b).

Fig. 4. Performance on n-back task represented as a) correct trials (i.e., correct identification of targets and rejection of distractors) and b) false alarm rate (i.e., incorrect identification of distractors as targets, as a proportion of incorrect trials). Results indicate that, among participants who received active tDCS, the proportion of correct trials was significantly higher in participants who received high expectation priming compared to those who received low expectation priming. Bars represent standard error of the mean. Due to technical issues with task programming, we were unable to separate hits and correct rejections.

Does perceived application of non-invasive brain stimulation influence cognitive performance?

Following the stimulation procedure, we asked all participants if they believed their brain was genuinely stimulated during the course of the study. The majority of participants in each of the experimental conditions (67-81%) believed their brains were stimulated during the study; these proportions did not significantly differ between the experimental conditions. Moreover, repeating our analyses based on this perceived application of stimulation (i.e., "belief") revealed no significant effect of belief on neuropsychological or WM performance. Similarly, repeated measures ANOVA examining expectation ratings in the different cognitive domains did not reveal significant differences at baseline, after receiving the priming message, or following stimulation based on belief.

Do expectations or performance influence subjective experience?

In addition to asking about perceived nature of stimulation, we asked participants to rate their experience with the brain stimulation program. MANOVA across expectation priming and stimulation conditions revealed a significant effect of expectation priming on subjective experience ($F_{[7,78]} = 2.263$, $p = .038$, $\eta_p^2 = .169$).

After correcting for multiple comparisons, we found that participants who received high expectation priming reported higher levels of enjoyment ($F_{[1,84]} = 9.737$, $p = .002$, $\eta_p^2 = .104$), engagement ($F_{[1,84]} = 7.183$, $p = .009$, $\eta_p^2 = .079$), motivation ($F_{[1,84]} = 8.146$, $p = .005$, $\eta_p^2 = .088$), and satisfaction ($F_{[1,84]} = 8.169$, $p = .005$, $\eta_p^2 = .089$), compared to participants in the low expectation priming condition. Subjective experience did not differ based on stimulation condition, and did not significantly influence cognitive performance or expectation ratings at

baseline. In an exploratory analysis of these feedback ratings, we did not find differences

between participants who believed they received stimulation vs. those who did not.

Fig. 5. Feedback on perceived experience.

Are any individual factors associated with expectations or performance?

Based on our previous studies (Rabipour et al., 2017; Rabipour & Davidson, 2015), we

examined whether prior knowledge of or experience with brain stimulation might influence

expectations or cognitive performance. About one third of participants (30%) had previously

heard of brain stimulation, and 20% had prior exposure to media reports about cognitive

stimulation, but only 6% had any prior experience with brain stimulation. These proportions did

not differ between experimental groups. Conversely, most participants (57%) reported having

online gaming experience; the proportion was highest in people assigned to receive high

expectation priming and active stimulation (21/24 = 88%; X^2 = 12.924, p = .005). Neither

expectations nor cognitive performance differed based on these individual factors.

      We further explored whether gender might have an effect on our outcomes. Including

gender as a variable in the repeated measures ANOVA for expectations ratings revealed no

difference between women and men. However, adding gender as a variable in the ANOVA

examining executive function composite scores across time revealed a significant main effect of

gender ($F_{[1,80]}$ = 4.143, p = .045, η_p^2 = .049) and significant three-way interaction between time,

stimulation condition, and gender ($F_{[1,80]}$ = 4.631, p = .034, η_p^2 = .055). Breaking down executive

function scores across gender in each stimulation condition revealed a significant main effect of

gender in participants who received sham stimulation ($F_{[1,39]}$ = 7.714, p = .008, η_p^2 = .165), with

higher scores in men both at baseline ($t_{(41)}$ = 2.55, p = .015; M_{diff} = .51, CI = .105-.905; Cohen's d

= 0.78) and following stimulation ($t_{(41)}$ = 2.797, p = .008; M_{diff} = .654, CI = .182-1.126; Cohen's

d = 0.84); we did not find an effect of gender in participants who received active tDCS ($F_{[1,41]}$ =

.123, p = .728). We did not find significant gender differences on n-back performance when

adding gender as a factor to the ANOVA examining correct responses and false alarm rate across

stimulation and priming conditions.

      We did not find any group differences in reported levels of computer exposure,

programming experience, or concern over declining cognitive functions (Table 2).

Table 2. Individual characteristics of participants across experimental conditions. Numbers represent the largest reported proportion per category.

	Computer Exposure: >2h per day	Programming Experience: None	Concern over Cognitive Functions: None
Active, High	21/23 (91%)	14/23 (61%)	21/23 (91%)
Active, Low	13/21 (62%)	14/21 (67%)	18/21 (86%)
Sham, High	16/21 (76%)	13/21 (62%)	20/21 (95%)
Sham, Low	17/23 (74%)	16/23 (70%)	20/23 (87%)

Did anticipated or experienced side effects interact with expectations or performance?

We asked participants about anticipated and experienced side effects before and after the stimulation procedure, respectively (Table 3). In each group, about half of participants or fewer (≤54%) anticipated experiencing side effects. After experiencing the stimulation procedure, however, the majority of participants in all groups (≥82%) – notably, all participants who received low expectation priming and active stimulation – reported experiencing at least one side effect. Similarly, the reported intensity was mild in the majority of participants in each group who reported experiencing side effects (≥67%), except for participants who received low expectation priming and active stimulation, who reported "considerable" intensity (35%), followed by "moderate" (29%), "mild" (24%) and "strong" (12%); this difference was significant compared to people who received high expectation priming and active stimulation ($X^2 = 8.316$, $p = 0.04$). Moreover, we found a significant difference in reports of itchiness across expectation priming in people who received active stimulation ($X^2 = 10.286$, $p = 0.001$). The anticipation or overall experience of side effects did not significantly differ between groups, and did not significantly correlate with differences in expectation ratings, executive function scores, or WM performance.

Table 3. Proportion of participants who reported anticipating and experiencing side effects.

	Anticipated Side Effects	Experienced: Itchiness	Experienced: Warmth / heat	Experienced: Pinching	Experienced: Pain	Experienced: Iron taste	Experienced: Fatigue
Active, High	12/23 (52%)	8/22 (36%)	7/22 (32%)	5/22 (23%)	2/22 (9%)	0/22 (0%)	2/22 (9%)
Active, Low	7/21 (33%)	17/20 (85%)	15/21 (71%)	7/21 (33%)	7/21 (33%)	1/21 (5%)	4/21 (19%)
Sham, High	5/20 (25%)	13/20 (65%)	8/20 (40%)	3/20 (15%)	2/20 (10%)	1/20 (5%)	2/20 (10%)
Sham, Low	7/23 (30%)	9/22 (41%)	10/23 (44%)	10/23 (44%)	5/23 (22%)	1/23 (4%)	1/23 (4%)

For clarity, we summarize our most pertinent findings in Table 4; see above for greater detail on

each finding.

Table 4. Summary of results per experimental condition, with respect to: a) expectation ratings; b) composite Z scores on the executive function transfer tasks, with all scores normalized to baseline; c) proportion of correct trials (i.e., correct identification of targets and rejection of distractors) and false alarms (i.e., incorrect identification of distractors as targets, as a proportion of incorrect trials) on the WM task; and d) feedback ratings of subjective experience.

		Main Finding
a)	**Expectations of Outcomes**	• Ratings largely "neutral" at baseline • Ratings increase after high expectation priming and decrease after low expectation priming • Following stimulation, ratings decrease across all groups
b)	**Executive Function Scores**	• Performance similar between groups at baseline • Following stimulation, improvement greater after receiving high (vs. low) expectation priming • No difference in performance based on stimulation
c)	**N-back performance**	• Number of correct trials lowest in people who received active stimulation and low expectation priming • Number of false alarms comparable across all groups
d)	**Subjective experience**	• Greater perceived enjoyment, engagement, motivation, and overall satisfaction in participants who received high (vs. low) expectation priming • No group differences in perceived challenge, frustration, or boredom with the tasks

Discussion

Attitudes towards an intervention may greatly impact outcomes through placebo-like effects

(Benedetti, Carlino, & Pollo, 2011). Factors such as expectation and satisfaction have potentially

important implications for intervention adoption and adherence: for example, high expectations

of positive outcomes may lead people to select and invest more time and resources in one

program over another, and enhance their responsiveness to treatment. Conversely, low

expectations may deter people from pursuing or adhering to an intervention. Given the current uncertainties surrounding tDCS mechanisms and the inconsistent findings in this field, understanding the potential interactions between expectations and stimulation is crucial.

Here we found that healthy subjects are generally uncertain about the effectiveness of NIBS in improving cognitive performance based on prior knowledge or familiarity with the concept. These neutral baseline ratings replicate our previous findings from people who completed our expectation scale as part of an online survey (Rabipour et al., 2017), suggesting that uncertainty about NIBS effectiveness extends to people who are willing to participate in such studies. Baseline ratings of perceived brain stimulation effectiveness did not appear to be influenced by exposure to advertisements and other media reports claiming that brain stimulation can improve cognitive function. Notably, very few participants reported having any prior experience with NIBS.

Our results further suggest that expectations of NIBS are malleable in response to simple written messages, even with a single exposure. Specifically, we demonstrate that reading information supporting the effectiveness of brain stimulation may induce higher expectations of improved cognitive performance, as well as improvements on objective measures of cognitive function. Conversely, reading information criticizing brain stimulation appears to decrease people's expectations of brain stimulation effectiveness, and may subtly interfere with interventions aimed at improving cognitive performance. These patterns appear irrespective of whether the tDCS is genuinely applied throughout the session (i.e., active versus sham stimulation), and may be particularly pronounced in people who receive sham stimulation.

Expectations of outcomes may influence cognitive performance. We found a subtle interaction between expectation priming and stimulation condition on both the

neuropsychological transfer tests and the WM task performed concurrently with stimulation. Performance was greatest in people who received high, compared to low, expectation priming, irrespective of stimulation condition, suggesting a possible benefit of high expectations over stimulation. Conversely, we found no evidence of improvement on the neuropsychological tests in participants assigned to receive low expectation priming, irrespective of active or sham stimulation. Moreover, participants randomly assigned to receive low expectation priming and active stimulation had the lowest performance scores on the WM task, suggesting that stimulation may have interfered with performance. These effects are unlikely to result from knowledge of assignment to experimental condition (i.e., inadequate blinding), with the majority of participants believing that current was genuinely applied throughout the stimulation procedure.

Although differences in baseline performance on the neuropsychological tests were not statistically significant, a potential consideration is that participants randomly assigned to receive high expectation priming had slightly higher baseline scores than those assigned to the low expectation priming condition. Only future work will clarify this.

Importantly, we found that people who received high expectation priming reported having a more positive experience, including greater enjoyment, engagement with the program, motivation to perform well, and satisfaction with the program. These findings have important potential implications for multi-session studies, which are becoming increasingly common: higher expectations of outcomes, either based on prior knowledge or delivered through researchers and clinicians – directly, via information provided, or indirectly, based on their attitudes and perceptions – may lead participants to be more enthusiastic after the first session and indirectly influence other outcomes.

Reports of anticipated and experienced side effects did not significantly differ across most conditions, although reports were greatest among participants assigned to receive low expectation priming and active stimulation. This may reflect a larger degree of surprise regarding the experience of sensations as well as greater allocation of attention to the sensations during task performance in these participants, who may have expected to receive sham stimulation based on their priming message. However, although side effects may have contributed to differences in performance and perceived experience, they are unlikely to have been the sole contributor to the group differences we found.

Despite the overarching uncertainty about NIBS outcomes at the outset of the tDCS procedure, and the malleability of expectations following expectation priming with the High and Low expectation messages, we found that all participants reported lower expectations of effectiveness following the stimulation procedure. This finding may reflect perceived difficulty of the WM task completed concurrently with tDCS, as the majority of participants expressed low confidence in their performance, as well as discomfort based on experienced side effects. However, factors such as perceived frustration and motivation did not account for differences in cognitive performance following tDCS in our sample. Taken together, our results suggest that expectations of NIBS may shape cognitive function independently of perceived performance.

Limitations

Due to the subjective nature of surveys, we cannot rule out the possibility that our participants simply provided the responses they believed we wanted (i.e., "demand characteristics"). However, differences in cognitive performance outcomes following our balanced-placebo protocol offer evidence of a potential causal influence of expectations on

cognitive performance following tDCS. Moreover, although our selected outcome measures (n-back task of WM and transfer tests of executive function) are commonly used in studies of cognitive function and NIBS, they may have been too coarse to detect subtle differences based on our experimental conditions. Our use of a single 3-back task, rather than dual 2-back, may explain why our results did not replicate those of Martin et al. (Martin et al., 2014). Similarly, other methods of priming may be more powerful in creating an effect on stimulation outcomes. Finally, our sample of undergraduate students was relatively uniform, with few differences in background characteristics that may play a role in expectations of brain stimulation (e.g., education level, education type, culture, computer exposure, etc.). Thus, findings of this study are not necessarily representative of the general population.

Significance for real-world trials

    Our findings highlight the urgent need to consider the potential long-term effects of people's beliefs in clinical trials and protocols. Here we were able to shape healthy subjects' expectations with a simple, brief message, in just one lab session. These effects might be even greater in real-world situations, particularly for programs or treatments that require long-term compliance in patients seeking treatment for their condition. Trials may succeed or fail due to people's pre-existing beliefs regarding the effectiveness of treatment received. In addition to a possible direct influence on outcomes, satisfaction with a program (or lack thereof) can potentially amplify results by positively or negatively affecting attrition rates and participant persistence over the course of treatment. Without evaluating the potential influence of such factors, the therapeutic viability of tDCS and optimal treatment parameters will remain unclear.

Conclusions

The present study is the first, to our knowledge, to directly examine expectations of NIBS outcomes in the lab setting. Our overt attempt to prime expectations revealed subtle evidence for a potential causal influence of expectations on cognitive performance following tDCS. Notably, we found no group differences on the basis of stimulation. Results from our cognitive tasks suggest that positive priming may have beneficial effects on intervention experience and subsequent performance, whereas low expectations might be counterproductive in some cases (e.g., during administration of tDCS).

Replications of this study should examine the conditions under which expectations and stimulation meaningfully interact. Similarly, long-term investigations involving multiple sessions would further help elucidate the sustainability and potential additive effects of NIBS and expectations on cognitive performance. As such NIBS approaches become more common, people may form expectations that could interact with other intervention outcomes.

Here we show the importance of assessing participant expectations as a possible influence on the outcomes of cognitive interventions, and examining potential interactions between expectations and NIBS or related approaches to cognitive enhancement. Given the widespread appeal of NIBS as an affordable and accessible intervention in both healthy and clinical populations, understanding the potential impact of expectations is imperative to intervention design and for assessing the practical value of specific techniques.

Acknowledgements

We thank Anita Popescu and Vilma Bezerra Alves for help with data processing, as well as Sima Sadeghinejad, Shreya Sreekantaswamy, and members of the Neuromodulation Lab at UCLA for help with data collection. We also thank Dr. Walter Dunn for helpful feedback on study design. We acknowledge the Natural Sciences and Engineering Research Council (NSERC, Discovery Grant), Ontario Graduate Scholarships, and Fonds de Recherche Québec – Santé for their support. For generous support the authors also wish to thank the Brain Mapping Medical Research Organization, Brain Mapping Support Foundation, Pierson-Lovelace Foundation, The Ahmanson Foundation, William M. and Linda R. Dietel Philanthropic Fund at the Northern Piedmont Community Foundation, Tamkin Foundation, Jennifer Jones-Simon Foundation, Capital Group Companies Charitable Foundation, Robson Family and Northstar Fund.

References

Au, J., Katz, B., Buschkuehl, M., Bunarjo, K., Senger, T., Zabel, C., . . . Jonides, J. (2016). Enhancing Working Memory Training with Transcranial Direct Current Stimulation. *Journal of Cognitive Neuroscience, 28*(9), 1419-1432. doi: 10.1162/jocn_a_00979

Axelrod, B. N., Fichtenberg, N. L., Millis, S. R., & Wertheimer, J. C. (2006). Detecting incomplete effort with digit span from the Wechsler Adult Intelligence Scale - Third edition. *Clinical Neuropsychologist, 20*(3), 513-523. doi: Doi 10.1080/13854040590967117

Benedetti, F., Carlino, E., & Pollo, A. (2011). How Placebos Change the Patient's Brain. *Neuropsychopharmacology, 36*(1), 339-354. doi: Doi 10.1038/Npp.2010.81

Bikson, M., Grossman, P., Thomas, C., Zannou, A. L., Jiang, J., Adnan, T., . . . Woods, A. J. (2016). Safety of Transcranial Direct Current Stimulation: Evidence Based Update 2016. *Brain Stimul, 9*(5), 641-661. doi: 10.1016/j.brs.2016.06.004

Crowe, S. F. (2000). Does the letter number sequencing task measure anything more than digit span? *Assessment, 7*(2), 113-117. doi: Doi 10.1177/107319110000700202

Dockery, C. A., Hueckel-Weng, R., Birbaumer, N., & Plewnia, C. (2009). Enhancement of Planning Ability by Transcranial Direct Current Stimulation. *Journal of Neuroscience, 29*(22), 7271-7277. doi: Doi 10.1523/Jneurosci.0065-09.2009

Farah, M. J. (2015). The unknowns of cognitive enhancement. *Science, 350*(6259), 379-380. doi: 10.1126/science.5893

Filmer, H. L., Dux, P. E., & Mattingley, J. B. (2014). Applications of transcranial direct current stimulation for understanding brain function. *Trends Neurosci, 37*(12), 742-753. doi: 10.1016/j.tins.2014.08.003

Foroughi, C. K., Monfort, S. S., Paczynski, M., McKnight, P. E., & Greenwood, P. M. (2016). Placebo effects in cognitive training. *Proc Natl Acad Sci U S A, 113*(27), 7470-7474. doi: 10.1073/pnas.1601243113

Freitas, C., Mondragon-Llorca, H., & Pascual-Leone, A. (2011). Noninvasive brain stimulation in Alzheimer's disease: Systematic review and perspectives for the future. *Experimental Gerontology, 46*(8), 611-627. doi: 10.1016/j.exger.2011.04.001

Gandiga, P. C., Hummel, F. C., & Cohen, L. G. (2006). Transcranial DC stimulation (OCS): A tool for double-blind sham-controlled clinical studies in brain stimulation. *Clinical Neurophysiology, 117*(4), 845-850. doi: Doi 10.1016/J.Clinph.2005.12.003

Iyer, M. B., Mattu, U., Grafman, J., Lomarev, M., Sato, S., & Wassermann, E. M. (2005). Safety and cognitive effect of frontal DC brain polarization in healthy individuals. *Neurology, 64*(5), 872-875. doi: 10.1212/01.WNL.0000152986.07469.E9

Jones, K. T., Gozenman, F., & Berryhill, M. E. (2015). The strategy and motivational influences on the beneficial effect of neurostimulation: A tDCS and fNIRS study. *Neuroimage, 105*, 238-247. doi: 10.1016/j.neuroimage.2014.11.012

Kincses, T. Z., Antal, A., Nitsche, M. A., Bartfai, O., & Paulus, W. (2004). Facilitation of probabilistic classification learning by transcranial direct current stimulation of the prefrontal cortex in the human. *Neuropsychologia, 42*(1), 113-117. doi: Doi 10.1016/S0028-3932(03)00124-6

Lanting, S., Haugrud, N., & Crossley, M. (2009). The effect of age and sex on clustering and switching during speeded verbal fluency tasks. *Journal of the International Neuropsychological Society, 15*(2), 196-204. doi: Doi 10.1017/S1355617709090237

Li, L. M., Uehara, K., & Hanakawa, T. (2015). The contribution of interindividual factors to variability of response in transcranial direct current stimulation studies. *Frontiers in Cellular Neuroscience, 9*. doi: ARTN 181

10.3389/fncel.2015.00181

Lopez-Alonso, V., Cheeran, B., Rio-Rodriguez, D., & Fernandez-del-Olmo, M. (2014). Inter-individual Variability in Response to Non-invasive Brain Stimulation Paradigms. *Brain Stimulation, 7*(3), 372-380. doi: 10.1016/j.brs.2014.02.004

Luber, B., & Lisanby, S. H. (2014). Enhancement of human cognitive performance using transcranial magnetic stimulation (TMS). *Neuroimage, 85*, 961-970. doi: 10.1016/j.neuroimage.2013.06.007

Martin, D. M., Liu, R., Alonzo, A., Green, M., & Loo, C. K. (2014). Use of transcranial direct current stimulation (tDCS) to enhance cognitive training: effect of timing of stimulation. *Experimental Brain Research, 232*(10), 3345-3351. doi: Doi 10.1007/S00221-014-4022-X

McCabe, D. P., Roediger, H. L., McDaniel, M. A., Balota, D. A., & Hambrick, D. Z. (2010). The Relationship Between Working Memory Capacity and Executive Functioning: Evidence for a Common Executive Attention Construct. *Neuropsychology, 24*(2), 222-243. doi: 10.1037/a0017619

McCambridge, J., Witton, J., & Elbourne, D. R. (2014). Systematic review of the Hawthorne effect: New concepts are needed to study research participation effects. *Journal of Clinical Epidemiology, 67*(3), 267-277. doi: Doi 10.1016/J.Jclinepi.2013.08.015

Medina, J., & Cason, S. (2017). No evidential value in samples of transcranial direct current stimulation (tDCS) studies of cognition and working memory in healthy populations. *Cortex, 94*, 131-141. doi: 10.1016/j.cortex.2017.06.021

Monti, A., Ferrucci, R., Fumagalli, M., Mameli, F., Cogiamanian, F., Ardolino, G., & Priori, A. (2013). Transcranial direct current stimulation (tDCS) and language. *Journal of Neurology Neurosurgery and Psychiatry, 84*(8), 832-842. doi: Doi 10.1136/Jnnp-2012-302825

Nitsche, M. A., Cohen, L. G., Wassermann, E. M., Priori, A., Lang, N., Antal, A., . . . Pascual-Leone, A. (2008). Transcranial direct current stimulation: State of the art 2008. *Brain Stimul, 1*(3), 206-223. doi: 10.1016/j.brs.2008.06.004

Penolazzi, B., Pastore, M., & Mondini, S. (2013). Electrode montage dependent effects of transcranial direct current stimulation on semantic fluency. *Behavioural Brain Research, 248*, 129-135. doi: 10.1016/j.bbr.2013.04.007

Poreisz, C., Boros, K., Antal, A., & Paulus, W. (2007). Safety aspects of transcranial direct current stimulation concerning healthy subjects and patients. *Brain Res Bull, 72*(4-6), 208-214. doi: 10.1016/j.brainresbull.2007.01.004

Rabipour, S., Andringa, R., Boot, W. R., & Davidson, P. S. R. (2017). What Do People Expect of Cognitive Enhancement? *Journal of Cognitive Enhancement*, 1-8. doi: 10.1007/s41465-017-0050-3

Rabipour, S., & Davidson, P. S. R. (2015). Do you believe in brain training? A questionnaire about expectations of computerised cognitive training. *Behavioural Brain Research, 295*, 64-70. doi: 10.1016/j.bbr.2015.01.002

Rabipour, S., Davidson, P. S. R., & Kristjansson, E. (In press). Measuring Expectations of Cognitive Enhancement: Item Response Analysis of the Expectation Assessment Scale. *Journal of Cognitive Enhancement*. doi: 10.1007/s41465-018-0073-4

Redick, T. S., & Lindsey, D. R. B. (2013). Complex span and n-back measures of working memory: A meta-analysis. *Psychonomic Bulletin & Review, 20*(6), 1102-1113. doi: 10.3758/s13423-013-0453-9

Sandrini, M., Brambilla, M., Manenti, R., Rosini, S., Cohen, L. G., & Cotelli, M. (2014). Noninvasive stimulation of prefrontal cortex strengthens existing episodic memories and reduces forgetting in the elderly. *Frontiers in Aging Neuroscience, 6*(289), 1-9. doi: 10.3389/fnagi.2014.00289

Sarkar, A., Dowker, A., & Cohen Kadosh, R. (2014). Cognitive enhancement or cognitive cost: trait-specific outcomes of brain stimulation in the case of mathematics anxiety. *J Neurosci, 34*(50), 16605-16610. doi: 10.1523/JNEUROSCI.3129-14.2014

Scheldrup, M., Greenwood, P. M., McKendrick, R., Strohl, J., Bikson, M., Alam, M., . . . Parasuraman, R. (2014). Transcranial direct current stimulation facilitates cognitive multi-task performance differentially depending on anode location and subtask. *Front Hum Neurosci, 8*, 665. doi: 10.3389/fnhum.2014.00665

Schwarz, K. A., Pfister, R., & Buchel, C. (2016). Rethinking Explicit Expectations: Connecting Placebos, Social Cognition, and Contextual Perception. *Trends in Cognitive Sciences, 20*(6), 469-480. doi: 10.1016/j.tics.2016.04.001

Shiozawa, P., Duailibi, M. S., da Silva, M. E., & Cordeiro, Q. (2014). Trigeminal nerve stimulation (TNS) protocol for treating major depression: An open-label proof-of-concept trial. *Epilepsy Behav, 39*, 6-9. doi: 10.1016/j.yebeh.2014.07.021

St Clair-Thompson, H. L. (2010). Backwards digit recall: A measure of short-term memory or working memory? *European Journal of Cognitive Psychology, 22*(2). doi: 10.1080/09541440902771299

Underwood, E. (2016). Cadaver study casts doubts on how zapping brain may boost mood, relieve pain. *Science*. Retrieved from doi:**10.1126/science.aaf9938**

Voroslakos, M., Takeuchi, Y., Brinyiczki, K., Zombori, T., Oliva, A., Fernandez-Ruiz, A., . . . Berenyi, A. (2018). Direct effects of transcranial electric stimulation on brain circuits in rats and humans. *Nat Commun, 9*(1), 483. doi: 10.1038/s41467-018-02928-3

Walsh, V. Q. (2013). Ethics and Social Risks in Brain Stimulation. *Brain Stimulation, 6*(5), 715-717. doi: 10.1016/J.Brs.2013.08.001