Drosophila egg-derived tyrosine phosphatase (EDTP): a novel target for improved survivorship to prolonged anoxia and cellular protein aggregates

Short title: *Drosophila EDTP* and improved survival

Chengfeng Xiao[a, *], Shuang Qiu[a, b], Xiao Li[c], Dan-Jun Luo[c], R Meldrum Robertson[a], Gong-Ping Liu[c, d, *]

[a] Department of Biology, Queen's University, Kingston, Ontario K7L 3N6 Canada

[b] Jiangsu Key Laboratory of Chemical Pollution Control and Resources Reuse, School of Environmental and Biological Engineering, Nanjing University of Science and Technology, Xiao Ling Wei 200, Nanjing 210094, Jiangsu, P. R. China

[c] Department of Pathophysiology, Key Laboratory of Neurological Diseases of Chinese Ministry of Education, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430030, P. R. China

[d] Co-Innovation Center of Neuroregeneration, Nantong University, Nantong JS 226001, P.R. China

[*] Corresponding authors: xiao.c@queensu.ca (C.X.) and liugp111@mail.hust.edu.cn (G-P. L.)

Abstract

Drosophila egg-derived tyrosine phosphatase (EDTP), a lipid phosphatase that removes 3-position phosphate at the inositol ring, has dual functions in the oogenesis and the muscle performance during adult stages. A mammalian homologous gene *MTMR14*, which encodes the myotubularin-related protein 14, negatively regulates autophagy. Mutation of *EDTP/MTMR14*, however, causes at least three deleterious consequences: (1) lethality in the early embryogenesis in *Drosophila*; (2) "jumpy" phenotype with apparently impaired motor functions; and (3) association with a rare genetic disorder called centronuclear myopathy. Here we show that flies carrying a heterozygous *EDTP* mutation had increased survivorship to prolonged anoxia; tissue-specific downregulation of *EDTP* in non-muscle tissues, particularly motoneurons, extended the lifespan; and tissue-specific downregulation of *EDTP* in motoneurons improved the survivorship to beta-amyloid peptides (Aβ42) and polyglutamine (polyQ) protein aggregates. MTMR14 expression was evident in the hippocampus and cortex in C57BL/6J and APP/PS1 mice. Compared with C57BL/6J mice, APP/PS1 mice had reduced MTMR14 in the cortex but not in the hippocampus. Hippocampal expression of MTMR14 was increased and plateaued at 9-17 months compared with 2-6 months in C57BL/6J mice. Aβ42 treatment increased the expression of MTMR14 in C57BL/6J primary culture neurons and Neuro2a neuroblasts. We demonstrated a novel approach of tissue-specific manipulation of the disease-associated gene *EDTP/MTMR14* for lifespan extension and the improvement of survivorship to cellular protein aggregates.

Keywords:

Egg-derived tyrosine phosphatase; myotubularin-related protein 14; survival; prolonged anoxia; cellular protein aggregate; tissue-specific expression; motoneuron; *Drosophila*

1. Introduction

Drosophila egg-derived tyrosine phosphatase (EDTP) is a lipid phosphatase that removes 3-position phosphate at the inositol ring of phosphatidylinositol 3-phosphate (PtdIns3P) and phosphatidylinositol (3, 5)-bi-phosphate [PtdIns(3,5)P2] (Tosch et al., 2006). EDTP possesses a function opposite to Vps34, a sole class III phosphoinositide 3-kinase (Hiles et al., 1992; Schu et al., 1993), in the regulation of the PtdIns3P pool. The most interesting characteristic of EDTP expression is that there are two peaks, one at oogenesis (Krauchunas et al., 2012; Yamaguchi et al., 2005), and another at adult stages (Seroude et al., 2002). The transcription of *MTMR14*, a mouse homolog of *EDTP*, also shows a peak at day five of differentiation in C2C12 myoblasts, followed by a decline (Tosch et al., 2006). Human *MTMR14* transcripts are detectable at the ages between 19 and 69 years in the tested tissues, including heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas (Tosch et al., 2006). Levels of *MTMR14* are highly coincident with those of *Vps34* in the human brain, heart, and skeletal muscles (Volinia et al., 1995), indicating a tightly regulated biological process which is associated with cellular PtdIns3P levels.

The decline of EDTP between early embryogenesis and young adult stages is accompanied with the *Drosophila* metamorphosis, a process requiring extensive autophagy and apoptosis for histolysis (Zirin et al., 2013). These observations suggest a role for EDTP in the regulation of autophagy. This is indeed supported by the findings that PtdIns3P stimulates autophagy in human HT-29 cells (Petiot et al., 2000), that MTMR14 negatively controls the autophagosome formation and maturation in mammalian cells (Vergne et al., 2009), and that the EDTP/MTMR14 inhibitor, AUTEN-99, activates autophagy in human HeLa cells and mouse tissues (Kovács et al., 2017). Interestingly, the latter group also shows that overexpression of a modified *EDTP* (by skipping the first exon) in the fat body antagonizes the effect of AUTEN-67 in inducing autophagy (Papp et al., 2016).

Despite the negative regulation of autophagy, deletion of *Drosophila EDTP* is lethal during embryogenesis or in the first instar, and germline clones with a null *EDTP* allele fail to produce mature oocytes (Yamaguchi et al., 2005). Homozygous flies carrying a hypomorphic *EDTP* allele are short-lived with impaired motor functions and reduced fecundity (Edelman, 2005). Additionally, muscles of the *MTMR14*-deficient mice have decreased force production, prolonged relaxation and exacerbated fatigue (Shen et al., 2009). A human MTMR14 missense variant (R336Q) is associated with centronuclear myopathy, a rare genetic disorder with muscle weakness and wasting (Tosch et al., 2006). Therefore, the function of autophagy initiation is likely overwhelmed by the lethality or disease-causing effects due to a ubiquitous loss of *EDTP/MTMR14*.

There are advantages of autophagy in degrading and recycling disrupted organelles, long-lived proteins, and denatured protein aggregates (Petiot et al., 2000; Takeshige et al., 1992). A strategy to maximize the potential benefit of *EDTP* is to manipulate its downregulation in favorable tissues while maintaining normal levels in oocytes and muscles at adult stages. This seems to be feasible by using the *Drosophila* Gal4/UAS expression system (Brand and Perrimon, 1993). We thus hypothesized that selective downregulation of *EDTP* in non-muscle tissues, particularly the central nervous system, extends lifespan and improves the survivorship to cellular protein aggregates in *Drosophila*.

In the current study, we demonstrate that heterozygous *EDTP* mutants had improved survival to prolonged anoxia, a condition mimicking the extreme hypoxia that induces autophagy in

mammalian and human cells (Bellot et al., 2009). We also show that selective downregulation of *EDTP* in the fly motoneurons extended lifespan and increased the survivorship to beta-amyloid peptides (Aβ42) or polyglutamine (polyQ) protein aggregates. The expression of MTMR14 in the hippocampus and cortex in mice was evident, promising a potential application by targeting *EDTP/MTMR14* for the treatment of neurodegeneration.

2. Materials and methods

2.1. Animals

Fly strains used in this study and their sources were: $EDTP^{DJ694}$ and w^{1118} (L. Seroude laboratory, Queen's University), D42-Gal4 (Bloomington Drosophila Stock Center (BDSC) #8816), UAS-EDTP-RNAi (II) (with an RNAi sequence CAGTAGTGTAATAGTAATCAA targeting exon 3 of *EDTP*, BDSC #41633), UAS-EDTP-RNAi (III) (with an RNAi sequence CAGCTACGACGAAGTCATCAA targeting exon 2, BDSC #36917), UAS-Aβ42 (BDSC #32038), UAS-Httex1-Q103-eGFP (Zhang et al., 2010). Flies were maintained with standard medium (cornmeal, agar, molasses, and yeast) at 21-23 °C with a 12:12 photoperiod.

Heterozygous mutants ($EDTP^{DJ694}$/+) and their sibling controls (w^{1118}) were prepared by two consecutive crosses between homozygous $EDTP^{DJ694}$ mutants and w^{1118} flies. We chose virgin female offspring ($EDTP^{DJ694}$/+) from the first mating and crossed them with w^{1118} males. Flies carrying the *EDTP* mutation (red-eyed) and their siblings (white-eyed) were collected for survival analysis.

C57BL/6J mice were purchased from the Experimental Animal Centre of Wuhan University. C57BL/6J mice are wild-type whereas APP/PS1 (stock number 004462, Jackson Laboratory) are the transgenic mouse models of Alzheimer's disease. APP/PS1 mice express a chimeric mouse/human amyloid precursor protein (Mo/HuAPP695swe) and a mutant human presenilin 1 (PS1-dE9) both directed to CNS neurons. The mice were housed in a light/dark (12:12) cycle in standard group cages (4-5 per cage) with accessible food and water *ad libitum*. The experimental procedures with mice were approved by the Ethics Committee of Tongji Medical College, Huazhong University of Science and Technology.

2.2. Post-anoxia survival in flies

Newly emerged flies were collected and aged to 6-8 days in regular culture conditions and exposed to an anoxia (generated by pure argon gas) during the light phase of the photoperiod. The percent survival at 12 h (% 12-h survival) after anoxia was examined. For the tests of post-anoxia survival time, flies were exposed to anoxia for 0, 1, or 6 h. The dead flies were scored daily during the first week of recovery, and twice a week thereafter until all the flies were counted. Throughout the long-term survival experiments, alive flies were transferred to fresh food vials twice a week.

2.3. Lifespan experiments

Newly emerged flies were collected and raised at a density of 20-25 flies per vial. They were transferred into fresh culture media twice a week. The lifespan was scored by counting dead flies

during each transfer until all the flies were counted. This procedure was also used for the examination of the lifespan of flies expressing β amyloid (Aβ) peptides N-42 (Aβ42) or polyglutamine (polyQ) protein aggregates (with 103 tandem repeats) in motoneurons, and the lifespan of flies with simultaneous expression of Aβ42 or polyQ aggregates and RNAi knockdown of *EDTP* in the same cells.

2.4. Cell Culture

For primary neuron cultures, hippocampal neurons of 18 days embryonic (E18) rat were seeded at 30,000 - 40,000 cells per well on 6-well plates coated with Poly-D-Lysine/Laminin (Bioscience) in neurobasal medium (Invitrogen), which was supplemented with 2% B27/0.5 mM glutamine/25 mM glutamate. Half of the culture medium was changed every 3 days with neurobasal medium supplemented with 2% B27 and 0.5 mM glutamine. All cultures were kept at 37 ℃ in a humidified 5% CO_2 cultural condition.

Neuro2a cells were cultured at 37 °C in a 1:1 mixture of DMEM (Dulbecco's modified Eagle's medium) and OPTI-MEM supplemented with 15% FBS (fetal bovine serum, Gibco), 100 units/ml penicillin, and 100 mg/ml streptomycin. A humidified atmosphere containing 5 % CO_2 was provided. The cells were plated on to six-well plates overnight and treated with 5 μM Aβ42 or vehicle control (DMSO) for 24 h.

2.5. Immunohistochemistry

Mice were anesthetized and perfused through aorta with 0.9% NaCl followed by phosphate buffer containing 4% paraformaldehyde. The brains were removed and post-fixed in perfusate overnight. Tissue sectioning (20 µm) was performed with a vibratome (Leica Biosystems). The sections were fixed with 0.3% H_2O_2 in the absolute methanol for 30 min and saturated with bovine serum albumin (BSA) for 30 min at room temperature. Tissues were then incubated with polyclonal anti-MTMR14 overnight at 4 °C. After wash, the tissues were incubated with appropriate secondary antibody. Immunoreaction was developed with diaminobenzidine using Histostain-SP Kits (Zymed, CA, USA). Images were taken using a light microscope (Olympus BX60, Tokyo, Japan).

2.6. Western Blotting

Western blotting was performed as described previously (Duan et al., 2013). The hippocampal region and cortex were quickly removed and homogenized in the sample buffer (Tris-HCl 50mM (pH 7.4), NaCl 150 mM, NaF 10mM, Na_3VO_4 1mM, EDTA 5 mM, benzamidine 2 mM, and phenylmethylsulphonyl fluoride 1 mM). The extracts were mixed with sample buffer (3:1, v/v, containing 200 mM Tris-HCl (pH 7.6), 8% SDS, 40% glycerol, 40mM dithiothreitol) and incubated with boiled water for 10 min. The proteins were separated by 10% SDS-polyacrylamide gel electrophoresis and transferred to nitrocellulose membrane. The membrane was blocked for 1 h with 5% skim milk (dissolved in TBSTween-20 containing 50mM Tris-HCl (pH 7.6), 150mM NaCl, 0.2% Tween-20), and probed with primary antibody at 4 °C overnight. Blots were incubated with anti-rabbit or anti-mouse IgG conjugated to IRDye (800 CW) at room

temperature and visualized using the Odyssey Infrared Imaging System (LI-COR Biosciences, Lincoln, NE, USA). Protein bands were analyzed using ImageJ (Schneider et al., 2012).

2.7. Statistics

   Statistical analysis was conducted by using R (R Development Core Team, 2016) and the following packages: gdata, survival, survminer, and ggplot2. Briefly, two-way ANOVA with Turkey's test was applied for the analysis of 12-h survival of different genotypes of flies which were subject to anoxia with varying durations. Log-rank test was performed for the comparison of two survival curves. Log-rank test with the Benjamini - Hochberg ("BH") adjustment was used for the comparison of three survival curves. One-way ANOVA with Bonferroni's multiple comparisons was performed to analyze the MTMR14 expression among three groups of cells. A student *t*-Test was conducted to compare the levels of MTMR14 between two groups of cells. A value of $P < 0.05$ was considered statistical significance.

3. Results

3.1. Improved 12-h survival to prolonged anoxia in male EDTP mutants

   $EDTP^{DJ694}$/+ flies and their sibling w^{1118}controls were exposed to an anoxia (generated by pure nitrogen gas) for 1, 3, or 6 h at the ages of 6-8 days. After 1 h anoxia, the average of percent survival at 12 h (% 12-h survival) of male flies was 100.0 ± 0.0 % (Mean $\pm$ SD) in both $EDTP^{DJ694}$/+ groups (n = 4) and controls (n = 3) (Fig. 1a). With 3 h exposure, the % 12-h survival in male $EDTP^{DJ694}$/+ groups (83.3 ± 10.8 %, n = 6) was higher than their controls (65.3 ± 18.4 %, n = 6) ($P = 0.0144$, Two-way ANOVA with Turkey's test). After 6 h anoxia, the % 12-h survival in male $EDTP^{DJ694}$/+ groups (57.6 ± 12.9 %, n = 7) was remarkably higher than their controls (25.4 ± 4.3 %, n = 7) ($P < 0.0001$, Two-way ANOVA with Turkey's test). The genotype, anoxia duration, and their interaction were all the factors contributing significantly to the % 12-h survival in male flies (Two-way ANOVA) (Fig. 1b).

   Female $EDTP^{DJ694}$/+ flies had the % 12-h survival statistically the same as w^{1118} flies with any of the 1, 3, or 6 h exposure. Two-way ANOVA indicated that while anoxia duration significantly contributed to % 12-h survival, the genotype or the interaction factor displayed an insignificant contribution (Fig. 1b).

   Therefore, male $EDTP^{DJ694}$/+ flies displayed improved 12-h survival to the prolonged anoxia (i.e. 3 - 6 h duration) but not the short-term exposure (i.e. 1 h).

3.2. Increased post-exposure survivorship to prolonged anoxia in EDTP mutants

   The increased 12-h survival to prolonged anoxia raised an immediate concern: whether *EDTP* mutants could have increased post-exposure survivorship. Young flies (6-8 days old) were exposed to 0, 1, or 6 h anoxia and the post-exposure survivorship was examined.

   Without anoxia, the median survival time was 69 d (95% confidence interval (CI) 65 - 69 d, n = 102) in $EDTP^{DJ694}$/+ males and 69 d (CI 65 - 69 d, n = 96) in controls. There was no statistical difference in survival time between these groups ($P = 0.945$, Log-rank test) (Fig. 2). There was also no significant difference in survival time between female $EDTP^{DJ694}$/+ (median

74 d, CI 69 - 79 d, n = 107) and their controls (median 74 d, CI 69 - 79 d, n = 97) (*P* = 0.869, Log-rank test). Thus, *EDTP* mutation had no effect on the survivorship without anoxia.

With 1 h anoxia, male *EDTP^{DJ694}*/+ flies had post-exposure survival (median 86 d, CI 80 - 86 d, n = 57) the same as controls (median 93 d, CI 86 - 99 d, n = 51) (*P* = 0.139, Log-rank test). Female *EDTP^{DJ694}*/+ showed the post-exposure survival (median 93 d, CI 93 - 99 d, n = 44) also the same as controls (median 93 d, CI 86 - 99 d, n = 58) (*P* = 0.167, Log-rank test). *EDTP* mutation displayed no effect on the post-exposure survivorship after 1 h anoxia.

With 6 h anoxia, male *EDTP^{DJ694}*/+ flies showed post-exposure survival (median 36 d, CI 1 - 72 d, n = 167) markedly longer than controls (median 1 d, CI 1 - 1 d, n = 163) (*P* = 0.0033, Log-rank test). Female *EDTP^{DJ694}*/+ also showed the post-exposure survival (median 63.5 d, CI 37 - 79 d, n = 150) remarkably longer than controls (median 11 d, CI 8 - 37 d, n = 148) (*P*= 0.0001, Log-rank test).

Thus, both male and female *EDTP^{DJ694}*/+ flies had increased post-exposure survivorship to 6 h anoxia.

3.3. RNAi knockdown of EDTP in motoneurons extended lifespan

Extreme hypoxia induces autophagy in mammalian and human cells (Bellot et al., 2009). The mammalian *MTMR14*, a homolog of *Drosophila EDTP*, negatively regulates the autophagic removal of protein aggregates (Vergne et al., 2009). These data together with our findings suggested that downregulation of *EDTP* could promote autophagy and thus have a beneficial effect on survivorship. We used the Gal4/UAS binary system (Brand and Perrimon, 1993) for selective downregulation of *EDTP* in motoneurons and examined its effect on lifespan. The rationale for selective targeting of motoneurons was to avoid affecting the abundant expression of EDTP in muscles, ovaries, spermatheca, and oocytes. The motoneuron-specific driver D42-Gal4 (Yeh et al., 1995) was used for RNAi knockdown of *EDTP*.

We used the UAS-EDTP-RNAi (III) line with a short interference sequence targeting the transcripts from exon 2 of *EDTP*. Flies with *EDTP* knockdown in motoneurons (D42-Gal4/UAS-EDTP-RNAi (III)) had a median lifespan of 113 d (CI 107 – 118 d, n = 132), which was longer than UAS controls (UAS-EDTP-RNAi (III)/+, median 89 d, CI 89 – 98 d, n = 63) (*P* < 0.0001, Log-rank test with "BH" adjustment), and also longer than Gal4 controls (D42-Gal4/+, median 73 d, CI 73 – 77 d, n = 137) (*P* < 0.0001, Log-rank test with "BH" adjustment) (Fig. 3a). Therefore, downregulation of *EDTP* in motoneurons resulted in extended lifespan.

To confirm this finding, we used a different RNAi line (UAS-EDTP-RNAi (II) with an interference sequence targeting the transcripts from exon 3) to knockdown *EDTP* in the cells. Flies with *EDTP* knockdown in motoneurons (UAS-EDTP-RNAi (II)/+; D42-Gal4/+) had the lifespan (median 74.5 d, CI 69 – 84 d, n = 114) significantly longer than UAS controls (median 58 d, CI 51 – 58 d, n = 79) (*P* < 0.0001, Log-rank test with "BH" adjustment), or Gal4 controls (median 51 d, CI 51 – 54 d, n = 87) (*P* < 0.0001, Log-rank test with "BH" adjustment) (Fig. 3b).

These data indicated that downregulation of *EDTP* in motoneurons extended the lifespan in *Drosophila*.

3.4. RNAi knockdown of EDTP in motoneurons improved the survivorship to beta-amyloid peptides or polyglutamine protein aggregates

We next asked whether downregulation of *EDTP* could increase the survivorship to cellular protein aggregates expressed in the motoneurons. We examined the survivorship of flies with simultaneous *EDTP* downregulation and the expression of toxic β amyloid (Aβ) peptides N-42 (Aβ42) (Kuo et al., 1996). The median lifespan in flies expressing Aβ42 in motoneurons (UAS-Aβ42/Y;; D42-Gal4/+) was 64 d (CI 64 - 64 d, n = 111). The median lifespan was increased to 94 d (CI 94 - 98 d, n = 86) in flies with Aβ42 expression and simultaneous RNAi knockdown of *EDTP* (UAS-Aβ42/Y;; D42-Gal4/UAS-EDTP-RNAi) (P < 0.0001, Log-rank test) (Fig. 4a).

We also examined the survivorship of flies with *EDTP* downregulation and simultaneous expression of polyglutamine (polyQ) protein aggregates in the motoneurons. The median lifespan was 38 d (CI 38 - 38 d, n = 45) in flies (UAS-Httex1-Q103-eGFP/Y;; D42-Gal4/+) expressing polyQ aggregates (each containing a polyQ tract of 103 tandem repeats of glutamine). The median lifespan was significantly increased to 44.5 d (CI 41 – 48 d, n = 30) in flies (UAS-Httex1-Q103-eGFP/Y;; D42-Gal4/UAS-EDTP-RNAi) with polyQ expression and simultaneous *EDTP* downregulation (P < 0.0001, Log-rank test) (Fig. 4b).

Therefore, *EDTP* downregulation in motoneurons improved the survivorship to the expression of Aβ42 or polyQ protein aggregates.

3.5. Expression of MTMR14 in the hippocampus and cortex in C57BL/6J and APP/PS1 mice

Human *MTMR14* is transcribed ubiquitously at relatively low levels in the brain (Fagerberg et al., 2014). MTMR14 is expressed in mouse muscle, liver, and fat but there is little evidence for expression in the brain (Lv et al., 2015). We examined the expression of MTMR14 in the hippocampus and cortex in C57BL/6J and APP/PS1 mice. C57BL/6J is a wild-type strain while APP/PS1 is a transgenic mouse model of Alzheimer's disease.

MTMR14 staining was observable in the CA3 pyramidal neurons and dentate gyrus (DG) in the hippocampus of a 3-month-old C57BL/6J mouse (Fig. 5a). MTMR14 expression was detectable in the hippocampus in both C57BL/6J and APP/PS1 mice. The levels in the two strains were not significantly different (Fig. 5b). MTMR14 expression was also detectable in the cortex in both strains. However, relative levels of MTMR14 in APP/PS1 mice were lower than those in C57BL/6J mice (P < 0.05, *t*-Test). These data suggested an association between transgenic expression of amyloid precursor protein/presenilin 1 and MTMR14 level.

Relative levels of MTMR14 in the C57BL/6J hippocampus increased and plateaued at 9, 12 and 17 months compared with the levels at 2 and 6 months (Fig. 5c), indicating an age-dependent kinetics of expression.

3.6. Increased MTMR14 induced by Aβ42 in primary hippocampal neurons and Neuro2a cells

Aβ42 causes microRNA deregulation in primary hippocampal neurons in C57BL/6J mice (Schonrock et al., 2010) and inhibits the viability of the mouse neuroblastoma Neuro2a cells (Dahlgren et al., 2002). We explored the effect of Aβ42 treatment on MTMR14 expression in primary hippocampal neurons and Neuro2a cells.

Relative MTMR14 levels were increased in hippocampal primary neurons after the 72-h inoculation with Aβ42 at 5 or 10 µM compared with controls. Levels of MTMR14 were higher in the neurons treated with 10 µM Aβ42 than the neurons treated with 5 µM Aβ42 ($P < 0.01$, One-way ANOVA with Bonferroni's multiple comparisons) (Fig. 6a). Data indicated a dosage-dependent induction of MTMR14 by Aβ42 treatment. We also examined the MTMR14 expression in Neuro2a cells with Aβ42 treatment. MTMR14 expression was increased in Neuro2a cells incubated with Aβ42 at 5 µM for 24 h compared with controls ($P < 0.01$, t-Test) (Fig. 6b). Therefore, Aβ42 treatment increased the expression of MTMR14 in primary hippocampal neurons and Neuro2a cells.

4. Discussion

We report in *Drosophila* a novel approach of fine-tuning the disease-associated gene *EDTP/MTMR14* for lifespan extension and improved survivorship to cellular protein aggregates. Cell-specific downregulation of *EDTP* in non-muscle tissues circumvents the deleterious consequences due to ubiquitous loss of *EDTP*. Specifically, downregulation of *EDTP* in motoneurons extends lifespan and increases the survivorship to Aβ42 or polyQ protein aggregates. The expression of the mouse *MTMR14* in the hippocampus and cortex, together with its age-dependent kinetics and the Aβ42-induced increase of MTMR14 in cultured neurons, promises a potential application by targeting *EDTP/MTMR14* in a tissue-specific manner for the treatment of neurodegeneration.

The proposed cell-specific downregulation of *EDTP/MTMR14* for beneficial consequences is formulated largely from the preliminary observation that heterozygous flies carrying a hypomorphic allele of *EDTP* have increased survivorship to prolonged anoxia. Homozygous flies carrying the same allele display a "jumpy" phenotype – impaired motor function associated with shortened lifespan and reduced fecundity (Edelman, 2005). These findings indicate that severe downregulation of *EDTP* in flies homozygous for mutant allele is overall deleterious whereas moderate downregulation of *EDTP* in flies heterozygous for the mutation is wild-type-like and protective from prolonged anoxia. Such a conditional benefit highlights the importance of subtle manipulation of EDTP/MTMR14 spatiotemporal expression to have the desired effect. *EDTP/MTMR14* has two main function: negative regulation of autophagy and disease-causing effect upon deficiency in the muscles (Edelman, 2005; Kovács et al., 2017; Papp et al., 2016; Shen et al., 2009; Tosch et al., 2006; Vergne et al., 2009). *Drosophila EDTP* has two peaks of expression, one at oogenesis and another at adult stages around 20-30 days (Krauchunas et al., 2012; Seroude et al., 2002; Yamaguchi et al., 2005). Both male and female adult flies have abundant EDTP in muscles and additionally, female flies express rich amounts of EDTP in the spermatheca and ovaries. The differential expression of EDTP in female flies might also be responsible for the lack of effect on short-term survival (i.e. % 12-h survival) to prolonged anoxia. The dual functions, multiple peaks over time, a broad range of expressing tissues, and sex difference together do not likely allow a complete deletion of *EDTP/MTMR14* to be associated with expected benefits. Precise control of EDTP/MTMR14 expression in the favorable cells or tissues could be essential for eliciting the expected consequences. The current study presents a strategy of cell-specific downregulation of *Drosophila EDTP* in motoneurons with beneficial effects in lifespan extension and improved survivorship to cellular protein aggregates.

Motoneurons are highly specialized, terminally differentiated, and have reduced ability for global genome repair (Aranda-Anzaldo, 2012; Nouspikel and Hanawalt, 2000). More importantly, neuronal cells utilize autophagy as an essential process for normal turnover of cytoplasmic contents (Hara et al., 2006). *Drosophila* motoneuronal expression of a human Cu-Zn superoxide dismutase (SOD1) leads to a marked extension of lifespan by up to 40% (Parkes et al., 1998). Motoneuronal overexpression of the heat shock protein 70 results in structural plasticity of axonal terminals which is associated with increased larval thermotolerance (Xiao et al., 2007). Motoneurons innervate muscle cells through neuromuscular junctions, making them an ideal target for *EDTP* downregulation while leaving muscular expression intact. Our findings that motoneuronal downregulation of *EDTP* extends lifespan and improves the survivorship to Aβ42 or polyQ protein aggregates firmly support motoneurons as a favorable targeting tissue.

EDTP/MTMR14 possesses a function opposite to Vps34 in the regulation of the PtdIns3P pool. Vps34 is the only phosphatidylinositol 3-kinases in yeast (Schu et al., 1993) and has been evolutionarily conserved through mammals. Vps34 plays an essential role in the process of autophagy (Backer, 2008; Jaber et al., 2012). Downregulation of *EDTP/MTMR14* shifts the focus from Vps34 to its functional opponent pertaining to the regulation of PtdIns3P and thus represents a novel approach to manipulate the process of autophagy.

Expression of MTMR14 in the mouse hippocampus and cortex provides a foundation from which the cell-specific manipulation of *EDTP/MTMR14* expression could be deployed in mammals. Interestingly, transgenic expression of a chimeric mouse/human amyloid precursor protein (Mo/HuAPP695swe) and a mutant human presenilin 1 (PS1-dE9) in APP/PS1 mice is associated with MTMR14 downregulation in the cortex but not in the hippocampus. These findings indicate the involvement of MTMR14 expression in the deposit of amyloid peptides and this involvement is tissue-specific. Additionally, MTMR14 expression displays an age-related kinetics and MTMR14 is inducible by Aβ42 treatment in primary hippocampal neurons as well as Neuro2a cells, suggesting a close relation between MTMR14 expression and aging. Notably, MTMR14 expression in response to the presence of amyloid peptides is different between *in vivo* animal experiments and cultured neurons. Perhaps the elevation of MTMR14 represents a short-term response to Aβ42, whereas MTMR14 downregulation in the animal studies is an active involvement for the removal of amyloid peptides. Nevertheless, our findings raise a promising direction by targeting MTMR14 for lifespan extension and survival improvement to cellular protein aggregates in mammals.

Conflict of interest

All authors declare no competing interests

Acknowledgements

This study was funded by Natural Sciences and Engineering Research Council of Canada (NSERC) grant (RGPIN 40930-09 to R.M.R.), and the Natural Science Foundation of China (81501211 to D.J. L.).

References

Aranda-Anzaldo, A., 2012. The post-mitotic state in neurons correlates with a stable nuclear higher-order structure. Commun. Integr. Biol. doi:10.4161/cib.18761

Backer, J.M., 2008. The regulation and function of Class III PI3Ks: novel roles for Vps34. Biochem. J. 410, 1–17. doi:10.1042/BJ20071427

Bellot, G., Garcia-Medina, R., Gounon, P., Chiche, J., Roux, D., Pouysségur, J., Mazure, N.M., 2009. Hypoxia-induced autophagy is mediated through hypoxia-inducible factor induction of BNIP3 and BNIP3L via their BH3 domains. Mol Cell Biol 29, 2570–2581.

Brand, A.H., Perrimon, N., 1993. Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. Development 118, 401–415.

Dahlgren, K.N., Manelli, A.M., Stine, W.B., Baker, L.K., Krafft, G.A., LaDu, M.J., 2002. Oligomeric and Fibrillar Species of Amyloid-β Peptides Differentially Affect Neuronal Viability. J. Biol. Chem. 277, 32046–32053. doi:10.1074/jbc.M201750200

Duan, D.-X., Chai, G.-S., Ni, Z.-F., Hu, Y., Luo, Y., Cheng, X.-S., Chen, N.-N., Wang, J.-Z., Liu, G.-P., 2013. Phosphorylation of tau by death-associated protein kinase 1 antagonizes the kinase-induced cell apoptosis. J. Alzheimers. Dis. 37, 795–808. doi:10.3233/JAD-130377

Edelman, S.W., 2005. Molecular Characterisation of the Age-regulated Jumpy Gene in Drosophila melanogaster. Library and Archives Canada= Bibliothèque et Archives Canada.

Fagerberg, L., Hallström, B.M., Oksvold, P., Kampf, C., Djureinovic, D., Odeberg, J., Habuka, M., Tahmasebpoor, S., Danielsson, A., Edlund, K., Asplund, A., Sjöstedt, E., Lundberg, E., Szigyarto, C.A.-K., Skogs, M., Takanen, J.O., Berling, H., Tegel, H., Mulder, J., Nilsson, P., Schwenk, J.M., Lindskog, C., Danielsson, F., Mardinoglu, A., Sivertsson, A., von Feilitzen, K., Forsberg, M., Zwahlen, M., Olsson, I., Navani, S., Huss, M., Nielsen, J., Ponten, F., Uhlén, M., 2014. Analysis of the human tissue-specific expression by genome-wide integration of transcriptomics and antibody-based proteomics. Mol. Cell. Proteomics 13, 397–406. doi:10.1074/mcp.M113.035600

Hara, T., Nakamura, K., Matsui, M., Yamamoto, A., Nakahara, Y., Suzuki-Migishima, R., Yokoyama, M., Mishima, K., Saito, I., Okano, H., Mizushima, N., 2006. Suppression of basal autophagy in neural cells causes neurodegenerative disease in mice. Nature 441, 885–889. doi:10.1038/nature04724

Hiles, I.D., Otsu, M., Volinia, S., Fry, M.J., Gout, I., Dhand, R., Panayotou, G., Ruiz-Larrea, F., Thompson, A., Totty, N.F., others, 1992. Phosphatidylinositol 3-kinase: structure and expression of the 110 kd catalytic subunit. Cell 70, 419–429.

Jaber, N., Dou, Z., Chen, J.-S., Catanzaro, J., Jiang, Y.-P., Ballou, L.M., Selinger, E., Ouyang, X., Lin, R.Z., Zhang, J., Zong, W.-X., 2012. Class III PI3K Vps34 plays an essential role in autophagy and in heart and liver function. Proc. Natl. Acad. Sci. 109, 2003–2008. doi:10.1073/pnas.1112848109

Kovács, T., Billes, V., Komlós, M., Hotzi, B., Manzéger, A., Tarnóci, A., Papp, D., Szikszai, F., Szinyákovics, J., Rácz, Á., others, 2017. The small molecule AUTEN-99 (autophagy enhancer-99) prevents the progression of neurodegenerative symptoms. Sci Rep 7.

Krauchunas, A.R., Horner, V.L., Wolfner, M.F., 2012. Protein phosphorylation changes reveal

new candidates in the regulation of egg activation and early embryogenesis in D. melanogaster. Dev Biol 370, 125–134.

Kuo, Y.M., Emmerling, M.R., Vigo-Pelfrey, C., Kasunic, T.C., Kirkpatrick, J.B., Murdoch, G.H., Ball, M.J., Roher, A.E., 1996. Water-soluble Abeta (N-40, N-42) oligomers in normal and Alzheimer disease brains. J. Biol. Chem. 271, 4077–81. doi:10.1074/jbc.271.8.4077

Lv, Y., Xue, L., Cai, C., Liu, Q.H., Shen, J., 2015. Deficiency of myotubularin-related protein 14 influences body weight, metabolism, and inflammation in an age-dependent manner. Cell Biosci. 5. doi:10.1186/s13578-015-0062-6

Nouspikel, T., Hanawalt, P.C., 2000. Terminally differentiated human neurons repair transcribed genes but display attenuated global DNA repair and modulation of repair gene expression. Mol. Cell. Biol. 20, 1562–70. doi:10.1128/MCB.20.5.1562-1570.2000

Papp, D., Kovács, T., Billes, V., Varga, M., Tarnóci, A., Hackler, L., Puskás, L.G., Liliom, H., Tárnok, K., Schlett, K., Borsy, A., Pádár, Z., Kovács, A.L., Hegedűs, K., Juhász, G., Komlós, M., Erdős, A., Gulyás, B., Vellai, T., 2016. AUTEN-67, an autophagy-enhancing drug candidate with potent antiaging and neuroprotective effects. Autophagy 12, 273–286. doi:10.1080/15548627.2015.1082023

Parkes, T.L., Elia, A.J., Dickinson, D., Hilliker, A.J., Phillips, J.P., Boulianne, G.L., 1998. Extension of Drosophila lifespan by overexpression of human SOD1 in motorneurons. Nat. Genet. 19, 171–174. doi:10.1038/534

Petiot, A., Ogier-Denis, E., Blommaart, E.F.C., Meijer, A.J., Codogno, P., 2000. Distinct classes of phosphatidylinositol 3'-kinases are involved in signaling pathways that control macroautophagy in HT-29 cells. J. Biol. Chem. 275, 992–998. doi:10.1074/jbc.275.2.992

R Development Core Team, 2016. R: A Language and Environment for Statistical Computing. R Found. Stat. Comput. Vienna Austria 0, {ISBN} 3-900051-07-0. doi:10.1038/sj.hdy.6800737

Schneider, C.A., Rasband, W.S., Eliceiri, K.W., 2012. NIH Image to ImageJ: 25 years of image analysis. Nat. Methods 9, 671–675. doi:10.1038/nmeth.2089

Schonrock, N., Ke, Y.D., Humphreys, D., Staufenbiel, M., Ittner, L.M., Preiss, T., Götz, J., 2010. Neuronal MicroRNA Deregulation in Response to Alzheimer's Disease Amyloid-β. PLoS One 5, e11070. doi:10.1371/journal.pone.0011070

Schu, P. V, Takegawa, K., Fry, M.J., Stack, J.H., Waterfield, M.D., Emr, S.D., 1993. Phosphatidylinositol 3-kinase encoded by yeast VPS34 gene essential for protein sorting. Science (80-.). 260, 88–91.

Seroude, L., Brummel, T., Kapahi, P., Benzer, S., 2002. Spatio-temporal analysis of gene expression during aging in *Drosophila melanogaster*. Aging Cell 1, 47–56. doi:10.1046/j.1474-9728.2002.00007.x

Shen, J., Yu, W.-M., Brotto, M., Scherman, J.A., Guo, C., Stoddard, C., Nosek, T.M., Valdivia, H.H., Qu, C.-K., 2009. Deficiency of MIP/MTMR14 phosphatase induces a muscle disorder by disrupting Ca2+ homeostasis. Nat Cell Biol 11, 769–776.

Takeshige, K., Baba, M., Tsuboi, S., Noda, T., Ohsumi, Y., others, 1992. Autophagy in yeast demonstrated with proteinase-deficient mutants and conditions for its induction. J Cell Biol

119, 301.

Tosch, V., Rohde, H.M., Tronchère, H., Zanoteli, E., Monroy, N., Kretz, C., Dondaine, N., Payrastre, B., Mandel, J.-L., Laporte, J., 2006. A novel PtdIns3P and PtdIns (3, 5) P2 phosphatase with an inactivating variant in centronuclear myopathy. Hum Mol Genet 15, 3098–3106.

Vergne, I., Roberts, E., Elmaoued, R.A., Tosch, V., Delgado, M.A., Proikas-Cezanne, T., Laporte, J., Deretic, V., 2009. Control of autophagy initiation by phosphoinositide 3-phosphatase jumpy. EMBO J. 28, 2244–2258. doi:10.1038/emboj.2009.159

Volinia, S., Dhand, R., Vanhaesebroeck, B., MacDougall, L.K., Stein, R., Zvelebil, M.J., Domin, J., Panaretou, C., Waterfield, M.D., 1995. A human phosphatidylinositol 3-kinase complex related to the yeast Vps34p-Vps15p protein sorting system. EMBO J 14, 3339.

Xiao, C., Mileva-Seitz, V., Seroude, L., Robertson, R.M., 2007. Targeting HSP70 to motoneurons protects locomotor activity from hyperthermia in *Drosophila*. Dev. Neurobiol. 67, 438–455. doi:10.1002/dneu.20344

Yamaguchi, S., Katagiri, S., Sekimizu, K., Natori, S., Homma, K.J., 2005. Involvement of EDTP, an egg-derived tyrosine phosphatase, in the early development of *Drosophila melanogaster*. J. Biochem. 138, 721–728. doi:10.1093/jb/mvi176

Yeh, E., Gustafson, K., Boulianne, G.L., 1995. Green fluorescent protein as a vital marker and reporter of gene expression in Drosophila. Proc Natl Acad Sci U A S 92, 7036–7040.

Zhang, S., Binari, R., Zhou, R., Perrimon, N., 2010. A genomewide RNA interference screen for modifiers of aggregates formation by mutant huntingtin in *Drosophila*. Genetics 184, 1165–1179. doi:10.1534/genetics.109.112516

Zirin, J., Cheng, D., Dhanyasi, N., Cho, J., Dura, J.-M., Vijayraghavan, K., Perrimon, N., 2013. Ecdysone signaling at metamorphosis triggers apoptosis of *Drosophila* abdominal muscles. Dev. Biol. 383, 275–284. doi:10.1016/j.ydbio.2013.08.029

FIGURE LEGENDS

Figure 1. Percent survival at 12 h (% 12-h survival) after anoxia in *EDTP* mutants. (**a**) % 12-h survival in heterozygous *EDTP* mutants (*EDTP^{DJ694}*/+, green) and their sibling controls (*w^{1118}*, grey) to anoxia. Flies at 6-8 days were exposed to 1, 3, or 6 h anoxia. Datasets are presented as scattered dots and Turkey-style boxes and whiskers. Left panel, male flies; right panel, female flies; * $P < 0.05$ and *** $P < 0.001$ from two-way ANOVA with Turkey's test for multiple comparisons. (b) Two-way ANOVA for the analysis of contributing significance of the genotype, anoxia duration and their interaction to the % 12-h survival.

Figure 2. Improved survivorship after prolonged anoxia in heterozygous *EDTP* mutants. Flies were exposed to 0, 1, or 6 h anoxia at 6-8 days. The post-exposure survival curves (days) between heterozygous *EDTP* mutants (*EDTP^{DJ694}*/+, green) and their sibling controls (*w^{1118}*, grey) were plotted. Anoxia durations (in red) are indicated. *P* values are from Log-rank test. Left panels, male flies; right panels, female flies.

Figure 3. Downregulation of *EDTP* in motoneurons improved the survivorship to protein aggregates expressed in the same cells. (**a**) The survival curves of flies expressing Aβ42 (red) and flies with Aβ42 expression and simultaneous knockdown of *EDTP* in the motoneurons (green). (**b**) The survival curves of flies expressing polyQ protein aggregates (red) and flies with polyQ expression and simultaneous knockdown of *EDTP* in the motoneurons (green). Notes: D42-Gal4, motoneuron-specific driver; UAS-Aβ42, a UAS line carrying a gene encoding human beta-amyloid peptide (N = 42); UAS-Httex1-Q103-eGFP, a UAS line for the expression of polyQ protein aggregates (containing in each molecule a polyglutamine tract with 103 tandem repeats); UAS-EDTP-RNAi, an RNAi line (#36917) targeting the exon 2 of *EDTP* gene. *P* values from log-rank tests.

Figure 4. Downregulation of *EDTP* in motoneurons extends lifespan. (**a**)The lifespan in flies with *EDTP* knockdown (D42-Gal4/UAS-EDTP-RNAi (III), green) and their controls (D42-Gal4/+, grey; UAS-EDTP-RNAi (III)/+, blue). (**b**)The lifespan in flies with *EDTP* knockdown (UAS-EDTP-RNAi (II)/+; D42-Gal4/+, green) and their controls (D42-Gal4/+, grey; UAS-EDTP-RNAi (II)/+, blue). Two UAS lines carrying independent RNAi constructs, one in the third chromosome and another in the second chromosome, were used. *** $P < 0.001$ from Log-rank test with "BH" adjustment method for pairwise comparison.

Figure 5. Expression of MTMR14 in the hippocampus and cortex in C57BL/6J and APP/PS1 mice. (**a**) Immunostaining of MTMR14 in the hippocampus in a C57BL/6J mouse. CA3, CA3 pyramidal neurons; DG, dentate gyrus. (**b**) Expression of MTMR14 in the hippocampus and cortex in C57BL/6J and APP/PS1 mice. α-Tubulin, loading control; * $P < 0.05$, student's *t*-test. (c) Expression of MTMR14 in the hippocampus in C57BL/6J mice at different ages (in months). Numbers in the bars indicate the ages of mice.

Figure 6. MTMR14 induction by Aβ42 in primary hippocampal neurons and Neuro2a cells.
(**a**) MTMR14 induction by Aβ42 treatment (0, 5, and 10 μM) for 72 h in the primary
hippocampal neurons of C57BL/6J mice. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ by one-way
ANOVA. (**b**) MTMR14 induction by Aβ42 treatment (0 and 5 μM) for 24 h in Neuro2a cells. **
$P < 0.01$, student's *t*-test.

a

% Survival at 12 h

EDTP^DJ694^/+   *w*^1118^

Male

Female

Anoxia duration (h)

b

Variables	Male	Female
Genotype	$P < 0.001$	$P = 0.383$
Anoxia duration	$P < 0.001$	$P = 0.004$
Genotype × Anoxia duration	$P = 0.002$	$P = 0.245$

Figure 1

Figure 2

Figure 3

Figure 4

a

C57BL/6J

CA3

DG

b

C57BL/6J
8-month

APP/PS1
8-month

Hippocampus

MTMR14

α-Tubulin

Cortex

MTMR14

α-Tubulin

MTMR14 relative level

Hippocampus

Cortex

*

C57BL/6J   APP/PS1

C57BL/6J   APP/PS1

c

C57BL/6J Hippocampus

2   6   9   12   17-mon

MTMR14

α-Tubulin

MTMR14 relative level

1.5

1.0

0.5

0.0

2   6   9   12   17

0   10   20

C57BL/6J age (months)

a C57BL/6J primary culture neurons

b Neuro2a cells

Figure 6