Full Title: High affinity Na$^+$ transport by wheat HKT1;5 is blocked by K$^+$

Running title: HKT1;5 proteins catalyse dual affinity Na$^+$ transporter in wheat

Authors: Bo Xu[1,2], Maria Hrmova[2] and Matthew Gilliham[1,2*]

[1]Australian Research Council Centre of Excellence in Plant Energy Biology,

University of Adelaide, Waite Research Precinct, Glen Osmond, South Australia

5064, Australia

[2]School of Agriculture, Food and Wine, and Waite Research Institute, University of

Adelaide, Waite Research Precinct, Glen Osmond, South Australia 5064, Australia

*Corresponding author: e-mail: matthew.gilliham@adelaide.edu.au

Abstract

The wheat sodium transporters TmHKT1;5-A and TaHKT1;5-D are encoded by

genes underlying major shoot Na^+ exclusion loci *Nax2* and *Kna1* from *Triticum*

monococcum (Tm) and *Triticum aestivum* (Ta), respectively. In contrast to HKT2

transporters that have been shown to exhibit high affinity K^+-dependent Na^+ transport,

HKT1 proteins have, with one exception, only been shown to catalyse low affinity

Na^+ transport and no K^+ transport. Here, using heterologous expression in *Xenopus*

laevis oocytes we show that both TmHKT1;5-A and TaHKT1;5-D encode dual (high

and low) affinity Na^+-transporters with the high-affinity component being abolished

when external K^+ is in excess of external Na^+. The low-affinity component for Na^+

transport of TmHKT1;5-A had a lower K_m than that of TaHKT1;5-D even when

blocked by external K^+. We use 3-D structural modelling to explain how K^+ block

may occur and propose potential physiological consequences of K^+ block. The

transport properties and localisation of wheat HKT1;5 proteins are well suited for

their role in a 'gatekeeper' process that secure shoot Na^+-exclusion and underpin

recent advances for improving crop plant salt tolerance.

Keywords: salinity; High-affinity K Transporters; HKT; wheat; membrane transport;

crop; 3-D structural modelling

Introduction

Crops suffer reduced growth and productivity under salinity stress. Salt, when it builds up to high concentrations in the growing medium (i.e. soil solution) imposes an osmotic limitation on water uptake, interferes with optimal nutrient homeostasis, and leads to the build up of leaf cellular sodium (Na^+) concentrations, which causes an ionic toxicity that limits photosynthesis and carbon assimilation in plants (Munns and Gilliham, 2015).

A major genetic mechanism contributing towards salinity tolerance of most crops including the cereals rice (*Oryza sativa*) and wheat (*Triticum aestivum* and *Triticum monococcum*) is Na^+ exclusion from leaves, with the High-affinity potassium (K^+) transporter (HKT) protein family having a major role in this trait (Ren *et al.* 2005; James *et al.* 2006; Munns *et al.* 2012; Munns and Gilliham, 2015; Campbell *et al.* 2017; Xu *et al.* 2018). This family of proteins are represented widely across the plant kingdom, and more broadly are members of the high affinity K^+/Na^+ transporting Ktr/TrK/HKT superfamily of proteins that are present in bacteria (Ktr and TrK), fungi (TrK) and plants (HKT) (Sentenac and Bonneaud 1992; Corratgé-Faillie *et al.* 2010).

These plant HKTs are divided into 2 clades: 1) class 1 HKT proteins (HKT1;x) are mostly Na^+-selective transporters; 2) whereas class 2 HKTs (HKT2;y) mostly function as K^+-Na^+ symporters and so far have been only identified in cereal monocots (Asins *et al.* 2013; Waters *et al.* 2013). This definition has been challenged on occasion, such as for OsHKT2;4 following its characterization in *Xenopus laevis* oocytes, where there are conflicting reports on its permeability to Ca^{2+}, Mg^{2+} and NH_4^+ in addition to K^+ and Na^+ (Lan *et al.* 2010; Horie *et al.* 2011; Sassi *et al.* 2012). Furthermore, there are two reports of K^+ permeability in the HKT1 clade . Firstly, Arabidopsis AtHKT1 could complement K^+ transport deficient *E. coli* although no K^+ permeability could be found when it was expressed in *X. laevis* oocytes in the same study (Uozumi *et al.*, 2000). Secondly, *Eucalyptus camaldulensis* EcHKT1;1 and EcHKT1;2, which appear to be the exception for the HKT1 proteins characterized so far in that as they transport both Na^+ and K^+ when expressed in *X. laevis* oocytes (Liu *et al.* 2001). Other HKT1 transporters have no reported K^+ permeability (Uozumi *et al.* 2000; Mäser *et al.* 2002; Platten *et al.* 2006; Jabnoune *et al.* 2009; Cotsaftis *et al.* 2012; Munns *et al.* 2012; Waters *et al.* 2013; Byrt *et al.* 2014; Campbell *et al.* 2017; Henderson et al., 2018).

Whilst K^+ transport is not a common feature of HKT1 transporters, several have

shown the property of K^+-regulated Na^+ transport (Ben Amar *et al.* 2013; Almeida *et*

al. 2014a; Almeida *et al.* 2014b; Byrt *et al.* 2014). For instance, the presence of 10

mM external K^+ ($[K^+]_{ext}$) reduced the inward Na^+ current of *X. laevis* oocytes

expressing *HKT1;5* from *T. aestivum* (*TaHKT1;5-D*) at -140 mV (Byrt *et al.* 2014),

and the outward Na^+ current of that expressing *HKT1;2* from *Solanum lycopersicum*

(*SlHKT1;2*) (Almeida *et al.* 2014a). Interestingly, $[K^+]_{ext}$ stimulated both inward and

outward Na^+ transport of *X. laevis* oocytes expressing two *HKT1;4* alleles from

Triticum turgidum L. subsp. *durum* when assayed with solutions of very low ionic

strength (Ben Amar *et al.* 2013).

Ktr/TrK/HKT proteins are predicted to consist of 8 transmembrane domains that fold

into 4 pseudo-tetramers around a central pore (Mäser *et al.*, 2002; Cotsaftis *et al.*,

2012; Xu *et al.*, 2018). The selectivity filter lining the pore of Ktr and TrK and class 2

HKT proteins is predicted to be composed of four glycine residues. In class 1 HKTs

the first glycine within the predicted selectivity filter is substituted with a serine

(Mäser *et al.*, 2002); the lack of K^+ permeability in this clade has been linked to the

presence of this glycine (Uozumi *et al.*, 2000; Mäser *et al.*, 2002; Platten *et al.*, 2006;

Cotsaftis *et al.*, 2012; Waters *et al.*, 2013); but the exact mechanism underlying this

block is to be determined.

Here, we further characterise HKT1;5 proteins underlying two major salt tolerance

associated loci, *TmHKT1;5-A* (*Nax2*) and *TaHKT1;5-D* (*Kna1)* from *T. monococcum*

and T. aestivum respectively. We find that both TmHKT1;5-A and TaHKT1;5-D

encode dual affinity Na^+-transporters and that their dual affinity Na^+ transport can be

blocked by raising external K^+ concentration, and we propose the residues within the

selectivity filter that cause this block. This is the first report that Na^+ transport has two

affinities in the HKT1 clade and demonstrates that dual affinity transport is a common

property of both clades of the HKT family.

Results

K$^+$ sensitivity of TmHTK1;5-A and TaHKT1;5-D

Previously, it was shown that inward Na$^+$ transport for both TmHKT1;5-A and

TaHKT1;5-D was inhibited by external K$^+$ (Munns *et al.* 2012; Byrt *et al.* 2014).

Here, we further compared the K$^+$-sensitivity of between TmHKT1;5-A and

TaHKT1;5-D (Fig. 1). An increase in [K$^+$]$_{ext}$ to 30 mM reduced the channel

conductance of TmHKT1;5-A at -140 mV by approximately 75% in a 1 mM Na$^+$

([Na$^+$]$_{ext}$) bath solution; similarly this amount of K$^+$ inhibited that of TaHKT1;5-D up

to 95 % (Fig. 1A, B and Fig. 2). However, such Na$^+$-inhibition by K$^+$ was gradually

decreased by increasing [Na$^+$]$_{ext}$ (Fig 1 and 2). For instance, 10 mM [K$^+$]$_{ext}$ suppressed

the channel conductance of TmHKT1;5-A by 43.5% in 1 mM [Na$^+$]$_{ext}$ and by 16.5%

in 10 mM [Na$^+$]$_{ext}$; it similarly reduced the TaHKT1;5-D conductance by 57% in 1

mM [Na$^+$]$_{ext}$ and by 23% in 10 mM [Na$^+$]$_{ext}$ (Fig. 1 and 2). When [Na$^+$]$_{ext}$ was

increased to 30 mM, the channel conductance of TmHKT1;5-A was insensitive to

[K$^+$]$_{ext}$, whereas that of TaHKT1;5-D was still inhibited (Fig. 2).

The increase in [K$^+$]$_{ext}$ from 0 to 10 mM did not significantly change the reversal

potential of TmHKT1;5-A or TaHKT1;5-D; all reversal potentials were close to the

theoretical equilibrium potential for Na$^+$ under all conditions (Fig. 3A,B). Moreover,

the observed shift in reversal potential was close to the theoretical Nernst shift for Na$^+$

regardless of the presence or absence of 10 mM [K$^+$]$_{ext}$ (Fig. 3). This suggests that

neither of protein were permeable to K$^+$.

Dual affinity transport of Na$^+$ by TmHTK1;5-A and TaHKT1;5-D

The affinity for Na$^+$ transport of TmHKT1;5-A was previously reported to be ~4-fold

higher than the Na$^+$-transport affinity of TaHKT1;5-D (Munns *et al.* 2012; Byrt *et al.*

2014; Xu *et al.* 2018), which we confirm here (Fig. 4A,B). On closer examination, we

found a property that has not been described before. When lower concentrations of

[Na$^+$]$_{ext}$ were present it was clear that the concentration dependence of inward Na$^+$

currents could be fitted by two components, for both TmHKT1;5-A and TaHKT1;5-D

(Fig. 4). Whilst both components were saturable, the concentration at which both

phases saturated were greater for TaHKT1;5-D compared to TmHKT1;5-A (Fig.

4A,B). For TmHKT1;5-A, the high affinity component of Na$^+$ transport had a K$_m$ of

34 µM ± 26 µM when [Na$^+$]$_{ext}$ was below 0.1 mM (K$_{m\,(<0.1\,Na)}$), whereas at

concentrations of Na^+ above 0.1 mM the K_m was 1.04 ± 0.17 mM ($K_{m\,(>0.1\,Na)}$) (Fig.

4A). The K_m for the higher affinity component of inward Na^+ transport for

TaHKT1;5-D was 29 µM ± 4 µM when $[Na^+]_{ext}$ was lower than 0.5 mM ($K_{m\,(<0.5\,Na)}$),

whilst the lower affinity phase had a K_m of 4.32 ± 0.5 mM when $[Na^+]_{ext}$ was higher

than 0.5 mM ($K_{m\,(>0.5\,Na)}$) (Fig. 4B).

We examined the extent of K^+-inhibition of both the lower and higher affinity

components of the inward Na^+ currents when $[K^+]_{ext}$ was 10 mM. The (normalised)

inward Na^+ current catalysed by both proteins was significantly suppressed by 10 mM

$[K^+]_{ext}$, resulting in negligible inward Na^+ transport by TmHKT1;5-A in presence of

$[Na^+]_{ext}$ below 0.1 mM, or by TaHKT1;5-D in presence of $[Na^+]_{ext}$ below 0.5 mM

(Fig. 4C and D). In the presence of 10 mM $[K^+]_{ext}$ it was now possible to fit the

concentration dependence of inward current through TmHKT1;5-A and TaHKT1;5-D

with one component. The K_m of both TmHKT1;5-A and TaHKT1;5-D were decreased

approximately by 3-fold by 10 mM $[K^+]_{ext}$ (Fig. 4E and F).

Structural modelling of TaHKT1;5-D

Three-dimensional structural models of TaHKT1;5-D were generated to explore why

K^+ does not traverse the pore but instead blocks the Na^+ currents, using the KtrB K^+

transporter from *B. subtilis* as a template (Vieira-Pires *et al.*, 2013; Xu *et al.*, 2018).

The KtrB K^+ protein was crystallised in the presence of KCl, hence the original

structure contained a K^+ ion in the selectivity filter region. This ion was substituted by

Na^+ during modelling of TaHKT1;5-D as it transports Na^+ but not K^+ (Xu *et el.*,

2018). Ramachandran analysis indicated that the template and TaHKT1;5-D models

generated in complex with Na^+, and K^+ had satisfactory stereo-chemical quality;

Ramachandran plots showed only two residues positioned in disallowed regions

(0.5% of all residues, except G and P residues). Average G-factors (measures of

correctness of dihedral angles and main-chain covalent forces of protein molecules)

calculated by PROCHECK of the template and TaHKT1;5-D models with K^+ and Na^+

were 0.06. -0.18 and -0.13, respectively. The ProSa 2003 analysis of z-scores (-8.4; -

6.0 and -5.8) indicated that template and modelled structures with K^+ and Na^+ had

acceptable conformational energies.

The TaHKT1;5-D model showed that the Na^+ ion is penta-hedrally coordinated with

V76, S77, S78, N231, C232 and H351 (all carbonyl oxygens), that lie near the

selectivity filter residues S78, G233, G353 and G457 (Fig. 5A). Notably, the carbonyl

oxygen one of the selectivity filter residues (S78) directly participates in Na^+ binding

(Fig. 5A). Conversely, the K^+ ion is hexa-hedrally coordinated with V76, N231,

C232, H351, N455 and V456, that also closely neighbour the selectivity filter residues

(Fig. 5B); here none of the selectivity filter residues participate in direct binding of

K^+. The ionic distances of residues neighbouring Na^+ and K^+ ions are within similar

ranges: for Na^+ they are between 2.3 and 2.4 Å; for K^+ they are between 2.6 and 2.9 Å

(Fig. 5, right panels).

Discussion

Oocytes expressing *TmHKT1;5-A* or *TaHKT1;5-D* have been previous reported to

undergo no obvious change in their reversal potential regardless of the presence or

absence of K^+ in bath solution. Interestingly, both transporters had differential

sensitivity to external $[K^+]_{ext}$. Whilst K^+ block occurs for TmHKT1;5-A catalysing

inward transport it was not evident at $[Na^+]_{ext}$ above 10 mM, whereas this blockage

did happen for TaHKT1;5-D up to 30 mM $[Na^+]_{ext}$ (Fig. 2) (Munns *et al.* 2012; Byrt

et al. 2014). This differential inhibition effect of K^+ upon Na^+ transport at high

$[Na^+]_{ext}$ by TaHKT1;5-D may result from a range of factors, that govern the rates of

Na^+ and K^+ ion binding and subsequent Na^+ transport. A variety of factors could

control the transport of Na^+ and preliminary displacement of competing K^+ with a

consequent energetic cost, before Na^+ passes bare or less-well hydrated than

competing K^+. Examples include free-energy variations between ions in binding sites

relative to the corresponding quantity in bulk water, differences in selectivity filters

solvent exposures (or dielectric constants), overall pore rigidity/stiffness (Dudev and

Lim 2010), structural changes within funnels and selectivity filters due to residue re-

orientation, differences in the hydration (coordination) numbers of Na^+ and K^+ (Na^+

and K^+ bind six and seven water molecules, respectively (Rowley and Roux 2012)

and the alterations on the rates of Na^+ or K^+ de-solvation (Degrève *et al.* 1996). It is

therefore plausible to expect that both ions would compete for binding sites within

HKT funnel regions, yielding ion-protein binding sites with differential binding

strengths.

Here, we refine the previous structural models of HKT1;5 to show that they contain a

pore that allows the passage of Na^+ but not K^+ (Cotsaftis *et al.*, 2012; Waters *et al.*,

2013; Xu *et al.*, 2018). This is consistent with the functional observations in this and

previous studies (Munns *et al.* 2012; Byrt *et al.* 2014). Previously it was suggested

that the Gly-Gly-Gly-Gly motif is important in coordinating the K^+ ion in Ktr/TrK

transporters and in conferring dual Na^+-K^+ transport in HKT2 transporters (Durrell

and Guy, 1999; Mäser *et al.* 2002). Here our modelling suggests that the serine residue

within the Ser-Gly-Gly-Gly selectivity motif directly interacts and binds with Na^+ (Fig

5B). Furthermore, our model suggests, despite ionic distances being shorter for Na^+

than those for K^+, that K^+ could be bound more strongly due to a higher coordination

pattern (Fig. 5A and 5B-right panels). Obviously, during binding, K^+ takes an

advantage of its larger ionic radius: 152 picometer (pm), compared to that of Na^+ (116

pm); these parameters reflects empirical atomic radii of K (220 pm) and Na (180 pm).

We assume that under the high concentrations of K^+, this ion may outcompete Na^+ and

would become bound in the selectivity filter, thus effectively blocking Na^+ transport.

This is illustrated by calculating permeation channels in both complexes, using the

Mole Voronoi algorithm, that predicts permeation trajectories and identifies path

bottlenecks. From the calculated permeation paths, it could be deduced that the K^+ ion

effectively blocks the permeation channel through the major gating protein pore,

contrary to Na^+. It is also obvious that Na^+ is likely to enter and exit the permeation

trajectory from several points on both sides of the transporter, but always by-passes

the selectivity filter constriction. Both ions arrive at the funnel of HKT proteins in

solvated (hydrated) forms. While the hydration number for Na^+ is 6, that for K^+ is 7 at

ambient temperature (Rowley and Roux 2012; Ma, 2016). The significance of the

seven-fold water coordination for K^+ is that this solvated complex may form a

stronger hydrogen-bond interaction pattern (in funnels of HKT proteins), and would

be de-solvated with a higher energy input and thus slower) that the Na^+ solvated

complex (Degrève et al., 1996). It is also plausible to expect that both solvated ion

complexes would compete for binding sites within HKT funnel regions with different

binding strengths. All in all, this would contribute to the blockage of the pore entry of

HKT transporters.

Potassium permeability for HKT1 transporters has only been reported once for the

proteins that have been characterised in *X. laevis* oocytes. The transporters EcHKT1;1

and EcHKT1;2 allow permeation of both K^+ and Na^+ (Liu *et al.*, 2001). Interestingly,

when 3D models of the Na^+ selective OsHKT1;5 and K^+ permeable EcHKT1;2 were

constructed and compared the pore size was predicted to be 0.2 Å larger in EcHKT1;2

(than OsHKT1;5 and TaHKT1;5-D), which would more than account for the weaker

interactions with K^+ and allow this larger ion to pass through the pore (Waters *et al.*,

2013). Without 3D modelling of each and every transporter in the HKT family to

show how the selectivity filter forms in the context of the rest of the protein it would

not be possible to predict which other HKT1s may allow passage of K^+. Predictions

based on sequence alone are insufficient to predict how the two sets of proteins

evolved their individual Na^+ and K^+ transport characteristics. An extension of the

model to include other HKT proteins, a greater survey of structure-function

relationships and mutations of key residues to predict functionally relevant residues

and evolutionary relationships will be the focus of further study.

Two affinity ranges for Na^+ transport have been detected in the HKT family

previously, but only for HKT2 transporters. OsHKT2;1, OsHKT2;2, OsHKT2;4 and

HvHKT2;1(Schachtman and Schroeder 1994; Rubio *et al.* 1995; Jabnoune *et al.* 2009;

Yao *et al.* 2010; Horie *et al.* 2011; Sassi *et al.* 2012). For instance, for OsHKT2;2,

when K^+ is present, the transport of Na^+ is K^+-dependent and displays a high affinity

($K_m = 0.077$ mM), whereas showing a lower affinity when absence of K^+ ($K_m= 16$

mM) (Rubio *et al.* 1995; Yao *et al.* 2010). OsHKT2;1 also acquires two phases of

Na^+-transport affinity, whose K_m for Na^+ is respectively 9.5 μM at $[Na^+]_{ext}$ below 1

mM and is 2.2 mM at $[Na^+]_{ext} > 1$ mM (Jabnoune *et al.* 2009). The K^+ transport

catalysed by HvHKT2;1 is affected by both $[Na^+]_{ext}$ and $[K^+]_{ext,}$ such as a decrease in

its K^+ transport affinity from 30 μM down to 3.5 mM by $[Na^+]_{ext}$ increased from 0.5

mM to 30 mM and a greatly reduced channel conductivity for K^+ by increasing $[K^+]_{ext}$

above ~3 mM (Mian *et al.* 2011). This is the first time that dual affinity characteristics

have been reported for the HKT1;*x* family (Fig. 4). Considering the considerably

homology between both clades (58% similarity/41% identity between TaHKT1;5 and

TaHKT2;1) it is perhaps not surprising that both clades can confer dual affinity

transport. How such ability is conferred structurally is still to be determined.

In summary, whilst the high-affinity component was similarly inhibited by 10 mM

$[K^+]_{ext}$ for both TaHKT1;5-D and TmHKT1;5-A, the low affinity component of Na^+

transport for TmHKT1;5-A was less inhibited by external K than TaHKT1;5-D (Fig.

4). Furthermore, both wheat HKT1;5 proteins facilitate Na^+ uptake at high affinity;

TmHKT1;5-A facilitates Na^+ uptake with a higher affinity than TaHKT1;5-D

regardless the absence or presence of $[K^+]_{ext,}$ (Fig. 4) (Munns *et al.* 2012; Byrt *et al.*

2014). TmHKT1;5-A has been shown to be more effective in conferring shoot Na^+

exclusion to wheat compared to TaHKT1;5-D (James *et al.*, 2012). We propose that

the higher affinity (with half-maximal activity nearer 1 mM compared to 4 mM), and

a lower sensitivity to $[K^+]_{ext}$ are both characteristics that can help confer better shoot

Na^+ exclusion, which is a property that can lead to greater salt tolerance in the field

for wheat (James *et al.* 2011; James *et al.* 2012; Munns *et al.* 2012).

Materials and Methods

Brief methods for cloning of *TmHKT1;5-A* and *TaHKT1;5-D* as well as its functional

characterisation in heterologous expression systems were described Munns et al.

(Munns *et al.* 2012) and Byrt et al. (Byrt *et al.* 2014). Further details are included

here.

Two-electrode voltage clamp recording in *X. laevis* oocytes

Oocyte recording followed the methods as described in Munns et al. (Munns *et al.*

2012)and Byrt et al. (Byrt *et al.* 2014). Briefly, 46 nl/23 ng of cRNA or equal

volumes of RNA-free water were injected into oocytes, followed by an incubation for

48 h before recording. Membrane currents were recorded in the HMg solution (6 mM

$MgCl_2$, 1.8 mM $CaCl_2$, 10 mM MES and pH 6.5 adjusted with a TRIS base) $\pm$ Na^+

glutamate and/or K^+ glutamate as indicated. All solution osmolarities were adjusted

using mannitol at 220-240 mOsmol kg^{-1}.

Construction of 3D models of TaHKT1;5D in complex with Na^+ and K^+ ions

The most suitable template for TaHKT1;5D, the KtrB K^+ transporter from *B. subtilis*

(Protein Data Bank accession 4J7C, chain I) (Vieira-Pires et al., 2013), was identified

as previously described (Xu et al., 2017). The KtrB K^+ protein was crystallised in the

presence of KCl, hence the structure contains a K^+ ion in the selectivity filter region.

This ion was substituted by Na^+ during modelling of TaHKT1;5D that transport Na^+ at

a greater rate than K^+ (Xu et el., 2017). 3D structural models in complex with Na^+, and

K^+ were generated using Modeller 9v16 (Sali and Blundell, 1993) on a Linux station

running the Fedora 12 operating system, as previously described (Cotsaftis et al.,
2012, Waters et al., 2013, Xu et al., 2017). During modelling, attention was paid to
ionic radii of Na$^+$, and K$^+$, whose topology parameters were taken from CHARMM
(Brooks et al., 2009). In each case, a total of 100 models were generated that were
scored by Modeller using the modeller objective function (Shen and Sali, 2006), the
discrete optimised protein energy function (Eswar et al., 2008), PROCHECK
(Laskowski et al., 1993), ProSa 2003 (Sippl, 1993) and FoldX (Schymkowitz et al.,
2005). Best scoring models constructed in Modeller 9v19 were further subjected to
energy minimisation (knowledge-based Yasara2 forcefield with parameters for bond
distances, planarity of peptide bonds, bond angles, Coulomb terms, dihedral angles,
and van der Waals forces) (Krieger et al., 2004) combined with the particle-mesh-
Ewald (PME) energy function for long range electrostatics (cutoff 8.0 Å), to obtain
smoothed electrostatic potentials. Structural images were generated with PyMOL
Molecular Graphics System V1.8.2.0 (Schrődinger LLC, Portland, OR, USA).
323

324 **Funding**
325 This work was supported by funding to M.G. from the Grains Research and
326 Development Corporation (UA00145, M.G.), and Australian Research Council
327 through the Centre of Excellence (CE1400008) and Future Fellowship
328 (FT130100709) schemes.

329
330

References

Almeida, P, de Boer, G-J, de Boer, AH (2014a) Differences in shoot Na^+ accumulation between two tomato species are due to differences in ion affinity of HKT1;2. *Journal of Plant Physiology* **171**, 438-447.

Almeida, P, Feron, R, de Boer, G-J, de Boer, AH (2014b) Role of Na^+, K^+, Cl^-, proline and sucrose concentrations in determining salinity tolerance and their correlation with the expression of multiple genes in tomato. *AoB Plants* **6**, plu039.

Asins, MJ, Villalta, I, Aly, MM, Olias, R, Álvarez De Morales, P, Huertas, R, Li, J, JAIME-PÉREZ, N, Haro, R, Raga, V (2013) Two closely linked tomato HKT coding genes are positional candidates for the major tomato QTL involved in Na^+/K^+ homeostasis. *Plant, Cell & Environment* **36**, 1171-1191.

Ben Amar, S, Brini, F, Sentenac, H, Masmoudi, K, Véry, A-A (2013) Functional characterization in Xenopus oocytes of Na^+ transport systems from durum wheat reveals diversity among two HKT1; 4 transporters. *Journal of experimental botany* **65**, 213-222.

Byrt, CS, Xu, B, Krishnan, M, Lightfoot, DJ, Athman, A, Jacobs, AK, Watson-Haigh, NS, Plett, D, Munns, R, Tester, M (2014) The Na^+ transporter, TaHKT1;5-D, limits shoot Na^+ accumulation in bread wheat. *Plant Journal* **80**, 516-526.

Campbell, MT, Bandillo, N, Al Shiblawi, FRA, Sharma, S, Liu, K, Du, Q, Schmitz, AJ, Zhang, C, Véry, A-A, Lorenz, AJ (2017) Allelic variants of OsHKT1; 1 underlie the divergence between indica and japonica subspecies of rice (*Oryza sativa*) for root sodium content. *PLoS genetics* **13**, e1006823.

Corratgé-Faillie, C, Jabnoune, M, Zimmermann, S, Véry, A-A, Fizames, C, Sentenac, H (2010) Potassium and sodium transport in non-animal cells: the Trk/Ktr/HKT transporter family. *Cellular and Molecular Life Sciences* **67**, 2511-2532.

Cotsaftis, O, Plett, D, Shirley, N, Tester, M, Hrmova, M (2012) A two-staged model of Na^+ exclusion in rice explained by 3D modeling of HKT transporters and alternative splicing. *PLoS One* **7**, e39865.

Degrève, L, Vechi, SM, Junior, CQ (1996) The hydration structure of the Na^+ and K^+ ions and the selectivity of their ionic channels. *Biochimica et Biophysica Acta (BBA)-Bioenergetics* **1274**, 149-156.

Dudev, T, Lim, C (2010) Factors governing the Na^+ vs K^+ selectivity in sodium ion channels. *Journal of the American Chemical Society* **132**, 2321-2332.

Henderson SW, Dunlevy JD, Wu Y, Blackmore DH, Walker RR, Edwards EJ, Gilliham M, Walker AR (2018) Functional differences in transport properties of natural HKT1;1 variants influence shoot Na^+ exclusion in grapevine rootstocks. *New Phytologist* **217**, 113-1127.

Horie, T, Brodsky, DE, Costa, A, Kaneko, T, Schiavo, FL, Katsuhara, M, Schroeder, JI (2011) K^+ transport by the OsHKT2; 4 transporter from rice with atypical Na^+ transport properties and competition in permeation of K^+ over Mg^{2+} and Ca^{2+} ions. *Plant Physiology* **156**, 1493-1507.

Jabnoune, M, Espeout, S, Mieulet, D, Fizames, C, Verdeil, J-L, Conéjéro, G, Rodríguez-Navarro, A, Sentenac, H, Guiderdoni, E, Abdelly, C (2009) Diversity in expression patterns and functional properties in the rice HKT transporter family. *Plant Physiology* **150**, 1955-1971.

James, RA, Blake, C, Byrt, CS, Munns, R (2011) Major genes for Na^+ exclusion, *Nax1* and *Nax2* (wheat *HKT1; 4* and *HKT1; 5*), decrease Na+ accumulation in

380   bread wheat leaves under saline and waterlogged conditions. *Journal of*
381   *experimental botany* **62**, 2939-2947.
382   James, RA, Blake, C, Zwart, AB, Hare, RA, Rathjen, AJ, Munns, R (2012) Impact of
383   ancestral wheat sodium exclusion genes *Nax1* and *Nax2* on grain yield of
384   durum wheat on saline soils. *Functional Plant Biology* **39**, 609-618.
385   James, RA, Davenport, RJ, Munns, R (2006) Physiological characterization of two
386   genes for Na^+ exclusion in durum wheat, *Nax1* and *Nax2*. *Plant Physiology*
387   **142**, 1537-1547.
388   Lan, W-Z, Wang, W, Wang, S-M, Li, L-G, Buchanan, BB, Lin, H-X, Gao, J-P, Luan,
389   S (2010) A rice high-affinity potassium transporter (HKT) conceals a calcium-
390   permeable cation channel. *Proceedings of the National Academy of Sciences,*
391   *U.S.A.* **107**, 7089-7094.
392   Liu, W, Fairbairn, DJ, Reid, RJ, Schachtman, DP (2001) Characterization of two
393   HKT1 homologues from *Eucalyptus camaldulensis* that display intrinsic
394   osmosensing capability. *Plant Physiol.* **127**, 283-294.
395   Mäser, P, Hosoo, Y, Goshima, S, Horie, T, Eckelman, B, Yamada, K, Yoshida, K,
396   Bakker, EP, Shinmyo, A, Oiki, S (2002) Glycine residues in potassium
397   channel-like selectivity filters determine potassium selectivity in four-loop-
398   per-subunit HKT transporters from plants. *Proceedings of the National*
399   *Academy of Sciences, U.S.A.* **99**, 6428-6433.
400   Mian, A, Oomen, RJ, Isayenkov, S, Sentenac, H, Maathuis, FJ, Véry, AA (2011)
401   Over-expression of an Na^+-and K^+-permeable HKT transporter in barley
402   improves salt tolerance. *Plant Journal* **68**, 468-479.
403   Munns, R, James, RA, Xu, B, Athman, A, Conn, SJ, Jordans, C, Byrt, CS, Hare, RA,
404   Tyerman, SD, Tester, M, Plett, D, Gilliham, M (2012) Wheat grain yield on
405   saline soils is improved by an ancestral Na^+ transporter gene. *Nature*
406   *Biotechnology* **30**, 360-364.
407   Munns, R, Gilliham, M (2015) Salinity tolerance of crops - What is the cost? *New*
408   *Phytologist* **208**, 668-673.
409   Platten, JD, Cotsaftis, O, Berthomieu, P, Bohnert, H, Davenport, RJ, Fairbairn, DJ,
410   Horie, T, Leigh, RA, Lin, H-X, Luan, S (2006) Nomenclature for HKT
411   transporters, key determinants of plant salinity tolerance. *Trends in Plant*
412   *Sciences* **11**, 372-374.
413   Ren, Z-H, Gao, J-P, Li, L-G, Cai, X-L, Huang, W, Chao, D-Y, Zhu, M-Z, Wang, Z-Y,
414   Luan, S, Lin, H-X (2005) A rice quantitative trait locus for salt tolerance
415   encodes a sodium transporter. *Nature Genetics* **37**, 1141-1146.
416   Rowley, CN, Roux, Bt (2012) The solvation structure of Na^+ and K^+ in liquid water
417   determined from high level ab initio molecular dynamics simulations. *Journal*
418   *of chemical theory and computation* **8**, 3526-3535.
419   Rubio, F, Gassmann, W, Schroeder, JI (1995) Sodium-driven potassium uptake by the
420   plant potassium transporter HKT1 and mutations conferring salt tolerance.
421   *Science* **270**, 1660.
422   Sassi, A, Mieulet, D, Khan, I, Moreau, B, Gaillard, I, Sentenac, H, Véry, A-A (2012)
423   The rice monovalent cation transporter OsHKT2; 4: revisited ionic selectivity.
424   *Plant Physiology* **160**, 498-510.
425   Schachtman, DP, Schroeder, JI (1994) Structure and transport mechanism of a high-
426   affinity potassium uptake transporter from higher plants. *Nature* **370**, 655-658.
427   Sentenac, H, Bonneaud, N (1992) Cloning and expression in yeast of a plant
428   potassium ion transport system. *Science* **256**, 663.

Uozumi, N, Kim, EJ, Rubio, F, Yamaguchi, T, Muto, S, Tsuboi, A, Bakker, EP,
Nakamura, T, Schroeder, JI (2000) The Arabidopsis HKT1 gene homolog
mediates inward Na$^+$ currents in *Xenopus laevis* oocytes and Na$^+$ uptake in
Saccharomyces cerevisiae. *Plant Physiology* **122**, 1249-1260.

Waters, S, Gilliham, M, Hrmova, M (2013) Plant high-affinity potassium (HKT)
transporters involved in salinity tolerance: structural insights to probe
differences in ion selectivity. *International Journal of Molecular Sciences* **14**,
7660-7680.

Xu, B, Waters, S, Byrt, CS, Plett, D, Tyerman, SD, Tester, M, Munns, R, Hrmova, M,
Gilliham, M (2018) Structural variations in wheat HKT1;5 underpin
differences in Na$^+$ transport capacity. *Cellular and Molecular Life Sciences* 1-
12.

Yao, X, Horie, T, Xue, S, Leung, H-Y, Katsuhara, M, Brodsky, DE, Wu, Y,
Schroeder, JI (2010) Differential sodium and potassium transport selectivities
of the rice OsHKT2;1 and OsHKT2;2 transporters in plant cells. *Plant*
Physiology **152**, 341-355.

 Figures

Figure 1. K$^+$-sensitivity of Na$^+$ currents carried by TmHKT1;5-A and

TaHKT1;5-D. A-D, current-voltage (I/V) curves of Na$^+$ currents recorded from *X.*

laevis oocytes expressing *TmHKT1*;5-*A* (A, C) and *TaHKT1*;5-*D* (B, D) in 1 mM Na$^+$

plus 0-30 mM K$^+$ (A,B), or 0-30 mM Na$^+$ solutions with or without 10 mM K$^+$ (C,D).

Mean ± S.E.M, n = 5-6.

Figure 2. Channel conductance of TmHKT1;5-A and TaHKT1;5-D. The channel

conductance of TmHKT1;5-A and TaHKT1;5-D was calculated from Figure 1, based

on the slope of curve between -140 mV and -120 mV, statistical difference was

determined by *Students' t-test*, *P < 0.05* and ** *P < 0.01*.

Figure 3. Shift in reversal potential of TmHKT1;5-A and TaHKT1;5-D in

response to changes in [Na$^+$]$_{ext}$ and [K$^+$]$_{ext}$.

A, Reversal potential of currents in *X. laevis* oocytes expressing *TmHKT1*;5-*A* and

TaHKT1;5-*D* derived from Figure. 1. B, Reversal potential of Na$^+$ currents plotted

against theoretical Nernst potential derived from internal concentrations measured in

Munns et al. (2012) and Byrt et al., (2014) at the used.

Nernst shift in oocytes expressing *TmHKT1*;5-*A* and *TaHKT1*;5-*D* from 1 mM to 10

mM Na$^+$ ([1-10]$_{ext}$ Na$^+$) and 10 mM to 30 mM Na$^+$ ([10-30]$_{ext}$ Na$^+$) with or without 10

mM [K$^+$]$_{ext}$, ΔExp.$_{Na}$ = experimentally measured shift in reversal potential for Na$^+$, the

theortectical equilibrium (or Nernst) potential for Na$^+$ was 57.7 mV and 26.9 mV,

respectively for [1-10]$_{ext}$ Na$^+$ and [10-30]$_{ext}$ Na$^+$, refers to Munns et al. (2012).

Figure 4. Transport affinity of TmHKT1;5-A and TaHKT1;5-D in *X. laevis*

oocytes.

A-B, dual Na$^+$-transport affinity of TmHKT1;5-A and TaHKT1;5-D in *X. laevis*

oocytes, plotted as normalised currents at -140 mV as shown against a series of Na$^+$

glutamate solutions. C-D, high affinity phase of TmHKT1;5-A and TaHKT1;5-D in

absence and presence of 10 mM [K$^+$]$_{ext}$. Data represented in A, B and C as mean ±

S.E.M (n = 5 for A, n = 4 for B, n = 7 for C). E-F, Na$^+$-transport affinity of

TmHKT1;5-A and TaHKT1;5-D in *X. laevis* oocytes, plotted as normalised currents

at -140 mV as shown against a series of Na$^+$ glutamate solutions with an additional 10

mM K$^+$ glutamate.

Figure 5. Molecular models of TaHKT1;5D in complex with Na$^+$ (A) and K$^+$ (B) ions.

(A-B; left panels) Cartoon representations of TaHKT1;5-D illustrate their overall protein folds with cylindrical α-helices, and permeation channels (black mesh). Na$^+$ (A) and K$^+$ (B) ions are shown as purple spheres located within the selectivity filter residues S78, G233, G353 and G457 (coloured in cpk magenta), and other neighbouring residues. Na$^+$ is likely to enter and exit the permeation trajectory from several points on both sides of the transporter, but always by-passes the selectivity filter constriction. K$^+$ enters the selectivity filter and effectively blocks the gating trajectory.

(A-B; right panels) Detailed views of bound Na$^+$ and K$^+$ ions that interact through respective penta-hedral (yellow dashed lines) and hexa-hedral (green dashed lines) coordination patterns with residues adjoining the selectivity filter residues S78, G233, G353 and G457. The ionic distances of 2.3-2.8 Å for Na$^+$ between V76, S77, S78, N231, C232 and H351, and of 2.6-2.9 Å for K$^+$ between V76, N231, C232, H351, N455 and V456, are formed between carbonyl oxygens of these residues (in sticks in atomic colours), located near selectivity filter residues. The K$^+$ ion effectively blocks the permeation channel through the protein pore, contrary to Na$^+$.

501   **Figure 1**

505 **Figure 2**

506
507

508 **Figure 3**

511 **Figure 4**

512
513
514

515   **Figure 5**

516
517