Full Title: High affinity Na^+ transport by wheat HKT1;5 is blocked by K^+

Running title: HKT1;5 proteins catalyse dual affinity Na^+ transporter in wheat

Authors: Bo Xu[1,2], Maria Hrmova[2] and Matthew Gilliham[1,2*]

[1]Australian Research Council Centre of Excellence in Plant Energy Biology,

University of Adelaide, Waite Research Precinct, Glen Osmond, South Australia

5064, Australia

[2]School of Agriculture, Food and Wine, and Waite Research Institute, University of

Adelaide, Waite Research Precinct, Glen Osmond, South Australia 5064, Australia

*Corresponding author: e-mail: matthew.gilliham@adelaide.edu.au

Keywords: salinity; High-affinity K Transporters; membrane transport; KtrB; dual

affinity; sodium exclusion

Abstract

The wheat sodium transporters TmHKT1;5-A and TaHKT1;5-D are encoded by

genes underlying major shoot Na^+ exclusion loci *Nax2* and *Kna1* from *Triticum*

monococcum (Tm) and *Triticum aestivum* (Ta), respectively. In contrast to HKT2

transporters that have been shown to exhibit high affinity K^+-dependent Na^+ transport,

HKT1 proteins have, with one exception, only been shown to catalyse low affinity

Na^+ transport and no K^+ transport. Here, using heterologous expression in *Xenopus*

laevis oocytes we show that both TmHKT1;5-A and TaHKT1;5-D encode dual (high

and low) affinity Na^+-transporters with the high-affinity component being abolished

when external K^+ is in excess of external Na^+. Based on 3-D structural modelling we

propose that tighter binding of K^+, compared to that of Na^+ in the selectivity filter

region by means of additional van der Waals forces, explains the K^+ block at the

molecular level. The low-affinity component for Na^+ transport of TmHKT1;5-A had a

lower K_m than that of TaHKT1;5-D and was less sensitive to external K^+. We propose

that these properties underpin the improvements in shoot Na^+-exclusion and crop

plant salt tolerance following the introgression of TmHKT1;5-A into diverse wheat

backgrounds.

Introduction

Crops suffer reduced growth and productivity under salinity stress. Salt, when it builds up to high concentrations in the growing medium (i.e. soil solution) imposes an osmotic limitation on water uptake, interferes with optimal nutrient homeostasis, and leads to the build up of leaf cellular sodium (Na^+) concentrations, which causes an ionic toxicity that limits photosynthesis and carbon assimilation in plants (1).

A major genetic mechanism contributing towards salinity tolerance of most crops including the cereals rice (*Oryza sativa*) and wheat (*Triticum aestivum* and *Triticum monococcum*) is Na^+ exclusion from leaves, with the High-affinity potassium (K^+) transporter (HKT) protein family having a major role in this trait (2-6). This family of proteins is present in all plants so far sequenced; more broadly the members of the high affinity K^+/Na^+ transporting Ktr/TrK/HKT superfamily of proteins are present in bacteria (Ktr and TrK), fungi (TrK) and plants (HKT) (7,8).

These plant HKTs are divided into 2 clades: 1) class 1 HKT proteins (HKT1;x) are mostly Na^+-selective transporters; 2) whereas class 2 HKTs (HKT2;y) mostly function as K^+-Na^+ symporters and so far have been only identified in cereal monocots (9,10). This definition has been challenged on occasion, such as for OsHKT2;4, where following its characterization in *Xenopus laevis* oocytes, there are conflicting reports on its permeability to Ca^{2+}, Mg^{2+} and NH_4^+ in addition to K^+ and Na^+ (11-13). Furthermore, there are two reports of K^+ permeability in the HKT1 clade. Firstly, *Arabidopsis* AtHKT1 could complement K^+ transport deficient *E. coli,* although no K^+ permeability could be found when *AtHKT1* was expressed in *X. laevis* oocytes in the same study (14). Secondly, *Eucalyptus camaldulensis* EcHKT1;1 and EcHKT1;2, which appear to be the exception for the HKT1 proteins characterized so far in that they transport both Na^+ and K^+ when expressed in *X. laevis* oocytes (15). Other HKT1 transporters have no reported K^+ permeability (4,5,10,14,16-20).

Whilst K^+ transport is not a common feature of HKT1 transporters, several have shown the property of K^+-regulated Na^+ transport (19,21-23). For instance, the presence of 10 mM external K^+ ($[K^+]_{ext}$) reduced the inward Na^+ current of *X. laevis* oocytes expressing *HKT1;5* from *T. aestivum* (*TaHKT1;5-D*) (19), and reduced outward Na^+ currents generated by expression of *HKT1;2* from *Solanum*

lycopersicum (*SlHKT1;2*) (21). Interestingly, $[K^+]_{ext}$ stimulated both inward and

outward Na^+ transport of *X. laevis* oocytes expressing two *HKT1;4* alleles from

Triticum turgidum L. subsp. *Durum*, when assayed with solutions of very low ionic

strength (23).

Ktr/TrK/HKT proteins are predicted to consist of 8 transmembrane α-helices that fold

into four pseudo-tetramers around a central pore (6,16,18). The selectivity filter lining

the pore of Ktr and TrK and class 2 HKT proteins is composed of four glycine

residues. In class 1 HKTs the first glycine within the predicted selectivity filter is

substituted with serine (16); the lack of K^+ permeability in this clade has been linked

to the presence of this glycine substitution (glycine for serine) (10,14,16-18); but the

exact molecular mechanism underlying this block is to be determined.

Here, we further characterise HKT1;5 proteins underlying two major salt tolerance

associated loci, *TmHKT1;5-A* (*Nax2*) and *TaHKT1;5-D* (*Kna1*) from *T. monococcum*

and T. aestivum respectively. We find that both TmHKT1;5-A and TaHKT1;5-D

encode dual affinity Na^+-transporters and that their dual affinity Na^+ transport can be

blocked by raising the external K^+ concentration. We further propose that the

adjoining residues within the selectivity filter cause this block by the virtue of a

higher number of van der Waals interactions with the K^+ ion, than with Na^+. Thus, the

Na^+ ion is capable of traversing the selectivity filter, while K^+ causes a block. This is

the first report that Na^+ transport has two affinities in the HKT1 clade and

demonstrates that dual affinity transport is a common property of both clades of the

HKT family.

Results

K^+ sensitivity of TmHTK1;5-A and TaHKT1;5-D

Previously, it was shown that inward Na^+ transport of both TmHKT1;5-A and

TaHKT1;5-D was inhibited by external K^+ ($[K^+]_{ext}$) (4,19). Here, we further compared

the K^+-sensitivity of TmHKT1;5-A and TaHKT1;5-D (Fig. 1). An increase in $[K^+]_{ext}$

to 30 mM reduced the channel inward conductance (i.e. <100 mV) of TmHKT1;5-A

by approximately 75% in a 1 mM Na^+ ($[Na^+]_{ext}$) bath solution; similarly this amount

of K^+ inhibited that of TaHKT1;5-D up to 95 % (Figs. 1A, 1B and 2). However, such

Na^+-inhibition by K^+ was gradually decreased by increasing $[Na^+]_{ext}$ (Figs. 1 and 2).

For instance, 10 mM $[K^+]_{ext}$ suppressed the channel inward conductance of

TmHKT1;5-A by 43.5% in 1 mM $[Na^+]_{ext}$ and by 16.5% in 10 mM $[Na^+]_{ext}$; it

similarly reduced the TaHKT1;5-D inward conductance by 57% in 1 mM $[Na^+]_{ext}$

and by 23% in 10 mM $[Na^+]_{ext}$ (Figs. 1 and 2). When $[Na^+]_{ext}$ was increased to 30

mM, the channel inwared conductance of TmHKT1;5-A was insensitive to $[K^+]_{ext}$,

whereas that of TaHKT1;5-D was inhibited (Fig. 2).

The increase in $[K^+]_{ext}$ from 0 to 10 mM did not significantly change the reversal

potential of TmHKT1;5-A or TaHKT1;5-D; all reversal potentials were close to the

theoretical equilibrium potential for Na^+ under all conditions (Fig. 3A, 3B).

Moreover, the observed shift in reversal potential were not significantly different

from the theoretical Nernst shift for Na^+ regardless of the presence or absence of 10

mM $[K^+]_{ext}$ (Fig. 3B). This suggests that neither of transporter proteins were

permeable to Na^+ and not to K^+.

Dual affinity transport of Na^+ by TmHTK1;5-A and TaHKT1;5-D

The affinity for Na^+ transport of TmHKT1;5-A was previously reported to be ~4-fold

higher than the Na^+-transport affinity of TaHKT1;5-D (4,6,19), which we confirm

here (Fig. 4A, 4B). On closer examination, we found a property that has not been

described before. When lower concentrations of $[Na^+]_{ext}$ were present it was clear that

the concentration dependence of inward Na^+ currents could be fitted by two

components, for both TmHKT1;5-A and TaHKT1;5-D (Fig. 4). For TmHKT1;5-A,

the high affinity component of Na^+ transport had a K_m of 18.4 μM ± 0.1 μM when

$[Na^+]_{ext}$ was below 0.1 mM ($K_{m\,(<0.1\,Na)}$), whereas at concentrations of Na^+ above 0.1

mM the K_m was 1.0 ± 0.2 mM ($K_{m\,(>0.1\,Na)}$) (Fig. 4A). The K_m for the higher affinity

component of inward Na^+ transport for TaHKT1;5-D was 15.3 μM ± 0.9 μM when

$[Na^+]_{ext}$ was lower than 0.1 mM ($K_{m\,(<0.1\,Na)}$), whilst the lower affinity phase had a K_m

of 4.3 ± 0.5 mM when $[Na^+]_{ext}$ was higher than 0.1 mM ($K_{m\,(>0.1\,Na)}$) (Fig. 4B).

We examined the extent of K^+-inhibition of both the lower and higher affinity

components of the inward Na^+ currents when $[K^+]_{ext}$ was 10 mM. The (normalised)

inward Na^+ current carried by both proteins was significantly suppressed by 10 mM

$[K^+]_{ext}$, resulting in negligible inward Na^+ transport by TmHKT1;5-A and TaHKT1;5-

D in presence of $[Na^+]_{ext}$ below 0.1 mM (Fig. 4C and 4D). In the presence of 10 mM

$[K^+]_{ext}$ it was now possible to fit the concentration dependence of inward current

through TmHKT1;5-A and TaHKT1;5-D with one component. The K_m values of both

TmHKT1;5-A and TaHKT1;5-D were decreased approximately by 2.5-fold by 10

mM $[K^+]_{ext}$ (Fig. 4E and 4F).

Structural modelling of TaHKT1;5-D

Three-dimensional structural models of TaHKT1;5-D were generated to explore why

K^+ does not traverse the pore but instead blocks the Na^+ currents, using the KtrB K^+

transporter from *B. subtilis* as a template (6,24). The KtrB K^+ protein was crystallised

in the presence of KCl, hence the original structure contained a K^+ ion in the

selectivity filter region. This ion was substituted by Na^+ during modelling of

TaHKT1;5-D as it transports Na^+ but not K^+ (6). Ramachandran analysis indicated

that the template and TaHKT1;5-D models generated in complex with Na^+, and K^+

had satisfactory stereo-chemical quality; Ramachandran plots showed only two

residues positioned in disallowed regions (0.5% of all residues, except glycine and

proline residues). Average G-factors (measures of correctness of dihedral angles and

main-chain covalent forces of protein molecules) calculated by PROCHECK of the

template and TaHKT1;5-D models with K^+ and Na^+ were 0.06. -0.18 and -0.13,

respectively. The ProSa 2003 analysis of z-scores (-8.4; -6.0 and -5.8) indicated that

template and modelled structures with K^+ and Na^+ had acceptable conformational

energies.

The TaHKT1;5-D model in complex with Na^+ showed that the Na^+ ion coordination

sphere had trigonal bipyramidal geometry, as defined by the Neighborhood algorithm

(25), with coordination number 5. Na^+ interacted with V76, S77, S78, N231, C232

and H351 (all carbonyl oxygens), that lies near the selectivity filter residues S78,

G233, G353 and G457 (Fig. 5A). Notably, the carbonyl oxygen one of the selectivity

filter residues (S78) directly participated in Na^+ binding (Fig. 5A). Conversely, the

TaHKT1;5-D model in complex with K^+ showed that the K^+ ion coordination sphere

had octahedral geometry (25) with coordination number 6; K^+ interacted with V76,

N231, C232, H351, N455 and V456, that were also in the close neighbour of

selectivity filter residues (Fig. 5B). However, in the second case, none of the

selectivity filter residues participate in direct binding of K^+. Ionic coordination

spheres and Na^+ and K^+ distances of residues were evaluated using the Metal Binding

Site Validation Server (25) that utilises more than 500 Protein Data Bank accession

numbers. Ionic coordination spheres and binding modes were found to be acceptable,

and ionic distances were within the following ranges: for Na^+ ranged between 2.3 and

2.4 Å; for K^+ were between 2.6 and 2.9 Å (Fig. 5, right panels).

Discussion

Oocytes expressing *TmHKT1;5-A* or *TaHKT1;5-D* have been previously reported to

undergo no obvious change in their reversal potential regardless of the presence or

absence of K^+ in bath solution. Interestingly, both transporters had differential

sensitivity to external $[K^+]_{ext}$. Whilst K^+ block occurs for TmHKT1;5-A mediating

inward transport it was not evident at $[Na^+]_{ext}$ above 10 mM, whereas this inhibition

did happen for TaHKT1;5-D up to 30 mM $[Na^+]_{ext}$ (Fig. 2) (4,19). This differential

inhibition effect of K^+ upon Na^+ transport at high $[Na^+]_{ext}$ by TaHKT1;5-D may result

from a range of factors, that govern the rates of Na^+ and K^+ ion binding and

subsequent Na^+ transport. A variety of factors could control the transport of Na^+ and

the preliminary displacement of competing K^+ with a consequent energetic cost,

before Na^+ passes desolvated or less-well hydrated than competing K^+. Examples

include free-energy variations between ions in binding sites relative to the

corresponding quantity in bulk water, differences in selectivity filters solvent

exposures (or dielectric constants), overall pore rigidity/stiffness (26), structural

changes within funnels and selectivity filters due to residue re-orientation, differences

in the hydration (coordination) numbers of Na^+ and K^+ (Na^+ and K^+ bind six and

seven water molecules, respectively (27)) and the alterations on the rates of Na^+ or K^+

de-solvation (28). It is therefore plausible to expect that both ions would compete for

binding sites within HKT funnel regions, yielding ion-protein binding sites with

differential binding strengths.

Here, we conduct advanced 3D homology modelling and include explicit ionic radii of

Na^+ and K^+ based on the CHARMM force field (29), rather than a traditional rigid

body-type modelling for ligands. The aim has been to refine the previous structural

model of HKT1;5 to show that the current model contains a pore that allows the

passage of Na^+ but not K^+ (6,10,18). This is consistent with the functional

observations in this and previous studies (4,19). Previously it was suggested that the

Gly-Gly-Gly-Gly motif is important in coordinating the K^+ ion in Ktr/TrK transporters

and in conferring dual Na^+-K^+ transport in HKT2 transporters (16,30). Here our

structural modelling suggests that the serine residue within the Ser-Gly-Gly-Gly

selectivity motif directly interacts and binds with Na^+ (Fig. 5B). Furthermore, our

model suggests, despite ionic distances being shorter for Na^+ than those for K^+, that

K^+ could be bound more strongly due to a higher coordination pattern (Fig. 5A and

5B-right panels). Obviously, during binding, K^+ takes an advantage of its larger ionic

radius: 152 picometer (pm), compared to that of Na^+ (116 pm); these parameters

reflects empirical atomic radii of K (220 pm) and Na (180 pm). We assume that under

the high concentrations of K^+, this ion may outcompete Na^+ and would become bound

in the selectivity filter, thus effectively blocking Na^+ transport. This is illustrated by

calculating permeation channels in both complexes, using the Mole Voronoi

algorithm, that predicts permeation trajectories and identifies path bottlenecks. From

the calculated permeation paths, it could be deduced that the K^+ ion effectively blocks

the permeation channel through the major gating protein pore, contrary to Na^+. It is

also obvious that Na^+ is likely to enter and exit the permeation trajectory from several

points on both sides of the transporter, but always by-passes the selectivity filter

constriction. Both ions arrive at the funnel of HKT proteins in solvated (hydrated)

forms. While the hydration number for Na^+ is 6, that for K^+ is 7 at ambient

temperature (27,31). The significance of the seven-fold water coordination for K^+ is

that this solvated complex may form a stronger hydrogen-bond interaction pattern (in

funnels of HKT proteins), and would be de-solvated with a higher energy input and

thus slower that the Na^+ solvated complex (28). It is also plausible to expect that both

solvated ion complexes would compete for binding sites within HKT funnel regions

with different binding strengths. All in all, this would contribute to the blockage of the

pore entry of HKT transporters by $K^{\pm}$.

Potassium permeability for HKT1 transporters has only been reported once for the

proteins that have been characterised in *X. laevis* oocytes. The transporters EcHKT1;1

and EcHKT1;2 allow permeation of both K^+ and Na^+ (15). Interestingly, when 3D

models of the Na^+ selective OsHKT1;5 and K^+ permeable EcHKT1;2 were

constructed and compared the pore size was predicted to be 0.2 Å larger in EcHKT1;2

(and that of OsHKT1;5 and TaHKT1;5-D), which would more than account for the

weaker interactions with K^+ and allow this larger ion to pass through the pore as well

as Na^+ (10). Without advanced 3D modelling of each and every transporter in the

HKT family to show how the selectivity filter forms in the context of the remainder of

the protein it would not be possible to predict which other HKT1s may allow passage

of K^+. Predictions based on sequence alone are insufficient to suggest how the two

sets of proteins evolved their individual Na^+ and K^+ transport characteristics. An

extension of the model to include other HKT proteins, a greater survey of structure-

function relationships and mutations of key residues to predict functionally relevant

residues and evolutionary relationships will be the focus of further studies.

Two affinity ranges for Na^+ transport have been detected in the HKT family

previously, but only for HKT2 transporters. OsHKT2;1, OsHKT2;2, OsHKT2;4 and

HvHKT2;1(12,13,20,32-34). For instance, for OsHKT2;2, when K^+ is present, the

transport of Na^+ is K^+-dependent and displays a high affinity (K_m = 0.077 mM),

whereas showing a lower affinity when absence of K^+ (K_m= 16 mM) (33,34).

OsHKT2;1 also acquires two phases of Na^+-transport affinity, whose K_m for Na^+ is

respectively 9.5 µM at $[Na^+]_{ext}$ below 1 mM and is 2.2 mM at $[Na^+]_{ext}$ > 1 mM (20).

The K^+ transport catalysed by HvHKT2;1 is affected by both $[Na^+]_{ext}$ and $[K^+]_{ext}$, such

as a decrease in its K^+ transport affinity from 30 µM down to 3.5 mM by $[Na^+]_{ext}$

increased from 0.5 mM to 30 mM and a greatly reduced channel conductivity for K^+

by increasing $[K^+]_{ext}$ above ~3 mM (35). This is the first time that dual affinity

characteristics have been reported for the HKT1;x family (Fig. 4). Considering the

relatively high homology between both clades (58% similarity/41% identity between

TaHKT1;5 and TaHKT2;1) it is perhaps not surprising that both clades can confer

dual affinity transport. How and why such ability has evolved and is conferred

structurally is still to be determined.

In summary, we have detected that TaHKT1;5-D and TmHKT1;5-A can both conduct

high and low affinity Na+ transport, both phases can be blocked by external K^+ with

the low affinity the low affinity component of Na^+ transport for TmHKT1;5-A being

less inhibited by external K^+ than TaHKT1;5-D (Fig. 4). The two K_m at the high

affinity phases were similar, but the K_m value for the low affinity component of

TmHKT1;5-A was always ~4 fold smaller regardless of the absence or presence of

$[K^+]_{ext}$ (Fig. 4) (4,19). TmHKT1;5-A has been shown to be more effective in

conferring shoot Na^+ exclusion to wheat compared to TaHKT1;5-D (36). We propose

that the higher affinity (with half-maximal activity nearer 1 mM compared to 4 mM),

and a lower sensitivity to $[K^+]_{ext}$ are both characteristics that can help confer better

shoot Na^+ exclusion, which is a property that can lead to greater salt tolerance in the

field for wheat (4,36,37).

Materials and Methods

282   Brief methods for cloning of *TmHKT1;5-A* and *TaHKT1;5-D* as well as its functional

283   characterisation in heterologous expression systems were described Munns et al. (4)

284   and Byrt et al. (19). Further details are included here.

Two-electrode voltage clamp recording in *X. laevis* oocytes

Oocyte recording followed the methods as described in Munns et al. (4) and Byrt et

al. (19). Briefly, 46 nl/23 ng of cRNA or equal volumes of RNA-free water were

injected into oocytes, followed by an incubation for 48 h before recording. Membrane

currents were recorded in the HMg solution (6 mM $MgCl_2$, 1.8 mM $CaCl_2$, 10 mM

MES and pH 6.5 adjusted with a TRIS base) ± Na^+ glutamate and/or K^+ glutamate as

indicated in relevant Figures. All solution osmolarities were adjusted using mannitol

at 220-240 mOsmol kg^{-1}.

Construction of 3D models of TaHKT1;5D in complex with Na^+ and K^+ ions

The most suitable template for TaHKT1;5D, the KtrB K^+ transporter from *B. subtilis*

(Protein Data Bank accession 4J7C, chain I) (24), was identified as previously

described (6). The KtrB K^+ protein was crystallised in the presence of KCl, hence the

structure contains a K^+ ion in the selectivity filter region. This ion was substituted by

Na^+ during modelling of TaHKT1;5D that transport Na^+ at a greater rate than K^+ (6).

3D structural models in complex with Na^+, and K^+ were generated using Modeller

9v16 (38) on a Linux station running the Fedora 12 operating system, as previously

described (6,10,18). During modelling, attention was paid to ionic radii of Na^+, and

K^+, whose topology parameters were taken from CHARMM (29). In each case, a total

of 100 models were generated that were scored by Modeller using the modeller

objective function (39), the discrete optimised protein energy function (40),

PROCHECK (41), ProSa 2003 (42) and FoldX (43). Best scoring models constructed

in Modeller 9v19 were further subjected to energy minimisation (knowledge-based

Yasara2 forcefield with parameters for bond distances, planarity of peptide bonds,

bond angles, Coulomb terms, dihedral angles, and van der Waals forces) (44),

combined with the particle-mesh-Ewald (PME) energy function for long range electrostatics (cutoff 8.0 Å), to obtain smoothed electrostatic potentials. Structural images were generated with PyMOL Molecular Graphics System V1.8.2.0 (Schrődinger LLC, Portland, OR, USA). Permeation channels and path bottlenecks in both TaHKT1;5D ionic complexes were calculated using the Mole algorithm, embedded in PyMol (45), but using the x-y-z coordinates of Na^+ or K^+ as specifications for starting points.

Funding

This work was supported by funding to M.G. from the Grains Research and Development Corporation (UA00145), the Australian Research Council through the Centre of Excellence (CE1400008) and Future Fellowship (FT130100709) schemes, and by funding to M.H. from the Australian Research Council (DP120100900).

References

1. Munns, R., and Tester, M. (2008) Mechanisms of salinity tolerance. *Annu. Rev. Plant Biol.* **59**, 651-681
2. Ren, Z.-H., Gao, J.-P., Li, L.-G., Cai, X.-L., Huang, W., Chao, D.-Y., Zhu, M.-Z., Wang, Z.-Y., Luan, S., and Lin, H.-X. (2005) A rice quantitative trait locus for salt tolerance encodes a sodium transporter. *Nat. Genet.* **37**, 1141-1146
3. James, R. A., Davenport, R. J., and Munns, R. (2006) Physiological characterization of two genes for Na^+ exclusion in durum wheat, *Nax1* and *Nax2*. *Plant Physiol.* **142**, 1537-1547
4. Munns, R., James, R. A., Xu, B., Athman, A., Conn, S. J., Jordans, C., Byrt, C. S., Hare, R. A., Tyerman, S. D., Tester, M., Plett, D., and Gilliham, M. (2012) Wheat grain yield on saline soils is improved by an ancestral Na^+ transporter gene. *Nat. Biotech.* **30**, 360-364
5. Campbell, M. T., Bandillo, N., Al Shiblawi, F. R. A., Sharma, S., Liu, K., Du, Q., Schmitz, A. J., Zhang, C., Véry, A.-A., and Lorenz, A. J. (2017) Allelic variants of OsHKT1;1 underlie the divergence between indica and japonica subspecies of rice (*Oryza sativa*) for root sodium content. *PLoS Genet.* **13**, e1006823
6. Xu, B., Waters, S., Byrt, C. S., Plett, D., Tyerman, S. D., Tester, M., Munns, R., Hrmova, M., and Gilliham, M. (2018) Structural variations in wheat HKT1;5 underpin differences in Na^+ transport capacity. *Cell. Mol. Life Sci.*, 1-12
7. Sentenac, H., and Bonneaud, N. (1992) Cloning and expression in yeast of a plant potassium ion transport system. *Science* **256**, 663
8. Corratgé-Faillie, C., Jabnoune, M., Zimmermann, S., Véry, A.-A., Fizames, C., and Sentenac, H. (2010) Potassium and sodium transport in non-animal cells: the Trk/Ktr/HKT transporter family. *Cell. Mol. Life Sci.* **67**, 2511-2532

9. Asins, M. J., Villalta, I., Aly, M. M., Olias, R., Álvarez De Morales, P.,
Huertas, R., Li, J., JAIME-PÉREZ, N., Haro, R., and Raga, V. (2013) Two
closely linked tomato HKT coding genes are positional candidates for the
major tomato QTL involved in Na$^+$/K$^+$ homeostasis. *Plant Cell Environ.* **36**,
1171-1191
10. Waters, S., Gilliham, M., and Hrmova, M. (2013) Plant high-affinity
potassium (HKT) transporters involved in salinity tolerance: structural insights
to probe differences in ion selectivity. *Int. J. Mol. Sci.* **14**, 7660-7680
11. Lan, W.-Z., Wang, W., Wang, S.-M., Li, L.-G., Buchanan, B. B., Lin, H.-X.,
Gao, J.-P., and Luan, S. (2010) A rice high-affinity potassium transporter
(HKT) conceals a calcium-permeable cation channel. *Proc. Natl. Acad. Sci.*
U.S.A. **107**, 7089-7094
12. Horie, T., Brodsky, D. E., Costa, A., Kaneko, T., Schiavo, F. L., Katsuhara,
M., and Schroeder, J. I. (2011) K$^+$ transport by the OsHKT2; 4 transporter
from rice with atypical Na$^+$ transport properties and competition in permeation
of K$^+$ over Mg^{2+} and Ca^{2+} ions. *Plant Physiol.* **156**, 1493-1507
13. Sassi, A., Mieulet, D., Khan, I., Moreau, B., Gaillard, I., Sentenac, H., and
Véry, A.-A. (2012) The rice monovalent cation transporter OsHKT2; 4:
revisited ionic selectivity. *Plant Physiol.* **160**, 498-510
14. Uozumi, N., Kim, E. J., Rubio, F., Yamaguchi, T., Muto, S., Tsuboi, A.,
Bakker, E. P., Nakamura, T., and Schroeder, J. I. (2000) The Arabidopsis
HKT1 gene homolog mediates inward Na$^+$ currents in *Xenopus laevis* oocytes
and Na$^+$ uptake in *Saccharomyces cerevisiae*. *Plant Physiol.* **122**, 1249-1260
15. Liu, W., Fairbairn, D. J., Reid, R. J., and Schachtman, D. P. (2001)
Characterization of two HKT1 homologues from *Eucalyptus camaldulensis*
that display intrinsic osmosensing capability. *Plant Physiol.* **127**, 283-294
16. Mäser, P., Hosoo, Y., Goshima, S., Horie, T., Eckelman, B., Yamada, K.,
Yoshida, K., Bakker, E. P., Shinmyo, A., and Oiki, S. (2002) Glycine residues
in potassium channel-like selectivity filters determine potassium selectivity in
four-loop-per-subunit HKT transporters from plants. *Proc. Natl. Acad. Sci.*
U.S.A. **99**, 6428-6433
17. Platten, J. D., Cotsaftis, O., Berthomieu, P., Bohnert, H., Davenport, R. J.,
Fairbairn, D. J., Horie, T., Leigh, R. A., Lin, H.-X., and Luan, S. (2006)
Nomenclature for HKT transporters, key determinants of plant salinity
tolerance. *Trends Plant Sci.* **11**, 372-374
18. Cotsaftis, O., Plett, D., Shirley, N., Tester, M., and Hrmova, M. (2012) A two-
staged model of Na$^+$ exclusion in rice explained by 3D modeling of HKT
transporters and alternative splicing. *PLoS One* **7**, e39865
19. Byrt, C. S., Xu, B., Krishnan, M., Lightfoot, D. J., Athman, A., Jacobs, A. K.,
Watson-Haigh, N. S., Plett, D., Munns, R., and Tester, M. (2014) The Na$^+$
transporter, TaHKT1;5-D, limits shoot Na$^+$ accumulation in bread wheat.
Plant J. **80**, 516-526
20. Jabnoune, M., Espeout, S., Mieulet, D., Fizames, C., Verdeil, J.-L., Conéjéro,
G., Rodríguez-Navarro, A., Sentenac, H., Guiderdoni, E., and Abdelly, C.
(2009) Diversity in expression patterns and functional properties in the rice
HKT transporter family. *Plant Physiol.* **150**, 1955-1971
21. Almeida, P., de Boer, G.-J., and de Boer, A. H. (2014) Differences in shoot
Na$^+$ accumulation between two tomato species are due to differences in ion
affinity of HKT1;2. *J. Plant Physiol.* **171**, 438-447

22. Almeida, P., Feron, R., de Boer, G.-J., and de Boer, A. H. (2014) Role of Na^+,
K^+, Cl^-, proline and sucrose concentrations in determining salinity tolerance
and their correlation with the expression of multiple genes in tomato. *AoB*
Plants **6**, plu039
23. Ben Amar, S., Brini, F., Sentenac, H., Masmoudi, K., and Véry, A.-A. (2013)
Functional characterization in *Xenopus oocytes* of Na^+ transport systems from
durum wheat reveals diversity among two HKT1; 4 transporters. *J. Exp. Bot.*
65, 213-222
24. Vieira-Pires, R. S., Szollosi, A., and Morais-Cabral, J. H. (2013) The structure
of the KtrAB potassium transporter. *Nature* **496**, 323-328
25. Zheng, H., Chordia, M. D., Cooper, D. R., Chruszcz, M., Müller, P.,
Sheldrick, G. M., and Minor, W. (2014) Validation of metal-binding sites in
macromolecular structures with the CheckMyMetal web server. *Nat. Protoc.*
9, 156
26. Dudev, T., and Lim, C. (2010) Factors governing the Na^+ vs K^+ selectivity in
sodium ion channels. *J. Am. Chem. Soc.* **132**, 2321-2332
27. Rowley, C. N., and Roux, B. t. (2012) The solvation structure of Na^+ and K^+
in liquid water determined from high level ab initio molecular dynamics
simulations. *J. Chem. Theory Comput.* **8**, 3526-3535
28. Degrève, L., Vechi, S. M., and Junior, C. Q. (1996) The hydration structure of
the Na^+ and K^+ ions and the selectivity of their ionic channels. *Biochim.*
Biophys. Acta-Bioenergetics **1274**, 149-156
29. Brooks, B. R., Brooks, C. L., MacKerell, A. D., Nilsson, L., Petrella, R. J.,
Roux, B., Won, Y., Archontis, G., Bartels, C., and Boresch, S. (2009)
CHARMM: the biomolecular simulation program. *Journal of Computational*
Chemistry **30**, 1545-1614
30. Durell, S. R., and Guy, H. R. (1999) Structural models of the KtrB, TrkH, and
Trk1, 2 symporters based on the structure of the KcsA K^+ channel. *Biophys. J.*
77, 789-807
31. Ma, H. (2014) Hydration structure of Na^+, K^+, F^-, and Cl^- in ambient and
supercritical water: A quantum mechanics/molecular mechanics study. *Int. J.*
Quantum Chem. **114**, 1006-1011
32. Schachtman, D. P., and Schroeder, J. I. (1994) Structure and transport
mechanism of a high-affinity potassium uptake transporter from higher plants.
Nature **370**, 655-658
33. Rubio, F., Gassmann, W., and Schroeder, J. I. (1995) Sodium-driven
potassium uptake by the plant potassium transporter HKT1 and mutations
conferring salt tolerance. *Science* **270**, 1660
34. Yao, X., Horie, T., Xue, S., Leung, H.-Y., Katsuhara, M., Brodsky, D. E., Wu,
Y., and Schroeder, J. I. (2010) Differential sodium and potassium transport
selectivities of the rice OsHKT2; 1 and OsHKT2; 2 transporters in plant cells.
Plant Physiol. **152**, 341-355
35. Mian, A., Oomen, R. J., Isayenkov, S., Sentenac, H., Maathuis, F. J., and
Véry, A. A. (2011) Over-expression of an Na^+-and K^+-permeable HKT
transporter in barley improves salt tolerance. *Plant J.* **68**, 468-479
36. James, R. A., Blake, C., Zwart, A. B., Hare, R. A., Rathjen, A. J., and Munns,
R. (2012) Impact of ancestral wheat sodium exclusion genes *Nax1* and *Nax2*
on grain yield of durum wheat on saline soils. *Funct. Plant Biol.* **39**, 609-618
37. James, R. A., Blake, C., Byrt, C. S., and Munns, R. (2011) Major genes for
Na^+ exclusion, *Nax1* and *Nax2* (wheat *HKT1; 4* and *HKT1; 5*), decrease Na+

accumulation in bread wheat leaves under saline and waterlogged conditions.
J. Exp. Bot. **62**, 2939-2947

38. Šali, A., and Blundell, T. L. (1993) Comparative protein modelling by
satisfaction of spatial restraints. *J. Mol. Biol.* **234**, 779-815

39. Shen, M. y., and Sali, A. (2006) Statistical potential for assessment and
prediction of protein structures. *Protein Sci.* **15**, 2507-2524

40. Eswar, N., Eramian, D., Webb, B., Shen, M.-Y., and Sali, A. (2008) Protein
structure modeling with MODELLER. in *Structural proteomics*, Springer. pp
145-159

41. Laskowski, R. A., MacArthur, M. W., Moss, D. S., and Thornton, J. M. (1993)
PROCHECK: a program to check the stereochemical quality of protein
structures. *J. Appl. Crystallogr.* **26**, 283-291

42. Sippl, M. J. (1993) Recognition of errors in three-dimensional structures of
proteins. *Proteins: Struct., Funct., Bioinf.* **17**, 355-362

43. Schymkowitz, J., Borg, J., Stricher, F., Nys, R., Rousseau, F., and Serrano, L.
(2005) The FoldX web server: an online force field. *Nucleic Acids Res.* **33**,
W382-W388

44. Krieger, E., Darden, T., Nabuurs, S. B., Finkelstein, A., and Vriend, G. (2004)
Making optimal use of empirical energy functions: force-field
parameterization in crystal space. *Proteins: Struct., Funct., Bioinf.* **57**, 678-
683

45. Petřek, M., Košinová, P., Koča, J., and Otyepka, M. (2007) MOLE: a Voronoi
diagram-based explorer of molecular channels, pores, and tunnels. *Structure*
15, 1357-1363

Figures and figure legends

Figure 1. K$^+$-sensitivity of Na$^+$ currents carried by TmHKT1;5-A and TaHKT1;5-D. A-D, current-voltage (I/V) curves of Na$^+$ currents recorded from *X. laevis* oocytes expressing *TmHKT1;5-A* (A, C) and *TaHKT1;5-D* (B, D) in 1 mM Na$^+$ plus 0-30 mM K$^+$ (A,B), or 0-30 mM Na$^+$ solutions with or without 10 mM K$^+$ (C,D). Mean ± S.E.M, n = 5-6.

Figure 2. Channel conductance of TmHKT1;5-A and TaHKT1;5-D. The channel conductance of TmHKT1;5-A and TaHKT1;5-D was calculated from Figure 1, based on the slope of curve between -140 mV and -120 mV, statistical difference was determined by *Students' t-test, *P < 0.05* and ** *P < 0.01*.

Figure 3. Shift in reversal potential of TmHKT1;5-A and TaHKT1;5-D in response to changes in [Na$^+$]$_{ext}$ and [K$^+$]$_{ext}$.
A, Reversal potential of currents in *X. laevis* oocytes expressing *TmHKT1;5-A* (red) and *TaHKT1;5-D* (blue) derived from Figure. 1. B, Reversal potential of Na$^+$ currents plotted with theoretical Nernst potential derived from internal concentrations measured in Munns et al. (2012) [19] and Byrt et al., (2014) [4]. Voltage at which current reversed in oocytes expressing *TmHKT1;5-A* and *TaHKT1;5-D* in solutions containing 1 mM to 10 mM Na$^+$ ([1-10]$_{ext}$ Na$^+$) and 10 mM to 30 mM Na$^+$ ([10-30]$_{ext}$ Na$^+$) with or without 10 mM [K$^+$]$_{ext}$, ΔV_{Na} of 0 mM or 10 mM K$^+$ = experimentally measured shift in reversal potential for Na$^+$ in 0 mM or 10 mM K$^+$ conditions as indicated; ΔV_{Na} of E$_{Na}$ = theoretical equilibrium (or Nernst) potential for Na$^+$ was 57.7 mV and 26.9 mV, respectively for [1-10]$_{ext}$ Na$^+$ and [10-30]$_{ext}$ Na$^+$. One way ANOVA was performed for Figure 3B and no statistical differences were found between the theoretical and experimentally measured values.

Figure 4. Transport affinity of TmHKT1;5-A and TaHKT1;5-D in *X. laevis* oocytes.
A-B, dual Na$^+$-transport affinity of TmHKT1;5-A and TaHKT1;5-D in *X. laevis* oocytes, plotted as normalised currents at -140 mV as shown against a series of Na$^+$ glutamate solutions. C-D, high affinity phase of TmHKT1;5-A and TaHKT1;5-D in absence and presence of 10 mM [K$^+$]$_{ext}$. Data represented in A, B and C as mean ± S.E.M (n = 4 for A, n = 7 for B, n = 4 of 0 mM K$^+$ and n = 5 of 10 mM K$^+$ for C). E-

F, Na$^+$-transport affinity of TmHKT1;5-A and TaHKT1;5-D in *X. laevis* oocytes,

plotted as normalised currents at -140 mV as shown against a series of Na$^+$ glutamate

solutions with an additional 10 mM K$^+$ glutamate, n = 5 for E and n = 4 for F.

Figure 5. Molecular models of TaHKT1;5D in complex with Na$^+$ (A) and K$^+$ (B)

ions.

(A-B; left panels) Cartoon representations of TaHKT1;5-D illustrate overall protein

folds with cylindrical α-helices, and translocating permeation channels (black mesh).

Na$^+$ (A) and K$^+$ (B) ions are shown as purple spheres located within the selectivity

filter residues S78, G233, G353 and G457 (coloured in cpk magenta), and other

neighbouring residues. Na$^+$ is likely to enter and exit the permeation trajectory from

several points on both sides of the transporter, but always by-passes the selectivity

filter constriction. K$^+$ enters the selectivity filter and effectively blocks the permeation

at the constriction bottleneck. For clarity, membrane α-helices are only shown.

(A-B; right panels) Detailed views of coordination spheres of Na$^+$ and K$^+$ ions, and

ionic interactions with residues adjoining the selectivity filter residues S78, G233,

G353 and G457. The ionic distances of 2.3-2.8 Å for Na$^+$ between V76, S77, S78,

N231, C232 and H351, and of 2.6-2.9 Å for K$^+$ between V76, N231, C232, H351,

N455 and V456, are formed between carbonyl oxygens of residues (in sticks in atomic

colours). The K$^+$ ion effectively blocks the permeation channel through the protein

pore at the constriction bottleneck point, contrary to Na$^+$.

Figure 1

535
536
537

538 **Figure 2**

539
540

541 **Figure 3**

544 **Figure 4**

545
546
547

548   **Figure 5**

549
550
551