Title:　Differential regulation of activation and differentiation processes

in CD4+ T cell populations and their single-cell level heterogeneity

Authors:

Alla Bradley[1], Tetsuo Hashimoto[1] and Masahiro Ono[2][§]

Affiliations:

[1] Faculty of Life and Environmental Sciences, University of Tsukuba,

Tsukuba, Japan

[2] Department of Life Sciences, Faculty of Natural Sciences, Imperial College

London, Sir Alexander Fleming Building, Imperial College Road, London,

SW7 2AZ, United Kingdom

[§] Address correspondence and reprint requests to Dr Masahiro Ono,

Department of Life Sciences, Sir Alexander Fleming Building, Imperial College

Road, South Kensington, London, SW7 2AZ, United Kingdom. E-mail

address: m.ono@imperial.ac.uk

20
21

Abstract

In T cells, T cell receptor (TCR) signalling initiates downstream transcriptional

mechanisms for T cell activation and differentiation. Foxp3-expressing

regulatory T cells (Treg) require TCR signals for their suppressive function

and maintenance in the periphery. It is, however, unclear how TCR signalling

controls the transcriptional programme of Treg. Since most of studies

identified the transcriptional features of Treg in comparison to naïve T cells,

the relationship between Treg and non-naïve T cells including memory-

phenotype T cells (Tmem) and effector T cells (Teff) is not well understood.

Here we dissect the transcriptomes of various T cell subsets from

independent datasets using the multidimensional analysis method Canonical

Correspondence Analysis (CCA). We show that resting Treg share gene

modules for activation with Tmem and Teff. Importantly, Tmem activate the

distinct transcriptional modules for T cell activation, which are uniquely

repressed in Treg. The activation signature of Treg is dependent on TCR

signals, and is more actively operating in activated Treg. Furthermore, by

analysing single cell RNA-seq data from tumour-infiltrating T cells, we

identified the common shared transcriptional modules for T cell activation,

including *CTLA-4*, in activated Treg and activated Teff. Moreover, we

identified distinct FOXP3-driven and T follicular helper-like transcriptional

modules in activated FOXP3+ Treg and FOXP3- Teff, respectively.

Collectively, we reveal the multidimensional identities and single cell-level

heterogeneity of Treg, identifying the differential regulation of the activation

and differentiation gene modules in Treg, Tmem and Teff during homeostasis

and in the tumour microenvironment.

1 Introduction

T cell receptor (TCR) signalling activates NFAT, AP-1, and NF-κB (1), which

induces the transcription of Interleukin (IL)-2 and IL-2 receptor (R) α−chain

(*Il2ra*, CD25). IL-2 signalling induces further T cell activation, proliferation and

differentiation (2). In addition, IL-2 signalling has key roles in immunological

tolerance (2). This is partly mediated through CD25-expressing regulatory T

cells (Treg), which suppress the activities of other T cells (3). Intriguingly, TCR

signalling also induces the transient expression of FoxP3, the lineage-specific

transcription factor of Treg (4), in any T cells in humans (5), and in mice in the

presence of IL-2 and TGF-β (6). These suggest that FoxP3 can be actively

induced as a negative feedback mechanism for the T cell activation process,

especially in inflammatory conditions in tissues (7). Thus, the T cell activation

processes may dynamically control Treg phenotype and function during

immune response and homeostasis.

In fact, TCR signalling plays a critical role in Treg. Studies using TCR

transgenic showed that Treg require TCR activation for *in vitro* suppression

(8). Foxp3 binds to the enhancer regions that have been opened by TCR

signals, which explains a major part of the Treg-type chromatin structure (9).

In fact, continuous TCR signals are required for Treg function, because the

conditional deletion of the TCR-α chain in Treg abrogates the suppressive

activity of Treg and eliminates their activated or effector-Treg type phenotype

(10, 11). It is, however, unclear how TCR signals contribute to the Treg-type

transcriptional programme, and whether TCR signals are operating in all Treg

cells or whether these are required only when Treg suppress the activity of

other T cells.

Heterogeneity of Treg has been previously addressed through classifying

Treg into subpopulations, according to the origin (thymic Treg, peripheral

Treg, visceral adipose tissue Treg (12)), the transcription factor expression

and ability to control inflammation (Th1-Treg (13) and Th2-Treg (14), and T

follicular regulatory T cells (15)), and their activation status (activated Treg/

effector Treg (eTreg), resting Treg, and memory-type Treg (16)). Among these

Treg subpopulations, of interest is eTreg, which are activated and functionally

mature Treg. Murine eTreg can be identified by memory/ activation markers

such as CD44, CD62L, and GITR (16, 17), and their differentiation is

controlled by the transcription factors Blimp-1, IRF4 and Myb (18, 19). Human

Treg can be classified into naïve Treg (CD25+CD45RA+Foxp3+) and eTreg

(eTreg, CD25+CD45RA-Foxp3+) (20). However, our recent computational

study showed that classical gating approach is not effective for understanding

multidimensional data, and that marker expression data may be rather

effectively analysed by the computational clustering approaches that aim to

understand the dynamics of marker expressions in Treg (21). Furthermore,

the recent advancement of single cell technologies has opened the door to

address the heterogeneity of Treg by their gene profiles at the single cell

level.

When addressing the single cell level heterogeneity, it is critical to analyse

activated effector T cells (Teff) and memory-like T cells (memory-phenotype T

cells; Tmem) together with Treg. The surface phenotype of Tmem is

CD44^{high}CD45RB^{low}CD25$^-$ (22), which is similar to CD25$^-$ Treg, apart from

Foxp3 expression and suppressive activity (23, 24). In addition, activated

effector T cells (Teff) express CD25 and CTLA-4 (25), the latter of which is

also known as a Treg marker (26). While Tmem may include both antigen-

experienced memory T cells (27) and self-reactive T cells (28). In fact,

CD44^{high}CD45RB^{low} Tmem do not develop in TCR Tg mice with the *Rag*

deficient background, indicating that they require agonistic TCR signals in the

thymus (29). In addition, a study using a fate-mapping approach showed that

a minority of Treg naturally lose Foxp3 expression and join the the Tmem

fraction (30). These suggest that, upon encountering with cognate self-

antigens, self-reactive T cells, which include Tmem and Treg, express and

sustain Foxp3 expression as a negative feedback mechanism for strong TCR

signals (7). Thus, Treg have a close relationship with Tmem and Teff.

However, since most studies used naïve T cells (Tnaïve) as the control for

Treg, many of known "Treg" features may be in fact shared with Tmem and

Teff.

Multidimensional analysis is an effective approach to address this problem,

allowing to systematically investigate relationships between more than 2

populations (e.g. based on transcriptional similarities) (31). The prototype

methods include Principal Component Analysis (PCA), Correspondence

Analysis (CA) (32) and Multidimensional Scaling (33). In the application to

genomic data, these methods measure distances (i.e. similarities) between

samples and/or genes using different metrics, and thereby visualise the

relationships between samples and/or genes in a reduced dimension, typically

either in 2- or 3-dimensions, providing means to explore and investigate data

(31). However, these multidimensional methods are often not sufficiently

powerful for hypothesis-driven research, and our previous studies developed

a transcriptome analysis method using a variant analysis of CA, Canonical

Correspondence Analysis (CCA) for microarray data (31) and RNA-seq data

(34). In this approach, two transcriptome data are canonically analysed: the

correlations between cell samples in one dataset and the immunological

processes in another dataset are analysed based on their correlations to

individual genes. Briefly, CCA uses a linear regression to identify the

interpretable part (constrained space) of main data by explanatory variables,

and visualises similarities between genes, cells, and explanatory variables

using a singular value decomposition (SVD) solution within the interpretable

space (34). Thus, CCA enables to investigate and identify the unique features

of each T cell population, visualising the relationship between T cell

populations.

In this study, we investigate the multidimensional features of Treg in

comparison to other CD4+ T cells including Teff, Tmem, and naïve T cells

under normal or pathological conditions. Here we aim to identify the

differential regulation of transcriptional modules for T cell activation and

differentiation in these populations, and to reveal systems and molecular

mechanisms behind the differential regulation. Furthermore, using a new

single cell CCA approach, we investigate the single-cell level heterogeneity of

CD4+ T cells including Treg and effector T cells, identifying the differentially

regulated gene modules and the dynamic and gradational changes in

transcriptomes of individual T cells.

Materials and Methods

Conventional CCA (Gene-oriented analysis)

CCA canonically analyses two independent microarray or RNA-seq datasets (34). Briefly, gene expression data of the standardised main dataset (**S**) is linearly regressed onto the explanatory variable(s) (**D**), which identifies the interpretable part of the main dataset ("Constrained data", **S***). When only one explanatory variable is used, the CA algorithm of CCA assigns numerical values to cell samples and genes so that the dispersion of samples is maximised (uncorrelated information components), providing a one-dimensional solution (34). In order to use the solution as a scoring system, CCA score (i.e. Axis 1 score) was multiplied by the single biplot value, which indicates positive or negative correlation to Axis 1, which ensures that cells and genes with high scores have high positive correlations to the explanatory variable. When two or more explanatory variables are used, the CA algorithm then performs SVD on **S***, creating new matrices (i.e. sample and gene score matrices). These scores are sorted into new uncorrelated axes α_k, along which the entire set of scores generated by SVD, are distributed. The first axis holds the greatest amount of information (largest variations, precisely, *inertia*). The map approach enables the comparison of more than two explanatory variables, while the regression process in CCA allows the analysis across two different experiments (34). Biplot values of the CCA result are shown by arrows on the CCA map. CCA provides a map that shows the correlations between samples of interest, explanatory variables, and genes. Highly correlated components are closely positioned on the map.

In the application of CCA to population data, transcriptomic datasets of peripheral CD4+ T cells (including Treg, naïve, memory, draining LN and non-dLN and tissue effector CD4+ T cells) were processed by CCA and the cross-level relationships between components at three different levels, namely, immunological process, gene and cell, were analysed. Note that the same genes must be used in both transcriptome dataset matrices (intersect). The main dataset is projected onto the explanatory variable dataset, thus the genes in common to both datasets comprise the interpretable part of the main data. Mathematical operation implemented in the CCA algorithm produces immunological process (explanatory variable), gene and cell sample scores. The results are visualized as the 3-dimensional CCA solution on the CCA map (i.e. CCA triplot) that shows the relationships between cell subsets, genes and immunological processes.

Single cell data pre-processing and single cell CCA (single-cell oriented analysis)

RNA-seq expression data of GSE72056 was obtained from single-cell suspension of tumour cells with unknown activation and differentiation statuses were sequenced by RNA-seq (35). Genes with low variances and low maximal values were excluded. In order to identify CD4+ T cells, single cell data were filtered by the expression of *CD4* and *CD3E* to obtain only the *CD4+CD3E+* single cells, and also by k-means clustering of PCA gene plot to exclude outlier cells (21) for subsequent analysis.

In the application of CCA to single cell data, importantly, the same single cells

are used in both main data and the explanatory variables (i.e. selected

genes). The main dataset is projected onto the explanatory variables,

visualising the relationships between single cells, genes and explanatory

variables, which represent major activation/differentiation processes in the

dataset.

Explanatory variables for conventional CCA

Explanatory variables for CCA were prepared as follows. Differentially-

expressed genes were selected by a moderated t-test result using the

Bioconductor package, *limma*. The top-ranked differentially expressed genes

(according to their *p*-values) were used for making the explanatory variables.

The T cell activation explanatory variable were defined by the difference in

gene expression between anti-CD3/CD28-stimulated (17 h) CD4$^+$ T cells and

untreated CD4$^+$ T cells from GEO42276 (36). Precisely, genes were selected

by FDR <0.01 and log 2 fold change (> 1 or < -1) in the comparison of the

gene expression profile of the activated and resting T cells. For Figure 1A, the

expression data of GSE15907 (37) was regressed onto the log2 fold change

of activated CD4+ T cells (17 h after activation) and naïve CD4+ T cells from

GSE42276 as 'T cell activation signature' explanatory variable, and

Correspondence Analysis was performed for the regressed data and the

correlation analysis was done between the new axis and the explanatory

variable. For Figure 1C-D, the expression data of GSE15907 (37) was

regressed onto the 'T cell activation signature' explanatory variable described

above, in combination with retrovirally gene-transduced T cells for Foxp3 and

Runx1 effects (GSE6939 (38)): the log2 fold change of Foxp3-transduced and

empty vector-transduced CD4+T cells as the 'Foxp3 effects' explanatory

variable, and a log2 fold change of Runx1-transduced and empty vector-

transduced CD4+T cells as 'Runx1 effects' explanatory variable.

Correspondence Analysis was performed for the regressed data and the

correlation analysis was done between the new axis and the explanatory

variable.

Choice of explanatory variables by SC4A

SC4A aims to identify a set of genes that make the dispersion of cell

populations maximum in the CCA solution. To achieve this, all the

combinations of genes will be used as explanatory variables and tested for

discriminating each two populations using CCA. During each combinatorial

cycle, two genes are chosen from the total selected genes for all defined

single-cell populations in the main dataset and tested for their correlations to

one defined cell population vs all other T cells. In the analysis of Figure 8, the

following two cell populations were analysed by the combinatorial CCA: (1)

Activated T cells vs Resting T cells; (2) FOXP3+ cells vs FOXP3- cells; (3)

BCL6+ cells (as Tfh-like T cells) vs BCL6- cells. The most correlated gene to

each population (Activated T cells, Resting T cells, FOXP3+ cells, or BCL6+

cells) was identified, and these 4 genes were used as explanatory variables in

the final output of SC4A in Figure 8.

Data pre-processing and other statistical methods

All microarray datasets were downloaded from GEO site, and normalized,

where appropriate using the Bioconductor package *Affy*. Data were arranged

into an expression matrix where each row corresponds with gene expression

for each gene and each column corresponds with cell phonotype (sample).

Data were log2-transformed and values above log2(10) were used for

analysis. Differentially expressed genes (DEG) the TCR KO dataset and the

aTreg dataset were identified by a moderated t-statistics. DEG for activated

CD44hi and resting CD44lo Treg were combined. The CRAN package *vegan*

was used for the computation of CCA. Gene scores used the *wa* scores of the

CCA output by vegan. The Bioconductor package *limma* was used to perform

a moderated t-test. RNA-seq data were preprocessed, normalised, and log-

transformed using standard techniques (34).

Heatmaps were generated the CRAN package *gplots*. Venn diagram was

generated using the R code, overLapper.R, which was downloaded from the

Girke lab at Institute for Integrative Genome Biology

(http://faculty.ucr.edu/~tgirke/Documents/R_BioCond/My_R_Scripts/overLapp

er.R). Gene lists were compared for enriched pathways in the REACTOME

pathway database using the Bioconductor packages *ReactomePA* and

clusterProfiler. Violin plots were generated by the Bioconductor package

ggplot2. The inside part of the violin plot shows the median and interquartile

range (IQR) of the original gene expression data.   The lineage curve was

constructed by clustering SC4A/CCA sample scores using an expectation–

maximization (EM) algorithm (39), and the nodes of these clusters were

identified by constructing a minimum spanning tree using the Bioconductor

package *Slingshot* (40).

<u>Results</u>

Identification of the Foxp3-independent activation signature in Treg and

memory-phenotype T cells

Firstly, we investigated how T cell activation-related genes are differentially

regulated in resting Treg and other CD4+ T cell populations including Tmem

and Teff. To address this multidimensional problem, we applied CCA to the

microarray dataset of various CD4$^+$ T cells using the explanatory variable for

the T cell activation process, which was obtained from the microarray dataset

that analysed resting and activated conventional T cells ("T cell subset data"

and "T cell activation data" in **Table 1**). Thus, we aimed to visualise the cross-

level relationships between genes, the T cell populations, and the T cell

activation process (**Figure 1A**). Using the single explanatory variable, the T

cell activation process, the solution of CCA is one-dimensional and the cell

sample scores of CCA (represented by Axis 1) provides "T cell activation

score" (see Methods), indicating the level of activation in each cell population

relative to the prototype signature of T cell activation, as defined by the

explanatory variable *Tact*. All the naïve T cell populations had low Axis 1

values (i.e. Foxp3- T naïve cells (Tnaive); Tnaive, and non-draining lymph

node (dLN) T cells from BDC TCR transgenic (Tg) mice, which develop type I

diabetes). In contrast, Foxp3$^+$ Treg, Tmem, and tissue-infiltrating Teff in the

pancreas from BDC Tg (i.e. with inflammation in the islets) had high scores

(**Figure 1B**). These results indicate that Treg are as "activated" as Tmem and

tissue-infiltrating activated Teff at the transcriptomic level by CCA.

Next, we addressed whether the highly "activated" status of Treg is dependent

on Foxp3. Since Foxp3 suppresses Runx1-mediated transcriptional activities

(38), we investigated the same T cell population dataset using the following

three explanatory variables: T cell activation (Tact), retroviral *Foxp3*

transduction (Foxp3) and *Runx1* transduction (Runx1) (see Methods). The

CCA solution was 3-dimensional, while the first two axes explained the

majority of variance (98.8%, **Figure 2A**). As expected, Tmem, tissue-

infiltrating Teff and Treg had low negative values and showed high

correlations to T cell activation (Tact) in Axis 1, whereas only Treg had high

correlations with the Foxp3 variable in Axis 2, while Tmem and Teff were

correlated with the Runx1 variable in Axis 2 (**Figure 2A**). By analysing the

gene space of the CCA solution, genes in the lower left quadrant (i.e. negative

in both Axes 1 and 2) were enriched with the genes that are involved in T cell

activation, effector functions, and T follicular helper cells (Tfh), including

Cxcr5, *Pdcd1(*PD-1) *Il21, Ifng, Tbx21* (T-bet)*, Mki67* (Ki-67) (**Figure 2B**). On

the other hand, genes in the upper left quadrant (i.e. negative in Axis 1 and

positive in Axis 2) were enriched with Treg-associated genes including *Ctla4*,

Il2ra (CD25), *Itgae* (CD103), *Tnfrsf9* (4-1BB) and *Tnfrsf4* (OX40) (**Figure 2B**).

These results indicate that a set of activation genes are operating in all the

three non-naïve T cell populations (i.e. Treg, Teff and Tmem), while some of

them are more specific to Treg.

The Treg transcriptome is characterized by the repression of a part of the

activation genes in Tmem transcriptome

Next, we determine the modules of genes that are differentially regulated

between Treg and Tmem, in order to reveal the multidimensional identity of

Treg and Tmem transcriptomes. Specifically, we asked if the Axis 2 captured

the differential transcriptional regulations between Tmem and Treg.

Importantly, Axis 2 represents Foxp3-driven and Runx1-driven transcriptional

effects, which are correlated with Treg and Tmem/Teff, respectively (**Figure**

3A). This suggests that Axis 2 provides a 'scoring system' for regulatory vs

effector functions. Thus, the genes in Axis 1-low (precisely, genes above 25

percentile for positive correlations with Tact) were identified as *Tact genes*,

which were subsequently classified into Axis 2-positive (i.e. positive

correlations with Foxp3 and Treg) [designated as "*Tact-Foxp3 genes*"; top left

quadrant of CCA gene space in Figure 1D] and Axis 2-negative genes (i.e.

positive correlations with Runx1 and Tmem/Teff) [designated as "*Tact-Runx1*

genes"; bottom left quadrant of CCA gene space in Figure 1D] (**Figure 3A**).

Tact-Runx1 genes contain genes linked to T cell activation (e.g. *Mki67*),

effector functions (e.g. *Tbx21*), and Tfh (e.g. *Bcl6*, *Pdcd1*), while Tact-Foxp3

genes contain "Treg markers" such as *Il2ra* (CD25) and *Tnfrsf18* (GITR)

(**Figure 2B**). These encouraged us to further analyse the gene space of the

CCA solution, aiming to capture the unique regulation of activation genes in

Foxp3+ Treg in comparison to Foxp3- non-naive T cells which include Tmem

and Teff.

Intriguingly, heatmap analysis showed that both Treg and Tmem expressed

Tact-Foxp3 genes at high levels, compared to naïve and effector T cells

(**Figure 3B**). On the other hand, Tact-Runx1 genes were selectively

downregulated in Treg, while their expressions were sustained in Tmem

(**Figure 3C**). In other words, the repression of Tact-Runx1 genes was the

major feature of Treg in comparison to Tmem (**Figure 2A**). Interestingly,

comparable selective downregulation of Tact-Runx1 genes was observed in

Teff as well (**Figure 3C**). This suggests that the set of activation genes

operating in Teff is different from the ones in Tmem, and that Tmem and Treg

share more activation genes than Treg-Teff and Tmem-Teff (**Figure 3B and**

3C). These collectively indicate that the Treg-ness is composed of the

induction of the Treg-Tmem shared activation genes and the repression of

Tmem-specific genes, defining the multidimensional identity of Treg.

While the overall activation levels of Treg and Tmem are similar to the ones of

Teff at transcriptional level (Figure 1B), when explained by the prototype

signature of activation in CD4+ T cells (i.e. the explanatory variable Tact), the

compositions of the activation genes are different between Treg, Tmem and

Teff (as captured by Figure 3B and 3C). Importantly, many of these activation

genes are shared between Treg and Tmem, but not with Teff. The closer

similarity between resting Treg and Tmem, compared to Teff, is not surprising,

considering that both resting Treg and Tmem are considered to be at the

resting steady-state, while Teff are more recently activated and executing

effector functions, which presumably require unique sets of genes. These

features were not captured by standard t-test analysis (**Supplementary Fig**

1), presumably due to the lack of multidimensional perspective.

Tact-Foxp3 genes included the transcription factors *Nfat5*, *Runx2*, and *Ahr,*

which were expressed by most of Tmem cells as well (**Figure 3D**). The Treg-

associated markers, *Il2ra* (CD25), *Itgae* (CD103), and *Tnfrsf18* (GITR) were

expressed not only by Treg but also by Tmem at moderate to high levels.

Notably, the expression of *Ctla4, Ccr4, and Lag3* was high in Treg and Tmem

cells, but it was repressed in Teff (**Figure 3D**). This suggests that Treg and

Tmem are in later stages of T cell activation, when the expression of CTLA-4

is induced as a negative feedback mechanism (41), while it is not induced in

tissue-infiltrating Teff, presumably because they are more recently activated

and actively proliferating.

Tact-Runx1 genes included many cell cycle-related genes (e.g. *Ccna1*,

Cdca2, and *Chek2*), suggesting that these cells are in cell cycle and

proliferating (**Figure 3E**). The higher expression of *Mki67* and *Fos* suggests

that these Tmem cells had been activated by TCR signals *in vivo* before the

analysis. Tact-Runx1 genes also included the transcription factors *Tbx21*,

Maf, *Hif1a*, and *Bcl6*, which have roles in Th1, Th2, Th17, and Tfh

differentiation, respectively (42-44). In accordance with this, the Tfh markers

Cxcr5 and *Pdcd1* were specifically expressed by Tmem. These results are

compatible with the model that Treg and Tmem constitute the self-reactive T

cell population that have constitutive activation status (7), and that the major

function of Foxp3 is to modify the constitutive activation processes by

repressing a part of the activation gene modules (i.e. Tact-Runx1 genes)

(**Figure 4**).

The activated status of Treg is TCR signal dependent

We next asked whether the constitutively "activated" status of Treg is

dependent on TCR signals. We applied CCA to the microarray data of

CD44^{hi}CD62L^{lo} activated Treg (CD44^{hi} activated Treg) and CD44^{lo}CD62L^{hi}

naïve-like Treg (CD44^{lo} naïve Treg) from inducible *TCRa* KO or WT (TCR KO

data, **Table 1, Figure 5A**) using the T cell activation variable as explanatory

variable. The CCA result showed that CD44^{hi} activated Treg from WT mice

only showed high activation scores, compared with all the other groups.

Interestingly, *TCRa* KO CD44^{lo} naïve-like Treg showed the lowest scores, and

were lower than WT CD44^{lo} naïve-like Treg (**Figure 5B**). These results

indicate that TCR signaling is required for the constitutive activation status of

Treg, especially CD44^{hi} activated Treg, and suggest that these activated Treg

are more enriched with the cells that received TCR signals recently,

compared to CD44^{lo} naïve-like Treg.

In order to further address whether the TCR signal-dependent activation

signature of Treg is constitutively maintained or specifically induced by *in vivo*

activation events (presumably as tonic TCR signals (7)), we analysed the

RNA-seq dataset of *in vivo* activated Treg (Ref. (16), **Table 1**). The dataset

was generated by depleting a part of Treg by Diphtheria toxin (DT) using bone

marrow chimera of *Foxp3^{GFPCreERT2}:Rosa26YFP* and *Foxp3$^{GFP\ DTR}$* (16). The

DT treatment depletes DT receptor (DTR)-expressing Treg from *Foxp3$^{GFP\ DTR}$*,

and thus induces a transient inflammation through the reduction of Treg. Van

der Veeken *et al* thus analysed resting Treg from untreated mice (rTreg),

activated Treg from mice with recent depletion (11 days before the analysis)

in an inflammatory condition (aTreg), and "memory" Treg (mTreg) from mice

with a distant depletion (60 days before the analysis) (**Figure 5C**). As

expected, the CCA analysis using the T cell activation variable showed that

aTreg had higher activation scores than both rTreg and mTreg (**Figure 5D**).

This indicates that the activation mechanisms are more actively operating in

activated Treg in an inflammatory environment.

In order to further dissect the activation signature of Treg, we obtained the

lists of differentially expressed genes (DEG) between WT Treg vs *TCRa* KO

Treg (designated as *TCR-dependent genes*), and between aTreg and rTreg

(designated as *aTreg-specific genes*, see Methods). Interestingly, 94/286

genes of Tact-Runx1 genes (Tmem-specific activation genes, repressed in

resting Treg) are also used during the activation of Treg (**Figure 6A**), while

only 8/119 of Tact-Foxp3 genes (used by Tmem and resting Treg) are

induced during the activation of Treg (**Figure 6B**). This indicates that the

activation of Treg does not enhance the genes that are used in resting Treg,

but induces the expression of the Tmem-specific genes that are suppressed

in resting Treg. On the other hand, 51/286 of Tact-Runx1 and 19/119 of Tact-

Foxp3 genes are regulated by TCR signalling by TCRa KO (**Figure 6A and**

6B), suggesting that the activation status of resting Treg and Tmem may be

sustained by TCR signals. Pathway analysis showed that Tact-Runx1 and

aTreg-specific genes were enriched for cell-cycle related pathways. In

contrast, Tact-Foxp3 genes were enriched for pathways related to signal

transduction only (**Figure 6C**). Collectively, the results above suggest that

resting Treg are maintained by TCR and cytokine signalling, and that the

activation of Treg induces the transcriptional activities of Tact-Runx1 genes,

which promote proliferation and cell division.

Tumour-infiltrating FOXP3+ Treg are more enriched in activated T cells than

resting cells by single cell CCA

The analyses above led us to hypothesise that the activation status of Treg is

variable at the single cell level in physiological settings. In order to address

this predicted heterogeneity in Treg, we investigated single cell data and

further addressed how the activation mechanisms are operating in Treg at the

single cell level. In order to address the differential regulations of activation

mechanisms in individual Treg and related T cells, firstly, the features of

individual cells needed to be characterized in a data-oriented manner, as no

annotation data were available for individual single cells. CCA is a powerful

method for identifying biological meanings, and we applied CCA to the single

cell RNA-seq data of tumour-infiltrating T cells from human patients (Ref. (35)

Table 1).

Firstly, we applied the standard CCA to the single cell RNA-seq data of

CD4$^+$CD3$^+$ T cells (single-cell T cell samples of Treg and CD4+ non-Treg cells

with unknown individual activation and differentiation statuses), using the

explanatory variables of activated conventional CD4+ T cells (Tact) and

resting T cells (Trest; GSE15390, **Table 1**), aiming to define activated T cells

and resting T cells by the correlations to these two variables (**Figure 7A**) in

Axis 1. Here we used these two variables, Tact and Trest, instead of the log2

fold changes between the two populations (i.e. T cell activation variable,

which produces a 1-dimensional CCA solution visualized as a single axis),

because we aimed to identify an additional major differentiation process(es) in

the Axis 2 (i.e. two explanatory variables produce a 2-dimensional result). In

the single cell space of the CCA solution, the majority of FOXP3+ T cells had

negative gene scores in the Axis 1, i.e. showing high positive correlations to

the T cell activation variable (**Figure 7B**). Here, CCA Axis 1 x (-1) is

designated as the *Activation Score*. Thus, using the Axis 1 score and FOXP3

expression, the following 4 subpopulations were defined: "Activated FOXP3+",

"Resting FOXP3+", "Activated FOXP3-", and "Resting FOXP3-" (**Figure 7B**).

Next, we aimed to determine whether activated Treg are more activated than

resting Treg at the single cell level, or whether Treg are enriched with

activated T cells. Establishing the T cell activation score by Axis 1 score (as

the correlation to Tact in Figure 4B, see Methods), $FOXP3^+$ Treg had

significantly higher scores than FOXP3- non-Treg on average, as indicated by

the higher median in the violin plots and greater density of samples with

higher CCA gene scores for T cell activation variable (**Figure 7C**). Using the

CCA definition of Activated and Resting Treg and non-Treg in Figure 4B, the

activation score neatly captured the activated status of single cells, allocating

high positive and negative scores to Activated and Resting cells, respectively

(**Figure 7D**). Importantly, there was no significant difference between

Activated FOXP3+ and Activated FOXP3- cells and between Resting FOXP3+

and Resting FOXP3- cells (**Figure 7D**), indicating that in tumour

microenvironment, Treg cells are as activated as non-Treg CD4+ T cells,

which may include Teff, Tmem, and Tfh. These results suggest that the Treg

population have an activated signature because FOXP3+ cells are enriched

with T cells that have recognised antigens and received TCR signals (i.e.

activated T cells), and that TCR signals can induce FOXP3 in these antigen-

experienced T cells. Alternatively, but not exclusively, FOXP3+ T cells may

have high-affinity TCRs to self-MHC and/or tumour antigens and be more

prone to be activated. In fact, strikingly, 32.5% of activated T cells by the CCA

result expressed FOXP3, while only 8.2% of resting T cells expressed

FOXP3. In other words, FOXP3 expression occurred more frequently in

activated T cells.

In the gene space of the CCA solution, genes with strong correlations to

activated FOXP3$^+$ T cells included *FOXP3* itself and common Treg markers

such as *CTLA4* and *IL2RA* (CD25), which were found in the upper left

quadrant (Axis 1-negative Axis 2-positive). Interestingly, the lower left

quadrant (Axis 1-negative Axis 2-negative) contained more Tfh-like or

effector-like molecules *PDCD1* (PD-1), *BCL6*, *IL21*, and *IFNG.* The

chemokine receptors *CCR5* and *CCR2* had negative scores in Axis 1 (i.e.

correlated with Tact), while *CCR7* had a high positive score in Axis 1 (i.e.

correlated with Trest) (**Figure 7E**).

Identification of Tfh-like differentiation and Foxp3-driven processes and the

common activation process in tumour-infiltrating T cells

Next, we aimed to identify major differentiation and activation processes in the

single cell transcriptomes above. To this end, we have employed a new CCA

approach using Single cell analysis (Single Cell Combinatorial CCA, SC4A),

which aims to construct a CCA model of single cell data and thereby to

identify major differentiation/activation processes and the underlying gene

regulations (**Figure 8A,** see Methods and Supplementary Text). Firstly, we

classified single cells into Activated and Resting cells, and FOXP3+ Treg and

FOXP3- non-Treg, and thereby identified the following 4 processes as

putative differentiation and activation processes in the dataset: T cell

activation (Activated cells), and naïve-ness (Resting cells), FOXP3-driven

process (Activated FOXP3+), and Tfh-like process (Activated FOXP3-)

(Figure 4). Secondly, based on their high scores in the CCA solution (i.e.

either high positive or high negative scores in either Axis 1 or 2 in Figure 4E)

and abundant expressions in major populations (Figure 4F-4L), we selected

12 candidate genes (*CCR7*, *CCR5*, *CCR4*, *IL2RA*, *IL2RB*, *CTLA4*, *ICOS*,

TNFRSF4, *TNFRSF9*, *FOXP3*, *BCL6*, *PDCD1*) as the candidate genes for the

four processes. From these genes, we identified the most positively correlated

gene to each of the 4 processes using the combinatorial CCA (i.e. test all the

combinations of the variables by CCA and obtain the most correlated gene for

each population; see Method). Thus, *PDCD1, FOXP3*, *CTLA4*, and *CCR7*

were identified as the most correlated gene for Activated FOXP3-, Activated

FOXP3+, Activated T cells, and Resting T cells, respectively (**Supplementary**

Figure 3), which represent the four immunological processes (see above).

Finally, using these 4 genes as explanatory variable, we applied CCA to the

single cell transcriptomes, obtaining the solution of the SC4A approach.

The single cell space of the SC4A solution showed that Activated and Resting

T cells had negative and positive scores, respectively (**Figure 8B**). This

indicates that Axis 1 represents the T cell activation vs naïve-ness. Single

cells were successfully clustered into Activated FOXP3$^+$ Treg, Activated

FOXP3- non-Treg, and Resting T cells. Resting FOXP3+ Treg and Resting

FOXP3- T cells were mostly overlapped (**Figure 8C**), indicating that the major

features in the dataset dominated the difference between these two resting T

cell groups. Importantly, the explanatory variable CTLA4, which represents

the T cell activation process, was highly correlated with both Activated

FOXP3$^+$ Treg and Activated FOXP3- non-Treg at the middle, indicating its

neutral position in terms of Tfh and Treg activation processes. As expected,

the variable CCR7, which represents naïve-ness, was correlated with both

Resting FOXP3+ Treg and Resting FOXP3- T cells. The explanatory variable

PDCD1, which represents the Tfh-like process, was highly correlated with

Activated FOXP3- non-Treg cells, while the variable FOXP3 was correlated

with Activated FOXP3+ Treg. Thus, the single cell transcriptomes were

modelled by the correlations between gene expression, single cells, and the

expression of the 4 key genes, which represent the 4 immunological

processes (**Figure 8B and 8C**). Principal Component Analysis and t-

distributed stochastic neighbor embedding (t-SNE) did not provide insights

into such cross-level relationships or clear separations of the populations

(**Supplementary Figure 4**).

Next, in order to understand the relationship between the T cell activation

signature and FOXP3-driven and Tfh-like processes (**Figure 8C**), we aimed to

identify and characterise genes with high correlations to these processes,

which were represented by CTLA4, FOXP3, and PDCD1, by analysing the

gene space of the final output of SC4A (see Methods). As expected, the Tfh

genes, *IL21* and *BCL6* (45), were highly correlated with *PDCD1*. *IL2RA*

(CD25) is a Treg marker (46) and was highly correlated with *FOXP3*. *IL7R*

and *BACH2* are known to be associated with naïve T cells (47, 48), and were

positively correlated with CCR7, which represents the naïve-ness (**Figure**

8C). Thus we identified *FOXP3-driven Treg genes* and *Tfh-like genes*

according to their high correlation to the *FOXP3* and the *PDCD1* explanatory

variables, respectively, while we designated as *Activation genes* the genes

that have high correlations with the *CTLA4* variable, including *LAG3* and

CCR5, and were positioned around 0 in Axis 2.

Identification of the bifurcation point of activated T cells that leads to Tfh-like

and Treg differentiation in tumour-infiltrating T cells

The analyses above strongly suggested that there are two major

differentiation pathways for those tumour-infiltrating T cells, which are

regulated by FOXP3-driven and Tfh-like processes. In order to identify these

lineages, we applied an unsupervised clustering algorithm to the sample

space of the SC4A/CCA result (Figure 8C), and identified 6 clusters, to which

a pseudotime method (49) was applied, "lineage" curves were constructed

(**Figure 8D;** see **Methods**). Importantly, the lineage curves had a bifurcation

point at the Cluster 2, which leads to the two distinct differentiation pathways,

Tfh-like and FOXP3-driven differentiation. Since cells may change and mature

their phenotypes in different dynamics between these two lineages, we

designated Tfh-like-associated and FOXP3-associated pseudotime as Tfh-

pseudotime and FOXP3-pseudotime (**Figure 8D**).

In fact, the expression of Activation genes was progressively increased in the

common clusters for the two pseudotime (i.e. the Clusters 1 and 2) and

throughout the rest of the FOXP3-pseudotime and the early phase of Tfh-like

differentiation in Tfh-pseudotime (i.e. Cluster 3), while it was suppressed

towards the end of Tfh-pseudotime (Cluster 4; **Figure 8E**). Given that Tfh-

pseudotime is correlated with PDCD1 expression (**Figure 8C**), this suggests

that *PDCD1* expression and the Tfh-effector process is induced during earlier

phases, and that the activation processes in *PDCD1*^high T cells with are

suppressed, presumably through PD1-PDL1 interactions in the tumour

environment (50). Interestingly, FOXP3-driven genes had similar dynamics to

Activation genes in both FOXP3- and Tfh-pseudotime (**Figure 8F**). In

contrast, Tfh-like genes were mostly suppressed throughout FOXP3-

pseudotime, while they were progressively induced throughout Tfh-

pseudotime (**Figure 8G**). These differential regulations of two gene modules

resonate with those of Tact-Foxp3 (which are expressed by both Treg and

Tmem) and Tact-Runx1 genes (which are expressed specifically by Tmem

only) (Figure 3). In fact, FOXP3 expression is weakly induced in some cells in

the bifurcating Cluster 2, and is progressively increased at and beyond the

Cluster 5 (**Figure 8H**). In contrast, RUNX1 is highly expressed in the common

Clusters 1 and 2, while it is specifically suppressed in the early phase of

FOXP3-pseudotime (Cluster 5, **Figure 8I**). By analysing other key genes used

as CCA explanatory variables, both CTLA4 and PDCD1 were induced at the

bifurcating point, Cluster 2, and onwards in both of the lineages, while PDCD1

expression was specifically suppressed througohout FOXP3-pseudotime

(**Figure 8J and 8K**). CCR7 is highly expressed in the relatively naïve cells,

Cluster 1, and moderately downregulated at the bifurcation point, Cluster 2,

and suppressed beyond that in both FOXP3- and Tfh-pseudotime (**Figure**

8L).

These results collectively support the model that constant activation

processes in the tumour microenvironment promote terminal differentiation of

the Treg- and Tfh-like lineages in both previously committed and non-

committed lineages of T cells. Interestingly, the Cluster 2 is the bifurcation

point, in which T cells show moderate activation and are engaged in decision-

making about their cell fate. This understanding was possible because SC4A

effectively annotated genes and cells and thereby allowed to identify new cell

populations.

Identification of markers for the differential regulation of Tfh-like and Treg

differentiation in activated T cells

Lastly, we aimed to identify new combinations of marker genes in order to

demonstrate the strength of the current approaches. The SC4A identified the

two lineages of T cells, and their potential differentiation dynamics, FOXP3-

and Tfh-pseudotime: the FOXP3-driven pathway differentiates Treg via the

clusters 1-2-5-6, while the Tfh-like pathway differentiates Teff via the clusters

1-2-3-4 (Figure 9A). Since Activation genes (Figure 8C) are shared by early

phases of Tfh-like and FOXP3-driven differentiation (Figure 8E), we took the

intersect of these genes and the Tact-Foxp3 genes, which were expressed by

both resting Tmem and resting Treg in the immgen dataset (Figure 3). Thus,

we obtained *DUSP4* and *NFAT5*, which were in fact induced in cells at the

activated bifurcating Cluster 2 and onwards (**Figure 9B**). Similarly, in order to

identify Treg-specific genes, we took the intersect of FOXP3-driven genes

(Figure 8C) and the Tact-Foxp3 gene, and thereby obtained *CCR8* and

IL2RA. These genes were induced highly and progressively in Treg-lineage

cells throughout FOXP3-pseudotime, while suppressed across Tfh-

pseudotime (**Figure 9C**). Next, in order to identify activated non-Treg (Tfh-

like)-specific genes, we took the intersect of Tfh-like genes (Figure 8C) and

the Tact-Runx1 genes, which are expressed in resting Tmem but suppressed

in resting Treg (Figure 2). These genes contained *BCL6* and *KCNK5*, which

were progressively induced across Tfh-pseudotime, while suppressed in

FOXP3-pseudotime (**Figure 9D**).

Lastly, in order to make the newly obtained knowledge easily accessible to

experimental biologists, we showed the expression of *NFAT5*, *IL2RA*, *CCR8*,

BCL6, and *KCKN5* in the tumour-infiltrating T cells (**Figure 9E**). The common

activation gene *NFAT5+* in fact captured 43% of Treg-lineage cells (i.e. cells

in the Clusters 5 and 6) and 52% of Tfh-like-lineage cells (i.e. cells in the

Clusters 3 and 4). The Treg-specific genes *IL2RA* and *CCR8* were expressed

by the majority of T cells, whether NFAT-positive or negative. In contrast, the

Tfh-like-specific genes *BCL6* and *KCKN5* were expressed by a majority of

Tfh-like-lineage cells and were not expressed in Treg-lineage cells (**Figure**

9E). Collectively, these results indicate that the SC4A analysis successfully

decomposed the gene regulations for T cell activation and Treg and effector T

cell differentiation, identifying new cell populations, which include activated

cells at the bifurcation point, early and late phases of Treg and Tfh-like

differentiation, and their feature genes. In addition, although there must be

considerable differences between resting T cells in the secondary lymphoid

organs and between human and mice, our study successfully identified the

shared activation processes and the conserved genes that are differentially

used between the Treg- and the Teff-lineage cells.

1 Discussion

Resting Treg showed an activated status, comparable to that of Teff and

Tmem at the population level. In addition, the activation signature of Treg was

more remarkable in CD44^{hi}CD62L^{lo} activated Treg than CD44^{lo}CD62L^{hi} naïve-

like Treg. CD44^{hi}CD62L^{lo} Treg are also identified as eTreg, which may have

enhanced immunosuppressive activities (51). The eTreg fraction includes the

GITR^{hi}PD-1^{hi}CD25^{hi} "Triple-high" eTreg that express high CD5 and Nur77

expressions, which indicate that they have received strong TCR signals (17).

In humans, CD25^{hi}CD45RA^{-}FOXP3^{hi} eTreg highly express Ki67 (52),

indicating that these cells were recently activated. Given that TCRs of Treg

have higher affinities to self-antigens (53), these eTreg may have the most

self-reactive TCRs during homeostasis. Alternatively, the eTreg subset may

have immediately recently received strong TCR signals and upregulated

activation markers, and such cells may acquire a resting status at later time

points. Future investigations by TCR repertoire analysis will answer this

question.

Our study revealed the heterogeneity of FOXP3+ Treg at the single cell level,

and showed that tumour-infiltrating Treg include FOXP3+ T cells with various

levels of activation (Figure 4B and Figure 6B). It is plausible that, in the

physiological polyclonal settings, the variations in the activated status of

individual Treg may be due to the TCR affinity to its cognate antigen, the

availability of cognate antigen, and the strength and duration of TCR signals.

Our SC4A analysis identified the FOXP3-driven genes, which are specific to

activated FOXP3+ cells and include the IL-2 and common gamma chain

cytokine receptors (i.e. *IL2RA*, *IL2RB*, *IL15RA*, *IL4R*, and *IL2RG*), DNA

replication licensing factor (e.g. MCM2), and transcription factors such as

PRDM1 (BLIMP1) and *IRF4* (which control the differentiation and function of

eTreg (19)). These gene modules are distinct from the Tfh-like genes and the

activation genes (Figure 8), and may be controlled specifically by FOXP3

under strong TCR signals. /These genes expressions/ The expression of

these genes is variable /are various/ within the FOXP3+ T cells, suggesting

that the transcriptional activities of these genes are dynamically regulated

over time in tumour-infiltrating Treg. Thus, single cell-level analysis is

becoming a key technology to address the heterogeneity of Treg. To our

knowledge, this study is one of the first single cell analyses of Treg

transcriptomes, while we find that, during the review process of this

manuscript, another study addressing Treg heterogeneity by single cell RNA-

seq was deposited at a preprint server (54)).

The shared activation genes between activated FOXP3+ Treg and FOXP3-

non-Treg contain apoptosis-related genes (e.g. *CASP3*, *BAD*), which may be

differentially controlled between Treg and non-Treg at the protein level. For

example, activated FOXP3- non-Treg express *DUSP6* (Figure 8E), which is a

negative regulator of JNK-induced apoptosis through BIM activation, while

FOXP3 suppresses *DUSP6* expression and promote the apoptosis

mechanism (55). In addition, the activation genes include transcription factors

such as *TBX21* (T-bet) and *BATF*. Although *TBX21* is sometimes thought to

be a Th1-specific gene, it is upregulated immediately after T cell activation

(56). *BATF* was identified as a critical factor for the differentiation and

accumulation of tissue-infiltrating Treg (57). These activation genes may be

required when T cells are activated and differentiate into either Treg or Teff.

Further studies are required to investigate the temporal sequences of these

differentiation events *in vivo*.

Although the effects of TCR signals on Tmem were not directly examined,

considering that Tmem are self-reactive and their differentiation is dependent

on the recognition of cognate antigens in the thymus (7), these results

collectively suggest that the activation signature of Tmem is also dependent

on TCR signals, as is the signature of Treg (Figure 2F). Intriguingly, some

Treg may lose their Foxp3 expression and become ex-Treg, which are

enriched in $CD44^{hi}$ effector T cells or Tmem (30). In contrast, a Tmem

population (precisely, $Foxp3^-CD44^{hi}CD73^{hi}FR4^{hi}$ T cells) efficiently express

Foxp3 during lymphopenia (58). These findings support the feedback control

model that Foxp3 expression can be induced in Tmem and sustained in Treg

as a regulatory feedback mechanism for TCR signals (7). Given the variations

in the activated status in individual Treg and Tmem, single cell analysis will be

required to address this problem. For example, although Samstein *et al.*

showed that DNA hypersensitivity sites in Treg are similar to those in

activated T cells (9), it is possible that DNA hypersensitivity sites are variable

between individual Treg, and that Tmem may have a similar chromatin

structure to Treg.

Importantly, our analysis showed that Tmem-specific activation-induced

genes (i.e. Tact-Runx1 genes) are uniquely repressed in Treg. The repression

is likely to be mediated by the interaction between Foxp3 and other

transcription factors that regulate the expression of the Tmem-specific

activation genes (Figure 2F). Interestingly, Runx1 was associated with these

Tmem-specific genes. In fact, Foxp3 interacts with Runx1 and thereby

represses IL-2 transcription and controls the regulatory function of Treg (38),

and a significant part of the Foxp3-binding to active enhancers occurs through

the Foxp3-Runx1 interaction (9). These suggest that Runx1 may have a

unique role in the differentiation and maintenance of Tmem.

While CTLA-4 is commonly recognised as a Treg marker, it is upregulated in

all activated T cells, thus CTLA-4 is also a marker of activated T cells (41).

CTLA-4 is in fact expressed by only a subset of resting Treg (59), which may

be more activated and proliferating *in vivo* (60). In fact, our study shows that

CTLA-4 is expressed by non-Treg activated T cells including resting Tmem

(Figure 2D) and FOXP3- Tfh-like effector T cells in the tumour

microenvironment (Figure 4G and 6C). Our SC4A analysis also identified

CTLA-4 as a molecule representing the activation process of CD4+ T cells.

These findings support that CTLA-4 is primarily a marker for general T cell

activation, rather than Treg-specific marker, and that Treg are highly activated

T cells with Foxp3 and CTLA-4 expression. In order to address this problem,

the *in vivo* dynamics of CTLA-4 expression need to be investigated. We

anticipate that single cell analysis will reveal the dynamics of CTLA-4

expression and T cell activation levels in resting Treg and other activated T

cells *in vivo*.

In contrast, PD-1 (*PDCD1*) was specifically expressed by activated FOXP3-

non-Treg cells in the tumour microenvironment. The co-expression of *BCL6*

and *IL21* in some of these PD-1+ cells indicates that Tfh differentiation occurs

in the tumour microenvironment, presumably through the repeated and

chronic exposure to quasi-self antigens (i.e. tumour antigens). Interestingly,

the Tfh signature has been identified in type-I diabetes including mice and

humans (61). Intriguingly, the Tfh-like genes include cell-cycle related genes

(e.g. *CDK6*), immediate early transcription factors (*NFATC1*, *EGR2/3*), and

RNA-processing genes (e.g. *DICER1*). The significance of these gene

modules should be addressed in the future studies. It should be emphasised,

however, that PD-1 in the tumour microenvironment may constitute

immunoregulatory mechanisms as well, which prevent effective tumour

immunity (50). Further experimental investigations are required to address

this problem.

SC4A is a useful method to identify distinct clusters of T cells and the

correlated genes to each cluster, and thereby to reveal characteristic cell

groups and their active gene modules, while retaining the single-cell level

variations. We also showed that SC4A and CCA results can be further

analysed by the pseudotime approach. Since SC4A/CCA provides functional

annotations to cell groups and gene clusters, the understanding of the

pseudotime axis is effective, as shown in the current study. However, it is

emphasised that pseudotime is not a measurement of the time-dependent

events, but rather is that of similarities between samples (62), and the

conclusions in this study require future studies with a new experimental

system to analyse time-dependent events in vivo. In order to make the current

SC4A/CCA approach accessible, we visualised single cell data using a flow-

cytometry style (Figure 9). Although currently reliable antibodies are not

available for those intracellular candidate genes, and the expression of protein

and transcripts may not be synchronized, the current study showed that the

power of single cell RNA-seq and the current SC4A/CCA approach. The

current limitation of SC4A is that it is computationally expensive (i.e. requires

several hours for each analysis using a standard desktop), and the

improvement of the computational algorithm using a low-level language will be

beneficial. Importantly, SC4A is most effective when used together with in-

depth knowledge of immunology and gene regulation, which will facilitate the

interpretation of CCA results and explanatory variable selection. Thus, it is

hoped that these tools will be used widely by experimental immunologists with

good understandings of the biological significance of results, as well as

adequate competence in computational analysis, which will enable to ask

questions involving multidimensional problems such as multiple T cell

subsets.

Data and code availability

All R codes are available upon request. Processed data will be provided upon

reasonable requests to the corresponding author.

Acknowledgement

We thank Dr David Bending for valuable comments on the manuscript. M.O.

is a David Phillips Fellow (BB/J013951/2) from the Biotechnology and

Biological Sciences Research Council (BBSRC), and is also supported by a

pump-priming grant from Cancer Research Centre of Excellence, Imperial

College London/the Institute of Cancer Research. A.B is supported by the

Japanese Ministry of Education, Culture, Sports, Science and Technology

(MEXT) PhD scholarship.

References

1. Macian, F., C. Lopez-Rodriguez, and A. Rao. 2001. Partners in transcription: NFAT and AP-1. *Oncogene* 20: 2476-2489.

2. Malek, T. R. 2008. The biology of interleukin-2. *Annu Rev Immunol* 26: 453-479.

3. Setoguchi, R., S. Hori, T. Takahashi, and S. Sakaguchi. 2005. Homeostatic maintenance of natural Foxp3(+) CD25(+) CD4(+) regulatory T cells by interleukin (IL)-2 and induction of autoimmune disease by IL-2 neutralization. *The Journal of experimental medicine* 201: 723-735.

4. Hori, S. 2014. Lineage stability and phenotypic plasticity of Foxp3(+) regulatory T cells. *Immunological reviews* 259: 159-172.

5. Tran, D. Q., H. Ramsey, and E. M. Shevach. 2007. Induction of FOXP3 expression in naive human CD4+FOXP3 T cells by T-cell receptor stimulation is transforming growth factor-beta dependent but does not confer a regulatory phenotype. *Blood* 110: 2983-2990.

6. Chen, W., W. Jin, N. Hardegen, K.-j. Lei, L. Li, N. Marinos, G. McGrady, and S. M. Wahl. 2003. Conversion of Peripheral CD4$^+$CD25$^-$ Naive T Cells to CD4$^+$CD25$^+$ Regulatory T Cells by TGF-β Induction of Transcription Factor *Foxp3*. *The Journal of experimental medicine* 198: 1875-1886.

7. Ono, M., and R. J. Tanaka. 2016. Controversies concerning thymus-derived regulatory T cells: fundamental issues and a new perspective. *Immunology and cell biology* 94: 3-10.

8. Thornton, A. M., and E. M. Shevach. 2000. Suppressor Effector Function of CD4$^+$CD25$^+$ Immunoregulatory T Cells Is Antigen Nonspecific. *The Journal of Immunology* 164: 183-190.

9. Samstein, R. M., A. Arvey, S. Z. Josefowicz, X. Peng, A. Reynolds, R. Sandstrom, S. Neph, P. Sabo, J. M. Kim, W. Liao, M. O. Li, C. Leslie, J. A. Stamatoyannopoulos, and A. Y. Rudensky. 2012. Foxp3 exploits a pre-existent enhancer landscape for regulatory T cell lineage specification. *Cell* 151: 153-166.

10. Levine, A. G., A. Arvey, W. Jin, and A. Y. Rudensky. 2014. Continuous requirement for the TCR in regulatory T cell function. *Nature immunology* 15: 1070-1078.

11. Vahl, J. C., C. Drees, K. Heger, S. Heink, Julius C. Fischer, J. Nedjic, N. Ohkura, H. Morikawa, H. Poeck, S. Schallenberg, D. Rieß, Marco Y. Hein, T. Buch, B. Polic, A. Schönle, R. Zeiser, A. Schmitt-Gräff, K.

Kretschmer, L. Klein, T. Korn, S. Sakaguchi, and M. Schmidt-Supprian.
2014. Continuous T Cell Receptor Signals Maintain a Functional
Regulatory T Cell Pool. *Immunity* 41: 722-736.

12. Feuerer, M., J. A. Hill, K. Kretschmer, H. von Boehmer, D. Mathis, and
C. Benoist. 2010. Genomic definition of multiple ex vivo regulatory T
cell subphenotypes. *Proceedings of the National Academy of Sciences
of the United States of America* 107: 5919-5924.

13. Koch, M. A., G. Tucker-Heard, N. R. Perdue, J. R. Killebrew, K. B.
Urdahl, and D. J. Campbell. 2009. The transcription factor T-bet
controls regulatory T cell homeostasis and function during type 1
inflammation. *Nature immunology* 10: 595-602.

14. Zheng, Y., A. Chaudhry, A. Kas, P. deRoos, J. M. Kim, T. T. Chu, L.
Corcoran, P. Treuting, U. Klein, and A. Y. Rudensky. 2009. Regulatory
T-cell suppressor program co-opts transcription factor IRF4 to control
T(H)2 responses. *Nature* 458: 351-356.

15. Linterman, M. A., W. Pierson, S. K. Lee, A. Kallies, S. Kawamoto, T. F.
Rayner, M. Srivastava, D. P. Divekar, L. Beaton, J. J. Hogan, S.
Fagarasan, A. Liston, K. G. Smith, and C. G. Vinuesa. 2011. Foxp3+
follicular regulatory T cells control the germinal center response.
Nature medicine 17: 975-982.

16. van der Veeken, J., A. J. Gonzalez, H. Cho, A. Arvey, S. Hemmers, C.
S. Leslie, and A. Y. Rudensky. 2016. Memory of Inflammation in
Regulatory T Cells. *Cell* 166: 977-990.

17. Wyss, L., B. D. Stadinski, C. G. King, S. Schallenberg, N. I. McCarthy,
J. Y. Lee, K. Kretschmer, L. M. Terracciano, G. Anderson, C. D. Surh,
E. S. Huseby, and E. Palmer. 2016. Affinity for self antigen selects
Treg cells with distinct functional properties. *Nature immunology* 17:
1093-1101.

18. Dias, S., A. D'Amico, E. Cretney, Y. Liao, J. Tellier, C. Bruggeman, F.
F. Almeida, J. Leahy, G. T. Belz, G. K. Smyth, W. Shi, and S. L. Nutt.
2017. Effector Regulatory T Cell Differentiation and Immune
Homeostasis Depend on the Transcription Factor Myb. *Immunity* 46:
78-91.

19. Cretney, E., A. Xin, W. Shi, M. Minnich, F. Masson, M. Miasari, G. T.
Belz, G. K. Smyth, M. Busslinger, S. L. Nutt, and A. Kallies. 2011. The
transcription factors Blimp-1 and IRF4 jointly control the differentiation
and function of effector regulatory T cells. *Nature immunology* 12: 304-
311.

20. Sakaguchi, S., M. Miyara, C. M. Costantino, and D. A. Hafler. 2010.
FOXP3+ regulatory T cells in the human immune system. *Nature
reviews. Immunology* 10: 490-500.

21. Fujii, H., J. Josse, M. Tanioka, Y. Miyachi, F. Husson, and M. Ono. 2016. Regulatory T Cells in Melanoma Revisited by a Computational Clustering of FOXP3+ T Cell Subpopulations. *Journal of immunology*.

22. Andersen, P., and B. Smedegaard. 2000. CD4+ T-Cell Subsets That Mediate Immunological Memory to Mycobacterium tuberculosis Infection in Mice. *Infection and Immunity* 68: 621-629.

23. Coleman, M. M., C. M. Finlay, B. Moran, J. Keane, P. J. Dunne, and K. H. G. Mills. 2012. The immunoregulatory role of CD4+FoxP3+CD25− regulatory T cells in lungs of mice infected with Bordetella pertussis. *FEMS Immunology & Medical Microbiology* 64: 413-424.

24. Ono, M., J. Shimizu, Y. Miyachi, and S. Sakaguchi. 2006. Control of autoimmune myocarditis and multiorgan inflammation by glucocorticoid-induced TNF receptor family-related protein(high), Foxp3-expressing CD25+ and CD25- regulatory T cells. *Journal of immunology* 176: 4748-4756.

25. Linsley, P. S., J. L. Greene, P. Tan, J. Bradshaw, J. A. Ledbetter, C. Anasetti, and N. K. Damle. 1992. Coexpression and functional cooperation of CTLA-4 and CD28 on activated T lymphocytes. *The Journal of experimental medicine* 176: 1595-1604.

26. Attridge, K., and L. S. Walker. 2014. Homeostasis and function of regulatory T cells (Tregs) in vivo: lessons from TCR-transgenic Tregs. *Immunological reviews* 259: 23-39.

27. Bunce, C., and E. B. Bell. 1997. CD45RC Isoforms Define Two Types of CD4 Memory T Cells, One of which Depends on Persisting Antigen. *The Journal of experimental medicine* 185: 767-776.

28. Min, B., R. McHugh, G. D. Sempowski, C. Mackall, G. Foucras, and W. E. Paul. 2003. Neonates Support Lymphopenia-Induced Proliferation. *Immunity* 18: 131-140.

29. Curotto de Lafaille, M. A., A. C. Lino, N. Kutchukhidze, and J. J. Lafaille. 2004. CD25- T cells generate CD25+Foxp3+ regulatory T cells by peripheral expansion. *Journal of immunology* 173: 7259-7268.

30. Miyao, T., S. Floess, R. Setoguchi, H. Luche, H. J. Fehling, H. Waldmann, J. Huehn, and S. Hori. 2012. Plasticity of Foxp3(+) T cells reflects promiscuous Foxp3 expression in conventional T cells but not reprogramming of regulatory T cells. *Immunity* 36: 262-275.

31. Ono, M., R. J. Tanaka, M. Kano, and T. Sugiman. 2013. Visualising the Cross-Level Relationships between Pathological and Physiological Processes and Gene Expression: Analyses of Haematological Diseases. *PLoS One* 8: e53544.

32. Greenacre, M. 2008. *Correspondence Analysis in Practice.* Chapman & Hall/CRC,, London.

33. Tzeng, J., H. H.-S. Lu, and W.-H. Li. 2008. Multidimensional scaling for large genomic data sets. *BMC Bioinformatics* 9: 179.

34. Ono, M., R. J. Tanaka, and M. Kano. 2014. Visualisation of the T cell differentiation programme by Canonical Correspondence Analysis of transcriptomes. *BMC genomics* 15: 1028.

35. Tirosh, I., B. Izar, S. M. Prakadan, M. H. Wadsworth, 2nd, D. Treacy, J. J. Trombetta, A. Rotem, C. Rodman, C. Lian, G. Murphy, M. Fallahi-Sichani, K. Dutton-Regester, J. R. Lin, O. Cohen, P. Shah, D. Lu, A. S. Genshaft, T. K. Hughes, C. G. Ziegler, S. W. Kazer, A. Gaillard, K. E. Kolb, A. C. Villani, C. M. Johannessen, A. Y. Andreev, E. M. Van Allen, M. Bertagnolli, P. K. Sorger, R. J. Sullivan, K. T. Flaherty, D. T. Frederick, J. Jane-Valbuena, C. H. Yoon, O. Rozenblatt-Rosen, A. K. Shalek, A. Regev, and L. A. Garraway. 2016. Dissecting the multicellular ecosystem of metastatic melanoma by single-cell RNA-seq. *Science* 352: 189-196.

36. Wakamatsu, E., D. Mathis, and C. Benoist. 2013. Convergent and divergent effects of costimulatory molecules in conventional and regulatory CD4+ T cells. *Proceedings of the National Academy of Sciences of the United States of America* 110: 1023-1028.

37. Painter, M. W., S. Davis, R. R. Hardy, D. Mathis, C. Benoist, and C. Immunological Genome Project. 2011. Transcriptomes of the B and T lineages compared by multiplatform microarray profiling. *Journal of immunology* 186: 3047-3057.

38. Ono, M., H. Yaguchi, N. Ohkura, I. Kitabayashi, Y. Nagamura, T. Nomura, Y. Miyachi, T. Tsukada, and S. Sakaguchi. 2007. Foxp3 controls regulatory T-cell function by interacting with AML1/Runx1. *Nature* 446: 685-689.

39. Chen, W., and R. Maitra. 2017. EMCluster: EM Algorithm for Model-Based Clustering of Finite Mixture Gaussian Distribution. R Package.

40. Street, K., D. Risso, R. B. Fletcher, D. Das, J. Ngai, N. Yosef, E. Purdom, and S. Dudoit. 2017. Slingshot: Cell lineage and pseudotime inference for single-cell transcriptomics. *bioRxiv*.

41. Walker, L. S., and D. M. Sansom. 2015. Confusing signals: recent progress in CTLA-4 biology. *Trends Immunol* 36: 63-70.

42. Szabo, S. J., B. M. Sullivan, C. Stemmann, A. R. Satoskar, B. P. Sleckman, and L. H. Glimcher. 2002. Distinct Effects of T-bet in T$_H$1 Lineage Commitment and IFN-γ Production in CD4 and CD8 T Cells. *Science* 295: 338-342.

43. Dang, E. V., J. Barbi, H. Y. Yang, D. Jinasena, H. Yu, Y. Zheng, Z. Bordman, J. Fu, Y. Kim, H. R. Yen, W. Luo, K. Zeller, L. Shimoda, S. L. Topalian, G. L. Semenza, C. V. Dang, D. M. Pardoll, and F. Pan. 2011.

Control of T(H)17/T(reg) balance by hypoxia-inducible factor 1. *Cell* 146: 772-784.

44. Ho, I. C., M. R. Hodge, J. W. Rooney, and L. H. Glimcher. 1996. The proto-oncogene c-maf is responsible for tissue-specific expression of interleukin-4. *Cell* 85: 973-983.

45. Vinuesa, C. G., M. A. Linterman, C. C. Goodnow, and K. L. Randall. 2010. T cells and follicular dendritic cells in germinal center B-cell formation and selection. *Immunological reviews* 237: 72-89.

46. Sakaguchi, S., T. Yamaguchi, T. Nomura, and M. Ono. 2008. Regulatory T cells and immune tolerance. *Cell* 133: 775-787.

47. Seddon, B., and R. Zamoyska. 2002. TCR and IL-7 Receptor Signals Can Operate Independently or Synergize to Promote Lymphopenia-Induced Expansion of Naive T Cells. *The Journal of Immunology* 169: 3752-3759.

48. Roychoudhuri, R., K. Hirahara, K. Mousavi, D. Clever, C. A. Klebanoff, M. Bonelli, G. Sciume, H. Zare, G. Vahedi, B. Dema, Z. Yu, H. Liu, H. Takahashi, M. Rao, P. Muranski, J. G. Crompton, G. Punkosdy, D. Bedognetti, E. Wang, V. Hoffmann, J. Rivera, F. M. Marincola, A. Nakamura, V. Sartorelli, Y. Kanno, L. Gattinoni, A. Muto, K. Igarashi, J. J. O'Shea, and N. P. Restifo. 2013. BACH2 represses effector programs to stabilize T(reg)-mediated immune homeostasis. *Nature* 498: 506-510.

49. Lonnberg, T., V. Svensson, K. R. James, D. Fernandez-Ruiz, I. Sebina, R. Montandon, M. S. Soon, L. G. Fogg, A. S. Nair, U. Liligeto, M. J. Stubbington, L. H. Ly, F. O. Bagger, M. Zwiessele, N. D. Lawrence, F. Souza-Fonseca-Guimaraes, P. T. Bunn, C. R. Engwerda, W. R. Heath, O. Billker, O. Stegle, A. Haque, and S. A. Teichmann. 2017. Single-cell RNA-seq and computational analysis using temporal mixture modelling resolves Th1/Tfh fate bifurcation in malaria. *Sci Immunol* 2.

50. Okazaki, T., S. Chikuma, Y. Iwai, S. Fagarasan, and T. Honjo. 2013. A rheostat for immune responses: the unique properties of PD-1 and their advantages for clinical application. *Nature immunology* 14: 1212-1218.

51. Rosenblum, M. D., S. S. Way, and A. K. Abbas. 2016. Regulatory T cell memory. *Nature reviews. Immunology* 16: 90-101.

52. Miyara, M., Y. Yoshioka, A. Kitoh, T. Shima, K. Wing, A. Niwa, C. Parizot, C. Taflin, T. Heike, D. Valeyre, A. Mathian, T. Nakahata, T. Yamaguchi, T. Nomura, M. Ono, Z. Amoura, G. Gorochov, and S. Sakaguchi. 2009. Functional delineation and differentiation dynamics of human CD4+ T cells expressing the FoxP3 transcription factor. *Immunity* 30: 899-911.

53.　Weissler, K. A., and A. J. Caton. 2014. The role of T-cell receptor
recognition of peptide:MHC complexes in the formation and activity of
Foxp3(+) regulatory T cells. *Immunological reviews* 259: 11-22.

54.　Miragaia, R. J., T. Gomes, A. Chomka, L. Jardine, A. Riedel, A. N.
Hegazy, I. Lindeman, G. Emerton, T. Krausgruber, J. Shields, M.
Haniffa, F. Powrie, and S. A. Teichmann. 2017. Single cell
transcriptomics of regulatory T cells reveals trajectories of tissue
adaptation. *bioRxiv*.

55.　Tai, X., B. Erman, A. Alag, J. Mu, M. Kimura, G. Katz, T. Guinter, T.
McCaughtry, R. Etzensperger, L. Feigenbaum, D. S. Singer, and A.
Singer. 2013. Foxp3 transcription factor is proapoptotic and lethal to
developing regulatory T cells unless counterbalanced by cytokine
survival signals. *Immunity* 38: 1116-1128.

56.　Sahni, H., S. Ross, A. Barbarulo, A. Solanki, C. I. Lau, A. Furmanski, J.
I. Saldana, M. Ono, M. Hubank, M. Barenco, and T. Crompton. 2015. A
genome wide transcriptional model of the complex response to pre-
TCR signalling during thymocyte differentiation. *Oncotarget* 6: 28646-
28660.

57.　Hayatsu, N., T. Miyao, M. Tachibana, R. Murakami, A. Kimura, T. Kato,
E. Kawakami, T. A. Endo, R. Setoguchi, H. Watarai, T. Nishikawa, T.
Yasuda, H. Yoshida, and S. Hori. 2017. Analyses of a Mutant Foxp3
Allele Reveal BATF as a Critical Transcription Factor in the
Differentiation and Accumulation of Tissue Regulatory T Cells.
Immunity 47: 268-283 e269.

58.　Kalekar, L. A., S. E. Schmiel, S. L. Nandiwada, W. Y. Lam, L. O.
Barsness, N. Zhang, G. L. Stritesky, D. Malhotra, K. E. Pauken, J. L.
Linehan, M. G. O'Sullivan, B. T. Fife, K. A. Hogquist, M. K. Jenkins,
and D. L. Mueller. 2016. CD4(+) T cell anergy prevents autoimmunity
and generates regulatory T cell precursors. *Nature immunology* 17:
304-314.

59.　Takahashi, T., T. Tagami, S. Yamazaki, T. Uede, J. Shimizu, N.
Sakaguchi, T. W. Mak, and S. Sakaguchi. 2000. Immunologic self-
tolerance maintained by CD25(+)CD4(+) regulatory T cells
constitutively expressing cytotoxic T lymphocyte-associated antigen 4.
The Journal of experimental medicine 192: 303-310.

60.　Tang, A. L., J. R. Teijaro, M. N. Njau, S. S. Chandran, A. Azimzadeh,
S. G. Nadler, D. M. Rothstein, and D. L. Farber. 2008. CTLA4
expression is an indicator and regulator of steady-state CD4+ FoxP3+
T cell homeostasis. *Journal of immunology* 181: 1806-1813.

61.　Kenefeck, R., C. J. Wang, T. Kapadi, L. Wardzinski, K. Attridge, L. E.
Clough, F. Heuts, A. Kogimtzis, S. Patel, M. Rosenthal, M. Ono, D. M.
Sansom, P. Narendran, and L. S. Walker. 2015. Follicular helper T cell

1    signature in type 1 diabetes. *The Journal of clinical investigation* 125:
2    292-303.

3 62. Haghverdi, L., M. Buttner, F. A. Wolf, F. Buettner, and F. J. Theis.
4    2016. Diffusion pseudotime robustly reconstructs lineage branching.
5    *Nat Methods* 13: 845-848.
6

Figure Legends

Figure 1. Identification of the activation signature in Treg and Tmem by CCA of T cell populations

The microarray dataset of peripheral CD4+ T cells, including naïve, effector and memory phenotype from various sites (GSE15907), was analysed using the T cell activation variable, which was obtained by the microarray dataset of conventional activated CD4+ T cells (GSE42276). (**A**) Schematic representation of CCA for the cross-level analysis of T cell population (Cell), immunological process, and genes. CD4+ T cell populations and immunological processes. (**B**) CCA was applied to the T cell population data using an explanatory variable for T cell activation, which was obtained as fold change between activated and resting conventional CD4+ T cells. The CCA solution is thus one-dimensional, and is used as "T cell activation score" (see Methods).

Figure 2. Identification of the Foxp3-independent activation signature in Treg by CCA of T cell populations

The microarray dataset of peripheral CD4+ T cells (GSE15907) was analysed using the T cell activation variable and the variables for retroviral *Foxp3* transduction and *Runx1* transduction as explanatory variables. (**A**) The CCA solution was visualised by a biplot where CD4+ T cell samples are shown by closed circles (see legend) and the explanatory variables are shown by blue arrows. Percentage indicates that of the variance accounted for by the inertia

of the axes (i.e. the amount of information (eigenvalue) retained in each axis).

(**B**) Gene biplot of the 2D CCA solution in (C) showing the relationships

between genes (grey circles) and the explanatory variables (blue arrows).

Selected key genes are annotated.

Figure 3. Differential regulations of transcriptional modules for

activation in Treg and Tmem by Foxp3 and Runx1.

(**A**) Definition of Tact-Foxp3 genes and Tact-Runx1 genes. In the gene plot of

the CCA solution in Figure 1D, Axis 1-low genes (25 percentile low) were

designated as *activation genes*, which were further classified into Tact-Foxp3

genes and Tact-Runx1 genes by Axis 2, which have high correlations to Treg

and Tmem samples, respectively, in the CCA cell space (c.f. Figure 1C). (**B**)

Heatmap analysis of all the Tact-Foxp3 genes. (**C**) Heatmap analysis of all the

Tact-Runx1 genes. (**D**) Heatmap analysis of selected Tact-Foxp3 genes. (**E**)

Heatmap analysis of selected Tact-Runx1 genes.

Figure 4. A model for the differential regulation of activation genes in

Treg and Tmem.

The proposed differential regulations of TCR signal downstream genes in

Treg and Tmem. Since both naturally-arising Treg and Tmem are self-reactive

T cells, they may frequently receive tonic TCR signals by recognising their

cognate antigens in the periphery. This results in the full activation of both the

Tact-Foxp3 and Tact-Runx1 gene modules in Tmem. However, in Treg,

Foxp3 represses Tact-Runx1 genes and sustains the expression of Tact-

Foxp3 genes, producing the characteristic Treg transcriptome.

Figure 5. The activation signature of Treg is dependent on TCR

signalling.

(**A**) The experimental design for the TCR dataset. CD44hi activated Treg and

CD44lo naïve-like Treg were obtained from TCRa KO or WT mice and

analysed for transcriptome analysis. (**B**) CCA was applied to the

transcriptome data of CD44^{lo}CD62^{hi} naïve-like and CD44^{hi}CD62^{lo} activated

Treg cell populations from inducible *TCRa* KO or WT (from the *TCR KO* data,

GSE61077), using the T cell activation variable as the explanatory variable.

This produces a 1D CCA solution, and the sample score was plotted

(representing "T cell activation score"). (**C**) The experimental design for the

activated Treg dataset. Bone marrow (BM) cells were obtained from

Foxp3^{GFPCreERT2}:Rosa26YFP mice (YFP mice), and transferred into *Foxp3^{GFP}*

^{DTR} mice (Foxp3-DTR mice), in order to make BM chimera, in which ~10% of

Treg expressed DTR. Subsequently, DT was administered to these BM

chimera, which depleted Foxp3-DTR cells but not donor cells. This treatment

induced a transient activation of T cells and inflammation *in vivo*. Activated

Treg (aTreg) were obtained from these mice with inflammation, while resting

Treg (rTreg) were from control mice, and memory Treg (mTreg) were from the

mice after the resolution of inflammation. (**D**) 1D CCA sample score plot of

transcriptomic data of resting Treg (rTreg), *in vivo* activated Treg (aTreg) and

memory Treg (mTreg) from the *aTreg* data (GSE83315), with T cell activation

signature as explanatory variable.

Figure 6. The comparative analysis of Tmem-specific and Treg-Tmem

shared activation genes and TCR-dependent and activated Treg-specific

genes.

(**A**) Venn diagram analysis of TCR genes (DEGs between TCRa KO and WT

Treg), activated Treg genes (DEGs between aTreg and rTreg), and Tact-

Foxp3 and Tact-Runx1 genes (see Figure 3). (**B**) Pie chart showing the

number of the genes in the intersect between the TCR genes and either Tact-

Foxp3 or Tact-Runx1 genes, and that of the rest (designated as TCR-specific

genes). (**C**) Pie chart showing the numbers of aTreg gene-specific genes and

the intersects with Tact-Foxp3 and Tact-Runx1 genes (**D**) Pathway analysis of

the gene clusters in (C). The colours of texts correspond to the ones in the

Venn Diagram in (C). The gene clusters that are not shown did not have any

enrichment.

Figure 7. Single cell CCA of melanoma-infiltrating T cells determines the

activation status of individual T cells and identifies a putative Tfh-like

process

(**A**) Schematic representation of CCA of CD4+ T cell single cell

transcriptomes analysed by two explanatory variables: activated naïve T cells

(Tact) and resting naïve T cells (Trest). (**B**) CCA biplot showing the

relationship between Treg and non-Treg T cells (sample scores) and the

explanatory variables (Tact and Trest). Axis 1 represents the difference

between Tact and Trest, and thus, Activated T cells and Resting T cells were

defined by the CCA Axis 1 score, and these cells were further classified into

Treg and non-Treg by their *FOXP3* expression (see legend). Percentage

indicates that of the variance (inertia) accounted for by the axis. (**C**) Violin plot

showing the CCA activation scores (Axis 1 score $\times$ -1) of FOXP3- and

FOXP3+ cell groups. Asterisk indicates statistical significance by Mann-

Whitney test (**D**) Violin plot showing the CCA activation scores of Activated

(Act.) and Resting (Rest.) FOXP3- and FOXP3+ cell groups. Asterisks

indicate the values of post-hoc Dunn's test following a Kruskal Wallis test. ***

$p < 0.005$. (**E**) Gene biplot of the CCA solution in (B) showing the relationships

between genes (grey circles) and the Tact and Trest explanatory variables

(blue arrows). Genes are shown by grey circles, and well-known genes that

are key for T cell activation processes are annotated.

13
Figure 8. SC4A identifies the bifurcation point of activated T cells that

leads to Tfh-like and Treg differentiation in tumour-infiltrating T cells

SC4A was applied to the single cell data of tumour-infiltrating T cells, and 4

genes (CTLA-4, CCR7, FOXP3, and PDCD1) were chosen as explanatory

variables to represent the T cell activation, resting, FOXP3-driven process,

and Tfh-like process. (**A**) The design of analysis. The single cell data from the

melanoma sample were analysed by SC4A to identify the most effective

combination of explanatory variables for dispersing the 4 presumptive T cell

populations identified in Figure 7. These genes were used as explanatory

variables to analyse the rest of the single cell data as main dataset. Thus, the

single-cell level dynamics of T cell differentiation and activation are modelled

by the key biological processes that are represented by the T cell populations

and explanatory variables. (**B**) Single cell sample space of the final SC4A

output showing correlations between single cell samples and the explanatory

variables **(C)** Gene space of the final SC4A output showing correlations

between genes and the explanatory variables. The genes that showed high

correlations to the PDCD1, CTLA4, and FOXP3 variables were identified as

Tfh-like genes, Activation genes, and FOXP3-driven genes, respectively. **(D)**

The identification of two differentiation processes as lineages and a

bifurcation point. The cells in the sample space of the SC4A output (B) were

classified into 6 clusters by an unsupervised clustering algorithm. These

clusters were further analysed for pseudotime inference. (**E-G**) The average

gene expression was plotted against each pseudotime (upper: FOXP3-

pseudotime; lower: Tfh-pseudotime). The bifurcation point (Cluster 2) is

emphasised by broken lines. The number in circle indicate the cluster number.

Gene expression was standardised, and the sum of the standardised

expression was obtained for (**E**) Activation genes, (**F**) FOXP3-driven genes,

and (**G**) Tfh-like genes (see C). (H-L) The expression of key genes was

plotted against each pseudotime.

Figure 9. Identification of the conserved genes for the differential

regulation of Tfh-like and Treg differentiation in activated T cells

(**A**) The identified lineage curves and the bifurcation point in the tumour-

infiltrating T cells. The number in circle indicates the cluster number in Figure

8D. (**B-D**) The expression of selected feature genes was plotted against each

pseudotime. Genes are from the intersect of (**B**) Activation genes (Figure 8C)

and Tact-Foxp3 genes (Figure 3), (**C**) FOXP3-driven genes (Figure 8C) and

Tact-Foxp3 genes, and (**D**) Tfh-like genes (Figure 8C) and Tact-Runx1 genes

(Figure 3). (E) The expression of selected genes in the tumour-infiltrating T

cells were shown by a 2-dimensional plot in a flow cytometric style. Data from

Treg-lineage cells (the cluster 5 and 6, upper panels) and Tfh-like lineage

cells (the cluster 3 and 4, lower panels). The gene in x-axis (NFAT5) is from

the activation gene group (B), while y-axis shows genes from either the

FOXP3-Treg group (C) or the Tfh-like/Tmem group (D). Thresholds and

quadrant gates were determined in an empirical manner using density plot.

Accession number	Short description	Reference	Description of animal models	Timing of cell harvest	Cell purification strategy and sorting markers	Tissue origin
GSE15907	T cell subsets	Immunological Genome Project; Painter *et al.*, 2011	Primary cells from multiple immune lineages are isolated *ex-vivo*, primarily from young adult B6 male mice, and double-sorted to >99% purity.	6 weeks	FACS; Treg: Foxp3, CD4, CD25, GFP CD4 Tmem (scLN): TCRb CD44^{high} CD122^{lo} CD4 CD4 Tmem (spleen): CD4+ CD8- CD25- CD44^{hi} CD122^{lo} CD4 Tmem ( ) CD44^{hi}62L^{o} Activated Tmem44^{hi}62L^{lo} (spleen): CD4 TCRb CD44^{hi} CD62L^{lo} Naïve CD4 (scLN): CD4+ CD8- CD25- 62L^{hi} 44lo Naïve CD4 (mLN): CD4+ CD8- CD25- 62L^{hi} 44^{lo} Naïve CD4 (PP): TCR+ CD4+ CD44^{lo} CD62L^{hi} Naïve CD4 (spleen): CD4+ CD8- CD25- 62L^{hi} 44^{lo} CD4+ BDC (scLN): CD4+ CD8- BDC+ CD4+ BDC (pancreas): CD4+ CD8- BDC+ CD4+ BDC (pancreatic LN): CD4+ CD8- BDC+ Foxp3-Tnaive (spleen): CD4+ CD8- GFP- CD44^{lo}	Spleen, subcutaneous LNs, mesenteric LN, Peyer's patches

GSE83315	aTreg data	Van der Veeken *et al.*, 2016	Diphtheria toxin (DT) was administered to Foxp3^{GFP-DTR} mice to deplete Treg cells and induce expansion/activation of pro-inflammatory effector CD4+ T cells. After DT clearance, new Treg cells are generated. Mixed bone marrow chimeras generated with 90% Foxp3-GFP-DTR /10% Foxp3-GFP-CRE-ERT2XR26Y bone marrow were used to deplete 90% of Treg (thereby inducing inflammation and generation of *de novo* activated Treg) and irreversibly label 10% remaining inflammation-experienced Treg (mTreg).	Day 0 ('resting Treg), day 11 (activated Treg), day 60 (memory Treg)	FACS; CD4, Foxp3, CD44, CD62L, GFP	Spleen and peripheral lymph nodes
GSE61077	TCR KO data	Levine *et al.*, 2014	8-10 week mice from TracFL x Foxp3^{YFP-Cre} (tamoxifen-inducible deletion of TCR in Treg. Tamoxifen was administered on days 0 and 1.	Day 9 after tamoxifen administration	FACS TCRβ, CD4, Foxp3, CD44, CD62L	Lymph nodes
GSE42276	T cell activation	Wakamatsu *et*	Conventional CD4+ T cells	8 weeks	FACS;	Spleen, lymph node

		al., 2013	from C57BL/6J male mice were stimulated by anti-CD3 and anti-CD28 for 20h and 48h and data were pooled. 0h unstimulated samples were used as control.		DAPI−CD45R−CD8a−CD11b/c−CD4+ GFP+	
GSE6939	RV-transduced T cells	Ono *et al.*, 2007	Primary CD4+ T cells were obtained from BALB/c mice and purified into CD4+ naive T cells (GITRlowCD25-CD4+). CD4+ naive T cells were activated by… and retrovirally gene transduced with Runx1 (AML1), wild type Foxp3, and empty vector as control. Mice: BALB/c, ~8 weeks old. Cells: CD25-GITRlowCD4+ cells were isolated from lymph nodes and spleen, and stimulated by mitomycin-treated Thy1(-) splenocytes and anti-CD3 antibody. On the following day, cells were transfected by retrovirus carrying either Foxp3 or Runx1. Two days after transduction, GFP+	60 hours after transfection	FACS ; GFP	Lymph nodes and spleen

			cells were sorted..			
GSE72056	Single cell analysis of tumour-infiltrating T cells	Tirosh *et al*., 2016	Single cell RNA-seq analysis of human melanoma tumour samples. Freshly resected samples were disaggregated to generate single cell suspensions of mixed cells of unknown identities. Individual viable immune (CD45+) and nonimmune (CD45–) cells (including malignant and stromal cells) were recovered from the single cell suspension by flow cytometry. Single cells were profiled by single-cell RNA-seq.	Single cells were obtained within 45min of tumour resection	FACS; CD45	
GSE15390	Human activated T cells and Treg	Beyer *et al*., 2011	Conventional CD4+ T cells were obtained from whole blood of healthy human donors. T cells were activated for 5 days with CD3+CD28-coated beads; unstimulated naïve T cells as control.	5 days	MACS/FACS; CD4, CD25, CD127	

Figure 1

A

Gene-oriented CCA of T cell populations
and immunological processes

Two transcriptome data

→ Visualise relationships between genes,
cell populations, and immunological processes

B

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A

Tact-Runx1 genes (Tmem-specific)

64
21
30
171

- aTreg
- TCR
- TCR&aTreg
- Tact-Runx1-specific

B

Tact-Foxp3 genes (Treg-Tmem shared)

5
16
3
95

- aTreg
- TCR
- TCR&aTreg
- Tact-Foxp3-specific

C

NGF signalling via TRKA from the plasma membrane
Signaling by EGFR
Downstream signal transduction
Signalling by NGF
Downstream signaling events of B Cell Receptor (BCR)
Cell Cycle, Mitotic
Cell Cycle
M Phase
Metabolism of proteins
Post-translational protein modification

Tact-Foxp3
Tact-Runx1
aTreg

p.adjust
0.03
0.02
0.01

GeneRatio
0.1
0.2
0.3
0.4
0.5

Figure 7

Figure 8

Figure 9