Title: Elucidating the activation mechanisms for bifurcation

of regulatory and effector T cell fates by multidimensional single cell

analysis

Authors:

Alla Bradley[1], Tetsuo Hashimoto[1] and Masahiro Ono[2][§]

Affiliations:

[1] Faculty of Life and Environmental Sciences, University of Tsukuba, Tsukuba, Japan

[2] Department of Life Sciences, Faculty of Natural Sciences, Imperial College London, Sir Alexander Fleming Building, Imperial College Road, London, SW7 2AZ, United Kingdom

[§] Address correspondence and reprint requests to Dr Masahiro Ono, Department of Life Sciences, Sir Alexander Fleming Building, Imperial College Road, South Kensington, London, SW7 2AZ, United Kingdom. E-mail address: m.ono@imperial.ac.uk

1   **Abstract**
2
In T cells, T cell receptor (TCR) signalling initiates downstream transcriptional

mechanisms for T cell activation and differentiation. Foxp3-expressing

regulatory T cells (Treg) require TCR signals for their suppressive function

and maintenance in the periphery. It is, however, unclear how TCR signalling

controls the transcriptional programme of Treg. Since most of studies

identified the transcriptional features of Treg in comparison to naïve T cells,

the relationship between Treg and non-naïve T cells including memory-

phenotype T cells (Tmem) and effector T cells (Teff) is not well understood.

Here we dissect the transcriptomes of various T cell subsets from

independent datasets using the multidimensional analysis method Canonical

Correspondence Analysis (CCA). We show that resting Treg share gene

modules for activation with Tmem and Teff. Importantly, Tmem activate the

distinct transcriptional modules for T cell activation, which are uniquely

repressed in Treg. The activation signature of Treg is dependent on TCR

signals, and is more actively operating in activated Treg. Furthermore, by

analysing single cell RNA-seq data from tumour-infiltrating T cells, we

revealed that FOXP3 expression occurs predominantly in activated T cells.

Moreover, we identified FOXP3-driven and T follicular helper (Tfh)-like

differentiation pathways in tumour microenvironments, and their bifurcation

point, which is enriched with recently activated T cells. Collectively, our study

reveals the activation mechanisms downstream of TCR signals for the

bifurcation of Treg and Teff differentiation and their maturation processes.

1 Introduction

T cell receptor (TCR) signalling activates NFAT, AP-1, and NF-κB (1), which

induces the transcription of Interleukin (IL)-2 and IL-2 receptor (R) α−chain

(*Il2ra*, CD25). IL-2 signalling induces further T cell activation, proliferation and

differentiation (2). In addition, IL-2 signalling has key roles in immunological

tolerance (2). This is partly mediated through CD25-expressing regulatory T

cells (Treg), which suppress the activities of other T cells (3). Intriguingly, TCR

signalling also induces the transient expression of FoxP3, the lineage-specific

transcription factor of Treg (4), in any T cells in humans (5), and in mice in the

presence of IL-2 and TGF-β (6). These suggest that FoxP3 can be actively

induced as a negative feedback mechanism for the T cell activation process,

especially in inflammatory conditions in tissues (7). Thus, the T cell activation

processes may dynamically control Treg phenotype and function during

immune response and homeostasis.

In fact, TCR signalling plays a critical role in Treg. Studies using TCR

transgenic mice showed that Treg require TCR activation for *in vitro*

suppression (8). The binding of Foxp3 protein to chromatin occurs mainly in

the enhancer regions that have been opened by TCR signals (9). In fact,

continuous TCR signals are required for Treg function, because the

conditional deletion of the TCR-α chain in Treg abrogates the suppressive

activity of Treg and eliminates their activated or effector-Treg (eTreg)

phenotype (10, 11). It is, however, unclear how TCR signals contribute to the

Treg-type transcriptional programme, and whether TCR signals are operating

in all Treg cells or whether these are required only when Treg suppress the

activity of other T cells.

Heterogeneity of Treg has been previously addressed through classifying

Treg into subpopulations, according to the origin (thymic Treg, peripheral

Treg, visceral adipose tissue Treg (12)), the transcription factor expression

and ability to control inflammation (Th1-Treg (13) and Th2-Treg (14), and T

follicular regulatory T cells (15)), and their activation status (activated

Treg/eTreg, resting Treg, and memory-type Treg (16)). Among these Treg

subpopulations, of interest is eTreg, which are activated and functionally

mature Treg. Murine eTreg can be identified by memory/activation markers

such as CD44, CD62L, and GITR (16, 17), and their differentiation is

controlled by the transcription factors Blimp-1, IRF4 and Myb (18, 19). Human

Treg can be classified into naïve Treg (CD25+CD45RA+Foxp3+) and eTreg

(CD25+CD45RA-Foxp3+) (20). However, our recent computational study

showed that classical gating approach is not effective for understanding

multidimensional data, and that marker expression data may be rather

effectively analysed by the computational clustering approaches that aim to

understand the dynamics of marker expressions in Treg (21). Furthermore,

the recent advancement of single cell technologies has opened the door to

address the heterogeneity of Treg by their gene profiles at the single cell

level.

When addressing the single cell level heterogeneity, it is critical to analyse

activated effector T cells (Teff) and memory-like T cells (memory-phenotype T

cells; Tmem) together with Treg. The surface phenotype of Tmem is

$CD44^{high}CD45RB^{low}CD25^-$ (22), which is similar to $CD25^-$ Treg, apart from

Foxp3 expression and suppressive activity (23, 24). In addition, Teff express

CD25 and CTLA-4 (25), the latter of which is also known as a Treg marker

(26). Tmem may include both antigen-experienced memory T cells (27) and

self-reactive T cells (28). In fact, $CD44^{high}CD45RB^{low}$ Tmem do not develop in

TCR transgenic mice with the *Rag* deficient background, indicating that they

require agonistic TCR signals in the thymus (29). In addition, a study using a

fate-mapping approach showed that a minority of Treg naturally lose Foxp3

expression and join the Tmem fraction (30). These suggest that, upon

encountering cognate self-antigens, self-reactive T cells, which include Tmem

and Treg, express and sustain Foxp3 expression as a negative feedback

mechanism for strong TCR signals (7). Thus, Treg have a close relationship

with Tmem and Teff. However, since most studies used naïve T cells (Tnaïve)

as the control for Treg, many of known Treg-associated features may be in

fact shared with Tmem and Teff.

Multidimensional analysis is an effective approach to address this problem,

allowing to systematically investigate the relationships between more than two

cell populations (e.g. based on transcriptional similarities) (31). The prototype

methods include Principal Component Analysis (PCA), Correspondence

Analysis (CA) (32) and Multidimensional Scaling (33). In the application to

genomic data, these methods measure distances (i.e. similarities) between

samples and/or genes using different metrics, and thereby visualise the

relationships between samples and/or genes in a reduced dimension, typically

either in two-dimensional (2D) or three-dimensional (3D) space, providing

means to explore and investigate data (31). However, these multidimensional

methods are often not sufficiently powerful for hypothesis-driven research,

and our previous studies developed a transcriptome analysis method using a

variant of CA, Canonical Correspondence Analysis (CCA) for microarray data

(31) and RNA-seq data (34). In this approach, two transcriptome datasets are

canonically analysed: the correlations between cell samples in one dataset

(main dataset) and the immunological processes (explanatory variables) in

another dataset (explanatory dataset) are analysed based on their

correlations to individual genes. Briefly, CCA uses a linear regression to

identify the interpretable part (constrained space) of main data by explanatory

variables, and visualises similarities between genes, cells, and explanatory

variables using a singular value decomposition (SVD) solution within the

interpretable space (34). Thus, CCA enables to investigate and identify the

unique features of each T cell population, visualising the relationships

between T cell populations.

In this study, we investigate the multidimensional features of Treg in

comparison to other CD4+ T cells including Teff, Tmem, and naïve T cells

under normal or pathological conditions. Here we aim to identify the

differential regulation of transcriptional modules for T cell activation and

differentiation in these populations, and to reveal systems and molecular

mechanisms behind the differential regulation. Furthermore, using our new

single cell combinatorial CCA (SC4A) approach, we investigate the single-cell

level heterogeneity of CD4+ T cells including Treg and effector T cells,

identifying the differentially regulated gene modules and the bifurcation point

for T cell fates.

Materials and Methods

Conventional CCA (Gene-oriented analysis)

CCA canonically analyses two independent microarray or RNA-seq datasets

(34). Briefly, gene expression data of the standardised main dataset (**S**) is

linearly regressed onto the explanatory variable(s) (**D**), which identifies the

interpretable part of the main dataset ("Constrained data", **S***). When only one

explanatory variable is used, the CA algorithm of CCA assigns numerical

values to cell samples and genes so that the dispersion of samples is

maximised (uncorrelated information components), providing a one-

dimensional solution (34). The correlation analysis of explanatory data and

gene scores in the CA solution generates a biplot value, which, in the one-

dimensional solution, represents the explanatory variable score. In the one-

dimensional CCA solution, the single biplot value can either be +1 or -1,

indicating the direction (increasing/decreasing) of correlated genes in the

explanatory variable against that in the main dataset. In order to use the one-

dimensional solution as a scoring system, the CCA score (i.e. Axis 1 score) is

multiplied by the single biplot value, which indicates positive or negative

correlation to Axis 1, ensuring that cells and genes with high scores have high

positive correlations to the explanatory variable. When two or more

explanatory variables are used, the CA algorithm then performs SVD on **S***,

creating new matrices (i.e. sample and gene score matrices). These scores

are sorted into new uncorrelated axes α_k, along which the entire set of scores

generated by SVD is distributed. The first axis has the largest variation

(*inertia*) and thereby explains the greatest amount of information extracted by

the analysis. The map approach enables the comparison of two or more

explanatory variables, while the regression process in CCA allows the

analysis across two different experiments (34). Biplot values of the CCA result

are shown by arrows on the CCA map. CCA provides a map that shows the

correlations between samples of interest, explanatory variables, and genes.

Highly correlated components are closely positioned on the map.

Note that the same genes must be used in both transcriptome dataset

matrices. The main dataset is projected onto the explanatory variable dataset,

thus the genes in common to both datasets comprise the interpretable part

(intersect) of the main data. Mathematical operation implemented in the CCA

algorithm produces immunological process (explanatory variable), gene and

cell sample scores. The results are visualized as the 3-dimensional CCA

solution on the CCA map (i.e. CCA triplot) that shows the relationships

between cell subsets, genes and immunological processes. For example, in

the application of CCA to population-level (bulk) data, transcriptome datasets

of peripheral CD4+ T cells (including Treg, naïve, memory, draining LN and

non-dLN and tissue effector CD4+ T cells) were processed by CCA using

indicated explanatory variables (e.g. T cell activation) and the cross-level

relationships between components at three different levels (immunological

process, gene and cell) were analysed.

Single cell data pre-processing and single cell CCA (single-cell oriented

analysis)

RNA-seq expression data of GSE72056 was obtained from single-cell

suspension of tumour cells with unknown activation and differentiation

statuses (35). Genes with low variances and low maximal values were

excluded. In order to identify CD4+ T cells, single cell data were filtered by the

expression of *CD4* and *CD3E* to obtain only the *CD4+CD3E+* single cells, and

also by k-means clustering of PCA gene plot to exclude outlier cells (21) for

subsequent analysis.

In the application of CCA to single cell data, importantly, the same single cells

are used in both main data and the explanatory variables (i.e. selected

genes). The main dataset is projected onto the explanatory variables,

visualising the relationships between single cells, genes and explanatory

variables, which represent major activation/differentiation processes in the

dataset.

Explanatory variables for conventional CCA

Explanatory variables for CCA were prepared as follows. Differentially-

expressed genes were selected by a moderated t-test result using the

Bioconductor package, *limma*. The top-ranked differentially expressed genes

(according to their *p*-values) were used for making the explanatory variables.

The T cell activation explanatory variable (*Tact*) was defined by the difference

in gene expression between anti-CD3/CD28-stimulated (17 h) CD4$^+$ T cells

and untreated naïve CD4$^+$ T cells from GSE42276 (36). Precisely, genes were

selected by FDR < 0.01 and log2 fold change (> 1 or < -1) in the comparison

of the gene expression profile of the activated and resting T cells. For the one-

dimensional CCA of T cell populations (Figure 1B), the expression data of

GSE15907 (37) was regressed onto gene values in *Tact* representing the

change in gene expression following T cell activation, and CA was performed

for the regressed data and the correlation analysis was done between the

new axis and the explanatory variable. For the two-dimensional CCA of T cell

populations (Figure 2A and 2B), the expression data of GSE15907 (37) was

regressed onto *Tact*, in combination with *Foxp3* and *Runx1* explanatory

variables representing the effects of *Foxp3* and *Runx1* expression on CD4+ T

cells (GSE6939 (38)). *Foxp3* explanatory variable is the log2 fold change of

Foxp3-transduced naïve CD4+ T cells and empty vector-transduced CD4+T

cells. *Runx1* explanatory variable is the log2 fold change of *Runx1*-transduced

naïve CD4+ T cells and empty vector-transduced CD4+T cells. Subsequently,

CA was applied to the regressed data and the correlation analysis was

performed between the new axes and the explanatory variables.

SC4A

SC4A is a composite approach to understand non-annotated single cell data

by identifying distinct populations of cells and the differentiation processes

that are correlated with these populations. Since T cell population is usually

identified by a single lineage specification factor, in the application of SC4A,

such a factor will represent the cell population and their differentiation process

(**Supplementary Figure 1A**). The advantage of this approach is that SC4A

uses the gene expression data of a part of the dataset analysed, and thus the

regression analysis of CCA becomes more efficient because of the absence

of between-experimental variation, which is usually significant in cross-dataset

analysis (34).

SC4A is performed by the following three steps: (1) identification of putative

cell populations and candidate genes for explanatory variables by standard

CCA; (2) combinatorial CCA to identify the top-ranked genes to be used as

explanatory variables; and (3) the final CCA solution using the selected genes

as explanatory variables.

1. Preliminary analysis

The aim of the preliminary analysis is to identify the putative cell populations

and candidate genes for explanatory variables, and standard CCA is a useful

method to do this, because candidate genes can be identified by their

correlation to each putative cell population. Considering that the final output is

most effectively understood by visualisation using 2-dimensional (showing

correlation between explanatory variable, samples and genes) or 3-

dimensional (showing correlation between explanatory variable and

samples/genes) plot, up to 4 cell populations will be identified, and up to 5- 10

genes for each population will be identified by their correlation to the

population (**Supplementary Figure 1B**).

2. Combinatorial CCA

Here SC4A aims to identify a set of genes that make the dispersion of cell

populations maximum in the CCA solution. To achieve this, all the

combinations of genes will be used as explanatory variables and tested for

discriminating each two populations using CCA. During each combinatorial

cycle, two genes are chosen from the total selected genes for all defined

single-cell populations in the main dataset and tested for their correlations to

one defined cell population vs all other T cells.

Correlation can be visualised as the degree angle measured between the

explanatory variable (gene) and the centroid of the defined cell population.

Out of the two genes, the gene with the smallest angle to the defined cell

population is the most correlated. All selected genes are tested in this

pairwise manner against all defined cell populations vs all other T cells to

identify the gene that is most highly and uniquely correlated to each defined

cell population. At each combinatorial CCA, the most correlated gene to each

cell population is identified using vector multiplication (**Supplementary Figure**

1C). The top-ranked genes are determined by F1 score (the harmonic mean

of precision and sensitivity) and the correlation to the population of interest.

When the top-ranked gene is different between F1 score and the correlation,

the most immunologically-meaningful gene can be chosen.

3. SC4A

Finally, CCA is performed using the genes that are selected by the

combinatorial CCA to be used as explanatory variables. Thus, the single cell

dataset will be explained by the expression of the set of chosen genes, each

of which uniquely correlates with a cell population and represents the

differentiation process of the cell population (**Supplementary Figure 1D**).

Algebraically, SC4A is defined as follows. Single cell RNA-seq data $\mathbf{X} \in \mathbf{R}^{p \times}$

^{m} is the measurement of m genes from p single cells. The j-th column $x_j = (x_{1j}$

$x_{2j} \dots x_{pj})^T$ is the expression data of the j-th gene from p single cells, where T

indicates transposed vector. In SC4A, by choosing a set of k genes for

explanatory variable, $\mathbf{X'} \in \mathbf{R}^{p \times (m-k)}$ will be analysed by CCA using $\mathbf{Z} \in \mathbf{R}^{p \times k}$

as explanatory variables. As in the algorithm for CCA, $\mathbf{X'}$ is standardised by

column sums (c) and row sums (r), i.e. $\mathbf{S} = \mathbf{D_r}^{-1/2} (\textit{1/n } \mathbf{X} - \mathbf{rc}T) \mathbf{D_c}^{-1/2}$, where n

is the grand total of gene expression data, $\mathbf{D_r}$ and $\mathbf{D_c}$ are the diagonal

matrices of $\mathbf{r}$ and $\mathbf{c}$, respectively. Meanwhile, $\mathbf{Z}$ is scaled and standardised

(i.e. mean = 0 and variance = 1). $\mathbf{S}$ is linearly regressed onto $\mathbf{Z}$ by the

projection matrix $\mathbf{Q} = \mathbf{D_r}^{1/2} \mathbf{Z} (\mathbf{Z}^T \mathbf{D_r} \mathbf{Z})^{-1} \mathbf{Z}^T \mathbf{D_r}^{1/2}$, and the constrained space $\mathbf{S^*}$

$= \mathbf{Q S}$. Next, CCA performs singular value decomposition (SVD) of $\mathbf{S^*} = \mathbf{U} \mathbf{D_\alpha}$

$\mathbf{VT}$, where $\mathbf{UT U} = \mathbf{VT V} = \mathbf{I}$, and $\mathbf{D_\alpha}$ is the diagonal matrix of singular values in

descending order ($\alpha 1 \geq \alpha 2 \geq \dots$). Thus, SVD analyses the constrained space

and provides new axes where the dispersion of samples and that of genes are

maximised in the first axes. Gene scores are defined as $\mathbf{D_r}^{-1/2} \mathbf{U} \mathbf{D_\alpha}$ or $\mathbf{D_r}^{-1/2} \mathbf{U}$,

and sample scores for single cells are defined by weighted average scores

(WA scores) $\mathbf{S V D_\alpha}$, or $\mathbf{S V}$.

4. Choice of explanatory variables by SC4A

SC4A aims to identify a set of genes that make the dispersion of cell

populations maximum in the CCA solution. To achieve this, all the

combinations of genes will be used as explanatory variables and tested for

discriminating each two populations using CCA. During each combinatorial

cycle, two genes are chosen from the total selected genes for all defined

single-cell populations in the main dataset and tested for their correlations to

one defined cell population vs all other T cells. In the analysis of Figure 8, the

following two cell populations were analysed by the combinatorial CCA: (1)

Activated T cells vs Resting T cells; (2) FOXP3+ cells vs FOXP3- cells; (3)

BCL6+ cells (as Tfh-like T cells) vs BCL6- cells. The most correlated gene to

each population (Activated T cells, Resting T cells, FOXP3+ cells, or BCL6+

cells) was identified, and these 4 genes were used as explanatory variables in

the final output of SC4A in Figure 8.

Data pre-processing and other statistical methods

All microarray datasets were downloaded from GEO site, and normalized,

where appropriate using the Bioconductor package *Affy*. Data were arranged

into an expression matrix where each row corresponds with gene expression

for each gene and each column corresponds with cell phonotype (sample).

Data were log2-transformed and values above log2(10) were used for

analysis. Differentially expressed genes (DEG) the TCR KO dataset and the

aTreg dataset were identified by a moderated t-statistics. DEG for activated

CD44hi and resting CD44lo Treg were combined. The CRAN package *vegan*

was used for the computation of CCA. Gene scores used the *wa* scores of the

CCA output by vegan. The Bioconductor package *limma* was used to perform

a moderated t-test. RNA-seq data were preprocessed, normalised, and log-

transformed using standard techniques (34).

Heatmaps were generated the CRAN package *gplots*. Venn diagram was

generated using the R code, overLapper.R, which was downloaded from the

Girke lab at Institute for Integrative Genome Biology

(http://faculty.ucr.edu/~tgirke/Documents/R_BioCond/My_R_Scripts/overLapp

er.R). Gene lists were compared for enriched pathways in the REACTOME

pathway database using the Bioconductor packages *ReactomePA* and

clusterProfiler. Violin plots shows kernel density plots (outside) and the

median and interquartile range (inside) of the original gene expression data,

and were generated by the Bioconductor package *ggplot2*. The lineage curve

was constructed by clustering SC4A/CCA sample scores using an

expectation–maximization (EM) algorithm (39), and the nodes of these

clusters were identified by constructing a minimum spanning tree using the

Bioconductor package *Slingshot* (40).

Results

Identification of the Foxp3-independent activation signature in Treg and
memory-phenotype T cells

Firstly, we investigated how T cell activation-related genes are differentially
regulated in resting Treg and other CD4+ T cell populations including Tmem
and Teff. To address this multidimensional problem, we applied CCA to the
microarray dataset of various $CD4^+$ T cells using the explanatory variable for
the T cell activation process, which was obtained from the microarray dataset
that analysed resting and activated conventional T cells ("T cell subset data"
and "T cell activation data" in **Table 1**). Thus, we aimed to visualise the cross-
level relationships between genes, the T cell populations, and the T cell
activation process (**Figure 1A**). Using the single explanatory variable, the T
cell activation process, the solution of CCA is one-dimensional and the cell
sample scores of CCA (represented by Axis 1) provides "T cell activation
score" (see Methods), indicating the level of activation in each cell population
relative to the prototype signature of T cell activation, as defined by the
explanatory variable *Tact*. All the naïve T cell populations had low Axis 1
values (i.e. Foxp3- T naïve cells (Tnaive); Tnaive, and non-draining lymph
node (dLN) T cells from BDC TCR transgenic (Tg) mice, which develop type I
diabetes). In contrast, $Foxp3^+$ Treg, Tmem, and tissue-infiltrating Teff in the
pancreas from BDC Tg (i.e. with inflammation in the islets) had high scores
(**Figure 1B**). These results indicate that Treg are as "activated" as Tmem and
tissue-infiltrating activated Teff at the transcriptomic level by CCA.

Next, we addressed whether the highly "activated" status of Treg is dependent

on Foxp3. Since Foxp3 suppresses Runx1-mediated transcriptional activities

(38), we investigated the same T cell population dataset using the following

three explanatory variables: T cell activation (Tact), retroviral *Foxp3*

transduction (Foxp3) and *Runx1* transduction (Runx1) (see Methods). The

CCA solution was 3-dimensional, while the first two axes explained the

majority of variance (98.8%, **Figure 2A**). As expected, Tmem, tissue-

infiltrating Teff and Treg had low negative values and showed high

correlations to T cell activation (Tact) in Axis 1, whereas only Treg had high

correlations with the Foxp3 variable in Axis 2, while Tmem and Teff were

correlated with the Runx1 variable in Axis 2 (**Figure 2A**). By analysing the

gene space of the CCA solution, genes in the lower left quadrant (i.e. negative

in both Axes 1 and 2) were enriched with the genes that are involved in T cell

activation, effector functions, and T follicular helper cells (Tfh), including

Cxcr5, *Pdcd1(*PD-1) *Il21, Ifng, Tbx21* (T-bet)*, Mki67* (Ki-67) (**Figure 2B**). On

the other hand, genes in the upper left quadrant (i.e. negative in Axis 1 and

positive in Axis 2) were enriched with Treg-associated genes including *Ctla4*,

Il2ra (CD25), *Itgae* (CD103), *Tnfrsf9* (4-1BB) and *Tnfrsf4* (OX40) (**Figure 2B**).

These results indicate that a set of activation genes are operating in all the

three non-naïve T cell populations (i.e. Treg, Teff and Tmem), while some of

them are more specific to Treg.

The Treg transcriptome is characterized by the repression of a part of the

activation genes for Tmem

Next, we determine the modules of genes that are differentially regulated

between Treg and Tmem, in order to understand the multidimensional identity

of Treg and Tmem transcriptomes (i.e. how these populations can be defined

in comparison to all relevant populations). Specifically, we asked if the Axis 2

captured the differential transcriptional regulations between Tmem and Treg.

Importantly, Axis 2 represents Foxp3-driven and Runx1-driven transcriptional

effects, which are correlated with Treg and Tmem/Teff, respectively (**Figure**

3A). This suggests that Axis 2 provides a 'scoring system' for regulatory vs

effector functions. Thus, the genes in Axis 1-low (precisely, genes above 25

percentile for positive correlations with Tact) were identified as *Tact genes*.

These genes were subsequently classified into Axis 2-positive (i.e. positive

correlations with Foxp3 and Treg) [designated as "*Tact-Foxp3 genes*"; top left

quadrant of CCA gene space in Figure 1D] and Axis 2-negative genes (i.e.

positive correlations with Runx1 and Tmem/Teff) [designated as "*Tact-Runx1*

genes"; bottom left quadrant of CCA gene space in Figure 1D] (**Figure 3A**).

Tact-Runx1 genes contain genes linked to T cell activation (e.g. *Mki67*),

effector functions (e.g. *Tbx21*), and Tfh (e.g. *Bcl6, Pdcd1*), while Tact-Foxp3

genes contain "Treg markers" such as *Il2ra* (CD25) and *Tnfrsf18* (GITR)

(**Figure 2B**).

Intriguingly, heatmap analysis showed that both Treg and Tmem expressed

Tact-Foxp3 genes at high levels, compared to naïve and effector T cells

(**Figure 3B**). On the other hand, Tact-Runx1 genes were selectively

downregulated in Treg, while their expressions were sustained in Tmem

(**Figure 3C**). In other words, the repression of Tact-Runx1 genes was the

major feature of Treg in comparison to Tmem, and Tact-Foxp3 genes are the

activation genes, the expression of which is induced by T cell activation in

both Treg and Tmem, and is sustained or enhanced even in the presence of

Foxp3. Interestingly, comparable selective downregulation of Tact-Runx1

genes was observed in Teff as well (**Figure 3C**). This suggests that the set of

activation genes operating in Teff is different from the ones in Tmem, and that

Tmem and Treg share more activation genes than Treg-Teff and Tmem-Teff

(**Figure 3B and 3C**). These results collectively indicate that the Treg-ness is

composed of the induction of the Treg-Tmem shared activation genes (i.e.

Tact-Foxp3 genes) and the Foxp3-mediated repression of Tmem-specific

genes (i.e. Tact-Runx1 genes), defining the multidimensional identity of Treg.

While the overall activation levels of Treg and Tmem are similar to the ones of

the tissue-infiltrating Teff at transcriptional level (Figure 1B), when explained

by the prototype signature of activation in CD4+ T cells (i.e. the explanatory

variable Tact), the compositions of the activation genes are different between

Treg, Tmem and Teff (as captured by Figure 3B and 3C). Importantly, many

of these activation genes are shared between Treg and Tmem, but not with

Teff. The closer similarity between resting Treg and Tmem, compared to Teff,

is not surprising, considering that both resting Treg and Tmem are at the

resting status, while Teff are more recently activated and executing effector

functions. In addition, the distinct features of Teff may also include their

capacity of tissue infiltration and the effects of the microenvironment. These

features were not captured by standard t-test analysis (**Supplementary Fig**

1).

Tact-Foxp3 genes included the transcription factors *Nfat5*, *Runx2*, and *Ahr,*

which were expressed by most of Tmem cells as well (**Figure 3D**). The Treg-

associated markers, *Il2ra* (CD25), *Itgae* (CD103), and *Tnfrsf18* (GITR) were

expressed not only by Treg but also by Tmem at moderate to high levels.

Notably, the expression of *Ctla4, Ccr4, and Lag3* was high in Treg and Tmem

cells, but it was repressed in Teff (**Figure 3D**). This suggests that Treg and

Tmem are in later stages of T cell activation, when the expression of CTLA-4

is induced as a negative feedback mechanism (41), while it is not induced in

tissue-infiltrating Teff, presumably because they are more recently activated

and actively proliferating.

Tact-Runx1 genes included many cell cycle-related genes (e.g. *Ccna1*,

Cdca2, and *Chek2*), suggesting that these cells are in cell cycle and

proliferating (**Figure 3E**). The higher expression of *Mki67* and *Fos* suggests

that these Tmem cells had been activated by TCR signals *in vivo* before the

analysis. Tact-Runx1 genes also included the transcription factors *Tbx21*,

Maf, *Hif1a*, and *Bcl6*, which have roles in Th1, Th2, Th17, and Tfh

differentiation, respectively (42-44). In accordance with this, the Tfh markers

Cxcr5 and *Pdcd1* were specifically expressed by Tmem, suggesting that

Tmem are heterogenous populations and composed of these Th and Tfh

cells. These results are compatible with the model that Treg and Tmem

constitute the self-reactive T cell population that have constitutive activation

status (7), and that the major function of Foxp3 is to modify the constitutive

activation processes by repressing a part of the activation gene modules (i.e.

Tact-Runx1 genes) (**Figure 4**).

The activated status of Treg is TCR signal dependent

We next asked whether the constitutively "activated" status of Treg is

dependent on TCR signals. We applied CCA to the microarray data of

CD44^{hi}CD62L^{lo} activated Treg (CD44^{hi} activated Treg) and CD44^{lo}CD62L^{hi}

naïve-like Treg (CD44^{lo} naïve Treg) from inducible *Tcra* KO or WT (TCR KO

data, **Table 1, Figure 5A**) using the T cell activation variable as explanatory

variable. The CCA result showed that CD44^{hi} activated Treg from WT mice

only showed high activation scores, compared with all the other groups.

Interestingly, *TCRa* KO CD44^{lo} naïve-like Treg showed the lowest scores, and

were lower than WT CD44^{lo} naïve-like Treg (**Figure 5B**). These results

indicate that TCR signaling is required for the constitutive activation status of

Treg, especially CD44^{hi} activated Treg, and suggest that these activated Treg

are more enriched with the cells that received TCR signals recently,

compared to CD44^{lo} naïve-like Treg.

In order to further address whether the TCR signal-dependent activation

signature of Treg is constitutively maintained or specifically induced by *in vivo*

activation events (presumably as tonic TCR signals (7)), we analysed the

RNA-seq dataset of *in vivo* activated Treg (Ref. (16), **Table 1**). The dataset

was generated by depleting a part of Treg by Diphtheria toxin (DT) using bone

marrow chimera of *Foxp3*^{GFPCreERT2}:*Rosa26YFP* and *Foxp3*$^{GFP\ DTR}$ (16). The

DT treatment depletes DT receptor (DTR)-expressing Treg from *Foxp3*$^{GFP\ DTR}$,

and thus induces a transient inflammation through the reduction of Treg.

Foxp3^{GFPCreERT2} allows to label Foxp3-expressing cells by YFP at the moment

of tamoxifen administration. Van der Veeken *et al* thus analysed resting Treg

from untreated mice (rTreg), activated Treg from mice with recent depletion

(11 days before the analysis) in an inflammatory condition (aTreg), and

"memory" Treg (mTreg) from mice with a distant depletion (60 days before the

analysis) (**Figure 5C**). As expected, the CCA analysis using the T cell

activation variable showed that aTreg had higher activation scores than both

rTreg and mTreg (**Figure 5D**). This indicates that the activation mechanisms

are more actively operating in activated Treg in an inflammatory environment.

In order to further dissect the activation signature of Treg, we obtained the

lists of differentially expressed genes (DEG) between WT Treg vs *Tcra* KO

Treg (designated as *TCR-dependent genes*), and between aTreg and rTreg

(designated as *aTreg-specific genes*, see Methods). Interestingly, 94/286

genes of Tact-Runx1 genes (Tmem-specific activation genes, repressed in

resting Treg) are also used during the activation of Treg (**Figure 6A**), while

only 8/119 of Tact-Foxp3 genes (used by Tmem and resting Treg) are

induced during the activation of Treg (**Figure 6B**). This indicates that the

activation of Treg does not enhance the genes that are used in resting Treg,

but induces the expression of the Tmem-specific genes that are suppressed

in resting Treg. On the other hand, 51/286 of Tact-Runx1 and 19/119 of Tact-

Foxp3 genes are regulated by TCR signalling (**Figure 6A and 6B**),

suggesting that the activation status of resting Treg and Tmem may be

sustained by TCR signals. Pathway analysis showed that Tact-Runx1 and

aTreg-specific genes were enriched for cell-cycle related pathways. In

contrast, Tact-Foxp3 genes were enriched for pathways related to signal

transduction only (**Figure 6C**). Collectively, the results above suggest that

resting Treg are maintained by TCR and cytokine signalling, and that the

activation of Treg induces the transcriptional activities of Tact-Runx1 genes,

which promote proliferation and cell division.

FOXP3 expression more frequently occurs in activated T cells than resting

cells by single cell CCA

The analyses above showed that Treg are on average more activated than

naïve T cells and that the activation status of Treg can be variable. However,

it is still unclear whether individual Treg are activated than any naïve T cells at

the single cell level. The alternative hypothesis is that Treg are enriched with

the T cells that have recognised their cognate antigens and been activated. In

order to determine this and thereby understand the dynamics of T cell

regulation in vivo, we investigated the single cell RNA-seq data of tumour-

infiltrating T cells from human patients (Ref. (35) **Table 1**), and further

enquired how the activation mechanisms are operating in Treg at the single

cell level.

Firstly, we *in silico*-sorted FOXP3+ and FOXP3- CD4$^+$CD3$^+$ T cells from

unannotated single cell data from tumours, which tissues were dispersed and

CD45+ cells were sorted by flow cytometry without the use of any lymphocyte

markers (GSE72056, **Table 1**). Thus, the identities of individual single cells

were needed to be identified in a data-oriented manner, and Treg and non-

Treg cells in these tumour tissues had unknown individual activation and

differentiation statuses. Thus, we applied CCA to the *in silico*-sorted single

cell T cell data using the explanatory variables of activated conventional CD4+

T cells (*Tact*) and resting T cells (*Trest*; GSE15390, **Table 1**), aiming to define

individual single cells according to their level of activation by their correlations

to these two variables (**Figure 7A**). Here we used these two variables, *Tact*

and *Trest*, in order to generate a two-dimensional CCA solution, instead of a

single explanatory variable that represents T cell activation by the log2 fold

change in gene expression between activated and resting CD4+ T cells (*c.f.*

Figures 1-6), which produces a one-dimensional CCA solution visualised as a

single axis), because we aimed to identify any additional major differentiation

process(es) in the Axis 2. The explanatory variables *Tact* and *Trest* are both

captured by Axis 1 because they represent two poles of one continuum – the

spectrum of activation – ranging from 'resting' to 'activated' cell state. Thus

the CCA aimed to sort the single cells according to their individual levels of

activation along the spectrum of activation, capturing the heterogeneity in

activation levels in single cells. Remarkably, in the cell sample space of the

CCA solution, the majority of FOXP3+ T cells were positively correlated with

the T cell activation variable *Tact* (**Figure 7B**), and thus had negative scores

in the Axis 1 (**Figure 7B**). Here, CCA Axis 1 $\times$ (-1) score is designated as the

T cell activation score. Thus, using the activation score and FOXP3

expression, the following four subpopulations were defined: "Activated

FOXP3+", "Resting FOXP3+", "Activated FOXP3-", and "Resting FOXP3-"

(**Figure 7B**).

Next, we aimed to determine whether individual activated FOXP3+ Treg are

more activated than activated FOXP3- non-Treg at the single cell level

According to the T cell activation score established by the CCA solution in

Figure 7B, FOXP3$^+$ Treg had significantly higher T cell activation scores than

FOXP3- non-Treg on average, as indicated by the higher median in the violin

plots and greater density of samples with higher T cell activation scores

(**Figure 7C**), confirming the results by bulk cell analysis (Figure 1). Using the

CCA definition of activated and resting Treg and non-Treg established in

Figure 7B, the T cell activation score neatly captured the activated status of

single cells, allocating high positive and negative scores to activated and

resting cells, respectively (**Figure 7D**). Importantly, there was no significant

difference between Activated FOXP3+ and Activated FOXP3- cells and

between Resting FOXP3+ and Resting FOXP3- cells (**Figure 7D**), indicating

that in tumour microenvironment, Treg cells are as activated as non-Treg

CD4+ T cells, which may be enriched with Teff. Strikingly, 32.5% of activated

T cells expressed FOXP3, while only 8.2% of resting T cells expressed

FOXP3 in Figure 7B. In other words, FOXP3 expression occurred more

frequently in activated T cells. Given that the activation signature of Treg is

dependent on TCR signals (Figure 5), these results suggest that FOXP3

expression occurs predominantly in the activated T cells that have recognised

the tumour antigens and received TCR signals, as a negative feedback

mechanism to suppress the effector response against tumour antigen (7).

Alternatively, but not exclusively, FOXP3+ T cells may have high-affinity TCRs

to self-MHC and/or tumour antigens and be more prone to activation (10).

In the gene space of the CCA solution, genes with strong correlations to

activated FOXP3$^+$ T cells included *FOXP3* itself and common Treg markers

such as *CTLA4* and *IL2RA* (CD25), which were found in the upper left

quadrant (Axis 1-negative Axis 2-positive). Interestingly, the lower left

quadrant (Axis 1-negative Axis 2-negative) contained more Tfh-like or

effector-like molecules *PDCD1* (PD-1), *BCL6*, *IL21*, and *IFNG.* The

chemokine receptors *CCR5* and *CCR2* had negative scores in Axis 1 (i.e.

correlated with *Tact*), while *CCR7* had a high positive score in Axis 1 (i.e.

correlated with *Trest*) (**Figure 7E**).

Identification of Tfh-like differentiation and Foxp3-driven processes and the

common activation process in tumour-infiltrating T cells

Next, we aimed to identify major differentiation and activation processes in the

single cell transcriptomes above. To this end, we have developed a new CCA

approach for single cell analysis (Single Cell Combinatorial CCA, SC4A),

which aims to visualise major differentiation/activation processes and the

underlying gene regulations (**Figure 8A,** see Materials and Methods). Firstly,

we classified single cells into the four populations (Activated and Resting

cells, and FOXP3+ Treg and FOXP3- non-Treg; Figure 7B), and thereby

identified the following four processes as putative differentiation and activation

processes in the dataset: T cell activation (Activated cells), and naïve-ness

(Resting cells), FOXP3-driven process (Activated FOXP3+), and Tfh-like

process (Activated FOXP3-) (Figure 7). Secondly, based on their high scores

in the CCA solution (i.e. either high positive or high negative scores in either

Axis 1 or 2 in Figure 7E) and abundant expressions in FOXP3+ and FOXP3-

cells (data not shown), we selected 12 candidate genes (*CCR7, CCR5,*

CCR4, IL2RA, IL2RB, CTLA4, ICOS, TNFRSF4, TNFRSF9, FOXP3, BCL6,

PDCD1) as the candidate genes for the four processes. From these genes,

we identified the most positively correlated gene to each of the four processes

using the combinatorial CCA, which tests all the combinations of variables by

CCA and obtains the most correlated gene for each population; see Materials

and Methods). Thus, *PDCD1, FOXP3*, *CTLA4*, and *CCR7* were identified as

the most correlated genes for Activated FOXP3-, Activated FOXP3+,

Activated T cells, and Resting T cells, respectively (**Supplementary Figure**

3), which represent the four immunological processes (see above). Finally,

using these four genes as explanatory variables, we applied CCA to the single

cell transcriptomes, obtaining the solution of the SC4A approach.

The single cell space of the SC4A solution showed that Activated and Resting

T cells had negative and positive scores, respectively (**Figure 8B**). This

indicates that Axis 1 represents T cell activation vs naïve-ness. Single cells

were successfully clustered into Activated FOXP3$^+$ Treg, Activated FOXP3-

non-Treg, and Resting T cells. Resting FOXP3+ Treg and Resting FOXP3- T

cells were mostly overlapped (**Figure 8C**), indicating that the major features in

the dataset dominated the difference between these two resting T cell groups.

Importantly, the explanatory variable CTLA4, which represents the T cell

activation process, was highly correlated with both Activated FOXP3$^+$ Treg

and Activated FOXP3- non-Treg at the middle, indicating its neutral position in

terms of Tfh and Treg activation processes. As expected, the variable CCR7,

which represents naïve-ness, was correlated with both Resting FOXP3+ Treg

and Resting FOXP3- T cells. The explanatory variable PDCD1, which

represents the Tfh-like process, was highly correlated with Activated FOXP3-

non-Treg cells, while the variable FOXP3 was correlated with Activated

FOXP3+ Treg. Thus, the single cell transcriptomes were modelled by the

correlations between gene expression, single cells, and the expression of the

four key genes, which represent the four immunological processes (**Figure 8B**

and 8C). PCA and t-distributed stochastic neighbor embedding (t-SNE) did

not provide insights into such cross-level relationships or clear separations of

the populations (**Supplementary Figure 4**).

Next, in order to understand the relationship between the T cell activation

signature and FOXP3-driven and Tfh-like processes (Figure 8B and 8C), we

aimed to identify and characterise genes with high correlations to these

processes, which were represented by CTLA4, FOXP3, and PDCD1

explanatory variables, by analysing the gene space of the final output of

SC4A (Figure 8C; see Methods). As expected, the Tfh genes, *IL21* and *BCL6*

(45), were highly correlated with PDCD1 explanatory variable. *IL2RA* (CD25)

is a Treg marker (46) and was highly correlated with FOXP3 explanatory

variable. *IL7R* and *BACH2* are known to be associated with naïve T cells (47,

48), and were positively correlated with CCR7 explanatory variable, which

represents the naïve-ness (**Figure 8C**). Thus we defined *FOXP3-driven Treg*

genes (magenta circles) and *Tfh-like genes* (blue circles) according to their

high correlation to the FOXP3 and the PDCD1 explanatory variables,

respectively, while we designated as *Activation genes* (red circles) the genes

that have high correlations with the CTLA4 variable, including *LAG3* and

CCR5, which were positioned around 0 in Axis 2 (Figure 8C).

Identification of the bifurcation point of activated T cells that leads to Tfh-like

and Treg differentiation in tumour-infiltrating T cells

The analyses above strongly suggested that there are two major

differentiation pathways for those tumour-infiltrating T cells, which are

regulated by FOXP3-driven and Tfh-like processes. In order to identify these

lineages, we applied an unsupervised clustering algorithm to the sample

space of the SC4A/CCA result (Figure 8B), and identified 6 clusters, to which

a pseudotime method (49) was applied, constructing "lineage curves" (**Figure**

8D; see **Methods**). Importantly, the lineage curves had a bifurcation point at

Cluster II, which leads to the two distinct differentiation pathways, Tfh-like and

FOXP3-driven differentiation. Since cells may change and mature their

phenotypes in different dynamics between these two lineages, we designated

Tfh-like-associated and FOXP3-associated pseudotime as Tfh-pseudotime

and FOXP3-pseudotime (**Figure 8D**).

In fact, the expression of *Activation genes* was progressively increased in the

shared clusters (i.e. Cluster I and II) for the two pseudotimes, and throughout

the rest of the FOXP3-pseudotime and the early phase of Tfh-like

differentiation (i.e. Cluster III) in Tfh-pseudotime, while it was suppressed

towards the end of Tfh-like differentiation **(**Cluster IV; **Figure 8E**) in Tfh-

pseudotime. Given that Tfh-pseudotime is correlated with *PDCD1* expression

(**Figure 8C**), this suggests that *PDCD1* expression and the Tfh-effector

process are induced during the earlier phases of effector T cell activity, and

that the activation processes in $PDCD1^{high}$ T cells are suppressed,

presumably through PD1-PDL1 interactions in the tumour environment (50).

Interestingly, *FOXP3-driven genes* had similar dynamics to *Activation genes*

in both FOXP3-pseudotime and Tfh-pseudotime (**Figure 8F**). In contrast, *Tfh-*

like genes were mostly suppressed throughout FOXP3-pseudotime, while

they were progressively induced throughout Tfh-pseudotime (**Figure 8G**).

These differential regulations of two gene modules resonate with those of

Tact-Foxp3 genes (which are expressed by both Treg and Tmem) and Tact-

Runx1 genes (which are expressed specifically in Tmem, and repressed in

Treg) (Figure 3). In fact, *FOXP3* expression is weakly induced in some cells in

the bifurcating Cluster II and the early phase of Tfh-like differentiation (Cluster

III) in Tfh-pseudotime, and is progressively increased at and beyond Cluster V

in FOXP3-pseudotime (**Figure 8H**).

RUNX1 is highly expressed in the common Clusters I and II, and is

downregulated in the transition from Cluster II to Cluster III in Tfh-pseudotime,

and from Cluster II to Cluster V in FOXP3-pseudotime **(Figure 8I)**, which is

compatible with the known dynamics of RUNX1 expression: Runx1 is

downregulated when naïve CD4+ T cells differentiate into activated/effector

cells following TCR signaling (51). By analysing other key genes used as CCA

explanatory variables, *CTLA4* was induced at the bifurcating point, Cluster II,

and onwards in both of the lineages at equivalent expression levels (**Figure**

8J), reflecting the activated status of both effector Tfh-type cells and Treg.

Importantly, CTLA4 is a marker of Treg as well as activated effector T cells,

where it acts as a negative regulator of T cell proliferation (52).

PDCD1 expression was also induced at the bifurcating point, and throughout

Tfh-pseudotime, but specifically suppressed in the early phase of FOXP3-

pseudotime (**Figure 8K**), which is compatible with the known dynamics that

PDCD1 is transiently upregulated in activated CD4+ T cells as a negative

regulatory mechanism to restrain proinflammatory immune responses and

maintain peripheral tolerance (53). Further supporting this dynamic

perspective, *IL2* expression occurs mainly in Cluster II, indicating that these

cells are enriched with the T cells that recently recognised antigens (54)

(**Figure 8L**). Consistently, the expression of the naïve T cell marker CCR7

was the highest in the cells with a relatively naïve phenotype in shared Cluster

I, and was moderately downregulated in the early and late phase of Tfh-

pseudotime, and suppressed in most Treg in FOXP3-pseudotime (**Figure**

8M).

These results collectively support the model that constant activation

processes in the tumour microenvironment promote terminal differentiation of

the Treg- and Tfh-like lineages in both previously committed and non-

committed lineages of T cells. Interestingly, Cluster II is the bifurcation point,

in which T cells show moderate activation and together with simultaneous

expression of FOXP3 and Tfh-like genes, as well as RUNX1 and PDCD1

expression. These cells are most probably engaged in decision-making about

their cell fate and the cell type-specific usage of these genes – whether their

transcriptional mechanisms would be used to generate a proinflamamtory or

regulatory response. This understanding was possible because SC4A

effectively annotated genes and cells and thereby allowed to identify new cell

populations.

Identification of markers for the differential regulation of Tfh-like and Treg

differentiation in activated T cells

Lastly, we aimed to demonstrate the utility of the current approach by

discovering exemplary marker genes that distinguish cells in FOXP3- and Tfh-

pseudotime (i.e. the FOXP3-driven pathway I-II-V-VI, and the Tfh-like pathway

I-II-III-IV) (**Figure 9A**), and identifying the T cell subpopulations by a flow

cytometric visualisation of single cell data. Since *Activation genes* (Figure 8C)

are shared by early phases of Tfh-like and FOXP3-driven differentiation

(Figure 8E), we took the intersect of these genes and the Tact-Foxp3 genes,

which were expressed by both resting Tmem and resting Treg in mice (Figure

3). *DUSP4* and *NFAT5* were such genes and in fact induced in cells at the

activated bifurcating Cluster II and onwards in both lineages (**Figure 9B**).

Similarly, in order to identify a marker to distinguish Treg and Tfh-like cells,

firstly, we identified *CCR8* and *IL2RA* in the intersect of FOXP3-driven genes

(Figure 8C) and the Tact-Foxp3 genes, which were induced highly and

progressively in Treg-lineage cells throughout FOXP3-pseudotime, while

mostly suppressed across Tfh-pseudotime (**Figure 9C**). In contrast, *BCL6* and

KCNK5 (found in the intersect of Tfh-like genes (Figure 8C) and the Tact-

Runx1 genes, which are expressed in resting Tmem but suppressed in resting

Treg (Figure 3)) were progressively induced across Tfh-pseudotime, while

suppressed in FOXP3-pseudotime (**Figure 9D**).

Lastly, in order to make the newly obtained knowledge easily accessible to

experimental immunologists, we showed the expression of *NFAT5*, *IL2RA*,

CCR8, *BCL6*, and *KCNK5* in the tumour-infiltrating T cells in a flow cytometric

format (**Figure 9E**). The common activation gene *NFAT5* in fact captured the

majority of Treg-lineage cells (i.e. cells in the Clusters V and VI) and Tfh-like-

lineage cells (i.e. cells in the Clusters III and IV). The Treg-specific genes the

expression of *IL2RA* and *CCR8* occurred in the majority of FOXP3+ Treg-

lineage cells, whether NFAT-positive or negative, but not in most of Tfh-like-

lineage cells. In contrast, the Tfh-like-specific genes *BCL6* and *KCNK5* were

expressed by a majority of Tfh-like-lineage cells and were not expressed in

Treg-lineage cells (**Figure 9E**).

Collectively, these results indicate that the SC4A analysis successfully

decomposed the gene regulations for T cell activation and Treg and effector T

cell differentiation, identifying new cell populations, which include activated

cells at the bifurcation point, early and late phases of Treg and Tfh-like

differentiation, and their feature genes. In addition, although there must be

considerable differences between resting T cells in the secondary lymphoid

organs and between humans and mice, our study successfully identified the

shared activation processes and the conserved genes that are differentially

used between the Treg- and the Teff-lineage cells, identifying a shared

systems-level mechanism for the differentiation regulation of activation and

differentiation processes in CD4+ T cell populations.

Discussion

Resting Treg showed an activated status, comparable to that of Teff and Tmem at the population level. In addition, the activation signature of Treg was more remarkable in CD44^{hi}CD62L^{lo} activated Treg than CD44^{lo}CD62L^{hi} naïve-like Treg. CD44^{hi}CD62L^{lo} Treg are also identified as eTreg, which may have enhanced immunosuppressive activities (55). The eTreg fraction includes the GITR^{hi}PD-1^{hi}CD25^{hi} "Triple-high" eTreg that have high CD5 and Nur77 expressions, which indicates that they have received strong TCR signals (17). In humans, CD25^{hi}CD45RA^{-}FOXP3^{hi} eTreg highly express Ki67 (56), indicating that these cells were recently activated. Given that TCRs of Treg have higher affinities to self-antigens (57), these eTreg may have the most self-reactive TCRs during homeostasis. Alternatively, the eTreg subset may have recently received strong TCR signals and upregulated activation markers, and such cells may acquire a resting status at later time points. Future investigations by TCR repertoire analysis will answer this question.

Our study revealed the heterogeneity of FOXP3+ Treg at the single cell level, and showed that tumour-infiltrating Treg include FOXP3+ T cells with various levels of activation (Figure 7B and Figure 8C). It is plausible that, in the physiological polyclonal settings, the variations in the activated status of individual Treg may be due to the TCR affinity to its cognate antigen, the availability of cognate antigen, and the strength and duration of TCR signals. Our SC4A analysis identified the *FOXP3*-driven genes, which are specific to activated FOXP3+ cells and include IL-2 and common gamma chain cytokine receptors (i.e. *IL2RA*, *IL2RB*, *IL15RA*, *IL4R*, and *IL2RG*), DNA replication

licensing factors (e.g. MCM2), and transcription factors such as *PRDM1*

(BLIMP1) and *IRF4* (which control the differentiation and function of eTreg

(19)). These gene modules are distinct from the Tfh-like genes and the

activation genes (Figure 8), and may be controlled specifically by *FOXP3*

under strong TCR signals. The expression of these genes is variable within

the FOXP3+ T cells, suggesting that the transcriptional activities of these

genes are dynamically regulated over time in tumour-infiltrating Treg. Thus,

single cell-level analysis is becoming a key technology to address the

heterogeneity of Treg. To our knowledge, this study is one of the first single

cell analyses of Treg transcriptomes, while we find that, during the review

process of this manuscript, another study addressing Treg heterogeneity by

single cell RNA-seq was deposited at a preprint server (58)).

The shared activation genes between activated FOXP3+ Treg and FOXP3-

non-Treg contain apoptosis-related genes (e.g. *CASP3*, *BAD*), which may be

differentially controlled between Treg and non-Treg at the protein level. For

example, activated FOXP3- non-Treg express *DUSP6* (Figure 9B), which is a

negative regulator of JNK-induced apoptosis through BIM activation, while

FOXP3 suppresses *DUSP6* expression and promotes the apoptosis

mechanism (59). In addition, the activation genes include transcription factors

such as *TBX21* (T-bet) and *BATF*. Although *TBX21* is sometimes thought to

be a Th1-specific gene, it is upregulated immediately after T cell activation

(60). *BATF* was identified as a critical factor for the differentiation and

accumulation of tissue-infiltrating Treg (61). These activation genes may be

required when T cells are activated and differentiate into either Treg or Teff.

Further studies are required to investigate the temporal sequences of these

differentiation events *in vivo*.

Although the effects of TCR signals on Tmem were not directly examined,

considering that Tmem are self-reactive and their differentiation is dependent

on the recognition of cognate antigens in the thymus (7), these results

collectively suggest that the activation signature of Tmem is also dependent

on TCR signals, as is the activation signature in Treg (Figure 5B). Intriguingly,

some Treg may lose their Foxp3 expression and become ex-Treg, which are

enriched in $CD44^{hi}$ effector T cells or Tmem (30). In contrast, a Tmem

population (precisely, $Foxp3^-CD44^{hi}CD73^{hi}FR4^{hi}$ T cells) efficiently express

Foxp3 during lymphopenia (62). These findings support the feedback control

model that Foxp3 expression can be induced in Tmem and sustained in Treg

as a regulatory feedback mechanism for TCR signals (7). Given the variations

in the activated status in individual Treg and Tmem, single cell analysis will be

required to address this problem. For example, although Samstein *et al.*

showed that DNA hypersensitivity sites in Treg are similar to those in

activated T cells (9), it is possible that DNA hypersensitivity sites are variable

between individual Treg, and that Tmem may have a similar chromatin

structure to Treg.

Importantly, our analysis showed that Tmem-specific activation-induced

genes (i.e. Tact-Runx1 genes) are uniquely repressed in Treg. The repression

is likely to be mediated by the interaction between Foxp3 and other

transcription factors that regulate the expression of the Tmem-specific

activation genes (Figure 3C). Interestingly, *Runx1* was associated with these

Tmem-specific genes. In fact, Foxp3 interacts with Runx1 and thereby

represses IL-2 transcription and controls the regulatory function of Treg (38),

and a significant part of the Foxp3-binding to active enhancers occurs through

the Foxp3-Runx1 interaction (9). These suggest that Runx1 may have a

unique role in the differentiation and maintenance of Tmem.

While CTLA-4 is commonly recognised as a Treg marker, it is upregulated in

all activated T cells, thus CTLA-4 is also a marker of activated T cells (41).

CTLA-4 is in fact expressed by only a subset of resting Treg (63), which may

be more activated and proliferating *in vivo* (64). In fact, our study shows that

CTLA-4 is expressed by non-Treg activated T cells including resting Tmem

(Figure 3D) and FOXP3- Tfh-like effector T cells in the tumour

microenvironment (Figure 7E and 8C). These findings support that CTLA-4 is

primarily a marker for general T cell activation, rather than Treg-specific

marker, and that Treg are highly activated T cells with *FOXP3* and *CTLA-4*

expression. Importantly, although both FOXP3+ and FOXP3- cells had the

same relative level of activation (Figure 7D), the absolute number of FOXP3+

cells expressing *CTLA4* was lower than that of Tfh-type cells (Figure 8J),

which suggests that therapeutic anti-CTLA4 antibodies (i.e. Ipilimumab)

primarily target activated Tfh-like effector cells and thereby directly enhance

their activities in tumour microenvironments. Future studies are required to

experimentally investigate the *in vivo* dynamics of CTLA-4 expression in mice

and humans.

In contrast, the expression of *PDCD1* was consistently high in all Tfh-like

cells, while it was sparse among FOXP3+ cells (Figure 8K). The co-

expression of *BCL6* and *IL21* in some of these PD-1+ cells indicates that Tfh

differentiation occurs in the tumour microenvironment, presumably through

the repeated and chronic exposure to quasi-self antigens (i.e. tumour

antigens). Interestingly, the Tfh signature has been identified in type-I

diabetes in both mice and humans (65). Intriguingly, the Tfh-like genes include

cell-cycle related genes (e.g. *CDK6*), immediate early transcription factors

(*NFATC1*, *EGR2/3*), and RNA-processing genes (e.g. *DICER1*). The

significance of these gene modules should be addressed in future studies.

However, the high *PDCD1* expression in Tfh-like cells may make them

vulnerable to the negative immunoregulatory effect of PD-1 in tumour

microenvironments (50). In fact, the most mature *PDCD1*^{high} Tfh-like cells

(cluster VI, Figure 8K) moderately decrease the expression levels of activation

genes (Figure 8E), suggesting that these cells may have started to be

regulated by PD1 ligands. Further experimental investigations are required to

reveal how dynamically PD1 regulates T cells during immune response.

Interestingly, *RUNX1* is completely repressed in the early phase of FOXP3-

pseudotime (Cluster V) but re-expressed in the late phase of FOXP3-

pseudotime the expression of *RUNX1* is significantly elevated (Cluster VI)

(**Figure 8I**). Similar to RUNX1, some cells appear to be expressing *PDCD1* in

the later phase of FOXP3-pseudotime in Treg-lineage cells. The

reappearance of effector phenotype genes *RUNX1* and *PDCD1* in FOXP3^{high}

cells may indicate that these Treg are highly activated effector Treg, which

may be actively participating in neutralizing the activities of effector T cells

and Tfh. Alternatively, these FOXP3^{high} cells may include ambivalent cells with

the characters of both regulatory and effector, or alternatively, be at the point

of conversion to Tmem (**Figure 8K**). In bulk resting Treg, *Pdcd1* was

expressed at low levels in some Treg (Figure 3E) as well as PD-L2-encoding

gene *Pdcd1lg2* (Figure 3D). Future studies are required to reveal the role of

these cells.

SC4A is a useful method to identify distinct clusters of T cells and the

correlated genes to each cluster, and thereby to reveal characteristic cell

groups and their active gene modules, while retaining the single-cell level

variations. We also showed that SC4A and CCA results can be further

analysed by the *pseudotime* approach (CCA-pseudotime). Since SC4A/CCA

provides functional annotations to cell groups and gene clusters, the

understanding of the pseudotime axis is effective, as shown in the current

study. However, given that pseudotime is not a direct measurement of the

time-dependent events, but rather is that of similarities between samples (66),

future studies are required to analyse time-dependent events *in vivo,* ideally

with a new experimental system to directly address the temporal dynamics. In

order to make the current SC4A/CCA approach accessible to experimental

immunologists, we visualised our single cell data findings using a flow-

cytometry style (Figure 9). Although reliable antibodies are currently not

available for those intracellular candidate genes, and the expression of protein

and transcripts may not be synchronized, the present study demonstrated the

power of the SC4A/CCA approach to extract biological meaning from

unannotated single cell RNA-seq data. The current limitation of SC4A is that it

is computationally expensive (i.e. requires several hours for each analysis

using a standard desktop), and the improvement of the computational

algorithm using a low-level language will be beneficial. Importantly, SC4A is

most effective when used together with in-depth knowledge of immunology

and gene regulation, facilitates the interpretation of CCA results and

explanatory variable selection. Thus, it is hoped that these tools will be widely

used by experimental immunologists with a sound understanding of the

biological significance of results, as well as adequate competence in

computational analysis, which will enable to ask questions involving

multidimensional problems such as multiple T cell subsets.

Data and code availability

All R codes are available upon request. Processed data will be provided upon

reasonable requests to the corresponding author.

Acknowledgement

We thank Dr Cristina Leslie for providing us the details of their dataset

GSE83315, Prof Lucy Walker for a useful discussion, and Dr David Bending

for valuable comments on the manuscript. M.O. is a David Phillips Fellow

(BB/J013951/2) from the Biotechnology and Biological Sciences Research

Council (BBSRC), and is also supported by a pump-priming grant from

Cancer Research Centre of Excellence, Imperial College London/the Institute

of Cancer Research. A.B is supported by the Japanese Ministry of Education,

Culture, Sports, Science and Technology (MEXT) PhD scholarship.

References

1. Macian, F., C. Lopez-Rodriguez, and A. Rao. 2001. Partners in transcription: NFAT and AP-1. *Oncogene* 20: 2476-2489.

2. Malek, T. R. 2008. The biology of interleukin-2. *Annu Rev Immunol* 26: 453-479.

3. Setoguchi, R., S. Hori, T. Takahashi, and S. Sakaguchi. 2005. Homeostatic maintenance of natural Foxp3(+) CD25(+) CD4(+) regulatory T cells by interleukin (IL)-2 and induction of autoimmune disease by IL-2 neutralization. *The Journal of experimental medicine* 201: 723-735.

4. Hori, S. 2014. Lineage stability and phenotypic plasticity of Foxp3(+) regulatory T cells. *Immunological reviews* 259: 159-172.

5. Tran, D. Q., H. Ramsey, and E. M. Shevach. 2007. Induction of FOXP3 expression in naive human CD4+FOXP3 T cells by T-cell receptor stimulation is transforming growth factor-beta dependent but does not confer a regulatory phenotype. *Blood* 110: 2983-2990.

6. Chen, W., W. Jin, N. Hardegen, K.-j. Lei, L. Li, N. Marinos, G. McGrady, and S. M. Wahl. 2003. Conversion of Peripheral CD4+CD25− Naive T Cells to CD4+CD25+ Regulatory T Cells by TGF-β Induction of Transcription Factor *Foxp3*. *The Journal of experimental medicine* 198: 1875-1886.

7. Ono, M., and R. J. Tanaka. 2016. Controversies concerning thymus-derived regulatory T cells: fundamental issues and a new perspective. *Immunology and cell biology* 94: 3-10.

8. Thornton, A. M., and E. M. Shevach. 2000. Suppressor Effector Function of CD4+CD25+ Immunoregulatory T Cells Is Antigen Nonspecific. *The Journal of Immunology* 164: 183-190.

9. Samstein, R. M., A. Arvey, S. Z. Josefowicz, X. Peng, A. Reynolds, R. Sandstrom, S. Neph, P. Sabo, J. M. Kim, W. Liao, M. O. Li, C. Leslie, J. A. Stamatoyannopoulos, and A. Y. Rudensky. 2012. Foxp3 exploits a pre-existent enhancer landscape for regulatory T cell lineage specification. *Cell* 151: 153-166.

10. Levine, A. G., A. Arvey, W. Jin, and A. Y. Rudensky. 2014. Continuous requirement for the TCR in regulatory T cell function. *Nature immunology* 15: 1070-1078.

11. Vahl, J. C., C. Drees, K. Heger, S. Heink, Julius C. Fischer, J. Nedjic, N. Ohkura, H. Morikawa, H. Poeck, S. Schallenberg, D. Rieß, Marco Y. Hein, T. Buch, B. Polic, A. Schönle, R. Zeiser, A. Schmitt-Gräff, K.

Kretschmer, L. Klein, T. Korn, S. Sakaguchi, and M. Schmidt-Supprian. 2014. Continuous T Cell Receptor Signals Maintain a Functional Regulatory T Cell Pool. *Immunity* 41: 722-736.

12. Feuerer, M., J. A. Hill, K. Kretschmer, H. von Boehmer, D. Mathis, and C. Benoist. 2010. Genomic definition of multiple ex vivo regulatory T cell subphenotypes. *Proceedings of the National Academy of Sciences of the United States of America* 107: 5919-5924.

13. Koch, M. A., G. Tucker-Heard, N. R. Perdue, J. R. Killebrew, K. B. Urdahl, and D. J. Campbell. 2009. The transcription factor T-bet controls regulatory T cell homeostasis and function during type 1 inflammation. *Nature immunology* 10: 595-602.

14. Zheng, Y., A. Chaudhry, A. Kas, P. deRoos, J. M. Kim, T. T. Chu, L. Corcoran, P. Treuting, U. Klein, and A. Y. Rudensky. 2009. Regulatory T-cell suppressor program co-opts transcription factor IRF4 to control T(H)2 responses. *Nature* 458: 351-356.

15. Linterman, M. A., W. Pierson, S. K. Lee, A. Kallies, S. Kawamoto, T. F. Rayner, M. Srivastava, D. P. Divekar, L. Beaton, J. J. Hogan, S. Fagarasan, A. Liston, K. G. Smith, and C. G. Vinuesa. 2011. Foxp3+ follicular regulatory T cells control the germinal center response. *Nature medicine* 17: 975-982.

16. van der Veeken, J., A. J. Gonzalez, H. Cho, A. Arvey, S. Hemmers, C. S. Leslie, and A. Y. Rudensky. 2016. Memory of Inflammation in Regulatory T Cells. *Cell* 166: 977-990.

17. Wyss, L., B. D. Stadinski, C. G. King, S. Schallenberg, N. I. McCarthy, J. Y. Lee, K. Kretschmer, L. M. Terracciano, G. Anderson, C. D. Surh, E. S. Huseby, and E. Palmer. 2016. Affinity for self antigen selects Treg cells with distinct functional properties. *Nature immunology* 17: 1093-1101.

18. Dias, S., A. D'Amico, E. Cretney, Y. Liao, J. Tellier, C. Bruggeman, F. F. Almeida, J. Leahy, G. T. Belz, G. K. Smyth, W. Shi, and S. L. Nutt. 2017. Effector Regulatory T Cell Differentiation and Immune Homeostasis Depend on the Transcription Factor Myb. *Immunity* 46: 78-91.

19. Cretney, E., A. Xin, W. Shi, M. Minnich, F. Masson, M. Miasari, G. T. Belz, G. K. Smyth, M. Busslinger, S. L. Nutt, and A. Kallies. 2011. The transcription factors Blimp-1 and IRF4 jointly control the differentiation and function of effector regulatory T cells. *Nature immunology* 12: 304-311.

20. Sakaguchi, S., M. Miyara, C. M. Costantino, and D. A. Hafler. 2010. FOXP3+ regulatory T cells in the human immune system. *Nature reviews. Immunology* 10: 490-500.

21. Fujii, H., J. Josse, M. Tanioka, Y. Miyachi, F. Husson, and M. Ono.
2016. Regulatory T Cells in Melanoma Revisited by a Computational
Clustering of FOXP3+ T Cell Subpopulations. *Journal of immunology*.

22. Andersen, P., and B. Smedegaard. 2000. CD4+ T-Cell Subsets That
Mediate Immunological Memory to Mycobacterium tuberculosis
Infection in Mice. *Infection and Immunity* 68: 621-629.

23. Coleman, M. M., C. M. Finlay, B. Moran, J. Keane, P. J. Dunne, and K.
H. G. Mills. 2012. The immunoregulatory role of CD4+FoxP3+CD25−
regulatory T cells in lungs of mice infected with Bordetella pertussis.
FEMS Immunology & Medical Microbiology 64: 413-424.

24. Ono, M., J. Shimizu, Y. Miyachi, and S. Sakaguchi. 2006. Control of
autoimmune myocarditis and multiorgan inflammation by
glucocorticoid-induced TNF receptor family-related protein(high),
Foxp3-expressing CD25+ and CD25- regulatory T cells. *Journal of
immunology* 176: 4748-4756.

25. Linsley, P. S., J. L. Greene, P. Tan, J. Bradshaw, J. A. Ledbetter, C.
Anasetti, and N. K. Damle. 1992. Coexpression and functional
cooperation of CTLA-4 and CD28 on activated T lymphocytes. *The
Journal of experimental medicine* 176: 1595-1604.

26. Attridge, K., and L. S. Walker. 2014. Homeostasis and function of
regulatory T cells (Tregs) in vivo: lessons from TCR-transgenic Tregs.
Immunological reviews 259: 23-39.

27. Bunce, C., and E. B. Bell. 1997. CD45RC Isoforms Define Two Types
of CD4 Memory T Cells, One of which Depends on Persisting Antigen.
The Journal of experimental medicine 185: 767-776.

28. Min, B., R. McHugh, G. D. Sempowski, C. Mackall, G. Foucras, and W.
E. Paul. 2003. Neonates Support Lymphopenia-Induced Proliferation.
Immunity 18: 131-140.

29. Curotto de Lafaille, M. A., A. C. Lino, N. Kutchukhidze, and J. J.
Lafaille. 2004. CD25- T cells generate CD25+Foxp3+ regulatory T cells
by peripheral expansion. *Journal of immunology* 173: 7259-7268.

30. Miyao, T., S. Floess, R. Setoguchi, H. Luche, H. J. Fehling, H.
Waldmann, J. Huehn, and S. Hori. 2012. Plasticity of Foxp3(+) T cells
reflects promiscuous Foxp3 expression in conventional T cells but not
reprogramming of regulatory T cells. *Immunity* 36: 262-275.

31. Ono, M., R. J. Tanaka, M. Kano, and T. Sugiman. 2013. Visualising the
Cross-Level Relationships between Pathological and Physiological
Processes and Gene Expression: Analyses of Haematological
Diseases. *PLoS One* 8: e53544.

32. Greenacre, M. 2008. *Correspondence Analysis in Practice.* Chapman
& Hall/CRC,, London.

33. Tzeng, J., H. H.-S. Lu, and W.-H. Li. 2008. Multidimensional scaling for large genomic data sets. *BMC Bioinformatics* 9: 179.

34. Ono, M., R. J. Tanaka, and M. Kano. 2014. Visualisation of the T cell differentiation programme by Canonical Correspondence Analysis of transcriptomes. *BMC genomics* 15: 1028.

35. Tirosh, I., B. Izar, S. M. Prakadan, M. H. Wadsworth, 2nd, D. Treacy, J. J. Trombetta, A. Rotem, C. Rodman, C. Lian, G. Murphy, M. Fallahi-Sichani, K. Dutton-Regester, J. R. Lin, O. Cohen, P. Shah, D. Lu, A. S. Genshaft, T. K. Hughes, C. G. Ziegler, S. W. Kazer, A. Gaillard, K. E. Kolb, A. C. Villani, C. M. Johannessen, A. Y. Andreev, E. M. Van Allen, M. Bertagnolli, P. K. Sorger, R. J. Sullivan, K. T. Flaherty, D. T. Frederick, J. Jane-Valbuena, C. H. Yoon, O. Rozenblatt-Rosen, A. K. Shalek, A. Regev, and L. A. Garraway. 2016. Dissecting the multicellular ecosystem of metastatic melanoma by single-cell RNA-seq. *Science* 352: 189-196.

36. Wakamatsu, E., D. Mathis, and C. Benoist. 2013. Convergent and divergent effects of costimulatory molecules in conventional and regulatory CD4+ T cells. *Proceedings of the National Academy of Sciences of the United States of America* 110: 1023-1028.

37. Painter, M. W., S. Davis, R. R. Hardy, D. Mathis, C. Benoist, and C. Immunological Genome Project. 2011. Transcriptomes of the B and T lineages compared by multiplatform microarray profiling. *Journal of immunology* 186: 3047-3057.

38. Ono, M., H. Yaguchi, N. Ohkura, I. Kitabayashi, Y. Nagamura, T. Nomura, Y. Miyachi, T. Tsukada, and S. Sakaguchi. 2007. Foxp3 controls regulatory T-cell function by interacting with AML1/Runx1. *Nature* 446: 685-689.

39. Chen, W., and R. Maitra. 2017. EMCluster: EM Algorithm for Model-Based Clustering of Finite Mixture Gaussian Distribution. R Package.

40. Street, K., D. Risso, R. B. Fletcher, D. Das, J. Ngai, N. Yosef, E. Purdom, and S. Dudoit. 2017. Slingshot: Cell lineage and pseudotime inference for single-cell transcriptomics. *bioRxiv*.

41. Walker, L. S., and D. M. Sansom. 2015. Confusing signals: recent progress in CTLA-4 biology. *Trends Immunol* 36: 63-70.

42. Szabo, S. J., B. M. Sullivan, C. Stemmann, A. R. Satoskar, B. P. Sleckman, and L. H. Glimcher. 2002. Distinct Effects of T-bet in T_H1 Lineage Commitment and IFN-γ Production in CD4 and CD8 T Cells. *Science* 295: 338-342.

43. Dang, E. V., J. Barbi, H. Y. Yang, D. Jinasena, H. Yu, Y. Zheng, Z. Bordman, J. Fu, Y. Kim, H. R. Yen, W. Luo, K. Zeller, L. Shimoda, S. L. Topalian, G. L. Semenza, C. V. Dang, D. M. Pardoll, and F. Pan. 2011.

Control of T(H)17/T(reg) balance by hypoxia-inducible factor 1. *Cell* 146: 772-784.

44. Ho, I. C., M. R. Hodge, J. W. Rooney, and L. H. Glimcher. 1996. The proto-oncogene c-maf is responsible for tissue-specific expression of interleukin-4. *Cell* 85: 973-983.

45. Vinuesa, C. G., M. A. Linterman, C. C. Goodnow, and K. L. Randall. 2010. T cells and follicular dendritic cells in germinal center B-cell formation and selection. *Immunological reviews* 237: 72-89.

46. Sakaguchi, S., T. Yamaguchi, T. Nomura, and M. Ono. 2008. Regulatory T cells and immune tolerance. *Cell* 133: 775-787.

47. Seddon, B., and R. Zamoyska. 2002. TCR and IL-7 Receptor Signals Can Operate Independently or Synergize to Promote Lymphopenia-Induced Expansion of Naive T Cells. *The Journal of Immunology* 169: 3752-3759.

48. Roychoudhuri, R., K. Hirahara, K. Mousavi, D. Clever, C. A. Klebanoff, M. Bonelli, G. Sciume, H. Zare, G. Vahedi, B. Dema, Z. Yu, H. Liu, H. Takahashi, M. Rao, P. Muranski, J. G. Crompton, G. Punkosdy, D. Bedognetti, E. Wang, V. Hoffmann, J. Rivera, F. M. Marincola, A. Nakamura, V. Sartorelli, Y. Kanno, L. Gattinoni, A. Muto, K. Igarashi, J. J. O'Shea, and N. P. Restifo. 2013. BACH2 represses effector programs to stabilize T(reg)-mediated immune homeostasis. *Nature* 498: 506-510.

49. Lonnberg, T., V. Svensson, K. R. James, D. Fernandez-Ruiz, I. Sebina, R. Montandon, M. S. Soon, L. G. Fogg, A. S. Nair, U. Liligeto, M. J. Stubbington, L. H. Ly, F. O. Bagger, M. Zwiessele, N. D. Lawrence, F. Souza-Fonseca-Guimaraes, P. T. Bunn, C. R. Engwerda, W. R. Heath, O. Billker, O. Stegle, A. Haque, and S. A. Teichmann. 2017. Single-cell RNA-seq and computational analysis using temporal mixture modelling resolves Th1/Tfh fate bifurcation in malaria. *Sci Immunol* 2.

50. Okazaki, T., S. Chikuma, Y. Iwai, S. Fagarasan, and T. Honjo. 2013. A rheostat for immune responses: the unique properties of PD-1 and their advantages for clinical application. *Nature immunology* 14: 1212-1218.

51. Wong, W. F., M. Kurokawa, M. Satake, and K. Kohu. 2011. Down-regulation of Runx1 expression by TCR signal involves an autoregulatory mechanism and contributes to IL-2 production. *J Biol Chem* 286: 11110-11118.

52. Buchbinder, E. I., and A. Desai. 2016. CTLA-4 and PD-1 Pathways: Similarities, Differences, and Implications of Their Inhibition. *Am J Clin Oncol* 39: 98-106.

53. McPherson, R. C., J. E. Konkel, C. T. Prendergast, J. P. Thomson, R. Ottaviano, M. D. Leech, O. Kay, S. E. Zandee, C. H. Sweenie, D. C.

Wraith, R. R. Meehan, A. J. Drake, and S. M. Anderton. 2014. Epigenetic modification of the PD-1 (Pdcd1) promoter in effector CD4(+) T cells tolerized by peptide immunotherapy. *Elife* 3.

54. Morgan, D. A., F. W. Ruscetti, and R. Gallo. 1976. Selective in vitro growth of T lymphocytes from normal human bone marrows. *Science* 193: 1007-1008.

55. Rosenblum, M. D., S. S. Way, and A. K. Abbas. 2016. Regulatory T cell memory. *Nature reviews. Immunology* 16: 90-101.

56. Miyara, M., Y. Yoshioka, A. Kitoh, T. Shima, K. Wing, A. Niwa, C. Parizot, C. Taflin, T. Heike, D. Valeyre, A. Mathian, T. Nakahata, T. Yamaguchi, T. Nomura, M. Ono, Z. Amoura, G. Gorochov, and S. Sakaguchi. 2009. Functional delineation and differentiation dynamics of human CD4+ T cells expressing the FoxP3 transcription factor. *Immunity* 30: 899-911.

57. Weissler, K. A., and A. J. Caton. 2014. The role of T-cell receptor recognition of peptide:MHC complexes in the formation and activity of Foxp3(+) regulatory T cells. *Immunological reviews* 259: 11-22.

58. Miragaia, R. J., T. Gomes, A. Chomka, L. Jardine, A. Riedel, A. N. Hegazy, I. Lindeman, G. Emerton, T. Krausgruber, J. Shields, M. Haniffa, F. Powrie, and S. A. Teichmann. 2017. Single cell transcriptomics of regulatory T cells reveals trajectories of tissue adaptation. *bioRxiv*.

59. Tai, X., B. Erman, A. Alag, J. Mu, M. Kimura, G. Katz, T. Guinter, T. McCaughtry, R. Etzensperger, L. Feigenbaum, D. S. Singer, and A. Singer. 2013. Foxp3 transcription factor is proapoptotic and lethal to developing regulatory T cells unless counterbalanced by cytokine survival signals. *Immunity* 38: 1116-1128.

60. Sahni, H., S. Ross, A. Barbarulo, A. Solanki, C. I. Lau, A. Furmanski, J. I. Saldana, M. Ono, M. Hubank, M. Barenco, and T. Crompton. 2015. A genome wide transcriptional model of the complex response to pre-TCR signalling during thymocyte differentiation. *Oncotarget* 6: 28646-28660.

61. Hayatsu, N., T. Miyao, M. Tachibana, R. Murakami, A. Kimura, T. Kato, E. Kawakami, T. A. Endo, R. Setoguchi, H. Watarai, T. Nishikawa, T. Yasuda, H. Yoshida, and S. Hori. 2017. Analyses of a Mutant Foxp3 Allele Reveal BATF as a Critical Transcription Factor in the Differentiation and Accumulation of Tissue Regulatory T Cells. *Immunity* 47: 268-283 e269.

62. Kalekar, L. A., S. E. Schmiel, S. L. Nandiwada, W. Y. Lam, L. O. Barsness, N. Zhang, G. L. Stritesky, D. Malhotra, K. E. Pauken, J. L. Linehan, M. G. O'Sullivan, B. T. Fife, K. A. Hogquist, M. K. Jenkins, and D. L. Mueller. 2016. CD4(+) T cell anergy prevents autoimmunity

and generates regulatory T cell precursors. *Nature immunology* 17: 304-314.

63. Takahashi, T., T. Tagami, S. Yamazaki, T. Uede, J. Shimizu, N. Sakaguchi, T. W. Mak, and S. Sakaguchi. 2000. Immunologic self-tolerance maintained by CD25(+)CD4(+) regulatory T cells constitutively expressing cytotoxic T lymphocyte-associated antigen 4. *The Journal of experimental medicine* 192: 303-310.

64. Tang, A. L., J. R. Teijaro, M. N. Njau, S. S. Chandran, A. Azimzadeh, S. G. Nadler, D. M. Rothstein, and D. L. Farber. 2008. CTLA4 expression is an indicator and regulator of steady-state CD4+ FoxP3+ T cell homeostasis. *Journal of immunology* 181: 1806-1813.

65. Kenefeck, R., C. J. Wang, T. Kapadi, L. Wardzinski, K. Attridge, L. E. Clough, F. Heuts, A. Kogimtzis, S. Patel, M. Rosenthal, M. Ono, D. M. Sansom, P. Narendran, and L. S. Walker. 2015. Follicular helper T cell signature in type 1 diabetes. *The Journal of clinical investigation* 125: 292-303.

66. Haghverdi, L., M. Buttner, F. A. Wolf, F. Buettner, and F. J. Theis. 2016. Diffusion pseudotime robustly reconstructs lineage branching. *Nat Methods* 13: 845-848.

Table 1. **Datasets used in this study**

Accession number	Short description	Reference	Description of animal models	Timing of cell harvest	Cell purification strategy and sorting markers	Tissue origin	Link to Figures	Link to materials and methods
GSE15907	T cell subsets	Immunological Genome Project; Painter *et al.*, 2011	Primary cells from multiple immune lineages are isolated *ex-vivo*, primarily from young adult B6 male mice (WT, Foxp3GFP or BDC Tg mice), and double-sorted to >99% purity.	6 weeks	Flow cytometric sorting; Treg (spleen): Foxp3GFP+ CD25+ CD4+ Tmem (subcutaneous (sc)LN, spleen): TCRβ+CD44^{high} CD122^{lo} CD25- CD4+ CD44^{hi}CD62L^{lo} Tmem (scLN, spleen): Foxp3GFP-TCRβ+ CD44^{hi} CD62L^{lo} CD4+ Naïve CD4 (scLN): CD25- 62L^{hi} 44lo CD4+ Naïve CD4 (mesenteric (m) LN): CD25- CD62L^{hi} CD44^{lo} CD4+ Naïve CD4 (Peyer's patches): TCRβ+ CD44^{lo} CD62L^{hi} CD4+ Naïve CD4 (spleen): CD25- CD62L^{hi} CD44^{lo} CD4+ Foxp3- Tnaive (spleen): Foxp3GFP- CD44^{lo} CD4+ Non-dLN, BDC (scLN): BDC+ CD4+ dLN BDC (pancreatic LN): CD4+ BDC+ Tissue-Teff, BDC (pancreas): BDC+ CD4+	Spleen, subcutaneous LNs, mesenteric LN, Peyer's patches, pancreatic LN, pancreas	Figure 1 Figure 2 Figure 3	NCBI GEO

					*Exclusion markers include PI, CD8, CD11b, CD11c, CD19, CD49b, Gr-1, Ter119			
GSE83315	aTreg data	Van der Veeken *et al.*, 2016	Mixed bone marrow chimeras were generated with 90% *Foxp3GFP-DTR* /10% *Foxp3-GFP-CRE-ERT2 Rosa26YFP* bone marrow. Diphtheria toxin (DT) was administered at day 0 to these chimeric mice in order to deplete *Foxp3GFP-DTR* Treg cells and induce expansion/ activation of effector CD4+ T cells and Treg, thereby inducing inflammation. Subsequently, tamoxifen was administered at days 3 and 4 to irreversibly label *Foxp3*-expressing *Foxp3-GFP-CRE-ERT2 Rosa26YFP* T cells with YFP. Resting Treg (rTr), activated Treg (aTr), and 'memory' Treg (mTr) were isolated at day 0, 11, and 60, respectively, based on the dynamics of inflammation (CD4+ T cell number is normalised by day 60).	Day 0 (resting Treg), day 11 (activated Treg), day 60 (memory Treg) after DT treatment	Flow cytometric sorting; rTr: CD4, Foxp3-GFP aTr and mTr: CD4, Foxp3-GFP, YFP	Spleen and peripheral LN	Figure 5C and 5D	PMC NCBI GEO

GSE61077	TCR KO data	Levine *et al.*, 2014	8-10 week mice from *Trac*^{flox/WT} x *Foxp3*^{ERT2-Cre} (tamoxifen-inducible deletion of TCRα in Treg. Tamoxifen was administered on days 0, 1 and 3.	Day 14 after the first tamoxifen administration	Flow cytometric sorting; TCRβ+ CD4+ Foxp3+ CD44high/low CD62Llow/high	LN			NCBI GEO
GSE42276	T cell activation	Wakamatsu *et al.*, 2013	Conventional CD4+ T cells from C57BL/6J male mice were stimulated by anti-CD3 and anti-CD28 for 20 h and 48 h and data were pooled. 0h unstimulated samples were used as control.	8 weeks	Flow cytometric sorting; DAPI−CD45R−CD8a−CD11b/c−CD4+ GFP+	Spleen, LN	Figure 1 Figure 2 Figure 3 Figure 5		NCBI GEO
GSE6939	RV-transduced T cells	Ono *et al.*, 2007	Cells: T cells from LN and spleen of 8 week-old BALB/c mice and purified into CD4+ naive T cells (GITR^{low}CD25$^-$CD4$^+$), which were subsequently activated by anti-CD3 and antigen-presenting cells (mitomycin-treated Thy1(-) splenocytes) in the presence of IL-2. On the following day, T cells were retrovirally gene transduced with Runx1 (AML1), wild type Foxp3, and empty vector as control.	60 hours after transfection	Flow cytometric sorting; CD4, GFP Exclusion marker: PI	Spleen, LN	Figure 2 Figure 3		NCBI GEO

GSE72056	Single cell analysis of tumour-infiltrating T cells	Tirosh *et al.*, 2016	Single cell RNA-seq analysis of human melanoma tumour samples. Freshly resected samples were disaggregated to generate single cell suspensions of mixed cells of unknown identities. Individual viable immune (CD45+) and nonimmune (CD45−) cells (including malignant and stromal cells) were recovered from the single cell suspension by flow cytometry. Single cells were profiled by single-cell RNA-seq.	Single cells were obtained within 45 min of tumour resection	Flow cytometric sorting; CD45	Human melanoma tissues	Figure 7 Figure 8 Figure 9	NCBI GEO
GSE15390	Human activated T cells and Treg	Beyer *et al.*, 2011	Resting T cells (CD25-CD4+ T cells; GSM386262, GSM386264, and GSM386266) were obtained from whole blood of healthy human donors. Activated T cells (GSM777695) were prepared by stimulating CD25-CD4+ T cells for 24 hours with CD3 and IL-2.	Freshly sorted from buffy coat; or cultured for 24 hours	Magnetic and flow cytometric sorting; CD4, CD25, CD127	Human PBMC	Figure 7	PMC

 Figure Legends

Figure 1. Identification of the activation signature in Treg and Tmem by

CCA of T cell populations

The microarray dataset of peripheral CD4+ T cells, including naïve, effector

and memory phenotype from various sites (GSE15907), was analysed using

the T cell activation variable, which was obtained by the microarray dataset of

conventional activated CD4+ T cells (GSE42276). (**A**) Schematic

representation of CCA for the cross-level analysis of T cell populations (cells),

immunological processes, and genes. (**B**) CCA was applied to the T cell

population data using an explanatory variable for T cell activation, which was

obtained as fold change between activated and resting conventional CD4+ T

cells. The CCA solution is thus one-dimensional, and is used as "T cell

activation score" (see Methods).

Figure 2. Identification of the Foxp3-independent activation signature in

Treg by CCA of T cell populations

The microarray dataset of peripheral CD4+ T cells (GSE15907) was analysed

using the T cell activation variable and the variables for retroviral *Foxp3*

transduction and *Runx1* transduction as explanatory variables. (**A**) The CCA

solution was visualised by a biplot where CD4+ T cell samples are shown by

closed circles (see legend) and the explanatory variables are shown by blue

arrows. Percentage indicates that of the variance accounted for by the inertia

of the axes (i.e. the amount of information (eigenvalue) retained in each axis).

(**B**) Gene biplot of the 2D CCA solution in (C) showing the relationships

between genes (grey circles) and the explanatory variables (blue arrows).

Selected key genes are annotated.

Figure 3. Differential regulations of transcriptional modules for

activation in Treg and Tmem by Foxp3 and Runx1

(**A**) Definition of Tact-Foxp3 genes and Tact-Runx1 genes. In the gene plot of

the CCA solution in Figure 2B, Axis 1-low genes (25 percentile low) were

designated as *activation genes*, which were further classified into Tact-Foxp3

genes and Tact-Runx1 genes by Axis 2, which have high correlations to Treg

and Tmem samples, respectively, in the CCA cell space (Figure 2A). (**B**)

Heatmap analysis of all the Tact-Foxp3 genes. (**C**) Heatmap analysis of all the

Tact-Runx1 genes. (**D**) Heatmap analysis of selected Tact-Foxp3 genes. (**E**)

Heatmap analysis of selected Tact-Runx1 genes.

Figure 4. A model for the differential regulation of activation genes in

Treg and Tmem

The proposed differential regulations of TCR signal downstream genes in

Treg and Tmem. Since both naturally-arising Treg and Tmem are self-reactive

T cells, they may frequently receive tonic TCR signals by recognising their

cognate antigens in the periphery. This results in the full activation of both the

Tact-Foxp3 and Tact-Runx1 gene modules in Tmem. However, in Treg,

Foxp3 represses Tact-Runx1 genes and sustains the expression of Tact-

Foxp3 genes, producing the characteristic Treg transcriptome.

Figure 5. The activation signature of Treg is dependent on TCR

signalling

(**A**) The experimental design for the TCR dataset. CD44hi activated Treg and

CD44lo naïve-like Treg were obtained from TCRa KO or WT mice and

analysed by transcriptome analysis. (**B**) CCA was applied to the transcriptome

data of CD44^{lo}CD62^{hi} naïve-like and CD44^{hi}CD62^{lo} activated Treg cell

populations from inducible *TCRa* KO or WT (from the *TCR KO* data,

GSE61077), using the T cell activation variable as the explanatory variable.

This produces a 1D CCA solution, and the sample score was plotted

(representing "T cell activation score"). (**C**) The experimental design for the

activated Treg dataset. Bone marrow (BM) cells were obtained from

Foxp3^{GFPCreERT2}:Rosa26YFP mice (YFP mice), and transferred into *Foxp3^{GFP}*

^{DTR} mice (Foxp3-DTR mice), in order to make BM chimera, in which ~10% of

Treg expressed DTR. Subsequently, DT was administered to these BM

chimera, which depleted Foxp3-DTR cells but not donor cells. This treatment

induced a transient activation of T cells and inflammation *in vivo*. Activated

Treg (aTreg) were obtained from these mice with inflammation, while resting

Treg (rTreg) were from control mice, and memory Treg (mTreg) were from the

mice after the resolution of inflammation. (**D**) 1D CCA sample score plot of

transcriptomic data of resting Treg (rTreg), *in vivo* activated Treg (aTreg) and

memory Treg (mTreg) from the *aTreg* data (GSE83315), with T cell activation

signature as explanatory variable.

Figure 6. The comparative analysis of Tmem-specific and Treg-Tmem shared activation genes and TCR-dependent and activated Treg-specific genes

Venn diagram analysis was used to obtain intersects of *TCR-dependent genes* (DEG between TCRa KO and WT Treg), *aTreg-specific genes* (DEG between aTreg and rTreg), and Tact-Foxp3 and Tact-Runx1 genes (see Figure 3). (**A**) Pie chart showing the number of genes in the intersects between *aTreg-specific genes*, *TCR-dependent genes,* and Tact-Runx1 genes. (**B**) Pie chart showing the numbers of *aTreg-specific genes*, *TCR-dependent genes,* and Tact-Foxp3 genes.

(**C**) Pathway analysis of Tact-Foxp3 genes, Tact-Runx1 genes and *aTreg-specific genes* showing enriched pathways in these gene lists.

Figure 7. Single cell CCA of melanoma-infiltrating T cells determines the activation status of individual T cells and identifies a putative Tfh-like process

(**A**) Schematic representation of CCA of CD4+ T cell single cell transcriptomes analysed by two explanatory variables: activated naïve T cells (Tact) and resting naïve T cells (Trest). (**B**) CCA biplot showing the relationship between Treg and non-Treg T cells (sample scores) and the explanatory variables (Tact and Trest). Axis 1 represents the difference between Tact and Trest, and thus, Activated T cells and Resting T cells were defined by the CCA Axis 1 score, and these cells were further classified into Treg and non-Treg by their *FOXP3* expression (see legend). Percentage

indicates that of the variance (inertia) accounted for by the axis. (**C**) Violin plot

showing the CCA activation scores (Axis 1 score $\times$ -1) of FOXP3- and

FOXP3+ cell groups. Asterisk indicates statistical significance by Mann-

Whitney test (**D**) Violin plot showing the CCA activation scores of Activated

(Act.) and Resting (Rest.) FOXP3- and FOXP3+ cell groups. Asterisks

indicate the values of post-hoc Dunn's test following a Kruskal Wallis test. ***

$p < 0.005$. (**E**) Gene biplot of the CCA solution in (B) showing the relationships

between genes (grey circles) and the Tact and Trest explanatory variables

(blue arrows). Genes are shown by grey circles, and well-known genes that

are key for T cell activation processes are annotated.

12
Figure 8. SC4A identifies the bifurcation point of activated T cells that

leads to Tfh-like and Treg differentiation in tumour-infiltrating T cells

SC4A was applied to the single cell data of tumour-infiltrating T cells, and 4

genes (CTLA-4, CCR7, FOXP3, and PDCD1) were chosen as explanatory

variables to represent the T cell activation, resting, FOXP3-driven process,

and Tfh-like process. (**A**) The design of analysis. The single cell data from the

melanoma samples were analysed by SC4A to identify the most effective

combinations of explanatory variables for dispersing the 4 presumptive T cell

populations identified in Figure 7. These genes were used as explanatory

variables to analyse the rest of the single cell data as main dataset. Thus, the

single-cell level dynamics of T cell differentiation and activation are modelled

by the key biological processes that are represented by the T cell populations

and explanatory variables. (**B**) Single cell sample space of the final SC4A

output showing correlations between single cell samples and the explanatory

variables **(C)** Gene space of the final SC4A output showing correlations

between genes and the explanatory variables. The genes that showed high

correlations to the PDCD1, CTLA4, and FOXP3 variables were identified as

Tfh-like genes, Activation genes, and FOXP3-driven genes, respectively. **(D)**

The identification of two differentiation processes as lineages and a

bifurcation point. The cells in the sample space of the SC4A output (B) were

classified into 6 clusters by an unsupervised clustering algorithm. These

clusters were further analysed for pseudotime inference. (**E-G**) The average

gene expression was plotted against each pseudotime (upper: FOXP3-

pseudotime; lower: Tfh-pseudotime). The bifurcation point (Cluster II) is

emphasised by broken lines. The numbers in circle indicate the cluster

number. Gene expression was standardised, and the sum of the standardised

expression was obtained for (**E**) Activation genes, (**F**) FOXP3-driven genes,

and (**G**) Tfh-like genes (see C). **(H-M)** The expression of key genes was

plotted against each pseudotime.

Figure 9. Identification of the conserved genes for the differential

regulation of Tfh-like and Treg differentiation in activated T cells

(**A**) The identified lineage curves and the bifurcation point in the tumour-

infiltrating T cells. The number in circle indicates the cluster number in Figure

8D. (**B-D**) The expression of selected feature genes was plotted against each

pseudotime. Genes are from the intersect of (**B**) Activation genes (Figure 8C)

and Tact-Foxp3 genes (Figure 3), (**C**) FOXP3-driven genes (Figure 8C) and

Tact-Foxp3 genes, and (**D**) Tfh-like genes (Figure 8C) and Tact-Runx1 genes

(Figure 3). **(E)** The expression of selected genes in the tumour-infiltrating T

cells was shown by a 2-dimensional plot in a flow cytometric style. Data from

Treg-lineage cells (Cluster V and VI, upper panels) and Tfh-like lineage cells

(Cluster III and IV, lower panels). The gene in x-axis (NFAT5) is from the

activation gene group (B), while y-axis shows genes from either the FOXP3-

Treg group (C) or the Tfh-like/Tmem group (D). Thresholds and quadrant

gates were determined in an empirical manner using density plot.

Figure 1

A

Gene-oriented CCA of T cell populations and immunological processes

Two transcriptome data

T cell populations

Explanatory variable(s)

Main data: T cell populations

Intersect

Explanatory variable: Prototype T cell data as immunological process

CCA solution = 3 levels:

Immunological process

Gene

Cell

→ Visualise relationships between genes, cell populations, and immunological processes

B

Main data: Peripheral CD4+ T cell populations

Intersect

Explanatory variable: T cell activation

Solution: CCA 1D space

Axis 1 (T cell activation)

Population

Foxp3- Tnaive, Foxp3+ Treg, Tnaive, Tmem, non-dLN, dLN, tissue-Teff

BDC Tg

Figure 2

A

B

Figure 3

Figure 4

Figure 5

A

B

Main data:
TCRa KO vs WT
Naive-like Treg
vs activated Treg

Intersect

Explanatory variable:
T cell activation

Solution: CCA 1D space

C

D

Main data:
rTreg
aTreg
mTreg

Intersect

Explanatory variable:
T cell activation

Solution: CCA 1D space

Figure 6

A

Tact-Runx1 genes (Tmem-specific)

B

Tact-Foxp3 genes (Treg-Tmem shared)

C

Figure 7

Figure 8

Figure 9