Evolution of the mutational process under relaxed selection in *Caenorhabditis elegans*

Ayush Shekhar Saxena[1], Matthew P. Salomon[1,2], Chikako Matsuba[1,2], Shu-Dan Yeh[1,3] and

Charles F. Baer[1,4]

1 – Department of Biology, University of Florida, Gainesville, FL, USA

2 – Department of Molecular Oncology, John Wayne Cancer Institute, Santa Monica, CA, USA

3 - Department of Life Sciences, National Central University, Taiwan

4 – University of Florida Genetics Institute

Correspondence to:

Ayush Shekhar Saxena

s.ayush@ufl.edu

or

Charles F. Baer

cbaer@ufl.edu

Abstract

The mutational process varies at many levels, from within genomes to among taxa. Many mechanisms have been linked to variation in mutation, but understanding of the evolution of the mutational process is rudimentary. Physiological condition is often implicated as a source of variation in microbial mutation rate and may contribute to mutation rate variation in multicellular organisms.

Deleterious mutations are a ubiquitous source of variation in condition. We test the hypothesis that the mutational process depends on the underlying mutation load in two groups of *Caenorhabditis elegans* mutation accumulation (MA) lines that differ in their starting mutation loads. "First-Order MA" (O1MA) lines maintained under minimal selection for ~250 generations were divided into high-fitness and low-fitness groups and sets of "second-order MA" (O2MA) lines derived from each O1MA line were maintained for ~150 additional generations. Genomes of 48 O2MA lines and their progenitors were sequenced. There is significant variation among O2MA lines in base-substitution rate (μ_{bs}), but no effect of initial fitness, whereas the indel rate is greater in high-fitness O2MA lines. Overall, μ_{bs} is positively correlated with recombination and proximity to short tandem repeats and negatively correlated with 10 bp and 1 Kb GC content. However, probability of mutation is well-predicted by the three-nucleotide motif. ~90% of the variance in standing nucleotide variation is explained by mutability. Total mutation rate increased in the O2MA lines, as predicted by the "drift barrier" model of mutation rate evolution. These data, combined with experimental estimates of fitness, suggest that epistasis is synergistic.

Introduction

The evolution of the mutation rate is of longstanding interest to evolutionary theorists (Fisher 1930; Sturtevant 1937; Lynch et al. 2016), and there is abundant empirical evidence that the overall rate, molecular spectrum, and phenotypic consequences of mutation - collectively, the mutational process - vary at many biological levels, from within an individual genome to among species and higher taxa (Drake et al. 1998; Conrad et al. 2011; Schrider et al. 2013; Long et al. 2016; Ness et al. 2016; Carlson et al. 2018). The mechanistic, environmental, ecological, and evolutionary factors that potentially influence variation in the mutational process are legion.

Microbiologists have appreciated for many decades that physiological stress is often associated with increased mutation rate (e.g., see Figure 6 of Ogur et al. 1960). Recently, Agrawal and his colleagues have undertaken a systematic investigation into the effects of physiological condition (~ "stress") on the mutational process in *Drosophila melanogaster*, motivated by theoretical findings that if the mutation rate is condition-dependent, the accumulation of deleterious mutations can have interesting and sometimes counterintuitive feedback effects on population mean fitness (Agrawal 2002; Shaw and Baer 2011). They manipulated physiological condition both exogenously, by manipulating food quality (Agrawal and Wang 2008) and endogenously, by allowing mutations to accumulate under relaxed selection on genomes that were initially identical except for the presence or absence of one or two mutations of large deleterious effect (Sharp and Agrawal 2012). Poorly-fed females transmitted ~30% more lethal mutations than did well-fed females (Agrawal and Wang 2008). Similarly, mutation accumulation (MA) lines beginning with a large genetic load declined in fitness more rapidly than lines with wild-type genomes, which is most simply explained by the low-fitness lines having a greater mutation rate than the high-fitness lines (Sharp and Agrawal 2012). Whole-genome sequencing revealed that the faster decline in fitness of the loaded MA lines can be attributed to an elevated rate of small deletions, plausibly as a result of different mechanisms of double-strand break repair (Sharp and Agrawal 2016). In an analogous study, Ávila et al. (2006) constructed a set of MA lines of *D. melanogaster* derived from a single MA line that had itself accumulated mutations under relaxed selection for 265 generations, a protocol that we call "second-order MA" (O2MA). The per-generation decline in fitness was greater in the second-order MA lines than in the ancestral ("first-order MA", O1MA) lines. That result is consistent with an increased mutation rate in the O2MA lines relative to the O1MA lines, but it is also consistent with mutational effects being greater in the second-order MA lines (i.e., synergistic epistasis; Dickinson 2008).

We report here the results of a second-order MA experiment in the nematode *C.*

elegans, specifically designed to assess the relationship between the initial genomic load of

spontaneous deleterious mutations and the subsequent effects on the mutational process

(Figure 1). Initially, a set of 100 MA lines derived from a single, highly inbred individual of the

N2 strain was allowed to accumulate mutations for approximately 250 generations under

minimal selection. From a subset of 67 O1MA lines assayed for absolute fitness (defined as

lifetime reproductive success under non-competitive conditions), we chose five lines with

consistently high fitness and five lines with consistently low fitness, from which we established

ten independent sets of 48 O2MA lines, referred to as "O2MA families", which were allowed to

accumulate mutations for an additional ~150 generations (Matsuba et al. 2012). Upon

completion of the second-order MA phase, five replicate O2MA lines from each of the ten O2MA

families were sequenced at ~25X average coverage, along with nine of the ten O1MA

progenitors and an additional 23 O1MA lines not included in the O2MA experiment.

The resulting data allow us to address several fundamental questions about the

evolution of the mutational process. First, does initial fitness affect the mutational process, and

if so, how? Second, how fast does genetic variation in the mutational process accumulate due

to the effects of new spontaneous mutations? Third, is the mutation rate itself subject to a

mutational bias, as predicted by the "drift barrier" hypothesis of mutation rate evolution? Fourth,

how does the mutational process depend on underlying features of the genome, e.g., local

sequence context, recombination rate or base composition? Fifth, to what extent does the local

mutational milieu predict standing nucleotide sequence variation?

Finally, in combination with experimental estimates of fitness, we can assess how the

average mutational effect depends on the underlying genetic load. If mutational effects are

greater in the low-fitness O2MA lines than in the high-fitness lines, it implies that deleterious

mutations interact synergistically; the converse implies diminishing-returns epistasis.

Results

Pooled over 48 (out of 50) O2MA lines and nine of the ten O2MA progenitors (three lines

yielded poor sequencing coverage), using our default pipeline we identified 1828 base

substitutions (1481 O2MA, 347 O1MA), 361 small deletions (293 O2 MA, 68 O1MA) and 236

small insertions (196 O2MA, 40 O1MA). We sequenced the genomes of an additional 23 O1MA

lines not included in the O2MA experiment, from which we identified 884 base substitutions, 142

deletions, and 93 insertions. Mutation rates for the different groups are summarized in Table 1,

and given for individual lines in

Figure 1. Schematic depiction of the second-order MA (O2MA) experiment. After ~250

generations of MA (unfilled worms), five low-fitness O1MA lines (light blue worms) and five high-

fitness O1MA lines (dark blue worms) were chosen as progenitors ("O2MA progenitor", circled

worm) for sets of 48 O2MA lines (orange worms).   Each set of 48 O2MA lines derived from an

O2MA progenitor is an "O2MA family". After ~150 additional generations (G400), five O2MA

lines were randomly chosen from each O2MA family for sequencing ("ACGT"). At the same

time, the O2MA progenitor of each O2MA family was sequenced, along with 23 randomly

chosen O1MA lines that were not progenitors of O2MA families (gray worms).

Table 1. Average mutation rates (SEM).

	O2MA progenitor (n=9)	O1MA other (n=23)	O1MA all (n=32)	O2MA, High (n=24)	O2MA, Low (n=24)	O2MA, all (n=48)
μ_{bs} ($\times10^9$)	2.26 (0.12)	2.35 (0.11)	2.33 (0.08)	2.58 (0.11)	2.49 (0.32)	2.57 (0.11)
μ_{INS} ($\times10^9$)	0.26 (0.04)	0.25 (0.03)	0.26^x (0.02)	0.35 (0.03)	0.32 (0.04)	0.34^y (0.03)
μ_{DEL} ($\times10^9$)	0.42 (0.05)	0.39 (0.04)	0.40 (0.03)	0.59^a (0.07)	0.41^b (0.04)	0.50 (0.04)
μ_{INDEL} ($\times10^9$)	0.68 (0.06)	0.64 (0.05)	0.66^x (0.04)	0.95^a (0.06)	0.73^b (0.05)	0.84^y (0.05)
μ_{Total} ($\times10^9$)	2.94 (0.11)	2.99 (0.13)	$2.98^{x,*}$ (0.10)	3.52 (0.13)	3.21 (0.31)	$3.37^{y,*}$ (0.16)
μ_{GENOME}	0.29	0.30	0.30	0.35	0.32	0.34

All mutation rates are per-site, per-generation except μ_{GENOME}. Abbreviations are: μ_{bs}, base substitution mutation rate; $\mu_{INS,}$ insertion rate; μ_{DEL}, deletion rate; μ_{INDEL}, indel rate; μ_{Total}, total mutation rate; μ_{GENOME}, haploid genome-wide mutation rate per-generation.

[a,b] superscripts are significantly different (P<0.05) within O1MA or O2MA groups.

[x,y] superscripts are significantly different (P<0.05) between O1MA and O2MA.

* P≈0.06.

Supplementary Table S1. Individual mutations and their properties are listed in Supplementary

Table S2. Raw sequence data are archived in the NCBI Short Read Archive, project numbers

PRJNA395568 (O2MA lines and O2MA progenitors) and PRJNA429972 (other O1MA lines).

Averaged over all 32 O1MA lines, the per-nucleotide base-substitution mutation rate μ_{bs} = 2.33

(± 0.08) x 10^{-9} per generation. μ_{bs} for the nine O2MA progenitors is 2.26 ± 0.12 x 10^{-9} per

generation). Averaged over the 48 O2MA lines, the base-substitution mutation rate over the

subsequent ~150 generations is estimated to be μ_{bs} = 2.57 (± 0.11) x10^{-9} per-generation, not

significantly different from the combined O1MA rate (general linear model, $F_{1,14.5}$ = 1.32, p>0.26;

see Supplementary Appendix A1.6 for details of the GLM). The short indel rate for the full set of

32 O1MA lines μ_{INDEL} =0.66 (±0.04) x 10^{-9} per-site per-generation. The indel rate of the nine

O2MA progenitors ($\mu_{INDEL,ANC}$ = 0.68 ± 0.06 x 10^{-9}/generation) does not differ from that of the

other 21 O1MA lines ($\mu_{INDEL,OTHER}$ = 0.64 ± 0.05 x 10^{-9}/generation). Averaged over all 48 O2MA

lines, μ_{INDEL} = 0.84 ± 0.05 x10^{-9}/generation, significantly greater than the combined O1MA rate

(GLM, $F_{1,24.2}$ = 8.27, p < 0.01) but not significantly greater than that of the nine O2MA

progenitors (GLM, $F_{1,45.4}$ = 1.61, p > 0.21).

There is a positive correlation between the number of callable sites and the estimated

base-substitution mutation rate μ_{bs} ($r_{HIGH,O2MA}$ = 0.25, p < 0.22, $r_{LOW,O2MA}$ = 0.58, p < 0.002, r_{O1MA} =

0.30, p < 0.09). In the O2MA lines, callable sites are confounded with both O2MA family and

fitness. However, the correlation is driven by a few lines with atypically low coverage (average

number of callable sites less than 60% of the genome). The indel rate is uncorrelated with

callable sites ($r_{HIGH,O2MA}$ = 0.16, p > 0.44, $r_{LOW,O2MA}$ = 0.12, p > 0.58, r_{O1MA} = 0.00, p > 0.99). We

elaborate on potential artifactual and/or biological factors that may contribute to the correlation

in the Extended Discussion in Supplementary Appendix A2.1. We do not believe the results are

meaningfully affected by the correlation between μ_{bs} and callable sites.

No evidence for fitness-dependent base-substitution mutation.

There is significantly more variation in μ_{bs} among O2MA lines than expected if the base-

substitution mutation rate is uniform across the full set of 48 O2MA lines (simulation P<0.0001;

Supplementary Figure S1; see Methods and Supplementary Appendix 3.1 for details of

simulations). Moreover, there is significant variation in μ_{bs} among O2MA families (LRT, chi-

square = 9.27, df=1, P<0.003; Figure 2). Similarly, there is more variation among O1MA lines

than predicted by a uniform mutation rate (simulation P<0.015). The simplest interpretation is

that some element(s) of the mutational process diverged over the course of the first ~250

generations of MA, and that the signal of the difference(s) carried through the next ~150

generations of O2MA.

However, there is no evidence that μ_{bs} differs consistently between the O2MA families

derived from high fitness and low fitness O2MA progenitors (GLM, $F_{1,6.24} = 0.18$, p>0.75;).

Averaged over the five O2MA families in each fitness class, the base substitution mutation rate

between G250 and G400 for High and Low fitness lines is estimated to be $\mu_{bs,HIGH} = 2.58\ (\pm0.11)$

x 10^{-9}/ gen, and $\mu_{bs,LOW} = 2.49\ (\pm0.32)$ x 10^{-9}/ gen (Figure 2).

As expected from the lack of differentiation of μ_{bs}, the base-substitution spectrum does

not differ significantly between the high-fitness and low-fitness O2MA lines (Supplementary

Figure S2; Monte Carlo Fisher's Exact Test, 10^{7} replicates, P>0.70), nor does it vary between

O2MA families (MC FET, P>0.25), between individual O2MA lines (MC FET, P>0.08), between

O2MA lines within any of the ten families (P>0.10 or greater in all ten cases) or between the

O2MA progenitors at G250 and the O2MA lines at G400 (Supplementary Figure S2; MC FET,

P>0.51).

Consistent with many previous studies (Lynch 2007), the average mutation rate from a C

or G to an A or T is significantly greater than the mutation rate from A or T to C or G ($\mu_{C/G\to A/T} =$

3.03 (±0.18) x 10^{-9}/gen; $\mu_{A/T\to C/G} = 0.93\ (\pm0.05)$ x 10^{-9}/gen). Extrapolating from these rates, the

expected base composition of the *C. elegans* genome at mutational equilibrium is ~76.5% AT,

greater than the actual AT fraction of ~64.5%. Inspection of the μ_{bs} data reveals two potential

outlying O2MA lines (Figure 2), one low fitness (O2MA line 508.34, $\mu_{bs} = 0.78$ x 10^{-9}/gen) and

one high fitness (O2MA line 579.36, $\mu_{bs} = 4.34$ x 10^{-9}/gen). When those two lines are omitted

from the analysis, the variance among O2MA lines is sufficiently explained by a single base-

substitution rate (simulation P>0.07).

However, O2MA lines derived from O2MA progenitor 508 have the lowest base-

substitution rate even with the extreme line omitted [$\mu_{bs} = 1.49\ (\pm1.90)$ x 10^{-9}/gen with line

508.34 included, 1.68 (±0.91) x 10^{-9}/gen) without line 508.34] and O2MA lines derived from

O2MA progenitor 579 have the highest base-substitution rate even with the extreme line omitted

[$\mu_{bs} = 3.53\ (\pm0.31)$ x 10^{-9}/gen with line 579.36 included, 3.26 ($\pm$ 0.22) x 10^{-9}/gen) without line

579.36]. The random chance that the most extreme high line comes from the family whose

other four members also have the highest average mutation rate (5/48) and that the most

extreme low line comes from the family whose other four members also have the lowest

average mutation rate (5/47) is approximately 1%. The most parsimonious explanation is that

the two outlying O2MA lines are simply the most extreme manifestations of a biological process

common to their respective O2MA progenitors rather than true outliers. The alternative is that

Figure 2. Distribution of mutation rate among O2MA families. Families derived from high-fitness O2MA progenitors are in orange,

families derived from low-fitness O2MA progenitors are in green. Families 522 and 547 contain four sequenced O2MA lines, the

other families contain five O2MA lines. The horizontal line denotes the mutation rate of the nine O2MA progenitors. Points shown

outside the box are beyond the 1.5X inter-quartile range of the family whereas whiskers represent data points within that range. that

the mutational process in the ten O2MA progenitors diverged by ~250 generations of evolution under MA conditions, and the signal

of the difference(s) was retained over the subsequent ~150 generations of O2MA. (A, left) Base-substitution mutation rate (μ_{bs}). (B,

right) Indel mutation rate (μ_{INDEL}).

the apparently extreme mutation rates are experimental artifacts, which we think is unlikely (see

Extended Discussion in Supplementary Appendix A2).

Fitness-dependence of the small indel rate.

In contrast to the base-substitution mutation rate, which does not differ between O2MA

lines derived from high fitness and low fitness O1MA lines, the O2MA short indel rate is

significantly greater in the high fitness group (μ_{INDEL} = 0.95 ± 0.06 x10^{-9}/generation) than in the

low fitness group (μ_{INDEL} = 0.73 ± 0.50 x10^{-}/generation: GLM, $F_{1,44.4}$ = 7.84, P<0.01). The

difference is primarily due to different rates of deletions ($\mu_{DEL,High}$ = 0.59 ± 0.07 x10^{-9}/generation,

$\mu_{DEL,Low}$ = 0.41 ± 0.04 x10^{-9}/generation; $F_{1,7.58}$ = 5.77, P < 0.05) rather than insertions ($\mu_{INS,High}$ =

0.35 ± 0.03 x10^{-9}/generation, $\mu_{INS,Low}$ = 0.32 ± 0.04 x10^{-9}/generation; $F_{1,17.5}$ = 0.51, P > 0.48).

The variance in μ_{INDEL} among O2MA lines within each fitness group is adequately explained by a

single, fitness-specific indel rate (high-fitness, simulation P>0.12; low-fitness, simulation P>0.2).

The distribution of indel lengths is given in Supplementary Figure S3.

The higher indel rate of high-fitness O2MA lines suggests that the indel rate of O1MA

lines should be greater than the overall O2MA rate, given the higher fitness of the G0 ancestor.

Interestingly, that is not what we observe. The overall O1MA indel rate, including all 32 O1MA

lines, is significantly lower than high-fitness O2MA indel rate ($\mu_{INDEL, O1MA}$ = 0.66 ± 0.04 x10^{-}

^{9}/generation, $\mu_{INDEL, O2MA\_High}$ = 0.95 ± 0.06 x10^{-10}/generation; GLM, $F_{1,41.7}$=15.61, P<0.0005), but

not significantly different from low fitness O2MA indel rate ($\mu_{INDEL, O1MA}$ = 0.66 ± 0.04 x10^{-}

^{9}/generation, $\mu_{INDEL, O2MA\_Low}$ = 0.73 ± 0.05 x10^{-9}/generation; $F_{1,47.6}$ = 1.10, P>0.29). The results

do not change if only the nine O2MA progenitors are used to calculate the indel rate.

The O2MA insertion and deletion rates are uncorrelated ($r_{INS,DEL}$ = 0.021, P>0.88, n=48),

suggestive of different factors underlying the two types of mutations. The base-substitution rate

is moderately positively correlated with the insertion rate ($r_{bs,INS}$ = 0.32, P<0.03, n=48) but

uncorrelated with the deletion rate ($r_{bs,DEL}$ = -0.016, P>0.91, n=48).

There are two potential evolutionary factors that can explain the difference in the

deletion rate between the different O2MA fitness groups. First, some element of the mutational

process may differ, e.g., DNA repair. Alternatively, selection may differ in either strength or

efficiency between the two treatments. In our MA protocol, differences in selection efficiency

between lines can result from different frequencies of going to backup (see Methods). However,

the mean effectively neutral selection coefficient (s_n =$1/4N_e$) is only slightly smaller in the low-

fitness lines ($\bar{s}_n$ = 0.228) than in the high-fitness lines ($\bar{s}_n$ = 0.239). The slightly lower efficiency

of selection in the high fitness group seems unlikely to account for the ~30% greater indel rate.

Alternatively, the strength of selection itself may be different between the two fitness

groups, such that some mutations that are only mildly deleterious – and thus effectively neutral

– in a high-fitness line are significantly more deleterious in a low-fitness line, i.e., epistasis is

synergistic (= negative; Phillips 2008). In that case, a larger fraction of mutations in the low-

fitness lines would exceed the threshold of effective neutrality and be removed by selection.

The mutational effect analysis, as implemented in snpEff 4.1 (see Methods), reveals no

significant difference in the frequency of potentially large-effect indels (20/303 among high-

fitness O2MA lines, 10/186 among low-fitness lines, Fisher's Exact Test, two-tailed P>0.70) or

SNPs (22/822 vs. 15/659, Yates' Chi-square, two-tailed P>0.75). Nor does the frequency of

potentially large-effect mutations differ between the high and low-fitness G250 O1MA lines,

between high-fitness O2MA and O1MA lines, or between low-fitness O2MA and O1MA lines

(P>0.12 in all cases). The preceding analysis provides no information about mutations that did

not fix. Nevertheless, synergistic epistasis, coupled with a slight overall increase in mutation

rate, seems a more parsimonious explanation than different indel rates in high-fitness and low-

fitness O2MA lines. Experimentally derived estimates of relative fitness (Matsuba et al. 2012)

combined with estimates of the average number of mutations carried by high-fitness and low-

fitness O2MA lines reinforce this conclusion (see Discussion).

Mutation is positively associated with recombination

Standing nucleotide diversity is almost always positively associated with recombination

rate, which could in principle be due either to the mutagenicity of recombination or to natural

selection (i.e., Hill-Robertson interference). There is good reason to think that HR interference

has an important causal role in this pattern (Lynch 2007). However, with the important

exception of humans, most of the data are derived from inferences drawn from comparisons

with a reference class of genomic sites believed to be free from selective constraints (e.g.,

processed pseudogenes or four-fold degenerate sites) rather than directly from *de novo*

mutations, and the extent to which recombination is mutagenic *per se* remains an open

question. Several large studies of *de novo* mutations in humans report a significant association

between recombination rate and mutation rate (Michaelson et al. 2012; Francioli et al. 2015;

Carlson et al. 2018), although recombination appears to affect nucleotide diversity beyond its

association with mutation. Two cohort studies in bees (Yang et al. 2015; Liu et al. 2017) found

a weak relationship between recombination and mutation, but the numbers of mutations were

small and the inferences somewhat circumstantial. Conversely, Ness et al. (2015) found no

association between recombination rate and mutation rate in the green alga *Chlamydomonas*

reinhardtii, although the data on recombination rate in *C. reinhardtii* are sparser than for

humans. In *C. elegans*, the relative proportions of different types of variants (SNPs, small and

large indels) in the standing genetic variation differ depending on the local recombination rate,

which has been suggested to reflect the signature of recombination-dependent mutation

(Hwang and Wang 2017).

To investigate the relationship between recombination and mutation, we determined the

association between μ_{bs} estimated from the O2MA lines and recombination rate using weighted

OLS regression, dividing each chromosome into the recombination rate bins reported by

Rockman and Kruglyak (2009) and weighting each bin by its size in Mb. Recombination rate is

nearly constant within each bin (see Figure 1 in Rockman and Kruglyak (2009)). Contrary to our

previous report from a different (and smaller) subset of G250 O1MA lines (Denver et al. 2009),

here we find a significant positive univariate association between recombination rate and μ_{bs}

(pseudo-r^2 = 0.26, P<0.003, Supplementary Figure S4); presumably the discrepancy between

this study and the previous report is due to the greater power afforded by the much greater

number of mutations included in the present study (316 vs. 1828).

The genomic correlates of mutability

Many features of the genome and epigenome influence the mutational process. The

effects of some such features are well-understood and seem to be relatively general to all living

organisms (e.g., short tandem repeats, G:C vs. A:T, 5'-methyl-C), whereas others remain

uncertain and/or appear to be taxon-specific. To more fully characterize the features of the *C.*

elegans genome that are associated with the mutational process, we employed a logistic

regression method, loosely following the approach of Michaelson et al. (2012) and Ness et al.

(2015). Since the deletion rate differs significantly between the two O2MA fitness groups, we

restricted the analysis to base-substitutions. Univariate logistic regression coefficients of the

features included in the full multiple regression are shown in Figure 3.

To assess the predictive power of the mutability model, we randomly sampled half of the

O2MA lines to train the model (24 lines; roughly 740 mutations) and tested the model on the

remaining 24 O2MA lines. All mutant sites and 100,000 randomly chosen non-mutant sites

were arbitrarily binned into 35 bins of uniform width, and the observed mutation rate for each

mutability bin was plotted against expected mutability. This analysis was repeated 100 times.

Of the factors initially included in the multiple regression (see Methods), the best model includes

only the 64 three-base motifs as a set of predictor variables (Figure 4a). Any combination of

other predictors, with or without the three-base motif, results in a poorer fit. The poorer fit

presumably results from either overfitting, multicollinearity and/or the inability of the logistic

regression model to accommodate non-linear relationships between predictor variables and

mutation rate, even when tuned with high penalty (λ) to drop predictor variables altogether

(Lasso). Logistic regression coefficients are tabulated in Supplementary Table S3.

It is reassuring but not surprising that the model provides a good fit to the MA data from

which it was trained. Of more interest is the relationship between mutability as predicted from

MA data and the genetic variation observed in nature. We obtained publicly-available whole

genome sequence data from 40 *C. elegans* wild isolates (Thompson et al. 2013) and identified

SNPs using the same pipeline that we used to call putative mutations in the MA data. We

identified ~537,000 SNPs by these criteria. Sites were categorized as variable or not variable,

without regard to allele frequency.

Mutability was assessed as described previously, except in this analysis the model was

trained on the full set of 1828 mutations. Figure 4b shows a plot of nucleotide diversity

(quantified as θ_W; (Watterson 1975)) at non-coding sites (288,585 intron sites, and 122,272

intergenic sites) against predicted mutability. Averaged over all bins, mutability strongly predicts

standing nucleotide diversity, although the variance is high at predicted high-mutability sites,

presumably because the sample sizes are small.

Short Tandem Repeats (STRs)

Short tandem repeat loci ("microsatellites") can mutate orders of magnitude faster than

other classes of loci, and potentially contribute a large fraction of the per-generation mutational

variance. We previously estimated the haploid per-genome mutation rate of dinucleotide STR

loci in the full set of O1MA lines to be ~0.12/generation (Phillips et al. 2009). That calculation

accounts for variation in mutation rate among repeat motifs (e.g., AT vs. AG, etc.) and at least

partly accounts for variation in mutation rate with repeat number, although there is substantial

uncertainty that cannot be easily quantified. Seyfert et al. (2008) found no significant effect of

repeat length (di, tri, or tetranucleotide repeat) on the rate of STR mutation in a different set of

N2-strain MA lines. Denver et al. (2004) investigated the mutational process of mononucleotide

repeats (= "homopolymers") in the same MA lines as Seyfert et al. and concluded that the

(haploid) per-genome mononucleotide mutation rate is about 0.8/generation, more than twice

the rate of all other classes of mutations combined. Mutational properties of STRs are

summarized by repeat type in Supplementary Table S4 and Supplementary Figures S5 and S6.

Figure 3. (A, Left panel) Univariate logistic regression coefficients for different genomic features (gray bars). Overlapping red bars depict the null expectation from sampling variance with 1828 randomly selected genomic sites treated as mutations. Variable abbreviations are: GC_10, 10 bp GC-content centered on the focal base; GC_1000, 1 kb GC content; mono_di, proximity to a mono or dinucleotide STR; motif_3bp, three-base motif with the mutant base at the 3' end; Rec, local recombination rate. The method used to condense several predictors into one, for 3-base motif and mono-di STR, is described in the Methods. (B, Right panel) Base-substitution mutation rate (μ_{bs}) of each 3-base motif, with mutations on the 3' end. Motifs are grouped by the mutant base.

Figure 4. Left panel (a) depicts the relationship between observed and predicted mutability

across mutability bins (n=35 bins). The mutability model includes only the 3-base motif. Right

panel (b) depicts the relationship between observed standing nucleotide variation, measured as

Watterson's θ, and predicted mutability. Predicted mutability of wild isolates is calculated from

the full complement of mutations in the MA lines (n=1828).

Our findings qualitatively recapitulate those of the previous studies. First, G:C

mononucleotides experience indel mutations at a tenfold greater rate than A:T

mononucleotides, as observed by Denver et al. (2004). Second, indel mutation rate differs only

about twofold between the three A:T containing dinucleotides (AC, AG, AT) and we detected no

mutations at CG dinucleotides, both as observed by Phillips et al. (2009). The ratio of

mononucleotide deletions (251) to insertions (139) is not significantly different from the 16:14

ratio reported by Denver et al. (Yates' chi-square = 1.02, df = 1, P > 0.31), and the ratio of

dinucleotide deletions (10) to insertions (32) is nearly identical to the 8:28 ratio observed by

Phillips et al. However, there are quantitative differences between the findings of this study and

the previously reported values. The mononucleotide indel rate is estimated to be 0.056

mutations per haploid genome per generation, about 10% the rate estimated by Denver et al.

(2004). The dinucleotide indel rate is estimated to be 0.003 mutations per haploid genome per

generation, about 5% of the estimate from Phillips et al. (2009).

We believe the differences between the STR mutation rates estimated in this study and

those reported previously are more apparent than real, and that the higher genome-wide

estimates from the previous studies are probably closer to the truth. The previous studies

included only a small number of STR loci, in which extremely long repeats were greatly

overrepresented. In contrast, long repeats are significantly underrepresented in our data.

There is no reason to doubt that mutation rate increases with repeat number. Probably, we

have missed a handful of highly mutable loci that contribute a disproportionately large fraction of

mutations to the genome-wide total. However, very long mono- and dinucleotide repeats

represent a miniscule fraction of the genome (e.g., mono- and dinucleotides > 20 repeats

represent approximately 0.02% of the genome), and it seems unlikely (although not impossible)

that those few highly mutable loci contribute meaningfully to fitness. Thus, although the true

genome-wide mutation rate may in fact approach one per haploid genome per generation, the

meaningful mutation rate may be much lower.

We elaborate on potential causes of the source(s) of the discrepancies between this

study and the previous studies in the Extended Discussion, Supplementary Appendix A2.3.

Copy Number Variants (CNVs)

CNVs were called using the read-depth based method implemented in the CNV-seq

software (Xie and Tammi 2009). The number of putative CNVs inferred is sensitive to the

parameters of the analysis, with different sets of input parameters leading to estimates of the

CNV rate that differ by an order of magnitude. Sensible interpretation of the results requires

understanding the details of the analysis, which we outline in the Methods and elaborate in

section 4 of the Extended Discussion, Appendix A2. There are two salient general results: (1) in

no case does the mean number of putative CNVs differ between the high and low-fitness O2MA

lines, but also (2) the number of putative CNVs in O2MA lines is consistently greater than in

O1MA lines (Supplementary Table S5). Given the uncertainty associated with the estimates,

we did not attempt to confirm CNVs (e.g., with qPCR). We report the results as a cautionary

note that, even with as "easy" a genome as *C. elegans* N2 strain MA lines, which are almost

completely homozygous and have an exceptionally well-characterized reference genome,

estimates of CNVs from short-read data are extremely sensitive to methodological details. We

expect that the CNV problem will eventually be resolved with accurate long-read sequencing at

high coverage (Chaisson et al. 2015; Tyson et al. 2018).

Discussion

The Effects of Fitness. (i) Mutation Rate

There is no evidence that the rate or molecular spectrum of base-substitution mutations

in the N2 strain of *C. elegans* depends on the fitness of the starting genotype. In contrast, the

short indel rate, especially the deletion rate, is fitness dependent, but not in the anticipated way:

low-fitness genotypes have a significantly lower deletion rate than do high-fitness genotypes.

This result differs from the finding in *Drosophila melanogaster* that low-fitness genotypes

experience significantly greater rates of small deletions than do high-fitness genotypes,

apparently because flies in poor physiological condition employ a different, more error-prone

mechanism of double-strand break repair than do flies in good condition (Sharp and Agrawal

2016).

The Effects of Fitness. (ii) Mutational Effects

In the abstract, existence of a robust system seems to imply redundancy of components,

which in the context of a genetic system implies that epistasis must be synergistic on average

(de Visser et al. 2003). However, empirical evidence concerning the average epistatic effect of

spontaneous deleterious mutations has been inconclusive (Halligan and Keightley 2009), and

whether deleterious mutations interact synergistically *on average* has vexed generations of

evolutionary biologists. .

It is certainly possible that worms with high-fitness genotypes incur more short deletions

than do worms with low-fitness genotypes and/or are worse at repairing them. If we use the

point estimates of the average number of mutations carried by O2MA lines derived from high-

fitness and low-fitness O2MA progenitors and the average mutational decline in relative fitness,

averaged over all O2MA lines (Supplementary Table S6), the average mutational effect on

relative fitness of the low-fitness O2MA lines is about twice that of the high fitness lines (0.48%

vs 0.25%). If we assume that only indels affect fitness, the mean effects on low-fitness and

high-fitness O2MA lines are about 2% and 1%, respectively. We further suppose that the "dark

matter" represented by the (assumed) missing fraction of indels in the low-fitness lines would

amplify the difference in average selective effects. These results represent some of the first

direct evidence that spontaneous mutations interact synergistically, on average (see also

Jasmin and Lenormand 2016).

Variation in the Mutational Process

430   The per-generation input of genetic variance for a trait, the mutational variance, V_M,

ultimately governs the evolvability of the trait and is a fundamental parameter in evolutionary

genetics (Lynch and Hill 1986). V_M can be estimated from MA data from the relationship

$V_M = \frac{V_L}{2t}$, where V_L is the among-line component of variance and t is the number of generations

of MA (Lynch and Walsh 1998). V_M is commonly standardized relative to the residual variance

V_E, called the mutational heritability $h_M^2 = \frac{V_M}{V_E}$. Among O2MA families, V_L for μ_{BS} = 0.196, V_E =

0.316, and $\bar{t}$ = 143, so $h_M^2 \approx$ 0.002/generation. To put that result in context, h_M^2 for a wide variety

of traits in disparate taxa averages about 0.001/generation (Houle et al. 1996), although h_M^2 for

some traits (notably gene expression) is consistently an order of magnitude less (e.g., Rifkin et

al. 2005; Landry et al. 2007). That the mutation rate is evolvable is not surprising (we know it is),

but the point estimate of h_M^2 suggests that the mutational target for mutation rate is substantial.

However, an estimate of a ratio of variances based on ten data points cannot be considered

robust.

Mutability and Genetic Variation

445   The observed positive univariate association between local recombination rate and

mutation rate in this experiment is almost surely not due to Hill-Robertson interference. The

expected time to fixation/loss of a new mutation is three generations (Keightley and Caballero

1997), and the average mutation rate (CNVs notwithstanding) is about one per genome every

three generations. Thus, the opportunity for H-R interference in our experiment is very low,

albeit not nonexistent. However, the causal factors underlying the relationship between local

recombination rate and base-substitution mutation rate remain uncertain. The cause is not as

simple as GC content (G:C being more mutable than A:T), because chromosome arms, which

have higher recombination rates, are AT-rich. The sufficiency of 3-bp motif as a predictor of

mutability suggests that the positive association of mutation with recombination is an

epiphenomenon resulting from the overrepresentation of highly mutable motifs in regions of high

recombination. Most clearly, mononucleotide runs are both mutagenic and positively associated

with local recombination rate.

The mutability model does a good job of predicting standing nucleotide variation in

natural isolates of *C. elegans*: sites that are more mutable are, on average, more variable

(Figure 4b). Again, this is reassuring, but not surprising; the same relationship has been

observed in *Chlamydomonas reinhardtii* (Ness et al. 2016) and humans (Francioli et al. 2015).

One difference between *C. elegans* and many other organisms (e.g., humans) is that, in *C.*

elegans, regions of low recombination (chromosome cores) are gene-rich rather than gene-

poor, so H-R interference is more important in regions of low recombination both for inherent

reasons and because the opportunity for selection is greater due to the larger target for

deleterious mutations. One possibility is that, because chromosome arms are gene-poor,

mutagenic features (e.g., specific motifs, such as mononucleotide runs) preferentially

accumulate because their background effects on linked loci are less important. Perhaps

paradoxically, the AT-richness of chromosome arms may be a signature of increased mutation

rate, because C:G mutates to A:T more often than the reverse.

Transition/Transversion ratio in MA and wild isolates

The ratio of transitions to transversions does not differ significantly between high fitness

O2MA and low fitness O2MA lines (Ts/Tv$_{High}$=0.72, Ts/Tv$_{Low}$=0.68, *t*-test, P>0.5), nor does it

differ between O2MA lines and O2MA progenitors (Ts/Tv$_{O1MA}$=0.70, Ts/Tv$_{O2MA}$=0.74, *t*-test,

P>0.8). As in previous studies with *C. elegans* MA lines, the Ts/Tv ratio in MA lines is less than

the Ts/Tv ratio in wild isolates (Denver et al. 2009; Denver et al. 2012). The difference could be

due to purifying selection against transversions (Babbitt and Cotter 2011), or it could be that the

mutational milieu in the lab environment differs from that in the wild. Ts/Tv of rare variants (<5%

variant allele frequency relative to the reference genome, Ts/Tv = 1.20) is similar to that of

common variants (>50% variant allele frequency, Ts/Tv = 1.14). That finding is not consistent

with purifying selection against transversions, which suggests that some feature of the lab

environment may cause the mutational spectrum of MA lines to differ from that in nature. An

independent test of this possibility is to compare Ts/Tv between our N2-strain MA ancestor and

the N2 reference genome; any variants will have arisen in the lab environment. In that

comparison, Ts/Tv = 0.78, consistent with the MA Ts/Tv ratio and different from that among wild

isolates. Denver et al. (2012) found a similarly lower Ts/Tv in MA lines of the congener *C.*
briggsae relative to wild isolates, suggesting that the phenomenon is not restricted to *C.*
elegans.
490
Evolution of the mutation rate.
The total mutation rate, μ_{Total}, of the O2MA lines is about 13% greater than that of the
O1MA lines (Table 1). Directional change in a trait under MA conditions ("mutational bias", ΔM)
suggests that the trait is under ongoing directional selection in the opposite direction, analogous
to the direction of phenotypic change upon inbreeding (Teotónio et al. 2017). This finding is
consistent with the "drift barrier" hypothesis of mutation rate evolution, which posits that
directional selection to reduce the mutation rate is opposed by a weak mutational bias (Lynch
2008). It is not consistent with the mutation rate being at an optimum established by a "cost of
fidelity", wherein direct selection to reduce the input of deleterious mutations is counterbalanced
by indirect selection to reduce the fitness cost of genome surveillance (Kimura 1967). If the
mutation rate is at an optimum imposed by countervailing components of selection, the overall
fitness function will be stabilizing. Provided that the fitness function is approximately
symmetrical around the optimum, the expectation is that the among-line variance in the trait will
increase but that the overall trait mean will not change. We emphasize that these findings do
not imply that there is no cost of fidelity, just that the mutation rate in *C. elegans* does not
appear to be at a global optimum.
The ~13% increase in mutation rate over the course of ~250 generations amounts to a
per-generation change ΔM ≈0.0005, compared to a per-generation decrease in competitive
fitness of ≈0.001 in the same lines (Yeh et al. 2018). It is difficult to believe that cumulative
selection on competitive fitness is only twice as strong as cumulative selection to decrease
mutation rate. The most logical conclusion is that mutations that increase mutation rate have
deleterious pleiotropic effects on fitness. However, this conclusion seems at odds with the
failure to observe a main effect of fitness in the O2MA lines. In principle, the discrepancy could
be resolved by determining the mutational correlation of mutation rate with fitness. The sample
sizes necessary to answer that question in multicellular organisms are currently prohibitive, but
it may be practical in a microbial system.
517
Conclusions. Sets of second-order MA lines that differed significantly in relative fitness were
allowed to accumulate mutations for ~150 generations, to test the hypothesis that physiological
stress imposed by mutation load causes the mutation rate to increase. Contrary to our

hypothesis, the base substitution rate did not differ significantly between high-fitness and low-

fitness lines, whereas the indel rate was greater in the high-fitness lines. Averaged over all

lines, the total mutation rate between generations 250-400 was significantly greater than the

mutation rate between generations 0 and 250, as predicted by the "drift barrier" hypothesis of

mutation rate evolution. The average mutational effect on fitness (α) is approximately twice as

great in the low-fitness lines, which implies that the epistatic effects of deleterious mutations are

synergistic.

Methods and Materials

Mutation Accumulation Protocol. A schematic depiction of the experimental design is presented

in Figure 1. Details of the first-order MA protocol and fitness assays are reported in Baer et al.

(2005); details of the second-order MA protocol are reported in Matsuba et al. (2012) and

summarized in the Extended Methods, Supplementary Appendix A1.1.

Genome sequencing and estimation of mutation rates. Five O2MA lines from the ten O2MA

families and 35 O1MA lines, including the ten O2MA progenitors, were sequenced at an

average of ~25X coverage depth. Sequencing was done using Illumina technology with 100 bp

paired-end reads. Protocols for DNA extraction and construction of sequencing libraries are

given in Supplementary Appendix A1.2; details of preliminary processing of raw sequence data

are given in Supplementary Appendix A1.3. The quality of sequence from two O2MA lines and

one O1MA line was poor and these samples were omitted from further analyses, leaving 48

O2MA line and nine of the ten O2MA progenitors.

Variants were called using GATK software (McKenna et al. 2010) with a minimum

coverage threshold of >10X. Variants were identified as putative mutations if (1) the variant was

identified as homozygous, and (2) it was present in one and only one O2MA line. Criterion (1)

means that any mutations that occurred in the last few generations of second-order MA that

were still segregating and/or occurred during population expansion for DNA extraction were

ignored. Because the O1MA progenitor was at mutation-drift equilibrium (Lynch and Hill 1986),

the segregating variation is expected to be the same in the O1MA progenitor and the O2MA

line, so ignoring heterozygotes results in an unbiased estimate of mutation rate. Criterion (2)

reduces the probability of mistakenly identifying a variant segregating at low frequency in the

expanded population of the O1MA progenitor as a new mutation. Two pairs of O1MA lines

shared multiple variants and were inferred to have experienced contamination at some point

during the MA phase; one line from each pair was arbitrarily omitted from subsequent analyses

(see Extended Discussion, Supplementary Appendix A2.1).

The mutation rate (per-site, per-generation) μ of each O2MA line was calculated as m/nt

where m is the number of mutations, n is the number of nucleotide sites observed and t is the

number of generations of MA (Denver et al. 2009). The average mutation rate of each O1MA

progenitor was calculated as the unweighted mean of the O2MA lines in that family.

We applied three additional variant-calling strategies, one with a more stringent set of

criteria which we refer to as the "trimmed genome" and two with more liberal criteria, which we

refer to as the "lenient genome" and "STR-relaxed", respectively. We also explored several

alternative GATK filters. Details of the alternative GATK filters are given in Supplementary

Appendix A1.3; justification and details of the additional variant-calling strategies are given in

Supplementary Appendix A1.4.

Mutation Confirmation - From each of the 48 O2MA lines, we randomly chose one putative

base-substitution and one putative indel mutation for confirmation by Sanger sequencing.

Details of the confirmation protocol are given in Supplementary Appendix A1.5. We confirmed

43/48 putative base-substitutions (zero false positives, five failures) and 36/48 putative indels

(zero false positives, 12 failures), consistent with a false positive rate below 2.5% based on the

upper 95% Poisson confidence limit. The number of failures did not differ significantly between

base-substitutions and indels (Fisher's exact test, two-tailed P>0.1).

Data Analysis

i) Variation among O1MA and O2MA lines - The simplest hypothesis regarding the mutational

process is that it remained constant over the course of the experiment subsequent to the

divergence of the O1MA lines from the common G0 ancestor. To test the hypothesis that a

uniform mutation rate sufficiently explains variation among MA lines, we simulated the

evolutionary process of 48 O2MA lines with a uniform base-substitution mutation rate equal to

the unweighted mean base-substitution mutation rate of the 48 O2MA lines, accounting for the

number of callable sites (Supplementary Table S2) and the number of generations of second-

order MA of each line (Supplementary Table S1). At each generation, the simulated genome

(100.25 Mb) of a line was assigned a number of mutations drawn from a Poisson distribution

with parameter equal to the unweighted mean mutation rate. Next, each simulated line was

assigned a number of generations of MA and a number of callable sites sampled with

replacement from the observed distribution. Sets of 48 simulated lines were generated 100,000

times, and the observed variance in mutation rate among the 48 O2MA lines was compared to

the distribution of simulated variances (Supplementary Figure S1). An analogous simulation

was done for the 34 O1MA lines. Details of the simulation and code are provided in

Supplementary Appendix A3.1.

To test for effects of fitness on mutation rate and to partition the variance in mutation

rate into within and among-group components, we consider the mutation rate itself as a

continuously-distributed dependent variable in a general linear model (GLM). Details of the

GLM analyses are given in Supplementary Appendix A1.6.

It is possible that the distribution of mutation types – the mutational spectrum - differs

among groups even if the overall mutation rate does not (Long et al. 2016), or if none of the

type-specific mutation rates achieves statistical significance. Spectra were compared among

groups by Fisher's Exact Test using Monte Carlo sampling as implemented in the FREQ

procedure of SAS v. 9.4.

ii) Mutability – Many factors potentially influence the probability that a site will mutate. Some of

these factors can be unambiguously characterized from pooled genomic DNA from a population

of multicellular organisms (local sequence motif, base composition at various scales, local

recombination rate), whereas other factors that potentially influence heritable mutation are only

relevant in the context of the germline or its embryonic precursors (e.g., chromatin state,

nucleosome occupancy, expression level). To elucidate the relationship between the various

genomic properties of a given site and the probability that a mutation occurs at that site, we

employed a logistic regression model in which the log odds-ratio that a mutation occurs at a

given site (success=mutation) is modeled as the sum of a set of linear predictor input variables

(Michaelson et al. 2012; Ness et al. 2016). We initially included as independent variables the

three-base motif with the focal nucleotide at the 3' end and at the 5' end (these need not be

redundant if the probability that a sequencing read is included is not identical for the two

strands), the five-base motif with the focal nucleotide in the center, local recombination rate (see

Results for details), presence or absence of the focal site in the vicinity of a (mono/di)nucleotide

run ($\pm$ 2 bp), and the 11-base and 1001-base GC-content, centered on the focal site. For cross-

validation, we trained our models using half of our O2MA lines (n=24), including all base

substitutions and 100,000 randomly chosen non-mutant sites, and tested the predictions on the

remaining 24 O2MA lines. Following model selection, we trained the model using all the

ancestral O1MA and O2MA mutations (n=1828). Including only SNPs from the O2MA lines

(n=1481) had no qualitative effect on the results.

Logistic regression was performed using the GLMnet package in R (Friedman et al.

2010). The two model parameters are the tuning penalty λ and the ridge/lasso penalty α. As

$\alpha \rightarrow 0$ (ridge), the model tends to shrink the coefficients of correlated predictor variables toward

each other without dropping any of the predictor variables. As $\alpha \rightarrow 1$ (lasso), when predictor

variables are correlated the model chooses one and discards the other(s). The tuning

parameter λ controls the overall strength of the penalty. For all the models we tested, λ was

chosen by the package's built-in cross validation function ("lambda-min"). The fit of our models

remain largely unchanged by the selection of α, with the exception of instances where $\alpha \rightarrow 0$

(ridge). For values of α sufficiently close to 0, we observed twice the slope (bias, or regression

coefficient; expected value = 1) expected when the observed mutation rate is regressed against

the predicted mutation rate. All results presented here used $\alpha = 0.05$.

Models including short tandem repeats (STRs), five base motif and/or 10bp GC content

together with three-base motif fit poorly. The poor fit could be due either to overfitting,

multicollinearity and/or a non-linear relationship between a predictor variable and mutability.

Moreover, models that included STRs as the only predictors fit reasonably well in terms of

calibration of the model, but suffer from poorer predictive discrimination, as they lack any

discrimination in the non-STR region of the genome (~97% of the genome). The final set of

models tested include only the three-base motif with the focal nucleotide at the 3' end, 1Kb GC

content and local recombination rate. For each set of input predictor variables and α, the dataset

was resampled 200 times, with a different randomly chosen set of 100,000 non-mutant sites,

and divided into halves, the training set and the test set. The training set consisted of half the

O2MA lines, and the test set consisted of the other half. For each data partition, the model was

trained and the fit to the test set was assessed by the fraction of the variance explained by the

linear regression of the observed (test) value on the predicted (training) value (R^2), the

regression coefficient (bias, expected value = 1), and the area under the ROC curve (AUC, or c-

statistic) for predictive discrimination (varies between 0 and 1). For each set of α and input

variables, we generated a distribution of 200 R^2 values, bias values and AUC values, and

retained model(s) that maximized R^2, and AUC while kept the regression coefficient to as close

to one as possible. Code and example files are provided in Supplementary Appendix A3.2. We

list the performance of all the models we tested in Supplementary Table S7.

GLMnet fits the following regression model utilizing every site in the training set, and

resulting mutability is the probability of mutation "Pr (mutant)" in the following equation.

$$\log \left(\frac{\Pr(mutant)}{\Pr(non-mutant)} \right) = \beta_0 + \beta^T x$$

A mutability value of (say) 0.08 for a specific set of features implies that 92 out of 100 sites with

the same features in the training set are non-mutant. The absolute value of mutability varies

positively with the number of mutations in the training set and negatively with the number of

non-mutant sites included during training. Mutability can only be interpreted in the relative

context of the study. For example, Ness et al. (2015) included ~6500 mutations in a training set

with 100,000 non-mutant sites, so the average mutability in that study would be ~3.5X greater

than the estimate in this study.

Although mutability potentially varies between 0 and 1, the highest mutability observed

rarely exceeded 0.2, as average mutability is the fraction of mutant sites as compared to the

total number of sites. During cross-validation, we trained the model on slightly under half of the

total mutations (~740 mutants; half of all O2MA mutants, and none of the O1MA mutants), and

100,000 randomly selected sites, yielding the expected value of mutability to be ~0.007

(740/(100000+740)). Following model selection, to assess the relationship between standing

genetic variation and mutability, we recalculated mutability using all 1828 mutations. Expected

mutability calculated from the full data set is ~0.017 (1828/(100000+1828)).

To assess the relative contribution to mutability of different genomic features, we took a

slightly different approach. It is not possible to condense 64 3-base motifs into a single

predictor in a standard logistic regression framework. Instead, we trained a logistic regression

model using only the 3-base motif, and used the predictions made from this model as predictors

in a second logistic regression, which yields a single coefficient for 3-base motif. We then

selected 1828 random sites from the genome, treated them as mutations, and repeated the

process 100 times to obtain the null distribution for these coefficients.

iii) Copy Number Variants - We called putative copy-number variants (CNVs) using a read-depth

(RD) based approach as implemented in the CNV-Seq software (Xie and Tammi 2009). Read-

depth outliers in an MA line (O1MA or O2MA) are identified relative to its progenitor. We

employed a sliding window approach, in which read-depth in a given window in the focal sample

is compared to read-depth in the reference (progenitor), by taking the ratio of total reads falling

in that window, normalized by the average read-depth for that chromosome in that sample. We

used two different signal thresholds (1.5x/2x) for calling duplications or deletions, and several

sets of sliding window sizes (see Supplementary Appendix 2, section 4 for details). The

minimum detectable size of a CNV is 1kb in all cases. A putative CNV is called only when all

sliding windows covering an interval meet the signal threshold. The smaller the window, the

more conservative the test, because more consecutive windows must meet the signal criterion.

Similarly, the greater the signal threshold, the more conservative the criterion. As with other

types of variants, only putative CNVs unique to a single O2MA line are considered potential

mutations. For reasons which we elaborate in the Results, we did not attempt to validate CNVs.

iv) Predicted Mutational Effects –To broadly categorize the distribution of predicted mutational

effects, we used the publicly available software snpEff 4.1 (Cingolani et al. 2012) to annotate

variants with respect to a set of characteristics potentially related to fitness

(http://snpeff.sourceforge.net/VCFannotationformat_v1.0.pdf). The snpEff software uses gene

lists (gtf/gff3) from the WS234 build of the *C. elegans* genome to assign a putative effect score

of high, moderate, or low based on these characteristics, where a "high" effect variant is most

likely to have a deleterious effect on fitness. Each variant can have multiple annotations due to

different alleles and/or different splice variants; we included only the largest potential effect of

each variant. Effects were parsed using custom AWK scripts and categorized as having low,

medium, or high maximum impact.

Acknowledgments

We thank Tim Crombie, Joanna Dembek, Joanna Joyner-Matos, Gigi Ostrow, Asher Shoucair

and the many undergraduate students who worked over the years to maintain and propagate

the MA lines. We thank Rob Ness and Jonathan Sebat for graciously explaining some details of

the analyses in their papers, and Mike Miyamoto, Henrique Teotónio, and the anonymous

reviewers for their helpful comments. This work was supported by NSF grant DEB-0717167 to

CFB and NIH grants R01GM072639 to CFB and D. R. Denver and R01GM107227 to CFB and

E. C. Andersen.

Literature Cited

Agrawal AF. 2002. Genetic loads under fitness-dependent mutation rates. *Journal of Evolutionary Biology* **15**(6): 1004-1010.

Agrawal AF, Wang AD. 2008. Increased transmission of mutations by low condition females: evidence for condition-dependent DNA repair. *PLoS Biology* **6**: 389-395.

Ávila V, Chavarrias D, Sanchez E, Manrique A, Lopez-Fanjul C, Garcia-Dorado A. 2006. Increase of the spontaneous mutation rate in a long-term experiment with *Drosophila melanogaster*. *Genetics* **173**(1): 267-277.

Babbitt GA, Cotter CR. 2011. Functional conservation of nucleosome formation selectively biases presumably neutral molecular variation in yeast genomes. *Genome Biology and Evolution* **3**: 15-22.

Baer CF, Shaw F, Steding C, Baumgartner M, Hawkins A, Houppert A, Mason N, Reed M, Simonelic K, Woodard W et al. 2005. Comparative evolutionary genetics of spontaneous mutations affecting fitness in rhabditid nematodes. *Proceedings of the National Academy of Sciences of the United States of America* **102**(16): 5785-5790.

Carlson J, Locke AE, Flickinger M, Zawistowski M, Levy S, Myers RM, Boehnke M, Kang HM, Scott LJ, Li JZ et al. 2018. Extremely rare variants reveal patterns of germline mutation rate heterogeneity in humans. *Nat Commun* **9**(1): 3753.

Chaisson MJP, Huddleston J, Dennis MY, Sudmant PH, Malig M, Hormozdiari F, Antonacci F, Surti U, Sandstrom R, Boitano M et al. 2015. Resolving the complexity of the human genome using single-molecule sequencing. *Nature* **517**(7536): 608-U163.

Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, Land SJ, Lu XY, Ruden DM. 2012. A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w(1118); iso-2; iso-3. *Fly* **6**(2): 80-92.

Conrad DF, Keebler JEM, DePristo MA, Lindsay SJ, Zhang YJ, Casals F, Idaghdour Y, Hartl CL, Torroja C, Garimella KV et al. 2011. Variation in genome-wide mutation rates within and between human families. *Nature Genetics* **43**(7): 712-U137.

de Visser J, Hermisson J, Wagner GP, Meyers LA, Bagheri HC, Blanchard JL, Chao L, Cheverud JM, Elena SF, Fontana W et al. 2003. Perspective: Evolution and detection of genetic robustness. *Evolution* **57**(9): 1959-1972.

Denver DR, Dolan PC, Wilhelm LJ, Sung W, Lucas-Lledo JI, Howe DK, Lewis SC, Okamoto K, Thomas WK, Lynch M et al. 2009. A genome-wide view of *Caenorhabditis elegans* base-substitution mutation

processes. *Proceedings of the National Academy of Sciences of the United States of America*

106(38): 16310-16314.

Denver DR, Morris K, Kewalramani A, Harris KE, Chow A, Estes S, Lynch M, Thomas WK. 2004.

Abundance, distribution, and mutation rates of homopolymeric nucleotide runs in the genome

of *Caenorhabditis elegans*. *Journal of Molecular Evolution* **58**(5): 584-595.

Denver DR, Wilhelm LJ, Howe DK, Gafner K, Dolan PC, Baer CF. 2012. Variation in base-substitution

mutation in experimental and natural lineages of Caenorhabditis nematodes. *Genome Biology*

and Evolution **4**(4): 513-522.

Dickinson WJ. 2008. Synergistic fitness interactions and a high frequency of beneficial changes among

mutations accumulated under relaxed selection in *Saccharomyces cerevisiae*. *Genetics* **178**(3):

1571-1578.

Drake JW, Charlesworth B, Charlesworth D, Crow JF. 1998. Rates of spontaneous mutation. *Genetics*

148(4): 1667-1686.

Fisher RA. 1930. *The Genetical Theory of Natural Selection*. Clarendon Press, Oxford.

Francioli LC, Polak PP, Koren A, Menelaou A, Chun S, Renkens I, van Duijn CM, Swertz M, Wijmenga C,

van Ommen G et al. 2015. Genome-wide patterns and properties of de novo mutations in

humans. *Nature Genetics* **47**(7): 822-+.

Friedman J, Hastie T, Tibshirani R. 2010. Regularization Paths for Generalized Linear Models via

Coordinate Descent. *Journal of Statistical Software* **33**(1): 1-22.

Halligan DL, Keightley PD. 2009. Spontaneous mutation accumulation studies in evolutionary genetics.

Annual Review of Ecology Evolution and Systematics **40**: 151-172.

Houle D, Morikawa B, Lynch M. 1996. Comparing mutational variabilities. *Genetics* **143**(3): 1467-1483.

Hwang HY, Wang J. 2017. Effect of mutation mechanisms on variant composition and distribution in

Caenorhabditis elegans. *PLoS Computational Biology* **13**(1).

Jasmin JN, Lenormand T. 2016. Accelerating Mutational Load Is Not Due to Synergistic Epistasis or

Mutator Alleles in Mutation Accumulation Lines of Yeast. *Genetics* **202**(2): 751-+.

Keightley PD, Caballero A. 1997. Genomic mutation rates for lifetime reproductive output and lifespan in

Caenorhabditis elegans. *Proceedings of the National Academy of Sciences of the United States of*

America **94**(8): 3823-3827.

Kimura M. 1967. On evolutionary adjustment of spontaneous mutation rates. *Genetical Research* **9**(1):

23-27.

Landry CR, Lemos B, Rifkin SA, Dickinson WJ, Hartl DL. 2007. Genetic properties influencing the

evolvability of gene expression. *Science* **317**(5834): 118-121.

Liu HX, Jia YX, Sun XG, Tian DC, Hurst LD, Yang SH. 2017. Direct Determination of the Mutation Rate in

the Bumblebee Reveals Evidence for Weak Recombination-Associated Mutation and an

Approximate Rate Constancy in Insects. *Molecular Biology and Evolution* **34**(1): 119-130.

Long HG, Behringer MG, Williams E, Te R, Lynch M. 2016. Similar Mutation Rates but Highly Diverse

Mutation Spectra in Ascomycete and Basidiomycete Yeasts. *Genome Biology and Evolution*

8(12): 3815-3821.

Lynch M. 2007. *The Origins of Genome Architecture*. Sinauer Associates, Inc., Sunderland, MA.

-. 2008. The Cellular, Developmental and Population-Genetic Determinants of Mutation-Rate Evolution.

Genetics **180**(2): 933-943.

Lynch M, Ackerman MS, Gout JF, Long H, Sung W, Thomas WK, Foster PL. 2016. Genetic drift, selection

and the evolution of the mutation rate. *Nature Reviews Genetics* **17**(11): 704-714.

Lynch M, Hill WG. 1986. Phenotypic evolution by neutral mutation. *Evolution* **40**(5): 915-935.

Lynch M, Walsh B. 1998. *Genetics and Analysis of Quantitative Traits*. Sinauer, Sunderland, MA.

Matsuba C, Lewis S, Ostrow DG, Salomon MP, Sylvestre L, Ungvari-Martin J, Baer CF. 2012. Invariance (?)

of mutational parameters for relative fitness over 400 generations of mutation accumulation in

Caenorhabditis elegans. *G3|Genes, Genomes, Genetics* **2**: 1497-1503.

McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K, Altshuler D, Gabriel

S, Daly M et al. 2010. The Genome Analysis Toolkit: A MapReduce framework for analyzing next-

generation DNA sequencing data. *Genome Research* **20**(9): 1297-1303.

Michaelson JJ, Shi YJ, Gujral M, Zheng HC, Malhotra D, Jin X, Jian MH, Liu GM, Greer D, Bhandari A et al.

2012. Whole-genome sequencing in autism identifies hot spots for *de novo* germline mutation.

Cell **151**(7): 1431-1442.

Ness RW, Kraemer SA, Colegrave N, Keightley PD. 2016. Direct Estimate of the Spontaneous Mutation

Rate Uncovers the Effects of Drift and Recombination in the Chlamydomonas reinhardtii Plastid

Genome. *Molecular Biology and Evolution* **33**(3): 800-808.

Ness RW, Morgan AD, Vasanthakrishnan RB, Colegrave N, Keightley PD. 2015. Extensive de novo

mutation rate variation between individuals and across the genome of Chlamydomonas

reinhardtii. *Genome Research* **25**(11): 1739-1749.

Ogur M, Ogur S, Stjohn R. 1960. Temperature dependence of the spontaneous mutation rate to

respiration deficiency in Saccharomyces. *Genetics* **45**(2): 189-194.

Phillips N, Salomon M, Custer A, Ostrow D, Baer CF. 2009. Spontaneous mutational and standing genetic
(co)variation at dinucleotide microsatellites in *Caenorhabditis briggsae* and *Caenorhabditis*
elegans. *Molecular Biology and Evolution* **26**(3): 659-669.

Phillips PC. 2008. Epistasis - the essential role of gene interactions in the structure and evolution of
genetic systems. *Nature Reviews Genetics* **9**(11): 855-867.

Rifkin SA, Houle D, Kim J, White KP. 2005. A mutation accumulation assay reveals a broad capacity for
rapid evolution of gene expression. *Nature* **438**(7065): 220-223.

Rockman MV, Kruglyak L. 2009. Recombinational Landscape and Population Genomics of Caenorhabditis
elegans. *PLoS Genetics* **5**(3).

Schrider DR, Houle D, Lynch M, Hahn MW. 2013. Rates and Genomic Consequences of Spontaneous
Mutational Events in *Drosophila melanogaster*. *Genetics* **194**: 937-954.

Seyfert AL, Cristescu MEA, Frisse L, Schaack S, Thomas WK, Lynch M. 2008. The rate and spectrum of
microsatellite mutation in Caenorhabditis elegans and Daphnia pulex. *Genetics* **178**(4): 2113-
2121.

Sharp NP, Agrawal AF. 2012. Evidence for elevated mutation rates in low-quality genotypes. *Proceedings*
of the National Academy of Sciences of the United States of America **109**(16): 6142-6146.

-. 2016. Low genetic quality alters key dimensions of the mutational spectrum. *PLoS Biology* **14**(3).

Shaw FH, Baer CF. 2011. Evolutionary consequences of fitness-dependent mutation rate in finite
populations. *Journal of Evolutionary Biology* **24**: 1677-1684.

Sturtevant AH. 1937. Essays on evolution. 1. On the effects of selection on mutation rate. *Quarterly*
Review of Biology **12**: 467-477.

Teotónio H, Estes S, Phillips PC, Baer CF. 2017. Experimental Evolution with Caenorhabditis Nematodes.
Genetics **206**(2): 691-716.

Thompson O, Edgley M, Strasbourger P, Flibotte S, Ewing B, Adair R, Au V, Chaudhry I, Fernando L,
Hutter H et al. 2013. The million mutation project: A new approach to genetics in Caenorhabditis
elegans. *Genome Research* **23**(10): 1749-1762.

Tyson JR, O'Neil NJ, Jain M, Olsen HE, Hieter P, Snutch TP. 2018. MinION-based long-read sequencing
and assembly extends the Caenorhabditis elegans reference genome. *Genome Research* **28**(2):
266-274.

Watterson GA. 1975. Number of segregating sites in genetic models without recombination. *Theoretical*
Population Biology **7**(2): 256-276.

Xie C, Tammi MT. 2009. CNV-seq, a new method to detect copy number variation using high-throughput

sequencing. *Bmc Bioinformatics* **10**.

Yang SH, Wang L, Huang J, Zhang XH, Yuan Y, Chen JQ, Hurst LD, Tian DC. 2015. Parent-progeny

sequencing indicates higher mutation rates in heterozygotes. *Nature* **523**(7561): 463-U187.

Yeh S-D, Saxena AS, Crombie TA, Feistel D, Johnson LM, Lam I, Lam J, Saber S, Baer CF. 2018. The

mutational decay of male-male and hermaphrodite-hermaphrodite competitive fitness in the

androdioecious nematode *C. elegans*. *Heredity* **120**(1): 1-12.

845