Coordinated cerebellar climbing fiber activity signals learned sensorimotor predictions

William Heffley II[1], Eun Young Song[1], Ziye Xu[1], Benjamin Taylor[1], Mary Anne Hughes[1], Andrew McKinney[1,2,3], Mati Joshua[1,4] and Court Hull[1]

[1]Department of Neurobiology, Duke University Medical School, Durham, United States
[2]Neuroscience Department, Baylor College of Medicine, Houston, United States
[3]Duncan Neurological Research Institute at Texas Children's Hospital, Houston, United States
[4]Department of Medical Neurobiology, The Hebrew University, Hadassah Medical School, Jerusalem, Israel

Correspondence:

Dr. Court Hull

Duke University Medical School

Dept. of Neurobiology

311 Research Drive

Durham, NC 27710

Email: Hull@neuro.duke.edu

Phone: 919-613-0927

Acknowledgements:

This work was supported by grants from the NIH NINDS (5R01NS096289-02), the Sloan Foundation, and the Whitehall Foundation. We thank Dr. Lindsey Glickfeld for helpful discussions and input on calcium imaging approaches and analyses, Drs. Stephen Lisberger, Greg Field, Kevin Franks and Fan Wang for feedback on early versions of this manuscript, and members of the Hull and Glickfeld labs for input and technical assistance throughout the project.

Abstract

The prevailing model of cerebellar learning states that climbing fibers (CFs) are both driven by, and serve to correct, erroneous motor output. However, this model is grounded largely in studies of behaviors that utilize hardwired neural pathways to link sensory input to motor output. To test whether this model applies to more flexible learning regimes that require arbitrary sensorimotor associations, we have developed a cerebellar-dependent motor learning paradigm compatible with both mesoscale and single dendrite resolution calcium imaging in mice. Here, we find that CFs are preferentially driven by, and more time-locked to correctly executed movements, exhibiting widespread correlated activity within parasagittal processing zones that is governed by predictions about the outcome of movement. Together, these CF activity patterns are well-suited to drive learning based on successful movement by providing predictive instructional input that does not rely exclusively on motor errors.

Introduction

A key role of the cerebellum is to form predictive associations between sensory inputs and motor outputs. These sensorimotor predictions are critical for generating well-timed and accurate movements, and in the absence of cerebellar function, the lack of such predictive motor output severely impairs our ability to generate coordinated responses to stimuli in the external world.

Classic models posit that the cerebellum generates sensorimotor predictions according to a supervised learning rule[1-3]. According to such models, projections from the inferior olive called climbing fibers are thought to signal motor errors, thus providing information to Purkinje cells about discrepancies between the expected consequences of a motor command and subsequent sensory feedback. To correct erroneous motor output, the climbing fibers instruct heterosynaptic long-term depression[4-6] by producing powerful regenerative calcium transients[7-9] in Purkinje cell dendrites called complex spikes[10, 11]. In so doing, climbing fibers are thought to appropriately update the cerebellar forward internal model with revised sensorimotor predictions.

This error-signaling framework provides a compelling explanation for climbing fiber activity in a variety of simple behaviors, such as classical conditioning (e.g. eyeblink conditioning) and adaptation (e.g. vestibulo-ocular reflex gain changes)[12-14] paradigms. Such behaviors typically rely on a yoked relationship between unconditioned sensory input and motor output, allowing the cerebellum to utilize signals from hardwired pathways to drive learning. Hence, the climbing fibers can instruct learning by responding to an unconditioned stimulus (e.g. periocular eye puff or retinal slip) that produces the same movement requiring modification (e.g. eyelid closure or eye movement).

However, many forms of motor learning do not involve modifications to motor programs linked directly to an unconditioned stimulus and response. Instead, the correct association between sensory input and motor output must be learned through experience, and the sensory information necessary for learning may have no direct relationship to the movement that requires modification. Such abstract associations necessitate that learning cannot be achieved by input from hardwired sensorimotor pathways alone. Under such conditions, where an unconditioned stimulus and response alone does not contain sufficient information to guide learning, it is unclear whether the current model of supervised error signaling can account for either climbing fiber activity or the cerebellar contribution to learning.

To test how the climbing fiber system is engaged under conditions where the sensory and motor signals necessary to drive learning are not innate, we have established a cerebellar-dependent behavioral paradigm compatible with population level *in vivo* calcium imaging, optogenetic and electrophysiological approaches. Using this paradigm, we reveal two key features of cerebellar climbing fiber (CF) driven complex spiking. First, we find that complex spiking cannot be accounted for by a simple error-based supervised learning model. Instead, complex spiking can signal learned, task specific predictions about the likely outcome of movement in a manner well suited for flexible control of complex motor behaviors. Second, population level recordings reveal that while complex spiking is correlated within and across parasagittal zones, these correlations also depend on behavioral context. While previous measurements have shown increased correlations in complex spiking in response to sensory input or motor output[15-18], and have suggested an important role for synchrony in downstream processing and motor learning[19], our results reveal that such modulation can vary for identical movements depending on behavioral relevance. Hence, these data reveal key features of cerebellar CF activity that differ significantly from classically studied cerebellar behaviors, and suggest an extension to current models of cerebellar learning in order to account for the role of complex spiking in tasks that require abstract sensorimotor associations.

Results:

To determine the conditions under which climbing fiber driven complex spiking occurs during a motor learning regime that requires an arbitrary sensorimotor association, we first designed an appropriate task. An important feature of the task design is to specifically engage neurons near the dorsal surface of the cerebellum, thereby allowing for visualization of complex spikes at the population level via calcium imaging. Hence, we used optogenetics to functionally map the midlateral dorsal surface of the mouse cerebellum according to the motor output produced by spatially defined populations of Purkinje cells (PCs). To do so, we expressed Archaerhodopsin (Arch) using a transgenic approach (methods) in cerebellar PCs (Fig. 1A), and

used an external fiber coupled laser to transiently suppress PCs near the dorsal surface of the

cerebellum (Fig. 1B). Consistent with previous work[20], we identified a population of superficial

PCs in lobule simplex capable of driving ipsilateral forelimb movements either during optogenetic

silencing with Arch or following a brief train of electrical simulation (Fig. 1, Supp. Fig. 1, Supp.

Movie 1). *In vivo* single unit recording confirmed that our optogenetic stimulation was effective in

silencing superficial PCs at the threshold for driving forelimb movement (~20 mW) up to a depth

of 500 μm (Fig. 1B).

A cerebellar dependent sensorimotor association learning task

Based on these results, we designed a sensorimotor association task involving forelimb

movement, with the rationale that the cerebellar contribution to learning in such a task should

necessarily involve the PCs capable of driving such movements. This task contains two discrete

phases; one where motor output is reactive ('cue reaction'), and another where motor output

becomes predictive with learning ('cue prediction'). In both task variations, head restrained, water

deprived mice are trained to self-initiate trials by depressing a lever, and to release the lever in

response to a visual cue (Fig. 1C,D).

In the cue reaction task, trial-to-trial variability in the timing of the visual cue (Fig. 1E,G-

"random cue delay") is imposed to necessitate that mice employ reactive forelimb responses. In

this regime, lever release reaction times are reflective of the latency for sensory integration[21] and

remain constant throughout each session (ΔRT = 1.1 $\pm$ 9.3 ms, n=26 sessions, 7 mice, Supp.

Movie 2).

In the cue prediction task, the visual cue is presented with the same delay on every trial.

In this regime, mice learn to predict the timing of the cue and adjust their motor responses to more

closely approximate the time of cue (Fig. 1F,G, ΔRT 500ms cue delay = 154.5 $\pm$ 13.6 ms, n=44

sessions, 10 mice). Moreover, learning occurs within a single training session (90 min., 309-700

trials), is retained across days (Supp. Fig. 2A), and can be extinguished by returning mice to the

cue reaction paradigm (Supp. Fig. 2C,D). In addition, we find that learning depends on the

duration of the cue delay: maximal learning occurs at short cue delays (e.g. 500 ms) and little or

no learning occurs for cue delays greater than 1.5 seconds (Fig. 1H; one-way ANOVA, main effect

of cue delay, p = 4.8×10^{-6}. Post hoc comparison to cue reaction: 500 ms, p=1.9×10^{-11}, 1 s,

p=8.4×10^{-11}, 1.5 s, p=0.359, 2 s, p=0.005). This dependence on a short interval delay is consistent

with the temporal characteristics of canonical cerebellar-dependent sensorimotor associative

learning[22].

To test whether animals adjust the timing or kinematics of their forelimb movement to

predict the timing of the visual cue, we measured lever kinematics during a subset of learning

sessions. Alignment of the lever trajectory to the onset of the cue revealed that animals altered

the time of lever release, releasing the lever sooner in anticipation of the visual cue (Fig. 1I; 20%

rise time occurs 116 ± 18 ms earlier, p = 1.2×10^{-5}, paired t-test). However, despite substantial

variability in the kinematics of release (Supp. Fig. 3), independent alignment of both press and

release demonstrated that there was no change in movement kinematics across learning

sessions (Fig 1I-J; First 1/3 of session 20-80% release rise time= 18.9 ± 0.6 ms, last 1/3 of session

= 19.3 ± 0.8 ms; p=0.513, paired t-test), indicating that mice release the lever sooner but with the

same kinematics after learning.

To test the necessity of the lobule simplex for this learning, we first locally applied lidocaine

in order to block spiking in cerebellar cortical neurons, including PCs. However, consistent with

the critical tonic inhibition PCs provide to the deep cerebellar nuclei, we observed significant motor

deficits including slowed movement, dramatically fewer initiated trials, and dystonic limb

contractions (not shown). Thus, this manipulation was not appropriate to define the necessity of

lobule simplex in our task.

To avoid significant impairment of motor output, we next used a pharmacological

manipulation to selectively disrupt excitatory synaptic transmission in lobule simplex without

abolishing the spontaneous pacemaking of PCs. For these experiments, we optimized

concentrations of the excitatory synaptic transmission blockers NBQX (300 μM), CPP (30 μM)

and MCPG (30 μM) to avoid visible impairment of motor output, and then tested this cocktail

during behavior. Following local drug application to lobule simplex (methods), mice retained the

ability to release the lever in response to the visual cue, maintaining stable kinematics across

sessions (Fig. 2A,B) that were not different from control sessions (Fig. 1I,J, and Supp. Fig. 3;

Press: control range = 28.97 ± 2.5 ms, NBQX range = 25.33 ± 1.9 ms, p=0.2612; Release: control

range = 6.16 ± 1.2 ms, NBQX range = 4.13 ± 1.06 ms, p=0.2215, unpaired t-test). These data

suggest that motor output was not significantly impaired as a result of this pharmacological

manipulation. However, despite stable movement kinematics, learning was severely impaired

when the cue was presented with a constant 500 ms delay (Fig 2A,C,D) (NBQX ΔRT with 500 ms

cue delay = 48.8 ± 26.7, control ΔRT with 500 ms cue delay = 154.5 ± 13.6 ms, ΔRT with cue

reaction = 1.1 ± 9.3, unpaired t-test NBQX vs control 500 ms p=4.6×10^{-4}, unpaired t-test NBQX

vs control cue reaction p=0.05). Acute single unit PC recordings during drug application revealed

that this manipulation significantly reduced complex spike rates, consistent with a reduction in the

amplitude of postsynaptic glutamatergic currents produced by climbing fiber input (control $0.60 \pm$

0.07 Hz, NBQX/CPP/MCPG 0.27 ± 0.04 Hz, n = 9, p=8.5×10^{-4}, paired t-test, Supp. Fig. 4). We

also found a modest reduction in simple spiking, consistent with reducing excitatory synaptic input

from the mossy fibers and parallel fibers (control 66.8 ± 17.8 Hz, NBQX/CPP/MCPG 45.9 ± 13.0

Hz, n = 8, n = 9, p=0.36, paired t-test). Hence, these data support that hypothesis that synaptic

transmission in lobule simplex is necessary for predictive sensorimotor learning to occur in the
cue reaction phase of our task.

Complex spiking is not driven by motor errors in a voluntary motor learning task

To test how the olivary system is engaged in this behavior, we imaged climbing fiber
evoked calcium signals in mice performing the cue reaction phase of our task. In this regime, all
sensorimotor signals are in place to drive learned, predictively timed forelimb movements, but no
learning can occur due to the variable cue timing (Fig. 1G,H). Hence, we can test what task
features drive complex spiking in a stable regime. We began by using a single photon microscope
to image populations of PCs virally expressing GCaMP6f (methods). Image segmentation in a
subset of experiments with the most superficial GCaMP expression revealed structures consistent
in size and shape with PC dendrites in cross section (Supp. Fig. 5). Calcium transients in these
structures occurred at 0.5 ± 0.01 Hz, consistent with the baseline rate of complex spiking we
measured with single unit electrophysiology (Supp. Fig. 5). These results are also consistent with
several *in vivo* calcium imaging studies using a variety of GCaMP variants showing that PC
responses are dominated by the large dendritic calcium transients produced by climbing fiber
driven complex spikes[15-18, 23-28].

We next identified regions of interest, defined by the expression pattern of GCaMP in each
animal, that were restricted to the region within lobule simplex capable of driving ipsilateral
forelimb movements (methods). Consistent with the role of this region in the control of forelimb
movements, we observed task modulated PC calcium transients associated with the timing of
lever press and release, with the largest modulation at the time of lever release (Fig. 3A-C).
Surprisingly, when we segregated trials according to outcome, we observed significantly larger
calcium transients associated with lever releases when animals correctly released the lever in
response to the visual cue as compared to trials where lever release occurred prior to the cue
(Correct = 0.156 ± 0.023 ΔF/F; Early = 0.092 ± 0.018 ΔF/F, n=17 sessions, 10 animals, p=4.60x10^{-5}
^{5}, paired t-test). These enhanced responses were distributed widely across dorsal, superficial
lobule simplex, and were consistent across sessions and animals (Fig. 3D). The same pattern of
enhanced complex spiking on correctly timed lever releases was observed in the cue prediction
task as measured by single unit PC electrophysiological recordings (Supp. Fig. 6). Thus, in this
motor task, our data suggest that complex spikes occur preferentially on correctly timed
movements, and not motor errors.

There are many differences between the correct and early release trials that could
contribute to the observed outcome-dependent difference in complex spiking. One possible
difference between trial types could be the kinematics of the forelimb movement. Hence, in a
subset of our experiments, we measured movement kinematics by comparing lever trajectories

on correct and early release trials (Fig. 4A-B). These data revealed closely matched movement

kinematics between correct and early lever release trials (Press: correct = 26.8 ± 1.6 ms, early =

25.1 ± 1.3 ms; Release: correct = 18.9 ± 0.8 ms, early: 22.1 ± 1.5 ms, n=14 sessions).

Another difference between correctly timed and early lever releases is that reward is only

delivered on correct trials, and this results in extended licking. Indeed, we find that lick rates were

significantly higher and prolonged on correct trials (Fig. 4C; correct = 10.5 ± 0.7 Hz, early = $1.5 \pm$

0.4 Hz, $p=3.74 \times 10^{-9}$, paired t-test, n=9 animals, 15 sessions). Moreover, in some recording

sessions we observed a late, sustained calcium transient that occurred during licking (Fig. 4D).

However, we found no correlation between the mean early, peak calcium transients and mean

lick rates across trial types (Fig. 4E,F; Correct $R^2=0.182$, p=0.11; Early $R^2=5.97 \times 10^{-5}$, p=0.98,

ANOVA) or within sessions across trials of each type (Fig. 4G; Correct $\rho = -0.002 \pm 0.028$, p=0.93;

Early $\rho = 0.054 \pm 0.113$, p=0.65, one sample t-test; Supp. Fig. 7D; Correct $\rho = 0.0053 \pm 0.0019$,

p=0.544; Early $\rho = 0.0912 \pm 0.0048$, p=0.428, n=447 dendrites, one sample t-test). Thus, we

conclude that the enhanced responses as measured by the early peak calcium transient on

correctly timed lever releases are not due to differences in forelimb movement kinematics or

licking, but are instead linked to the context of the movement.

Behavioral context determines the probability and synchrony of complex spiking

To investigate the dynamics of complex spiking at the single cell level, we used video-rate

two-photon microscopy to image the dendrites of PCs during behavior. Using this approach, we

isolated the dendrites of individual PCs[24] (Fig. 5, Supp. Figs. 5, 8), and measured both

spontaneous and task-evoked calcium transients. As with the single photon imaging experiments,

spontaneous calcium transients measured during inter-trial intervals occurred at a rate of $0.61 \pm$

0.01 Hz, consistent with our electrophysiological measurements of spontaneous complex spiking

(Supp. Fig. 5E,F). The two-photon data also replicated the main finding that the mean calcium

transients were significantly larger on correct release trials within individual PC dendrites (Correct

$= 0.101 \pm 0.004$ $\Delta F/F$; Early = 0.035 ± 0.002 $\Delta F/F$, n=1146 dendrites, $p=9.79^{-76}$, paired t-test) (Fig.

5A,B). Moreover, lever releases that were rewarded but occurred too fast to be responses to the

visual cue (reaction times less than 200 ms) produced calcium transients that were significantly

smaller than those produced by correct lever release trials ('Too fast' 0.029 ± 0.002 $\Delta F/F$, $p=3.91^{-}$

^{104}, paired t-test) and not significantly different from responses on early release trials (p=0.086).

Hence, these data support the analysis in Figure 4 that the enhanced complex spiking on correct

trials is not due to licking or reward per se.

In addition to larger mean responses across trials, population level analysis revealed that

more PCs were driven on correct release trials (Fig 5C). Of the total population of PCs that

responded to any lever press or release (1146 of 1302), most (88%) PCs exhibited complex

spikes on correct release trials, whereas less than half (41%) responded during early lever releases (Fig. 5D, *top*). Approximately half (48%) of PCs exhibited complex spikes in response to lever press (Fig. 5D, *bottom*). The increased number of active dendrites on correct trials is unlikely to be a thresholding artifact, since there was no significant difference in the average response amplitude of correct-only and correct+early responsive dendrites to correct trials (Correct only 0.112 ± 0.006 ΔF/F, n=578; Correct + Early 0.119 ± 0.005 ΔF/F, n=457 dendrites, p=0.44 unpaired t-test). These data also indicate that the measured increase in number of active dendrites is not due to contamination of fluorescent signals from neighboring dendrites. Hence, these single dendrite measurements explain our single photon results by demonstrating that both more PCs exhibit complex spikes on correct trials, and those dendrites that exhibit complex spikes in response to both trial types have larger mean calcium transients on correct trials.

The difference in the average release-evoked response on correct versus early trials may be due to a difference in the amplitude of the calcium transients on each trial. For instance, complex spikes are not "all-or-none" events[29], and the number of spikelets can vary from event to event[30-32], altering the size of the evoked calcium transient[33]. Thus, we extracted single trial events associated with lever releases across trial types, and compared these with each other and to the spontaneous calcium transients. However, this analysis revealed no significant difference between the amplitude of calcium transients evoked on single trials in response to either correct or incorrect lever releases (Correct 0.266 ± 0.004 ΔF/F n=19,550 events, Early 0.263 ± 0.004 ΔF/F n=9,167 events), and between movement-evoked and spontaneous activity (Spontaneous 0.272 ± 0.004 ΔF/F n=286,964 events) (p=0.17, one-way ANOVA) (Fig. 5E). Thus, if there are variable numbers of evoked spikelets across trials, such differences are not systematically correlated with either trial outcome or task engagement. Moreover, the equivalent amplitude of lever-evoked and spontaneous events suggests that the majority of the measured signal is from calcium influx due to complex spiking, and is not substantially influenced by additional input from the parallel fibers. Indeed, the size and shape of our segmented ROIs is consistent with several studies showing that complex spikes produce a calcium signal throughout the entire PC dendrite[7, 9], whereas parallel fiber calcium signals are much smaller and confined to isolated regions of spiny dendritic branchlets[34-37].

To assess how equivalent single event responses can produce larger mean responses on correct trials, we generated a peristimulus time histogram of complex spike events for each trial type aligned to the time of release (Fig. 6A). This analysis revealed a significantly enhanced complex spike rate at the time of lever release for trials when movement was correctly timed (Fig. 6A,B; correct spike rate=2.04 ± 0.05 Hz, early spike rate= 1.33 ± 0.04 Hz, p=8.14x10^{-15}, n=1146 dendrites, paired t-test). In addition, complex spiking on correct trials in single PC dendrites occurred with shorter latency (Fig 6C; correct latency=35.86 ± 1.19 ms, early latency=46.49 ±

1.25 ms, p=2.65x10^{-11}, n=1146 dendrites, paired t-test) and less temporal jitter across trials (Fig

6D; correct jitter (S.D of latency) = 53.79 ± 1.39 ms S.D., early jitter 95.16 ± 1.88 ms S.D.,

p=6.16x10^{-45}, n=1146 dendrites, unpaired t-test). Notably, these measures of jitter are likely to

be an overestimate since our measure of event time is limited by our sampling interval (33 ms),

and the true onset might have occurred between samples. However, when accounting for the

overall spontaneous event rate of ~1 Hz, the 50 ms jitter of events on correct trials is comparatively

precise.

These results also suggest the possibility of enhanced synchrony at the population level

on correct release trials.   Indeed, in some sessions, we measured enhanced population

synchrony across dendrites (Fig 6E; 13 out of 30 sessions have significantly elevated synchrony

on correct trials, open circles, p<0.05 for each session, mean across the population is not

significant, p=0.098, paired t-test.).   However, in many experiments, we observed no such

increase in population synchrony on correct trials.   We thus considered the possibility that

enhanced synchrony was location specific, and might obey spatial structure that was not well

demarcated at the scale of the field of view in our two-photon experiments.

Context determines correlated population activity at the mesoscale level

To investigate whether correlated complex spiking was spatially organized, we returned

to single photon imaging and analyzed experiments that exhibited widespread GCaMP

expression across superficial lobule simplex (n=5 animals, 7 sessions). Using an unsupervised,

iterative pixel-clustering approach[26] (methods), we first assessed correlations between pixels

across all trials (Fig. 7A). Despite non-uniform, unpatterned GCaMP expression, this analysis

revealed spatial patterns of correlated activity that were organized across parasagittal bands

oriented in the rostro-caudal axis. These bands were 221 ± 15 μm wide on average across 39

measured zones, consistent with previous anatomical and physiological measurements of

cerebellar microzones[38, 39].   Hence, these data support the longstanding hypothesis that the

anatomical pattern of climbing fiber projections into the cerebellar cortex establishes functionally

distinct parasagittal processing modules.

To test how complex spiking is modulated during behavior within and across parasagittal

zones, we divided trials according to outcome and analyzed brief epochs surrounding the time of

lever release (Fig. 7B). While this analysis lacks the fine temporal resolution of the two-photon

event based analysis in Figure 6, it nonetheless revealed that activity amongst pixels within a

zone exhibited higher correlations for most zones on correct lever release trials as compared to

early release trials ($r_{correct}$ = 0.77 ± 0.01, r_{early} = 0.69 ± 0.017, p = 1.60x10^{-5}, n=39 clusters, paired

t-test) (Fig. 7C).   Because higher spike rates can cause larger correlations, we tested whether

the higher complex spike rates on correct trials could account for the higher correlations. While

increased complex spiking did produce larger correlations, early release trials still had weaker

correlations than correct release trials across equivalent activity levels as assessed by linear fits

to bootstrapped distributions paired from each trial type (methods) (Fig. 7D). Hence, the

enhanced correlations on correct trials are not an artifact of enhanced complex spike rates on

correct trials. We also found enhanced correlations between neighboring and non-neighboring

zones on correct lever release trials (Fig. 7E,F) (Correct $r_{neighboring}$ = 0.69 ± 0.03, n=31 cluster

pairs, p=0.004 vs early, Correct $r_{non\text{-}neighboring}$ = 0.65 ± 0.03, n=50 cluster pairs, p=6.2x10^{-12} vs

early, paired t-test). However, these cross-zone correlations were significantly lower on average

than those within zones (within zone vs neighboring, p=0.004; within zone vs. non-neighboring

p=3.4x10^{-4}, unpaired t-test). Hence, these data reveal a precise spatial organization of climbing

fiber activity, with highly correlated complex spiking within parasagittal zones that is task specific,

differing for movements with the same kinematics depending on behavioral context.

Complex spiking signals learned sensorimotor predictions

Our data suggests that complex spiking does not signal motor errors in this behavioral

paradigm. Instead, our results are consistent with the possibility that the climbing fibers signal

either 1) correctly timed motor output, 2) a temporal-difference (TD) prediction error driven by the

unexpected visual cue that instructs lever release and predicts upcoming reward delivery, or 3)

an expectancy signal related to the probable outcome of movement. To distinguish between these

possibilities, we took advantage of the temporal structure of the task, wherein the probability of

cue appearance, and thus reward delivery, increases with lever hold time (i.e. the hazard function

is not flat). Thus, we tested whether complex spiking depends on lever hold time for correct and

early lever releases.

We find that while complex spiking is independent of lever hold time for correctly timed

movements (1P: ρ = -0.07, p=0.76, n=17 sessions ; 2P: correct ρ = -0.01, p=0.98 n=30 sessions;

Fig. 8 A,B), there is a strong positive relationship between lever hold time and complex spiking

for early lever releases across both single and multiphoton imaging sessions (1P: 0.91 p<1.0x10^{-16},

n=17 sessions; 2P: early ρ = 0.90, p=4.58x10^{-6}, n=30 sessions ; Fig. 8 A,B). The elevated

complex spiking associated with longer lever hold times is due to both an increase in spiking at

the time of lever release (Window 1: short duration early ΔF/F = 0.03 ± 9.55x10^{-4}, long duration

early ΔF/F = 0.04 ± 0.00, p=3.84x10^{-10}, n=1146 dendrites, paired t-test; Figure 8C), and an

additional late response not present on short duration early release trials (Window 2: short

duration early ΔF/F = 0.04 ± 0.00, long duration early ΔF/F = 0.07 ± 0.00, p=1.33x10^{-25}; Figure

8C). The late response occurred on single trials and within individual dendrites, indicating that it

was not generated by a diversity of response timings across trials or by a separate population of

PC dendrites (Fig. 8 D). In addition, the late response did not correlate with an increase in licking

on long duration trials, and occurred approximately 200 ms after lever release when lick rates began to decrease and reward delivery was no longer possible (Fig 8 H,I). These results are consistent with an expectancy model, where complex spiking signals both an initial prediction about movement outcome and a readout of whether the initial prediction was violated. In this model, an initial prediction accompanies lever release that depends primarily on cue presentation (Window 1: long duration correct $\Delta F/F = 0.07 \pm 0.00$, long duration early $\Delta F/F = 0.04 \pm 0.00$, p = 3.16×10^{-7}, n=1146 dendrites, paired t-test). While the visual cue provides the strongest prediction of trial outcome, trial duration also contributes to expectation according to the task structure. In the case that initial expectation is unmet, the climbing fibers also signal this unexpected outcome through an increase in spiking.

To further test this model, we performed a second set of experiments to probe the role of violated expectations in driving complex spiking. In these experiments, we omitted reward delivery on 20% of correctly timed lever releases (Fig. 8 J-N). Consistent with the cue appearance establishing expectation at the time of lever release, both rewarded and omission trials resulted in calcium transients that were nearly equivalent at the time of lever release (Window 1: rewarded = 0.07 ± 0.01, omission = 0.07 ± 0.01 ; p=0.04, n=81 dendrites, paired t-test; Figure 8L). However, omission trials resulted in an additional, late response at the time when lick rates began to decrease and reward delivery was no longer possible (Window 2: rewarded = 0.06 ± 0.01, omission = 0.08 ± 0.01; p=2.54×10^{-9}, n=81 dendrites, paired t-test; Figure 8M). The timing of this late response was similar to the late response following long duration early releases, and was present on single trials and within individual dendrites (Figure 8J,N). Hence, these experiments support the hypothesis that climbing fibers can both signal and evaluate predictions about the likely outcome of movements.

Discussion

We have established a sensorimotor association task that involves PCs near the dorsal surface of lobule simplex. This behavior has several hallmarks of cerebellar learning, including a dependence on short delay intervals for generating the learned sensorimotor association and the requirement for excitatory synaptic transmission in the cerebellar cortex. In this behavior, however, climbing fibers do not signal erroneous motor output as described by classical models of supervised learning. Specifically, we demonstrate that climbing fiber driven complex spiking occurs with higher probability, shorter latency, and less jitter when movement is correctly timed.

Our data support the hypothesis that these enhanced climbing fiber responses are related to the predicted outcome of movement, which in this behavior constitutes delivery of a water reward. Evidence supporting this model is threefold: First, the highest probability of complex spiking occurs on correctly timed lever releases when the visual cue instructs behavior and thus

a high reward expectation. Second, complex spiking is modulated by the temporal features of the

task that dictate expectation. Specifically, complex spiking in response to lever release increases

with increasing hold duration, matching the increased likelihood of reward as trial length

increases. Third, complex spiking is also driven when the expectation of reward is violated, both

when the reward is omitted on correctly executed movements and on longer hold duration early

releases when a correct outcome of lever release is more probable. Together, these experiments

suggest that climbing fibers carry instructional signals that both predict and evaluate the expected

outcome of movement. Such responses are consistent with the known role of cerebellar circuits

in generating predictive motor output, and we suggest that they could thus provide a substrate for

generating and testing the type forward models that have long been hypothesized to govern

cerebellar processing.

The climbing fiber activity observed in response to violated expectations has some

similarities to the motor error signals seen in classic cerebellar behaviors. However, in our

behavioral paradigm, we emphasize that these climbing fiber responses do not reflect motor

errors: the movements on trials in which rewards are omitted are identical to those in which

rewards are earned. Thus, these results indicate that the climbing fibers can respond differentially

to the very same movement according to its expected outcome. These context-specific climbing

fiber responses may be due to aspects of our task design. The behavior described here differs

from most cerebellar dependent learning regimes in that the movement requiring modification

(lever release) is not directly related to an unconditioned stimulus (reward) or response (licking).

As a consequence, the cerebellum cannot harness sensorimotor input from hardwired pathways

to enable learned changes in motor output. Instead, the necessary forward model must define

and evaluate the relationship between a neutral visual stimulus, a forelimb movement, and

reward.

It remains unclear whether the climbing fiber activity that enables such a forward model is

generated at the level of the olive, or if it is inherited from upstream brain regions such as the

neocortex, colliculus, or elsewhere. However, evidence suggests that the olive may have access

to different information depending on task requirements. Anatomical and physiological work in the

rodent has demonstrated that the pathway from the forelimb region of motor cortex to the IO is

independent of the pathway providing ascending sensorimotor input to the olive from the

periphery[40]. Such data argue that the olive has access to unique information in tasks that involve

a descending motor command, and further suggest that the olive may have access to diverse

cortical computations. Indeed, the presence of abstract task timing information in the activity of

the climbing fibers suggests that the olive has access to higher-order signals. In support of this

view, evidence from a different forelimb movement task in non-human primates that also requires

an abstract sensorimotor association has shown similar climbing fiber responses[41]. Specifically,

this study demonstrated that complex spikes can signal both the predicted destination of reaching
as well as deviations from the expected destination.

Importantly, the climbing fiber activity patterns described here are appropriate to drive
motor learning under conditions that require flexible sensorimotor associations. Specifically, the
higher probability complex spiking and enhanced correlations at the population level are both
context dependent and spatio-temporally organized. At the population level, our single photon
imaging of cerebellar complex spiking has revealed correlated activity within parasagittal bands
of approximately 200 μm, likely corresponding to "microzones"[16, 26, 39, 42-46]. Because PCs across
microzones can respond to different sensorimotor input, and have different spiking[47] and
synaptic[48] properties, these zones are thought to constitute discrete computational units in the
cerebellum. Recent measurements have revealed that complex spiking in microzones can
become more synchronous during both motor output[18] and sensory input[15, 16, 27]. Our population
imaging results provide a fundamental extension to this idea by demonstrating that correlated
complex spiking within parasagittal zones is not simply enhanced by sensorimotor input, but
rather can be enhanced for the same action according to behavioral context. Precisely timed
spiking across a large population of nearby PCs that receive common parallel fiber input and
converge to the same DCN neurons would provide an ideal substrate to maximize the impact of
plasticity. However, other circuits could also play a role in learning. In particular, complex spiking
can robustly inhibit nuclear neurons[49, 50], and enhanced climbing fiber synchrony would magnify
this effect via convergence of simultaneously active PC axons in the DCN[51]. Because
synchronous inhibition has been shown to play a key role in synaptic plasticity at nuclear neuron
synapses[52, 53], correlated complex spiking across populations of nearby PCs could place the DCN
as the central site of learning under such conditions[54]. In either case, we note that the climbing
fiber responses following correct lever releases are well-timed to promote learning in our cue
prediction task by instructing movements that more closely approximate earliest time of reward
delivery.

Finally, it is notable that ours are not the first results to suggest alternate learning rules
instructed by climbing fiber activity. Recently, conditioned eyeblink learning data has pointed
toward climbing fibers implementing temporal difference (TD) reinforcement learning rule by
signaling sensory prediction errors in order to instruct motor learning[55]. Because we do not
observe evidence of negative prediction errors on reward omission trials or long duration early
release trials, our results are incompatible with this model as strictly interpreted. Instead, our data
show some similarities with the type of unsigned prediction signals observed in serotonergic[56]
and other neurons[57, 58] that have been also thought to enable associations between unexpected
outcomes and novel cues[59]. Evidence for both models can often be found in neural activity within
the same brain region, and it has been suggested that TD and unsigned prediction mechanisms

may in fact be linked, and act together to promote learning[60]. Hence, as has been done in other

brain regions, it will be necessary to systematically vary task parameters such as the valence of

instructional stimuli and the requirements for learned associations in order to resolve how the

cerebellum implements discrete learning rules according to task demands. For the present, the

results described here provide an important new demonstration that complex spiking can signal

learned, task specific information necessary for flexible control of complex motor behaviors in a

manner that does not depend exclusively on motor errors.

Author Contributions

WH, ES, AM, MJ, and CH designed experiments, WH, ES, AM, BT, MAH, and MJ conducted

experiments, WH, ES, ZX, BT, AM, MJ, and CH analyzed data, and CH wrote the manuscript.

Figure Legends

Figure 1. Cerebellar sensorimotor task for head-fixed mice. Ai) Confocal image of a sagittal

section from PCP2-Cre x Arch mouse showing expression of the inhibitory opsin Arch in PCs at

low (left) and high (right) resolution. ML; molecular layer, PCL; Purkinje cell layer, GCL; granule

cell layer. Aii) Schematics of a cranial window implanted over lobule simplex (left) and

configuration for optogenetic stimulation (right). Bi) *Left*, representative single unit recording of

PC simple spike raster (top) and spike rate histogram (bottom) aligned to onset of Arch activation.

Right, Location of the stimulation sites that evoked ipsilateral forelimb movements (green). Bii)

Average normalized firing rates from superficial (top) or deep (bottom) PCs at low (left) or high

(right) stimulation powers (n=5 mice; 10-20 mW – superficial: n=6 cells; deep: n=5 cells; 30-50

mW – superficial: n=10 cells; deep: n=4 cells). Note that the lower powers used to map motor

responses were only sufficient to strongly suppress neurons above 500 μm. C) Schematic of

configuration of behavioral task. Head-fixed mice were trained to release a lever in response to a

visual cue, with reward delivered immediately upon correctly timed movement. D) Schematic of

trial structure. The delay between lever press and release cue was either randomized (cue

reaction) or constant (cue prediction) from trial to trial. E) Average reaction time (from cue onset)

as a function of trial number for an example cue reaction session. Each point is the average of 5

trials. F) Same as E, for a cue prediction session. G) Summary of reaction times for all cue

prediction sessions with a 0.5 s cue delay (gray; n=44 sessions, 10 mice) and all cue reaction

sessions (black; n=26 sessions, 7 mice). Error bars are ±SEM across sessions. H) Average change in reaction time from the beginning to the end of cue prediction and reaction sessions (methods). I) *Top*, average lever kinematic traces from an example 0.5 s cue prediction session aligned to the cue show that lever releases occur sooner in the last 1/3 of trials (black) as compared to the first 1/3 of trials (gray). Shaded area is ±SEM across trials. *Bottom,* same example session as above aligned separately to press and release showing no difference in kinematics from beginning (gray) to end (black) of the session. J) Summary of average lever release times (20-80% rise time) across sessions (n=15 sessions, 4 mice)

Figure 2. Learning requires synaptic transmission in lobule simplex. A) Average lever kinematic traces from an example 0.5 s cue prediction session after application of NBQX (300 μM), CPP (30 μM) and MCPG (30 μM). *Top*, traces aligned separately to press and release showing no difference in kinematics from beginning (blue, first 1/3 of trials) to end (black, last 1/3 of trials) of the session. *Bottom*, traces aligned to the cue show that lever releases do not occur sooner in the last 1/3 of trials (black) as compared to the first 1/3 of trials (blue). Shaded area is ±SEM across trials. B) Summary of average lever release times (20-80% rise time) across NBQX 0.5s cue prediction sessions (n=10 sessions, 6 mice). C) Summary of all 0.5 s cue prediction sessions where NBQX (n=14) CPP (n=14) and MCPG (n=8) were applied locally to LS (blue; n=6 mice) compared to summary of cue reaction sessions (black; replotted from Fig 1G). D) Average change in reaction time from the beginning to the end of NBQX 0.5s cue prediction sessions (blue; n=10 sessions, 6 mice) compared to cue prediction and reaction sessions (replotted from Fig. 1H)

Figure 3. Single photon imaging during cue reaction sessions. A) Example ΔF/F single trial timecourses for a correctly timed lever release (left) and an early lever release (right). RW = reward, delivered immediately on correct release. B) Field of view (images separated by 100 ms) showing the average fractional change in fluorescence aligned to lever release for correct (top) and early (bottom) trials from an example session (same session as example from A. C) Calcium transient timecourses averaged across trials from the experiment in A,B) for correct (black) and early trials (red). Error bars are ±SEM across trials. D) Summary of the average peak calcium transients (measured on single trial basis, methods) for all correct and early trials across sessions (n=10 animals, 17 sessions). Colors represent different animals, and each point represents an imaging session. Note that all points lie below the diagonal. *Inset*, schematic of ROIs for all sessions. Outline color corresponds to mouse from scatter plot, and fill color indicates imaging depth identified by post-hoc histology.

Figure 4. Lever dynamics and licking do not explain differences in complex spiking across trial types. A) Average lever kinematics from a representative cue reaction session for correct (black) and early (red) trials aligned to threshold crossing for both press and release. B) Summary of rise times for press (left) and release (right) on correct and early trials across sessions (n=14 sessions, 3 mice). C) Average timecourse of licking for correct (top) and early (bottom) release trials across sessions (n=15 sessions, 9 mice). Error bars are ±SEM across sessions. D) Two example sessions illustrating the relationship between release-evoked calcium transients (black and red lines) and licking (blue bars) for correct (left) and early (right) release trials. E) Summary of the relationship between lick rates and the magnitude of fluorescent transients for correct (black) and early (red) release trials (n= 15 sessions, 9 mice). Error bars are ±SEM across trials. F) Summary of the mean ratio of early and correct trial-evoked peak calcium transient and lick rates for the same sessions in E. G) Summary of the Spearman's correlation (ρ) between lick rate and peak ΔF/F on correct (black) and early (red) lever releases across trials for each session (n=8 sessions, 7 animals). Filled circles are mean and SEM across sessions.

Figure 5. Complex spiking produces larger mean response in individual dendrites and enhanced population responses when movement is correctly timed. A) *Left*, average GCaMP fluorescence from an example 2-photon imaging session (30 Hz acquisition rate). Scale bar = 100 μm. *Middle*, pixel mask of individual PC dendrites extracted from the same session. Right, average calcium responses measured from the highlighted dendrite segregated by trial event. Dotted line represents time of lever press (blue) or lever release (correct: black; early: red). Shaded area is ±SEM across trials. B) Left, average timecourse of calcium response for all significantly responsive dendrites aligned to lever press (blue), or release (correct releases (black), early releases (red), and too-fast releases (green) that occurred < 200 ms following the visual cue; n=17 animals, 30 sessions, 1146 dendrites). Shaded area is ±SEM across dendrites. *Middle*, summary scatter plot comparing the average peak amplitude of the calcium transient for each significantly responsive PC dendrite (gray) and the average of all dendrites (red) for correct and early release trials. *Right*, same as middle for correct and too fast release trials. C) Pixel masks of dendrites significantly responsive to lever release on correct trials, lever release on early trials, and lever press for the example session in A. Color map represents average ΔF/F. D) Summary across experiments of the fraction of dendrites responsive to correct lever releases vs early releases (top, left), to only correct or only early releases (top, right), to lever press vs release (bottom, left), and only to press or release (bottom, right). E) *Left,* average amplitude of correct and early release-evoked calcium events across all dendrites normalized to the amplitude of spontaneous calcium events. *Right,* mean spontaneous (gray) and correct (black) and early (red)

release lever-evoked calcium events across all responsive dendrites Error bars are ±SEM across

cells.

Figure 6. Complex spiking occurs with higher peak rates and greater synchrony when movements

are correctly timed. A) Peri-release time histogram of calcium events on correct (black) and early

(red) release trials (n=17 animals, 30 sessions, 1146 dendrites). Shaded area is ±SEM across

dendrites. B) Summary of average event rate for correct and early releases across sessions.

Error bars are ±SEM across dendrites. C) Same as B for average event latency relative to release.

D) Summary of average standard deviation (S.D.) across events within single dendrites across

trials. Error bars are ±SEM across dendrites. E) Same as D for measurement of S.D. across

dendrites within each experiment. Open circles are statistically significant for reduced correct trial

jitter across dendrites. Error bars are ±SEM across trials.

Figure 7. Complex spiking is correlated across parasagittal zones, with higher correlations on

correct lever releases. A) *Left*, Average GCaMP fluorescence for three example experiments.

ROIs from lobule simplex (white boxes) were selected for unsupervised clustering analysis (meta

K-means, methods). *Right*, voxels are colored according to cluster identity for each example

experiment. M=medial, R=rostral, C=caudal, L=lateral. Note the alignment of the clustered zones

along the rostro-caudal axis. B) Within cluster voxel-by-voxel correlations between correct and

early lever release trials from an example session in A (bottom) for each of the six identified

clusters (yellow). C) Summary of average correlation coefficient for all voxel pairs within an area

across experiments for correct and early release trials. (n=5 animals, 7 sessions, 39 clusters).

Colors denote clusters from the same session. Error bars are ±SEM across voxel pairs. D)

Summary of the relationship between peak $\Delta F/F$ and correlation coefficients for correct (black)

and early (red) trials. Lines are linear fits to each cluster. E) Same as C for pairs of voxels across

neighboring clusters. F) Same as C for pairs of voxels across non-neighboring clusters.

Figure 8. Complex spiking is modulated by learned sensorimotor predictions. A) Summary of

peak calcium transients in a 500 ms window at the time of release (methods) across all single

photon experiments for correct (black) and early (red) release trials binned according to hold time

(250ms bins). Linear fits were applied to data from each trial type (n=10 animals, 17 sessions).

B) Same as in A, for 2-photon imaging (n=17 animals, 30 sessions). C) Mean timecourse of

calcium transients for an example session divided according to early trials with hold durations of

< 1.0s (red) and > 3.5s (dark red). Note the late, second response for long hold duration trials. D)

Single trial example timecourses from an individual dendrite from the session in C). Gray bars

represent analysis windows for subsequent plots E-G) Summary of peak calcium transients

measured in the shaded windows from D for all dendrites (n=1146) according to lever hold time

(E,F) and trial outcome (G). H) Mean calcium transient timecourses averaged across all dendrites

on short and long hold early release trials. I) Mean lick rates for the subset of experiments where

licking was measured in H (n= 6 animals, 10 sessions)). J) Example timecourses of mean calcium

transients from a single session for correct (black), and reward omission (dark blue) trials. Note

the late, second response on omission trials. K) Single trial example timecourses from an

individual dendrite from the session in J. L-M) Summary of peak calcium transients measured in

the shaded windows in K on correct and reward omission lever trials (n=81 dendrites, 3 animals,

3 sessions). N) Mean timecourses of calcium transients averaged across all dendrites for correct

(black) and reward omission (dark blue) trials. Vertical bars represent the lick rate averaged

across trials for all reward omission sessions.

Supplemental Figures

Figure 1. Mapping of motor output driven by Arch and electrical stimulation of superficial

cerebellar cortex. A) Schematics of the results of Arch stimulation over a grid of sites in lobule

simplex (Sim) and Crus I for all five mice. Colors denote outcome of stimulation. B) Same as A

for three mice tested with electrical microstimulation.

Figure 2. Retention and extinguishment of learning for cue prediction regime. A) Average

reaction times relative to cue for the first (left) and second (right) sessions where animals

performed 0.5 s cue prediction on consecutive training days (n=17 sessions, 5 mice). Error bars

are ±SEM across sessions. B) Same as A for 2 s cue prediction sessions (n=10 sessions, 6

mice). C) *Left,* example cumulative distributions of reaction times for cue reaction sessions

immediately surrounding a cue prediction session (session 0). Trials from the cue prediction

session include only the last 1/3 of the session when the animal was anticipating the cue timing.

Right, summary of distribution skewness across cue reaction sessions immediately surrounding

cue prediction sessions. D) Same as A for pairs of 0.5 s prediction sessions separated by 3 or

more cue reaction sessions (n=20 sessions, 10 mice).

Figure 3. Performance on cue prediction task in control and NBQX sessions. A-E) Summary

across experiments of average percent correct lever releases, baseline reaction time from the

visual cue, mean reaction time variance, mean duration to press, number of trials performed, and

the duration of behavioral sessions. n=44 0.5 s, n=9 1.0 s, n=6 1.5 s, n=30 2.0 s, n=26 Δt, n=14

0.5 s NBQX. For each cue delay, the minimum and maximum (x,y) number of trials performed in

single sessions was: 0.5 s: (309, 700) 1 s: (252, 900) 1.5 s: (202, 775) 2 s: (278,1185) Δt: (313,

772) 0.5 s NBQX: (227, 429). F_i) Individual (top) and average (bottom) lever trajectories aligned

to lever press (left) and release(right) segregated by duration into quartiles for a representative 0.5 s cue prediction session. F_{ii}) Same as F_i for a representative 0.5 s cue prediction session in NBQX. G) Summary of press (left) and release (right) quartiles across experiments for control (top; n=15 sessions) and NBQX (bottom; n=14 sessions) experiments. H) Summary of the ratio between press and release ranges (difference between 1^{st} and 4^{th} quartile) for control (black) and NBQX (blue) experiments.

Figure 4. Pharmacological modulation of complex spiking in lobule simplex. A) Single unit recordings during NBQX application to LS. Top, summary of complex spike rates in control and after application of NBQX for individual cells (black; n=9 cells, 5 mice) and the average of all cells (red). Error bars are ±SEM across cells. Bottom, same as top for simple spike rates. Note that NBQX strongly reduces complex but not simple spike rates. B) (top) Raw epifluorescence images from parasagittal cerebellar sections showing fluorescein labeling after surface application to lobule simplex *in vivo* (methods, 3 mice). (bottom) Pixel masks from the same images above. Pixels were thresholded at 30% of maximum value to visualize and quantify fluorescein labeling. C) Summary of mediolateral fluorescein spread across sections. (n=3 animals).

Figure 5. Segmentation of PC dendrites from imaging experiments. A) Example single photon imaging field of view (left) is segmented to mask individual dendrites (right) used for subsequent single cell analysis. Scale bar = 100 μm. B) Example timecourse of raw fluorescence (top) from the circled dendrite in A showing individual calcium transients identified (gray circles) according to peaks in the first derivative of the fluorescence trace (bottom). C) Same as A for an example two photon field of view. Note that while PC somata are visible (round, top right), image segmentation does not extract activity from these structures. Scale bar = 100 μm. D) Same as B for the experiment in C. E) Top, example time course of a single unit PC recording from an awake mouse illustrating detection of complex spikes (gray circles). Bottom, overlay of individual simple (left) and complex (right) spike waveforms (gray) and the average waveform (black). F) Summary of spike rates across all two photon dendrite imaging sessions (blue; n=1051 dendrites) and acute single unit recordings (n=11 units).

Figure 6. Single unit PC recordings of complex spiking during the cue prediction task. A) Schematic of the trial structure schematic, where a constant cue delay of 500ms was imposed on each trial. B) Mean normalized complex spike rate (methods) from single unit recordings (40 PCs, 8 mice) aligned to lever release for correct (black) and early (red) trials during the cue prediction task.

Figure 7. Differential complex spiking on correct and early release trials is not driven by motor signals due to licking. A) Example imaging data from an inter-trial interval (ITI) showing a lick responsive cell (top, dark blue) and a lick unresponsive cell (bottom, black) in the same field of view. B) Averaged lick triggered calcium transients for the lick responsive cell (dark blue) and a lick unresponsive cell (black) in A). C) Mean lick triggered calcium transient across all lick responsive (dark blue) and lick unresponsive (black) cells. D) Summary of the Spearman's correlation between lick rate and amplitude of the calcium transient in each frame across trials for lick responsive (dark blue) and lick unresponsive cells (grey). Red points indicate the mean $\pm$ SEM across dendrites (447 dendrites, 6 animals and 10 sessions) E) Mean lick triggered calcium transients for lick responsive neurons only in response to either a single lick (dark blue) or a lick bout (3 or more licks with $\leq$ 300 ms between licks, blue). F) Mean lever release-evoked calcium transients across neurons for correct (black) and early (red) lever releases for lick unresponsive neurons only. G) Summary comparing the amplitude of calcium transients on correct and early lever releases for lick unresponsive cells only.

Figure 8. Example single trial and session average calcium transients from 2-photon imaging. A) Example single trial data from an individual dendrite showing the normalized calcium transient across an early release trial from before lever press to after the lever release. B) Same as A) but for a correct trial, including the time of cue presentation. C) Mean calcium transients across trials for the dendrite in A and B. D) Single trial example, aligned to lever release, showing all neurons from the experiment illustrated in Fig. 5C on a correct trial over 1 second surrounding release. Lever press is off scale. E) Same as D, but for an early release trial from the same imaging session.

Figure 9. Mice have full control over the lever trajectory. A) Individual (top) and average (bottom) lever trajectories aligned to lever press (left) and release (right) for manual lever depression followed by unperturbed gravity return. B) Summary of press (left) and release (right) quartiles across experiments for manual lever experiments (n=4). Note the lack of variability across quartiles. C) Summary scatterplot comparing 20-80% rise times of lever press and release for experiments where the lever was controlled by mice (black) and manual press (green) D) Summary comparison measuring the difference from the mean of the slowest release quartile and the fastest release quartile for experiments where the lever was controlled by mice (black) and manual press (green). Lever trajectories were sorted according to duration in a window from 200ms before threshold crossing to the time of a threshold crossing half way between the top and bottom of the total lever displacement. The slowest and fastest 25 trials were extracted for comparison, and their average rise time was normalized by subtracting that of the mean trajectory

717 of the whole session. These results show that mice produce movements both slower and faster
718 than the mean, indicating full control over the lever trajectory.
719

Methods:

Mice

All experimental procedures using animals were carried out with the approval of the Duke
University Animal Care and Use Committee. All experiments were performed during light cycle
using adult mice (>p60) of both sexes, randomly selected from breeding litters. All mice were
housed in a vivarium with normal light/dark cycles in cages with 1-5 mice. Imaging experiments
were performed using Tg(PCP2-Cre)3555Jdhu mice (Jackson Labs, 010536; n=12). Optogenetic
mapping experiments were conducted in PCP2-Cre animals crossed with Ai35(RCL-Arch/GFP)
(Jax 012735; n=5 mice). Single unit recording during the cue prediction task was performed in
wild type C57BL/6J mice (n= 8 mice). Additional behavioral experiments where imaging was not
performed were conducted in wild type C57BL/6J mice (n=18). We used two exclusion criteria for
animals in this study: (1) poor recovery or other health concerns following surgical intervention or
(2) missed virus injection, as determined by *in vivo* imaging and post-hoc histological analysis.
733

Surgical Procedures

3-10 hours prior to surgery, animals received dexamethasone (3mg/kg) and ketoprofen (5mg/kg).
Surgical procedures were performed under anesthesia, using an initial dose of ketamine/xylazine
(50mg/kg and 5mg/kg) 5 minutes prior to surgery and sustained during surgery with 1.0-2.0%
isoflurane. Toe pinches and breathing were used to monitor anesthesia levels throughout
surgeries. Body temperature was maintained using a heating pad (TC-1000 CWE Inc.). Custom-
made titanium headplates (HE Parmer) were secured to the skull using Metabond (Parkell). For
imaging experiments, a 3mm diameter craniotomy was made over the lobule simplex
approximately 1.4mm lateral and 2.8mm posterior to lambda, and glass cover slips consisting of
two 3mm bonded to a 5mm coverslip (Warner Instruments No. 1) with index matched adhesive
(Norland No. 1) were secured in the craniotomy using Metabond. Buprenex (0.05mg/kg) and
cefazolin (50mg/kg) were administered following surgery twice a day for two days. Following a
minimum of 4 recovery days, animals were water deprived for 3 days, or until body weight
stabilized at 85% of initial weight, and were habituated to head restraint (3-5 days) prior to
behavioral training.
For imaging experiments, the glass cover slip was removed following behavioral training,
and mice were injected (WPI UMP3) with AAV1.CAG.Flex.GCaMP6f.WPRE.SV40 (UPenn vector
core, titer = 9.40×10^{12} or 7.60×10^{12}). 150 nL virus diluted 1:1-1:5 in ACSF was injected at a rate of

30nl/min and a depth of 150 μm at 1-3 sites in dorsal lobule simplex. Imaging was performed

beginning 14 days following injection.

For *in vivo* pharmacology experiments, a 3mm craniotomy was performed over lobule

simplex of headposted mice, and the dura mater was peeled back at the center of the craniotomy.

10uL of combined NBQX (300 μM) and CPP (30 μM) was applied into a well surrounding the

craniotomy on the surface of the cerebellum 20 minutes prior to behavioral training. In 8 of 14

experiments, MCPG (30 μM) was also included to block metabotropic glutamate receptors.

Because there no significant differences in performance across these two groups, all data were

pooled for analysis. During behavior, a second drug application of 10uL was administered 20 and

40 minutes into the task, and depending on the duration of training, a third dose was applied after

1 hour. Craniotomies were subsequently covered with silicone elastomer (WPI, Inc.) prior to

returning animals to their home cage. To quantify the spread of pharmacological agents in the

cerebellum, 10 μL of fluorescein dye (Sigma-Aldrich #F6377, 1mM in aCSF) was applied to the

craniotomy in 3 animals with the same method and timecourse as for drug applications. Following

standard histological processing and imaging (below), labeling was quantified by creating a binary

pixel mask thresholded at 30% of the maximum fluorescence value for each experiment. Dye

labeling was then measured using the area of these pixel masks (Supp. Fig. 4 B,C).

Behavior

During behavioral training, animals were head-fixed and placed in front of a computer monitor,

lever and reward delivery tube. Animals were trained to self-initiate trials by depressing the lever

using their right forepaw, and required to successfully hold the lever in the down position for

randomized intervals ranging between 500ms and 5 s on the cue reaction paradigm before

performing the cue prediction paradigm. For both paradigms, a high contrast hold cue was present

at all times, including the intertrial interval (ITI), and transitioned 90 degrees to the release cue on

each trial at the instructed time of release until the animal either released the lever or 1 second

had passed. Lever releases within one second of the release cue were rewarded immediately at

the time of lever release (0.01 M saccharine). Immediately following any lever release, a solenoid

was engaged to prevent lever press during the ITI. Following the ITI (3-6 s), the solenoid is

lowered, allowing the mouse to self initiate a new trial. During training, a 1-3 second 'timeout'

was implement to punish early lever releases. No timeouts were used in fully trained animals for

imaging or behavior data collection sessions. For reward omission sessions, 20% of randomly

determined correct lever releases were unrewarded. Animals used for imaging experiments

performed the cue reaction task with a mean peak percent correct of 79.7 ± 0.23% achieved in

26 ± 2 training days. On imaging days animals performed a range of 150 to 580 trials per session

with a mean of 273.9 ±13.1 trials. Behavioral parameters including lever press, lever release and

licking were monitored using Mworks (http://mworks-project.org) and custom software written in
MATLAB (Mathworks). To assess the degree of lever control by the mice, the dynamics of lever
press and release trajectories were compared to lever presses initiated by the experimenter
where the lever was allowed to return to the rest position on its own (Supp. Fig. 9). These results
demonstrate that mice moved the lever both faster and slower than its intrinsic kinematics, and
thus had full control over its trajectory. Licking was measured with electrical contact circuit. Inter-
trial intervals ranged from 3 to 5 seconds.
795
<u>Calcium Imaging</u>
Wide-field imaging: Single photon imaging was performed using a customized microscope (Sutter
SOM) affixed with a 5x objective (Mitutoyo, 0.14NA) and CMOS camera (Qimaging, Rolera em-
c^2). Excitation (470 nm) was provided by an LED (ThorLabs, M470L3), and data were collected
through a green filter (520-536 nm band pass, Edmund Optics) at a frame rate of 10Hz, with a
field of view of 3.5x3.5mm at 1002x1004 pixels.
802
Two-Photon Imaging: Two-photon imaging was performed with a resonant scanning microscope
(Neurolabware) using a 16x water immersion objective (Nikon CFI75 LWD 16xW 0.80NA).
Imaging was performed using a polymer to stabilize the immersion solution (MakingCosmetics,
0.4% Carbomer 940). A Ti:Sapphire laser tuned to 920nm (Spectra Physics, Mai Tai eHP
DeepSee) was raster scanned via a resonant galvanometer (8 kHz, Cambridge Technology) onto
the brain at a frame rate of either 30 Hz with a field of view of either 1030 μm x 581 μm (796 x
264 pixels) or 555 μm x 233 μm (796 x 264 pixels), or 15.5Hz with a field of view of 555 μm x 452
μm and (796 x 512 pixels). Data were collected through a green filter (510 ± 42 nm band filter
(Semrock)) onto GaAsP photomultipliers (H10770B-40, Hamamatsu) using Scanbox software
(Neurolabware). A total of 12 mice were used for imaging experiments (10 mice for wide-field, 11
mice for two-photon, 12 mice total).
814
<u>Single Unit Recordings and Optogenetics</u>
Acute single unit recordings were performed in awake animals by performing a craniotomy over
lobule simplex and inserting a multi-electrode silicone probe (Neuronexus, A4x8-5mm-100-400-
177-A32, 4 shanks, 8 site/shank at 100 µm spacing) using a Cerebus multichannel acquisition
system (Blackrock Microsystems, Salt Lake City). For single unit recordings obtained in the cue
prediction task (Supp. Fig. 6), chronically implanted electrode arrays were used (Dual drive
movable electrode bundles with 8 tungsten electrodes (23 µm) in each cannula or 16 tungsten
electrodes bundle in one cannula, Innovative Neurophysioloy Inc). Electrode arrays were
implanted using stereotaxic coordinates to target lobule simplex at AP 6.2; ML 2.0. Electrode

bundles were inserted into lobule simplex at a depth of approximately 0.2-0.3 mm. The implant
was encased in Metabond for stability.

For both recording conditions, continuous recording data was bandpass filtered with a 2-
pole Butterworth between 250 Hz and 5 kHz and referenced against an electrode with no spikes
using Spike2. Single units were isolated by amplitude thresholding. Template waveforms were
defined and characterized by their width and peak, and PCA of waveforms was done in off-line in
Spike2.

Complex spikes and simple spikes were discriminated as in de Solages et al.[61], first on
the basis of their stereotypical waveform. Complex spikes typically had a multi-wavelet form,
including a large positive peak within 6ms following spike initiation. Manual identification of 10-
20 complex spikes was used to generate a mean template waveform, which was then compared
to all other spikes for a given unit using the Spearman rank order correlation coefficient. The
combination of Spearman coefficients and the magnitude of the positive waveform deflection was
used to segregate complex and simple spikes. The presence of a post-complex spike pause (20
ms or more) in simple spike firing was verified by cross-correlogram for all isolated single units.
Complex spike rates for the cue prediction task were normalized to the baseline firing rate
determined in a one-second window during the ITI according to: (FR-baseline)/baseline) (Supp.
Fig. 6).

For optogenetic mapping of the dorsal cerebellum, an optical fiber (0.39 NA, 400 μm core
multimode, ThorLabs FT400EMT) coupled to a 532 nm laser (Optoengine, MGL-III-532) was
positioned above the cranial window using a micromanipulator (Scientifica PatchStar).

Histology
Mice were deeply anesthetized with ketamine/xylazine (200mg/kg & 30mg/kg respectively, IP)
then perfused with PBS then 4% paraformaldehyde. 100um sagittal sections were cut using a
vibratome (Pelco 102). Cerebellar slices were then mounted using a mounting medium (Southern
Biotech Fluoromount-G or DAPI Fluoromount-G) and imaged with a fluorescence microscope
(Nikon Eclipse 80i).

Data Analysis
Behavior: Behavior sessions were only analyzed within the time range of active task
performance. Accordingly, the last trial of each session was determined by the occurrence of
either two consecutive failed trials (in which the mouse did not release the lever in response to
the cue) or two consecutive trials with post-ITI duration to press longer than 1.5x the session
average.

Reaction times were measured using lever releases within the reward window as well as

those occurring 200ms prior to the reward cue to account for predictive responses. Reaction times

are plotted as a binned average of five trials. Initial reaction times were calculated according to

the y-intercept of the linear regression of the first fifty trials. Final reaction time at the end of each

session was calculated by averaging the reaction times of trials 200-250.

The symmetry of reaction time data about the sample mean was measured to test the

extinguishment of learning after following cue prediction sessions according to skewness.

Sample skewness was defined by

868
$$s = \frac{E(\mu - X)^3}{\sigma^3}$$

870 where μ is the mean of X, $E(*)$ is the expected value of the quantity $*$, and σ is the standard

871 deviation of X. Thus, for a given sample x, this calculates:

872
$$\frac{\frac{1}{n}\sum_{i=1}^{n}(x_i - \bar{x})^3}{\left(\sqrt{\frac{1}{n}\sum_{i=1}^{n}(x_i - \bar{x})^2}\right)^3}$$

for $x_i \in x$, where $\bar{x}$ is the sample mean.

Movement trajectories were measured using a lever affixed to a rotary encoder (US

Digital) with a 60mm radius and 1250 pulses per revolution. Encoder values were collected at

1000 Hz, and trajectories were calculated by up-sampling at 200 µs (5000 Hz) using nearest-

neighbor interpolation. Pulse number was converted to degrees, and the vertical displacement in

millimeters was calculated using the chord length of the leaver displacement angle. 20-80% rise

times were calculated using normalized average trajectories. Presses and releases were sorted

according to duration in a window from 200ms before threshold crossing to the time of a threshold

crossing half way between the top and bottom of the total lever displacement.

Wide-field imaging: Imaging analysis were performed using custom MATLAB code. Regions of

interest (ROIs) were selected within empirically defined forelimb movement region of lobule

simplex according to the spread of GCaMP expression. Window location and the lobule identity

were identified according to folia patterns visible through the cranial window, landmarks recorded

during surgery, and post-hoc histology. Baseline fluorescence (F) was measured on a trial-by-trial

basis during the inter-trial interval (ITI) as the mean fluorescence 900 to 200ms before trial

initiation. Normalized fluorescence (ΔF/F) was calculated according to the cumulative activity

within an ROI, or on a pixel by pixel basis. Lever hold times <200ms in total, or > 1 s after the

visual cue were exceedingly rare, and thus excluded from analysis. Lever releases between

200ms and 1000ms following the visual cue were classified as correct, and releases prior to visual

cue were classified as early. Lever releases $\leq$200 ms following the visual cue were considered

too fast to be reactions to the cue based on average reaction time distributions. Peak ΔF/F was

measured on a trial by trial basis in the time window from 100ms before to 400 ms after lever

release. Note that differences in the timing of single trial calcium transient peaks produce smaller

amplitudes for ΔF/F timecourses as compared to reported peak ΔF/F measurements.

Spearman's correlation between lick rate and peak ΔF/F within sessions was calculated according

to the lick rate within 500 ms following each lever release and the peak calcium transient for each

trial. For this correlation analysis, trials without licking were removed, and only sessions with 7 or

more trials in each condition were included.

Meta K-means analysis and Cluster Correlations: Images were first registered to reduce motion

artifacts, and then thresholded at 70% of maximum intensity for each frame to remove background

noise. Images were downsampled 5-fold in both X and Y. Baseline F was defined as the averaged

fluorescence across the entire movie for each pixel, and used to normalize change in fluorescence

(ΔF/F). ΔF/F was then re-normalized to the maximum ΔF/F during the entire movie for each pixel.

The repeated k-means clustering algorithm (meta-k-means) separated pixels to cluster centroids

based on a pairwise correlation distance function using Pearson's linear correlation coefficient (r).

The final clusters were determined by first thresholding all the k-means results at 800 out of 1000

runs, and then by merging highly correlated clusters based on Dunn's index. Clusters occupying

less than 3% of the total imaging field were excluded from further analysis. Intra-cluster and inter-

cluster correlation coefficients were calculated between 100 ms before and 300 ms after lever

release for both correct and early trials on a frame by frame basis. To test for differences in the

relationship between spike rates and correlation coefficients between trial types, paired clusters

from each distribution were resampled with replacement and fit with a line to measure the y-

intercept 1000 times. Statistical significance was computed according to the 95% confidence

interval of the distribution of y-intercept difference values (difference between y-intercepts 0.0375

$\pm$ 0.0013, 95% confidence interval [0.0349, 0.0402]).

Two-photon imaging: Motion in the XY plane was corrected by sub-pixel image registration. To

isolate signals from individual PC dendrites, we utilized principal component analysis (PCA)

followed by independent component analysis (ICA). Final dendrite segmentation was achieved

by thresholding the smoothed spatial filters from ICA. A binary mask was created by combining

highly correlated pixels (correlation coefficient > 0.8) and removing any overlapped regions

between segmented dendrites. Notably, image segmentation using these criteria did not extract

PC soma, which were clearly visible in many single- and two-photon imaging experiments.

Fluorescence changes (ΔF) were normalized to a window of baseline fluorescence (F) between

500ms and 100ms preceding trial initiation. Responses were categorized as significant (p<0.05)

according to a one-tailed t-test if the ΔF/F in a 200 ms window surrounding lever press or release

was larger than that from a 500 ms window immediately preceding lever movement. To extract

events on single trials used for estimation of complex spike rates and event amplitudes, the first

derivative of the raw fluorescence trace was thresholded at 2.5 standard deviations from baseline.

The same methods were used to extract events for the subset of single photon experiments

described in Supp. Fig. 5. For single trial amplitude measurements, only events well separated

in time (greater than 650ms from the next event) were considered; however, for measurements

of rates, all events separated by at least one frame were included. Note that rate estimates are

thus likely to be lower than actual rates, particularly for single photon imaging where data were

collected at 100 ms intervals. Peak calcium transients on reward omission trials and on early

release trials were quantified in two time windows (Fig. 8). Window 1 spanned the first 100 ms

after lever release, and window 2 spanned 175 ms that began 165ms after lever release.

Spearman's correlation was calculated between lick rate and the peak calcium transient within

sessions on a frame by frame basis for each trial in a window spanning 500 ms after lever release.

Correlation values for correct and early release trials for each cell were averaged separately.

Some neurons showed responses to licking, as defined by a mean lick-triggered calcium transient

that was significantly larger (p<0.05) with respect to baseline (Supp. Fig. 7). Lick triggered

averages were constructed from licks presumed to be spontaneous that occurred during the inter-

trial interval (Supp. Fig. 7).

Additional statistics and data sharing: Data are presented as mean ± S.E.M., unless stated

otherwise. Statistical tests were two-sided, and analyses of variance (ANOVA) were performed

when more than two groups were tested. Differences were considered statistically significant

when $P < 0.05$. No correction for multiple comparisons was applied. No statistical methods were

used to predetermine sample sizes. Data collection and analysis were not performed blind to the

conditions of the experiments, but data collection relied on automatized measurements and

subsequent analysis was based on code uniformly applied across experimental conditions. Data

analysis code are available on GitHub (access available upon request). The data that support the

findings of this study are available from the corresponding author upon reasonable request.

References

1.   Albus, J.S. A theory of cerebellar function. *Mathematical Biosciences* **10**, 25-61 (1971).

2.   Ito, M. Neural design of the cerebellar motor control system. *Brain Res* **40**, 81-84 (1972).

3.   Marr, D. A theory of cerebellar cortex. *J Physiol* **202**, 437-470 (1969).

4.   Ito, M., Sakurai, M. & Tongroach, P. Climbing fibre induced depression of both mossy fibre
responsiveness and glutamate sensitivity of cerebellar Purkinje cells. *J Physiol* **324**, 113-134
(1982).

5.   Ekerot, C.F. & Kano, M. Long-term depression of parallel fibre synapses following
stimulation of climbing fibres. *Brain Res* **342**, 357-360 (1985).

6.   Sakurai, M. Synaptic modification of parallel fibre-Purkinje cell transmission in in vitro
guinea-pig cerebellar slices. *J Physiol* **394**, 463-480 (1987).

7.   Miyakawa, H., Lev-Ram, V., Lasser-Ross, N. & Ross, W.N. Calcium transients evoked by
climbing fiber and parallel fiber synaptic inputs in guinea pig cerebellar Purkinje neurons. *J
Neurophysiol* **68**, 1178-1189 (1992).

8.   Tank, D.W., Sugimori, M., Connor, J.A. & Llinas, R.R. Spatially resolved calcium dynamics
of mammalian Purkinje cells in cerebellar slice. *Science* **242**, 773-777 (1988).

9.   Konnerth, A., Dreessen, J. & Augustine, G.J. Brief dendritic calcium signals initiate long-
lasting synaptic depression in cerebellar Purkinje cells. *Proc Natl Acad Sci U S A* **89**, 7051-7055
(1992).

10.   Llinas, R. & Sugimori, M. electrophysiological properties of in vitro purkinje cell somata in
mammalian cerebellar slices. *J. Physiol.*, 171-195 (1980).

11.   Eccles, J.C., Llinas, R. & Sasaki, K. The excitatory synaptic action of climbing fibres on the
Purkinje cells of the cerebellum. *J Physiol* **182**, 268-296 (1966).

12.   Medina, J.F., Nores, W.L., Ohyama, T. & Mauk, M.D. Mechanisms of cerebellar learning
suggested by eyelid conditioning. *Current opinion in neurobiology* **10**, 717-724 (2000).

13.   Ohyama, T.N.W.M.M. & Mauk, M. What the cerebellum computes. *Trends in
Neurosciences* **26**, 222-227 (2003).

14.   Raymond, J.L., Lisberger, S.G. & Mauk, M.D. The cerebellum: a neuronal learning
machine? *Science* **272**, 1126-1131 (1996).

15.   Schultz, S.R., Kitamura, K., Post-Uiterweer, A., Krupic, J. & Hausser, M. Spatial pattern
coding of sensory information by climbing fiber-evoked calcium signals in networks of
neighboring cerebellar Purkinje cells. *J Neurosci* **29**, 8005-8015 (2009).

16. Tsutsumi, S.*, et al.* Structure-function relationships between aldolase C/zebrin II 996 expression and complex spike synchrony in the cerebellum. *J Neurosci* **35**, 843-852 (2015).

17. Ozden, I., Dombeck, D.A., Hoogland, T.M., Tank, D.W. & Wang, S.S. Widespread state-998 dependent shifts in cerebellar activity in locomoting mice. *PLoS One* **7**, e42650 (2012).

18. De Gruijl, J.R., Hoogland, T.M. & De Zeeuw, C.I. Behavioral correlates of complex spike 1000 synchrony in cerebellar microzones. *J Neurosci* **34**, 8937-8947 (2014).

19. Tang, T., Suh, C.Y., Blenkinsop, T.A. & Lang, E.J. Synchrony is Key: Complex Spike Inhibition 1002 of the Deep Cerebellar Nuclei. *Cerebellum* **15**, 10-13 (2016).

20. Lee, K.H.*, et al.* Circuit mechanisms underlying motor memory formation in the 1004 cerebellum. *Neuron* **86**, 529-540 (2015).

21. Histed, M.H., Carvalho, L.A. & Maunsell, J.H. Psychophysical measurement of contrast 1006 sensitivity in the behaving mouse. *J Neurophysiol* **107**, 758-765 (2012).

22. Chettih, S.N., McDougle, S.D., Ruffolo, L.I. & Medina, J.F. Adaptive timing of motor output 1008 in the mouse: the role of movement oscillations in eyelid conditioning. *Front Integr Neurosci* **5**, 1009 72 (2011).

23. Gaffield, M.A., Amat, S.B., Bito, H. & Christie, J.M. Chronic imaging of movement-related 1011 Purkinje cell calcium activity in awake behaving mice. *J Neurophysiol* **115**, 413-422 (2016).

24. Mukamel, E.A., Nimmerjahn, A. & Schnitzer, M.J. Automated analysis of cellular signals 1013 from large-scale calcium imaging data. *Neuron* **63**, 747-760 (2009).

25. Najafi, F., Giovannucci, A., Wang, S.S. & Medina, J.F. Coding of stimulus strength via analog 1015 calcium signals in Purkinje cell dendrites of awake mice. *Elife* **3**, e03663 (2014).

26. Ozden, I., Lee, H.M., Sullivan, M.R. & Wang, S.S. Identification and clustering of event 1017 patterns from in vivo multiphoton optical recordings of neuronal ensembles. *J Neurophysiol* **100**, 1018 495-503 (2008).

27. Ozden, I., Sullivan, M.R., Lee, H.M. & Wang, S.S. Reliable coding emerges from 1020 coactivation of climbing fibers in microbands of cerebellar Purkinje neurons. *J Neurosci* **29**, 1021 10463-10473 (2009).

28. Najafi, F., Giovannucci, A., Wang, S.S. & Medina, J.F. Sensory-driven enhancement of 1023 calcium signals in individual Purkinje cell dendrites of awake mice. *Cell Rep* **6**, 792-798 (2014).

29. Najafi, F. & Medina, J.F. Beyond "all-or-nothing" climbing fibers: graded representation of 1025 teaching signals in Purkinje cells. *Front Neural Circuits* **7**, 115 (2013).

30.   Yang, Y. & Lisberger, S.G. Purkinje-cell plasticity and cerebellar motor learning are graded
by complex-spike duration. *Nature* **510**, 529-532 (2014).

31.   Rudolph, S., Overstreet-Wadiche, L. & Wadiche, J.I. Desynchronization of multivesicular
release enhances Purkinje cell output. *Neuron* **70**, 991-1004 (2011).

32.   Maruta, J., Hensbroek, R.A. & Simpson, J.I. Intraburst and interburst signaling by climbing
fibers. *J Neurosci* **27**, 11263-11270 (2007).

33.   Kitamura, K. & Hausser, M. Dendritic calcium signaling triggered by spontaneous and
sensory-evoked climbing fiber input to cerebellar Purkinje cells in vivo. *J Neurosci* **31**, 10847-
10858 (2011).

34.   Brenowitz, S.D. & Regehr, W.G. Associative short-term synaptic plasticity mediated by
endocannabinoids. *Neuron* **45**, 419-431 (2005).

35.   Eilers, J., Augustine, G.J. & Konnerth, A. Subthreshold synaptic Ca2+ signalling in fine
dendrites and spines of cerebellar Purkinje neurons. *Nature* **373**, 155-158 (1995).

36.   Hartell, N.A. Strong activation of parallel fibers produces localized calcium transients and
a form of LTD that spreads to distant synapses. *Neuron* **16**, 601-610 (1996).

37.   Wang, S.S., Denk, W. & Hausser, M. Coincidence detection in single dendritic spines
mediated by calcium release. *Nat Neurosci* **3**, 1266-1273 (2000).

38.   Andersson, G. & Oscarsson, O. Climbing fiber microzones in cerebellar vermis and their
projection to different groups of cells in the lateral vestibular nucleus. *Exp Brain Res* **32**, 565-579
(1978).

39.   Apps, R. & Garwicz, M. Anatomical and physiological foundations of cerebellar
information processing. *Nat Rev Neurosci* **6**, 297-311 (2005).

40.   Ackerley, R., Pardoe, J. & Apps, R. A novel site of synaptic relay for climbing fibre pathways
relaying signals from the motor cortex to the cerebellar cortical C1 zone. *J Physiol* **576**, 503-518
(2006).

41.   Kitazawa, S., Kimura, T. & Yin, P.B. Cerebellar complex spikes encode both destinations
and errors in arm movements. *Nature* **392**, 494-497 (1998).

42.   Sugihara, I. & Shinoda, Y. Molecular, topographic, and functional organization of the
cerebellar cortex: a study with combined aldolase C and olivocerebellar labeling. *J Neurosci* **24**,
8771-8785 (2004).

43.   Sugihara, I., Wu, H.S. & Shinoda, Y. The entire trajectories of single olivocerebellar axons

in the cerebellar cortex and their contribution to Cerebellar compartmentalization. *J Neurosci* **21**,

7715-7723 (2001).

44.   Voogd, J., Pardoe, J., Ruigrok, T.J. & Apps, R. The distribution of climbing and mossy fiber

collateral branches from the copula pyramidis and the paramedian lobule: congruence of

climbing fiber cortical zones and the pattern of zebrin banding within the rat cerebellum. *J*

Neurosci **23**, 4645-4656 (2003).

45.   Ekerot, C.F., Garwicz, M. & Schouenborg, J. Topography and nociceptive receptive fields

of climbing fibres projecting to the cerebellar anterior lobe in the cat. *J Physiol* **441**, 257-274

(1991).

46.   Oberdick, J., Baader, S.L. & Schilling, K. From zebra stripes to postal zones: deciphering

patterns of gene expression in the cerebellum. *Trends Neurosci* **21**, 383-390 (1998).

47.   Zhou, H*., et al.* Cerebellar modules operate at different frequencies. *Elife* **3**, e02536

(2014).

48.   Paukert, M., Huang, Y.H., Tanaka, K., Rothstein, J.D. & Bergles, D.E. Zones of enhanced

glutamate release from climbing fibers in the mammalian cerebellum. *J Neurosci* **30**, 7290-7299

(2010).

49.   Blenkinsop, T.A. & Lang, E.J. Synaptic action of the olivocerebellar system on cerebellar

nuclear spike activity. *J Neurosci* **31**, 14708-14720 (2011).

50.   Bengtsson, F., Ekerot, C.F. & Jorntell, H. In vivo analysis of inhibitory synaptic inputs and

rebounds in deep cerebellar nuclear neurons. *PLoS One* **6**, e18822 (2011).

51.   Person, A.L. & Raman, I.M. Purkinje neuron synchrony elicits time-locked spiking in the

cerebellar nuclei. *Nature* **481**, 502-505 (2012).

52.   McElvain, L.E., Bagnall, M.W., Sakatos, A. & du Lac, S. Bidirectional plasticity gated by

hyperpolarization controls the gain of postsynaptic firing responses at central vestibular nerve

synapses. *Neuron* **68**, 763-775 (2010).

53.   Pugh, J.R. & Raman, I.M. Potentiation of mossy fiber EPSCs in the cerebellar nuclei by

NMDA receptor activation followed by postinhibitory rebound current. *Neuron* **51**, 113-123

(2006).

54.   Miles, F.A. & Lisberger, S.G. Plasticity in the vestibulo-ocular reflex: a new hypothesis.

Annu Rev Neurosci **4**, 273-299 (1981).

55.   Ohmae, S. & Medina, J.F. Climbing fibers encode a temporal-difference prediction error

during cerebellar learning in mice. *Nat Neurosci* **18**, 1798-1803 (2015).

56.   Matias, S., Lottem, E., Dugue, G.P. & Mainen, Z.F. Activity patterns of serotonin neurons

underlying cognitive flexibility. *Elife* **6** (2017).

57.   Belova, M.A., Paton, J.J., Morrison, S.E. & Salzman, C.D. Expectation modulates neural

responses to pleasant and aversive stimuli in primate amygdala. *Neuron* **55**, 970-984 (2007).

58.   Hayden, B.Y., Heilbronner, S.R., Pearson, J.M. & Platt, M.L. Surprise signals in anterior

cingulate cortex: neuronal encoding of unsigned reward prediction errors driving adjustment in

behavior. *J Neurosci* **31**, 4178-4187 (2011).

59.   Pearce, J.M. & Hall, G. A model for Pavlovian learning: variations in the effectiveness of

conditioned but not of unconditioned stimuli. *Psychol Rev* **87**, 532-552 (1980).

60.   Roesch, M.R., Esber, G.R., Li, J., Daw, N.D. & Schoenbaum, G. Surprise! Neural correlates

of Pearce-Hall and Rescorla-Wagner coexist within the brain. *Eur J Neurosci* **35**, 1190-1200 (2012).

61.   de Solages, C.*, et al.* High-frequency organization and synchrony of activity in the purkinje

cell layer of the cerebellum. *Neuron* **58**, 775-788 (2008).

1102

Figure 1

Figure 2

A Single trials

Press Cue Release Press Release
                + RW

Correct   Early

0.05 ΔF/F
500 ms

B Session Avg.

1 mm

Correct

0.15

ΔF/F

-0.5   0   0.5   1.0

Early

-0.15

-0.5   0   0.5   1.0

Time from lever release (s)

C Session Avg.

Correct
Early

press   release

0.02 ΔF/F
200 ms

D

Early trial response (ΔF/F)

depth (μm)

1mm

Correct trial response (ΔF/F)

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

A

B

Arch: 5 mice
10-50mW, one Step pulse for 1sec

Right arm •
Right & Left arms ●
Very weak arm movements •
Whiskers or mouth or nose •
nothing •

Electrical Stim. 3 mice
0.1mA-1mA
50Hz for 500ms or 1sec

Right arm •
Right & Left arms ●
Very weak arm movements •
Whiskers or mouth or nose •
nothing •

supplemental figure 1

supplemental figure 2

supplemental figure 3

A

CS rate (Hz)

control
0.60 ± 0.07

NBQX
0.27 ± 0.04

SS rate (Hz)

control
66.8 ± 17.8

NBQX
45.9 ± 13.0

B

Fluorescein

dorsal →

4/5
6
7

lateral →

LS
Crus1
IntA

Crus1
Crus2
PM

Pixel masks

dorsal →

lateral →

C

pixel mask area (mm²)

-2   -1   0   -1

← medial   lateral →

Distance from LS center (mm)

supplemental figure 4

A
mesoscale imaging

Field of view   1P dendrites

B
raw F trace

1st derivative

500 ms

C
2P imaging

Field of view

2P dendrites

D
raw F trace

1st derivative

1 s

E
50 µV
100 ms

400 µV
1 ms

F
0.6 ± 0.01 Hz   0.6 ± 0.06 Hz
n = 1051   n = 11

2P dendrites
single units

Event rate (Hz)

supplemental figure 5

supplemental figure 6

supplemental figure 7

A Early, single trial

press   release

0.1 ΔF/F
500 ms

B Correct, single trial

press   cue release

0.1 ΔF/F
500 ms

C Session Avg.

press   release

correct
early

n = 115   0.05 ΔF/F
500 ms

D Correct, single trial

neurons

release

E Early, single trial

neurons

press   release

1.0 ΔF/F
200 ms

supplemental figure 8

supplemental figure 9

Reviewer Figure