1   **Early-life social environment alters juvenile behavior and neuroendocrine function in a**

2   **highly social cichlid fish**

6   Tessa K. Solomon-Lane[1,2*] & Hans A. Hofmann[1-3]

8   [1]Department of Integrative Biology, [2]Institute for Neuroscience, [3]Center for Computational

9   Biology and Bioinformatics, The University of Texas at Austin, Austin, TX 78712

*Corresponding author:

Tessa Solomon-Lane

University of Texas at Austin

Department of Integrative Biology

1 University Station #C0930

Austin, TX 78712

tksolomonlane@utexas.edu

512-475-7318

Abstract

Early-life experiences can shape adult behavior, with consequences for fitness and health, yet

fundamental questions remain unanswered about how early social environments and experiences

are translated into variation in brain and behavior. The African cichlid fish *Astatotilapia burtoni*,

a model system in social neuroscience, is well known for its highly plastic social phenotypes in

adulthood. Here, we rear juveniles in either social groups or pairs to investigate the effects of

early-life social environments on behavior and neuroendocrine gene expression. We find that

both juvenile behavior and neuroendocrine function are sensitive to early-life social effects.

Behavior robustly co-varies across multiple contexts (open field, social cue investigation, and

dominance behavior assays) to form a behavioral syndrome. Rearing environment shifts pair-

reared juveniles towards the end of syndrome that is less active and socially interactive. Pair-

reared juveniles also submit more readily as subordinates. In a separate cohort, we then measured

neural expression for stress and sex hormone genes, signaling systems known to be

developmentally plastic and involved in translating environmental conditions into biological

responses and regulating adult social behavior. Rearing environment causes striking differences

in neuroendocrine gene co-expression networks. Specifically, expression was tightly integrated

in pair-reared juveniles, but not group-reared or isolated juveniles. Glucocorticoid receptor

subtypes 1a, 1b, and 2, as well as androgen receptor α, drive the significant differences between

treatment groups, which supports a highly conserved role for the stress axis mediating early-life

effects. Together, this research demonstrates the important developmental origins of behavioral

phenotypes and identifies potential behavioral and neuroendocrine mechanisms.

Introduction

Ontogeny has long been recognized as essential to understanding phenotype (Tinbergen,

1963), yet the early-life origins of individual behavioral variation remain understudied.

Development reveals the proximate mechanisms by which genes interact with the environment

during early life to sculpt the 'machinery of behavior' (Stamps, 2003; Tinbergen, 1963). Current

or predicted environmental conditions can trigger developmental plasticity, and the resulting

changes are often long-lasting, or even permanent, and can facilitate locally-adapted (e.g.,

predator resistant, Gilbert, 2001) phenotypes (Kasumovic and Brooks, 2011; Lummaa and

Clutton-Brock, 2002; Piersma and Drent, 2003; Snell-Rood, 2013; Stamps, 2003; Stearns, 1989;

West-Eberhard, 1989). The developmental mechanisms that shape social behavior via underlying

neural regulatory mechanisms should be a particularly important target for natural selection

(Taborsky, 2016) because of the direct consequences of social behavior for fitness and health

(e.g., Bennett et al., 2006; Meyer-Lindenberg and Tost, 2012; Silk, 2007; Solomon-Lane et al.,

2015; Wilson, 1980).

Social stimuli are among the most important attributes of the early-life environment

(Taborsky, 2016). Although maternal (and, to a lesser extent, paternal) interactions have largely

been the focus (e.g., Champagne & Curley, 2005; McClelland, Korosi, Cope, Ivy, & Baram,

2011), the broader early-life social environment is increasingly recognized for its role in

behavioral and neural plasticity (Buist et al., 2013; Creel et al., 2013; Jonsson and Jonsson, 2014;

Kasumovic and Brooks, 2011; Taborsky, 2016; White, 2010). For example, the early presence of

brood care helpers, unrelated adult males, and multiple mothers and litters have long-term effects

on social behavior in the Daffodil cichlid fish *Neolamprologus pulcher* (Arnold and Taborsky,

2010; Taborsky et al., 2012), brown-headed cowbirds (White et al., 2002), and laboratory mice

(Branchi et al., 2013, 2006; D'Andrea et al., 2007), respectively. These features of the social

environment alter the quality and quantity of social experiences and sensory cues perceived,

which together influence neural function and behavior (Taborsky, 2016). Developmental

plasticity may be limited to a single behavior or extend to an entire suite of behaviors (i.e., a

behavioral syndrome), and the effects may be context-specific (Bell, 2007; Snell-Rood, 2013;

Stamps, 2003; Stamps and Groothuis, 2010).

Neuroendocrine signaling is a primary mechanism by which environmental conditions

and experience are translated into physiological responses (Crespi and Denver, 2005; Remage-

Healey and Romero, 2000; Wingfield et al., 1990). Hormones are also important sources of

individual variation in social behavior (e.g., across seasons, sexes, reproductive tactics) and

underlie developmental plasticity relevant to adult behavior. The stress axis, or hypothalamic-

pituitary-adrenal (interrenal in fish; HPA/I) axis, is widely implicated as a highly-conserved

mechanism of early-life effects (Champagne and Curley, 2005; Francis et al., 1999; McClelland

et al., 2011; Taborsky, 2016). In response to an environmental stressor, which includes any

external condition that disrupts or threatens to disrupt homeostasis, the HPA/I axis integrates

relevant internal and external cues and coordinates a response, such as changes in behavior and

physiology. The stress response is initiated by the release of corticotropin-releasing factor (CRF)

from the hypothalamus, which signals to the pituitary to release adrenocorticotropic hormone,

which then signals the adrenal glands to release glucocorticoids (e.g., cortisol in fish) (Denver,

2009; Lowry and Moore, 2006).

Effects of early-life experiences on HPA/I axis function have been demonstrated in every

major vertebrate lineage (e.g., birds: Banerjee, Arterbery, Fergus, & Adkins-Regan, 2012;

mammals: Champagne & Curley, 2005; amphibians: Crespi & Denver, 2005; fish: Jonsson &

Jonsson, 2014). For example, the presence of brood helpers during early-life affects *N. pulcher*

cichlid social behavior via changes in neural expression levels of CRF and glucocorticoid

receptor (GR), as well as receptor ratios (Taborsky et al., 2013). Stress axis mechanisms can also

mediate the effects of the early-life social environment on human health (e.g., Turecki &

Meaney, 2016). Sex steroid hormones (e.g., androgens, estrogens) also play a role mediating the

long-term effects of early-life experiences (Adkins-Regan, 2009; Brown and Spencer, 2013;

Shepard et al., 2009) and regulating social behavior (Goodson, 2005; Newman, 1999). For

example, neural estrogen receptor expression is associated with maternal behavior in mother rats

and offspring (Champagne et al., 2003; Champagne and Meaney, 2007), and socially stressed

pre- and postnatal female guinea pigs have upregulated neural estrogen and androgen receptor

levels, elevated testosterone, and masculinized behavior (Kaiser et al., 2003). Together, these and

other neuroendocrine systems interact to affect behavior.

To investigate the effects of the early-life social environment on behavior and its

neuroendocrine mechanisms, we used the highly social African cichlid *Astatotilapia burtoni*, a

model system in social neuroscience (Fernald and Maruska, 2012; Hofmann, 2003). Adults of

this species form mixed-sex, hierarchical communities with males of dominant or subordinate

status and females. Dominant males are territorial, reproductively active, and colorful. In

comparison, subordinate males shoal with females, are reproductively suppressed, and drab in

coloration. Male status is socially regulated, and individuals regularly transition between status

phenotypes (Fernald and Maruska, 2012; Hofmann, 2003). Adults, and juveniles (Fernald and

Hirata, 1979), express a suite of highly conserved social behaviors, including aggression,

affiliation, courtship, and cooperation (Fernald, 2012; Hofmann, 2003; Weitekamp et al., 2017).

Substantial progress has also been made towards understanding variation in stress and sex steroid

hormone signaling, including in the regulation of social behavior (Chen and Fernald, 2008; Fox

et al., 1997; Greenwood et al., 2003; Munchrath and Hofmann, 2010; O'Connell and Hofmann,

2012a). All GRs (Greenwood et al., 2003), estrogen receptors (ER), and androgen receptors (AR)

(Munchrath and Hofmann, 2010) have been studied in the adult *A. burtoni* brain, and

neuroendocrine function can vary substantially. Subordinate males, for example, have lower

levels of whole brain CRF and GR2 (Chen and Fernald, 2008), higher cortisol, and lower

testosterone than dominants (Fox et al., 1997; O'Connell and Hofmann, 2012a), although these

patterns can vary dynamically (Maguire and Hofmann, in prep.). The transcriptomic response in

the preoptic area (POA) to pharmacological manipulation, such as an ER antagonist, is also

status-specific (O'Connell and Hofmann, 2012a).

Given this rich literature on adult *A. burtoni*, it may seem surprising that the

developmental origins of adult phenotypic variation remain largely unknown. The few studies

that have investigated juveniles demonstrate the importance of early-life. For example, the

development of male behavior and nuptial coloration, as well as reproductive maturation, are

affected by the early-life social environment (Fernald and Hirata, 1979; Fraley and Fernald,

1982). Gestational cues (e.g., maternal social crowding) also have lasting effects on methylation

and transcription of the *gnrh*1 gene in offspring (Alvarado et al., 2015). This result is particularly

interesting given that preoptic GnRH1 neurons, which regulate gonadotropin release from the

pituitary, are socially modulated in adults (Davis and Fernald, 1990; Hofmann and Fernald,

2001). However, studies of the effects of different early-life experiences on other neuroendocrine

139 pathways or behavior are lacking.

140   In the present study, we conducted two experiments to test the hypothesis that the early-

141 life social environment generates variation in juvenile behavior through neuroendocrine gene

142 expression. We manipulated the early-life social environment, and consequently social

143 experience, by rearing juveniles in either social groups or pairs. In the group condition, social

144 experience implies interactions with more social partners, who also vary in size, sex, experience,

145 and patterns of behavior. Interactions in groups can also involve more than two individuals, and

146 it is possible to observe and learn from interactions of group members as a bystander. Although

147 it has not been tested in juveniles, adults are capable of gaining important social information as a

148 bystander (Desjardins et al., 2012, 2010; Grosenick et al., 2007). In the pair condition, juveniles

149 occupy only one social role in a relationship with just one other individual. Similar

150 manipulations of early-life social complexity have been important for behavioral and neural

151 development in other species (reviewed in Taborsky, 2016). We predicted that rearing

152 environment would affect a suite of social behaviors across contexts, including social

153 investigation, dominant, and subordinate behavior. In the brain, we predicted effects on whole

154 brain gene expression of neuroendocrine systems that mediate early-life experiences. Related to

155 the HPA/I axis, we measured glucocorticoid receptor 1a (GR1a), glucocorticoid receptor 1b

156 (GR1b), glucocorticoid receptor 2 (GR2) (nomenclature from Maruska & Fernald, 2010),

157 mineralocorticoid receptor (MR), and CRF. For sex steroid hormone signaling, we quantified

158 androgen receptor α (ARα) and estrogen receptor α (ERα). By investigating these early-life

159 effects in juveniles, we can identify important intermediary steps that inform how developmental

160 plasticity may shape the adult phenotype.

Methods

Animals

Juvenile *A. burtoni* came from a laboratory population descended from a wild-caught stock. The adults that bred the juveniles were housed in naturalistic social groups of males and females. Dominant males court gravid females that then lay eggs in his territory. The female then scoops up the eggs into her mouth, where the male fertilizes them. The mother orally incubates the larvae as they develop for 10-13 days. Under natural (and some laboratory) conditions, juveniles remain close to their mother for the 2-4 weeks following their initial release from her mouth. As they age, juveniles seek shelter in her mouth less and less often. In the first two weeks, juveniles primarily school together, with overt social interactions beginning at 2-3 weeks old (Fernald and Hirata, 1979; Renn et al., 2009). Social behaviors, such as chasing, nipping, territorial displays, emerge in a predictable sequence as juveniles approach reproductive maturity, which can occur as early as 15 weeks, depending on the early-life social conditions (Fernald and Hirata, 1979; Fraley and Fernald, 1982).

We removed juveniles from the mother's mouth 6-12 days post-fertilization. Once sufficiently developed (~day 12, freely swimming with no remaining yolk), juveniles were transferred into experimental rearing environments. Juveniles are all silver (drab) in coloration, and none developed coloration during the study, which would indicate reproductive maturity for males. Sex cannot be determined anatomically until maturation; therefore, the sex ratios of our rearing environments, and the sex of the focal individuals, is unknown. The sex ratio of *A. burtoni* broods is approximately 1:1. All work was done in compliance with the Institutional Animal Care and Use Committee at The University of Texas at Austin.

Experimental rearing conditions (Experiments 1 & 2)

As the first study of this kind in this species, we opted to quantify behavior and gene

expression in separate experiments in order to capture different developmental time points. In

Experiment 1, juveniles for the behavioral assays were reared in social groups of 16 fish (n=12

groups) or in pairs (n=9 pairs) for 58-73 days (average 65.76 ± 0.81; ~8-10 weeks), as long of a

duration that could be used without juveniles reaching reproductive maturity. In Experiment 2,

neural gene expression was measured in a separate cohort of juveniles reared in social groups of

16 fish, pairs, or in isolation for 1 week (groups: n=8; pairs: n=8; isolates: n=8) or 5 weeks

(groups: n=14; pairs: n=10). Here, we aimed to capture early changes in gene expression that

might set individuals along different developmental trajectories. Isolation was included because

we expected it to impact gene expression in this highly social species, not as a social control. We

cannot distinguish between the effects of chronological age from the treatment duration (i.e., 1

vs. 5 weeks) in this study.

For both Experiments, juveniles from multiple clutches of the same age and

developmental stage (day 12-14 fry) were divided among treatment groups. Group-reared fish

were housed in 35 L aquaria with three shards of terracotta pots for a shelter and/or territory.

Pairs and isolated fish were housed in small aquaria (22.9 x 15.2 x 15.2 cm) with one terracotta

pot shard. The volume of water per fish was similar for the group (2.6 L) and paired (2.7 L)

treatments. Juveniles were fed daily with Hikari plankton (Pentair Aquatic Eco-Systems, Cary,

NC). The food was mixed in water, and a transfer pipette was used to deliver a set volume to

each tank. Groups received eight times more food than pairs. Pairs and isolated fish received the

same amount. All juveniles were maintained on a 12:12 light/dark cycle.

Experiment 1: Behavioral assays

Behavior for both members of the pairs (n=18 individuals) and two fish from each group

(n=24 individuals) was analyzed. To choose focal individuals from the groups, we removed all

fish from the aquarium and selected the largest fish. Because size is a strong predicator of social

dominance (Alcazar et al., 2014), this individual was very likely to have dominance experience,

similar to the larger fish in the pair. A smaller fish was then chosen such that the ratio of large-

to-small fish standard length (SL, mm) was approximately equal in the group and a pair from the

same cohort of juveniles (same age). Standard length was recorded for all focal fish. Behavior

was observed in novel, small aquaria (22.9 x 15.2 x 15.2 cm) without covers. For analysis, the

aquaria were divided into 4 zones (Fig 1), delineated with permanent marker. In the middle of

each short side, a circle was drawn (28 mm diameter) to indicate the placement of the

scintillation vial (see below: social cue investigation). An arc 2.54 cm from the edge of that

circle was drawn to form a semicircle. One semicircle was designated the "territory" zone and

had a terracotta pot shard for a shelter and/or territory. The other semicircle was designated the

"investigate" zone. The "close" zone was between the territory zone and halfway along the long

side of the tank. The "far" zone was between the halfway mark and the investigate zone (Fig 1).

Video cameras recorded behavior from above so that all areas of the tank, except under the

terracotta pot shard, were visible. Solomon Coder was used for analysis

(www.solomoncoder.com).

We quantified behavior in four assays, which were always presented in the same

sequence (Fig. 1): an open field test that is commonly used in other species to assess activity and

anxiety (e.g., Cachat et al., 2010; Prut & Belzung, 2003); a social cue investigation as a measure

of social motivation or preference (e.g., Bonuti & Morato, 2018; Moy et al., 2004); and social

interactions within either dominant or subordinate status contexts, which regularly occur in

social communities of *A. burtoni* (Hofmann, 2003). All observations were made by the same

observer who was blind to treatment.

<u>Open field test</u>: The focal fish was transferred to the test aquarium with a hand net and

remained in the tank alone for 30 min. Movement around the tank was observed from minutes 20

to 30. We recorded the number of times a fish crossed into each zone (frequency) and the time

(s) spent in each zone. <u>Social cue investigation</u>: Novel juveniles were collected from a

community tank and placed into scintillation vials (20 mL). The top of the vial was covered with

parafilm with holes to allow water through. A vial containing one cue fish was placed into each

test aquarium (n=16 group-reared, n=13 pair-reared). Cue fish were 0-6.4 mm SL (average 3.37

± 0.27) smaller than their focal fish. An empty vial was used as a control (n=8 group-reared, n=5

pair-reared). The social cues were in the aquarium for 30 min. Movement around the tank

(frequency and time in each zone) was scored from minutes 2 to 12.

<u>Dominance behavior</u>: The scintillation vials were removed from the aquaria and a novel

smaller fish (by 1-6.4 mm SL, average 3.37 ± 0.25) was immediately added to each aquarium,

freely swimming with the focal fish. The pair remained together for 30 minutes, and behavior

was scored from minutes 2 to 12. <u>Subordinate behavior</u>: The small cue fish was removed from

the aquaria and a novel, larger fish (by 2.4-12 mm SL, average 5.74 ± 0.34) was immediately

added to each aquarium, freely swimming with the focal fish. The pair remained together for 30

minutes, and behavior was scored from minutes 2 to 12. In the dominance and subordinate

behavior assays, we analyzed agonistic interactions between the pair. An approach was defined

as one fish swimming directly towards any part of the other fish's body, within 3 body lengths. If

the approached fish responded by moving away, in any direction, the behavior was recorded as a

displacement for the initiator and a submission for the responder. From these measures, we

calculated agonistic efficiency, or the proportion of approaches that led to a displacement

(Solomon-Lane et al., 2014), for focal and cue fish. The difference in agonistic efficiency

between the focal and cue fish was used as a measure of agonistic asymmetry, which

characterizes status relationships (Drews, 1993). We also recorded the frequency of entering and

the time spent in the territory, for the focal fish, cue fish, and both together.

Importantly, after 8-10 weeks in their respective treatment condition, group-reared

juveniles were significantly larger than pair-reared juveniles (see Results). This size difference

influenced the size of the fish selected to be the social stimuli. Specifically, the difference in SL

(in mm) between the focal fish and the social cue (t-test: t=3.38, p=0.0016), as well as the focal

fish and the small cue fish (t=3.476, p=0.0013), was significantly greater for group-reared

juveniles. The size difference (SL) between the focal fish and the large cue fish was significantly

greater for pair-reared juveniles (t=-3.22, p=0.0025). Relative size differences followed the same

pattern as absolute size differences (data not shown).

Experiment 2: Whole brain gene expression

Gene expression for two fish from each group (1 week: n=8; 5 weeks: n=14), both

members of the pairs (1 week: n=8; 5 weeks: n=10), and every isolate (1 week: n=8) was

analyzed. Focal individuals from the group condition were selected haphazardly. Juveniles were

removed from their rearing environments with a hand net and rapidly decapitated. The brains

were dissected immediately, flash frozen on dry ice, and stored at -80° C until processing. Gene

expression was quantified using qPCR and previously validated primers (Supplemental Table 1,

Chen and Fernald, 2008; Greenwood et al., 2003; O'Connell and Hofmann, 2012a) for GR1,

GR2a, GR2b, MR, CRF, ARα, and ERα, as well as control genes 18S and G3PDH. RNA was

extracted using the Maxwell 16 LEV simplyRNA Tissue Kit (Promega, Madison, WI), and the

Promega GoScript Reverse Transcription System (Promega, Madison, WI) was used for reverse

transcription. PowerUp SYBR Green Master Mix (ThermoFisher Scientific, Waltham, MA) was

used for quantitative PCR. All standard kit protocols were followed. Relative gene expression

levels were quantified using $\Delta\Delta$CT analysis, using 18S and G3PDH as reference genes. The

results are largely concordant independent of the reference gene used. Here, we present the

analyses for 18S, as this gene has shown very little expression variation across social phenotypes

in transcriptome studies of *A. burtoni* (O'Connell and Hofmann, 2012a; Renn et al., 2008).

Statistical analyses

All statistical analyses were conducted using R Studio (version 1.0.143). Results were

considered significant at the $p < 0.05$ level, and averages $\pm$ standard error of the mean are included

in the text. The box of the box and whisker plots show the median and the first and third

quartiles. The whiskers extend to the largest and smallest observations within or equal to 1.5

times the interquartile range. Comparisons between group- and pair-reared juveniles were

conducted using t-tests for fish SL, time and frequency in each tank zone, and rates of agonistic

behavior. Mann-Whitney-Wilcoxon tests were used for data that did not meet the assumptions of

parametric statistics. Regression analysis was used to identify significant associations between

SL and frequency and time in a zone and between SL and agonistic behavior. We used a false

discovery rate correction for regressions with focal fish SL (Benjamini and Hochberg, 1995).

Two-way ANOVAs were used to identify significant effects of rearing environment, presence of

the social cue, or an interaction, on the frequency and time spent in each zone of the tank. We

used Principal Components Analysis (PCA) to identify how behaviors clustered across the four

assays and for each assay individually. T-tests were used to compare principal component scores

between group- and pair reared juveniles. Correlation analysis was used to identify significant

associations among principal components (PCs).

Gene expression data did not meet the assumptions of parametric statistics; therefore, the

effects of rearing environment (group, pair, isolated) and treatment duration (1 week, 5 weeks)

were analyzed separately using Kruskal-Wallis tests or one-way ANOVA and Mann-Whitney-

Wilcoxon tests, respectively. To get a tentative estimate for possible interaction effects, we

conducted two-way ANOVAs. These analyses, along with visual inspection of the data, do not

suggest any interactions between rearing environment and treatment duration. We also used PCA

to identify how the expression of candidate genes clustered. T-tests, or Mann-Whitney-Wilcoxon

tests if appropriate, were used to compare group- and pair-reared juveniles and expression

following 1 vs. 5 weeks in rearing environments. Dunn's test was used for *post hoc* analysis of

significant results. Partial correlation networks were calculated using the "ppcor" package in R

and visualized using "qgraph." The nodes of the networks represent the gene. The edges are the

partial correlation coefficient, with thicker edges indicating stronger correlations. Only

significant correlations are shown. Mantel tests were used to test for pairwise differences

between the gene expression networks. A non-significant p-value (> 0.05) indicates that the

partial correlation matrices are not related.

Results

Experiment 1

<u>Open field test and social cue investigation</u>

In the open field test (and subsequent assays), juveniles of both treatment groups moved

readily around the novel environment with minimal acclimation. We present the data for the

frequency of entering each zone. There were no significant effects for the time spent in each

zone ($p > 0.05$). Group-reared juveniles entered the territory (Mann-Whitney-Wilcoxon test:

$W = 299$, $p = 0.034$), close ($W = 293.5$, $p = 0.049$), and investigate zones ($W = 293.5$, $p = 0.049$)

significantly more frequently than pair-reared juveniles. There was no significant difference for

the far zone ($W = 289$, $p = 0.064$).

Next, we used a social cue investigation task to examine whether and how rearing

environment and/or the presence of the social cue affect locomotor activity. Two-way ANOVA

revealed that, following the addition of the social cue, juveniles entered the investigate zone

significantly more frequently than controls ($F_{1,36} = 4.91$, $p = 0.033$). There was no effect of rearing

environment ($F_{1,36} = 1.69$, $p = 0.20$) and no interaction ($F_{1,36} = 0.046$, $p = 0.83$). There was no effect

of rearing environment ($F_{1,36} = 2.68$, $p = 0.11$), social cue ($F_{1,36} = 0.87$, $p = 0.36$), or an interaction

($F_{1,36} = 0.84$, $p = 0.37$) on frequency of entering the far zone. Group-reared juveniles entered the

close zone significantly more than pair-reared juveniles ($F_{1,35} = 4.47$, $p = 0.042$), but there was no

effect of the social cue ($F_{1,35} = 0.11$, $p = 0.74$) and no interaction ($F_{1,35} = 0.44$, $p = 0.52$). There was no

effect of rearing environment ($F_{1,35} = 3.28$, $p = 0.079$), social cue ($F_{1,35} = 0.17$, $p = 0.68$) and no

interaction ($F_{1,35} = 0.83$, $p = 0.37$) on the frequency of entering the territory zone. Linear regression

analyses of the relationship between SL and the frequency of entering zones of the tank are

shown in Table 1.

<u>Dominant and subordinate behavior</u>

Interestingly, rearing environment did not affect rates of focal fish behavior. As the

dominant fish, there were no differences in approaching (W=242.5, p=0.20) or displacing

(W=253, p=0.12) the small cue fish. As the subordinate, there were no differences in

approaching (W=205.5, p=0.85), displacing (W=214.5, p=0.62), or submitting to (W=217.5,

p=0.56) the large cue fish. In the dominance assay, rearing environment did not affect agonistic

efficiency for the focal fish (t=0.83, p=0.41), small cue fish (W=115.5, p= 0.97), or the

difference between the pair (t=1.03, p=0.32). In the subordinate assay, although there was no

effect of rearing environment on agonistic efficiency for the focal fish (W=169.5, p=0.28) or the

large cue fish (W=112.5, p=0.061), the difference in agonistic efficiency was significantly higher

for pair-reared juveniles (p=0.022). See Table 1 for linear regression analyses of SL with

behavior.

<u>Multivariate analysis across assays</u>

In order to gain more insight into this multivariate dataset, we employed PCA to

determine which measures of morphology (i.e., size) and behavior act in concert to explain

different aspects of the variability across individuals. We first conducted a PCA that included

variables from each of the four assays (focal fish SL; frequency of entering each zone in the open

field test and social cue investigation; focal fish social approaches and displacements as a

dominant towards the small cue fish; and focal fish approaches, displacements, and submissions

as a subordinate with the larger cue fish). We found that principal component (PC) 1 accounts

for 43.3% of the variation and differs significantly between group- and pair-reared juveniles

(p=0.029, Fig 2A). As the vector plot in Fig 2B shows, variables from the open field test, social

cue investigation, and dominance behavior assay all load on PC1, along with focal fish SL.

Measures of subordinate behavior do not contribute. There were no significant treatment

differences in higher order PCs except for PC6, which accounted for 5.0% of the variation in the

data and contained significantly higher values for group-reared compared to pair-reared juveniles

(p= 4.082e-05, Fig 2C). Focal fish SL loads most strongly on PC6 (data not shown). This is

entirely in accordance with the finding that after 8-10 weeks in their respective early-life

environments, group-reared juveniles (16.85 ± 0.32 mm SL) were significantly larger than pair-

reared juveniles (13.76 ± 0.40 mm SL) (t=6.00, p=7.25e-07).

To better understand how rearing environment affected behavior within the assays that

contributed to the treatment difference, we conducted PCAs for the open field, social cue

investigation, and dominance behavior assays separately. We expanded these analyses to include

all of the measured variables, for the focal and cue fish. The open field test analysis included

focal fish SL and the frequency of entering and time in each zone of the tank. The social cue

investigation included the same measures, as well as the SL of the cue fish. Finally, the

dominance behavior analysis included SL of the focal fish and small cue fish, approaches and

displacements of both fish, and the frequency of entering and time spent in the territory by either

or both fish. For each analysis, PC1 differed significantly between group- and pair-reared

juveniles: open field (43.4% variation, p=0.04, Fig 3A), social cue investigation (37.2%

variation, W=102, p = 0.0032, Fig 3B), and dominance behavior (29.8% variation, W=128,

p=0.025, Fig 3C). The PC1s were also significantly and linearly correlated with each other (Fig

3D, open field x social cue: r^2=0.46, p=5.33e-07; open field x dominance: r^2=0.33, p= 4.69e-05;

social cue x dominance: r^2=0.46, p= 4.97e-07, Supplemental Figure 1). We found no significant

differences for any higher order PCs in the three analyses. See Supplemental Figure 2 for the

proportion of variation explained by each PC.

For the open field test, all variables loaded on PC1 except time in the territory and

investigate zones. For the social cue investigation, all variables loaded on PC1 except time in the

territory, investigate, and close zones. Finally, for dominance behavior, the strongest loadings for

PC1 include approaches and displacements by the focal and small cue fish, the frequency of the

focal fish entering the territory, the time spent in the territory by the small cue fish, and the SL of

both the focal and small cue fish (Supplemental Figure 3).

Experiment 2

<u>Neural gene expression patterns</u>

Neuroendocrine signaling is a primary mechanism by which early-life experiences are

translated into biological changes. To identify potential mediators of the behavioral effects we

identified, we measured mRNA levels of genes involved in the stress axis and in sex steroid

signaling in the brains of a separate cohort of juveniles. We compared relative expression across

rearing environments (isolation, pairs, groups) and time in rearing environment (1 week, 5

weeks). Overall, there was little significant variation in neural gene expression with regards to

either rearing environment or treatment duration ($p>0.05$, Fig 4, Supplemental Fig 4,

Supplemental Table 2). There was a significant effect of rearing environment on GR1a

expression (Kruskal-Wallis: $\chi^2_2=16.58$, $p=0.00025$). *Post hoc* analysis showed that expression

was significantly higher in groups-reared juveniles than pair-reared ($p=0.0015$) or isolated

($p=0.0015$) juveniles, which did not differ from each other ($p=0.39$). Expression of GR2

($F_{2,43}=4.22$, $p=0.021$) and CRF ($\chi^2_2=6.17$, $p=0.046$) also differed significantly among rearing

environments; however, *post hoc* analyses showed there were no significant pair-wise

differences ($p>0.05$).

Genes function within regulatory networks, rather than in isolation, and they can affect

each other's expression. Similarly, a common upstream regulator may control multiple

functional networks of genes. Because of their known effects on physiology and behavior, these

candidate genes are likely to function in pathways that interact with each other. To quantify how

rearing environment affects gene co-expression, we calculated partial correlation networks (Fig

5). Partial correlations show the associations between gene pairs, independent of other

correlations in the network. Comparing the group and pair networks (Mantel test: $p=0.31$), the

group and isolate networks ($p=0.61$), and the pair and isolate networks ($p=0.12$) revealed that

there was no evidence that any of these networks were similar to any other.

To gain a more holistic understanding of how rearing environment and/or treatment

duration affect variation in neuroendocrine gene expression, we used PCA. PC1 accounts for

69.1% of the variation in the data. While there were no differences in PC1 based on rearing

environment ($W=113$, $p=0.13$), there was a trend for differences based on treatment duration

($W=99$, $p=0.053$; Fig 6A). There were no differences due to rearing environment for any higher

order PCs except for PC4, which accounted for 5.7% of the variation in the data and differed

significantly according to rearing environment ($p=0.011$; Fig 6B). Fig 6C shows how the

different candidate genes load onto PC1 and PC4.

Discussion

In the present study, we demonstrate that juvenile *A. burtoni* behavior and

neuroendocrine gene expression are both sensitive to early-life social effects. By rearing

juveniles in different social environments—either in a social group or as a pair, both of which

allow individuals to interact freely at all times—we altered the quality and quantity of social

experiences and sensory cues perceived and set individuals along different developmental

trajectories. Behaviorally, the early-life environment shifted juveniles in a predictable manner

along a continuum of a novel behavioral syndrome (i.e., correlated behaviors across contexts, see

below) comprised of open field, social cue investigation, and dominance behaviors (Fig 2, Fig 3)

and affected patterns of subordinate behavior, a critically important social role for young

individuals. In the brain, rearing environment caused striking changes in neuroendocrine gene

co-expression patterns, differences driven by the expression of GRs and AR. Together, these

experiments provide an essential step towards understanding how developmental plasticity

generates the individual variation in behavior and neuroendocrine function that has fitness and

health consequences in adulthood (e.g., Champagne, 2010; Turecki and Meaney, 2016). Our

results also contribute to an important and growing literature on the impact of early-life social

environments beyond parental interactions (Champagne and Curley, 2005; Taborsky, 2016),

using a species that, despite its prominence in social neuroscience (Fernald and Maruska, 2012;

Hofmann, 2003), has rarely been studied during development (Alvarado, Lenkov, Williams, &

Fernald, 2015; Fernald & Hirata, 1979; Fraley & Fernald, 1982).

<u>Juvenile behavior forms a syndrome affected by early-life social environment</u>

Using a battery of four behavioral assays to gain a comprehensive understanding of

behavioral phenotype, within and across contexts (Fig 1), we discovered that open field, social

cue investigation, and dominance behavior together formed a behavioral syndrome (Fig 3).

Syndromes are a population-level metric defined as the correlation between rank-order

differences between individuals, across contexts and/or over time (Bell, 2007). The presence of a

syndrome indicates consistency in patterns of individual behavior across contexts and/or over

time (Bell, 2007; Sih et al., 2004a, 2004b). Our data suggest that how individuals move around

in space is relevant to the social role they play. Specifically, juveniles that were more active in

the open field test were more likely to be active in the social cue investigation and more

interactive in the dominance assay. Interestingly, behavior from the subordinate assay does not

contribute to the treatment effect or syndrome, likely because subordinate focal individuals

respond primarily to the dominant fish's behavior. To our knowledge, this is the first behavioral

syndrome to be identified in *A. burtoni* at any developmental stage.

Behavior patterns may coalesce into a syndrome due to shared mechanisms (e.g.,

neuroendocrine regulation), early-life experiences that set individuals along developmentally

plastic trajectories, or correlational selection (Bell, 2007; Ketterson and Nolan, Jr., 1999; Stamps,

2003). We found that the behavior of all juveniles was described by the same syndrome,

indicating that how the behaviors are related across experimental contexts (i.e., assays) was

maintained independently of the early-life social environment. Whether an individual was reared

in a group or pair then dictates where along the continuum of the syndrome they fall (Fig 3D).

Pair-reared juveniles appear restricted to one end, whereas group-reared juveniles are represented

along the full range of behavioral variation. That there are group-reared juveniles that

behaviorally resemble the pair-reared individuals suggests there may be social environments

within a group (Saltz et al., 2016) that share key elements with the paired experience.

Conversely, the range of possible social roles seems much more restricted in the paired

treatment. To identify the causal behavioral and/or sensory cues, it will be necessary to conduct

detailed observations of individuals within the rearing environments (Taborsky, 2016). Based on

our pilot observations, we hypothesize that the complexity of interactions and/or abundance of

social sensory cues in groups cause these treatment differences (Taborsky, 2016, e.g., Arnold &

Taborsky, 2010).

Activity and social interaction are common components of syndromes in other species,

along with bold-shy and proactive-reactive behaviors (Bell, 2007; Conrad et al., 2011; Groothuis

and Carere, 2005; Koolhaas et al., 1999; Sih et al., 2004b; Verbeek et al., 1994). For example,

large juvenile brown trout are more active and aggressive (Näslund and Johnsson, 2016), similar

to our results. Activity-aggression syndromes are also found in a number of other fish species

(reviewed in Conrad et al., 2011). For *A. burtoni* juveniles, locomotor activity and social

interaction may be causally related. First, active individuals may encounter conspecifics more

frequently and, as a result, initiate more interactions. Second, juvenile social interactions appear

to be prosocial in that they increase the likelihood of future proximity and interaction. In the

dominance behavior assay, approaches and displacements for both the focal and subordinate cue

fish load in the same direction on PC1. Correlation analysis (data not shown) confirms that, as

one member of the pair initiates social interactions, the other member also initiates, potentially

leading to more activity. This may be beneficial by increasing shoaling and reducing the risk of

predation. Interestingly, adult dominance behavior does not lead to a prosocial response in

subordinates, suggesting that although social behavior appears similar across life history stages

(Fernald and Hirata, 1979; Fraley and Fernald, 1982), there are important differences.

<u>Size plays a secondary role in determining juvenile behavioral phenotype</u>

Size is central to understanding juvenile activity, social interactions, and the effects of the

early-life social environment. Group-reared juveniles were larger than those reared in pairs, and

SL was positively associated with activity in the open field and social cue investigation assays,

approaching and displacing as a dominant fish, as well as entering the territory zone, alone or

with the cue fish, in the dominance behavior assay. The importance of size in juveniles is

consistent with the research showing that adult *A. burtoni* are highly sensitive to size during

social interactions (Alcazar et al., 2014; Weitekamp & Hofmann, 2017). Growth is also socially

regulated in both juveniles and adults (Fraley and Fernald, 1982; Hofmann et al., 1999; this

study). However, the effect of the early social environment is much larger and more complex

than size alone. First, the PCA of behavior from all four assays shows that focal fish SL

contributes only moderately to the significant treatment difference for PC1 (Fig 2B), as many

other variables load much more strongly on PC1 (i.e., open field, social cue investigation, and

dominance behaviors) (see also: Supplemental Fig 3). Second, SL is the strongest contributing

variable for PC6, which represents the significant size difference between group-reared and pair-

reared individuals discussed above, yet contributes only 5% to the overall variation in the data.

Finally, the group-reared juveniles that fall within the range of pair-reared juveniles along the

continuum of the behavioral syndrome (i.e., high PCA scores, Fig 3) are not the smallest

individuals. While size may be secondary for understanding early-life effects in juveniles, it

remains to be tested how individual behavior changes over time in relation to both size and

developmental stage, which can be decoupled from chronological age in fish (Jonsson and

Jonsson, 2014).

<u>Early-life social experience affects social dynamics when focal juveniles are subordinate</u>

Developmental plasticity can shift behavior in ways that ultimately benefit fitness (Smith

and Blumstein, 2008), in part because social behavior has direct consequences for reproductive

success (Wilson, 1980, e.g., Henry et al., 2013; Robbins et al., 2007; Young et al., 2006). A

majority (64%) of studies show that experimentally increasing the frequency, diversity, or

complexity of early-life social experiences enhances social skills or competence (Taborsky,

2016). For example, juvenile *N. pulcher* cichlids reared with brood helpers demonstrated more

context-appropriate behavior when establishing status, integrating into novel groups, and

competing for a resource (Arnold and Taborsky, 2010; Fischer et al., 2015; Taborsky et al.,

2013, 2012). We have no evidence of an advantage for group-reared juveniles; however,

juveniles may fill the subordinate role differently. While nearly all focal fish successfully

established themselves as subordinate (88%) in the assay, and there were no treatment

differences in approaches or displacements, there was a significantly larger asymmetry in

agonistic efficiency for pair-reared juveniles. There was also a trend for pair-reared juveniles to

submit more readily (measured as large fish agonistic efficiency). Status relationships are

defined by asymmetrical agonistic displays (Drews, 1993); therefore, pair-reared juveniles may

behave more submissively. The subordinate role is a critically important one for juveniles

because all will enter adult communities as subordinates. It will be necessary to measure

behavior and reproductive success of these juveniles once they are adults in order to determine

whether these phenotypes persist or if one is more successful than another (Pradhan, Solomon-

Lane, & Grober, 2015).

<u>Early-life social environment affects neuroendocrine gene co-expression networks</u>

We have shown that early-life environments can determine where individuals will end up

along the axis of the newly discovered behavioral syndrome, which raises questions about the

underlying mechanisms (e.g., pleiotropic genes and/or neuroendocrine regulation). The

behavioral effects we detect as a result of the early-life social environment suggest important

variation in the underlying neural regulatory mechanisms. Neuroendocrine stress and sex

hormone signaling are likely sites of developmental plasticity in *A. burtoni* because they are

sensitive to early-life effects (Champagne & Curley, 2005; Shepard et al., 2009), translate

environmental conditions and experiences into biological responses (Crespi & Denver, 2005;

Wingfield et al., 1990), and regulate behavior (Adkins-Regan, 2009; Solomon-Lane, Crespi, &

Grober, 2013). We focused on steroid hormone nuclear receptors specifically because they

regulate the transcription of target genes with a diversity of physiological and behavioral roles

(Rochette-Egly, 2005). Overall, gene expression was highly variable, especially among group-

reared juveniles, possibly reflecting the considerable variability in experiences within groups.

With the exception of GR1a, we found no significant differences when comparing the expression

of single genes across rearing environments or treatment durations (Fig 4, Supplemental Fig 4),

suggesting individual variation is strongly influenced by other factors, which we did not measure

here (e.g., social status, body size, and sex). For example, social experience and status can both

affect gene expression (Li et al., 2014). In adult male *A. burtoni*, the expression of ARα, MR,

GR1a, and GR2 in the POA of the hypothalamus is higher in dominants, whereas GR1b is higher

in subordinates (Korzan et al., 2014). Our behavior data suggest that an individual's position

along the behavioral syndrome, as well as its social status, may be critical for understanding gene

expression variation.

The expression of other genes can also contribute to expression variation.

Neuroendocrine systems are dynamic and interact on multiple biological levels, including within

gene regulatory networks (e.g., Huffman et al., 2012; Korzan, Fernald, & Grone, 2014;

O'Connell & Hofmann, 2012). Based on their co-localization in the POA of *A. burtoni* (Korzan

et al., 2014), co-localization and correlation in other species (e.g., Meyer & Korz, 2013), and

overlapping physiological effects (Crespi & Denver, 2005; Wingfield et al., 1990), the

neuroendocrine pathways represented by our candidate genes are likely to functionally interact.

We identified striking differences in co-expression networks among juveniles reared in different

environments. Expression was highly correlated in pair-reared juveniles (Fig 5A), such that

every candidate gene was significantly correlated with at least two others. At the center of the

network, ARα shares five significant connections. The two sex steroid hormone genes (ARα,

ERα) are also integrated with the stress axis genes, which form distinct smaller networks: CRF-

GR1a-GR1b and GR2-MR. In contrast, group-reared juveniles have only one significant partial

correlation between ERα and GR1b, a connection that is not present in the pair-reared network

(Fig 5B). There are no significant partial correlations for isolated juveniles, suggesting that the

neuroendocrine regulatory network is dysregulated, possibly due to isolation acting as a stressor

(Galhardo and Oliveira, 2014). These network differences might underlie the behavioral

differences we identified in the behavioral syndrome, subordinate behavior, or more broadly

related to stress response (see below). The differential co-regulation could also serve to make

behavior more similar in the face of other neural differences caused by rearing environment, as is

the case for some neural sex differences and behavior (De Vries, 2004). These hypotheses can be

tested directly using central pharmacological manipulation.

Complex regulation of neural stress and sex steroid signaling by the early-life social environment

From the gene co-expression networks alone, it is challenging to determine whether

specific genes drive the significant differences across treatment groups. This is where

decomposing the total variance into principal components offers a powerful approach that

allowed us to show that the HPI axis plays a central, and likely highly-conserved (Crespi and

Denver, 2005), role in responding to the early-life social environment in juvenile *A. burtoni*.

First, PCA revealed that the neural expression of ERα, MR, and CRF strongly loaded on PC1,

which explains 69.1% of the variance (Fig 6C). The trend for differences in PC1 based on

treatment duration (Fig 6A) suggests that we cannot rule out that neuroendocrine gene

expression patterns change in important ways over the course of development. Because all

juveniles entered treatment at the same age and developmental stage, future work is needed to

distinguish between the effects of treatment duration and age (if any), and to identify possible

critical periods for early-life effects.

Second, scores for PC4 (5.7%) were significantly different between group- and pair-

reared juveniles (Fig 6B). All of the GRs, as well as AR, load on PC4 and contribute to the

treatment effect (Fig 6C). Many teleosts, including *A. burtoni*, have three glucocorticoid

receptors: MR, GR1, and GR2. Receptor 1 has splice variants 1a and 1b, which differ by a nine

amino acid insertion in the DNA-binding domain of 1b that reduces transcriptional response

(Greenwood et al., 2003; Korzan et al., 2014). Consistent with the distinct roles for the different

receptors and splice variants (Greenwood et al., 2003), GR1a and GR2 load in the opposite

direction from GR1b and ARα (Fig 6C), suggesting their expression may be antagonistically

regulated (e.g., Fig 5A). Changes in HPA/I axis function typically manifest as altered baseline

levels of circulating glucocorticoids, a higher or lower glucocorticoid 'peak' in response to an

acute stressor, and/or altered efficiency of the negative feedback loop that returns the system to

baseline. Negative feedback, in particular, is regulated by neural GR expression and can be

affected by early-life experience (Champagne and Curley, 2005; Francis et al., 1999). This

suggests either group- or pair-reared juveniles, or both, may have altered negative feedback

mechanisms via differential GR expression. In particular, the hippocampus and amygdala (and

their non-mammalian homologs), brain regions important in spatial cognition and emotional

processing, respectively, are central to negative feedback (Denver, 2009).

Finally, none of the endpoints we measured loaded on PC2 (11.3%) or PC3 (7.17%),

suggesting that there are likely other molecular pathways involved that we did not assay, which

is to be expected for a complex phenotype such as social behavior. A genome-scale analysis of

gene expression using RNA-seq can provide novel candidates here.

<u>Integrating the effects of early-life social environments on behavior and brain</u>

Our work demonstrates that early-life social environments shape behavioral phenotype

and neuroendocrine gene expression in powerful ways for *A. burtoni* juveniles. In this study, we

quantified behavior and gene expression in separate experiments in order to focus on different

developmental time points. However, understanding the full scope and consequences of early-

life effects requires measuring brain and behavior in the same individuals, throughout

development and into adulthood. This work can begin to address the fact that across species,

remarkably little is known about the mechanisms that shape the ontogeny of behavior (Taborsky,

2016). Our results suggest that brain regions that express GRs and ARα (Greenwood et al., 2003;

Korzan et al., 2014; Munchrath and Hofmann, 2010), along with brain regions of the social

decision-making network (SDMN) that together regulate social behavior (O'Connell and

Hofmann, 2012b), are likely to be sensitive to early-life effects and could cause the observed

changes in behavior. Interestingly, the POA—a critical node in the SDMN (O'Connell and

Hofmann, 2012b)—contains GR1a, GR1b, GR2, and ARα in adult *A. burtoni* (Korzan et al.,

2014; Munchrath and Hofmann, 2010). Additional SDMN nodes, such as the hippocampus and

amygdala are also likely sites of overlap. Interactions between the HPI axis and androgen

signaling, including in the POA, could be a mechanism for the social regulation of development

(Fraley and Fernald, 1982; Korzan et al., 2014; Solomon-Lane et al., 2013; Wada, 2008).

Another hypothesis is that the behavior patterns sensitive to early-life effects and HPI axis

function will together form a specific kind of syndrome called a coping style, which ranges from

proactive to reactive copers. Proactive copers tend to be more active, aggressive, and less

responsive to stress (i.e., lower baseline glucocorticoid levels, faster negative feedback) than

reactive copers (Koolhaas et al., 1999). Overall, this research can uncover the neuroendocrine

mechanisms by which early-life social experience gives rise to individual variation in adults,

which is critical to understanding subsequent disparities in fitness and health.

Acknowledgements

We thank Savannah Clapp, Pamela Del Valle, and Najah Hussain for assistance with data

collection and fish maintenance and care. We thank Becca Young for helpful comments on

earlier versions of this manuscript and members of the Hofmann Lab for discussion and

feedback.

Funding

This work was supported by NSF grant IOS-1354942 to HAH and the BEACON Center for the

Study of Evolution in Action awards #947 (2016) and #1081 (2017) to TKSL.

Declarations of interest: None.
664

665   **References**

666   Adkins-Regan, E., 2009. Neuroendocrinology of Social Behavior. ILAR J. 50, 5–14.
667   https://doi.org/10.1093/ilar.50.1.5
668   Alcazar, R.M., Hilliard, A.T., Becker, L., Bernaba, M., Fernald, R.D., 2014. Brains over brawn:
669   experience overcomes a size disadvantage in fish social hierarchies. J. Exp. Biol. 217,
670   1462–1468. https://doi.org/10.1242/jeb.097527
671   Alvarado, S.G., Lenkov, K., Williams, B., Fernald, R.D., 2015. Social crowding during
672   development causes changes in GnRH1 DNA methylation. PLoS One 10, 1–13.
673   https://doi.org/10.1371/journal.pone.0142043
674   Arnold, C., Taborsky, B., 2010. Social experience in early ontogeny has lasting effects on social
675   skills in cooperatively breeding cichlids. Anim. Behav. 79, 621–630.
676   https://doi.org/10.1016/j.anbehav.2009.12.008
677   Banerjee, S.B., Arterbery, A.S., Fergus, D.J., Adkins-Regan, E., 2012. Deprivation of maternal
678   care has long-lasting consequences for the hypothalamic-pituitary-adrenal axis of zebra
679   finches. Proc. R. Soc. B Biol. Sci. 279, 759–766. https://doi.org/10.1098/rspb.2011.1265
680   Bell, A.M., 2007. Future directions in behavioural syndromes research. Proc. R. Soc. B Biol. Sci.
681   274, 755–761. https://doi.org/10.1098/rspb.2006.0199
682   Benjamini, Y., Hochberg, Y., 1995. Controlling the False Discovery Rate: A Practical and
683   Powerful Approach to Multiple Testing. J. R. Stat. Soc. Ser. B.
684   https://doi.org/10.2307/2346101
685   Bennett, D. a, Schneider, J. a, Tang, Y., Arnold, S.E., Wilson, R.S., 2006. The effect of social
686   networks on the relation between Alzheimer's disease pathology and level of cognitive
687   function in old people: a longitudinal cohort study. Lancet. Neurol. 5, 406–12.
688   https://doi.org/10.1016/S1474-4422(06)70417-3
689   Bonuti, R., Morato, S., 2018. Proximity as a predictor of social behavior in rats. J. Neurosci.
690   Methods 293, 37–44. https://doi.org/10.1016/j.jneumeth.2017.08.027
691   Branchi, I., D'Andrea, I., Fiore, M., Di Fausto, V., Aloe, L., Alleva, E., 2006. Early Social
692   Enrichment Shapes Social Behavior and Nerve Growth Factor and Brain-Derived
693   Neurotrophic Factor Levels in the Adult Mouse Brain. Biol. Psychiatry 60, 690–696.
694   https://doi.org/10.1016/j.biopsych.2006.01.005
695   Branchi, I., Santarelli, S., D'Andrea, I., Alleva, E., 2013. Not all stressors are equal: Early social
696   enrichment favors resilience to social but not physical stress in male mice. Horm. Behav.
697   63, 503–509. https://doi.org/10.1016/j.yhbeh.2013.01.003
698   Brown, G.R., Spencer, K.A., 2013. Steroid hormones, stress and the adolescent brain: A
699   comparative perspective. Neuroscience 249, 115–128.
700   https://doi.org/10.1016/j.neuroscience.2012.12.016
701   Buist, K.L., Deković, M., Prinzie, P., 2013. Sibling relationship quality and psychopathology of
702   children and adolescents: A meta-analysis. Clin. Psychol. Rev. 33, 97–106.
703   https://doi.org/10.1016/j.cpr.2012.10.007
704   Cachat, J., Stewart, A., Grossman, L., Gaikwad, S., Kadri, F., Chung, K.M., Wu, N., Wong, K.,
705   Roy, S., Suciu, C., Goodspeed, J., Elegante, M., Bartels, B., Elkhayat, S., Tien, D., Tan, J.,
706   Denmark, A., Gilder, T., Kyzar, E., Dileo, J., Frank, K., Chang, K., Utterback, E., Hart, P.,
707   Kalueff, A. V., 2010. Measuring behavioral and endocrine responses to novelty stress in
708   adult zebrafish. Nat. Protoc. 5, 1786–1799. https://doi.org/10.1038/nprot.2010.140
709   Champagne, F.A., 2010. Early adversity and developmental Outcomes: Interaction between

Genetics, Epigenetics, and social experiences across the Life Span. Perspect. Psychol. Sci. 5, 564–574. https://doi.org/10.1177/1745691610383494

Champagne, F.A., Curley, J.P., 2005. How social experiences influence the brain. Curr. Opin. Neurobiol. 15, 704–709. https://doi.org/10.1016/j.conb.2005.10.001

Champagne, F.A., Meaney, M.J., 2007. Transgenerational effects of social environment on variations in maternal care and behavioral response to novelty. Behav. Neurosci. 121, 1353–1363. https://doi.org/10.1037/0735-7044.121.6.1353

Champagne, F.A., Weaver, I.C.G., Diorio, J., Sharma, S., Meaney, M.J., 2003. Natural Variations in Maternal Care Are Associated with Estrogen Receptor α Expression and Estrogen Sensitivity in the Medial Preoptic Area. Endocrinology 144, 4720–4724. https://doi.org/10.1210/en.2003-0564

Chen, C.-C., Fernald, R.D., 2008. Sequences, expression patterns and regulation of the corticotropin-releasing factor system in a teleost. Gen. Comp. Endocrinol. 157, 148–155. https://doi.org/10.1016/j.ygcen.2008.04.003

Conrad, J.L., Weinersmith, K.L., Brodin, T., Saltz, J.B., Sih, A., 2011. Behavioural syndromes in fishes: A review with implications for ecology and fisheries management. J. Fish Biol. 78, 395–435. https://doi.org/10.1111/j.1095-8649.2010.02874.x

Creel, S.R., Dantzer, B., Goymann, W., Rubenstein, D.R., 2013. The ecology of stress: effects of the social environment. Funct. Ecol. 27, 66–80. https://doi.org/10.1111/j.1365-2435.2012.02029.x

Crespi, E.J., Denver, R.J., 2005. Ancient origins of human developmental plasticity. Am. J. Hum. Biol. 17, 44–54. https://doi.org/10.1002/ajhb.20098

D'Andrea, I., Alleva, E., Branchi, I., 2007. Communal nesting, an early social enrichment, affects social competences but not learning and memory abilities at adulthood. Behav. Brain Res. 183, 60–66. https://doi.org/10.1016/j.bbr.2007.05.029

Davis, M.R., Fernald, R.D., 1990. Social control of neuronal soma size. J. Neurobiol. 21, 1180–1188. https://doi.org/10.1002/neu.480210804

De Vries, G.J., 2004. Minireview: Sex Differences in Adult and Developing Brains: Compensation, Compensation, Compensation. Endocrinology 145, 1063–1068. https://doi.org/10.1210/en.2003-1504

Denver, R.J., 2009. Structural and functional evolution of vertebrate neuroendocrine stress systems. Ann. N. Y. Acad. Sci. 1163, 1–16. https://doi.org/10.1111/j.1749-6632.2009.04433.x

Desjardins, J.K., Hofmann, H.A., Fernald, R.D., 2012. Social Context Influences Aggressive and Courtship Behavior in a Cichlid Fish. PLoS One 7, e32781. https://doi.org/10.1371/journal.pone.0032781

Desjardins, J.K., Klausner, J.Q., Fernald, R.D., 2010. Female genomic response to mate information. Proc. Natl. Acad. Sci. U. S. A. 107, 21176–21180. https://doi.org/10.1073/pnas.1010442107

Drews, C., 1993. The concept and definition of dominance in animal behaviour. Behaviour 125, 283–313. https://doi.org/10.1163/156853993X00290

Fernald, R.D., 2012. Social Control of the Brain. Annu. Rev. Neurosci. 35, 133–151. https://doi.org/10.1146/annurev-neuro-062111-150520

Fernald, R.D., Hirata, N.R., 1979. The Ontogeny of Social behavior and body coloration in the african cichlid fish haplochromis burtoni. Zeitschrif für Tierpsychologie. https://doi.org/10.1111/j.1439-0310.1979.tb01025.x

Fernald, R.D., Maruska, K.P., 2012. Social information changes the brain. Proc. Natl. Acad. Sci.
109, 17194–17199. https://doi.org/10.1073/pnas.1202552109

Fischer, S., Bessert-Nettelbeck, M., Kotrschal, A., Taborsky, B., 2015. Rearing-Group Size
Determines Social Competence and Brain Structure in a Cooperatively Breeding Cichlid.
Am. Nat. 186, 123–140. https://doi.org/10.1086/681636

Fox, H.E., White, S. a, Kao, M.H., Fernald, R.D., 1997. Stress and dominance in a social fish. J.
Neurosci. 17, 6463–6469.

Fraley, N.B., Fernald, R.D., 1982. Social control of developmental rate in the African cichlid,
Haplochromis burtoni. Z.Tierpsychol. 60, 66–82. https://doi.org/10.1111/j.1439-
0310.1982.tb01077.x

Francis, D.D., Caldji, C., Champagne, F.A., Plotsky, P.M., Meaney, M.J., 1999. The role of
corticotropin-releasing factor–norepinephrine systems in mediating the effects of early
experience on the development of behavioral and endocrine responses to stress. Biol.
Psychiatry 46, 1153–1166. https://doi.org/10.1016/S0006-3223(99)00237-1

Galhardo, L., Oliveira, R.F., 2014. The effects of social isolation on steroid hormone levels are
modulated by previous social status and context in a cichlid fish. Horm. Behav. 65, 1–5.
https://doi.org/10.1016/j.yhbeh.2013.10.010

Gilbert, S.F., 2001. Ecological Developmental Biology: Developmental Biology Meets the Real
World. Dev. Biol. 233, 1–12. https://doi.org/10.1006/dbio.2001.0210

Goodson, J.L., 2005. The vertebrate social behavior network: Evolutionary themes and
variations. Horm. Behav. 48, 11–22. https://doi.org/10.1016/j.yhbeh.2005.02.003

Greenwood, A.K., Butler, P.C., White, R.B., Demarco, U., Pearce, D., Fernald, R.D., 2003.
Multiple corticosteroid receptors in a teleost fish: Distinct sequences, expression patterns,
and transcriptional activities. Endocrinology 144, 4226–4236.
https://doi.org/10.1210/en.2003-0566

Groothuis, T.G.G., Carere, C., 2005. Avian personalities: Characterization and epigenesis.
Neurosci. Biobehav. Rev. 29, 137–150. https://doi.org/10.1016/j.neubiorev.2004.06.010

Grosenick, L., Clement, T.S., Fernald, R.D., 2007. Fish can infer social rank by observation
alone. Nature 445, 429–432. https://doi.org/10.1038/nature05646

Henry, M.D., Hankerson, S.J., Siani, J.M., French, J.A., Dietz, J.M., 2013. High rates of
pregnancy loss by subordinates leads to high reproductive skew in wild golden lion
tamarins (Leontopithecus rosalia). Horm. Behav. 63, 675–683.
https://doi.org/10.1016/j.yhbeh.2013.02.009

Hofmann, H.A., 2003. Functional genomics of neural and behavioral plasticity. J. Neurobiol. 54,
272–282. https://doi.org/10.1002/neu.10172

Hofmann, H.A., Benson, M.E., Fernald, R.D., 1999. Social status regulates growth rate:
consequences for life-history strategies. Proc. Natl. Acad. Sci. U. S. A. 96, 14171–6.
https://doi.org/10.1073/pnas.96.24.14171

Hofmann, H.A., Fernald, R.D., 2001. What cichlids tell us about the social regulation of brain
and behavior. J. Aquaric. Aquat. Sci. 9, 17–31.

Huffman, L.S., Mitchell, M.M., O'Connell, L.A., Hofmann, H.A., 2012. Rising StARs:
Behavioral, hormonal, and molecular responses to social challenge and opportunity. Horm.
Behav. 61, 631–641. https://doi.org/10.1016/j.yhbeh.2012.02.016

Jonsson, B., Jonsson, N., 2014. Early environment influences later performance in fishes. J. Fish
Biol. 85, 151–188. https://doi.org/10.1111/jfb.12432

Kaiser, S., Kruijver, F.P.M., Swaab, D.F., Sachser, N., 2003. Early social stress in female guinea

pigs induces a masculinization of adult behavior and corresponding changes in brain and
neuroendocrine function. Behav. Brain Res. 144, 199–210. https://doi.org/10.1016/S0166-
4328(03)00077-9
Kasumovic, M.M., Brooks, R.C., 2011. It's All Who You Know: The Evolution Of Socially
Cued Anticipatory Plasticity As A Mating Strategy. Q. Rev. Biol. 86, 181–197.
https://doi.org/10.1086/661119
Ketterson, E.D., Nolan, Jr., V., 1999. Adaptation, Exaptation, and Constraint: A Hormonal
Perspective. Am. Nat. 154, S4–S25. https://doi.org/10.1086/303280
Koolhaas, J.M., Korte, S.M., De Boer, S.F., Van Der Vegt, B.J., Van Reenen, C.G., Hopster, H.,
De Jong, I.C., Ruis, M. a W., Blokhuis, H.J., 1999. Coping styles in animals: Current status
in behavior and stress- physiology. Neurosci. Biobehav. Rev. 23, 925–935.
https://doi.org/10.1016/S0149-7634(99)00026-3
Korzan, W.J., Fernald, R.D., Grone, B.P., 2014. Social regulation of cortisol receptor gene
expression. J. Exp. Biol. 3221–3228. https://doi.org/10.1242/jeb.104430
Li, C.-Y., Earley, R.L., Huang, S.-P., Hsu, Y., 2014. Fighting experience alters brain androgen
receptor expression dependent on testosterone status. Proc. R. Soc. B Biol. Sci. 281,
20141532–20141532. https://doi.org/10.1098/rspb.2014.1532
Lowry, C. a., Moore, F.L., 2006. Regulation of behavioral responses by corticotropin-releasing
factor. Gen. Comp. Endocrinol. 146, 19–27. https://doi.org/10.1016/j.ygcen.2005.12.006
Lummaa, V., Clutton-Brock, T., 2002. Early development, survival and reproduction in humans.
Trends Ecol. Evol. 17, 141–147. https://doi.org/10.1016/s0169-5347(01)02414-4
Maruska, K.P., Fernald, R.D., 2010. Steroid receptor expression in the fish inner ear varies with
sex, social status, and reproductive state. BMC Neurosci. 11, 58.
https://doi.org/10.1186/1471-2202-11-58
McClelland, S., Korosi, A., Cope, J., Ivy, A., Baram, T.Z., 2011. Emerging roles of epigenetic
mechanisms in the enduring effects of early-life stress and experience on learning and
memory. Neurobiol. Learn. Mem. 96, 79–88. https://doi.org/10.1016/j.nlm.2011.02.008
Meyer-Lindenberg, A., Tost, H., 2012. Neural mechanisms of social risk for psychiatric
disorders. Nat. Neurosci. 15, 663–668. https://doi.org/10.1038/nn.3083
Meyer, K., Korz, V., 2013. Estrogen receptor α functions in the regulation of motivation and
spatial cognition in young male rats. PLoS One 8.
https://doi.org/10.1371/journal.pone.0079303
Moy, S.S., Nadler, J.J., Perez, A., Barbaro, R.P., Johns, J.M., Magnuson, T.R., Piven, J.,
Crawley, J.N., 2004. Sociability and preference for social novelty in five inbred strains: an
approach to assess autistic-like behavior in mice. Genes, Brain Behav. 3, 287–302.
https://doi.org/10.1111/j.1601-1848.2004.00076.x
Munchrath, L.A., Hofmann, H.A., 2010. Distribution of sex steroid hormone receptors in the
brain of an african cichlid fish, astatotilapia burtoni. J. Comp. Neurol. 518, 3302–3326.
https://doi.org/10.1002/cne.22401
Näslund, J., Johnsson, J.I., 2016. State-dependent behavior and alternative behavioral strategies
in brown trout (Salmo trutta L.) fry. Behav. Ecol. Sociobiol. 70, 2111–2125.
https://doi.org/10.1007/s00265-016-2215-y
Newman, S., 1999. The medial extended amygdala in male reproductive behavior. Ann NY Acad
Sci 242–257.
O'Connell, L.A., Hofmann, H.A., 2012a. Social Status Predicts How Sex Steroid Receptors
Regulate Complex Behavior across Levels of Biological Organization. Endocrinology 153,

1341–1351. https://doi.org/10.1210/en.2011-1663

O'Connell, L.A., Hofmann, H.A., 2012b. Evolution of a Vertebrate Social Decision-Making
Network. Science (80-.). 336, 1154–1157. https://doi.org/10.1126/science.1218889

Piersma, T., Drent, J., 2003. Phenotypic flexibility and the evolution of organismal design.
Trends Ecol. Evol. 18, 228–233. https://doi.org/10.1016/S0169-5347(03)00036-3

Pradhan, D.S., Solomon-Lane, T.K., Grober, M.S., 2015. Contextual modulation of social and
endocrine correlates of fitness: insights from the life history of a sex changing fish. Front.
Neurosci. 9, 1–21. https://doi.org/10.3389/fnins.2015.00008

Prut, L., Belzung, C., 2003. The open field as a paradigm to measure the effects of drugs on
anxiety-like behaviors: A review. Eur. J. Pharmacol. 463, 3–33.
https://doi.org/10.1016/S0014-2999(03)01272-X

Remage-Healey, L., Romero, L.M., 2000. Daily and seasonal variation in response to stress in
captive starlings (sturnus vulgaris): Glucose. Gen. Comp. Endocrinol. 119, 60–68.
https://doi.org/10.1006/gcen.2000.7492

Renn, S., Carleton, J.B., Magee, H., Nguyen, M.L.T., Tanner, A.C.W., 2009. Maternal care and
altered social phenotype in a recently collected stock of Astatotilapia burtoni cichlid fish.
Integr. Comp. Biol. 49, 660–673. https://doi.org/10.1093/icb/icp085

Renn, S.C.P., Aubin-Horth, N., Hofmann, H.A., 2008. Fish and chips: functional genomics of
social plasticity in an African cichlid fish. J. Exp. Biol. 211, 3041–3056.
https://doi.org/10.1242/jeb.018242

Robbins, M.M., Robbins, A.M., Gerald-Steklis, N., Steklis, H.D., 2007. Socioecological
influences on the reproductive success of female mountain gorillas (Gorilla beringei
beringei). Behav. Ecol. Sociobiol. 61, 919–931. https://doi.org/10.1007/s00265-006-0321-y

Rochette-Egly, C., 2005. Dynamic combinatorial networks in nuclear receptor-mediated
transcription. J. Biol. Chem. 280, 32565–32568. https://doi.org/10.1074/jbc.R500008200

Saltz, J.B., Geiger, A.P., Anderson, R., Johnson, B., Marren, R., 2016. What, if anything, is a
social niche? Evol. Ecol. 30, 349–364. https://doi.org/10.1007/s10682-015-9792-5

Shepard, K.N., Michopoulos, V., Toufexis, D.J., Wilson, M.E., 2009. Genetic, epigenetic and
environmental impact on sex differences in social behavior. Physiol. Behav. 97, 157–170.
https://doi.org/10.1016/j.physbeh.2009.02.016

Sih, A., Bell, A.M., Johnson, J.C., 2004a. Behavioral syndromes: an ecological and evolutionary
overview. Trends Ecol. Evol. 19, 372–378. https://doi.org/10.1016/j.tree.2004.04.009

Sih, A., Bell, A.M., Ziemba, R.E., 2004b. Behavioural syndromes: An integrative overview. Q.
Rev. Biol. 80, 1–1. https://doi.org/10.1086/516403

Silk, J.B., 2007. Social Components of Fitness in Primate Groups. Science (80-.). 317, 1347–
1351. https://doi.org/10.1126/science.1140734

Smith, B.R., Blumstein, D.T., 2008. Fitness consequences of personality: A meta-analysis.
Behav. Ecol. 19, 448–455. https://doi.org/10.1093/beheco/arm144

Snell-Rood, E.C., 2013. An overview of the evolutionary causes and consequences of
behavioural plasticity. Anim. Behav. 85, 1004–1011.
https://doi.org/10.1016/j.anbehav.2012.12.031

Solomon-Lane, T.K., Crespi, E.J., Grober, M.S., 2013. Stress and serial adult metamorphosis :
multiple roles for the stress axis in socially regulated sex change. Front. Neuroendocrinol. 7,
1–12. https://doi.org/10.3389/fnins.2013.00210

Solomon-Lane, T.K., Pradhan, D.S., Willis, M.C., Grober, M.S., 2015. Agonistic reciprocity is
associated with reduced male reproductive success within haremic social networks. Proc. R.

Soc. B Biol. Sci. 282, 20150914. https://doi.org/10.1098/rspb.2015.0914

Solomon-Lane, T.K., Pradhan, D.S., Willis, M.C., Grober, M.S., 2014. Female, but not male, agonistic behaviour is associated with male reproductive success in stable bluebanded goby (Lythrypnus dalli) hierarchies. Behaviour 151, 1367–1387. https://doi.org/10.1163/1568539X-00003188

Stamps, J.A., 2003. Behavioural processes affecting development: Tinbergen's fourth question comes of age. Anim. Behav. 66, 1–13. https://doi.org/10.1006/anbe.2003.2180

Stamps, J.A., Groothuis, T.G.G., 2010. The development of animal personality: Relevance, concepts and perspectives. Biol. Rev. 85, 301–325. https://doi.org/10.1111/j.1469-185X.2009.00103.x

Stearns, S.C., 1989. The Evolutionary Significance of Phenotypic Plasticity. BioScience, 39, 436–445. https://doi.org/10.2307/1311135

Taborsky, B., 2016. Opening the Black Box of Developmental Experiments: Behavioural Mechanisms Underlying Long-Term Effects of Early Social Experience. Ethology 122, 267–283. https://doi.org/10.1111/eth.12473

Taborsky, B., Arnold, C., Junker, J., Tschopp, A., 2012. The early social environment affects social competence in a cooperative breeder. Anim. Behav. 83, 1067–1074. https://doi.org/10.1016/j.anbehav.2012.01.037

Taborsky, B., Tschirren, L., Meunier, C., Aubin-Horth, N., 2013. Stable reprogramming of brain transcription profiles by the early social environment in a cooperatively breeding fish. Proc. Biol. Sci. 280, 20122605. https://doi.org/10.1098/rspb.2012.2605

Tinbergen, N., 1963. On aims and methods of Ethology. Anim. Biol. 55, 297–321. https://doi.org/10.1163/157075605774840941

Turecki, G., Meaney, M.J., 2016. Effects of the Social Environment and Stress on Glucocorticoid Receptor Gene Methylation: A Systematic Review. Biol. Psychiatry 79, 87–96. https://doi.org/10.1016/j.biopsych.2014.11.022

Verbeek, M.E.M., Drent, P.J., Wiepkema, P.R., 1994. Consistent individual differences in early exploratory behaviour of male great tits. Anim. Behav. https://doi.org/10.1006/anbe.1994.1344

Wada, H., 2008. Glucocorticoids: mediators of vertebrate ontogenetic transitions. Gen. Comp. Endocrinol. 156, 441–53. https://doi.org/10.1016/j.ygcen.2008.02.004

Weitekamp, C.A., Hofmann, H.A., 2017. Neuromolecular correlates of cooperation and conflict during territory defense in a cichlid fish. Horm. Behav. 89, 145–156. https://doi.org/10.1016/j.yhbeh.2017.01.001

Weitekamp, C.A., Solomon-Lane, T.K., Del Valle, P., Triki, Z., Nugent, B.M., Hofmann, H.A., 2017. A Role for Oxytocin-Like Receptor in Social Habituation in a Teleost. Brain. Behav. Evol. 89, 153–161. https://doi.org/10.1159/000464098

West-Eberhard, M.J., 1989. Phenotypic plasticity and the origins of diversity. Annu. Rev. Ecol. Syst. 20, 249–278. https://doi.org/10.1146/annurev.es.20.110189.001341

White, D.J., 2010. The Form and Function of Social Development - Insights From a Parasite. Curr. Dir. Psychol. Sci. 19, 314–318. https://doi.org/10.1177/0963721410383384

White, D.J., King, A.P., West, M.J., 2002. Facultative development of courtship and communication in juvenile male cowbirds (Molothrus ater). Behav. Ecol. 13, 487–496. https://doi.org/10.1093/beheco/13.4.487

Wilson, E.O., 1980. Sociobiology: the Abridged Edition. The Belknap Press of Harvard University Press, Cambridge.

Wingfield, J.C., Hegner, R.E., Dufty, Jr., A.M., Ball, G.F., 1990. The "Challenge Hypothesis":
Theoretical Implications for Patterns of Testosterone Secretion, Mating Systems, and
Breeding Strategies. Am. Nat. 136, 829. https://doi.org/10.1086/285134
Young, A.J., Carlson, A. a, Monfort, S.L., Russell, A.F., Bennett, N.C., Clutton-Brock, T., 2006.
Stress and the suppression of subordinate reproduction in cooperatively breeding meerkats.
Proc. Natl. Acad. Sci. 103, 12005–12010. https://doi.org/10.1073/pnas.0510038103

946
947

948 **Table 1:** Linear regressions of focal fish standard length with behavior in the open field, social

949 cue investigation, dominance, and subordinate behavior assays. The zones of the tank refer to the

950 frequency of entering that zone. Adjusted R^2 values are reported. Significant results following a

951 false discovery rate correction are bolded.

Assay	Behavior	r-squared	p-value
Open field	Investigate zone	0.17	**0.0036**
	Far zone	0.15	**0.0059**
	Close zone	0.11	**0.017**
	Territory zone	0.18	**0.0028**
Social cue investigation	Investigate zone	0.035	0.13
	Far zone	0.085	**0.038**
	Close zone	0.096	**0.031**
	Territory zone	0.11	**0.021**
Dominance behavior	Approach	0.089	**0.035**
	Displace	0.10	**0.024**
	Territory zone (focal)	0.11	**0.022**
	Territory zone (subordinate fish)	0.14	**0.011**
	Territory zone (both)	0.11	**0.02**
Subordinate behavior	Approach	-0.02	0.57
	Displace	-0.0044	0.37
	Submit	-0.023	0.72
	Territory zone (focal)	-0.0015	0.34
	Territory zone (subordinate fish)	-0.02	0.67
	Territory zone (both)	-0.024	0.87

Figure 1: Experimental setup for behavior assays. Juvenile behavior was observed in a novel

experimental tank in four sequential assays administered in the same order, each lasting 30 min.

A terracotta shard served as a shelter and/or territory. The black lines (dotted, solid) were drawn

on the tank bottom in permanent maker, dividing the tank into four zones: territory, close, far,

and investigate. The center dividing line (white) was not drawn (A). The focal fish was alone in

the tank for the open field assay, and the time in each zone and frequency of entered each zone

was recorded (B, assay 1). For the social cue investigation, a juvenile inside of a scintillation vial

was placed in the circle within the investigate zone (see C). The time in and frequency of

entering each zone was recorded (B, assay 2). The social cue was removed and a freely

swimming, novel cue fish (smaller than the focal) was added to the tank for the dominance

behavior assay (B, assay 3). The small cue fish was then removed and a freely swimming, novel

cue fish (larger than the focal) was added to the tank for the subordinate behavior assay (B, assay

4). Social interactions were recorded for the dominant and subordinate behavior assays. The time

in and frequency of entering the territory zone was also recorded for both fish.

Figure 2: Principal component analysis (PCA) of focal fish behavior from all four assays (open

field, social cue investigation, dominance, subordinate behavior). Differences in PC1 between

group- and pair-reared juveniles (p=0.029) (A). Vector plot showing the PCA variables that load

on PC1 (B). Differences in PC6 between group- and pair-reared juveniles (p= 4.082e-05) (C).

Percentages refer to the amount of variation explained by that component. Pair (n=18

individuals). Group (n=24 individuals). Social cue investigation (SCI). Open field exploration

(OF). *p<0.05, ****p<0.001.

Figure 3: Separate principal component analyses performed for the open field (A), social cue

investigation (B), and dominance behavior (C) assays. Both focal and non-focal fish variables

(behavior, size). The significant, positive correlations about the PC1s are shown in a three-

dimensional plot (D). Percentages refer to the amount of variation explained by that component.

Pair (n=18 individuals). Group (n=24 individuals).

Figure 4: Relative gene expression calculated using ΔΔCT analysis (reference gene 18S) for

juveniles reared in isolation (1 week, n=8), pairs (1 week or 5 weeks, n=18), and groups (1 week

or 5 weeks, n=22). Androgen receptor α (ARα). Estrogen receptor α (ERα). Glucocorticoid

receptors (GR). Mineralocorticoid receptor (MR). Corticotropin-releasing factor (CRF). Letters

indicate significant *post hoc* differences within a gene (p<0.05).

Figure 5: Partial correlation network of gene expression in pair-reared juveniles (n=18) (A) and

group-reared juveniles (n=22) (B). Nodes are the candidate genes. Edges represent partial

correlations between nodes. Only significant partial correlations are shown (p<0.05), and edge

thickness indicates correlation strength. There were no significant partial correlations for

juveniles reared in isolation (n=8) (p>0.05). Androgen receptor α (AR). Estrogen receptor α

(ER). Glucocorticoid receptors (GR). Mineralocorticoid receptor (MR). Corticotropin-releasing

factor (CRF).

1005

Figure 6: Principal component analysis of relative expression of candidate genes in whole brain from group- (n=22) and pair-reared (n=18) juveniles. Differences in PC1 between juveniles in treatment groups for 1 vs. 5 weeks (A). Differences in PC4 between juveniles reared in groups vs. pairs (p=0.011) (B). Vector plot showing how candidate genes load on PC1 and PC4 (C). Percentages refer to the amount of variation explained by that component. Androgen receptor α (AR). Estrogen receptor α (ER). Glucocorticoid receptors (GR). Mineralocorticoid receptor (MR). Corticotropin-releasing factor (CRF).