1 **Early-life social environment alters juvenile behavior and neuroendocrine function in a**

2 **highly social cichlid fish**

6 Tessa K. Solomon-Lane[1-3]* & Hans A. Hofmann[1-3]

8 [1]Department of Integrative Biology, [2]Institute for Neuroscience, [3]Center for Computational

9 Biology and Bioinformatics, The University of Texas at Austin, Austin, TX 78712

*Corresponding author:

Tessa Solomon-Lane

University of Texas at Austin

Department of Integrative Biology

1 University Station #C0930

Austin, TX 78712

tksolomonlane@utexas.edu

512-475-7318

Abstract

Early-life experiences can shape adult behavior, with consequences for fitness and health, yet fundamental questions remain unanswered about how early-life social experiences are translated into variation in brain and behavior. The African cichlid fish *Astatotilapia burtoni*, a model system in social neuroscience, is well known for its highly plastic social phenotypes in adulthood. Here, we rear juveniles in either social groups or pairs to investigate the effects of early-life social environments on behavior and neuroendocrine gene expression. We find that both juvenile behavior and neuroendocrine function are sensitive to early-life effects. Behavior robustly co-varies across multiple contexts (open field, social cue investigation, and dominance behavior assays) to form a behavioral syndrome, with pair-reared juveniles towards the end of syndrome that is less active and socially interactive. Pair-reared juveniles also submit more readily as subordinates. In a separate cohort, we measured whole brain expression of stress and sex hormone genes. Expression of glucocorticoid receptor (GR) 1a was elevated in group-reared juveniles, supporting a highly-conserved role for the stress axis mediating early-life effects. The effect of rearing environment on androgen receptor (AR) α and estrogen receptor (ER) α expression was mediated by treatment duration (1 vs. 5 weeks). Finally, expression of corticotropin-releasing factor (CRF) and GR2 decreased significantly over time. Rearing environment also caused striking differences in gene co-expression, such that expression was tightly integrated in pair-reared juveniles, but not group-reared or isolates. Together, this research demonstrates the important developmental origins of behavioral phenotypes and identifies potential behavioral and neuroendocrine mechanisms.

Key words: early environment, ontogeny, social behavior, behavioral syndrome, hypothalamic-pituitary-adrenal axis, stress hormones, sex hormones

Introduction

Ontogeny has long been recognized as essential to understanding phenotype (Tinbergen, 1963), yet the early-life origins of individual behavioral variation remain understudied. Development reveals the proximate mechanisms by which genes interact with the environment during early life to sculpt the 'machinery of behavior' (Stamps, 2003; Tinbergen, 1963). Current or predicted environmental conditions can trigger developmental plasticity, and the resulting changes are often long-lasting, or even permanent, and can facilitate locally-adapted (e.g., predator resistant, Gilbert, 2001) phenotypes (Kasumovic and Brooks, 2011; Langenhof and Komdeur, 2018; Lummaa and Clutton-Brock, 2002; Piersma and Drent, 2003; Snell-Rood, 2013; Stamps, 2003; Stearns, 1989; West-Eberhard, 1989). The developmental mechanisms that shape social behavior via underlying neural regulatory mechanisms should be a particularly important target for natural selection (Taborsky, 2016) because of the direct consequences of social behavior for fitness and health (e.g., Bennett et al., 2006; Meyer-Lindenberg and Tost, 2012; Silk, 2007; Solomon-Lane et al., 2015; Wilson, 1980).

Social stimuli are among the most important attributes of the early-life environment (Taborsky, 2016). Although maternal (and, to a lesser extent, paternal) interactions have largely been the focus (e.g., Champagne & Curley, 2005; McClelland, Korosi, Cope, Ivy, & Baram, 2011), the broader early-life social environment is increasingly recognized for its role in behavioral and neural plasticity (Buist et al., 2013; Creel et al., 2013; Jonsson and Jonsson, 2014; Kasumovic and Brooks, 2011; Taborsky, 2016; White, 2010). For example, the early presence of brood care helpers, unrelated adult males, and multiple mothers and litters have long-term effects on social behavior in the Daffodil cichlid fish *Neolamprologus pulcher* (Arnold and Taborsky, 2010; Taborsky et al., 2012), brown-headed cowbirds (White et al., 2002), and laboratory mice

(Branchi et al., 2013, 2006; D'Andrea et al., 2007), respectively. These features of the social

environment alter the quality and quantity of social experiences and sensory cues perceived,

which together influence neural function and behavior (Taborsky, 2016). Developmental

plasticity may be limited to a single behavior or extend to an entire suite of behaviors (i.e., a

behavioral syndrome), and the effects may be context-specific (Bell, 2007; Snell-Rood, 2013;

Stamps, 2003; Stamps and Groothuis, 2010).

Neuroendocrine signaling is a primary mechanism by which environmental conditions

and experience are translated into physiological responses (Crespi and Denver, 2005; Remage-

Healey and Romero, 2000; Wingfield et al., 1990). Hormones are also important sources of

individual variation in social behavior (e.g., across seasons, sexes, reproductive tactics) and

underlie developmental plasticity relevant to adult behavior. The stress axis, or hypothalamic-

pituitary-adrenal (interrenal in fish; HPA/I) axis, is widely implicated as a highly-conserved

mechanism of early-life effects (Champagne and Curley, 2005; Francis et al., 1999; McClelland

et al., 2011; Taborsky, 2016). In response to an environmental stressor, which includes any

external condition that disrupts or threatens to disrupt homeostasis, the HPA/I axis integrates

relevant internal and external cues and coordinates a response, such as changes in behavior and

physiology. The stress response is initiated by the release of corticotropin-releasing factor (CRF)

from the hypothalamus, which signals to the pituitary to release adrenocorticotropic hormone,

which then signals the adrenal glands to release glucocorticoids (e.g., cortisol in fish) (Denver,

2009; Lowry and Moore, 2006; Wendelaar Bonga, 1997).

Effects of early-life experiences on HPA/I axis function have been demonstrated in every

major vertebrate lineage (e.g., birds: Banerjee, Arterbery, Fergus, & Adkins-Regan, 2012;

mammals: Champagne & Curley, 2005; amphibians: Crespi & Denver, 2005; fish: Jonsson &

Jonsson, 2014). For example, the presence of brood helpers during early-life affects social

behavior in the cooperatively breeding *N. pulcher* cichlid via changes in neural expression levels

of CRF and glucocorticoid receptor (GR), as well as the ratio of the mineralocorticoid receptor

(MR) to GR1 (Taborsky et al., 2013). Stress axis mechanisms can also mediate the effects of the

early-life social environment on human health (e.g., Turecki & Meaney, 2016). Sex steroid

hormones (e.g., androgens, estrogens) also play a role mediating the long-term effects of early-

life experiences (Adkins-Regan, 2009; Brown and Spencer, 2013; Shepard et al., 2009) and

regulating social behavior (Goodson, 2005; Newman, 1999). For example, neural estrogen

receptor expression is associated with maternal behavior in mother rats and offspring (Cameron

et al., 2008; Champagne et al., 2003; Champagne and Meaney, 2007), and socially stressed pre-

and postnatal female guinea pigs have upregulated neural estrogen and androgen receptor levels,

elevated testosterone, and masculinized behavior (Kaiser et al., 2003). Together, these and other

neuroendocrine systems interact (e.g., Acevedo-Rodriguez et al., 2018) to affect behavior.

To investigate the effects of the early-life social environment on behavior and its

neuroendocrine mechanisms, we used the highly social African cichlid *Astatotilapia burtoni*, a

model system in social neuroscience (Fernald and Maruska, 2012; Hofmann, 2003; Stevenson et

al., 2017). Adults of this species form mixed-sex, hierarchical communities with males of

dominant or subordinate status and females. Dominant males are territorial, reproductively

active, and colorful. In comparison, subordinate males shoal with females, are reproductively

suppressed, and drab in coloration. Male status is socially regulated, and individuals regularly

transition between status phenotypes (Fernald and Maruska, 2012; Hofmann, 2003). Adults, and

juveniles (Fernald and Hirata, 1979), express a suite of highly evolutionarily conserved social

behaviors, including aggression, affiliation, courtship, and cooperation (Fernald, 2012; Hofmann,

2003; Weitekamp et al., 2017). Substantial progress has also been made towards understanding

variation in stress and sex steroid hormone signaling, including in the regulation of social

behavior (Chen and Fernald, 2008; Fox et al., 1997; Greenwood et al., 2003; Munchrath and

Hofmann, 2010; O'Connell and Hofmann, 2012a). All GRs (Greenwood et al., 2003), estrogen

receptors (ER), and androgen receptors (AR) (Munchrath and Hofmann, 2010) have been studied

in the adult *A. burtoni* brain, and neuroendocrine function can vary substantially. Subordinate

males, for example, have lower levels of whole brain CRF and GR2 (Chen and Fernald, 2008),

higher cortisol, and lower testosterone than dominants (Fox et al., 1997; O'Connell and

Hofmann, 2012a), although these patterns can vary dynamically (Maguire and Hofmann, in

prep.). The transcriptomic response in the preoptic area (POA) to pharmacological manipulation,

such as an ER antagonist, is also status-specific (O'Connell and Hofmann, 2012a).

Given this rich literature on adult *A. burtoni*, it may seem surprising that the

developmental origins of adult phenotypic variation remain largely unknown. The few studies

that have investigated juveniles demonstrate the importance of early-life. For example, the

development of male behavior and nuptial coloration, as well as reproductive maturation, are

affected by the early-life social environment (Fernald and Hirata, 1979; Fraley and Fernald,

1982). Gestational cues (e.g., maternal social crowding) also have lasting effects on methylation

and transcription of the *gnrh*1 gene in offspring (Alvarado et al., 2015). This result is particularly

interesting given that POA GnRH1 neurons, which regulate gonadotropin release from the

pituitary, are socially modulated in adults (Davis and Fernald, 1990; Hofmann and Fernald,

2001). However, studies of the effects of different early-life experiences on other neuroendocrine

pathways or behavior are lacking.

In the present study, we conducted two experiments to test the hypothesis that the early-

life social environment generates variation in juvenile behavior through neuroendocrine gene

expression. We manipulated the early-life social environment, and consequently social

experience, by rearing juveniles in either social groups or pairs. The natural distribution of

territories and shoals across shallow shore pools and river estuaries (Fernald and Hirata, 1977;

Rajkov et al., 2018) suggest that *A. burtoni* encounter a variety of dynamic social environments,

including during development, although the degree of variation across individuals and over time

has not been quantified. By directly manipulating group size, we can experimentally enhance the

frequency, diversity, and/or complexity of early-life social experiences. Similar manipulations

impact behavioral and neural development in a variety of species (reviewed in Taborsky, 2016).

In the group condition, social experience implies interactions with more social partners, who also

vary in size, sex, experience, and patterns of behavior. Interactions in groups can also involve

more than two individuals, and it is possible to observe and learn from interactions of group

members as a bystander. Although it has not been tested in juveniles, adults are capable of

gaining important social information as a bystander (Desjardins et al., 2012, 2010; Grosenick et

al., 2007). In the pair condition, juveniles occupy only one social role in a relationship with just

one other individual. We predicted that rearing environment might affect social behavior—

including social investigation, dominant, and subordinate behavior—possibly in a consistent

manner across contexts. We also predicted effects on the expression of various genes that are

part of candidate neuroendocrine systems known to mediate early-life experiences in other

systems. Specifically, related to the HPA/I axis, we measured glucocorticoid receptor 1a (GR1a),

glucocorticoid receptor 1b (GR1b), glucocorticoid receptor 2 (GR2) (nomenclature from

Maruska & Fernald, 2010), mineralocorticoid receptor (MR), and CRF. For sex steroid hormone

signaling, we quantified androgen receptor α (ARα) and estrogen receptor α (ERα). By

investigating these early-life effects in juveniles, we can identify important intermediary steps

that inform how developmental plasticity may shape the adult phenotype.

Methods

Animals

Juvenile *A. burtoni* came from a laboratory population descended from a wild-caught

stock. The adults that bred the juveniles were housed in naturalistic social groups of males and

females. Dominant males court gravid females that then lay eggs in his territory. The female then

scoops up the eggs into her mouth, where the male fertilizes them. The mother orally incubates

the larvae as they develop for 10-13 days. Under natural (and some laboratory) conditions,

juveniles remain close to their mother for the 2-4 weeks following their initial release from her

mouth. As they age, juveniles seek shelter in her mouth less and less often. In the first two

weeks, juveniles primarily school together, with overt social interactions beginning at 2-3 weeks

old (Fernald and Hirata, 1979; Renn et al., 2009). Social behaviors, such as chasing, nipping,

territorial displays, emerge in a predictable sequence as juveniles approach reproductive

maturity, which can occur as early as 15 weeks, depending on the early-life social conditions

(Fernald and Hirata, 1979; Fraley and Fernald, 1982).

We removed juveniles from the mother's mouth 6-12 days post-fertilization. Once

sufficiently developed (~day 12, freely swimming with no remaining yolk), juveniles were

transferred into experimental rearing environments. Juveniles are all silver (drab) in coloration,

and none developed coloration during the study, which would indicate reproductive maturity for

males. Sex cannot be determined anatomically until maturation; therefore, the sex ratios of our

rearing environments, and the sex of the focal individuals, is unknown. The sex ratio of *A*.

burtoni broods is approximately 1:1 (Heule et al., 2014). All work was done in compliance with

the Institutional Animal Care and Use Committee at The University of Texas at Austin.

Experimental rearing conditions (Experiments 1 & 2)

As the first study of this kind in this species, we opted to quantify behavior and gene

expression in separate experiments in order to capture different developmental time points. In

Experiment 1, juveniles for the behavioral assays were reared in social groups of 16 fish (n=12

groups) or in pairs (n=9 pairs) for 58-73 days (average 65.76 ± 0.81; ~8-10 weeks). This is the

longest duration that could be used without juveniles reaching reproductive maturity. In

Experiment 2, neural gene expression was measured in a separate cohort of juveniles reared in

social groups of 16 fish, pairs, or in isolation for 1 week (groups: n=8; pairs: n=8; isolates: n=8)

or 5 weeks (groups: n=14; pairs: n=10). Here, we aimed to capture early changes in gene

expression that might set individuals along different developmental trajectories. Isolation was

included because we expected it to impact gene expression in this highly social species, not as a

social control. We cannot distinguish between the effects of chronological age from the treatment

duration (1 vs. 5 weeks) in this study.

For both Experiments, juveniles from multiple clutches of the same age and

developmental stage (day 12-14 fry) were divided among treatment groups. Group-reared fish

were housed in 35 L aquaria with three terracotta pot shards for shelter and/or territory. Pairs and

isolated fish were housed in small aquaria (22.9 x 15.2 x 15.2 cm) with one terracotta pot shard.

The volume of water per fish was similar for the group (2.6 L) and paired (2.7 L) treatments.

Juveniles were fed daily with Hikari plankton (Pentair Aquatic Eco-Systems, Cary, NC). The

food was mixed in water, and a transfer pipette was used to deliver a set volume to each tank.

Groups received eight times more food than pairs. Pairs and isolated fish received the same

amount. All juveniles were maintained on a 12:12 light/dark cycle.

Experiment 1: Behavioral assays

We quantified behavior in four assays, which were always presented in the same

sequence (Fig 1): an open field test that is commonly used in other species to assess activity and

anxiety (e.g., Cachat et al., 2010; Prut & Belzung, 2003); a social cue investigation as a measure

of social motivation or preference (e.g., Bonuti & Morato, 2018; Moy et al., 2004); and social

interactions within either dominant or subordinate status contexts, which individuals regularly

experience in social communities of *A. burtoni* (Hofmann, 2003). Behavioral neuroscientists

employ a wide range of different assays across different model systems, and we explored which

assays juvenile *A. burtoni* would participate in in a series of pilot experiments. We decided on

this combination of assays because each assay has been used with multiple species, thus allowing

for cross-species comparisons, and the target behaviors (e.g., locomotion, space use, social

approach, social interaction) are all expressed by *A. burtoni* in natural contexts and directly

relevant to adult social status and reproduction (e.g., via territoriality, aggression). Including

multiple assays in combination also provides a more comprehensive understanding of behavioral

phenotype, which is complex and expressed in context-specific ways.

Behavior for both members of the pairs (n=18 individuals) and two fish from each group

(n=24 individuals) was analyzed. To choose focal individuals from the groups, we removed all

fish from the aquarium and selected, by eye, one of the largest fish. A smaller fish was then

chosen such that the ratio of large-to-small fish standard length (SL, mm) was approximately

equal in the group and a pair from the same cohort of juveniles (same age). These smaller fish

were never the smallest in their groups. Because size is a strong predicator of social status

(Alcazar et al., 2014), the larger fish was very likely to have dominance experience, similar to

the larger fish in the pair. The smaller fish were very likely to have subordinate and dominant

interactions with larger and smaller individuals in the group, respectively. Standard length was

recorded for all focal fish.

Behavior was observed in novel, small aquaria (22.9 x 15.2 x 15.2 cm) without covers.

For analysis, the aquaria were divided into 4 zones (Fig 1), delineated with permanent marker. In

the middle of each short side, a circle was drawn (28 mm diameter) to indicate the placement of

the scintillation vial (see below: social cue investigation). An arc 2.54 cm from the edge of that

circle was drawn to form a semicircle. One semicircle was designated the "territory" zone and

had a terracotta pot shard for a shelter and/or territory. The other semicircle was designated the

"investigate" zone. The "close" zone was between the territory zone and halfway along the long

side of the tank. The "far" zone was between the halfway mark and the investigate zone (Fig 1).

Video cameras recorded behavior from above so that all areas of the tank, except under the

terracotta pot shard, were visible. Solomon Coder was used for analysis

(www.solomoncoder.com). All observations were made by the same observer who was blind to

treatment. Ten minutes of behavior was analyzed from each behavior assay for a total of 40 min

of behavior scored for each individual.

Open field test: The focal fish was transferred to the test aquarium with a hand net and

remained in the tank alone for 30 min. Movement around the tank was observed from minutes 20

to 30. We recorded the number of times a fish crossed into each zone (frequency) and the time

(s) spent in each zone. Social cue investigation: Novel juveniles were collected from a

community tank and placed into scintillation vials (20 mL). The top of the vial was covered with

parafilm with holes to allow water through. A vial containing one cue fish was placed into each

test aquarium (n=16 group-reared, n=13 pair-reared). Cue fish were 0-6.4 mm SL (average 3.37

$\pm$ 0.27) smaller than their focal fish. An empty vial was used as a control (n=8 group-reared, n=5

pair-reared). The social cues were in the aquarium for 30 min. Movement around the tank

(frequency and time in each zone) was scored from minutes 2 to 12.

Dominance behavior: The scintillation vials were removed from the aquaria and a novel

smaller fish (by 1-6.4 mm SL, average 3.37 $\pm$ 0.25) was immediately added to each aquarium,

freely swimming with the focal fish. The pair remained together for 30 minutes, and behavior

was scored from minutes 2 to 12. Subordinate behavior: The small cue fish was removed from

the aquaria and a novel, larger fish (by 2.4-12 mm SL, average 5.74 $\pm$ 0.34) was immediately

added to each aquarium, freely swimming with the focal fish. The pair remained together for 30

minutes, and behavior was scored from minutes 2 to 12. In the dominance and subordinate

behavior assays, we analyzed agonistic interactions between the pair. An approach was defined

as one fish swimming directly towards any part of the other fish's body, within 3 body lengths. If

the approached fish responded by moving away, in any direction, the behavior was recorded as a

displacement for the initiator and a submission for the responder. From these measures, we

calculated agonistic efficiency, or the proportion of approaches that led to a displacement

(Solomon-Lane et al., 2014), for focal and cue fish. The difference in agonistic efficiency

between the focal and cue fish was used as a measure of agonistic asymmetry, which

characterizes status relationships (Drews, 1993). We also recorded the frequency of entering and

the time spent in the territory, for the focal fish, cue fish, and both together.

Experiment 2: Whole brain gene expression

Whole brain gene expression for two fish from each group (1 week: n=8; 5 weeks: n=14),

both members of the pairs (1 week: n=8; 5 weeks: n=10), and every isolate (1 week: n=8) was

analyzed. Because the present study is the first to examine the neuromolecular substrates

associated with early life social experience in *A. burtoni*, we did not have an *a priori* expectation

as to which brain regions or cell types might be the most critical to examine. Therefore, we

decided to analyze expression in whole brain, even though important differences in circuits and

brain regions may not be identified using this approach. It should also be noted that recent

evidence suggests that patterns of expression specific to brain region, or even cell-type, can be

inferred from bulk tissue samples (Kelley et al., 2018), such as whole brain.

Focal individuals from the group condition were selected haphazardly. Juveniles were

removed from their rearing environments with a hand net and rapidly decapitated. The brains

were dissected immediately, flash frozen on dry ice, and stored at -80° C until processing. Gene

expression was quantified using qPCR and previously validated primers (Supplemental Table 1,

Chen and Fernald, 2008; Greenwood et al., 2003; O'Connell and Hofmann, 2012a) for GR1,

GR2a, GR2b, MR, CRF, ARα, and ERα, as well as control genes 18S and G3PDH. With regards

to (nuclear) sex steroid receptors, we chose subtypes ARα and ERα because of their

demonstrated role in the regulation of adult *A. burtoni* social and reproductive behavior

(Burmeister et al., 2007; Korzan et al., 2014; Maruska, 2015; O'Connell and Hofmann, 2012a).

Other subtypes (e.g., ARβ, ERβa,b), as well as progesterone receptor, also have distinct

distributions and regulatory roles (Burmeister et al., 2007; Munchrath and Hofmann, 2010) and

certainly warrant investigation in future studies. RNA was extracted using the Maxwell 16 LEV

simplyRNA Tissue Kit (Promega, Madison, WI), and the Promega GoScript Reverse

Transcription System (Promega, Madison, WI) was used for reverse transcription. PowerUp

SYBR Green Master Mix (ThermoFisher Scientific, Waltham, MA) was used for quantitative

PCR. All standard kit protocols were followed. Relative gene expression levels were quantified

using ΔΔCT analysis, using 18S and G3PDH as reference genes. The results are largely

concordant independent of the reference gene used. Here, we present the analyses for 18S, as this

gene has shown very little expression variation across social phenotypes in transcriptome studies

of *A. burtoni* (O'Connell and Hofmann, 2012a; Renn et al., 2008).

Statistical analyses

All statistical analyses were conducted using R Studio (version 1.0.143). Results were

considered significant at the $p<0.05$ level, and averages ± standard error of the mean are included

in the text. Cohen's d is reported to estimate effect size (small effect: $0.2<d<0.5$; medium:

$0.5<d<0.8$; large: $0.8<d$). The box of the box and whisker plots show the median and the first and

third quartiles. The whiskers extend to the largest and smallest observations within or equal to

1.5 times the interquartile range. Comparisons between group- and pair-reared juveniles were

conducted using t-tests for fish SL, time and frequency in each tank zone, and rates of agonistic

behavior. Mann-Whitney-Wilcoxon tests were used for data that did not meet the assumptions of

parametric statistics. Regression analysis was used to identify significant associations between

SL and frequency and time in a zone and between SL and agonistic behavior. We used a false

discovery rate correction for regressions with focal fish SL (Benjamini and Hochberg, 1995).

Two-way ANOVAs were used to identify significant effects of rearing environment, presence of

the social cue, or an interaction, on the frequency and time spent in each zone of the tank. We

used Principal Components Analysis (PCA) to identify how behaviors clustered across the four

assays and for each assay individually. Independent t-tests were used to compare principal

component scores between group- and pair reared juveniles. Paired t-tests (or Mann-Whitney-

Wilcoxon tests) were used to compare principal component scores between the larger and

smaller fish sampled from groups and pairs. Correlation analysis was used to identify significant

associations among principal components (PCs).

We used two-way ANOVAs to identify significant effects of rearing environment (group,

pair, isolated), treatment duration (1 week, 5 weeks), or an interaction on the expression of

individual candidate genes. All gene expression data were log transformed to meet the

assumptions of parametric statistics. Partial correlation networks were calculated using the

"ppcor" package in R and visualized using "qgraph." The nodes of the networks represent the

gene. The edges are the partial correlation coefficient, with thicker edges indicating stronger

correlations. Only significant correlations are shown. Mantel tests were used to test for pairwise

differences between the gene expression networks. A non-significant p-value (> 0.05) indicates

that the partial correlation matrices are not related.

Results

Experiment 1

Standard length

After 8-10 weeks in their respective treatment condition, group-reared juveniles ($16.85 \pm$

0.32 mm SL) were significantly larger than pair-reared juveniles (13.76 ± 0.40 mm SL) ($t=6.00$,

$p=7.25e-7$, $d=1.89$). This size difference influenced the size of the fish selected to be the social

stimuli. Specifically, the difference in SL between the focal fish and the social cue ($t=3.38$,

$p=0.0016$, $d=1.02$), as well as the focal fish and the small cue fish ($t=3.48$, $p=0.0013$, $d=1.09$),

was significantly greater for group-reared juveniles. The SL difference between the focal fish

and the large cue fish was significantly greater for pair-reared juveniles (t=-3.22, p=0.0025,

d=0.95). Relative size differences followed the same pattern as absolute size differences (data

not shown).

Open field test and social cue investigation

In the open field test (and subsequent assays), juveniles of both treatment groups moved

readily around the novel environment with minimal acclimation. We present the data for the

frequency of entering each zone (Supplemental Fig 1A-D). There were no significant effects for

the time spent in each zone (p>0.05). Group-reared juveniles entered the territory (Mann-

Whitney-Wilcoxon test: W=299, p=0.034, d=0.51), close (W=293.5, p=0.049, d=0.41), and

investigate zones (W=293.5, p=0.049, d=0.60) significantly more frequently than pair-reared

juveniles. There was no significant difference for the far zone (W=289, p=0.064).

Next, we used a social cue investigation task to examine whether and how rearing

environment and/or the presence of the social cue affect locomotor activity (Supplemental Fig

1E-H). Two-way ANOVA revealed that, following the addition of the social cue, juveniles

entered the investigate zone significantly more frequently than controls ($F_{1,36}$= 4.91, p=0.033,

d=0.96). There was no effect of rearing environment ($F_{1,36}$=1.69, p=0.20) and no interaction

($F_{1,36}$=0.046, p=0.83). There was no effect of rearing environment ($F_{1,36}$=2.68, p=0.11), social

cue ($F_{1,36}$=0.87, p=0.36), or an interaction ($F_{1,36}$=0.84, p=0.37) on frequency of entering the far

zone. Group-reared juveniles entered the close zone significantly more than pair-reared juveniles

($F_{1,35}$=4.47, p=0.042, d=0.71), but there was no effect of the social cue ($F_{1,35}$=0.11, p=0.74) and

no interaction ($F_{1,35}$=0.44, p=0.52). There was no effect of rearing environment ($F_{1,35}$=3.28,

p=0.079), social cue ($F_{1,35}$=0.17, p=0.68) and no interaction ($F_{1,35}$=0.83, p=0.37) on the

frequency of entering the territory zone. Linear regression analyses show that SL is not

associated with the frequency of entering zones of the tank for group- or pair-reared juveniles

(Supplemental Table 2).

<u>Dominant and subordinate behavior</u>

Rearing environment did not affect rates of focal fish behavior (Supplemental Fig 2). As

the dominant fish, there were no differences in approaching (W=242.5, p=0.20) or displacing

(W=253, p=0.12) the small cue fish. As the subordinate, there were no differences in

approaching (W=205.5, p=0.85), displacing (W=214.5, p=0.62), or submitting to (W=217.5,

p=0.56) the large cue fish. In the dominance assay, rearing environment did not affect agonistic

efficiency for the focal fish (t=0.83, p=0.41), small cue fish (W=115.5, p= 0.97), or the

difference between the pair (t=1.03, p=0.32). In the subordinate assay, although there was no

effect of rearing environment on agonistic efficiency for the focal fish (W=169.5, p=0.28) or the

large cue fish (W=112.5, p=0.061), the difference in agonistic efficiency was significantly higher

for pair-reared juveniles (t=-2.42, p=0.022, d=0.81). Linear regression analyses show that SL is

not associated with social behavior for group- or pair-reared juveniles (Supplemental Table 2).

<u>Multivariate analysis of behavior across assays</u>

In order to gain more insight into this multivariate dataset, we employed PCA to

determine which measures of morphology (i.e., size) and behavior might act in concert to explain

different aspects of the variability across individuals, including based on rearing environment

and whether the focal individual was the larger or smaller fish sampled from the group or pair.

Given that body size serves as a reliable proxy for social status experience in adults, we refer to

the larger and smaller juvenile as dominant and subordinate, respectively. We first conducted a

PCA that included variables from each of the four assays: focal fish SL; frequency of entering

each zone in the open field test and social cue investigation; focal fish social approaches and

displacements as a dominant towards the small cue fish; and focal fish approaches,

displacements, and submissions as a subordinate with the larger cue fish. We found that principal

component (PC) 1 accounts for 43.3% of the total variance and differs significantly between

group- and pair-reared juveniles (t=-2.30, p=0.029, d=0.75, Fig 2A). There was a trend for PC2

(16.4%; z= -1.96, p=0.05, d=0.39, Fig 2B) to differ based on status experience (or relative size),

and the difference was significant for PC5 (6.6%; t=- -2.16, p=0.043, d=0.53, Fig 2C). PC6

(5.0%) also differed significantly between group- and pair-reared juveniles (t=4.66, p= 4.082e-5,

d=1.46, Fig 2D). No significant differences were identified for other PCs (p>0.05). As the vector

plot in Fig 2E shows, variables from the open field test, social cue investigation, and dominance

behavior assay all load on PC1, along with focal fish SL, while measures of behavior during the

subordinate assay load on PC2. The vector plot in Fig 2F shows that a number of behaviors load

on PC5, the strongest of which relate movement around the tank during the open field and social

cue investigation assays. Focal fish SL loads most strongly on PC6.

409   To disentangle the possible effects of SL and rearing environment on behavior, we re-ran

the PCA without focal fish SL. In this analysis, PC1 (44.8% of the variance) still differs

significantly between group-reared and pair-reared juveniles (W=126, p=0.022, d=0.64).

Although focal fish SL is significantly and positively correlated with PC1 (r^2=0.19, p=0.0026),

SL does not correlate with PC1 for group-reared (p=0.16) or pair-reared juveniles (p=0.096)

separately.

To better understand how rearing environment affected behavior within the assays that

contributed to the treatment difference in PC1 (Fig 2E), we conducted PCAs for the open field,

social cue investigation, and dominance behavior assays separately. We expanded these analyses

to include all of the measured variables, for the focal and cue fish. The open field test analysis

included focal fish SL and the frequency of entering and time in each zone of the tank. The

social cue investigation included the same measures, as well as the SL of the cue fish. Finally,

the dominance behavior analysis included SL of the focal fish and small cue fish, approaches and

displacements of both fish, and the frequency of entering and time spent in the territory by either

or both fish. For each analysis, we focused on PC1, which differed significantly between group-

and pair-reared juveniles: open field (accounting for 43.4% of the total variance; t=-2.14, p=0.04,

d=0.71, Fig 3A), social cue investigation (37.2%; W=102, p = 0.0032, d=0.92, Fig 3B), and

dominance behavior (29.8%; W=128, p=0.025, d=0.71, Fig 3C). The PC1s were also

significantly and linearly correlated with each other (Fig 3D, open field x social cue: r^2=0.46,

p=5.33e-7; open field x dominance: r^2=0.33, p= 4.69e-5; social cue x dominance: r^2=0.46, p=

4.97e-7, Supplemental Fig 3). See Supplemental Figure 4 for the proportion of the variance

explained by each PC. Vector plots in Supplemental Figure 5 shows the variables that load on

PC1 (and PC2) for each included assay.

Experiment 2

<u>Neural gene expression patterns</u>

Neuroendocrine signaling is a primary mechanism by which early-life experiences are

translated into biological changes. To identify potential mediators of the behavioral effects we

identified, we measured mRNA levels of genes involved in the stress axis and in sex steroid

signaling in the brains of a separate cohort of juveniles. We compared relative expression across

rearing environments (isolation, pairs, groups) and time in rearing environment (1 week, 5

weeks) (Fig 4) using two-way ANOVAs. The sex steroid hormones, ARα and ERα, were the

only genes to have significant interactions between rearing environment and treatment duration.

For ARα, there was no significant effect of treatment ($F_{2,42}$=2.23, p=0.12), but there was a

significant effect of treatment duration ($F_{1,42}$=7.89, p=0.0075) and a significant interaction

($F_{1,42}$=4.95, p=0.032). *Post hoc* analysis of the simple main effects revealed that for the 5 week

juveniles, ARα expression was significantly higher in group-reared fish (t=3.67, p=0.0015).

There were no treatment differences after 1 week ($F_{2,21}$=1.15, p=0.34). In pair-reared juveniles,

ARα expression was significantly higher after 1 week in treatment compared to after 5 weeks

(t=4.72, p=0.00038). There were no treatment duration differences among group-reared juveniles

(t=0.42, p=0.68), and isolates were only analyzed following 1 week in treatment, so comparison

was not possible (Fig 4A). For ERα, there was no significant effect of treatment ($F_{2,42}$=0.73,

p=0.49) or treatment duration ($F_{1,42}$=0.71, p=0.41), but there was a significant interaction

($F_{1,42}$=4.89, p=0.032). *Post hoc* analysis of the simple main effects revealed a pattern similar to

ARα. For juveniles in treatment groups for 5 weeks, ERα expression was significantly higher for

group-reared juveniles (t=2.59, p=0.018). There were no differences after 1 week in treatment

groups ($F_{2,21}$=0.63, p=0.54). In pair reared juveniles, ERα was significantly higher after 1 week

in treatment compared to after 5 weeks (t=3.49, p=0.0031). There were no treatment differences

among group-reared juveniles (t=-0.73, p=0.48) (Fig 4B).

For genes related to the stress response, we found significant main effects for CRF,

GR1a, and GR2. For CRF, there was a significant effect of treatment duration, where week 1

expression was significantly higher than after 5 weeks in treatment $F_{1,42}$=5.77, p=0.021). There

was no effect of treatment ($F_{2,42}$=2.45, p=0.099) and no interaction effect ($F_{1,42}$=0.27, p=0.61)

(Fig 4C). For GR1a, there was a significant effect of treatment ($F_{2,42}$=12.47, p=5.63e-5), and *post*

hoc analysis showed that group-reared juveniles had significantly higher expression than pair-

reared (p=0.0008) and isolated (p=0.00034) juveniles. Expression for pair-reared juveniles was

not significantly different from isolates (p=0.49). There was no main effect of treatment duration

($F_{1,42}$=2.32, p=0.14), and there was no interaction ($F_{1,42}$=0.38, p=0.54) (Fig 4D). For GR2, there

was a significant main effect of treatment duration ($F_{1,42}$=4.10, p=0.049), and similar to CRF,

expression was significantly higher after 1 week in treatment. There was also a significant main

effect of treatment ($F_{2,42}$=3.40, p=0.026); however, *post hoc* analysis revealed that none of the

pairwise differences were significant (group vs. isolates: p=0.20; group vs. pair: p=0.084; pair

vs. isolate: p=0.85). The interaction effect was not significant ($F_{1,42}$=3.25, p=0.079) (Fig 4F).

There were no significant differences for GR1b (treatment: $F_{2,42}$=0.70, p=0.50; treatment

duration: $F_{1,42}$=0.01, p=0.92; interaction: $F_{1,42}$=2.38, p=0.13; Fig 4E) or MR (treatment:

$F_{2,42}$=1.32, p=0.28; treatment duration: $F_{1,42}$=3.28, p=0.077; interaction: $F_{1,42}$=2.91, p=0.095; Fig

4G).

Genes function within regulatory networks, rather than in isolation, and they can affect

each other's expression. A common upstream regulator may also control multiple functional

networks of genes. Because of their known effects on physiology and behavior, these candidate

genes are likely to function in pathways that interact with each other. To quantify how rearing

environment affects gene co-expression, we calculated partial correlation networks (Fig 5).

Partial correlations show the associations between gene pairs, independent of other correlations

in the network. Comparing the group and pair networks (Mantel test: p=0.31), the group and

isolate networks (p=0.61), and the pair and isolate networks (p=0.12) revealed that there was no

evidence that any of these networks were similar to any other.

Discussion

In the present study, we demonstrate that juvenile *A. burtoni* behavior and

neuroendocrine gene expression are both sensitive to early-life social effects. By rearing

juveniles in different social environments—either in a social group or as a pair, both of which

allow individuals to interact freely at all times—we altered the quality and quantity of social

experiences and sensory cues perceived and set individuals along different developmental

trajectories. Behaviorally, the early-life environment shifted juveniles in a predictable manner

along a continuum of a novel behavioral syndrome (i.e., correlated behaviors across contexts, see

below) comprised of open field, social cue investigation, and dominance behaviors (Fig 2, Fig 3)

and affected patterns of subordinate behavior, a critically important social role for young

individuals. In the brain, rearing environment caused significant changes in the expression of key

neuroendocrine genes, including ARα, ERα, and GR1a (Fig 4), and led to striking differences in

patterns of co-expression (Fig 5). The significant effects of treatment duration also provide

important insights into developmental processes (Fig 4). Together, these experiments provide an

essential step towards understanding how developmental plasticity generates the individual

variation in behavior and neuroendocrine function that has fitness and health consequences in

adulthood (e.g., Champagne, 2010; Turecki and Meaney, 2016). Our results also contribute to an

important and growing literature on the impact of early-life social environments beyond parental

interactions (Champagne and Curley, 2005; Taborsky, 2016), using a species that, despite its

prominence in social neuroscience (Fernald and Maruska, 2012; Hofmann, 2003), has rarely

been studied during development (Alvarado, Lenkov, Williams, & Fernald, 2015; Fernald &

Hirata, 1979; Fraley & Fernald, 1982).

<u>Juvenile behavior forms a syndrome affected by early-life social environment</u>

Using a battery of four behavioral assays to gain a comprehensive understanding of

behavioral phenotype, within and across contexts (Fig 1), we discovered that open field, social

cue investigation, and dominance behavior together formed a behavioral syndrome (Fig 3).

Syndromes are a population-level metric defined as the correlation between rank-order

differences between individuals, across contexts and/or over time (Bell, 2007). The presence of a

syndrome indicates consistency in patterns of individual behavior across contexts and/or over

time (Bell, 2007; Sih et al., 2004b, 2004a). Our data suggest that how individuals move around

in space is relevant to the social role they play. Specifically, juveniles that were more active in

the open field test were more likely to be active in the social cue investigation and more

interactive in the dominance assay (Fig 3). Interestingly, behavior from the subordinate assay

does not contribute to the treatment effect or syndrome, likely because subordinate focal

individuals primarily respond to the dominant fish's behavior. To our knowledge, this is the first

behavioral syndrome to be identified in *A. burtoni* at any developmental stage.

Behavior patterns may coalesce into a syndrome due to shared mechanisms (e.g.,

neuroendocrine regulation), early-life experiences that set individuals along developmentally

plastic trajectories, or correlational selection (Bell, 2007; Ketterson and Nolan, Jr., 1999; Stamps,

2003). We found that the behavior of all juveniles was described by the same syndrome,

indicating that how the behaviors are related across experimental contexts (i.e., assays) was

maintained independently of the early-life social environment. Whether an individual was reared

in a group or pair then dictates where along the continuum of the syndrome they fall (Fig 3D).

Pair-reared juveniles appear restricted to one end, whereas group-reared juveniles are represented

along the full range of behavioral variation. That there are group-reared juveniles that

behaviorally resemble the pair-reared individuals suggests there may be social environments

within a group (Saltz et al., 2016) that share key elements with the paired experience. In contrast,

the range of possible social roles seems much more restricted in the paired treatment. To identify

the causal behavioral and/or sensory cues, it will be necessary to conduct detailed observations

of individuals within the rearing environments (Taborsky, 2016). We hypothesize that the

complexity of interactions and/or abundance of social sensory cues in groups cause these

treatment differences (Taborsky, 2016, e.g., Arnold & Taborsky, 2010). Directly quantifying the

range of experience, behavior, and growth within and across early-life environments will be

critical to understanding the nature and magnitude of individual phenotypic variation. It can also

inform more nuanced selection criteria and analysis methods for comparing focal fish across

treatments and tanks than based on size or size ratios alone, as we did in this study.

Activity and social interaction are common components of syndromes in other species,

along with bold-shy and proactive-reactive behaviors (Bell, 2007; Conrad et al., 2011; Groothuis

and Carere, 2005; Koolhaas et al., 1999; Sih et al., 2004b; Verbeek et al., 1994). For example,

large juvenile brown trout are more active and aggressive (Näslund and Johnsson, 2016), similar

to our results. Activity-aggression syndromes are also found in a number of other fish species

(reviewed in Conrad et al., 2011). For *A. burtoni* juveniles, locomotor activity and social

interaction may be causally related. First, active individuals may encounter conspecifics more

frequently and, as a result, initiate more interactions. Second, juvenile social interactions appear

to be prosocial in that they increase the likelihood of future proximity and interaction. In the

dominance behavior assay, approaches and displacements for both the focal and subordinate cue

fish load in the same direction on PC1. Correlation analysis (data not shown) confirms that, as

one member of the pair initiates social interactions, the other member also initiates, potentially

leading to more activity. This may be beneficial by increasing shoaling and reducing the risk of

predation. Interestingly, adult dominance behavior does not lead to a prosocial response in

subordinates, suggesting that although social behavior appears similar across life history stages

(Fernald and Hirata, 1979; Fraley and Fernald, 1982), there are important differences.

<u>Size plays a secondary role in determining juvenile behavioral phenotype</u>

Size is central to understanding the effects of the early-life social environment. Group-

reared juveniles were larger than those reared in pairs, which is consistent with previous work

showing growth is socially regulated in both juveniles and adults (Fraley and Fernald, 1982;

Hofmann et al., 1999). Adult *A. burtoni* are also highly sensitive to size during social interactions

(Alcazar et al., 2014; Weitekamp & Hofmann, 2017); therefore, size differences could cause

differences in behavior. In this study, however, the effect of the early social environment appears

larger and more complex than size alone. First, the PCA of behavior from all four assays shows

that focal fish SL contributes only moderately to the significant treatment difference for PC1 (Fig

2E), as many other variables load much more strongly on PC1 (i.e., open field, social cue

investigation, and dominance behaviors) (see also: Supplemental Fig 5). Second, SL is the

strongest contributing variable for PC6, which differs significantly between group- and pair-

reared juveniles (Fig 2F). The proportion of the variance described by PC6 (5%) compared to

PC1 (43.3%) suggests that size contributes relatively less to the overall treatment effect than

behaviors in the open field, social cue investigation, and dominance behavior assays. This is

further supported by the finding that in a PCA excluding focal fish SL, PC1 still differs

significantly between group- and pair-reared juveniles. In this analysis, PC1 is not associated

with SL for either group- or pair-reared juveniles, suggesting size does not drive behavior. The

significant, positive association between PC1 and SL for all juveniles results from group-reared

juveniles being larger than pair-reared juveniles. Third, SL is also not associated with behavior in

any of the four behavior assays (Supplemental Table 2). Finally, the group-reared juveniles that

fall within the range of pair-reared juveniles along the continuum of the behavioral syndrome

(i.e., high PCA scores, Fig 3) are not the smallest individuals. Together, this evidence suggests

that size is secondary in understanding early-life effects on behavior. In future studies, it will be

important to test how individual behavior changes over time in relation to both size and

developmental stage, which can be decoupled from chronological age in fish (Jonsson and

Jonsson, 2014).

<u>Early-life social experience affects social dynamics when focal juveniles are subordinate</u>

Developmental plasticity can shift behavior in ways that ultimately benefit fitness (Smith

and Blumstein, 2008), in part because social behavior has direct consequences for reproductive

success (Wilson, 1980, e.g., Henry et al., 2013; Robbins et al., 2007; Young et al., 2006). A

majority (64%) of studies show that experimentally increasing the frequency, diversity, or

complexity of early-life social experiences enhances social skills or competence (Taborsky,

2016). For example, juvenile *N. pulcher* cichlids reared with brood helpers demonstrated more

context-appropriate behavior when establishing status, integrating into novel groups, and

competing for a resource (Arnold and Taborsky, 2010; Fischer et al., 2015; Taborsky et al.,

2013, 2012). We have no evidence yet of an advantage for group-reared juveniles; however,

juveniles appear fill the subordinate role differently based on rearing environment, as well as

social status experience. While nearly all focal fish successfully established themselves as

subordinate (88%) in the assay, and there were no treatment differences in approaches or

displacements, there was a significantly larger asymmetry in agonistic efficiency for pair-reared

juveniles. There was also a trend for pair-reared juveniles to submit more readily (measured as

large fish agonistic efficiency). Status relationships are defined by asymmetrical agonistic

displays (Drews, 1993); therefore, pair-reared juveniles may behave more submissively.

We also found that the larger juveniles sampled from the groups and pairs, which we are

confident accrued more dominance experience during development given the importance of size

for juvenile (and adult, Weitekamp and Hofmann, 2017) social interactions (this study), differed

in their patterns of behavior compared to the smaller juveniles. Behaviors from the subordinate

assay load on PC2 (16.4% of variance, Fig 2E), and there is a trend for PC2 to differ between the

larger and smaller sampled fish (Fig 2B). PC5 (6.6% of variance) differs significantly between

the larger and smaller fish. A variety of behaviors load on PC5, including activity in the open

field and social cue investigation assays, suggesting that space use is also influenced by status

experience and/or relative size within a rearing environment. Overall, the subordinate role is

critically important for juveniles because all juveniles will enter adult communities as

subordinates. It will be necessary to measure behavior and reproductive success of these

juveniles once they are adults in order to determine whether these phenotypes persist or if one is

more successful than another (Pradhan, Solomon-Lane, & Grober, 2015).

Early-life social environment and treatment duration affect neuroendocrine gene expression

We have shown that early-life environments can determine where individuals will fall

along the continuum of a newly discovered behavioral syndrome, which raises questions about

the underlying mechanisms (e.g., pleiotropic genes and/or neuroendocrine regulation). The

behavioral effects we detect as a result of the early-life social environment suggest important

variation in the underlying neural regulatory mechanisms. Neuroendocrine stress and sex steroid

signaling are likely sites of developmental plasticity in *A. burtoni* because they are sensitive to

early-life effects (Champagne & Curley, 2005; Shepard et al., 2009), translate environmental

conditions and experiences into biological responses (Crespi & Denver, 2005; Wingfield et al.,

1990), and regulate behavior (Adkins-Regan, 2009; Solomon-Lane, Crespi, & Grober, 2013).

We focused on steroid hormone nuclear receptors, with the addition of CRF, specifically because

they regulate the transcription of target genes with a diversity of physiological and behavioral

roles (Rochette-Egly, 2005). We found that both the early-life social environment and treatment

duration—which corresponds to age, in this study—had a significant effect on gene expression in

whole brain. GR1a was the only gene to respond exclusively to treatment, while CRF and GR2

changed significantly over time. Early-life environment and treatment duration interacted to

affect the expression of sex steroid hormone receptors ARα and ERα. Finally, although GR1b

and MR expression varied across individuals, these genes were not significantly affected by

treatment or treatment duration. Factors that we did not measure here (e.g., social status, body

size, sex), including individual behavior and position along the behavioral syndrome, are also

likely to contribute to important variation in gene expression.

The HPA/I axis has a highly-conserved role in responding to early-life environments

(Crespi and Denver, 2005). Our results suggest that developmental plasticity can "tune" the HPI

axis in nuanced ways via changes in the density and distribution of different receptors and by

affecting circulating glucocorticoid levels (Bernier et al., 2009), over developmental time (e.g.,

CRF, GR2, Fig 4C, E) and in response to different environments (e.g., GR1a, Fig 4D). Many

teleosts, including *A. burtoni*, have four glucocorticoid receptors: MR, GR1a, GR1b, and GR2.

Receptor 1 has subtypes 1a and 1b, which differ by a nine amino acid insertion between the two

zinc fingers in the DNA-binding domain (Bury, 2017; Greenwood et al., 2003; Korzan et al.,

2014). These receptors differ substantially in their affinity for cortisol. In adult *A. burtoni*, MR is

100-fold more sensitive to cortisol than the GRs and is likely to be occupied with cortisol at

basal levels (in fish and tetrapods). GR2 has the next highest sensitivity, followed by GR1a, then

GR1b (Arterbery et al., 2011; Bury, 2017; Greenwood et al., 2003). Changes in HPA/I axis

function typically manifest as altered baseline levels of circulating glucocorticoids, a higher or

lower glucocorticoid 'peak' in response to an acute stressor, and/or altered efficiency of the

negative feedback loop that returns the system to baseline. Negative feedback, in particular, is

regulated by neural GR expression (Bernier et al., 2009; Bury, 2017; Denver, 2009; Kiilerich et

al., 2018; Wendelaar Bonga, 1997) and can be affected by early-life experience (Champagne and

Curley, 2005; Francis et al., 1999).

Consistent with the distinct roles for different components of the stress axis (Greenwood

et al., 2003), our results show differences in expression patterns across HPI axis candidate genes

(Fig 4C-G). GR1, specifically, appears to respond to the early-life social environment in *A.*

burtoni and other teleost species (Fokos et al., 2017; Nyman et al., 2018, 2017; Taborsky et al.,

2013). In the group-living cichlid *N. pulcher*, for example, increased early-life social complexity

led to altered GR1 expression, but not GR2 or MR expression, in whole brain and telencephalon

(Nyman et al., 2018, 2017; Taborsky et al., 2013). In *A. burtoni*, higher expression of GR1a in

group-reared juveniles (Fig 4D) might increase sensitivity to cortisol and result in more efficient

negative feedback, making these individuals less susceptible to stress. Alternatively, given that

A. burtoni naturally live in groups (Fernald and Hirata, 1977), paired rearing or isolation may

actually decrease efficiency. Juvenile stress physiology should be tested directly because

negative feedback mechanisms are complex and involve multiple receptors (Bury, 2017;

Kiilerich et al., 2018). Overall, little is known about the differential roles of GR1a and GR1b,

and the differences that have been demonstrated appear to be species-specific (Bury, 2017). For

A. burtoni, sensitivity to the early-life social environment may be a defining difference (Fig 4D,

F). That GR2 and MR also do not respond to the early environment may be consistent with their

roles in baseline glucocorticoid signaling rather than the stress response (Greenwood et al.,

2003), although whole brain expression of GR2 (and CRF) is lower adult subordinate males

compared to dominants (Chen and Fernald, 2008). Finally, the stress axis undergoes important

changes throughout development (Alsop and Vijayan, 2008; Barry et al., 1995; Jeffrey and

Gilmour, 2016; Tsalafouta et al., 2018), and lower levels of CRF and GR2 after 5 weeks (Fig 4C,

E) could indicate a developmental shift towards lower stress axis activity. Alternatively,

familiarity with or predictability of a social environment (e.g., treatment duration) could shift

HPI axis function. Future research testing these HPI axis hypotheses promises to uncover

important mechanisms of early-life effects on neuroendocrine and behavioral development.

The sex steroid hormone receptors ARα and ERα were unique among our candidate

genes in that effect of rearing environment on gene expression was mediated by treatment

duration. These genes are also not a part of the HPI axis. For both receptors, expression in pair-

reared juveniles was significantly lower after 5 weeks in the rearing environments compared to

pair-reared juveniles after 1 week and group-reared juveniles after both 1 week and 5 weeks (Fig

4A, B). Compared to the HPA/I axis, less is known about sex steroid hormones in the context of

early-life effects, but there are multiple, non-mutually exclusive mechanisms that could explain

these expression patterns. First, it is well-established that the HPI axis interacts with the

hypothalamic-pituitary-gonadal axis that regulates reproduction, in part through neural ARs and

ERs (Acevedo-Rodriguez et al., 2018; Huffman et al., 2012; Schreck, 2010). While we do not

expect HPI axis plasticity to entirely drive the changes in ARα and ERα expression, interactions

are likely between these important neuroendocrine axes (see below, Fig 5). Second, early-life

social experiences can exert lasting changes in sex steroid hormone receptor expression via

epigenetic mechanisms. In rats, for example, ERα in the medial POA is critical to the

neuroendocrine regulation of maternal licking and grooming. The rates of maternal care received

by female pups subsequently affects their future maternal behavior. The mechanism for this

early-life maternal effect has been described in detail and involves brain region-specific

epigenetic methylation of the ERα promoter (Cameron et al., 2008). Similar epigenetic

mechanisms may regulate ARα and ERα (as well as GR, Turecki and Meaney, 2016) in juvenile

A. burtoni, such that epigenetic marks accrue over time in particular early-life social

environments. In our study, expression differences were evident after 5 weeks but not yet after 1

week (Fig 4A, B).

Finally, ARα and ERα are found throughout the social decision-making network, a

highly-conserved set of brain regions that, together, are involved in the regulation of social

behavior across vertebrates, including *A. burtoni* (O'Connell and Hofmann, 2012b, 2011). Sex

steroid hormone receptors regulate and respond to social behavior and context (Burmeister et al.,

2007; Maruska, 2015; O'Connell and Hofmann, 2012a); therefore, the expression patterns in

juveniles could reflect social interactions. ARα and ERα mRNA may increase early in

development to facilitate sociality, possibly along with other relevant neuromodulators and

receptors. After 5 weeks in pairs, the decrease in transcription (Fig 4A, B) may reflect a less

dynamic social decision-making network as a consequence of a social environment that is highly

predictable. Conversely, expression for juveniles in groups, the most complex environment in

this study, remains high over time. Expression for isolated juveniles, an environment absent of

social stimuli, is closest to the expression of pair-reared juveniles after 5 weeks. It is noteworthy

that all of the candidate genes show a similar pattern (Fig 4): the expression of pair-reared

juveniles after 5 week is the lowest compared to other treatments and time points, and the most

similar group is isolated juveniles after 1 week. Future work is needed to understand the

functional significance of this downregulation.

An important consideration in interpreting these results, and the co-expression networks

below, is that gene expression was measured in whole brain. Although the brain is a

heterogeneous tissue made up multiple cell types (e.g., neurons, glia) and regions with distinct

functionality, we chose this approach because we did not have an *a priori* expectation as to

which brain regions or cell types might be the most critical to examine in our study. We

recognize that important variation in gene expression might not be detected using this approach;

therefore, future research should use approaches that allow for increased spatial resolution (see

below), as well as unbiased (rather than candidate) gene expression analysis (e.g., via RNA-Seq).

A genome-scale analysis of expression can provide insight into large numbers of genes

simultaneously and suggest novel candidate pathways. However, recent analyses have

demonstrated that brain region-specific, or even cell type-specific, gene expression patterns can

be inferred from bulk tissue samples (e.g., whole brain) (Kelley et al., 2018).

<u>Complex co-expression of stress and sex steroid signaling by the early-life social environment</u>

Neuroendocrine systems are dynamic and interact on multiple biological levels (e.g.,

Acevedo-Rodriguez et al., 2018), including within gene regulatory networks (e.g., Huffman et

al., 2012; Korzan, Fernald, & Grone, 2014; O'Connell & Hofmann, 2012); therefore, the

expression of other genes can also contribute to the variation in a gene of interest. Based on their

co-localization in the POA of *A. burtoni* (Korzan et al., 2014), co-localization and correlation in

other species (e.g., Meyer & Korz, 2013), and overlapping physiological effects (Crespi &

Denver, 2005; Wingfield et al., 1990), the neuroendocrine pathways represented by our

candidate genes are likely to functionally interact. We identified striking differences in co-

expression networks among juveniles reared in different environments. Expression was highly

correlated in pair-reared juveniles (Fig 5A), such that every candidate gene was significantly

correlated with at least two others. At the center of the network, ARα shares five significant

connections. The two sex steroid hormone genes (ARα, ERα) are also integrated with the stress

axis genes, which form distinct smaller networks: CRF-GR1a-GR1b and GR2-MR. In contrast,

group-reared juveniles have only one significant partial correlation between ERα and GR1b, a

connection that is not present in the pair-reared network (Fig 5B). There are no significant partial

correlations for isolated juveniles, suggesting that the neuroendocrine regulatory network is

dysregulated, possibly due to isolation acting as a stressor (Galhardo and Oliveira, 2014). These

network differences, together with other relevant genes not included in our candidate analysis,

might underlie the behavioral differences we identified in the behavioral syndrome, subordinate

behavior, or more broadly related to stress response. The differential co-regulation could also

serve to make behavior more similar in the face of other neural differences caused by rearing

environment, as is the case for some neural sex differences and behavior (De Vries, 2004). These

hypotheses can be tested directly using central pharmacological manipulation.

<u>Integrating the effects of early-life social environments on behavior and brain</u>

Our work demonstrates that early-life social environments shape behavioral phenotype

and neuroendocrine gene expression in powerful ways for *A. burtoni* juveniles. In the present

study, we quantified behavior and gene expression in separate experiments in order to focus on

different developmental time points. Taken together, our results allow us to generate strong

hypotheses about the mechanisms, consequences, and developmental time course of early-life

social effects. By focusing on early time points for gene expression—after 1 and 5 weeks in the

social environments—we aimed to identify highly sensitive components of these neuroendocrine

systems. Based on our results, we expect the HPI axis via GR1a expression to respond rapidly to

different social environments. In contrast, the treatment differences in ARα and ERα emerge

over time. After 8-10 weeks, when we analyzed behavior, we hypothesize that altered stress

physiology aligns with the behavior patterns sensitive to early-life effects, together forming a

specific kind of syndrome called a coping style. Individuals range in coping style from proactive

to reactive, such that proactive copers are more active, aggressive, and less responsive to stress

(i.e., lower baseline glucocorticoid levels, faster negative feedback) than reactive copers

(Koolhaas et al., 1999). As juveniles approach reproductive maturity (as early as 12 weeks old,

Fraley and Fernald, 1982), we expect social environment and age to continue to interact for ARα

and ERα. Maturation is socially regulated, including by the early-life social environment;

therefore, we predict that expression will reflect ongoing social demands, as well as emerging

reproductive behavior (Fraley and Fernald, 1982). This work can begin to address the fact that

across species, remarkably little is known about the mechanisms that shape the ontogeny of

behavior (Taborsky, 2016).

Understanding the full scope and consequences of early-life effects ultimately requires

measuring brain and behavior in the same individuals, throughout development and into

adulthood. Our results suggest that brain regions that express GR1a, ARα, and ERα (Greenwood

et al., 2003; Korzan et al., 2014; Munchrath and Hofmann, 2010), along with brain regions of the

social decision-making network (O'Connell and Hofmann, 2012b), are likely to be sensitive to

early-life effects and could cause the observed changes in behavior. Interestingly, the POA—a

critical node in the social decision-making network (O'Connell and Hofmann, 2012b)—contains

GR1a, GR1b, GR2, MR, ARα, and ERα in adult *A. burtoni* (Korzan et al., 2014; Munchrath and

Hofmann, 2010). Additional nodes, such as the hippocampus and amygdala are also likely sites

of overlap. These brain regions are important in spatial cognition and emotional processing,

respectively, and are central to HPI axis negative feedback (Denver, 2009). Interactions between

the HPI axis and sex steroid hormone signaling, including in the POA, could be a mechanism for

the social regulation of development (Fraley and Fernald, 1982; Korzan et al., 2014; Solomon-

Lane et al., 2013; Wada, 2008). Overall, this research can uncover the neuroendocrine

mechanisms by which early-life social experience gives rise to individual variation in adults,

which is critical to understanding subsequent disparities in fitness and health.

Acknowledgements

We thank Savannah Clapp, Pamela Del Valle, and Najah Hussain for assistance with data

collection and fish maintenance and care. We thank Dr. Becca Young for helpful comments on

earlier versions of this manuscript and members of the Hofmann Lab for discussion and

feedback.

Funding

This work was supported by NSF grant IOS-1354942 to HAH and the BEACON Center for the

Declarations of interest: None.
809

References

Acevedo-Rodriguez, A., Kauffman, A.S., Cherrington, B.D., Borges, C.S., Roepke, T.A.,
Laconi, M., 2018. Emerging insights into hypothalamic-pituitary-gonadal axis regulation
and interaction with stress signalling. J. Neuroendocrinol. 1–11.
https://doi.org/10.1111/jne.12590
Adkins-Regan, E., 2009. Neuroendocrinology of Social Behavior. ILAR J. 50, 5–14.
https://doi.org/10.1093/ilar.50.1.5
Alcazar, R.M., Hilliard, A.T., Becker, L., Bernaba, M., Fernald, R.D., 2014. Brains over brawn:
experience overcomes a size disadvantage in fish social hierarchies. J. Exp. Biol. 217,
1462–1468. https://doi.org/10.1242/jeb.097527
Alsop, D., Vijayan, M.M., 2008. Development of the corticosteroid stress axis and receptor
expression in zebrafish. AJP Regul. Integr. Comp. Physiol. 294, R711–R719.
https://doi.org/10.1152/ajpregu.00671.2007
Alvarado, S.G., Lenkov, K., Williams, B., Fernald, R.D., 2015. Social crowding during
development causes changes in GnRH1 DNA methylation. PLoS One 10, 1–13.
https://doi.org/10.1371/journal.pone.0142043
Arnold, C., Taborsky, B., 2010. Social experience in early ontogeny has lasting effects on social
skills in cooperatively breeding cichlids. Anim. Behav. 79, 621–630.
https://doi.org/10.1016/j.anbehav.2009.12.008
Arterbery, A.S., Fergus, D.J., Fogarty, E.A., Mayberry, J., Deitcher, D.L., Kraus, W.L., Bass,
A.H., 2011. Evolution of ligand specificity in vertebrate corticosteroid receptors. BMC
Evol. Biol. 11, 1–15. https://doi.org/10.1186/1471-2148-11-14
Banerjee, S.B., Arterbery, A.S., Fergus, D.J., Adkins-Regan, E., 2012. Deprivation of maternal
care has long-lasting consequences for the hypothalamic-pituitary-adrenal axis of zebra
finches. Proc. R. Soc. B Biol. Sci. 279, 759–766. https://doi.org/10.1098/rspb.2011.1265
Barry, T.P., Malison, J.A., Held, J.A., Parrish, J.J., 1995. Ontogeny of the Cortisol Stress
Response in Larval Rainbow Trout. Gen. Comp. Endocrinol. 97, 57–65.
Bell, A.M., 2007. Future directions in behavioural syndromes research. Proc. R. Soc. B Biol. Sci.
274, 755–761. https://doi.org/10.1098/rspb.2006.0199
Benjamini, Y., Hochberg, Y., 1995. Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. J. R. Stat. Soc. Ser. B.
https://doi.org/10.2307/2346101
Bennett, D. a, Schneider, J. a, Tang, Y., Arnold, S.E., Wilson, R.S., 2006. The effect of social
networks on the relation between Alzheimer's disease pathology and level of cognitive
function in old people: a longitudinal cohort study. Lancet. Neurol. 5, 406–12.
https://doi.org/10.1016/S1474-4422(06)70417-3
Bernier, N.J., Flik, G., Klaren, P.H.M., 2009. Regulation And Contribution Of Corticotropic,
Melanotropic And Thyrotropic Axes To The Stress Response In Fishes, 1st ed, Fish
Physiology. Elsevier Inc. https://doi.org/10.1016/S1546-5098(09)28006-X
Bonuti, R., Morato, S., 2018. Proximity as a predictor of social behavior in rats. J. Neurosci.
Methods 293, 37–44. https://doi.org/10.1016/j.jneumeth.2017.08.027
Branchi, I., D'Andrea, I., Fiore, M., Di Fausto, V., Aloe, L., Alleva, E., 2006. Early Social
Enrichment Shapes Social Behavior and Nerve Growth Factor and Brain-Derived
Neurotrophic Factor Levels in the Adult Mouse Brain. Biol. Psychiatry 60, 690–696.
https://doi.org/10.1016/j.biopsych.2006.01.005

Branchi, I., Santarelli, S., D'Andrea, I., Alleva, E., 2013. Not all stressors are equal: Early social
enrichment favors resilience to social but not physical stress in male mice. Horm. Behav.
63, 503–509. https://doi.org/10.1016/j.yhbeh.2013.01.003
Brown, G.R., Spencer, K.A., 2013. Steroid hormones, stress and the adolescent brain: A
comparative perspective. Neuroscience 249, 115–128.
https://doi.org/10.1016/j.neuroscience.2012.12.016
Buist, K.L., Deković, M., Prinzie, P., 2013. Sibling relationship quality and psychopathology of
children and adolescents: A meta-analysis. Clin. Psychol. Rev. 33, 97–106.
https://doi.org/10.1016/j.cpr.2012.10.007
Burmeister, S.S., Kailasanath, V., Fernald, R.D., 2007. Social dominance regulates androgen and
estrogen receptor gene expression. Horm. Behav. 51, 164–170.
https://doi.org/10.1016/j.yhbeh.2006.09.008
Bury, N.R., 2017. The evolution, structure and function of the ray finned fish (Actinopterygii)
glucocorticoid receptors. Gen. Comp. Endocrinol. 251, 4–11.
https://doi.org/10.1016/j.ygcen.2016.06.030
Cachat, J., Stewart, A., Grossman, L., Gaikwad, S., Kadri, F., Chung, K.M., Wu, N., Wong, K.,
Roy, S., Suciu, C., Goodspeed, J., Elegante, M., Bartels, B., Elkhayat, S., Tien, D., Tan, J.,
Denmark, A., Gilder, T., Kyzar, E., Dileo, J., Frank, K., Chang, K., Utterback, E., Hart, P.,
Kalueff, A. V., 2010. Measuring behavioral and endocrine responses to novelty stress in
adult zebrafish. Nat. Protoc. 5, 1786–1799. https://doi.org/10.1038/nprot.2010.140
Cameron, N.M., Shahrokh, D., Del Corpo, A., Dhir, S.K., Szyf, M., Champagne, F.A., Meaney,
M.J., 2008. Epigenetic programming of phenotypic variations in reproductive strategies in
the rat through maternal care. J. Neuroendocrinol. 20, 795–801.
https://doi.org/10.1111/j.1365-2826.2008.01725.x
Champagne, F.A., 2010. Early adversity and developmental Outcomes: Interaction between
Genetics, Epigenetics, and social experiences across the Life Span. Perspect. Psychol. Sci.
5, 564–574. https://doi.org/10.1177/1745691610383494
Champagne, F.A., Curley, J.P., 2005. How social experiences influence the brain. Curr. Opin.
Neurobiol. 15, 704–709. https://doi.org/10.1016/j.conb.2005.10.001
Champagne, F.A., Meaney, M.J., 2007. Transgenerational effects of social environment on
variations in maternal care and behavioral response to novelty. Behav. Neurosci. 121, 1353–
1363. https://doi.org/10.1037/0735-7044.121.6.1353
Champagne, F.A., Weaver, I.C.G., Diorio, J., Sharma, S., Meaney, M.J., 2003. Natural
Variations in Maternal Care Are Associated with Estrogen Receptor α Expression and
Estrogen Sensitivity in the Medial Preoptic Area. Endocrinology 144, 4720–4724.
https://doi.org/10.1210/en.2003-0564
Chen, C.-C., Fernald, R.D., 2008. Sequences, expression patterns and regulation of the
corticotropin-releasing factor system in a teleost. Gen. Comp. Endocrinol. 157, 148–155.
https://doi.org/10.1016/j.ygcen.2008.04.003
Conrad, J.L., Weinersmith, K.L., Brodin, T., Saltz, J.B., Sih, A., 2011. Behavioural syndromes in
fishes: A review with implications for ecology and fisheries management. J. Fish Biol. 78,
395–435. https://doi.org/10.1111/j.1095-8649.2010.02874.x
Creel, S.R., Dantzer, B., Goymann, W., Rubenstein, D.R., 2013. The ecology of stress: effects of
the social environment. Funct. Ecol. 27, 66–80. https://doi.org/10.1111/j.1365-
2435.2012.02029.x
Crespi, E.J., Denver, R.J., 2005. Ancient origins of human developmental plasticity. Am. J.

Hum. Biol. 17, 44–54. https://doi.org/10.1002/ajhb.20098

D'Andrea, I., Alleva, E., Branchi, I., 2007. Communal nesting, an early social enrichment,
affects social competences but not learning and memory abilities at adulthood. Behav. Brain
Res. 183, 60–66. https://doi.org/10.1016/j.bbr.2007.05.029

Davis, M.R., Fernald, R.D., 1990. Social control of neuronal soma size. J. Neurobiol. 21, 1180–
1188. https://doi.org/10.1002/neu.480210804

De Vries, G.J., 2004. Minireview: Sex Differences in Adult and Developing Brains:
Compensation, Compensation, Compensation. Endocrinology 145, 1063–1068.
https://doi.org/10.1210/en.2003-1504

Denver, R.J., 2009. Structural and functional evolution of vertebrate neuroendocrine stress
systems. Ann. N. Y. Acad. Sci. 1163, 1–16. https://doi.org/10.1111/j.1749-
6632.2009.04433.x

Desjardins, J.K., Hofmann, H.A., Fernald, R.D., 2012. Social Context Influences Aggressive and
Courtship Behavior in a Cichlid Fish. PLoS One 7, e32781.
https://doi.org/10.1371/journal.pone.0032781

Desjardins, J.K., Klausner, J.Q., Fernald, R.D., 2010. Female genomic response to mate
information. Proc. Natl. Acad. Sci. U. S. A. 107, 21176–21180.
https://doi.org/10.1073/pnas.1010442107

Drews, C., 1993. The concept and definition of dominance in animal behaviour. Behaviour 125,
283–313. https://doi.org/10.1163/156853993X00290

Fernald, R.D., 2012. Social Control of the Brain. Annu. Rev. Neurosci. 35, 133–151.
https://doi.org/10.1146/annurev-neuro-062111-150520

Fernald, R.D., Hirata, N.R., 1979. The Ontogeny of Social behavior and body coloration in the
african cichlid fish haplochromis burtoni. Zeitschrif für Tierpsychologie.
https://doi.org/10.1111/j.1439-0310.1979.tb01025.x

Fernald, R.D., Hirata, N.R., 1977. Field study of Haplochromis burtoni: Quantitative behavioural
observations. Anim. Behav. 25, 964–975. https://doi.org/10.1016/0003-3472(77)90048-3

Fernald, R.D., Maruska, K.P., 2012. Social information changes the brain. Proc. Natl. Acad. Sci.
109, 17194–17199. https://doi.org/10.1073/pnas.1202552109

Fischer, S., Bessert-Nettelbeck, M., Kotrschal, A., Taborsky, B., 2015. Rearing-Group Size
Determines Social Competence and Brain Structure in a Cooperatively Breeding Cichlid.
Am. Nat. 186, 123–140. https://doi.org/10.1086/681636

Fokos, S., Pavlidis, M., Yiotis, T., Tsalafouta, A., Papandroulakis, N., Dermon, C.R., 2017.
Early life low intensity stress experience modifies acute stress effects on juvenile brain cell
proliferation of European sea bass (D. Labrax). Behav. Brain Res. 317, 109–121.
https://doi.org/10.1016/j.bbr.2016.09.026

Fox, H.E., White, S. a, Kao, M.H., Fernald, R.D., 1997. Stress and dominance in a social fish. J.
Neurosci. 17, 6463–6469.

Fraley, N.B., Fernald, R.D., 1982. Social control of developmental rate in the African cichlid,
Haplochromis burtoni. Z.Tierpsychol. 60, 66–82. https://doi.org/10.1111/j.1439-
0310.1982.tb01077.x

Francis, D.D., Caldji, C., Champagne, F.A., Plotsky, P.M., Meaney, M.J., 1999. The role of
corticotropin-releasing factor–norepinephrine systems in mediating the effects of early
experience on the development of behavioral and endocrine responses to stress. Biol.
Psychiatry 46, 1153–1166. https://doi.org/10.1016/S0006-3223(99)00237-1

Galhardo, L., Oliveira, R.F., 2014. The effects of social isolation on steroid hormone levels are

947 modulated by previous social status and context in a cichlid fish. Horm. Behav. 65, 1–5.
948 https://doi.org/10.1016/j.yhbeh.2013.10.010
949 Gilbert, S.F., 2001. Ecological Developmental Biology: Developmental Biology Meets the Real
950 World. Dev. Biol. 233, 1–12. https://doi.org/10.1006/dbio.2001.0210
951 Goodson, J.L., 2005. The vertebrate social behavior network: Evolutionary themes and
952 variations. Horm. Behav. 48, 11–22. https://doi.org/10.1016/j.yhbeh.2005.02.003
953 Greenwood, A.K., Butler, P.C., White, R.B., Demarco, U., Pearce, D., Fernald, R.D., 2003.
954 Multiple corticosteroid receptors in a teleost fish: Distinct sequences, expression patterns,
955 and transcriptional activities. Endocrinology 144, 4226–4236.
956 https://doi.org/10.1210/en.2003-0566
957 Groothuis, T.G.G., Carere, C., 2005. Avian personalities: Characterization and epigenesis.
958 Neurosci. Biobehav. Rev. 29, 137–150. https://doi.org/10.1016/j.neubiorev.2004.06.010
959 Grosenick, L., Clement, T.S., Fernald, R.D., 2007. Fish can infer social rank by observation
960 alone. Nature 445, 429–432. https://doi.org/10.1038/nature05646
961 Henry, M.D., Hankerson, S.J., Siani, J.M., French, J.A., Dietz, J.M., 2013. High rates of
962 pregnancy loss by subordinates leads to high reproductive skew in wild golden lion
963 tamarins (Leontopithecus rosalia). Horm. Behav. 63, 675–683.
964 https://doi.org/10.1016/j.yhbeh.2013.02.009
965 Heule, C., Göppert, C., Salzburger, W., Böhne, A., 2014. Genetics and timing of sex
966 determination in the East African cichlid fish Astatotilapia burtoni. BMC Genet. 15, 1–17.
967 https://doi.org/10.1186/s12863-014-0140-5
968 Hofmann, H.A., 2003. Functional genomics of neural and behavioral plasticity. J. Neurobiol. 54,
969 272–282. https://doi.org/10.1002/neu.10172
970 Hofmann, H.A., Benson, M.E., Fernald, R.D., 1999. Social status regulates growth rate:
971 consequences for life-history strategies. Proc. Natl. Acad. Sci. U. S. A. 96, 14171–6.
972 https://doi.org/10.1073/pnas.96.24.14171
973 Hofmann, H.A., Fernald, R.D., 2001. What cichlids tell us about the social regulation of brain
974 and behavior. J. Aquaric. Aquat. Sci. 9, 17–31.
975 Huffman, L.S., Mitchell, M.M., O'Connell, L.A., Hofmann, H.A., 2012. Rising StARs:
976 Behavioral, hormonal, and molecular responses to social challenge and opportunity. Horm.
977 Behav. 61, 631–641. https://doi.org/10.1016/j.yhbeh.2012.02.016
978 Jeffrey, J.D., Gilmour, K., 2016. Correction: Programming of the hypothalamic-pituitary-
979 interrenal axis by maternal social status in zebrafish (*Danio rerio*). J. Exp. Biol. 219,
980 2963–2963. https://doi.org/10.1242/jeb.148452
981 Jonsson, B., Jonsson, N., 2014. Early environment influences later performance in fishes. J. Fish
982 Biol. 85, 151–188. https://doi.org/10.1111/jfb.12432
983 Kaiser, S., Kruijver, F.P.M., Swaab, D.F., Sachser, N., 2003. Early social stress in female guinea
984 pigs induces a masculinization of adult behavior and corresponding changes in brain and
985 neuroendocrine function. Behav. Brain Res. 144, 199–210. https://doi.org/10.1016/S0166-
986 4328(03)00077-9
987 Kasumovic, M.M., Brooks, R.C., 2011. It's All Who You Know: The Evolution Of Socially
988 Cued Anticipatory Plasticity As A Mating Strategy. Q. Rev. Biol. 86, 181–197.
989 https://doi.org/10.1086/661119
990 Kelley, K.W., Nakoa-Inoue, H., Molofsky, A. V., Oldham, M.C., 2018. Variation among intact
991 tissue samples revelas the core transcriptional features of human CNS cell classes. Nat.
992 Neurosci. 21, 1171–1184.

Ketterson, E.D., Nolan, Jr., V., 1999. Adaptation, Exaptation, and Constraint: A Hormonal
Perspective. Am. Nat. 154, S4–S25. https://doi.org/10.1086/303280
Kiilerich, P., Servili, A., Péron, S., Valotaire, C., Goardon, L., Leguen, I., Prunet, P., 2018.
Regulation of the corticosteroid signalling system in rainbow trout HPI axis during
confinement stress. Gen. Comp. Endocrinol. 258, 184–193.
https://doi.org/10.1016/j.ygcen.2017.08.013
Koolhaas, J.M., Korte, S.M., De Boer, S.F., Van Der Vegt, B.J., Van Reenen, C.G., Hopster, H.,
De Jong, I.C., Ruis, M. a W., Blokhuis, H.J., 1999. Coping styles in animals: Current status
in behavior and stress- physiology. Neurosci. Biobehav. Rev. 23, 925–935.
https://doi.org/10.1016/S0149-7634(99)00026-3
Korzan, W.J., Fernald, R.D., Grone, B.P., 2014. Social regulation of cortisol receptor gene
expression. J. Exp. Biol. 3221–3228. https://doi.org/10.1242/jeb.104430
Langenhof, M.R., Komdeur, J., 2018. Why and how the early-life environment affects
development of coping behaviours. Behav. Ecol. Sociobiol. 72, 34.
https://doi.org/10.1007/s00265-018-2452-3
Lowry, C. a., Moore, F.L., 2006. Regulation of behavioral responses by corticotropin-releasing
factor. Gen. Comp. Endocrinol. 146, 19–27. https://doi.org/10.1016/j.ygcen.2005.12.006
Lummaa, V., Clutton-Brock, T., 2002. Early development, survival and reproduction in humans.
Trends Ecol. Evol. 17, 141–147. https://doi.org/10.1016/s0169-5347(01)02414-4
Maruska, K.P., 2015. Social Transitions Cause Rapid Behavioral and Neuroendocrine Changes.
Integr. Comp. Biol. 55, 294–306. https://doi.org/10.1093/icb/icv057
Maruska, K.P., Fernald, R.D., 2010. Steroid receptor expression in the fish inner ear varies with
sex, social status, and reproductive state. BMC Neurosci. 11, 58.
https://doi.org/10.1186/1471-2202-11-58
McClelland, S., Korosi, A., Cope, J., Ivy, A., Baram, T.Z., 2011. Emerging roles of epigenetic
mechanisms in the enduring effects of early-life stress and experience on learning and
memory. Neurobiol. Learn. Mem. 96, 79–88. https://doi.org/10.1016/j.nlm.2011.02.008
Meyer-Lindenberg, A., Tost, H., 2012. Neural mechanisms of social risk for psychiatric
disorders. Nat. Neurosci. 15, 663–668. https://doi.org/10.1038/nn.3083
Meyer, K., Korz, V., 2013. Estrogen receptor α functions in the regulation of motivation and
spatial cognition in young male rats. PLoS One 8.
https://doi.org/10.1371/journal.pone.0079303
Moy, S.S., Nadler, J.J., Perez, A., Barbaro, R.P., Johns, J.M., Magnuson, T.R., Piven, J.,
Crawley, J.N., 2004. Sociability and preference for social novelty in five inbred strains: an
approach to assess autistic-like behavior in mice. Genes, Brain Behav. 3, 287–302.
https://doi.org/10.1111/j.1601-1848.2004.00076.x
Munchrath, L.A., Hofmann, H.A., 2010. Distribution of sex steroid hormone receptors in the
brain of an African cichlid fish, Astatotilapia burtoni. J. Comp. Neurol. 518, 3302–3326.
https://doi.org/10.1002/cne.22401
Näslund, J., Johnsson, J.I., 2016. State-dependent behavior and alternative behavioral strategies
in brown trout (Salmo trutta L.) fry. Behav. Ecol. Sociobiol. 70, 2111–2125.
https://doi.org/10.1007/s00265-016-2215-y
Newman, S., 1999. The medial extended amygdala in male reproductive behavior. Ann NY Acad
Sci 242–257.
Nyman, C., Fischer, S., Aubin-Horth, N., Taborsky, B., 2018. Evolutionary conserved neural
signature of early life stress affects animal social competence. Proc. R. Soc. B Biol. Sci.

1039   285. https://doi.org/10.1098/rspb.2017.2344

1040 Nyman, C., Fischer, S., Aubin-Horth, N., Taborsky, B., 2017. Effect of the early social
1041   environment on behavioural and genomic responses to a social challenge in a cooperatively
1042   breeding vertebrate. Mol. Ecol. 26, 3186–3203. https://doi.org/10.1111/mec.14113

1043 O'Connell, L.A., Hofmann, H.A., 2012a. Social Status Predicts How Sex Steroid Receptors
1044   Regulate Complex Behavior across Levels of Biological Organization. Endocrinology 153,
1045   1341–1351. https://doi.org/10.1210/en.2011-1663

1046 O'Connell, L.A., Hofmann, H.A., 2012b. Evolution of a Vertebrate Social Decision-Making
1047   Network. Science (80-.). 336, 1154–1157. https://doi.org/10.1126/science.1218889

1048 O'Connell, L.A., Hofmann, H.A., 2011. The Vertebrate mesolimbic reward system and social
1049   behavior network: A comparative synthesis. J. Comp. Neurol. 519, 3599–3639.
1050   https://doi.org/10.1002/cne.22735

1051 Piersma, T., Drent, J., 2003. Phenotypic flexibility and the evolution of organismal design.
1052   Trends Ecol. Evol. 18, 228–233. https://doi.org/10.1016/S0169-5347(03)00036-3

1053 Pradhan, D.S., Solomon-Lane, T.K., Grober, M.S., 2015. Contextual modulation of social and
1054   endocrine correlates of fitness: insights from the life history of a sex changing fish. Front.
1055   Neurosci. 9, 1–21. https://doi.org/10.3389/fnins.2015.00008

1056 Prut, L., Belzung, C., 2003. The open field as a paradigm to measure the effects of drugs on
1057   anxiety-like behaviors: A review. Eur. J. Pharmacol. 463, 3–33.
1058   https://doi.org/10.1016/S0014-2999(03)01272-X

1059 Rajkov, J., Weber, A.A.T., Salzburger, W., Egger, B., 2018. Adaptive phenotypic plasticity
1060   contributes to divergence between lake and river populations of an East African cichlid fish.
1061   Ecol. Evol. 8, 7323–7333. https://doi.org/10.1002/ece3.4241

1062 Remage-Healey, L., Romero, L.M., 2000. Daily and seasonal variation in response to stress in
1063   captive starlings (sturnus vulgaris): Glucose. Gen. Comp. Endocrinol. 119, 60–68.
1064   https://doi.org/10.1006/gcen.2000.7492

1065 Renn, S., Aubin-Horth, N., Hofmann, H.A., 2008. Fish and chips: functional genomics of social
1066   plasticity in an African cichlid fish. J. Exp. Biol. 211, 3041–3056.
1067   https://doi.org/10.1242/jeb.018242

1068 Renn, S., Carleton, J.B., Magee, H., Nguyen, M.L.T., Tanner, A.C.W., 2009. Maternal care and
1069   altered social phenotype in a recently collected stock of Astatotilapia burtoni cichlid fish.
1070   Integr. Comp. Biol. 49, 660–673. https://doi.org/10.1093/icb/icp085

1071 Robbins, M.M., Robbins, A.M., Gerald-Steklis, N., Steklis, H.D., 2007. Socioecological
1072   influences on the reproductive success of female mountain gorillas (Gorilla beringei
1073   beringei). Behav. Ecol. Sociobiol. 61, 919–931. https://doi.org/10.1007/s00265-006-0321-y

1074 Rochette-Egly, C., 2005. Dynamic combinatorial networks in nuclear receptor-mediated
1075   transcription. J. Biol. Chem. 280, 32565–32568. https://doi.org/10.1074/jbc.R500008200

1076 Saltz, J.B., Geiger, A.P., Anderson, R., Johnson, B., Marren, R., 2016. What, if anything, is a
1077   social niche? Evol. Ecol. 30, 349–364. https://doi.org/10.1007/s10682-015-9792-5

1078 Schreck, C.B., 2010. Stress and fish reproduction: The roles of allostasis and hormesis. Gen.
1079   Comp. Endocrinol. 165, 549–556. https://doi.org/10.1016/j.ygcen.2009.07.004

1080 Shepard, K.N., Michopoulos, V., Toufexis, D.J., Wilson, M.E., 2009. Genetic, epigenetic and
1081   environmental impact on sex differences in social behavior. Physiol. Behav. 97, 157–170.
1082   https://doi.org/10.1016/j.physbeh.2009.02.016

1083 Sih, A., Bell, A.M., Johnson, J.C., 2004a. Behavioral syndromes: an ecological and evolutionary
1084   overview. Trends Ecol. Evol. 19, 372–378. https://doi.org/10.1016/j.tree.2004.04.009

Sih, A., Bell, A.M., Ziemba, R.E., 2004b. Behavioural syndromes: An integrative overview. Q.
Rev. Biol. 80, 1–1. https://doi.org/10.1086/516403
Silk, J.B., 2007. Social Components of Fitness in Primate Groups. Science (80-.). 317, 1347–
1351. https://doi.org/10.1126/science.1140734
Smith, B.R., Blumstein, D.T., 2008. Fitness consequences of personality: A meta-analysis.
Behav. Ecol. 19, 448–455. https://doi.org/10.1093/beheco/arm144
Snell-Rood, E.C., 2013. An overview of the evolutionary causes and consequences of
behavioural plasticity. Anim. Behav. 85, 1004–1011.
https://doi.org/10.1016/j.anbehav.2012.12.031
Solomon-Lane, T.K., Crespi, E.J., Grober, M.S., 2013. Stress and serial adult metamorphosis :
multiple roles for the stress axis in socially regulated sex change. Front. Neuroendocrinol. 7,
1–12. https://doi.org/10.3389/fnins.2013.00210
Solomon-Lane, T.K., Pradhan, D.S., Willis, M.C., Grober, M.S., 2015. Agonistic reciprocity is
associated with reduced male reproductive success within haremic social networks. Proc. R.
Soc. B Biol. Sci. 282, 20150914. https://doi.org/10.1098/rspb.2015.0914
Solomon-Lane, T.K., Pradhan, D.S., Willis, M.C., Grober, M.S., 2014. Female, but not male,
agonistic behaviour is associated with male reproductive success in stable bluebanded goby
(Lythrypnus dalli) hierarchies. Behaviour 151, 1367–1387.
https://doi.org/10.1163/1568539X-00003188
Stamps, J.A., 2003. Behavioural processes affecting development: Tinbergen's fourth question
comes of age. Anim. Behav. 66, 1–13. https://doi.org/10.1006/anbe.2003.2180
Stamps, J.A., Groothuis, T.G.G., 2010. The development of animal personality: Relevance,
concepts and perspectives. Biol. Rev. 85, 301–325. https://doi.org/10.1111/j.1469-
185X.2009.00103.x
Stearns, S.C., 1989. The Evolutionary Significance of Phenotypic Plasticity. BioScience, 39,
436–445. https://doi.org/10.2307/1311135
Stevenson, T.J., Alward, B.A., Ebling, F.J.P., Fernald, R.D., Kelly, A., Ophir, A.G., 2017. The
Value of Comparative Animal Research: Krogh's Principle Facilitates Scientific
Discoveries. Policy Insights from Behav. Brain Sci. 237273221774509.
https://doi.org/10.1007/978-3-540-68279-0_15
Taborsky, B., 2016. Opening the Black Box of Developmental Experiments: Behavioural
Mechanisms Underlying Long-Term Effects of Early Social Experience. Ethology 122,
267–283. https://doi.org/10.1111/eth.12473
Taborsky, B., Arnold, C., Junker, J., Tschopp, A., 2012. The early social environment affects
social competence in a cooperative breeder. Anim. Behav. 83, 1067–1074.
https://doi.org/10.1016/j.anbehav.2012.01.037
Taborsky, B., Tschirren, L., Meunier, C., Aubin-Horth, N., 2013. Stable reprogramming of brain
transcription profiles by the early social environment in a cooperatively breeding fish. Proc.
Biol. Sci. 280, 20122605. https://doi.org/10.1098/rspb.2012.2605
Tinbergen, N., 1963. On aims and methods of Ethology. Anim. Biol. 55, 297–321.
https://doi.org/10.1163/157075605774840941
Tsalafouta, A., Sarropoulou, E., Papandroulakis, N., Pavlidis, M., 2018. Characterization and
Expression Dynamics of Key Genes Involved in the Gilthead Sea Bream (Sparus aurata)
Cortisol Stress Response during Early Ontogeny. Mar. Biotechnol. 20, 611–622.
https://doi.org/10.1007/s10126-018-9833-5
Turecki, G., Meaney, M.J., 2016. Effects of the Social Environment and Stress on

1131   Glucocorticoid Receptor Gene Methylation: A Systematic Review. Biol. Psychiatry 79, 87–
1132   96. https://doi.org/10.1016/j.biopsych.2014.11.022
1133 Verbeek, M.E.M., Drent, P.J., Wiepkema, P.R., 1994. Consistent individual differences in early
1134   exploratory behaviour of male great tits. Anim. Behav.
1135   https://doi.org/10.1006/anbe.1994.1344
1136 Wada, H., 2008. Glucocorticoids: mediators of vertebrate ontogenetic transitions. Gen. Comp.
1137   Endocrinol. 156, 441–53. https://doi.org/10.1016/j.ygcen.2008.02.004
1138 Weitekamp, C.A., Hofmann, H.A., 2017. Neuromolecular correlates of cooperation and conflict
1139   during territory defense in a cichlid fish. Horm. Behav. 89, 145–156.
1140   https://doi.org/10.1016/j.yhbeh.2017.01.001
1141 Weitekamp, C.A., Solomon-Lane, T.K., Del Valle, P., Triki, Z., Nugent, B.M., Hofmann, H.A.,
1142   2017. A Role for Oxytocin-Like Receptor in Social Habituation in a Teleost. Brain. Behav.
1143   Evol. 89, 153–161. https://doi.org/10.1159/000464098
1144 Wendelaar Bonga, S.E., 1997. The stress response in fish. Physiol. Rev. 77, 591–625.
1145   https://doi.org/stress physiologie comportement cortisol cerveau hypophyse catecholamine
1146   osmoregulation branchie
1147 West-Eberhard, M.J., 1989. Phenotypic plasticity and the origins of diversity. Annu. Rev. Ecol.
1148   Syst. 20, 249–278. https://doi.org/10.1146/annurev.es.20.110189.001341
1149 White, D.J., 2010. The Form and Function of Social Development - Insights From a Parasite.
1150   Curr. Dir. Psychol. Sci. 19, 314–318. https://doi.org/10.1177/0963721410383384
1151 White, D.J., King, A.P., West, M.J., 2002. Facultative development of courtship and
1152   communication in juvenile male cowbirds (Molothrus ater). Behav. Ecol. 13, 487–496.
1153   https://doi.org/10.1093/beheco/13.4.487
1154 Wilson, E.O., 1980. Sociobiology: the Abridged Edition. The Belknap Press of Harvard
1155   University Press, Cambridge.
1156 Wingfield, J.C., Hegner, R.E., Dufty, Jr., A.M., Ball, G.F., 1990. The "Challenge Hypothesis":
1157   Theoretical Implications for Patterns of Testosterone Secretion, Mating Systems, and
1158   Breeding Strategies. Am. Nat. 136, 829. https://doi.org/10.1086/285134
1159 Young, A.J., Carlson, A. a, Monfort, S.L., Russell, A.F., Bennett, N.C., Clutton-Brock, T., 2006.
1160   Stress and the suppression of subordinate reproduction in cooperatively breeding meerkats.
1161   Proc. Natl. Acad. Sci. 103, 12005–12010. https://doi.org/10.1073/pnas.0510038103
1162
1163

Figures & legends

1165

Figure 1: Experimental setup for behavior assays. Juvenile behavior was observed in a novel

experimental tank in four sequential assays administered in the same order, each lasting 30 min.

A terracotta shard served as a shelter and/or territory. The black lines (dotted, solid) were drawn

on the tank bottom in permanent maker, dividing the tank into four zones: territory, close, far,

and investigate. The center dividing line (white) was not drawn (A). The focal fish was alone in

the tank for the open field assay, and the time in each zone and frequency of entered each zone

was recorded (B, assay 1). For the social cue investigation, a juvenile inside of a scintillation vial

was placed in the circle within the investigate zone (see C). The time in and frequency of

entering each zone was recorded (B, assay 2). The social cue was removed and a freely

swimming, novel cue fish (smaller than the focal) was added to the tank for the dominance

behavior assay (B, assay 3). The small cue fish was then removed and a freely swimming, novel

cue fish (larger than the focal) was added to the tank for the subordinate behavior assay (B, assay

4). Social interactions were recorded for the dominant and subordinate behavior assays. The time

in and frequency of entering the territory zone was also recorded for both fish.

Figure 2: Principal component analysis (PCA) of focal fish behavior from all four assays (open field, social cue investigation, dominance, subordinate behavior). Differences in PC1 between group- and pair-reared juveniles (A). Differences in PC2 (B) and PC5 (C) between the larger / dominant (Dom) fish and smaller / subordinate (Sub) fish selected from the group and pair. The larger fish is very likely to have more dominance experience, while the smaller fish has more subordinate experience. Differences in PC6 between group- and pair-reared juveniles (D). Vector plot showing the PCA variables that load on PC1 and PC2 (E). Vector plot showing the PCA variables that load on PC5 and PC6 (F). Percentages refer to the amount of variance explained by that component. Pair (n=18 individuals). Group (n=24 individuals). Social cue investigation (SCI). Open field exploration (OF). *p<0.05, ****p<0.001.

1191

1192

1193 **Figure 3:** Separate principal component analyses performed for the open field (A), social cue

1194 investigation (B), and dominance behavior (C) assays. Both focal and non-focal fish variables

1195 (behavior, size). The significant, positive correlations about the PC1s are shown in a three-

1196 dimensional plot (D). Percentages refer to the amount of variance explained by that component.

1197 Pair (n=18 individuals). Group (n=24 individuals). *p<0.05, **p<0.01.

1198

1199

1200 **Figure 4:** Relative gene expression calculated using ΔΔCT analysis (reference gene 18S) for

1201 juveniles reared in isolation (1 week, n=8), pairs (1 week or 5 weeks, n=18), and groups (1 week

1202 or 5 weeks, n=22). Androgen receptor α (ARα). Estrogen receptor α (ERα). Glucocorticoid

1203 receptors (GR). Mineralocorticoid receptor (MR). Corticotropin-releasing factor (CRF). Letters

1204 indicate significant differences across treatment groups (p<0.05). *p<0.05, **p<0.01.

1205

Figure 5: Partial correlation network of gene expression in pair-reared juveniles (n=18) (A) and

group-reared juveniles (n=22) (B). Nodes are the candidate genes. Edges represent partial

correlations between nodes. Only significant partial correlations are shown (p<0.05), and edge

thickness indicates correlation strength. There were no significant partial correlations for

juveniles reared in isolation (n=8) (p>0.05). Androgen receptor α (AR). Estrogen receptor α

(ER). Glucocorticoid receptors (GR). Mineralocorticoid receptor (MR). Corticotropin-releasing

factor (CRF).