Impulsive choice in mice lacking paternal expression of *Grb10* provides evidence for intra-

genomic conflict in behavior

Claire L. Dent*, Trevor Humby[†], Katie Lewis[*], Andrew Ward[‡], Reiner Fischer-Colbrie[§],

Lawrence S. Wilkinson[*,†], Jon F Wilkins[**] & Anthony R. Isles[*,††]

*Behavioural Genetics Group, MRC Centre for Neuropsychiatric Genetics and Genomics,

Neuroscience and Mental Health Research Institute, Cardiff University, Hadyn Ellis Building,

Maindy Road, Cardiff, United Kingdom

[†]Behavioural Genetics Group, School of Psychology, Cardiff University, Tower Building,

Cardiff, United Kingdom

[‡]Department of Biology and Biochemistry, University of Bath, Bath, United Kingdom

[§]Department of Pharmacology, Innsbruck Medical University, Innsbruck, Austria

[**]Ronin Institute, Montclair, NJ 07043, USA

[††]Author for correspondence: Anthony R. Isles, E-mail IslesAR1@cardiff.ac.uk

Running title: **Intragenomic conflict and impulsivity**

Keywords: genomic imprinting; Nesp55; delayed reinforcement; intra-genomic conflict;

Grb10

1   Abstract

Imprinted genes are expressed from one parental allele only as a consequence of epigenetic

events that take place in the mammalian germ line and are thought to have evolved through

intra-genomic conflict between parental alleles. We demonstrate, for the first time,

oppositional effects of imprinted genes on brain and behavior. Specifically, here we show

that mice lacking paternal *Grb10* make fewer impulsive choices, with no dissociable effects

on a separate measure of impulsive action. Taken together with previous work showing that

mice lacking maternal Nesp55 make more impulsive choices this suggests that impulsive

choice behavior is a substrate for the action of genomic imprinting. Moreover, the

contrasting effect of these two genes suggests impulsive choices are subject to intra-

genomic conflict and that maternal and paternal interests pull this behavior in opposite

directions. Finally, these data may also indicate that an imbalance in expression of imprinted

genes contributes to pathological conditions such as gambling and drug addiction, where

impulsive behavior becomes maladaptive.

INTRODUCTION

Imprinted genes are expressed from one parental allele only as a consequence of epigenetic events that take place in the mammalian germ line (FERGUSON-SMITH 2011). Functionally, imprinted genes converge on specific biological processes that have prominent importance in mammals, such as *in utero* growth, metabolism and behavior (PETERS 2014; WILKINS *et al.* 2016). Many of the effects that maternally- and paternally-expressed imprinted genes have on *in utero* growth and during the early post-natal period are oppositional in direction (HAIG AND GRAHAM 1991; MADON-SIMON *et al.* 2014), lending support to the theory that imprinting has evolved as a consequence of intra-genomic conflict (MOORE AND HAIG 1991; WILKINS AND HAIG 2003).

Grb10 is currently a unique example of an imprinted gene that is expressed from the maternal allele only in some tissues, and the paternal allele only in others (ARNAUD *et al.* 2003; HIKICHI *et al.* 2003; MONK *et al.* 2009). Consequently, different parental alleles of *Grb10* mediate distinct physiological functions (GARFIELD *et al.* 2011). Maternal *Grb10* is expressed in many peripheral body tissues, whilst being excluded from the central nervous system (GARFIELD *et al.* 2011), and plays a prominent role in controlling placental function (CHARALAMBOUS *et al.* 2003; CHARALAMBOUS *et al.* 2010) and metabolism (SMITH *et al.* 2007). In contrast, paternal *Grb10* is essentially restricted to the brain alone and impacts on behavioral phenotypes, including social dominance (GARFIELD *et al.* 2011). *Grb10* expression in the brain is neuronal, showing a pattern of expression in discrete brain regions that is shared with the maternally expressed imprinted gene *Nesp* (PLAGGE *et al.* 2005). This overlap in brain expression of maternal *Nesp* (encoding Nesp55) and paternal *Grb10* has led to speculation that these two genes may influence common adult behaviors, possibly in an

opposite manner (Garfield *et al.* 2011; Dent and Isles 2014), which would be consistent with

the prediction from the intra-genomic conflict theory of genomic imprinting evolution (Haig

2000).

Here, we address this question directly. We have shown that mice lacking maternal Nesp55

(*Nesp*$^{m/+}$) make more impulsive choices (Dent *et al.* 2016). In contrast, here we show that

Grb10$^{+/p}$ mice make less impulsive choices relative to their littermate controls. Furthermore,

there were no dissociable effects on a separate measure of impulsive action, completely

matching the specificity of effects seen previously in *Nesp*$^{m/+}$ mice (Dent *et al.* 2016). Our

data provide new findings on *Grb10* function in brain and reveal opposite effects of

manipulating *Grb10* and *Nesp* expression on discrete aspects of impulsive responding. Taken

together these data are consistent with the conflict theory of the evolution of imprinted

genes and as such provide the first direct evidence for intra-genomic conflict impacting

adult behavior.

RESULTS AND DISCUSSION

***Grb10*$^{+/p}$ mice make less impulsive choices**

We used the delayed-reinforcement task to examine impulsive choice behavior (Isles *et al.*

2003), where subjects made a choice between receiving a small food reward after a short

(1s) delay, or a larger reward after a longer delay (1, 8 or 16s). The extent to which subjects

make impulsive choices is indexed by a preference for choosing the immediate but smaller

reward, versus a larger delayed reward (delayed gratification). Increasing the delay to the

larger reward within session decreased the likelihood of choosing that reward across all

subjects (Figure 1A; main effect of DELAY, $F_{2,42}$=9.20, *p*=0.001, partial η^2=0.37). However,

Grb10[+/p] mice were significantly more tolerant of an increased delay to the larger reward

than wild-type (WT) littermate mice (Figure 1A; main effect of GENOTYPE, $F_{1,21}=6.45$,

p=0.019, partial η^2=0.25). However, whilst the data indicate that the experimental group

diverges from wild-type increasingly as the delay increases, this effect failed to result in a

significant interaction between GENOTYPE and DELAY (Figure 1A; $F_{2,42}=0.56$, *p*=0.946, partial

η^2=0.003), due largely to a pre-existing group difference in the first 1s delay block.

A standard control measure to assess the extent to which behavior is sensitive to delay is a

task manipulation where any differential delay associated with the larger and smaller

reward was equalized (1s) across all three blocks. Here, all subjects now demonstrated the

expected preference for the larger reward throughout the session (Figure 1B; no main effect

of DELAY, $F_{1.7,35.1}=1.88$, *p*=0.17, partial η^2=0.08); that is under these task conditions there

were no differences in choice-bias between *Grb10*[+/p] and WT mice (no main effect of

GENOTYPE, $F_{1,21}=0.63$, *p*=0.44, partial η^2=0.03). Furthermore, there was no difference in the

sessions taken to achieve a stable baseline in these conditions (WT 9 ±0.8; *Grb10*[+/p] 8 ±0.5;

t_{21}=1.51, p=0.15).

The contrasting patterns of choice behavior between the *Grb10*[+/p] and WT mice was also not

due to any differences between the groups in terms of basic motivation to carry out the task

as *Grb10*[+/p] animals acquired the task at the same rate as WT littermates (sessions to last

day of baseline; WT 50.4 ±1.1 SEM, *Grb10*[+/p] 47.7 ±1.2 SEM t_{21}=1.73, *p*=0.10). Additionally,

variability between *Grb10*[+/p] and WT mice was not related to differences in experiencing the

information trial contingencies. In the 'forced' trials (where no choice was available), both

Grb10[+/p] and their WT littermates made equal responses to the large and small reward-

related stimuli at all delays (no interaction GENOTYPE x DELAY x CHOICE, $F_{2,42}=0.44$,

p=0.649; data not shown). Finally, there was no difference between $Grb10^{+/p}$ and WT mice

on general measures of task performance (see Supplemental Information).

Delay to large reward (s)

Figure 1 $Grb10^{+/p}$ mice make less impulsive choices on a delayed-reinforcement task.

Behavior of both $Grb10^{+/p}$ and WT mice changed across session blocks (blk) with increasing

delay, such that choice bias moved away from the response leading to the large reward

towards the small reward, with increasing delay (**A**). However, there were systematic

differences between the groups in their behavior, such that $Grb10^{+/p}$ animals switched their

choice to the small, less delayed reward less quickly than WT mice.

When the delay associated with the large and small rewards was equal (1s) throughout the

session (**B**), choice bias was consistently high (large reward chosen approximately 80% of

the time). Under these task conditions there were no differences in choice bias between

$Grb10^{+/p}$ and WT mice. Data shows mean ±SEM of three consecutive stable sessions; *

represents P<0.05 main effect of GENOTYPE; ## represents P<0.01 main effect of DELAY.

In large part choice bias behavior in delayed reinforcement tasks is governed by three

potentially interacting psychological processes; the perceived value of the rewards, and the

perceived length and aversive nature of the delay (Ho *et al.* 1999). The difference in choice

between *Grb10*^{+/p} and WT mice in the first block of the task (where delay for the larger and

smaller rewards are both 1s) may suggest a contribution to behavior in the *Grb10*^{+/p} from a

relatively more less pronounced generalized conditioned place aversion to the delayed

larger reward response (see Isles *et al.* 2004). However, importantly, such an effect did not

lead to an inflexible bias in responding as demonstrated by the continued sensitivity to

delay, both within a baseline session and in response to the equal-delay probe

manipulation. Therefore, whilst the between-group behavioral differences were clearly

influenced to a large degree by a differential tolerance to delay the present data do not rule

out influences due to aversion and reward perception

Grb10^{+/p} mice show no difference in impulsive action

In contrast to impulsive choice, there was no difference between *Grb10*^{+/p} and WT mice on a

measure of impulsive action. The stop-signal reaction time (SSRT) task measures the ability

to stop an action once initiated by presenting a stop-signal (in this case an auditory tone)

during a rapid response between two stimuli locations (the 'go-response'). Correctly

inhibiting the go response will earn reward in a 'stop' trial, whereas reward was also

presented on completing a go-response in trials where no stop-signal was presented.

Throughout the training stages of the SSRT task, all subjects showed show equivalent

behavior in learning the task and *Grb10*^{+/p} and WT mice acquired the task at the same rate

(sessions taken to complete the task, *Grb10*^{+/p}: 42.4± 9.8, WT: 40.8±4.5, t_{20}=0.54, *p*=0.59).

Figure 2 No difference beween *Grb10*$^{+/p}$ and WT mice in performance on a stop-signal reaction time (SSRT) task. Both WT and *Grb10*$^{+/p}$ mice showed an equivalent ability to perform the SSRT task showing the expected change in percentage correct responding during a 'stop' trial (**A**) as the position of the stop-signal was altered, but there were no differences between *Grb10*$^{+/p}$ and WT mice. *Grb10*$^{+/p}$ and WT mice also showed equivalent SSRTs at 50% correct stopping (**B**). There were no genotype differences for the Go-response for both groups of mice, in terms of percentage correct responding (**C**) or response speed (**D**). Data shows mean ±SEM; ## represents P<0.01 main effect of stop-signal position

2   As expected (Humby *et al.* 2013; Davies *et al.* 2014; Dent *et al.* 2016), presenting the stop-

3   signal progressively closer to the execution of the response (10, 40, 50, 60 and 90% into the

4   individualised go response) led to systematic reductions in the ability to stop for all mice

5   (Figure 2A; ANOVA, main effect of STOP, $F_{4,80}$=24.37, *p*<0.001, partial η^2=0.55). However,

there were no differences in stopping efficiency between $Grb10^{+/p}$ and WT mice (Figure 2A, main effect of GENOTYPE, $F_{1,20}=0.94$, $p=0.34$, partial $\eta^2=0.05$). The lack of genotype differences in stopping ability was further demonstrated by another measure of response inhibition, the speed of stopping or stop-signal reaction time (SSRT) derived in sessions where the subjects exhibited 50% correct stopping (BARI et al. 2009; HUMBY et al. 2013; DAVIES et al. 2014; DENT et al. 2016). Equivalent SSRTs were observed in both $Grb10^{+/p}$ and their WT littermates (Figure 2B; $t=17$, $p=0.87$).

There were also no differences in the Go-response between $Grb10^{+/p}$ and WT mice, in terms of the amount (Figure 2C, main effect of GENOTYPE, $F_{1,20}=1.14$, $P=0.30$) or speed (Fig. 2D, main effect of GENOTYPE, $F_{1,20}=0.03$, $P=0.86$), of correct responding in Go trials. These parameters were not affected in sessions when the stop-signal position was moved from baseline (main effect of STOP-SIGNAL POSITION, $F_{4,80}=0.28$, $P=0.89$ and $F_{4,80}=0.85$, $P=0.50$, for the amount and speed of correct responding in Go trials, respectively). Additionally, there was no difference between $Grb10^{+/p}$ and WT mice on general measures of task performance during individualized SSRT session (see Supplemental Information), indicating a high degree of stimulus control for both groups in the task.

Opposite effects to maternal *Nesp*

The current data obtained from knocking out the paternal copy of *Grb10* mirror are essentially opposite to our previously published effects of knocking out the maternal copy of Nesp55 (DENT et al. 2016). That is, under identical task conditions, mice lacking maternal Nesp55 make more impulsive choices, whereas mice lacking paternal *Grb10* make fewer impulsive choices, relative to their littermate controls. These effects were highly specific in that they occurred in the absence of deficits in either model on responding in the SSRT task.

The idea that these oppositely imprinted genes converge on discrete aspects of impulse

control is further supported by their general overlapping expression patterns in the brain

(PLAGGE *et al.* 2005; GARFIELD *et al.* 2011). In the present work we extend these

neuroanatomical data in showing that Nesp55 and Grb10 are co-localised in cells in the

locus coeruleus (Figure 3a-c), hypothalamus (Figure 3d-f), and the dorsal raphe nucleus

(Figure 3g-i). Together, these studies indicate that impulsive choice behavior is a substrate

for the action of genomic imprinting, and an extension of this idea is the speculation that an

imbalance in expression of imprinted genes, such as *GRB10* or *NESP*, may also contribute to

pathological conditions, such as pathological gambling, where choices become highly

maladaptive. In this regard it should be noted that neither *GRB10* nor *NESP* have associated

SNPs identified as significant variants in recent GWAS studies of gambling behaviour (LANG

et al. 2016), or indeed delay discounting (SANCHEZ-ROIGE *et al.* 2018). However, it is worth

noting also that genes with known roles in impulsive behaviour (eg *DAT, DRD1* or *DRD2*)

were not associated either, suggesting that a lack of signal in GWAS does not necessarily

diminish the suggestion of a role for *GRB10* and/or *NESP* in these disorders of impulse

control.

Figure 3 Dual-labelling immunofluorescence histochemistry of Nesp55 and Grb10 in coronal sections of adult brain. Sections were dual-labelled with antibodies against Nesp55 and β-gal, where the reporter gene LacZ is expressed in place of Grb10 in tissue from Grb10$^{+/p}$ mice, and can be used to identify Grb10-positive cells. Images were viewed at different light intensities (568nm and 488nm for Nesp55 and b-gal, respectively), and were then merged to gauge cellular co-localisation of the two target proteins, depicted by white arrows in the merged figures. The majority of cells showed evidence for co-localisation: within the locus coeruleus (LC) (a-c), the hypothalamus (HYP) (d-f) and the dorsal raphe nuclei (DRN) (g-i). LC and HYP images at x40 magnification, DRN at x20.

We also suggest these data are consistent with the intra-genomic conflict theory of the

evolution of imprinted genes, providing the first direct evidence for oppositional effects of

imprinted genes on an adult behavior. In this context, the examples of Nesp55- and *Grb10*-

null mice are particularly important, as these models are free from confounding effects on

in utero or pre-weaning growth (PLAGGE *et al.* 2005; GARFIELD *et al.* 2011). Hence, this

strengthens the argument that imprinted gene effects on adult behavior have adaptive

significance and are not simply epiphenomena, that is, enduring into adulthood but

resulting from the more familiar imprinted gene substrate of resource allocation between

mother and offspring (CONSTANCIA *et al.* 2004).

Imprinted genes are thought to have evolved as a consequence of conflicting phenotypic

"interests" between maternal and paternal genes that causes an escalating arms race in

relation to allelic expression, eventually leading to silencing of one or other parental allele

(Moore & Haig, 1991, Wilkins & Haig, 2003). Evidence of this parental conflict is seen in the

contrasting action of maternal and paternal imprinted genes during *in utero* growth and

early post-natal life (Haig, 2004). Although the present work is the first demonstration of

opposing parental interests on behavior, the specificity of imprinting effects on impulsive

choice behavior described here would fit with previous suggestions that maternal and

paternal genomes may have a contrasting impact on decision-making in social animals (HAIG

2000; BRANDVAIN *et al.* 2011). Patterns of impulsive choice behavior with increasing delay, as

measured by delay-discounting in the delayed-reinforcement task here, are often thought of

as 'irrational', in that the shape of behavior is not consistent with simple models of

optimization (such as expected utility maximization) (SCHONBERG *et al.* 2011). It may be that

the so-called "parliament of the mind" (HAIG 2000), caused by opposing parental genomes

pulling impulsive choice in different directions, is an additional factor that should be taken

into consideration in the context of more complex models of apparently 'irrational'

behaviors (FAWCETT *et al.* 2014).

MATERIALS AND METHODS

Animals

All procedures were conducted in accordance with the UK Animals (Scientific Procedures)

Act 1986. Subjects were male mice, and were 4 months old at the start of testing, which was

completed after 6-months. Standard laboratory chow was available *ad libitum*, but during

the experiment water was restricted to 2h access per day. This regime maintained the

subjects at ≈90% of free-feeding body weight. Due to potentially confounding phenotypes

associated with *Grb10*$^{m/+}$ mice, comparisons of *Grb10*$^{+/p}$ were made with wild-type (WT)

littermate controls derived from 8 separate litters. The *Grb10* null line was maintained on an

F1-hybrid (C57/Bl6 X CBA) background. The same cohort was used throughout testing but a

number of animals were lost (due to death or inability to perform training stages) as testing

progressed.

Behavioral testing

Operant testing apparatus

All the sessions of the delayed reinforcement task and SSRT task were performed in 9-hole

operant chambers (Cambridge Cognition Ltd, U.K) modified for use in mice, as described

previously (HUMBY *et al.* 1999; ISLES *et al.* 2003; HUMBY *et al.* 2013). For the delayed

reinforcement task holes 3, 5 and 7 were open; whereas only holes 4 and 6 were open for

the SSRT task. The mice were presented with a visual stimuli (light) recessed into the holes

and were trained to respond to this stimulus with a nose-poke as recorded by infra-red

beams spanning the hole. Reward was presented in a recessed compartment on the wall

opposite to the nose-poke/stimulus array. The control of the stimuli and recording of the

responses were managed by an Acorn Archimedes computer with additional interfacing by

ARACHNID (Cambridge Cognition Ltd). For all operant testing, animals were maintained on a

restricted water access schedule, water provided for two hours immediately after testing.

Delayed reinforcement task

Details of the shaping procedures and basic aspects of the delayed-reinforcement task itself

can be found elsewhere (Isles *et al.* 2003). Briefly, the task comprised of three sequential

blocks of 12 trials, with each trial consisting of an initial nose-poke to the centrally located

stimulus, followed by a second nose-poke to either the left or right apertures. Trials 1-4 in

any block were 'forced' information trials, where the initial nose-poke resulted in

presentation of only one of the two choice options. This measure was designed to provide

the subjects with prior notice of the extent of any delay associated with choosing the large

reward. In the remaining 8 trials of each block, designated as 'choice' trials, the initial

centre nose-poke led to the option of a second nose-poke response to either the left or right

apertures. One response resulted in the delivery of a large reward (50µL 10% solution of

condensed milk; Nestle Ltd, UK), and the other in the delivery of small reward (25µL 10%

solution of condensed milk). The response contingencies were kept constant for each

mouse, but were counterbalanced between subjects. In block 1 both responses led to the

delivery of reward after a 1s delay. In blocks 2 and 3 increasing delays were introduced

between the response and the delivery of the large reward (8s, 16s, respectively) whereas

the delay between response and delivery of the small reward was fixed at 1s. As a probe to

test the effect of the delays on behavior, sessions were conducted where the delay

associated with the large reward was fixed at 1s, equivalent to that associated with the small reward, throughout all three blocks of the session.

The bias in choice of the larger reward at each block (whereby always choosing the large reward=1; never choosing the large reward=0) was the main measure used to determine impulsive responding. Additional measurements that related to general motoric competence and motivation within the task were also monitored, including the 'Start' and 'Choice' latencies, the time taken to initiate a trial and the time taken to make a choice once a trial was initiated, respectively. Also measured were the number of 'Non-started' (no initial, central nose-poke) and 'Omitted' (no secondary, choice nose-poke, following central nose-poke initiating trial) trials.

Stop-signal reaction time (SSRT) task

Details of the shaping procedures and basic aspects of the main SSRT task can also be found elsewhere (HUMBY *et al.* 2013). The SSRT task itself consisted of sessions of 100 trials, which involved both 'go' and 'stop' trials. Go trials consisted of rapid double nose-pokes (a 'go' response) between two separate stimuli locations, which were rewarded with reinforcement (22ul, 10% solution of condensed milk, Nestle Ltd, UK). 20% of trials were stop trials, pseudo-randomly distributed throughout each session, where a stop-signal (65db white noise for 0.3s) was presented between the first and second nose-poke responses. The aim of the stop-signal was to inhibit ('Stop') the mouse from making the second ('Go') nose-poke, and then wait for the reward. Failure to refrain from making this pre-potent response was punished by the absence of reward and 5 second time out (chamber light on). At baseline, the stop-signals were presented concurrently with the initial nose-poke response. To maintain high levels of performance of both go and stop responding, the go stimulus duration and wait period to reward delivery in a stop-signal trial were determined

individually for each subject. To assess the ability to stop once an action had been initiated, sessions were implemented in which the onset of the stop-signal was presented at different positions within the individualised go response of each mouse. Thus, the stop-signal was pseudo-randomly presented 10, 40, 50, 60 and 90% from the onset of the go response of each subject, with the assumption that stopping would be more difficult the closer the stop-signal presentation was to the termination of the go response.

The amount of correct stopping in stop-signal trials and the SSRT were the main measures of impulsive responding in this task. The SSRT was calculated by determining the 50% stopping ability for each subject from the range of sessions in which the stop-signal onset was varied from baseline (full details of this calculation can be found in the Supporting Information and (DAVIES *et al.* 2014)). The proportion of correct go-responses, and latency to respond, were also assessed. Additional measurements that related to general motoric competence and motivation within the task were also monitored, including the "initiation" and "magazine" latencies, the time taken to initiate a trial and the time taken to collect the reward. Also measured was the number of trials completed for any given session.

Immunohistochemistry

Dual-labelling immunofluorescence analysis of Nesp55 co-localisation with Grb10 was carried out on brain sections from *Grb10$^{+/p}$* mice in order to stain for the Lac-Z reporter gene, which is expressed in place of *Grb10* and has been used as a faithful proxy for Grb10 expression previously (GARFIELD *et al.* 2011). *Grb10$^{+/p}$* mice were transcardially perfused using 10% formalin (Sigma-Aldrich, UK) and whole brains dissected, post-fixed and equilibrated with 30% sucrose in PBS. Brains were sectioned into 40μm coronal slices using

a freezing microtome. Sections were washed three times for 10 min each in 0.1% PBS before

being incubated for 15 min in 0.3 M glycine in 0.1% PBS at room temperature, to neutralise

endogenous aldehyde groups. Sections were washed, as before, in 0.1% PBS and then

incubated at room temperature for 1 hour in 10% blocking solution; 0.5% BSA (BB

International, Cardiff, UK), 0.5% Triton X-100 (v/v, Sigma Aldrich) in 0.1% PBS. Sections were

then transferred to a 1% blocking solution containing a β-galactosidase (β-gal) specific

antibody (Abcam, U.K.), used at a 1:1000 dilution; and a Nesp55 primary antibody (1:1000).

The Nesp-55 primary antibody was generated in house, and is a rabbit anti-Nesp55

polycolonal antibody recognising the free terminal end (GAIPIRRH) of Nesp55. It has been

successfully characterised and used previously (Isсhɪᴀ *et al.* 1997). Sections were incubated

overnight at 4°C whilst gently shaking. Sections were then washed three times for 10

minutes in 0.1% PBS and then incubated with the appropriate fluorescent secondary

antibodies (Alexa Fluor; Life technologies) (1:1000) in 1% blocking solution in the dark at

room temperature for 2 hours, whilst gently shaking. Sections were then washed in 0.1%

PBS as before (in the dark) and transferred to polysine coated slides and allowed to dry

over-night in a dark dust-free environment. The mounted slides were then dehydrated

through a process of incubation in a rising concentration of alcohol, followed by xylene,

then cover-slipped and sealed using DPX (Raymond Lamb DPX), and allowed to dry over-

night. To control for non-specific binding of the secondary antibodies, secondary-only

negative controls were carried out alongside all experiments. Immunofluorescence slides

were viewed and images captured using an upright fluorescence microscope (Leica DM5000

B). Dual-labelled immunofluorescence images were acquired through separate channels for

different wavelengths (488 and 568 nm) then subsequently merged using ImageJ

(Image>colour>merge channels).

Data analysis and statistics

All behavioral data were analysed using SPSS 20 (SPSS, USA). Data were assessed for

normality and then analysed by Student's t-test or mixed ANOVA, with between-subjects

factors of GENOTYPE ($Grb10^{+/p}$ vs. WT), and within-subject factors DELAY (1s,8s,16s, or

1s,1s,1s), CHOICE (choice of large or small reward during forced trials of the delayed

reinforcement task), and STOP-SIGNAL POSITION (position of stop-signal relative to

individualized Go-response). For repeated-measures analyses, Mauchly's test of sphericity

of the covariance matrix was applied; significant violations from the assumption of

sphericity were subject to the Huynh–Feldt correction to allow more conservative

comparisons through adjusted degrees of freedom. Variables that were expressed as a

percentage (Delayed reinforcement, choice bias; SSRT, % correct) were subjected to an

arcsine transformation in order to limit the effect of an artificially imposed ceiling. All

significance tests were performed at alpha level of 0.05. Effect sizes (partial η^2) were

reported for the main measures in all tasks.

ACKNOWLEDGMENTS

This work was funded by a Leverhulme Trust project grant (F/00 407/BF) to ARI and JFW,

which supported CLD. The work was also funded by the MRC Centre for Neuropsychiatric

Genetics and Genomics (G0801418), of which ARI, TH and LSW are members, and which also

supported KL via a PhD studentship.

REFERENCES

Arnaud, P., D. Monk, M. Hitchins, E. Gordon, W. Dean *et al.*, 2003 Conserved methylation imprints in the human and mouse GRB10 genes with divergent allelic expression suggests differential reading of the same mark. Hum Mol Genet 12**:** 1005-1019.

Bari, A., D. M. Eagle, A. C. Mar, E. S. Robinson and T. W. Robbins, 2009 Dissociable effects of noradrenaline, dopamine, and serotonin uptake blockade on stop task performance in rats. Psychopharmacology (Berl) 205**:** 273-283.

Brandvain, Y., J. Van Cleve, F. Ubeda and J. F. Wilkins, 2011 Demography, kinship, and the evolving theory of genomic imprinting. Trends in genetics : TIG 27**:** 251-257.

Charalambous, M., M. Cowley, F. Geoghegan, F. M. Smith, E. J. Radford *et al.*, 2010 Maternally-inherited Grb10 reduces placental size and efficiency. Dev Biol 337**:** 1-8.

Charalambous, M., F. M. Smith, W. R. Bennett, T. E. Crew, F. Mackenzie *et al.*, 2003 Disruption of the imprinted Grb10 gene leads to disproportionate overgrowth by an Igf2-independent mechanism. Proc Natl Acad Sci U S A 100**:** 8292-8297.

Constancia, M., G. Kelsey and W. Reik, 2004 Resourceful imprinting. Nature 432**:** 53-57.

Davies, W., T. Humby, S. Trent, J. B. Eddy, O. A. Ojarikre *et al.*, 2014 Genetic and pharmacological modulation of the steroid sulfatase axis improves response control; comparison with drugs used in ADHD. Neuropsychopharmacology 39**:** 2622-2632.

Dent, C. L., T. Humby, K. Lewis, A. Plagge, R. Fischer-Colbrie *et al.*, 2016 Impulsive choices in mice lacking imprinted Nesp55. Genes Brain Behav 15**:** 693-701.

Dent, C. L., and A. R. Isles, 2014 Brain-expressed imprinted genes and adult behaviour: the example of Nesp and Grb10. Mamm Genome 25**:** 87-93.

Fawcett, T. W., B. Fallenstein, A. D. Higginson, A. I. Houston, D. E. Mallpress *et al.*, 2014 The evolution of decision rules in complex environments. Trends Cogn Sci 18**:** 153-161.

Ferguson-Smith, A. C., 2011 Genomic imprinting: the emergence of an epigenetic paradigm. Nature reviews. Genetics 12**:** 565-575.

Garfield, A. S., M. Cowley, F. M. Smith, K. Moorwood, J. E. Stewart-Cox *et al.*, 2011 Distinct physiological and behavioural functions for parental alleles of imprinted Grb10. Nature 469**:** 534-538.

Haig, D., 2000 Genomic imprinting, sex-biased dispersal, and social behavior. Ann N Y Acad Sci 907**:** 149-163.

Haig, D., and C. Graham, 1991 Genomic imprinting and the strange case of the insulin-like growth-factor II receptor. Cell 64**:** 1045-1046.

Hikichi, T., T. Kohda, T. Kaneko-Ishino and F. Ishino, 2003 Imprinting regulation of the murine Meg1/Grb10 and human GRB10 genes; roles of brain-specific promoters and mouse-specific CTCF-binding sites. Nucleic acids research 31**:** 1398-1406.

Ho, M. Y., S. Mobini, T. J. Chiang, C. M. Bradshaw and E. Szabadi, 1999 Theory and method in the quantitative analysis of "impulsive choice" behaviour: implications for psychopharmacology. Psychopharmacology (Berl) 146**:** 362-372.

Humby, T., J. B. Eddy, M. A. Good, A. C. Reichelt and L. S. Wilkinson, 2013 A novel translational assay of response inhibition and impulsivity: effects of prefrontal cortex lesions, drugs used in ADHD, and serotonin 2C receptor antagonism. Neuropsychopharmacology 38**:** 2150-2159.

Humby, T., F. M. Laird, W. Davies and L. S. Wilkinson, 1999 Visuospatial attentional functioning in mice: interactions between cholinergic manipulations and genotype. European Journal of Neuroscience 11**:** 2813-2823.

Ischia, R., P. Lovisetti-Scamihorn, R. Hogue-Angeletti, M. Wolkersdorfer, H. Winkler *et al.*, 1997 Molecular cloning and characterization of NESP55, a novel chromogranin-like precursor of a peptide with 5-HT1B receptor antagonist activity. J Biol Chem 272**:** 11657-11662.

Isles, A. R., T. Humby, E. Walters and L. S. Wilkinson, 2004 Common genetic effects on variation in impulsivity and activity in mice. J Neurosci 24**:** 6733-6740.

Isles, A. R., T. Humby and L. S. Wilkinson, 2003 Measuring impulsivity in mice using a novel operant delayed reinforcement task: effects of behavioural manipulations and d-amphetamine. Psychopharmacology (Berl) 170**:** 376-382.

Lang, M., T. Lemenager, F. Streit, M. Fauth-Buhler, J. Frank *et al.*, 2016 Genome-wide association study of pathological gambling. Eur Psychiatry 36**:** 38-46.

Madon-Simon, M., M. Cowley, A. S. Garfield, K. Moorwood, S. R. Bauer *et al.*, 2014 Antagonistic roles in fetal development and adult physiology for the oppositely imprinted Grb10 and Dlk1 genes. BMC Biol 12**:** 771.

Monk, D., P. Arnaud, J. Frost, F. A. Hills, P. Stanier *et al.*, 2009 Reciprocal imprinting of human GRB10 in placental trophoblast and brain: evolutionary conservation of reversed allelic expression. Hum Mol Genet 18**:** 3066-3074.

Moore, T., and D. Haig, 1991 Genomic imprinting in mammalian development - a parental tug-of-war. Trends in Genetics 7**:** 45-49.

Peters, J., 2014 The role of genomic imprinting in biology and disease: an expanding view. Nat Rev Genet 15**:** 517-530.

Plagge, A., A. R. Isles, E. Gordon, T. Humby, W. Dean *et al.*, 2005 Imprinted nesp55 influences behavioral reactivity to novel environments. Mol Cell Biol 25**:** 3019-3026.

Sanchez-Roige, S., P. Fontanillas, S. L. Elson, A. Pandit, E. M. Schmidt *et al.*, 2018 Genome-wide association study of delay discounting in 23,217 adult research participants of European ancestry. Nat Neurosci 21**:** 16-18.

Schonberg, T., C. R. Fox and R. A. Poldrack, 2011 Mind the gap: bridging economic and naturalistic risk-taking with cognitive neuroscience. Trends in cognitive sciences 15**:** 11-19.

Smith, F. M., L. J. Holt, A. S. Garfield, M. Charalambous, F. Koumanov *et al.*, 2007 Mice with a disruption of the imprinted Grb10 gene exhibit altered body composition, glucose homeostasis, and insulin signaling during postnatal life. Molecular and cellular biology 27**:** 5871-5886.

Wilkins, J. F., and D. Haig, 2003 What good is genomic imprinting: the function of parent-specific gene expression. Nat Rev Genet 4**:** 359-368.

Wilkins, J. F., F. Ubeda and J. Van Cleve, 2016 The evolving landscape of imprinted genes in humans and mice: Conflict among alleles, genes, tissues, and kin. Bioessays 38**:** 482-489.

Supplementary 1. Results for additional control measures in the Delay discounting task (Figure 1 and Table1) and SSRT task (Figure 2); further Methodological detail relating to SSRT measure.

Impulsive choice in mice lacking paternal expression of *Grb10* provides evidence for intra-genomic conflict in behaviour

Claire L. Dent[a], Trevor Humby[b], Katie Lewis[a], Andrew Ward[c], Reiner Fischer-Colbrie[d], Lawrence S. Wilkinson[a,b], Jon F Wilkins[e] & Anthony R. Isles[a]

[a]Behavioural Genetics Group, MRC Centre for Neuropsychiatric Genetics and Genomics, Neuroscience and Mental Health Research Institute, Cardiff University, Hadyn Ellis Building, Maindy Road, Cardiff, United Kingdom

[b]Behavioural Genetics Group, School of Psychology, Cardiff University, Tower Building, Cardiff, United Kingdom

[c]Department of Biology and Biochemistry, University of Bath, Bath, United Kingdom

[d]Department of Pharmacology, Innsbruck Medical University, Innsbruck, Austria

[e]Ronin Institute, Montclair, NJ 07043, USA

*Author for correspondence: Anthony R. Isles, E-mail IslesAR1@cardiff.ac.uk

SUPPLEMENTARY MATERIALS AND METHODS

Calculating the Stop Signal Reaction Time

Correct go reaction times were determined directly, however the stop-signal reaction time (SSRT) had to be derived from the distribution of correct go reaction times and the proportion of correctly stopped trials. SSRTs in the task were estimated employing the standard procedure previously described (LOGAN et al. 1984), using data from where the proportion of correct stop responses was ~50%. For each subject, data from the sessions in which the stop-signal positions were varied relative to the individualised go reaction time, were ranked by the proportion of correct stop responses, and data from sessions in which this value was between 40% and 60% (i.e. 50%±10%) were averaged. The latency of stopping as defined by the SSRT was derived from the distribution of correct go reaction times and the proportion of correctly stopped trials as previously described (LOGAN 1994; EAGLE AND ROBBINS 2003). Hence, for each of the sessions determined above, the correct go reaction times were rank ordered from smallest to largest and the nth value found, where n is the rank order position based on the proportion of failing to stop correctly in stop trials was corrected for the occurrence of omitted go trials. Although omitted go trials were a rare occurrence, this correction was implemented based on the rationale that they could alter the observed inhibition function and affect determination of the nth correct go reaction time value and hence the final SSRT (TANNOCK et al. 1989; SOLANTO et al. 2001; EAGLE AND ROBBINS 2003). To determine the SSRT, the time the stop-signal was presented (i.e. 'mean correct go reaction time' x '% mean stop-signal position') was subtracted from the nth correct go reaction time value.

SUPPLEMENTARY RESULTS

Additional measures in the delayed reinforcement task

In addition to choice, a number of other measures of basic behaviour within the delayed reinforcement task were obtained. These relate to general motoric competence and motivation within the task. As expected, $Grb10^{+/p}$ and wild-type (WT) littermates demonstrated a general main effect of DELAY on all of these measures as follows: Start (Figure S1a; $F_{1.4,26.8}=11.76$, $p=0.001$) and choice latencies (Figure S1b; $F_{1.8,33.5}=31.5$, $p<0.001$); and non-started trials (Figure S1c; $F_{1.6,30}=21.95$, $p<0.001$) and omitted trials ($F_{1.3,24.6}=12.48$, $p<0.001$). In all of these measures $Grb10^{+/p}$ and WT littermate mice showed a high degree of stimulus control, with no GENOTYPE differences or interactions between GENOTYPE and DELAY (see Table S1). Data for Start latency, Choice latency and non-started trials can be found in Figure S1. Omitted trials occurred very infrequently (Grand mean 0.21, ±0.05 SE), and are not shown.

Figure S1 Additional behavioural measures of performance of $Grb10^{+/p}$ and WT mice in the delayed reinforcement task. a Start latency; **b** Choice latency; **c** Non-started trials; **d** Panel pushes. All data represent means ±s.e.m.

Measure	Main effect of GENOTYPE		Interaction: GENOTYPE and DELAY	
	F	p	F	p
Start latency	2.63	0.12	0.58	0.51
Choice latency	0.45	0.51	0.10	0.88
Non-started trials	0.35	0.56	0.06	0.94
Omitted trials	0.01	0.93	0.62	0.48

Table S1 Statistical values for additional measures of performance of _Grb10_$^{+/p}$ and WT littermate controls within the delayed-reinforcement task. Analysis of main effect of GENOTYPE, and interaction between GENOTYPE and DELAY.

Additional measures in the SSRT task

In addition to the main measures, a number of other measures of basic behavior within the SSRT task were obtained (Figure S3). There were no differences between _Grb10_$^{+/p}$ and WT littermates in the number of trials completed (t_{20}=0.01, p=0.99), latency to initiate a trial (t_{20}=-0.54, p=0.60) and magazine latency (t_{20}=0.99, p=0.33).

Figure S2 Additional behavioural measures of performance of *Grb10*[+/p] and WT mice in the SSRT task. a Number of trials in a session with individualised SSRT (when the stop cue is presented 50% into the individual's Go-reaction time); **b** Initiation and magazine latency in a session with individualised SSRT. All data represent means ±s.e.m.

REFERENCES

Eagle, D. M., and T. W. Robbins, 2003 Lesions of the medial prefrontal cortex or nucleus accumbens core do not impair inhibitory control in rats performing a stop-signal reaction time task. Behav Brain Res 146**:** 131-144.

Logan, G., 1994 On the ability to inhibit thought and action: A users' guide to the stop signal paradigm., pp. 189-239 in *Inhibitory processes in attention, memory, and language.* , edited by D. Dagenbach and T. Carr. Academic Press, San Diego.

Logan, G. D., W. B. Cowan and K. A. Davis, 1984 On the ability to inhibit simple and choice reaction time responses: a model and a method. J Exp Psychol Hum Percept Perform 10**:** 276-291.

Solanto, M. V., H. Abikoff, E. Sonuga-Barke, R. Schachar, G. D. Logan *et al.*, 2001 The ecological validity of delay aversion and response inhibition as measures of impulsivity in AD/HD: a

supplement to the NIMH multimodal treatment study of AD/HD. J Abnorm Child Psychol 29**:** 215-228.

Tannock, R., R. J. Schachar, R. P. Carr, D. Chajczyk and G. D. Logan, 1989 Effects of methylphenidate on inhibitory control in hyperactive children. J Abnorm Child Psychol 17**:** 473-491.